# Fourier expansion along geodesics on Riemann surfaces

Central European Journal of Mathematics 12, No 4, 559-573 (2014)

#### Anton Deitmar

Institute of Mathematics
University of Tuebingen
Auf der Morgenstelle 10
72076 Tuebingen, GERMANY.

**Abstract.** For an eigenfunction of the Laplacian on a hyperbolic Riemann surface, the coefficients of the Fourier expansion are described as intertwining functionals. All intertwiners are classified. A refined growth estimate for the coefficients is given and a summation formula is proved.

## Contents

1	Generalized period integrals	3
2	Intertwining functionals	6
3	Fourier expansion of Maaß forms	17
4	Triple products	18

#### Introduction

For an automorphic function, the invariance under parabolic elements is used to give the standard Fourier expansion, the coefficients of which define the L-function of the form. In this paper, we instead consider the Fourier-expansion along a hyperbolic element. In other terms, let Y be a hyperbolic Riemann surface and let c be a closed geodesic in Y. We are interested in the Fourier coefficients

$$c_k(f) = \int_0^1 f(c(l_c t)e^{2\pi ikt} dt)$$

of a smooth function  $f \in C^{\infty}(Y)$ . Here  $l_c$  is the length of the geodesic c. Under the assumption that f be an eigenfunction of the Laplace operator on Y with eigenvalue  $\alpha$ , one can relate  $c_k$  to an intertwining integral  $I_k^{\alpha}(f)$ , which depends on  $\alpha$  and f, but not on c. There is an automorphic coefficient  $a_k \in \mathbb{C}$ , such that

$$c_k = a_k I_k^{\alpha}$$
.

The present paper contains three main results:

- in Theorem 2.3 one finds a classification of all intertwining functionals on the dual of the group  $PGL_2(\mathbb{R})$ .
- In Theorem 2.5 there is given the growth estimate

$$a_k = O(|k|^{\frac{1}{2}})$$

as  $|k| \to \infty$ . The proof uses the technique of analytic continuation developed by Bernstein and Reznikov in [BR99].

• In Theorem 4.2 finally, a summation formula is proved, which involves the coefficients  $a_k$  and the spectral decomposition in the compact case. The proof relies on the uniqueness of invariant trilinear forms as in [BR04]. The sum formula is of the form

$$\sum_{k \in \mathbb{Z}} |a_k|^2 \hat{\alpha}(k) = \sum_j c_j \int_{\mathbb{R}^2} W_j(t, x) \alpha(\hat{t}_x) dt dx,$$

where  $\alpha$  is a test function, the decomposition of the G-representation on  $L^2(\Gamma \backslash G)$  is  $\bigoplus_j \pi_j$  and the constants  $c_j$  and the explicit functions  $W_j$  depend on  $\pi_j$ . Finally  $\hat{t}_x = \frac{1}{2} \log \left| \frac{(e^{2t} + x)(x-1)}{(e^{2t} + x - 1)x} \right|$ . It is hoped that the choice of specific test functions will lead to more precise growth estimates for the  $a_k$ .

We explain the construction of the factors  $a_k$  in a bit more detail. Let X be the universal covering of Y and  $\Gamma$  its fundamental group. Then  $\Gamma$  acts on X by isometries and Y is the quotient  $\Gamma \backslash X$ . So  $\Gamma$  injects into the isometry group G of X, which acts transitively on X, i.e.,  $X \cong G/K$  for a maximal compact subgroup K. Let  $(\pi, V_{\pi})$  be an irreducible unitary representation of the group G and let  $\eta: V_{\pi} \to L^2(\Gamma \backslash G)$  be an isometric linear G-map. Let  $P_K: L^2(\Gamma \backslash G) \to L^2(\Gamma \backslash G)^K = L^2(\Gamma \backslash G/K) = L^2(Y)$  denote the orthogonal projection onto the subspace of K-invariants. Demanding that  $f \in C^{\infty}(Y)$  be an eigenfunction of the Laplacian amounts to the same as demanding f to lie in the image of  $P_K \circ \eta$  for some  $\pi$  and some  $\eta$ . The functional  $I_k^{\gamma} = c_k \circ P_K \circ \eta$  on  $V_{\pi}$  then has an intertwining property with respect to a split torus A inside G. By a uniqueness result, proven in Section 2, this implies that  $I_k^{\gamma}$  is a multiple of a standard intertwiner  $I_{\pi,k}^{\text{st}}$  on  $V_{\pi}$ , which we named  $I_k^{\alpha}$  above. So we get the existence of a factor  $a_k \in \mathbb{C}$  with  $I_k^{\gamma} = a_k I_{\pi,k}^{\text{st}} = a_k I_k^{\alpha}$  as above.

In Section 1 we describe the setting in greater precision. In Section 2 we classify the intertwining functionals that show up in the context and define the standard intertwiners that give rise to the factors  $a_k$  above. We also show the growth estimate of the factors  $a_k$ . In Section 3 we show how the Fourier expansion along a geodesic expands to an expansion on the whole space and in Section 4 we show how to derive the summation formula from the uniqueness of triple products.

## 1 Generalized period integrals

In this paper we use the group  $\operatorname{GL}_2(\mathbb{R})$ , the elements of which we write as matrices  $\binom{a\ b}{c\ d}$  and the group  $G = \operatorname{PGL}_2(\mathbb{R}) = \operatorname{GL}_2(\mathbb{R})/\mathbb{R}^{\times}$ , the elements of which we write in the form  $\begin{bmatrix} a\ b\\ c\ d \end{bmatrix}$ . Here we will usually arrange the determinant of  $\binom{a\ b}{c\ d}$  to be 1 or -1. The connected component of G is  $G^0 = \operatorname{PSL}_2(\mathbb{R}) = \operatorname{SL}_2(\mathbb{R})/(\pm 1)$ . The group  $G^0$  acts on the upper half plane  $\mathbb{H}$  in  $\mathbb{C}$  by linear fractionals and this action extends to an action of G in a way that G is identified with the group of all hyperbolic isometries on  $\mathbb{H}$ . The stabilizer in G of the point  $i \in \mathbb{H}$  is the maximal compact subgroup  $K = \operatorname{PO}(2) = \operatorname{O}(2)/\pm 1$  of G. So  $\mathbb{H}$  is identified with G/K.

Let  $\Gamma$  be a discrete subgroup of the group G. Later we will assume  $\Gamma$  to be of finite covolume. For simplicity, we will assume  $\Gamma$  to be torsion-free and that  $\Gamma \subset G^0$ . This implies that  $\Gamma$  is the fundamental group of  $Y = \Gamma \backslash \mathbb{H}$  and

the latter is a Riemann surface equipped with the hyperbolic metric.

For a closed geodesic c in Y let l(c) denote its length. The period integral  $I_c(f) = \int_0^{l(c)} f(c(t)) dt$  is the zeroth coefficient of the Fourier-expansion of the function  $t \mapsto f(c(t))$ . Therefore the higher coefficients can be viewed as "generalised period integrals".

The real Lie-algebra of G is  $\mathfrak{g}_{\mathbb{R}} = sl_2(\mathbb{R})$ , the Lie-algebra of all real  $2 \times 2$  matrices of trace zero. For  $X,Y \in \mathfrak{g}_{\mathbb{R}}$  let  $b(X,Y) = \frac{1}{2}\mathrm{tr}(XY)$ . Then b is an invariant symmetric bilinear form. Let  $\mathfrak{k}_{\mathbb{R}} \subset \mathfrak{g}_{\mathbb{R}}$  be the Lie algebra of K. Then b is negative definite on  $\mathfrak{k}_{\mathbb{R}}$  and positive definite on its orthogonal complement  $\mathfrak{p}_{\mathbb{R}}$ . Let  $\mathfrak{a}_{\mathbb{R}} = \mathbb{R} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ , and let  $A = \exp(\mathfrak{a}_{\mathbb{R}})$  be the corresponding subgroup of G. Let  $\mathfrak{a}_{\mathbb{R}}^+ = \mathbb{R}_{>0} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$  be the positive cone and let  $A^+ = \exp(\mathfrak{a}_{\mathbb{R}}^+)$ . Then A is closed and non-compact and its centralizer in G is the group AM of all diagonal matrices in G. Here M is the two element group generated by  $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ . Let  $\mathfrak{a}$  be the complexification of  $\mathfrak{a}_{\mathbb{R}}$  and  $\mathfrak{a}^*$  the dual space of  $\mathfrak{a}$ . Then  $\mathfrak{a}^*$  can be identified with the set of continuous homomorphisms from A to  $\mathbb{C}^{\times}$ . For  $\lambda \in \mathfrak{a}^*$  we write  $a \mapsto a^{\lambda} = e^{\lambda(\log a)}$  for the corresponding homomorphism. It is known that  $G^0 \to G/K$  can be identified with the sphere-bundle of  $\mathbb{H} = G/K$  in a way that the geodesic flow is given by

$$\phi_t(g) = g \exp(tH_1), \qquad g \in G^0,$$

where  $H_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ . The sphere bundle SY equals  $\Gamma \backslash G^0$ .

A closed geodesic c in Y gives rise to a conjugacy class  $[\gamma]$  in  $\Gamma$  of elements which "close" c. Any such  $\gamma \in \Gamma$  is hyperbolic in the sense that it is conjugate in G to an element of the form  $a_{t_0} = \operatorname{diag}(e^{t_0}, e^{-t_0}) \in A \setminus \{1\}$ . We insist that  $a_{t_0} \in A^+$ , i.e.,  $t_0 > 0$  to make it unique. We now assume that  $\gamma$  be primitive, i.e.,  $\gamma$  is no power  $\tau^n$  for any  $\tau \in \Gamma$  and  $n \geq 2$ . This is equivalent to the geodesic c being primitive, i.e., c is no power of any shorter geodesic. The characters of the compact abelian group  $A/\langle a_{\gamma} \rangle$  are given by those  $\mu \in \mathfrak{a}^*$  with  $a_{\gamma}^{\mu} = 1$ , i.e.,  $\mu(\log a_{\gamma}) \in 2\pi i\mathbb{Z}$ . Let  $\mu_{\gamma}$  be the unique element of  $\mathfrak{a}^*$  with  $\mu_{\gamma}(\log a_{\gamma}) = 2\pi i$ . Then  $A/\langle a_{\gamma} \rangle = \mathbb{Z}\mu_{\gamma}$ . Later we will use the notation

$$\tilde{\mu} = \frac{1}{2\pi i} \mu.$$

Fix an element  $\sigma = \sigma_{\gamma} \in G^0$  with  $\gamma = \sigma^{-1}a_{t_0}\sigma$ . Let  $f \in C^{\infty}(\Gamma \backslash G) = C^{\infty}(\Gamma \backslash G)$  and set

$$f^{\sigma}(x) = f(\sigma x).$$

Then the map  $t \mapsto f^{\sigma}(a_t x)$  with  $a_t = \text{diag}(e^t, e^{-t})$  is periodic of period  $t_0$  and thus has a Fourier-expansion

$$f^{\sigma}(a_t x) = \sum_{k \in \mathbb{Z}} e^{2\pi i t/t_0} \frac{1}{t_0} \int_0^{t_0} f^{\sigma}(a_t x) e^{-2\pi i t/t_0} dt.$$

For  $k \in \mathbb{Z}$  let

$$I_k^{\gamma}: C^{\infty}(\Gamma \backslash G) \to \mathbb{C}$$

$$f \mapsto \frac{1}{t_0} \int_0^{t_0} f^{\sigma}(a_t) e^{-2\pi i k t / t_0} dt.$$

Note that  $I_k^{\gamma}$  depends on the choice of  $\sigma$ . Geometrically, this corresponds to choosing a base-point on the closed orbit c. This dependence is not severe, as  $\sigma$  is determined up to multiplication from the right by elements of A. If we replace  $\sigma$  by  $\sigma a_0$ , then  $I_k^{\gamma}$  is replaced by  $a_0^{k\mu\gamma}I_k^{\gamma}$ . So in particular, the absolute value  $|I_k^{\gamma}|$  is uniquely determined by k and  $\gamma$ . Further, if  $\gamma$  is replaced by a  $\Gamma$ -conjugate, say  $\gamma' = \tau \tau^{-1}$  then one can choose  $\sigma_{\gamma'}$  to be equal to  $\tau \sigma_{\gamma}$  and with this choice one gets  $I_k^{\gamma'} = I_k^{\gamma}$ .

The form b determines a Haar measure dg on G. Let R denote the unitary G-representation on  $L^2(\Gamma \backslash G)$  given by right translations. We are particularly interested in the subspace  $L^2_{\text{cusp}}(\Gamma \backslash G)$  of cusp forms. This representation space decomposes discretely,

$$L^2_{\text{cusp}}(\Gamma \backslash G) \cong \bigoplus_{\pi \in \hat{G}} N_{\Gamma}(\pi)\pi,$$

where the sum runs over the unitary dual  $\hat{G}$  of G and the multiplicities  $N_{\Gamma}(\pi)$  are finite. Here and later we understand the direct sum to be a completed direct sum in the appropriate topology. We define  $C_{\text{cusp}}^{\infty}(\Gamma \backslash G)$  to be the intersection of  $C^{\infty}(\Gamma \backslash G)$  with the space of cusp forms. Then it turns out that  $C_{\text{cusp}}^{\infty}(\Gamma \backslash G)$  is the set of smooth vectors in the representation space  $L_{\text{cusp}}^2(\Gamma \backslash G)$ , i.e.,

$$C_{\mathrm{cusp}}^{\infty}(\Gamma \backslash G) = L_{\mathrm{cusp}}^{2}(\Gamma \backslash G)^{\infty} = \bigoplus_{\pi \in \hat{G}} N_{\Gamma}(\pi)\pi^{\infty},$$

where  $\pi^\infty$  is the representation on the Fréchet space of smooth vectors. The linear functional  $I_k^\gamma$  satisfies

$$I_k^{\gamma}(R(a)\varphi) = a^{k\mu\gamma} I_k^{\gamma}(\varphi)$$

for every  $a \in A$ . This means that  $I_k^{\gamma}$  is an intertwining functional.

## 2 Intertwining functionals

We first shall give a description of the admissible and unitary duals of the group G. For any topological group G, let  $\widehat{G}$  denote the *unitary dual* of G, that is, the set of unitary equivalence classes of irreducible unitary representations of G.

Next let G denote a semisimple Lie group with finite center and finitely many connected components. Then G has a maximal compact subgroup K which is unique up to conjugation. A representation  $(\pi, V_{\pi})$  of G is called admissible, if for every  $\tau \in \widehat{K}$  the isotype  $V_{\pi}(\tau)$  is finite-dimensional. In that case the space  $V_{\pi,K}$  of K-finite vectors in  $V_{\pi}$  forms a  $(\mathfrak{g},K)$ -module, where  $\mathfrak{g}$  is the complexified Lie algebra of G. Two admissible representations are called infinitesimally equivalent if their  $(\mathfrak{g},K)$ -modules of K-finite vectors are isomorphic. The admissible dual  $\widehat{G}_{\mathrm{adm}}$  of G is the set of infinitesimal equivalence classes of irreducible admissible representations of G.

Due to results of Harish-Chandra, every irreducible unitary representation of G is admissible and two unitary admissible representations are unitarily equivalent if and only if they are infinitesimally equivalent. Thus the unitary dual  $\widehat{G}$  can be considered a subset of the admissible dual  $\widehat{G}_{\rm adm}$ .

Now consider  $G = \operatorname{PGL}_2(\mathbb{R})$ . There is a canonical character

$$\chi: G \to \{\pm 1\}; \qquad g \mapsto \operatorname{sign}(\det(g)),$$

taking the values  $\pm 1$  and having the connected component  $G^0$  for kernel. For a representation  $\pi$  of G we define the  $\chi$ -twist  $\chi\pi$  of  $\pi$ , also written  $\chi\otimes\pi$  as the representation with the same space  $V_{\pi}$  as  $\pi$  but defined as

$$\chi \pi(x) = \chi(x) \pi(x)$$
.

Let P denote the parabolic subgroup of G consisting of all upper triangular matrices. Then P=MAN, where N is the group of all upper triangular matrices with ones on the diagonal. For  $\lambda\in\mathbb{C}$  and  $a=\left[\begin{smallmatrix}e^t\\e^{-t}\end{smallmatrix}\right]\in A$  we write

$$a^{\lambda} = e^{\lambda t}$$
.

Let  $\pi_{\lambda}$  be the corresponding principal series representation, which we normalize to live on the space of functions  $\varphi: G \to \mathbb{C}$  satisfying  $\varphi(manx) = a^{\lambda+1}\varphi(x)$ . Let  $V_{\lambda}$  be the space of  $\pi_{\lambda}$  and let  $V_{\lambda}^{\infty}$  be the space of smooth

7

vectors in it. These can be viewed as smooth sections of the line bundle  $E_{\lambda}$  over  $P \backslash G$  given by the P-representation  $(man) \mapsto a^{\lambda+1}$ . Note that restriction of functions to K identifies  $V_{\lambda}^{\infty}$  with the space  $C^{\infty}(M \backslash K)$ , so in particular, for  $\varphi \in V_{\lambda}^{\infty}$ , the function  $\varphi|_{K}$  is independent of  $\lambda$ .

**Proposition 2.1.** Let  $G = \operatorname{PGL}_2(\mathbb{R})$ . The admissible dual  $\widehat{G}_{\operatorname{adm}}$  of G consists of

- (a)  $\pi_{\lambda}$ ,  $\lambda \in \mathbb{C}$ ,  $\lambda \notin 1 + 2\mathbb{Z}$ ,
- (b)  $\mathcal{D}_{2n}$  for  $n = 1, 2, 3, \ldots$  the standard discrete series representations,
- (c)  $\delta_m$  for  $m = 0, 2, 4, \ldots$  where  $\delta_m$  is the (m+1)-dimensional representation on the space of all homogeneous polynomials p(X,Y) of degree m,

together with their  $\chi$ -twists. The only isomorphisms occurring are  $\pi_{it} \cong \pi_{-it}$  and  $\chi \pi_{it} \cong \chi \pi_{-it}$  for  $t \in \mathbb{R}$ .

The unitary dual  $\widehat{G}$  consists of

- (a)  $\pi_{\lambda}$  for  $\lambda \in i[0, \infty) \cup (0, 1)$ ,
- (b)  $\mathcal{D}_{2n}$  for  $n=1,2,3,\ldots$  the discrete series representations,
- (c)  $\delta_0$ ,

together with their  $\chi$ -twists.

For n = 1, 2, 3, ... we have the exact sequence of representations

$$0 \to \mathcal{D}_{2n} \to \pi_{2n-1} \to \delta_{2n-2} \to 0$$
,

and

$$0 \to \delta_{2n-2} \to \pi_{1-2n} \to \mathcal{D}_{2n} \to 0.$$

*Proof.* This can be deduced from the description of the unitary dual of  $SL_2(\mathbb{R})$  given, for example, in [Kna01].

Let  $T = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  be the non-trivial element of M. As  $T^2 = 1$ , for every representation  $(\pi, V_{\pi})$ , the space  $V_{\pi}$  splits as a direct sum  $V_{\pi} = V_{\pi}^+ \oplus V_{\pi}^-$ ,

where  $V_{\pi}^{\pm}$  is the  $\pm 1$ -eigenspace of  $\pi(T)$ . Twisting by  $\chi$  interchanges the roles of  $V_{\pi}^{+}$  and  $V_{\pi}^{-}$ .

Let  $\mu \in \mathfrak{a}^*$  and let  $(\pi, V_{\pi})$  be a representation of G. A continuous linear functional  $l: V_{\pi}^{\infty} \to \mathbb{C}$  is called a  $\mu$ -intertwiner, if

$$l(\pi(a)v) = a^{\mu}l(v)$$

holds for every  $a \in A$  and every  $v \in V_{\pi}$ . Let  $V_{\pi}^{\infty}(\mu)$  be the space of all  $\mu$ -intertwiners. Note that  $V_{\pi}^{\infty}(\mu) = V_{\chi\pi}^{\infty}(\mu)$ , where we consider the  $\chi$ -twist  $\chi\pi$  as a representation with the same representation space as  $\pi$ .

Let  $w_0 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  and  $n_0 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . Then  $w_0$  is a representative of the non-trivial element of the Weyl-group W(G, A).

The base space  $P \setminus G$  of the bundle  $E_{\lambda}$  consists of three orbits under the group AM, namely the open orbit  $[w_0n_0]$ , and the two closed orbits [1],  $[w_0]$  which are indeed points.

We now define a standard intertwiner on the representation  $\pi_{\lambda}$  for Re( $\lambda$ ) > -1. Let

$$I_{\lambda,\mu}^{\mathrm{st}}(\varphi) = I_{\pi_{\lambda},\mu}^{\mathrm{st}}(\varphi) = \int_{A} \varphi(w_0 n_0 a) a^{-\mu} da, \qquad \varphi \in V_{\lambda}^{\infty}.$$

If supp  $\varphi \subset [w_0 n_0]$ , then the integral  $I_{\lambda,\mu}^{\rm st}$  is extended over a compact set, hence it exists.

**Lemma 2.2.** Let  $\mu \in \mathbb{C}$ . If  $\operatorname{Re}(\lambda) > -1 - \operatorname{Re}(\mu)$ , then the integral  $I_{\lambda,\mu}^{\operatorname{st}}(\varphi)$  exists for every  $\varphi \in V_{\lambda}^{\infty}$  and defines a  $\mu$ -intertwiner. The map  $\lambda \mapsto I_{\lambda,\mu}^{\operatorname{st}}$  extends to a meromorphic operator-valued function with poles exactly at

$$\lambda = -\mu - 1 - 4k, \qquad k \in \mathbb{N}_0$$

and

$$\lambda = \mu - 3 - 4k, \qquad k \in \mathbb{N}_0.$$

Outside the poles,  $I_{\lambda,\mu}^{\text{st}}$  spans the one dimensional space of intertwiners supported on the open orbit. At the poles this space is zero.

*Proof.* Let  $\underline{a}: G \to A$ ,  $\underline{n}: G \to N$ ,  $\underline{k}: G \to K$  be the smooth maps defined by the Iwasawa decomposition  $g = ank = \underline{a}(g)\underline{n}(g)\underline{k}(g)$  for  $g \in G$ . For

 $\varphi \in V_{\lambda}^{\infty}$  one has

$$I_{\lambda,\mu}^{\text{st}}(\varphi) = \int_{A} \underline{a}(w_0 n_0 a)^{\lambda+1} \varphi(\underline{k}(w_0 n_0 a)) a^{-\mu} da$$
$$= \int_{A} \underline{a}(w_0 n_0^a)^{\lambda+1} a^{-\lambda-1-\mu} \varphi(\underline{k}(w_0 n_0^a)) da,$$

where  $n_0^a = a^{-1}n_0a$ . Noting that if  $a = \begin{bmatrix} e^t \\ e^{-t} \end{bmatrix}$ , then we have  $n_0^a = \begin{bmatrix} 1 & e^{-2t} \\ 1 \end{bmatrix}$ , we get  $w_0n_0^a = \begin{bmatrix} 0 & -1 \\ 1 & e^{-2t} \end{bmatrix}$  and thus  $\underline{k}(w_0n_0^a) = \frac{1}{\sqrt{e^{-4t}+1}} \begin{bmatrix} e^{-2t} & -1 \\ 1 & e^{-2t} \end{bmatrix}$ . We now define special test functions. For  $k, l \in \mathbb{N}_0$  let  $\varphi_{k,l} : \mathrm{SO}(2) \to \mathbb{C}$  be defined by

$$\varphi_{k,l}\begin{pmatrix} d & -c \\ c & d \end{pmatrix} = c^k d^l.$$

If k+l is even,  $\varphi_{k,l}$  defines an element of  $V_{\lambda}^{\infty}$ . The theory of Taylor-series tells us that every  $\varphi \in V_{\lambda}^{\infty}$  can be written as

$$\varphi = \sum_{0 \le k, l \le N} c_{k,l} \varphi_{k,l} + R_N(\varphi),$$

where  $c_{k,l} \in \mathbb{C}$  and the function  $R_N(\varphi) \in V_\lambda^\infty \cong C^\infty(M \setminus K)$  vanishes to order N at 1 and  $w_0$ . The integral  $I_{\lambda,\mu}^{\mathrm{st}}(\varphi)$ , as written above, makes sense for  $\varphi = \varphi_{k,l}$  also in the case when k+l is not even. We use this fact for convenience. For  $k,l \in \mathbb{N}_0$  we compute

$$I_{\lambda,\mu}^{\text{st}}(\varphi_{k,l}) = \int_{A} \underline{a}(w_{0}n_{0}^{a})^{\lambda+1} a^{-\lambda-1-\mu} \varphi_{k,l}(\underline{k}(w_{0}n_{0}^{a})) da$$

$$= \int_{\mathbb{R}} \underline{a} \begin{pmatrix} 0 & -1 \\ 1 & e^{-2t} \end{pmatrix}^{\lambda+1} e^{-(\lambda+1+\mu)t} \varphi_{k,l} \left(\underline{k} \begin{pmatrix} 0 & -1 \\ 1 & e^{-2t} \end{pmatrix} \right) dt$$

$$= \int_{\mathbb{R}} (e^{-4t} + 1)^{-\frac{\lambda+1}{2}} e^{-(\lambda+1+\mu)t} \varphi_{k,l} \left(\frac{1}{\sqrt{e^{-4t} + 1}} \begin{pmatrix} e^{-2t} & -1 \\ 1 & e^{-2t} \end{pmatrix} \right) dt$$

$$= \int_{\mathbb{R}} (e^{-4t} + 1)^{-\frac{\lambda+k+l+1}{2}} e^{-(\lambda+2l+1+\mu)t} dt$$

$$= \frac{1}{4} \int_{\mathbb{R}} (e^{-t} + 1)^{-\frac{\lambda+k+l+1}{2}} e^{-\frac{\lambda+2l+1+\mu}{4}t} dt$$

$$= \frac{1}{4} B \left(\frac{\lambda+2l+1+\mu}{4}, \frac{\lambda+2k+1-\mu}{4}\right),$$

where  $B(x,y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$  is Euler's Beta-function. We conclude

$$I_{\lambda,\mu}^{\mathrm{st}}(\varphi_{k,l}) = \frac{\Gamma\left(\frac{\lambda+2l+1+\mu}{4}\right)\Gamma\left(\frac{\lambda+2k+1-\mu}{4}\right)}{\Gamma\left(\frac{\lambda+k+l+1}{2}\right)}.$$

Note that

$$I_{\lambda,\mu}^{\mathrm{st}}(\varphi_{k+1,l+1}) = I_{\lambda+2,\mu}^{\mathrm{st}}(\varphi_{k,l}).$$

The space  $\varphi_{N,N}C^{\infty}(M\backslash K)$  is the space of all  $\varphi\in V_{\lambda}^{\infty}$  which vanish to order  $\geq N$  at 1 and  $w_0$ . For any  $\varphi$  in this space, the integral  $I_{\lambda,\mu}^{\rm st}(\varphi)$  converges if  $\mathrm{Re}(\lambda) > -N - 1 - \mathrm{Re}(\mu)$ . Therefore we get analytic continuation of the map  $\lambda \mapsto I_{\lambda,\mu}^{\rm st}$  as claimed. The lemma follows.

We next consider intertwiners which are supported on the closed orbits [1] and  $[w_0]$ . Let  $S_{[1],0}:V_\lambda^\infty\to\mathbb{C}$  denote the distribution

$$S_{[1],0}(\varphi) = \varphi(1).$$

Then  $S_{[1],0} \circ R(a) = a^{\lambda+1} S_{[1],0}$ , so  $S_{[1],0}$  is an  $\mu$ -intertwiner for  $\mu = \lambda + 1$ .

We next consider higher derivatives of this distribution. For  $X \in \mathfrak{g}$ , the Lie algebra of G, and  $f \in C^{\infty}(G)$ , we let

$$R_X f(y) = \frac{d}{dt}\Big|_{t=0} f(y \exp(tX)).$$

Let  $\bar{N}=\theta(N)$  and let  $\bar{\mathfrak{n}}_{\mathbb{R}}$  be its Lie algebra. Then the tangent space of  $P\backslash G$  at the unit is isomorphic to  $\bar{n}_{\mathbb{R}}$ . Let  $X=\left(\begin{smallmatrix} 0&1\\0&0\end{smallmatrix}\right)$  and  $\bar{X}=\left(\begin{smallmatrix} 0&0\\1&0\end{smallmatrix}\right)$ . Then  $\mathfrak{n}_{\mathbb{R}}=\mathbb{R}X$  and  $\bar{\mathfrak{n}}_{\mathbb{R}}=\mathbb{R}\bar{X}$ . For  $k\in\mathbb{N}$  set

$$S_{[1],k}(\varphi) = R_{\bar{X}}^k \varphi(1).$$

Then  $S_{[1],k} \circ R(a) = a^{\lambda+1+2k} S_{[1],k}$ . Since these span the space of all distributions supported at 1 we see that we get a non-zero  $\mu$ -intertwiner supported on 1 if and only if

$$\mu = \lambda + 1 + 2k$$

for some  $k \in \mathbb{N}_0$ . If this condition is satisfied, then the space of intertwiners supported on [1] is one dimensional.

We turn to the other closed orbit  $[w_0]$ . In this case we define

$$S_{[w_0],k}(\varphi) = R_X^k \varphi(w_0).$$

Then  $S_{[w_0],k} \circ R(a) = a^{-\lambda-1-2k} S_{[w_0],k}$  and we conclude that there exists a non-zero intertwiner supported on  $[w_0]$  if and only if

$$\mu = -\lambda - 1 - 2k, \qquad k \in \mathbb{N}_0,$$

in which case the space of intertwiners supported on  $[w_0]$  is one dimensional.

**Theorem 2.3.** (a) For  $\lambda, \mu \in \mathbb{C}$  with  $\mu \neq 0$  we have

$$\dim V_{\lambda}^{\infty}(\mu) = 1.$$

In this case, if  $\lambda \notin \pm \mu - 1 - 2\mathbb{N}_0$ , then  $V_{\lambda}^{\infty}(\mu)$  is spanned by  $I_{\lambda,\mu}^{\mathrm{st}}$ . If  $\lambda = \mu - 1 - 2k$  with  $k \in \mathbb{N}_0$ , then  $V_{\lambda}^{\infty}(\mu)$  is spanned by  $S_{[1],k}$  and if  $\lambda = -\mu - 1 - 2k$  with  $k \in \mathbb{N}_0$ , then  $V_{\lambda}^{\infty}(\mu)$  is spanned by  $S_{[w_0],k}$ .

The same holds for the  $\chi$ -twist.

(b) For  $\lambda \in \mathbb{C}$  and  $\mu = 0$  we have

$$\dim V_{\lambda}^{\infty}(0) = \begin{cases} 2 & \lambda \in -1 - 2\mathbb{N}_0, \\ 1 & otherwise. \end{cases}$$

If  $\lambda \notin -1 - 2\mathbb{N}_0$ , then  $V_{\lambda}^{\infty}(0)$  is spanned by  $I_{\lambda,0}^{\mathrm{st}}$ . If  $\lambda = -1 - 2k$  with  $k \in \mathbb{N}_0$ , then  $V_{\lambda}^{\infty}(0)$  is spanned by  $S_{[w_0],k}, S_{[1],k}$ .

The same holds for the  $\chi$ -twist.

(c) For the finite-dimensional representations we have dim  $\delta_{2n}(\mu) = 0$  if  $\mu \neq -2n, -2n + 2, \dots, 2n$  and

$$\dim \delta_{2n}(\mu) = 1$$
 if  $\mu \in \{-2n, -2n + 2, \dots, 2n\}$ .

(d) Let  $\mu \in \mathbb{C}$ ,  $n \in \mathbb{N}$ . Then we have an exact sequence

$$0 \to \mathcal{D}_{2n}^{\infty}(\mu) \to V_{1-2n}^{\infty}(\mu) \to \delta_{2n-2}(\mu) \to 0.$$

If  $\mu \notin \{-2n+2,\ldots,2n-2\}$ , then  $\delta_{2n-2}(\mu) = 0$  and therefore  $\mathcal{D}_{2n}^{\infty}(\mu) \cong V_{1-2n}^{\infty}(\mu)$  is one dimensional.

If  $\mu \in \{-2n+2,\ldots,2n-2\}$  but  $\mu \neq 0$ , then  $\mathcal{D}_{2n}^{\infty}(\mu) = 0$  and finally  $\mathcal{D}_{2n}^{\infty}(0)$  is one-dimensional and is spanned by  $S_{[1],n-1} - S_{[w_0],n-1}$ .

*Proof.* (a) and (b) are clear by the above. For (c) recall that  $\delta_{2n}$  has the basis  $e_{2j-2n}=X^jY^{2n-j}$  for  $j=0,\ldots 2n$  and the group A acts by  $\delta_{2n}(a)e_{2j-2n}=a^{2j-2n}e_{2j-2n}$ . This proves (c).

For (d) we consider the exact sequence

$$0 \to \delta_{2n-2} \to \pi_{1-2n} \to \mathcal{D}_{2n} \to 0$$
,

which induces the exact sequence of intertwiners

$$0 \to \mathcal{D}_{2n}^{\infty}(\mu) \to V_{1-2n}^{\infty}(\mu) \to \delta_{2n-2}(\mu).$$

This proves the first assertion, i.e., the case  $\delta_{2n-2}(\mu) = 0$ . If  $\mu \in \{-2n + 2, \dots, 2n-2\}$ , which means  $\delta_{2n-2}(\mu) \neq 0$ , ten we have to show that the map  $V_{1-2n}^{\infty}(\mu) \to \delta_{2n-2}(\mu)$  is non-zero, for it is automatically onto then, as the target space is one-dimensional. The above exact sequence dualizes to the exact sequence

$$0 \to \mathcal{D}_{2n} \to \pi_{2n-1} \to \delta_{2n-2} \to 0$$
,

which yields an exact sequence

$$0 \to \delta_{2n-2}(\mu) \to V_{2n-1}^{\infty}(\mu) \to \mathcal{D}_{2n}^{\infty}(\mu).$$

So the arrow  $\delta_{2n-2}(\mu) \to V_{2n-1}^{\infty}(\mu)$  is non-zero, hence its dual  $V_{1-2n}^{\infty}(\mu) \to \delta_{2n-2}(\mu)$  likewise.

Finally, for  $\mu=0$  we show that the kernel of the restriction map  $V_{1-2n}^{\infty}(0) \to \delta_{2n-2}(0)$  is spanned by  $S_{[w_0],n-1}-S_{[1],n-1}$ . Recall that  $R_X$  and  $R_{\bar{X}}$  are the weight-change operators in highest weight theory, so  $R_X^{n-1}e_0$  is a multiple of  $e_{2n-2}$  and  $R_{\bar{X}}^{n-1}e_0$  is a multiple of  $e_{2-2n}$ . Let  $\theta(x)=x^{-t}$  be the transpose followed by the inversion, i.e.,  $\theta$  is the Cartan-involution on G with fixed point set K. We have  $\theta(X)=\bar{X}$ . It is readily verified that the map  $\Psi$  with  $\Psi(f)(x)=f(w_0\theta(x))$  is an isomorphism between the representations  $\pi_\lambda$  and  $\pi_\lambda\circ\theta$ . Thus it follows that  $S_{[w_0],n-1}(\Psi(\varphi))=S_{[1],n-1}(\varphi)$ . As  $\Psi(\pi(a)\varphi)=\pi(a^{-1})\Psi(\varphi)$  and  $\Psi^2=\mathrm{Id}$  it follows that  $\Psi(e_0)=\pm e_0$ . We want to show  $\Psi(e_0)=e_0$ . For this we let  $\binom{d-c}{c-d}\in\mathrm{SO}(2)$  with c,d>0. The A-invariance of  $e_0$  shows that for a>0 we have

$$e_0\left(\frac{1}{\sqrt{a^2c^2+d^2/a^2}} \left( \frac{ad-c/a}{ac-d/a} \right) \right) = (c^2a^2+d^2/a^2)^{\frac{2-n}{2}} e_0 \left( \frac{d-c}{c-d} \right).$$

So the A-orbit is mapped to positive multiples of  $e_0 \begin{pmatrix} d & -c \\ c & d \end{pmatrix}$ . By the M-invariance we get on the other hand,

$$e_0\left(\left(\begin{smallmatrix} & -1 \\ 1 & \end{smallmatrix}\right)\left(\begin{smallmatrix} d & -c \\ c & d \end{smallmatrix}\right)\right) = e_0\left(\begin{smallmatrix} -c & -d \\ d & -c \end{smallmatrix}\right) = e_0\left(\begin{smallmatrix} c & -d \\ d & c \end{smallmatrix}\right).$$

As  $\binom{c-d}{d-c}$  lies in the same A-orbit as  $\binom{d-c}{c-d}$ , the claim follows. We therefore conclude that  $S_{[w_0],n-1}(e_0)=S_{[1],n-1}(e_0)$ .

A sequence  $(c_j)_{j\in\mathbb{N}}$  of complex numbers is said to be of moderate growth, if there exist  $N\in\mathbb{N}$  such that

$$|c_j| = O(j^N),$$

as  $j \to \infty$ . The sequence is called *rapidly decreasing*, if for every  $N \in \mathbb{N}$  one has

$$|c_j| = O(j^{-N})$$

as  $j \to \infty$ . The product of two moderately growing sequences is moderately growing and the product of a moderately growing sequence and a rapidly decreasing sequence is rapidly decreasing.

**Proposition 2.4.** Let  $\operatorname{Re}(\lambda) > -1$  and  $f \in V_{\lambda}^{\infty}$  as well as  $\mu \in i\mathbb{R} \setminus \{0\}$  be given. Then the sequence  $(I_{\lambda,k\mu}^{\operatorname{st}}(f))_{k\in\mathbb{Z}}$  is rapidly decreasing.

*Proof.* Recall  $H_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ . Using integration by parts, we compute for  $N \in \mathbb{N}$ ,

$$I_{\lambda,k\mu}^{\text{st}}(\varphi) = \int_{A} \varphi(w_0 n_0 a) a^{-k\mu} da$$

$$= \int_{\mathbb{R}} \varphi(w_0 n_0 \exp(tH_1) e^{-k\mu t} dt$$

$$= \left(\frac{1}{k\mu}\right)^N \int_{\mathbb{R}} R_{H_1}^N \varphi(w_0 n_0 a) a^{-k\mu} da.$$

As  $Re(\mu) = 0$  we have  $|a^{-k\mu}| = 1$  and the claim follows.

By an automorphic representation  $(\pi, V_{\pi}, \eta)$  we mean an irreducible unitary representation  $(\pi, V_{\pi})$  of G together with an isometric G-equivariant linear map  $\eta \colon V_{\pi} \to L^2(\Gamma \backslash G)$ . Then  $\eta$  maps the space  $V_{\pi}^{\infty}$  of smooth vectors into  $C^{\infty}(\Gamma \backslash G)$ . Let  $k \in \mathbb{Z}$ . The map  $I_k^{\gamma} \circ \eta$  is an intertwiner for  $k\mu_{\gamma}$ . So, for instance, let  $\pi = \pi_{\lambda} \in \widehat{G}$ , then the space of intertwiners on  $V_{\lambda}$  is spanned by  $I_{\lambda,k\mu_{\gamma}}^{\mathrm{st}}$ . Therefore there exists  $a_k^{\eta,\gamma} \in \mathbb{C}$  such that

$$I_k^{\gamma} \circ \eta = a_k^{\eta,\gamma} I_{\lambda,k\mu_{\gamma}}^{\mathrm{st}}.$$

However, if  $\pi = \mathcal{D}_{2n}$  is a discrete series representation and k = 0, then there are  $b_0^{\eta,\gamma}, c_0^{\eta,\gamma} \in \mathbb{C}$  such that

$$I_0^{\gamma} \circ \eta = b_0^{\eta,\gamma} S_{[1],n-1} + c_0^{\eta,\gamma} S_{[w_0],n-1},$$

where  $c_0^{\eta,\gamma} = -b_0^{\eta,\gamma}$  if n is even and analogously for the  $\chi$ -twist. Finally, If  $\pi$  is the trivial representation, we consider  $\pi$  as a subrepresentation of  $\pi_{-1}$ , so this case does not need extra treatment.

**Theorem 2.5.** (a) There exists a constant  $C_{\gamma} > 0$ , depending on  $\gamma$ , such that

$$|a_k^{\eta,\gamma}| \le C_\gamma \left(1 + |\lambda|^{\frac{1}{4}}\right) \left(1 + |\lambda^2 - k^2 \mu^2|^{\frac{1}{4}}\right) e^{\frac{\pi}{4}|k\mu|}$$

holds for every cuspidal automorphic representation  $\eta: V_{\lambda} \to L^2_{\text{cusp}}(\Gamma \backslash G)$ and every  $k \in \mathbb{Z}$ .

(b) For a fixed automorphic representation  $\eta: V_{\lambda} \to L^2_{\text{cusp}}(\Gamma \backslash G)$  there exists a constant  $C_{\eta,\gamma} > 0$  such that

$$|a_k^{\eta,\gamma}| \le C_{\eta,\gamma} (1 + |k|^{\frac{1}{2}})$$

holds for every  $k \in \mathbb{Z}$ .

(c) There exists a constant  $D_{\gamma} > 0$ , depending on  $\gamma$ , such that

$$|b_0^{\eta,\gamma}|, |c_0^{\eta,\gamma}| \le D_{\gamma} n^{5-n}$$

holds for every cuspidal automorphic representation  $\eta: \mathcal{D}_{2n} \to L^2_{\text{cusp}}(\Gamma \backslash G)$ .

*Proof.* (a) Let  $\varphi = \varphi_{\lambda} \in V_{\lambda}$  be the unique K-invariant function which on K takes the value 1 and let  $f = \eta(\varphi)$ . Then

$$I_k^{\gamma}(f) = a_k^{\eta,\gamma} I_{\lambda}^{k\mu}(\varphi) = a_k^{\eta,\gamma} I_{\lambda}^{k\mu}(f_{0,0}) = a_k^{\eta,\gamma} \frac{\Gamma\left(\frac{\lambda+1+k\mu}{4}\right) \Gamma\left(\frac{\lambda+1-k\mu}{4}\right)}{\Gamma\left(\frac{\lambda+1}{2}\right)}$$

If  $\Gamma$  is cocompact, by [SS89] the sup norm of f satisfies

$$||f||_{\infty} = O\left(|\lambda|^{1/4}\right).$$

If  $\Gamma$  is not cocompact, one finds in [Iwa02], Sec. 13.2, that

$$|f(z)| = O_z(|\lambda|^{1/4}),$$

where the implied constant depends continuously on z. Therefore, this estimate holds uniformly on the closed geodesic attached to  $\gamma$ . According to [GR07] 8.328.1, for fixed real x and for  $|y| \to \infty$  one has

$$|\Gamma(x+iy)| \sim \sqrt{2\pi}e^{-\frac{\pi}{2}|y|}|y|^{x-\frac{1}{2}}.$$

This implies the claim.

(b) We start out as in the above proof, except that we don't use the estimate  $|f(z)| \ll_z |\lambda|^{\frac{1}{4}}$ . We thus get

$$|a_k^{\eta,\gamma}| \ll |I_k^{\gamma}(f)|(1+|\lambda^2-k^2\mu^2|^{\frac{1}{4}})e^{\frac{\pi}{4}|k\mu|}.$$

It suffices to show that for fixed  $\gamma$  and  $\eta$  one has

$$|I_k^{\gamma}(f)| \ll e^{-\frac{\pi}{4}|k\mu|}.$$

We will show this using the technique of analytic continuation of representations from [BR99]. Let  $X \in sl_2(\mathbb{R})$  with  $\gamma = \exp(X)$ . After conjugating  $\Gamma$ , we may assume  $X = \operatorname{diag}(A, -A)$  for some A > 0. Then  $\gamma = \operatorname{diag}(e^A, e^{-A})$  and  $\mu = \frac{2\pi i}{A}$ . We have

$$I_k^{\gamma}(f) = \int_0^1 f(\exp(tX))e^{-2\pi ikt} dt.$$

It is easy to see that the function  $t \mapsto \pi_{\lambda}(a_t)\varphi$  with  $a_t = \left[\begin{smallmatrix} e^{At} \\ e^{-At} \end{smallmatrix}\right]$  extends to a holomorphic function from  $\{|\mathrm{Im}z| < \frac{\pi}{4A}\}$  to  $V_{\lambda}$ . It follows that the function  $f(\exp(tX)) = \eta(\pi_{\lambda}(a_t)\varphi)(1)$  extends to a holomorphic function on the set of all  $z = x + iy \in \mathbb{C}$  with  $|y| < \frac{\pi}{4A}$ . We get a continuous extension to  $|y| \leq \frac{\pi}{4A}$ . For  $k \geq 0$  we get by a shift of the contour integral that

$$\begin{split} I_k^{\gamma}(f) &= \int_0^1 f(\exp(tX)) e^{-2\pi i k t} \, dt \\ &= \int_0^1 f(\exp((t - i \frac{\pi}{4A}) X) e^{-2\pi i k t} \, dt \, e^{-\frac{\pi^2}{2A} k} = \operatorname{const} \cdot e^{-\frac{\pi}{4} |k\mu|}. \end{split}$$

For k < 0 we similarly move the contour to  $i\frac{\pi}{4A}$ .

(c) Let  $n \in \mathbb{N}$  and let  $\varphi_n : K \to \mathbb{C}$  be given by

$$\varphi_n\left(\left[\begin{smallmatrix} d & -c \\ c & d \end{smallmatrix}\right]\right) = (ci+d)^{2n}.$$

Then  $\varphi_n$  and its complex conjugate span the lowest K-type in  $\mathcal{D}_{2n} \cong \pi_{1-2n}/\delta_{2n-2}$ . A computation shows

$$R_{\bar{X}}\varphi_n = in(\varphi_n + \varphi_{n+1}),$$
  

$$R_X\varphi_n = -in(\varphi_n - \varphi_{n+1}).$$

We use induction in  $k \in \mathbb{N}_0$  to show that

$$S_{[1],k}(\varphi_n) = (-1)^{k+n} S_{[w_0],k}(\varphi_n).$$

For k = 0 we have

$$S_{[1],k}(\varphi_n) = \varphi_n(1) = 1 = (-1)^n \varphi(w_0) = (-1)^{k+n} S_{[w_0],k}(\varphi_n).$$

The step  $k \mapsto k+1$  is

$$\begin{split} S_{[1],k+1}(\varphi_n) &= R_{\bar{X}}^{k+1} \varphi_n(1) \\ &= in(R_{\bar{X}}^k(\varphi_n + \varphi_{n+1})(1) \\ &= in(S_{[1],k}(\varphi_n) + S_{[1],k}(\varphi_{n+1}) \\ &= in((-1)^{k+n} S_{[w_0],k}(\varphi_n) + (-1)^{k+n+1} S_{[w_0],k}(\varphi_{n+1}) \\ &= (-1)^{k+n+1} \left( -in(S_{[w_0],k}(\varphi_n) + S_{[w_0],k}(\varphi_{n+1}) \right) \\ &= (-1)^{k+n+1} \left( -in(R_X^k \varphi_n(w_0) + R_X^k \varphi_{n+1}(w_0)) \right) \\ &= (-1)^{k+n+1} \left( R_X^{k+1} \varphi_n(w_0) \right) \\ &= (-1)^{k+n+1} \left( S_{[w_0],k+1}(\varphi_n) \right). \end{split}$$

Next we show that

$$|S_{[1],n-1}(\varphi_n)| \ge n^{n-1}$$
 and  $|S_{[1],n-1}(\varphi_{n+1})| \ge n^{n-1}$ 

This follows from  $\varphi_m(1) = 1$  and the fact that

$$R_{\bar{X}}^k \varphi_n = i^k (n^k \varphi_n + (*)),$$

where (\*) denotes a linear combination of  $\varphi_m$ ,  $m \geq n+1$  with positive coefficients. Now the equality

$$I_0^{\gamma}(\eta(\varphi_n)) = b_0^{\eta,\gamma} S_{[1],n-1}(\varphi_n) + c_0^{\eta,\gamma} S_{[w_0],n-1}(\varphi_n)$$

implies  $I_0^{\gamma}(\eta(\varphi_n))=(b_0^{\eta,\gamma}-c_0^{\eta,\gamma})S_{[1],n-1}(\varphi_n)$  and thus

$$|b_0^{\eta,\gamma} - c_0^{\eta,\gamma}| \ll \frac{|I_0^{\gamma}(\eta(\varphi_n))|}{n^{n-1}}.$$

Similarly, we have  $I_0^{\gamma}(\eta(\varphi_{n+1})) = (b_0^{\eta,\gamma} + c_0^{\eta,\gamma}) S_{[1],n-1}(\varphi_{n+1})$ , and thus

$$|b_0^{\eta,\gamma} + c_0^{\eta,\gamma}| \ll \frac{|I_0^{\gamma}(\eta(\varphi_n))|}{n^{n-1}}.$$

So the same estimate holds for  $b_0^{\eta,\gamma}$  and  $c_0^{\eta,\gamma}$  separately. We finish the proof of the lemma by estimating the period integral

$$|I_0^{\gamma}(\eta(\varphi_n))| \ll n^4,$$

independent of  $\eta$ . To see this, let  $\Delta$  denote the group Laplacian, i.e.,  $\Delta = -C + 2C_K$ , where C is the Casimir-operator and  $C_K$  is the Casimir operator of the group K. Then for  $\eta: \mathcal{D}_{2n} \to L^2_{\text{cusp}}(\Gamma \backslash G)$  a computation shows  $\Delta \eta(\varphi_n) = n(n-1)\eta(\varphi_n)$ . Since  $\Delta$  is elliptic, positive definite and of order two on the 3-dimensional manifold  $\Gamma \backslash G$ , the operator  $(1 + \Delta)^{-2}$  has a continuous kernel k(x,y). It acts on the space of cusp forms through the continuous kernel

$$k_0(x,y) = \sum_{j=1}^{\infty} (1 + \lambda_j)^{-2} \phi_j(x) \overline{\phi_j(y)},$$

where  $\phi_j$  is an orthonormal basis of  $L^2_{\text{cusp}}(\Gamma \backslash G)$  consisting of  $\Delta$ -eigenfunctions and  $\lambda_j$  is the eigenvalue of  $\phi_j$ . Selberg has shown in the Göttingen lectures, that  $k_0$  actually is an  $L^2$ -kernel. We have

$$|\eta(\varphi_n)(x)| = (1 + n(n-1))^2 |(1 + \Delta)^{-2} \eta(\varphi_n)(x)|$$

$$= (1 + n(n-1))^2 \left| \int_{\Gamma \setminus G} k_0(x, y) \eta(\varphi_n)(y) \, dy \right|$$

$$\leq (1 + n(n-1))^2 \left( \int_{\Gamma \setminus G} |k_0(x, y)|^2 \, dy \right)^{1/2}.$$

As  $k_0$  is an  $L^2$ -kernel, the latter integral is finite almost everywhere in x and is locally bounded outside a set of measure zero. Hence the continuous function  $\eta(\varphi_n)$  is locally bounded by a constant times  $(1 + n(n-1))^2$ , so it is locally  $O(n^4)$  and the same holds for the period integral  $I_0^{\gamma}(\eta(\varphi_n))$ .

# 3 Fourier expansion of Maaß forms

Let  $f: \Gamma\backslash\mathbb{H} \to \mathbb{C}$  be a Maaß form, i.e.,  $f \in L^2(\Gamma\backslash\mathbb{H})$  is an eigenform of the hyperbolic Laplacian, say  $\Delta f = (\frac{1}{4} - \lambda^2)f$  for  $\lambda \in \mathbb{C}$ . Then there exists an automorphis representation  $(\pi, V_{\pi}, \eta)$  with  $\pi = \pi_{\lambda}$  such that  $f = \eta(\varphi_0)$ , where  $\varphi_0 \in V_{\pi}$  is a K-invariant function. By scaling, one can achieve  $\varphi_0(K) = \{1\}$ . One then gets the Fourier expansion

$$f(x) = \sum_{k \in \mathbb{Z}} a_k^{\eta, \gamma} I_{\lambda, k\mu_{\gamma}}^{\text{st}}(\pi(\sigma^{-1}x)\varphi_0).$$

Note that the automorphic representation  $\eta$ , which determines the form f, enters on the right hand side only through the coefficients  $(a_k^{\eta,\gamma})_{k\in\mathbb{Z}}$ . In particular, if  $x = \sigma a_t$  for  $t \in \mathbb{R}$ , one has  $I_{\lambda,k\mu_{\gamma}}^{\mathrm{st}}(\pi(\sigma^{-1}x)\varphi_0) = I_{\lambda,k\mu_{\gamma}}^{\mathrm{st}}(\pi(a_t)\varphi_0) = I_{\lambda,k\mu_{\gamma}}^{\mathrm{st}}(\pi(a_t)\varphi_0)$ 

$$a_t^{\mu} I_{\lambda,k\mu_{\gamma}}^{\mathrm{st}}(\varphi_0)$$
, and therefore  $f\left(\sigma\left(\begin{smallmatrix}e^t\\e^{-t}\end{smallmatrix}\right)\right)$  equals 
$$\frac{1}{\Gamma\left(\frac{\lambda+1}{2}\right)} \sum_{i=\pi} a_k^{\eta,\gamma} e^{2\pi i \tilde{\mu} k t} \Gamma\left(\frac{\lambda+2\pi i k \tilde{\mu}+1}{4}\right) \Gamma\left(\frac{\lambda-2\pi i k \tilde{\mu}+1}{4}\right).$$

#### 4 Triple products

In this section we assume  $\Gamma$  to be cocompact. Then  $L^2_{\text{cusp}}(\Gamma \backslash G) = L^2(\Gamma \backslash G)$ . Let  $(\pi, V_{\pi}, \eta)$  be an automorphic representation with  $\pi = \pi_{\lambda}$  for some  $\lambda \in \mathbb{C}$ . Since  $\pi$  is unitary, there is an anti-linear isomorphism to the dual  $c: V_{\pi} \to V_{\pi}^*$ . Let  $\check{\pi}$  denote the dual representation on  $V_{\pi}^* = V_{\check{\pi}}$ . Let  $\bar{\cdot}$  be the complex conjugation on  $L^2(\Gamma \backslash G)$  and let  $\check{\eta}$  be the composition of the maps

$$V_{\breve{\pi}} \xrightarrow{c^{-1}} V_{\pi} \xrightarrow{\eta} L^2(\Gamma \backslash G) \xrightarrow{:} L^2(\Gamma \backslash G)$$

Then  $\check{\eta}$  is a G-equivariant linear isometry of  $V_{\check{\pi}}$  into  $L^2(\Gamma \backslash G)$ , so  $(\check{\pi}, V_{\check{\pi}}, \check{\eta})$  is an automorphic representation as well.

Let  $\Delta: \Gamma \backslash G \to \Gamma \backslash G \times \Gamma \backslash G$  be the diagonal map. Let  $\Delta^*: C^{\infty}(\Gamma \backslash G \times \Gamma \backslash G) \to C^{\infty}(\Gamma \backslash G)$  be the corresponding pullback map and let  $E = V_{\pi}^{\infty} \hat{\otimes} V_{\tilde{\pi}}^{\infty}$ , where  $\hat{\otimes}$  denotes the projective completion of the algebraic tensor product. Let

$$\eta_E : E \hookrightarrow C^{\infty}(\Gamma \backslash G) \hat{\otimes} C^{\infty}(\Gamma \backslash G) \cong C^{\infty}(\Gamma \backslash G \times \Gamma \backslash G)$$

be given by  $\eta \otimes \check{\eta}$ . For  $\gamma$  as in the first section and  $k \in \mathbb{Z}$  we get an induced functional on E,

$$l_{\Delta(\gamma)}^k = I_k^{\gamma} \circ \Delta^* \circ \eta_E.$$

In other words, for  $w \in E$  we have

$$l_{\Delta(\gamma)}(w) = \frac{1}{l(\gamma)} \int_{A/\langle a_{\gamma} \rangle} \eta_E(w) (\sigma_{\gamma} a, \sigma_{\gamma} a) \, da.$$

This has the Fourier series expansion,

$$\begin{split} l_{\Delta(\gamma)}(w) &= \sum_{k \in \mathbb{Z}} I_k^{\gamma} \circ \eta \otimes I_{-k}^{\gamma} \circ \check{\eta}(w) \\ &= \sum_{k \in \mathbb{Z}} a_k^{\eta, \gamma} a_{-k}^{\check{\eta}, \gamma} I_{\lambda, k \mu_{\gamma}}^{\text{st}} \otimes I_{-\lambda}^{-k \mu_{\gamma}}(w) \\ &= \sum_{k \in \mathbb{Z}} |a_k^{\eta, \gamma}|^2 \, \hat{w}(k, -k), \end{split}$$

where the last line defines  $\hat{w}$  and we have used the fact that  $a_{-k}^{\check{\eta},\gamma} = \overline{a_k^{\eta,\gamma}}$ .

Let  $(\tau, V_{\tau})$  be another element of  $\hat{G}$ . According to [Mol79] there is a canonical G-invariant continuous functional

$$T_{\tau}^{\mathrm{st}}: E \,\hat{\otimes} V_{\breve{\tau}}^{\infty} \,\to\, \mathbb{C},$$

and any other such functional is a scalar multiple of  $T_{\tau}^{\text{st}}$ . This induces a canonical G-equivariant continuous map

$$T_{\tau}^{\mathrm{st}}: E \to V_{\tau}^{\infty}.$$

On the other hand we have  $\Delta^* \circ \eta_E \colon E \to L^2(\Gamma \backslash G)^{\infty}$ . For an automorphic representation  $(\tau, V_{\tau}, \eta_{\tau})$  we have an orthogonal projection  $\Pr_{\eta_{\tau}} \colon L^2(\Gamma \backslash G) \to V_{\tau}$  and thus we get a map

$$T_{\eta_{\tau}}^{\mathrm{aut}} = \mathrm{Pr}_{\eta_{\tau}} \circ \Delta^* \circ \eta_E$$

from E to  $V_{\tau}^{\infty}$ . Hence there is a coefficient  $c(\eta, \eta_{\tau}) \in \mathbb{C}$  such that

$$T_{\eta_{\tau}}^{\text{aut}} = c(\eta, \eta_{\tau}) T_{\tau}^{\text{st}}.$$

Fix a complete family  $(\eta_j)$  of normalized, pairwise orthogonal automorphic representations  $\eta_j: \pi_j \to L^2(\Gamma \backslash G)$ . Then the spectral expansion of  $\Delta^* \circ \eta_E$  is

$$\Delta^* \circ \eta_E = \sum_j c(\eta, \eta_j) T_{\pi_j}^{\text{st}}.$$

And hence, for  $w \in E$ ,

$$\begin{split} l_{\Delta(\gamma)}(w) &= \sum_{j: \pi_j \notin \hat{G}_{\mathrm{ds}}} c(\eta, \eta_j) a_0^{\eta_j, \gamma} I_{\pi_j}^0(T_{\pi_j}^{\mathrm{st}}(w)) \\ &+ \sum_{j: \pi_j \in \hat{G}_{\mathrm{ds}} \atop \pi_j \cong \mathcal{D}_{2n}} c(\eta, \eta_j) b_0^{\eta_j, \gamma} \left( S_{[1], n-1}(T_{\pi_j}^{\mathrm{st}}(w)) - S_{[w_0], n-1}(T_{\pi_j}^{\mathrm{st}}(w)) \right), \end{split}$$

where  $\hat{G}_{ds} \subset \hat{G}$  is the set of all discrete series representations of G. So we conclude

#### Lemma 4.1.

$$\sum_{k \in \mathbb{Z}} |a_k^{\eta,\gamma}|^2 \, \hat{w}(k,-k) = \sum_{j:\pi_j \notin \hat{G}_{ds}} c(\eta,\eta_j) a_0^{\eta_j,\gamma} I_{\pi_j}^0(T_{\pi_j}^{st}(w))$$

$$+ \sum_{\substack{j:\pi_j \in \hat{G}_{ds} \\ \pi_j \cong \mathcal{D}_{2n}}} c(\eta,\eta_j) b_0^{\eta_j,\gamma} \left( S_{[1],n-1}(T_{\pi_j}^{st}(w)) - S_{[w_0],n-1}(T_{\pi_j}^{st}(w)) \right).$$

We make this a bit more explicit by plugging in special test functions. The result is the following Theorem.

**Theorem 4.2** (Summation formula). For  $\alpha \in C_c^{\infty}(\mathbb{R}^{\times})$  we have

$$\sum_{k \in \mathbb{Z}} |a_k^{\eta, \gamma}|^2 \hat{\alpha}(\tilde{\lambda} + k\tilde{\mu}_{\gamma}) = \sum_{j: \pi_j \notin \hat{G}_{\mathrm{ds}}} c(\eta, \eta_j) a_0^{\eta_j, \gamma} \int_{\mathbb{R}^2} W_j(t, x) \alpha(\hat{t}_x) dt dx,$$

where

$$W_{j}(t,x) = \left( \left| \frac{(e^{2t} + x)x}{(e^{2t} + x - 1)(x - 1)} \right|^{\frac{1}{2}} + \left| \frac{(e^{2t} + x)x}{(e^{2t} + x - 1)(x - 1)} \right|^{-\frac{1}{2}} \right)^{\frac{\lambda_{j} - 1}{2}} \times \left| \frac{(e^{2t} + x)x}{(e^{2t} + x - 1)(x - 1)} \right|^{\frac{1}{2}} e^{(\lambda_{j} + 1)t} |x|^{\lambda - 1} |e^{2t} + x|^{-\lambda - 1},$$

$$\hat{t}_{x} = \frac{1}{2} \log \left| \frac{(e^{2t} + x)(x - 1)}{(e^{2t} + x - 1)x} \right|,$$

*Proof.* Let  $H_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ . For  $x \in \mathbb{R}$  write  $n(x) = \begin{pmatrix} 1 & x \\ 1 & 1 \end{pmatrix}$ . For  $\varphi \in C_c^{\infty}(\mathbb{R})$  we set

$$f_{\varphi}(w_0 n(x)) = \varphi(-\frac{1}{2}\log|x|).$$

For given  $\lambda \in \mathfrak{a}^*$  the function  $f_{\varphi}$  extends uniquely to an element of  $V_{\lambda}^{\infty}$ . For  $a = \exp(tH_1) \in A$  we have  $a^{\lambda} = e^{\lambda t}$ . We define  $\tilde{\lambda} = \frac{\lambda}{2\pi i}$ , so that  $a^{\lambda} = e^{2\pi i \tilde{\lambda} t}$ . We normalize the Haar measure da on A such that  $\int_A g(a) da = \int_{\mathbb{R}} g(\exp(tH_1) dt)$ . Assuming that  $n_0 = \begin{pmatrix} 1 & 1 \\ 1 & \end{pmatrix}$ , we compute for  $k \in \mathbb{Z}$ ,

$$I_{\lambda,k\mu_{\gamma}}^{\mathrm{st}}(f_{\varphi}) = \int_{A} f_{\varphi}(w_{0}n_{0}^{a})a^{-\lambda - 1 - k\mu_{\gamma}}da$$
$$= \int_{\mathbb{R}} \varphi(t) e^{-t(2\pi i\tilde{\lambda} + 1 + 2\pi ik\tilde{\mu}_{\gamma})}dt.$$

Likewise, for  $\phi \in C_c^{\infty}(\mathbb{R}^2)$  let  $w_{\phi} \in E$  be defined by  $w_{\phi}(w_0 n(x), w_0 n(y)) = \phi(-\frac{1}{2}\log|x|, -\frac{1}{2}\log|y|)$ . Here we identify E with  $V_{\lambda}^{\infty} \hat{\otimes} V_{-\lambda}^{\infty}$ . We get

$$\hat{w}_{\phi}(k, -k) = \int_{\mathbb{R}^2} \phi(t, s) e^{(s-t)(2\pi i\tilde{\lambda} + 2\pi i k\tilde{\mu}_{\gamma}) - (s+t)} dt ds$$

Note that  $w_{\phi}(w_0n(x), w_0n(y))$  vanishes in a neighborhood of  $\{xy = 0\}$  as well as in a neighborhood of  $\{x = \infty\} \cup \{y = \infty\}$ , which means that  $w_{\phi}$  indeed lies in E and that

$$S_{[1],n-1]}(T_{\pi_j}^{\text{st}}(w_\phi)) = S_{[w_0],n-1}(T_{\pi_j}^{\text{st}}(w_\phi)) = 0$$

for every  $n \in \mathbb{N}$  and every  $\pi_j \cong \mathcal{D}_{2n}$ . So for these test functions ony the first sum on the right hand side of Lemma 4.1 is present.

**Lemma 4.3.** Let  $g = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  and  $g' = \begin{bmatrix} a' & b' \\ c' & d' \end{bmatrix}$  be in G. If cc'dd' = 0, then  $w_{\phi}(g, g') = 0$ . Otherwise,

$$w_{\phi}(g, g') = |c|^{-\lambda - 1} |c'|^{\lambda - 1} \phi\left(\frac{1}{2} (\log \left| \frac{c}{d} \right|, \frac{1}{2} \log \left| \frac{c'}{d'} \right|\right).$$

*Proof.* Replacing g by  $\begin{bmatrix} 1 \end{bmatrix} g$  or g' by  $\begin{bmatrix} 1 \end{bmatrix} g'$  does not change  $w_{\phi}(g,g')$ . We therefore can restrict our attention to the case  $g,g'\in G^0$ . The Ansatz  $g=anw_0n(x)$  with  $a=\begin{pmatrix} y \\ 1/y \end{pmatrix}$  and  $n=\begin{pmatrix} 1 & z \\ 1 \end{pmatrix}$  leads to y=1/c and x=d/c as well as z=ac. This gives the claim.

With  $w = w_{\phi}$  as above, we want to compute  $I_0^{\pi_j}(T_{\pi_j}^{\mathrm{st}}(w))$ . We first consider the case  $\pi_j = \pi_{\lambda_j}$  for some  $\lambda_j \in \mathbb{C}$ . Recall that the functional  $T^{\mathrm{st}} : \pi \otimes \check{\pi} \otimes \check{\pi}_j \to \mathbb{C}$  maps a given  $\varphi = w \otimes f$  to

$$T^{\text{st}}(\varphi) = \int_{G} \varphi(w_0 n_0 y, w_0 y, y) \, dy$$
$$= \int_{ANK} w(w_0 n_0 ank, w_0 ank) f(ank) \, da \, dn \, dk$$
$$= \int_{K} \int_{AN} a^{-\lambda_j + 1} w(w_0 n_0 ank, w_0 ank) da \, dn \, f(k) \, dk.$$

The induced map  $T_{\pi_j}^{\mathrm{st}}: E \to \pi_j$  is defined via the pairing

$$\breve{\pi}_j \otimes \pi_j = \pi_{-\lambda_j} \otimes \pi_{\lambda_j} \to \mathbb{C}$$

given by

$$(f \otimes h) \mapsto \int_K f(k)h(k) dk.$$

The resulting map  $T^{\mathrm{st}}_{\pi_j}:\pi\otimes \breve{\pi}\to \pi_j$  therefore is given by

$$T_{\pi_j}^{\text{st}}(w)(k) = \int_{AN} a^{-\lambda_j + 1} w(w_0 n_0 ank, w_0 ank) da dn.$$

We have

$$I_0^{\pi_j}(T_{\pi_j}^{\mathrm{st}}(w)) = \int_A \underline{a}(w_0 n_0 a_1)^{\lambda_j + 1} T_{\pi_j}^{\mathrm{st}}(w) (\underline{k}(w_0 n_0 a_1)) da_1$$

which equals

$$\int_A \int_{AN} \underline{a}(w_0 n_0 a_1)^{\lambda_j + 1} a^{-\lambda_j + 1} w(w_0 n_0 a n \underline{k}(w_0 n_0 a_1), w_0 a n \underline{k}(w_0 n_0 a_1)) \, da \, dn \, da_1.$$

We write  $\underline{k}(w_0n_0a_1) = \underline{an}(w_0n_0a_1)^{-1}w_0n_0a_1$  and use the change of variables  $an \mapsto an\underline{an}(w_0n_0a_1)$  in the AN-integral. For this we use the formula  $\int_H f(xy) dx = \Delta(y^{-1}) \int_H f(x) dx$  for Haar integration over the group H = AN and the fact that the modular function  $\Delta$  of AN equals  $\Delta(a_tn) = e^{-2t}$ . In this way we see that  $I_0^{\pi_j}(T_{\pi_j}^{\rm st}(w))$  equals

$$\int_{A} \int_{AN} \underline{a}(w_{0}n_{0}a_{1})^{\lambda_{j}-1} a^{\lambda-\lambda_{j}} w(w_{0}n_{0}anw_{0}n_{0}a_{1}, w_{0}nw_{0}n_{0}a_{1}) da dn da_{1}.$$

Writing  $\overline{n}_x = \begin{pmatrix} 1 \\ -x & 1 \end{pmatrix} = w_0 n_x w_0$  this equals

$$\int_{A} \int_{AN} \underline{a}(w_0 n_0 a_1)^{\lambda_j - 1} a^{\lambda - \lambda_j} w(\overline{n}_0 a^{-1} \overline{n} n_0 a_1, \overline{n} n_0 a_1) da dn da_1.$$

Writing 
$$a = \begin{bmatrix} e^t \\ e^{-t} \end{bmatrix}$$
,  $n = \begin{bmatrix} 1 & x \\ 1 \end{bmatrix}$ , and  $a_1 = \begin{bmatrix} e^s \\ e^{-s} \end{bmatrix}$  we get

$$\overline{n}_0 a^{-1} \overline{n} n_0 a_1 = \begin{bmatrix} -e^{-t} & -e^{-t} \\ e^s (e^t x + e^{-t}) & e^{-s} (e^t (x - 1) + e^{-t}) \end{bmatrix}$$

and

$$\overline{n}n_0a_1 = \begin{bmatrix} -e^s & -e^{-s} \\ e^s x & e^{-s}(x-1) \end{bmatrix}.$$

By Lemma 4.3 we conclude that  $w_{\phi}(\overline{n}_0 a^{-1} \overline{n} n_0 a_1, \overline{n} n_0 a_1)$  equals

$$e^{-2s}|e^tx + e^{-t}|^{-\lambda - 1}|x|^{\lambda - 1}\phi\left(s + \frac{1}{2}\log\left|\frac{e^{-2t} + x}{e^{-2t} + x - 1}\right|, s + \frac{1}{2}\log\left|\frac{x}{x - 1}\right|\right).$$

We now pick  $\alpha, \beta \in C_c^{\infty}(\mathbb{R})$  and set

$$\phi(x,y) = \alpha(x-y)\beta\left(\frac{x+y}{2}\right).$$

With this choice, we get

$$\hat{w}_{\phi}(k, -k) = \hat{\alpha}(\tilde{\lambda} + k\tilde{\mu}_{\gamma})\hat{\beta}\left(\frac{1}{\pi i}\right),$$

and  $w_{\phi}(\overline{n}_0 a^{-1} \overline{n} n_0 a_1, \overline{n} n_0 a_1)$  equals

$$e^{-2s}|e^{t}x + e^{-t}|^{-\lambda - 1}|x|^{\lambda - 1}\alpha \left(\frac{1}{2}\log\left|\frac{(e^{-2t} + x)(x - 1)}{(e^{-2t} + x - 1)x}\right|\right) \times \beta \left(s + \frac{1}{4}\log\left|\frac{(e^{-2t} + x)x}{(e^{-2t} + x - 1)(x - 1)}\right|\right).$$

We conclude that  $I_0^{\pi_j}(T_{\pi_j}^{\mathrm{st}}(w))$  equals

$$\begin{split} & \int_{\mathbb{R}^3} (e^{2s} + e^{-2s})^{\frac{\lambda_j - 1}{2}} e^{(\lambda - \lambda_j)t} e^{-2s} |e^t x + e^{-t}|^{-\lambda - 1} |x|^{\lambda - 1} \\ & \times \alpha \left( \frac{1}{2} \log \left| \frac{(e^{-2t} + x)(x - 1)}{(e^{-2t} + x - 1)x} \right| \right) \\ & \times \beta \left( s + \frac{1}{4} \log \left| \frac{(e^{-2t} + x)x}{(e^{-2t} + x - 1)(x - 1)} \right| \right) \, dx \, ds \, dt. \end{split}$$

After the change of variables  $s\mapsto s-\frac{1}{4}\log\left|\frac{(e^{-2t}+x)x}{(e^{-2t}+x-1)(x-1)}\right|$  we end up with

$$\int_{\mathbb{R}^{3}} \left( e^{2s} \left| \frac{(e^{-2t} + x)x}{(e^{-2t} + x - 1)(x - 1)} \right|^{\frac{1}{2}} + e^{-2s} \left| \frac{(e^{-2t} + x)x}{(e^{-2t} + x - 1)(x - 1)} \right|^{-\frac{1}{2}} \right)^{\frac{\lambda_{j} - 1}{2}} \\
\times \left| \frac{(e^{-2t} + x)x}{(e^{-2t} + x - 1)(x - 1)} \right|^{\frac{1}{2}} e^{(\lambda - \lambda_{j})t} |x|^{\lambda - 1} |e^{t}x + e^{-t}|^{-\lambda - 1} \\
\times \alpha \left( \frac{1}{2} \log \left| \frac{(e^{-2t} + x)(x - 1)}{(e^{-2t} + x - 1)x} \right| \right) \beta(s) \, ds \, dt \, dx.$$

Assuming  $\beta \geq 0$  and  $\int_{\mathbb{R}} \beta(s) ds = 1$  we replace  $\beta$  with  $\beta_T(x) = T\beta(Tx)$  and let  $T \to \infty$  to get the claim. The interchange of limit and sum resp. integral is justified by the Theorem of dominated convergence as follows. Using results from [BR04] one deduces that

$$c(\eta, \eta_j) = O(|\lambda_j|^{2+\varepsilon}).$$

Weyl's asymptotic law says that  $|\lambda_j| = O(j)$  and with Theorem 2.5 we get

$$a_0^{\eta_j,\gamma}c(\eta,\eta_j) = O(\lambda_j)^3 = O(j^3).$$

So we need an estimate  $O(|\lambda_j|^{-5})$  of the above integral, with  $\beta = \beta_T$ , which is uniform in T. One achieves this by iterated use of the fact that  $2\lambda_j e^{2t\lambda_j} = \frac{d}{dt}e^{2t\lambda_j}$  and using integration by parts. This, actually, is the place where it is needed that  $\alpha$  be vanishing in a neighborhood of zero, as otherwise there would appear boundary terms of this integration by parts. The Theorem follows.

### References

- [BR99] Joseph Bernstein and Andre Reznikov, Analytic continuation of representations and estimates of automorphic forms, Ann. of Math. (2) **150** (1999), no. 1, 329–352, DOI 10.2307/121105. MR1715328 (2001h:11053)
- [BR04] \_\_\_\_\_\_, Estimates of automorphic functions, Mosc. Math. J. 4 (2004), no. 1, 19–37, 310 (English, with English and Russian summaries). MR2074982 (2005f:11097)
- [Dei06] Anton Deitmar, Invariant triple products, Int. J. Math. Math. Sci., posted on 2006, Art. ID 48274, 22, DOI 10.1155/IJMMS/2006/48274. MR2251763 (2007g:22013)
- [GV88] Ramesh Gangolli and V. S. Varadarajan, Harmonic analysis of spherical functions on real reductive groups, Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas], vol. 101, Springer-Verlag, Berlin, 1988. MR954385 (89m:22015)
- [GR07] I. S. Gradshteyn and I. M. Ryzhik, Table of integrals, series, and products, 7th ed., Elsevier/Academic Press, Amsterdam, 2007. Translated from the Russian; Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger; With one CD-ROM (Windows, Macintosh and UNIX). MR2360010 (2008g:00005)
- [HC75] Harish-Chandra, Harmonic analysis on real reductive groups. I. The theory of the constant term, J. Functional Analysis 19 (1975), 104–204. MR0399356 (53 #3201)
- [IS95] H. Iwaniec and P. Sarnak,  $L^{\infty}$  norms of eigenfunctions of arithmetic surfaces, Ann. of Math. (2) **141** (1995), no. 2, 301–320, DOI 10.2307/2118522. MR1324136 (96d:11060)
- [Iwa02] Henryk Iwaniec, Spectral methods of automorphic forms, 2nd ed., Graduate Studies in Mathematics, vol. 53, American Mathematical Society, Providence, RI, 2002. MR1942691 (2003k:11085)
- [Kna01] Anthony W. Knapp, Representation theory of semisimple groups, Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 2001. An overview based on examples; Reprint of the 1986 original. MR1880691 (2002k:22011)
- [Lok01] Hung Yean Loke, Trilinear forms of  $\mathfrak{gl}_2$ , Pacific J. Math. **197** (2001), no. 1, 119–144, DOI 10.2140/pjm.2001.197.119. MR1810211 (2002b:22028)
- [Mol79] V. F. Molčanov, Tensor products of unitary representations of the threedimensional Lorentz group, Izv. Akad. Nauk SSSR Ser. Mat. 43 (1979), no. 4, 860–891, 967 (Russian). MR548507 (80i:22030)
- [Oks73] A. I. Oksak, Trilinear Lorentz invariant forms, Comm. Math. Phys. 29 (1973), 189-217. MR0340478 (49 #5231)
  - [SS89] A. Seeger and C. D. Sogge, Bounds for eigenfunctions of differential operators, Indiana Univ. Math. J. 38 (1989), no. 3, 669–682, DOI 10.1512/iumj.1989.38.38031. MR1017329 (91f:58097)

Mathematisches Institut, Auf der Morgenstelle 10, 72076 Tübingen, Germany, deitmar@uni-tuebingen.de

22.6.2021