

Actions of finitely generated groups on \mathbb{R} -trees.

Vincent Guirardel

October 4, 2018

Abstract

We prove a structure theorem for actions of finitely generated groups on \mathbb{R} -trees. We show that the action can be obtained by gluing together actions of simple types: actions on simplicial trees, actions on lines, and actions coming from measured foliations on 2-orbifolds. This extends results by Sela and Rips-Sela. However, their results are misstated, and we give a counterexample to their statements.

The proof relies on an extended version of Scott's Lemma of independent interest. This statement claims that if a group G is a direct limit of groups having suitably compatible splittings, then G splits.

Actions of groups on \mathbb{R} -trees are an important tool in geometric group theory. For instance, actions on \mathbb{R} -trees are used to compactify sets of hyperbolic structures ([MS84, Pau88]) or Culler-Vogtmann's Outer space ([CM87, BF, CL95]). Actions on \mathbb{R} -trees are also a main ingredient in Sela's approach to acylindrical accessibility [Sel97] (see [Del99] for an alternative approach) and in some studies of morphisms of a given group to a (relatively) hyperbolic group [Sel99, BS, DS]. Limit groups and limits of (relatively) hyperbolic groups [Sel01a, Sel, Gui04, Ali, Gro05], and Sela's approach to Tarski's problem ([Sel01b]) are also studied using \mathbb{R} -trees as a basic tool.

To use \mathbb{R} -trees as a tool, one needs to understand the structure of a group acting on an \mathbb{R} -tree. The main breakthrough in this analysis is due to Rips. He proved that any finitely generated group acting freely on an \mathbb{R} -tree is a free product of surface groups and of free abelian groups (see [GLP94, BF95]).

More general results involve *stability* hypotheses. Roughly speaking, these hypotheses prohibit infinite sequences of nested arc stabilizers. A simple version of these stability hypotheses is the *ascending chain condition*: for any decreasing sequence of arcs $I_1 \supset I_2 \supset \dots$ whose lengths converge to 0, the sequence of their pointwise arc stabilizers $G(I_1) \subset G(I_2) \subset \dots$ stabilizes. The ascending chain condition implies BF-stability used below. See Definition 1.1 for other versions of stability and relations between them.

Another theorem by Rips claims that if G is finitely presented and has a BF-stable action on an \mathbb{R} -tree T , then either T has an invariant line or G splits over an extension of a subgroup fixing an arc by a cyclic group (see [BF95]). Using methods in the spirit of [GLP94], the author proved under the same hypotheses that one can *approximate* the action $G \curvearrowright T$ by a sequence of actions on simplicial trees $G \curvearrowright T_k$ converging to T ([Gui98b]). Edge stabilizers of T_k are extensions by finitely generated free abelian groups of subgroups fixing an arc in T . The convergence is in the length functions topology.

Next results give a very detailed structure of the action on the \mathbb{R} -tree. This is the basis for Sela's shortening argument (see for instance [RS94, Sel97]). These results say that the action on the \mathbb{R} -tree can be obtained by gluing together simple building blocks. The building blocks are actions on simplicial trees, actions on a line, and actions coming from a measured foliation on a 2-orbifold. Two building blocks are glued together along one point, and globally, the combinatorics of the gluing is described by a simplicial tree

(see section 1.3 for more details). When $G \curvearrowright T$ arises in this fashion, we say that T splits as a graph of actions on \mathbb{R} -trees.

In [RS94] and [Sel97], Rips-Sela and Sela give a structure theorem for actions of finitely presented groups and of finitely generated groups on \mathbb{R} -trees. The proof uses the following super-stability hypothesis: an arc $I \subset T$ is *unstable* if there exists $J \subset I$ with $G(I) \subsetneq G(J)$; an action $G \curvearrowright T$ is *super-stable* if the stabilizer of any unstable arc is trivial. In particular, super-stability implies that chains of arc stabilizers have length at most 2. Sela's result is as follows: consider a minimal action $G \curvearrowright T$ of a finitely generated group on an \mathbb{R} -tree, and assume that this action is super-stable, and has trivial tripod stabilizers. Then either G splits as a free product, or T can be obtained from simple building blocks as above ([Sel97, Theorem 3.1]).

Actually, Sela's result is stated under the ascending chain condition instead of super-stability. However, the proof really uses the stronger hypothesis of super-stability. In section 6, we give a counterexample to the more general statement. This does not affect the rest of Sela's papers since super-stability is satisfied in the cases considered. Since the counterexample is an action of a finitely presented group, this also shows that one should include super-stability in the hypotheses of Rips-Sela's statement [RS94, Theorem 10.8] (see also [Sel97, Theorem 2.3]).

Our main result generalizes Sela's result in two ways. First, we don't assume that tripod stabilizers are trivial. Moreover, we replace super-stability by the ascending chain condition together with weaker assumptions on stabilizers of unstable arcs.

Theorem (Main Theorem). *Consider a non-trivial minimal action of a finitely generated group G on an \mathbb{R} -tree T by isometries. Assume that*

1. T satisfies the ascending chain condition;
2. for any unstable arc J ,
 - (a) $G(J)$ is finitely generated;
 - (b) $G(J)$ is not a proper subgroup of any conjugate of itself i. e. $\forall g \in G, G(J)^g \subset G(J) \Rightarrow G(J)^g = G(J)$.

Then either G splits over the stabilizer of an unstable arc or over the stabilizer of an infinite tripod, or T has a decomposition into a graph of actions where each vertex action is either

1. *simplicial: $G_v \curvearrowright Y_v$ is a simplicial action on a simplicial tree;*
2. *of Seifert type: the vertex action $G_v \curvearrowright Y_v$ has kernel N_v , and the faithful action $G_v/N_v \curvearrowright Y_v$ is dual to an arational measured foliation on a closed 2-orbifold with boundary;*
3. *axial: Y_v is a line, and the image of G_v in $\text{Isom}(Y_v)$ is a finitely generated group acting with dense orbits on Y_v .*

If one assumes triviality of stabilizers of tripods and of unstable arcs (super-stability) in Main Theorem, one gets Sela's result.

The non-simplicial building blocks are canonical. Indeed, non-simplicial building blocks are *indecomposable* which implies that they cannot be split further into a graph of action (see Definition 1.17 and Lemma 1.18). If T is not a line, simplicial building blocks can also be made canonical by imposing that each simplicial building block is an arc which intersects the set of branch points of T exactly at its endpoints.

For simplicity, we did not state the optimal statement of Main Theorem, see Theorem 5.1 for more details. In particular, one can say a little more about the tripod stabilizer on which G splits. This statement also includes a relative version: G is only assumed to be finitely generated relative to a finite set of elliptic subgroups $\mathcal{H} = \{H_1, \dots, H_p\}$, and each

H_i is conjugate into a vertex group in the splittings of G produced.

When T has a decomposition into a graph of actions as in the conclusion of Main Theorem, then G splits over an extension of an arc stabilizer by a cyclic group except maybe if T is a line. Thus we get:

Corollary 5.2. *Under the hypotheses of Main Theorem, either T is a line, or G splits over a subgroup H which is an extension of a cyclic group by an arc stabilizer.*

Remark. In the conclusion of the corollary, H is an extension by a *full* arc stabilizer, and not of a subgroup fixing an arc. This contrasts with [BF95, Theorem 9.5].

Here is a simple setting where the main theorem applies. A group H is *small* (resp. *slender*) if it contains no non-abelian free group (resp. if all its subgroups are finitely generated). The following corollary applies to the case where G is hyperbolic relative to slender groups.

Corollary 5.3. *Let G be a finitely generated group for which any small subgroup is finitely generated. Assume that G acts on an \mathbb{R} -tree T with small arc stabilizers.*

Then either G splits over the stabilizer of an unstable arc or over a tripod stabilizer, or T has a decomposition into a graph of actions as in Main Theorem. In particular, G splits over a small subgroup.

In some situations, one can control arc stabilizers in terms of tripod stabilizers. For instance, we get:

Corollary 5.4. *Consider a finitely generated group G acting by isometries on an \mathbb{R} -tree T . Assume that*

1. *arc stabilizers have a nilpotent subgroup of bounded index (maybe not finitely generated);*
2. *tripod stabilizers are finitely generated (and virtually nilpotent);*
3. *no group fixing a tripod is a proper subgroup of any conjugate of itself;*
4. *any chain $H_1 \subset H_2 \dots$ of subgroups fixing tripods stabilizes.*

Then either G splits over a subgroup having a finite index subgroup fixing a tripod, or T has a decomposition as in the conclusion of Main Theorem.

Our proof relies on a particular case of Sela's Theorem, assuming triviality of arc stabilizers. For completeness, we give a proof of this result in appendix A. To reduce to this theorem, we prove that under the hypotheses of Main Theorem, the action is piecewise stable. Piecewise stability implies that T splits into a graph of actions where each vertex action $G_v \curvearrowright T_v$ has trivial arc stabilizers up to some kernel (Theorem 4.1). Because G_v itself need not be finitely generated, we need to extend Sela's result to finitely generated pairs (Proposition 4.4).

So the main step in the proof consists in proving piecewise stability (Theorem 3.1). All studies of actions on \mathbb{R} -trees use resolutions by foliated 2-complexes. Such foliated 2-complexes have a dynamical decomposition into two kinds of pieces: *simplicial* pieces where each leaf is finite, and *minimal components* where every leaf is dense. These resolutions give sequence of actions $G_k \curvearrowright T_k$ converging strongly to T , where G is the inductive limit of G_k .

Maybe surprisingly, the main difficulty in the proof arises from the simplicial pieces. This is because minimal components give large stable subtrees (Lemma 1.20). In particular, a crucial argument is a result saying that if all the groups G_k split in some nice compatible way, then so does G . In the case of free splittings, this is Scott's Lemma. In our setting, we need an extended version of Scott's Lemma, which is of independent interest.

Theorem 2.1 (Extended Scott’s Lemma). *Let $G_k \curvearrowright S_k$ be a sequence of minimal actions of finitely generated groups on simplicial trees, and $(\varphi_k, f_k) : G_k \curvearrowright S_k \rightarrow G_{k+1} \curvearrowright S_{k+1}$ be epimorphisms. Assume that f does not increase edge stabilizers in the following sense:*

$$\forall e \in E(S_k), \forall e' \in E(S_{k+1}), \quad e' \subset f_k(e) \Rightarrow G_{k+1}(e') = \varphi_k(G_k(e))$$

Then the inductive limit $G = \lim_{\rightarrow} G_k$ has a non-trivial splitting over the image of an edge stabilizer of some S_k .

The paper is organized as follows. Section 1 is devoted to preliminaries, more or less well known, except for the notion of indecomposability which seems to be new. Section 2 deals with Extended Scott’s Lemma. Section 3 proves piecewise stability. Section 4 shows how piecewise stability and Sela’s Theorem for actions with trivial arc stabilizers allow to conclude. Section 5 gives the proof of the corollaries. Section 6 contains our counterexample to Sela’s misstated result. Appendix A gives a proof of the version of Sela’s result we need.

1 Preliminaries

1.1 Basic vocabulary

An \mathbb{R} -tree is a 0-hyperbolic geodesic space. In all this section, we fix an isometric action of a group G on an \mathbb{R} -tree T . A subgroup $H \subset G$ is *elliptic* if it fixes a point in T . The action $G \curvearrowright T$ is *trivial* if G is elliptic. The action $G \curvearrowright T$ is *minimal* if T has no proper G -invariant subtree. We consider the trivial action of G on a point as minimal. When G contains a hyperbolic element, then T contains a unique minimal G -invariant subtree T_{\min} , and T_{\min} is the union of axes of all hyperbolic elements.

An *arc* is a set homeomorphic to $[0, 1]$. A subtree is *non-degenerate* if it contains an arc.

Say that an action on an \mathbb{R} -tree $G \curvearrowright T$ is *simplicial* if it can be obtained from an action on a combinatorial tree by assigning equivariantly a positive length for each edge. If S is a simplicial tree, we denote by $V(S)$ its set of vertices, and $E(S)$ its set of oriented edges.

A *morphism of \mathbb{R} -trees* $f : T \rightarrow T'$ is a 1-Lipschitz map such that any arc of T can be subdivided into a finite number of sub-arcs on which f is isometric.

For basic facts about \mathbb{R} -trees, see [Sha91, Chi01].

1.2 Stabilities

Stability hypotheses say how arc stabilizers are nested. By stabilizer, we always mean *pointwise* stabilizer. When we will talk about the global stabilizer, we will mention it explicitly. If G acts on T , and $X \subset T$, we will denote by $G(X)$ the (pointwise) stabilizer of X .

Definition 1.1 (Stabilities). *Consider an action of a group G on an \mathbb{R} -tree T .*

- *A non-degenerate subtree $Y \subset T$ is called stable if for every arc $I \subset Y$, $G(I) = G(Y)$.*
- *T is BF-stable (in the sense of Bestvina-Feighn [BF95]) if every arc of T contains a stable arc.*
- *T satisfies the ascending chain condition if for any sequence of arcs $I_1 \supset I_2 \supset \dots$ whose lengths converge to 0, the sequence of stabilizers $G(I_1) \subset G(I_2) \subset \dots$ is eventually constant.*
- *T is piecewise stable if any arc of T can be covered by finitely many stable arcs. Equivalently, the action is piecewise stable if for all $a \neq b$, the arc $[a, b]$ contains a stable arc of the form $[a, c]$ (one common endpoint).*

- T is super-stable if any arc with non-trivial stabilizer is stable.

Remark. Clearly, super-stability or piecewise stability imply the ascending chain condition which implies BF-stability. If T is a stable subtree of itself, then any arc stabilizer fixes T ; in other words, if N is the kernel of the action $G \curvearrowright T$ (i. e. N is the set of elements acting as the identity), G/N acts with trivial arc stabilizers on T .

1.3 Graphs of actions on \mathbb{R} -trees and transverse coverings

Graphs of actions on \mathbb{R} -trees are a way of gluing equivariantly \mathbb{R} -trees. They were introduced by Levitt in [Lev94]. Here, we rather follow [Gui04, section 4]. One main difference is that we assign length zero to every edge of the skeleton.

Definition 1.2. A graph of actions on \mathbb{R} -trees $\mathcal{G} = (S, (Y_v)_{v \in V(S)}, (p_e)_{e \in E(S)})$ consists of the following data:

- a simplicial tree S called the skeleton with a simplicial action without inversion of a group G ;
- for each vertex $v \in S$, an \mathbb{R} -tree Y_v (called vertex tree or vertex action);
- for each oriented edge e of S with terminal vertex v , an attaching point $p_e \in Y_v$.

All this data should be invariant under G : G acts on the disjoint union of the vertex trees so that the projection $Y_v \mapsto v$ is equivariant; and for every $g \in G$, $p_{g \cdot e} = g \cdot p_e$.

Remark. Some vertex trees may be reduced to a point (but they are not allowed to be empty). The definition implies that Y_v is $G(v)$ -invariant and that p_e is $G(e)$ -invariant.

To a graph of actions \mathcal{G} corresponds an \mathbb{R} -tree $T_{\mathcal{G}}$ with a natural action of G . Informally, $T_{\mathcal{G}}$ is obtained from the disjoint union of the vertex trees by identifying the two attaching points of each edge of S (see [Gui04] for a formal definition).

Alternatively, one can define a graph of actions as a graph of groups Γ with an isomorphism $G \simeq \pi_1(\Gamma)$ together with an action of each vertex group G_v on an \mathbb{R} -tree Y_v and for each oriented edge e with terminus vertex v , a point of Y_v fixed by the image of G_e in G_v . This is where the terminology comes from.

Definition 1.3. Say that T splits as a graph of actions \mathcal{G} if there is an equivariant isometry between T and $T_{\mathcal{G}}$.

Transverse coverings are very convenient when working with graphs of actions.

Definition 1.4. A transverse covering of an \mathbb{R} -tree T is a covering of T by a family of subtrees $\mathcal{Y} = (Y_v)_{v \in V}$ such that

- every Y_v is a closed subtree of T
- every arc of T is covered by finitely many subtrees of \mathcal{Y}
- for $v_1 \neq v_2 \in V$, $Y_{v_1} \cap Y_{v_2}$ contains at most one point

When T has an action of a group G , we always require the family \mathcal{Y} to be G -invariant.

Remark. In the definition above, if some subtrees Y_v are reduced to a point, we may as well forget them in \mathcal{Y} . Moreover, given a covering by subtrees Y_v which are not closed, but which satisfy the two other conditions for a transverse covering, then the family $(\overline{Y_v})_{v \in V}$ of their closure is a transverse covering.

The relation between graphs of actions and transverse coverings is contained in the following result.

Lemma 1.5 ([Gui04, Lemma 4.7]). Assume that T splits as a graph of actions with vertex trees $(Y_v)_{v \in V(S)}$. Then the family $\mathcal{Y} = (Y_v)_{v \in V(S)}$ is a transverse covering of T .

Conversely, if T has a transverse covering by a family $\mathcal{Y} = (Y_v)_{v \in V}$ of non-degenerate trees, then T splits as a graph of actions whose non-degenerate vertex trees are the Y_v .

We recall for future use the definition of the graph of actions induced by a transverse covering \mathcal{Y} . We first define its skeleton S . Its vertex set $V(S)$ is $V_0(S) \cup V_1(S)$ where $V_1(S)$ is the set of non-degenerate subtrees $Y \in \mathcal{Y}$, and $V_0(S)$ is the set of points $x \in T$ lying in the intersection of two distinct non-degenerate trees of \mathcal{Y} . There is an edge $e = (x, Y)$ between $x \in V_0(S)$ and $Y \in V_1(S)$ if $x \in Y$. The vertex tree of $v \in V(S)$ is the corresponding subtree of T (reduced to a point if and only if $v \in V_0(S)$). The two attaching points the edge (x, Y) are the copies of x in $\{x\}$ and Y respectively.

Example. Let $G \curvearrowright T$ be a simplicial action. Then the family of its edges is a transverse covering of T . If T has no terminal vertex, then the skeleton of this transverse covering is the barycentric subdivision of T .

Here is another general example which will be useful.

Lemma 1.6. *Assume that $G \curvearrowright$ contains a hyperbolic element and denote by T_{\min} be the minimal subtree of T . Let \mathcal{Y}_0 be the set of closures of connected components of $T \setminus \overline{T_{\min}}$. Then $\{\overline{T_{\min}}\} \cup \mathcal{Y}_0$ is a transverse covering of T .*

Remark. If $Y_v \in \mathcal{Y}_0$, then its global stabilizer G_v fixes the point $Y_v \cap \overline{T_{\min}}$. In particular, this lemma says that a finitely supported action can be decomposed into a graph of actions where each vertex action is either trivial or has a dense minimal subtree.

Proof. Any arc $I \subset T$ is covered by $\overline{T_{\min}}$ and at most two elements of \mathcal{Y}_0 , containing the endpoints of I . The other properties of the transverse covering are clear. \square

Consider a graph of actions $\mathcal{G} = (S, (Y_v)_{v \in V(S)}, (p_e)_{e \in E(S)})$, and $T_{\mathcal{G}}$ the corresponding \mathbb{R} -tree. Given a G -invariant subset $V' \subset V(S)$, we want to define an \mathbb{R} -tree by collapsing all the trees $(Y_v)_{v \in V'}$. For each $v \in V'$, replace Y_v by a point, and for each edge e incident on $v \in V'$, change p_e accordingly. Let $\mathcal{G}' = (S, (Y'_v)_{v \in V(S)}, (p'_e)_{e \in E(S)})$ the corresponding graph of actions, and $T'_{\mathcal{G}}$ the corresponding \mathbb{R} -tree.

Definition 1.7. *We say that the tree $T_{\mathcal{G}'}$ is obtained from $T_{\mathcal{G}}$ by collapsing the trees $(Y_v)_{v \in V'}$.*

Let $p_v : Y_v \rightarrow Y'_v$ the natural map (either the identity or the constant map), and $p : T_{\mathcal{G}} \rightarrow T'_{\mathcal{G}}$ the induced map.

Definition 1.8. *We say that a map $f : T \rightarrow T'$ preserves alignment the three following equivalent conditions hold:*

- for all $x \in [y, z]$ $f(x) \in [f(y), f(z)]$.
- the preimage of a convex set is convex,
- the preimage of a point is convex.

See for instance [Gui05, Lemma 1.1] for the equivalence.

Lemma 1.9. *The map $p : T_{\mathcal{G}} \rightarrow T'_{\mathcal{G}}$ preserves alignment. In particular, if $T_{\mathcal{G}}$ is minimal, so is $T'_{\mathcal{G}}$.*

Proof. Consider a point $x \in T_{\mathcal{G}'}$. Let $E'_v = \{x\} \cap Y'_v$. If u, v are such that $E'_u, E'_v \neq \emptyset$ for $i = 1, 2$, there is a path $u = v_0, \dots, v_n = v$ such that the attaching points of the edge $v_i v_{i+1}$ coincide with the point in E'_{v_i} and $E'_{v_{i+1}}$ (in particular $E'_{v_i} \neq \emptyset$).

Consider $E_v = p_v^{-1}(E'_v)$. For each v , either $E_v = Y_v$ or E_v consists of at most one point. The existence of the path above shows that the image of E_v in $T_{\mathcal{G}}$ is connected. This proves that $p^{-1}(\{x\})$ is convex so p preserves alignment.

Let $Y' \subset T'_{\mathcal{G}}$ be a non-empty G -invariant subtree. Then $p^{-1}(Y')$ is a non-empty G -invariant subtree, so $p^{-1}(Y') = T_{\mathcal{G}}$. Since p is onto, $Y' = T_{\mathcal{G}'}$, and minimality follows. \square

1.4 Actions of pairs, finitely generated pairs

By a pair of groups (G, \mathcal{H}) , we mean a group G together with a finite family of subgroups $\mathcal{H} = \{H_1, \dots, H_p\}$. We also say that the groups H_i are peripheral subgroups of G . In fact, each peripheral subgroup is usually only defined up to conjugacy so it would be more correct to define \mathcal{H} as a finite set of conjugacy class of subgroups.

An action of a pair (G, \mathcal{H}) on a tree is an action of G in which each subgroup H_i is elliptic. When the tree is simplicial, we also say that this is a splitting of (G, \mathcal{H}) or a splitting of G relative to \mathcal{H} .

Definition 1.10. *A pair (G, \mathcal{H}) is finitely generated if there exists a finite set $F \subset G$ such that $F \cup H_1 \cup \dots \cup H_p$ generates G . We also say in this case that G is finitely generated relative to \mathcal{H} .*

We say that F is a generating set of the pair (G, \mathcal{H}) .

By Serre's Lemma, if $(G, \mathcal{H}) \curvearrowright T$ is a non-trivial action of a finitely generated pair, then G contains a hyperbolic element.

Remark. If $(G, \{H_1, \dots, H_p\})$ is finitely generated, then for any $g_i \in G$, so is $(G, \{H_1^{g_1}, \dots, H_p^{g_p}\})$: just add g_i to F . Of course, if G is finitely generated, then (G, \mathcal{H}) is finitely generated for any \mathcal{H} .

If G_v is a vertex group in a finite graph of groups Γ , then it has a natural structure of pair (G_v, \mathcal{H}_v) consisting of the images in G_v of the incident edge groups. We call this pair the *peripheral structure* of G_v in Γ .

Lemma 1.11. *Consider a graph of groups Γ and $G = \pi_1(\Gamma)$.*

If Γ is finite and G is finitely generated, then the peripheral structure (G_v, \mathcal{H}_v) of each vertex group G_v is finitely generated.

Proof. We use the notations of [Ser77]. Choose a maximal subtree τ of Γ . For each vertex v , the vertex group G_v is now identified with a subgroup of $\pi_1(\Gamma, \tau)$. Let F be a finite generating set of $\pi_1(\Gamma, \tau)$. Since we can write each $g \in F$ as a product of edges of Γ and of elements of the vertex groups, there is a generating set of $\pi_1(\Gamma, \tau)$ consisting of the edges of Γ and of a finite set F' of elements of the vertex groups of Γ .

For each vertex $v \in \Gamma$, let $H_v \subset G_v$ be the subgroup generated by the elements of F' which lie in G_v , and by the image in G_v of incident edge groups. Of course, $\pi_1(\Gamma, \tau)$ is generated by the groups H_v and the edges of Γ .

Fix $v \in \Gamma$. We shall prove that $G_v = H_v$. Take $g \in G_v$, and write g as $g = g_0 e_1 g_1 e_2 \dots e_k g_k$ where $v_0 e_1 v_1 \dots e_k v_k$ is a loop based at v , and each g_i is in H_{v_i} . If the word has length 0, there is nothing to prove. If not, we shall find a shorter word of this form representing g , and the lemma will be proved by induction. If this word has not length 0, it is not reduced as a word in the graph of groups. Therefore, there are two consecutive edges e_i, e_{i+1} such that $e_{i+1} = \bar{e}_i$, and $g_i \in j_{e_i}(G_{e_i})$ (where j_e denotes the edge morphisms of Γ). Thus, we can write $h = g_0 e_1 \dots e_{i-1} g_{i-1} g'_i g_{i+1} e_{i+2} \dots e_k g_k$ where $g'_i \in j_{e_{i+1}}(G_{e_{i+1}})$. Note that $g_{i-1} g'_i g_{i+1} \in H_{v_{i-1}} = H_{v_{i+1}}$ hence we found a shorter word of the required form representing g . \square

Lemma 1.12. *Let (G, \mathcal{H}) be a finitely generated pair, and consider a (relative) splitting of (G, \mathcal{H}) . Let v be a vertex of the corresponding graph of groups.*

Then G_v is finitely generated relative to a family \mathcal{H}_v consisting of the images in G_v of the incident edge groups together with at most one conjugate of each H_i .

Proof. Let $G \simeq \pi_1(\Gamma)$ be a splitting of G as a graph of groups. We can assume that $H_i \subset G_{v_i}$ for some vertex v_i of Γ . For each index i , consider a finitely generated group \hat{H}_i containing H_i . Consider the graph of groups $\hat{\Gamma}$ obtained from Γ by adding for each i a

new edge e_i edge carrying H_i incident on a new vertex \hat{v}_i carrying \hat{H}_i , and by gluing the other side of e_i on v_i . Thus, e_i carries the amalgam $G_{v_i} *_{H_i} \hat{H}_i$. The fundamental group of $\hat{\Gamma}$ is finitely generated. By Lemma 1.11, the peripheral structure of G_v in \hat{G}_v is a finitely generated pair. The lemma follows. \square

1.5 Finitely supported actions

A *finite tree* in an \mathbb{R} -tree is the convex hull of a finite set.

Definition 1.13. *A set $K \subset T$ spans T if any arc $I \subset T$ is covered by finitely many translates of K . The tree T is finitely supported if it is spanned by a finite tree K .*

If T is a simplicial tree, then T is finitely supported if and only if T/G is a finite graph. Any minimal action of a finitely generated group is finitely supported. More generally we have:

Lemma 1.14. *A minimal action of a finitely generated pair on an \mathbb{R} -tree is finitely supported.*

Proof. Consider $(G, \{H_1, \dots, H_p\})$ a finitely generated pair acting minimally on T . Consider a finite generating set $\{f_1, \dots, f_q\}$ of the pair $(G, \{H_1, \dots, H_p\})$. Let $x_i \in T$ be a point fixed by H_i and $x \in T$ be any point. Let K be the convex hull of $\{x, f_1.x, \dots, f_q.x, x_1, \dots, x_p\}$. Then $G.K$ is connected because for each generator $g \in \{f_1, \dots, f_q\} \cup H_1 \cup \dots \cup H_p$, $g.K \cap K \neq \emptyset$. By minimality, $G.K = T$ and T is finitely supported. \square

Lemma 1.15. *Consider an action of a finitely generated pair $(G, \mathcal{H}) \curvearrowright T$, \mathcal{Y} a transverse covering of T , and S the skeleton of \mathcal{Y} .*

Then any $H \in \mathcal{H}$ is elliptic in S . Moreover, if $G \curvearrowright T$ is minimal (resp. finitely supported) then so is $G \curvearrowright S$.

Proof. The statement about minimality is proved in [Gui04, Lemma 4.9].

We use the description of the skeleton described after Lemma 1.5. We can assume that every subtree $Y \in \mathcal{Y}$ is non-degenerate. Let K be a finite tree spanning T . It is covered by finitely many trees of \mathcal{Y} . It follows that $\mathcal{Y}/G = V_1(S)/G$ is finite. Choose a lift of each element of $V_1(S)/G$ in $V_1(S)$, and consider the convex hull $L \subset S$ of those vertices. Let $x_i \in T$ be a point fixed by H_i . Each $H_i \in \mathcal{H}$ fixes a vertex $v_i \in S$, lying in $V_0(\mathcal{Y})$ or $V_1(\mathcal{Y})$ according to whether x_i lies in two distinct elements of \mathcal{Y} or not.

Let $\{f_1, \dots, f_q\}$ be a finite generating set of (G, \mathcal{H}) . Let L_0 be the convex hull of $L \cup f_1.L \cup \dots \cup f_q.L \cup \{v_1, \dots, v_p\}$. The set $S_0 = G.L_0$ is a subtree of S containing $V_1(S)$. It is finitely supported. If $S_0 \neq S$, consider $x \in V_0(S) \setminus S_0$. Recall that the vertex $x \in V_0(S)$ corresponds to a point $x \in T$. Any edge of S incident on x corresponds to a subtree $Y \in V_1(S)$ such that $x \in Y$. Since S_0 is a subtree and contains $V_1(S)$, there is exactly one such edge. It follows that x belongs to exactly one $Y \in \mathcal{Y}$, contradicting the definition of $V_0(S)$. Therefore, $S = S_0$ is finitely supported. \square

If T splits as a graph of actions, vertex actions may fail to be minimal, even if T is minimal. However, the property of being finitely supported is inherited by vertex actions:

Lemma 1.16. *Assume that T splits as a graph of actions and that T is finitely supported. Then each vertex action is finitely supported.*

Proof. Let K be a finite tree spanning of T . Let \mathcal{Y} be a transverse covering of T , and $Y_0 \in \mathcal{Y}$ be a vertex tree. Consider a finite subset F of \mathcal{Y} which covers K , and let $\{g_1.Y_0, \dots, g_p.Y_0\}$ be the set of elements of F lying in the orbit of Y_0 . Consider the convex hull K_0 of the finite trees $Y_0 \cap g_i^{-1}K$ for $i \in \{1, \dots, p\}$. It is easy to check that the finite tree K_0 spans Y_0 . \square

The conjunction of indecomposability and stability has a nice consequence:

Lemma 1.20. *If T is BF-stable, and if $Y \subset T$ is indecomposable, then Y is a stable subtree.*

Proof. First, one easily checks that if K_1, K_2 are two stable subtrees of T such that $K_1 \cap K_2$ is non-degenerate, then $K_1 \cup K_2$ is a stable subtree. Consider a stable arc $I \subset Y$, and any other arc $J \subset Y$. Since J is contained in a union $g_1.I \cup \dots \cup g_n.I$ with $g_i.I \cap g_{i+1}.I$ non-degenerate, J is stable. Since this holds for any arc J , Y is a stable subtree of T . \square


1.7 Geometric actions and strong convergence

We review some material from [LP97] where more details can be found.

Strong convergence

Given two actions $G \curvearrowright T, G' \curvearrowright T'$, we write $(\varphi, f) : G \curvearrowright T \rightarrow G' \curvearrowright T'$ when $\varphi : G \rightarrow G'$ is a morphism, and $f : T \rightarrow T'$ is a φ -equivariant map. We say that (φ, f) is onto if both f and φ are. We say that (φ, f) is morphism of \mathbb{R} -trees if f is.

A *direct system* of actions on \mathbb{R} -trees is a sequence of actions of finitely generated groups $G_k \curvearrowright T_k$ and an action $G \curvearrowright T$, with surjective morphisms of \mathbb{R} -trees $(\varphi_k, f_k) : G_k \curvearrowright T_k \rightarrow G_{k+1} \curvearrowright T_{k+1}$ and $(\Phi_k, F_k) : G_k \curvearrowright T_k \rightarrow G \curvearrowright T$ such that the following diagram commutes:



For convenience, we will use the notation $f_{kk'} = f_{k'-1} \circ \dots \circ f_k : T_k \rightarrow T_{k'}$ and $\varphi_{kk'} = \varphi_{k'-1} \circ \dots \circ \varphi_k : G_k \rightarrow G_{k'}$.

Definition 1.21 (Strong convergence). *A direct system of finitely supported actions of groups on \mathbb{R} -trees $G_k \curvearrowright T_k$ converges strongly to $G \curvearrowright T$ if*

- G is the direct limit of the groups G_k
- for every finite tree $K \subset T_k$, there exists $k' \geq k$ such that $F_{k'}$ restricts to an isometry on $f_{kk'}(K)$,

In practice, strong convergence allows to lift the action of a finite number of elements on a finite subtree of T to T_k for k large enough.

As an application of the definition, we prove the following useful lemma.

Lemma 1.22. *Assume that $G_k \curvearrowright T_k$ converges strongly to $G \curvearrowright T$. Assume that $G \curvearrowright T$ is minimal.*

Then for k large enough, $G_k \curvearrowright T_k$ is minimal.

Proof. By definition of strong convergence, T_0 is finitely supported. Let $K_0 \subset T_0$ be a finite tree spanning T_0 . Let K (resp. K_k) be the image of K_0 in T (resp. in T_k). Choose some hyperbolic elements $g_1, \dots, g_p \in G$ whose axes cover K . Let K' be the convex hull of $K \cup g_1.K \cup \dots \cup g_p.K$. Choose $g_1^0, \dots, g_p^0 \in G_0$ some preimages of g_1, \dots, g_p and let g_i^k be the image of g_i^0 in G_k . Let $K'_0 \subset T_0$ be the convex hull of $K_0 \cup g_1^0.K_0 \cup \dots \cup g_p^0.K_0$. Let K'_k be the image of K'_0 in T_k . Take k large enough so that F_k induces an isometry between K'_k and K' . Then the axes of g_1^k, \dots, g_p^k cover K_k . Since K_k spans T_k , T_k is a union of axes, so T_k is minimal. \square

Geometric actions

A *measured foliation* \mathcal{F} on a 2-complex X consists of the choice, for each closed simplex σ of X of a non-constant affine map $f_\sigma : \sigma \rightarrow \mathbb{R}$ defined up to post-composition by an isometry of \mathbb{R} , in a way which is consistent under restriction to a face: if τ is a face of σ , then $f_\tau = \varphi \circ (f_\sigma)|_\tau$ for some isometry φ of \mathbb{R} . Level sets of f_σ define a foliation on each closed simplex. Leaves of the foliations on X are defined as the equivalence classes of the equivalence relation generated by the relation x, y belong to a same closed simplex σ and $f_\sigma(x) = f_\sigma(y)$.

This also defines a transverse measure as follows: the transverse measure $\mu(\gamma)$ of a path $\gamma : [0, 1] \rightarrow \sigma$ to the foliation is the length of the path $f_\sigma \circ \gamma$. The transverse measure of a path which is a finite concatenation of paths contained in simplices is simply the sum of the transverse measures of the pieces. The transverse measure is invariant under the holonomy along the leaves. We also view this transverse measure as a metric on each transverse edge.

For simplicity, we will say *foliated 2-complex* to mean a 2-complex endowed with a measured foliation.

The pseudo-metric

$$\delta(x, y) = \inf\{\mu(\gamma) \text{ for } \gamma \text{ joining } x \text{ to } y\}$$

is zero on each leaf of X . By definition, the *leaf space made Hausdorff* X/\mathcal{F} of X is the metric space obtained from X by making δ Hausdorff, i. e. by identifying points at pseudo-distance 0. In nice situations, $\delta(x, y) = 0$ if and only if x and y are on the same leaf. In this case, X/\mathcal{F} coincides with the space of leaves of the foliation, and we say that the *leaf space is Hausdorff*.

Theorem 1.23 ([LP97, Proposition 1.7]). *Let (X, \mathcal{F}) be a foliated 2-complex. Assume that $\pi_1(X)$ is generated by free homotopy classes of curves contained in leaves.*

Then X/\mathcal{F} is an \mathbb{R} -tree. □

Definition 1.24 (Geometric action). *An action of a finitely generated group G on an \mathbb{R} -tree T is geometric if there exists a foliated 2-complex X endowed with a free, properly discontinuous, cocompact action of G , such that*

- *each transverse edge of X isometrically embeds into X/\mathcal{F}*
- *T and X/\mathcal{F} are equivariantly isometric.*

In this case, we say that T is dual to X .

Remark. The definition in [LP97] is in terms of a compact foliated 2-complex Σ (here $\Sigma = X/G$) and of a Galois covering $\overline{\Sigma}$ with deck group G (here $\overline{\Sigma} = X$). The two points of view are clearly equivalent.

This definition requires G to be finitely generated. For finitely generated pairs, one should weaken the cocompactness hypothesis, but we won't enter into this kind of consideration.

Decomposition of geometric actions

Proposition 1.25. *Let $G \curvearrowright T$ be a geometric action dual to a 2-complex X whose fundamental group is generated by free homotopy classes of curves contained in leaves.*

Then T has a decomposition into a graph of actions where each non-degenerate vertex action is either indecomposable, or is an arc containing no branch point of T except possibly at its endpoints.

The arcs in the lemma above are called *edges*, and the indecomposable vertex actions are called *indecomposable components*.

The basis of the proof of this proposition is a version of a theorem by Imanishi giving a dynamical decomposition of a compact foliated 2-complex Σ ([Ima79]). A leaf of Σ is *regular* if it contains no vertex of Σ . More generally, a leaf segment (i. e. a path contained in a leaf) is *regular* if it contains no vertex of Σ . A leaf or leaf segment which is not regular is *singular*.

Let $\Sigma^* = \Sigma \setminus V(\Sigma)$ be the complement of the vertex set of Σ . It is endowed with the restriction of the foliation of Σ . Let $C^* \subset \Sigma^*$ be the union of leaves of Σ^* which are closed but not compact.

Definition 1.26. *We call the set $C = C^* \cup V(\Sigma)$ the cut locus of Σ .*

The cut locus is a finite union of leaf segments joining two vertices of Σ . In particular, C is compact, and $\Sigma \setminus C$ consists of finitely many connected components. Each component of $\Sigma \setminus C$ is a union of leaves of Σ^* .

The dynamical decomposition of Σ is as follows:

Proposition 1.27 ([LP97, Proposition 1.5]). *Let U be a component of $\Sigma \setminus C$. Then either every leaf of U is compact, or every leaf of U is dense in U .* \square

Proof of Proposition 1.25. Let X be a foliated 2-complex such that T is dual to X . Denote by $q : X \rightarrow T$ the natural map (recall that T is the leaf space made Hausdorff of X). The quotient $\Sigma = X/G$ is a compact foliated 2-complex and the quotient map $\pi : X \rightarrow \Sigma$ is a covering map. Let C be the cut locus of Σ and \tilde{C} its preimage in X .

Let $(U_v)_{v \in V}$ be the family of connected components of $X \setminus \tilde{C}$. Let \overline{U}_v be the closure of U_v in X , and $Y_v = q(\overline{U}_v) \subset T$. Note that $\overline{U}_v \setminus U_v \subset \tilde{C}$ is contained in a union of singular leaves and that any leaf segment in U_v is regular. We shall first prove that the family $\mathcal{Y} = (Y_v)_{v \in V}$ is a transverse covering of T . We will need the following result.

Proposition 1.28 ([LP97, Lemma 3.4]). *Assume that $\pi_1(X)$ is generated by free homotopy classes of curves contained in leaves.*

Then there exists a countable union of leaves \mathcal{S} such that for all $x, y \in X \setminus \mathcal{S}$, $q(x) = q(y)$ if and only if x, y are in the same leaf. \square

Since there are finitely many orbits of singular leaves in X , we can choose such a set \mathcal{S} containing every singular leaf.

Assume that $Y_v \cap Y_w$ is non-degenerate. There is an uncountable number of regular leaves of X meeting both $\overline{U}_v \cap \overline{U}_w$. Since $\overline{U}_v \setminus U_v \subset \tilde{C}$ is contained in a union of singular leaves, there is a regular leaf meeting both U_v and U_w , so $U_v = U_w$, and $v = w$.

We denote by $X^{(t)}$ the subset of the 1-skeleton of X consisting of the union of all closed transverse edges. All the paths we consider are chosen as a concatenation of leaf segments and of arcs in $X^{(t)}$. Let I be an arc in T . Let $a, b \in \Sigma$ be two preimages of the endpoints of I in $X^{(t)}$. Choose a path γ in X joining a to b . We view such a path both as a subset of X and as a map $[0, 1] \rightarrow X$. Since T is an \mathbb{R} -tree, $q(\gamma) \supset I$. Since the cut locus C is a finite union of leaf segments, $\gamma \cap \tilde{C}$ consists of a finite number of points and of a finite number of leaf segments. In particular, γ is a concatenation of a finite number of paths which are contained in \tilde{C} or in some \overline{U}_v . Since a path contained in \tilde{C} is mapped to a point in T , we get that I is covered by finitely many elements of \mathcal{Y} .

To prove that the subtree Y_v is closed in T , it is sufficient to check that given a semi-open interval $[a_0, b_0)$ contained in Y_v , b_0 is contained in Y_v . Consider $a, b \in X^{(t)}$ some preimages of a_0, b_0 , and a path $\gamma : [0, 1] \rightarrow X$ joining a to b . By restricting γ to a smaller interval, we can assume that for all $t < 1$, $q \circ \gamma(t) \neq b_0$. Since the image of each transverse edge of X is an arc in T , $K = q(\gamma)$ is a finite tree, and b_0 is a terminal vertex of K . In particular, there exists $\varepsilon > 0$ such that for $q([1 - \varepsilon, 1]) \subset [a_0, b_0) \subset Y_v$. We can also assume

that $\gamma([1 - \varepsilon, 1])$ is contained in a transverse edge of X . For all but countably many t 's, $\gamma(t) \notin \mathcal{S}$ and lies in a regular leaf. Since $q \circ \gamma(t) \in Y_v$, the leaf l through $\gamma(t)$ meets \overline{U}_v . Since \tilde{C} consists of singular leaves, $l \subset U_v$. In particular, $\gamma(t) \in U_v$, so $\gamma(1) \in \overline{U}_v$ and $b_0 \in Y_v$. We have proved that \mathcal{Y} is a transverse covering of T .

Assume that v is such that any leaf of $\pi(U_v)$ is compact. By definition of the cut locus, this means that every leaf of U_v is regular. In particular, U_v is a union of leaves of X and $q(U_v)$ is open in T . Since every transverse edge of X is embedded into T , the holonomy along any regular leaf is trivial. It follows that U_v is a foliated product: is homeomorphic to $U_v \simeq J \times l$ foliated by $\{*\} \times l$, where J is an open interval in a transverse edge. Therefore, $q(U_v)$ is an open interval isometric to J . Since $q(U_v)$ is open in T , Y_v is an arc containing no branch point of T , except possibly at its endpoints.

Assume that leaves of $\pi(U_v)$ are dense. We shall prove that Y_v is indecomposable. First, we claim that any point in Y_v has a preimage in U_v . This will follow from the fact that for any $x \in \overline{U}_v \setminus U_v$, the leaf through x intersects U_v . Assume on the contrary that there is a leaf l of X which meets \overline{U}_v but does not intersect U_v . Consider l_0 a connected component of $l \cap \overline{U}_v$. This is a leaf of the foliated 2-complex \overline{U}_v . Let G_v be the global stabilizer of U_v , and consider the foliated 2-complex \overline{U}_v/G_v . The natural map $\overline{U}_v/G_v \rightarrow \Sigma$ restricts to an isomorphism between U_v/G_v and a connected component $\Sigma \setminus C$ (namely $\pi(U_v)$). Since U_v/G_v consists of finitely many open cells, \overline{U}_v/G_v is compact. Moreover $\overline{U}_v/G_v \setminus U_v/G_v$ consists of finitely many vertices and finitely many edges contained in leaves. The image λ_0 of l_0 in \overline{U}_v/G_v is a leaf which does not intersect U_v/G_v . Therefore, this leaf is compact and so is any leaf of \overline{U}_v/G_v close to λ_0 . This contradicts the fact that every leaf of $\pi(U_v)$ is dense.

Let's prove indecomposability. Let I_0, J_0 be two arcs in Y_v . By taking I_0 smaller, one can assume that $I_0 = q(I)$ for some arc $I \subset U_v$ contained in a transverse edge. Consider a, b some preimages of the endpoints of J_0 in $U_v \cap X^{(t)}$. Consider a path $\gamma \subset U_v$ joining a to b , and write γ as a concatenation $l_0.\tau_1.l_1.\tau_2 \dots \tau_p.l_p$ where τ_i is contained in a transverse edge and l_i is a leaf segment. Since every leaf is dense in $\pi(U_v)$, for all $x \in \tau_1 \cup \dots \cup \tau_p$, there exists $g_x \in G$ and a leaf segment l_x in U_v joining x to $g_x.I$. Since any leaf segment in U_v is regular, this is still valid with the same g_x on a neighbourhood of x . By compactness, one can choose each g_x in a finite set $F = \{g_1, \dots, g_n\}$. Clearly, J_0 is contained in $g_1.q(I) \cup \dots \cup g_n.q(I)$. Denote by x and y the endpoints of l_i . There is a (regular) leaf segment in U_v joining $g_x.I$ to $g_y.I$, so $q(g_x.I) \cap q(g_y.I)$ is non-degenerate. Indecomposability follows. \square

Remark 1.29. Each indecomposable vertex action Y_v is geometric, and dual to the 2-complex \overline{U}_v . Moreover, if $\pi_1(X)$ is generated by free homotopy classes of curves contained in leaves, then so is $\pi_1(\overline{U}_v)$ (this will be useful in section A). Indeed, consider a curve $\gamma \subset \overline{U}_v$, and a map f from a planar surface S to X such that one boundary component is mapped to γ , and every other boundary component is mapped to a leaf. Let $S_0 \subset S$ be the connected component of $f^{-1}(\overline{U}_v)$ containing \overline{U}_v . Since every connected component of $\partial\overline{U}_v$ is contained in a leaf, each component of ∂S_0 is mapped into a leaf, which proves that $\pi_1(\overline{U}_v)$ is generated by free homotopy classes of curves contained in leaves.

Strong approximation by geometric actions

We now quote a result from [LP97].

Theorem 1.30 ([LP97, Theorem 3.7]). *Consider a minimal action of a finitely generated group G on an \mathbb{R} -tree T .*

Then $G \curvearrowright T$ is a strong limit of a direct system of geometric actions $G_k \curvearrowright T_k$ such that

- Φ_k is one-to-one in restriction to each arc stabilizer of T_k ,

- T_k is dual to a 2-complex X whose fundamental group is generated by free homotopy classes of curves contained in leaves.

□

Remark. The second statement follows from the construction in [LP97, Theorem 2.2].

Since T_k is geometric, it has a decomposition into a graph of actions as described above. We show how to adapt this result to actions of finitely generated pairs.

Proposition 1.31. *Consider a minimal action $(G, \mathcal{H}) \circ T$ of a finitely generated pair on an \mathbb{R} -tree. Then there exists a direct system of minimal actions $(G_k, \mathcal{H}_k) \circ T_k$ converging strongly to $G \circ T$ and such that*

- φ_k and Φ_k are one-to-one in restriction to arc stabilizers of T_k
- φ_k (resp. Φ_k) restricts to an isomorphism between \mathcal{H}_k and \mathcal{H}_{k+1} (resp. \mathcal{H}).
- T_k splits as a graph of actions where each non-degenerate vertex action is indecomposable, or is an arc containing no branch point of T_k except at its endpoints.

Proof. We follow Theorem 2.2 and 3.5 from [LP97]. Let $S \subset G$ be a finite set such that $S_\infty = S \cup H_1 \cdots \cup H_p$ generates G . Let S_k be an exhaustion of S_∞ by finite subsets. Let K_k be an exhaustion of T by finite trees. We assume that for each $i \in \{1, \dots, p\}$, K_k contains a point a_i fixed by H_i .

For each $s \in S_k$, consider $A_s = K_k \cap s^{-1}K_k$ and $B_s = s.A_s = K_k \cap s.K_k$. We obtain a foliated 2-complex Σ_k as follows. For each $s \in S_k$, consider a band $A_s \times [0, 1]$ foliated by $\{*\} \times [0, 1]$, glue $A_s \times \{0\}$ on K using the identity map on A_s , and glue $A_s \times \{1\}$ using the restriction of $s : A_s \rightarrow K_k$. For each $s \in H_i \setminus S_k$, add an edge and glue its endpoints on a_i ; this edge is contained in a leaf.

The fundamental group of Σ_k is isomorphic to the free group $F(S_\infty)$. Let $\varphi : F(S_\infty) \rightarrow G$ be the natural morphism. Let N_k be the subgroup of $\ker \varphi$ generated by free homotopy classes of curves contained in leaves (in particular, any relation among the elements of some H_i lies in N_k). This is a normal subgroup of $F(S_\infty)$, and let $G_k = F(S_\infty)/N_k$. We denote by $\varphi_k : G_k \rightarrow G$ and $\Phi_k : G_k \rightarrow G$ the natural morphisms. Clearly, G is the direct limit of G_k . Let $H_i^{(k)}$ be the image in G_k of the subgroup of $F(S_\infty)$ generated by the elements of H_i . Let $\mathcal{H}_k = \{H_1^{(k)}, \dots, H_p^{(k)}\}$. The pair (G_k, \mathcal{H}_k) is finitely generated, and φ_k and Φ_k restrict to isomorphisms between $H_i^{(k)}$, $H_i^{(k+1)}$, and H_i .

Let $\tilde{\Sigma}_k$ be the Galois covering of Σ_k corresponding to N_k . By definition of N_k , $\pi_1(\tilde{\Sigma}_k)$ is generated by free homotopy classes of curves contained in leaves. Let T_k be the leaf space made Hausdorff of $\tilde{\Sigma}_k$. This is an \mathbb{R} -tree by Theorem 1.23. Each $H_i^{(k)}$ fixes a point in T_k . By Proposition 1.28, each arc stabilizer of T_k embeds into G . Finally, the argument of [LP97, Th. 2.2] applies to prove that $G_k \circ T_k$ converges strongly to $G \circ T$. Minimality follows from Lemma 1.22.

The 2-complex Σ_k is a finite foliated complex, with a finite set of infinite roses attached. In particular, Σ_k has the same dynamical decomposition as a finite foliated 2-complex so one can repeat the argument of Proposition 1.25. □

2 Extended Scott's Lemma

Scott's lemma claims that if G is a direct limit of groups G_k having compatible decompositions into free products, then G itself has such a decomposition (this follows from [Sco73, Th 1.7]). Scott's Lemma is usually proved using Scott's complexity ([Sco73, Th 1.7]). Delzant has defined a refinement of this complexity for morphisms which has many important applications ([Swa]).

The main result of this section is an extension of Scott's Lemma for more general splittings. This will be an essential tool to prove piecewise stability of T .

Theorem 2.1 (Extended Scott's Lemma). *Let $G_k \curvearrowright S_k$ be a sequence of minimal actions of finitely generated groups on simplicial trees, and $(\varphi_k, f_k) : G_k \curvearrowright S_k \rightarrow G_{k+1} \curvearrowright S_{k+1}$ be epimorphisms. Assume that f does not increase edge stabilizers in the following sense:*

$$\forall e \in E(S_k), \forall e' \in E(S_{k+1}), \quad e' \subset f_k(e) \Rightarrow G_{k+1}(e') = \varphi_k(G_k(e)) \quad (*)$$

Then $\varinjlim G_k$ has a non-trivial splitting over the image of an edge stabilizer of some S_k .

Remark. When edge stabilizers of S_k are trivial, we obtain Scott's Lemma.

2.1 Decomposition into folds.

Collapses, folds, and group-folds. We recall and adapt definitions of [Dun98, BF91]. Consider a finitely generated group G acting on a simplicial tree S , without inversion (no element of G flips an edge). Given an edge e of S , collapsing all the edges in the orbit of e defines a new tree S' with an action of G . We say that S' is a *collapse* of S and we call the natural map $S \rightarrow S'$ a *collapse*.

Consider two distinct oriented edges $e_1 = uv_1$, $e_2 = uv_2$ of S having the same origin u ; assume that uv_1 and v_2u are not in the same orbit (as oriented edges). Identifying $g.e_1$ with $g.e_2$ for every $g \in G$ defines a new tree S' on which G acts (without inversion). We say that S' is obtained by *folding* e_1 on e_2 and we call the natural map $S \rightarrow S'$ a *fold*.

Consider a vertex $v \in S$, and N_v a normal subgroup of $G(v)$. Let N be the normal subgroup of G generated by N_v and let $G' = G/N$. The graph $S' = S/N$ is a tree (see fact 2.3 below) on which G' acts. We say that S' is obtained from S by a *group-fold* and we call the natural map $S \rightarrow S'$ a *group-fold*. This map is φ -equivariant where $\varphi : G \rightarrow G'$ is the quotient map.

Decomposition into folds. The following result is a slight variation on a result by Dunwoody ([Dun98, Theorem 2.1]). For previous results of this nature, see [BF91, Sta83].

Proposition 2.2. *Let $(\Phi, F) : G \curvearrowright T \rightarrow G' \curvearrowright T'$ be an epimorphism between finitely supported simplicial actions. Assume that*

$$\forall e \in E(T), \forall e' \in E(T'), \quad e' \subset F(e) \Rightarrow G'(e') = \Phi(G(e)) \quad (*)$$

Then we may subdivide T and T' so that there exists a finite sequence of simplicial actions $G \curvearrowright T = G_0 \curvearrowright T_0, \dots, G_n \curvearrowright T_n = G' \curvearrowright T'$, some epimorphisms $(\varphi_i, f_i) : G_i \curvearrowright T_i \rightarrow G_{i+1} \curvearrowright T_{i+1}$ such that $\Phi = \varphi_{n-1} \circ \dots \circ \varphi_0$, each (φ_i, f_i) satisfies $()$, and is either*

1. *a collapse*
2. *a group-fold*
3. *or a fold between two edges uv, uv' of T_i such that $G_i(u)$ injects into G' under Φ_i .*

Remark. We don't claim that $F = f_{n-1} \circ \dots \circ f_1$.

Proof. We may change F so that it is linear in restriction to each edge of T . Then we can subdivide T and T' so that F maps each vertex to a vertex and each edge to an edge or a vertex. The new map satisfies $(*)$. Set $T_0 = T$, $G_0 = G$.

We describe an iterative procedure. We assume that the construction has begun, and that we are given $(\Phi_i, F_i) : G_i \curvearrowright T_i \rightarrow G' \curvearrowright T'$ satisfying $(*)$.

Step 1. Assume that F_i maps an edge e of T_i to a point. Let T_{i+1} be the tree obtained by collapsing e and let $G_{i+1} = G_i$, $\varphi_i = \text{id}$ and $\Phi_{i+1} = \Phi_i$. Define $f_i : T_i \rightarrow T_{i+1}$ as the collapse map, and $F_{i+1} : T_{i+1} \rightarrow T'$ as the map induced by F_i . Then return to step 1.

Clearly, (id, f_i) and (Φ_{i+1}, F_{i+1}) satisfy $(*)$. Since T_{i+1} has fewer orbits of edges than T_i , we can repeat step 1 until F_i does not collapse any edge of T_i .

Step 2. Assume that there is a vertex $v \in T_i$ such that the kernel N_v of $\Phi_{i|G_i(v)} : G_i(v) \rightarrow G'$ is non-trivial. Let N the normal closure of N_v in G_i , and consider $G_{i+1} = G_i/N \circ T_{i+1} = T_i/N$ the action obtained by the corresponding group-fold. Define $f_i : T_i \rightarrow T_{i+1}$ and $\varphi_i : G_i \rightarrow G_{i+1}$ as the quotient maps. Define $F_{i+1} : T_{i+1} \rightarrow T'$ and $\Phi_{i+1} : G_{i+1} \rightarrow G'$ as the maps induced by F_i and Φ_i . Then return to step 2.

The group fold (φ_i, f_i) automatically satisfies (*). Moreover, (Φ_{i+1}, F_{i+1}) inherits property (*) from (Φ_i, F_i) . This step will repeated only a finite number of times since step 2 decreases the number of orbits of vertices with $N_v \neq \{1\}$.

Step 3. Assume that there exists two edges $e_1 = uv_1, e_2 = uv_2$ of T_i such that $F_i(e_1) = F_i(e_2)$. Define T_{i+1} as the tree obtained by folding e_1 on e_2 , $f_i : T_i \rightarrow T_{i+1}$ as the folding map, and $F_{i+1} : T_{i+1} \rightarrow T'$ as the induced map. Define $G_{i+1} = G_i$, $\varphi_i = \text{id}$ and $\Phi_{i+1} = \Phi_i$. Then return to step 2.

Denote by ε the common image of e_1 and e_2 in T_{i+1} . Since at the beginning of step 3, no group fold can be done, $\Phi_{i|G_i(u)}$ is one-to-one. We claim that e_1 and e_2 cannot be in the same orbit. Indeed, if $e_1 = g.e_2$ for some $g \in G$, then $g \in G_i(u)$ (because T' has no inversion) so $\Phi_i(G_i(e_1)) \subsetneq \Phi_i(\langle G_i(e_1), g \rangle) = \Phi_{i+1}(G_{i+1}(\varepsilon)) \subset G'(F_i(e_1))$, contradicting (*). It follows that step 3 will be repeated only finitely many times because it decreases the number of orbits of edges, and step 2 does not change it. Since F_i satisfies (*) and $\Phi_{i|G_i(u)}$ is one-to-one, we get $\Phi_i(G_i(e_1)) = \Phi_i(G_i(e_2))$ so $G_i(\varepsilon) = G_i(e_1)$ and f_i satisfies (*). The fact that F_{i+1} inherits (*) is clear.

When step 3 cannot be repeated any more, F_i is an isometry. It follows that any $g \in \ker \Phi_i$ fixes T_i pointwise. Since $\Phi_{i|G_i(u)}$ is one-to-one for every vertex, Φ_i is an isomorphism. \square

2.2 Proof of Extended Scott's Lemma

Proof. By Proposition 2.2, we may assume that each map f_k is a collapse, a group-fold or a fold. Let $\text{Ell}(S_k)$ be the subset of G_k consisting of elements fixing a point in S_k . The first Betti number of the graph S_k/G_k coincides with the rank of the free group $G_k/\langle \text{Ell}(S_k) \rangle$. In particular, this Betti number is non-increasing and we can assume that it is constant.

We work at the level of quotient graph of groups $\overline{S}_k = S_k/G_k$ and we denote by $x \mapsto \overline{x}$ the quotient map. Consider an oriented edge \overline{e} of \overline{S}_k with terminal vertex $\overline{v} = t(\overline{e})$. Say that \overline{e} carries the symbol $=$ if the edge morphism $i_{\overline{e}} : G_{\overline{e}} \rightarrow G_{\overline{v}}$ is onto. Otherwise, we say that \overline{e} carries \neq . Define W_k as the set of vertices $\overline{v} \in \overline{S}_k$ such that there is an oriented edge \overline{e} with $t(\overline{e}) = \overline{v}$ carrying \neq . At the level of the tree, W_k is the set of orbits of vertices $v \in S_k$ for which there exists an edge e incident on v with $G_k(e) \subsetneq G_k(v)$. Note that $\#W_k$ is invariant under subdivision.

We claim that $\#W_k$ is non-increasing, and that folds and collapses which don't decrease $\#W_k$ don't change $\text{Ell}(S_k)$.

Assume that $f_k : S_k \rightarrow S_{k+1}$ is induced by the collapse of an edge $e = uv$. Let $\overline{f}_k : \overline{S}_k \rightarrow \overline{S}_{k+1}$ be the induced map. The endpoints of \overline{e} are distinct since otherwise, the first Betti number would decrease. If both \overline{u} and \overline{v} are outside W_k , then the stabilizer of u coincides with that of $f_k(u)$; in particular, $\overline{f}_k(\overline{u}) \notin W_k$ and $\#W_k$ is non-increasing. If $\text{Ell}(S_k)$ increases under the collapse, then both orientations of \overline{e} carry \neq , in which case both \overline{u} and \overline{v} belong to W_k , so $\#W_{k+1} < \#W_k$.

Now assume that $f_k : S_k \rightarrow S_{k+1}$ is the fold of $e_1 = uv_1$ with $e_2 = uv_2$ (in particular $G_k = G_{k+1}$). Denote by v' (resp e') the common image of v_1, v_2 (resp. e_1, e_2) in S_{k+1} , and u' the image of u . Since f_k satisfies (*), e_1 and e_2 are in distinct orbits and $G_k(e') = G_k(e_1) = G_k(e_2)$. Since the first Betti number of \overline{S}_k is constant, $\overline{v}_1 \neq \overline{v}_2$. It may happen that $\overline{u} = \overline{v}_i$ for some $i \in \{1, 2\}$. By subdividing e_1 and e_2 and replacing the original fold by two consecutive folds, we can ignore this case. Thus we assume that $\overline{u}, \overline{v}_1, \overline{v}_2$ are distinct. One has $G_k(v') = \langle G_k(v_1), G_k(v_2) \rangle$. It follows that if both \overline{v}_1 and \overline{v}_2

lie outside W_k , then $\bar{v} \notin W_{k+1}$. Since $G_k(u) = G_k(u')$, $\bar{u} \in W_k$ if and only if $\bar{u}' \in W_{k+1}$. It follows that $\#W_k$ is non-increasing. If $\text{Ell}(S_k)$ increases, then $G_k(v) = \langle G_k(v_1), G_k(v_2) \rangle$ is distinct from both $G_k(v_1)$ and $G_k(v_2)$. Therefore, \bar{e}_1 can't carry = at \bar{v}_1 , since this would imply $G_k(v_1) = G_k(e_1) = G_k(e_2) \subset G_k(v_2)$. Similarly, \bar{e}_2 carries \neq at \bar{v}_2 . Therefore, $\bar{v}_1, \bar{v}_2 \in W_k$, and $\#W_{k+1} < \#W_k$. This proves the claim.

Without loss of generality, we assume that $\#W_k$ is constant. For each k , either f_k is a group-fold, or $\text{Ell}(S_k) = \text{Ell}(S_{k+1})$. Let $N_k = \ker \varphi_{k-1} \circ \dots \circ \varphi_1$. We prove by induction on k that S_0/N_k is a tree and $\text{Ell}(S_0/N_k) = \text{Ell}(S_k) \subset G_k$. We will use the following standard fact:

Fact 2.3. *Let T be a simplicial tree and N a group of isometries. Then T/N is a tree if and only if N is generated by elliptic elements.*

Proof. If the graph of groups T/N is a tree, then its fundamental group is generated by its vertex groups. If T/N is not a tree, killing its vertex groups gives a non-trivial free group. \square

Assume that S_0/N_k is a tree. If f_k is not a group-fold, then $G_k = G_{k+1}$, $N_k = N_{k+1}$ and $\text{Ell}(S_k) = \text{Ell}(S_{k+1})$ so we are done. If f_k is a group fold, then G_{k+1} is the quotient of G_k by a normal subgroup K generated by subset of $\text{Ell}(S_k)$ and $S_{k+1} = S_k/K$. It follows that $S_0/N_{k+1} = (S_0/N_k)/K$ is a tree. For $g \in G_k$, denote by \bar{g} its image in $G_{k+1} = G_k/K$. We get: $\bar{g} \in \text{Ell}(S_k/K) \Leftrightarrow \exists k \in K, gk \in \text{Ell}(S_k) \Leftrightarrow \exists k \in K, gk \in \text{Ell}(S_0/N_k) \Leftrightarrow \bar{g} \in \text{Ell}(S_0/N_{k+1})$. The induction follows.

Since S_0/N_k is a tree for all k , N_k is generated by elliptic elements, and so is $N = \cup_k N_k$. This way, we get an action of $G = \lim_{\rightarrow} G_k = G_0/N$ on the tree S_0/N . Assume that this action has a global fix point and argue towards a contradiction. Let $\{g_1, \dots, g_p\}$ be a generating set of G_0 . There exists $x \in S_0$ and $n_1, \dots, n_p \in N$ such that $g_i n_i x = x$. Choose k large enough so that $n_1, \dots, n_p \in N_k$. Then G_k has a global fix point in S_0/N_k . Since $\text{Ell}(S_0/N_k) = \text{Ell}(S_k)$, G_k fixes a point in S_k , a contradiction.

Finally, an edge stabilizer in $G \circ S_0/N$ is the image of an edge stabilizer in $G_0 \circ S_0$, which concludes the proof. \square

2.3 Relative version

Theorem 2.4 (Extended Scott's Lemma, relative version). *Consider $(G_k, \{H_1^k, \dots, H_p^k\}) \circ S_k$ a sequence of minimal actions of finitely generated pairs on simplicial trees. Let $(\varphi_k, f_k) : G_k \circ S_k \rightarrow G_{k+1} \circ S_{k+1}$ be epimorphisms mapping H_i^k onto H_i^{k+1} . Consider $G = \lim_{\rightarrow} G_k$ the inductive limit and $\Phi_k : G_k \rightarrow G$ the natural map.*

Assume that

1. $\forall e \in E(S_k), \forall e' \in E(S_{k+1}), e' \subset f_k(e) \Rightarrow G_{k+1}(e') = \varphi_k(G_k(e))$ (*)
2. $\forall e \in E(S_k), \forall i, \Phi_k(H_i^k) \not\subset \Phi_k(e)$

Then the pair (G, \mathcal{H}) has a non-trivial splitting over the image of an edge stabilizer of some S_k . Moreover, any subgroup $H \subset G$ fixing a point in some S_k fixes a point in the obtained splitting of G .

Remark 2.5. The additional assumption is necessary as shows the following example. Let A be an unsplittable finitely generated group containing a finitely generated free group F . Let $\{a_1, a_2, \dots\} \subset F$ be an infinite basis of a free subgroup of F , and $F' = F(a'_1, a'_2, \dots)$ another free group with infinite basis. Consider

$$G_k = \langle F * F' \mid a_1 = a'_1, \dots, a_k = a'_k \rangle *_F A.$$

The group G_k is finitely generated relative to F' . This sequence of splittings (over F) satisfies condition $(*)$ but not the additional assumption. The inductive limit of G_k is the unsplittable A .

Proof. Since an action of a finitely generated pair is finitely supported, we can use the decomposition into folds of Proposition 2.2. The proof of Extended Scott's Lemma does not use finite generation until the proof of the non-triviality of the obtained splitting. Recall the notations of the end of the proof of Theorem 2.1: $\Phi_k : G_k \rightarrow G$ is the natural morphism, $N = \ker \Phi_0 : G_0 \rightarrow G$, $N_k = \ker \varphi_{0k} : G_0 \rightarrow G_k$ so that $N = \bigcup_k N_k$.

At this point, we know that we can forget finitely many terms in our sequence of actions so that S_0/N is a tree, and that for all k , the action $G_k \curvearrowright S_0/N_k$ is non-trivial. We assume that $G \curvearrowright G_0/N$ has a global fix point and argue towards a contradiction. Let $S^0 \subset G_0$ be a finite set such that $S^0 \cup H_1^0 \cup \dots \cup H_p^0$ generates G . Consider $\{g_1, \dots, g_q\}$ such that $\{g_1, \dots, g_q\} \cup H_1^0 \cup \dots \cup H_p^0$ generates G_0 . There exists $a \in S_0$ and $n_1, \dots, n_q \in N$ such that $g_j n_j \cdot a = a$ for all $j \in \{1, \dots, q\}$. Let $b_i \in S_0$ be a fix point of H_i^0 . By the second hypothesis, H_i^0 fixes no edge in S_0/N so it fixes a unique point in S_0/N . In particular, the images of b_i and a in S_0/N coincide. Therefore, $b_i = n'_i \cdot a$ for some $n'_i \in N$. Choose k large enough so that N_k contains all those elements n_j, n'_i . Then G_k fixes the image of a in S_0/N_k . This contradicts the non-triviality of $G_k \curvearrowright S_0/N_k$. \square

3 Getting piecewise stability

Theorem 3.1. *Consider a minimal action of a finitely generated pair (G, \mathcal{H}) on an \mathbb{R} -tree T . Assume that*

1. T satisfies the ascending chain condition;
2. there exists a finite family of arcs I_1, \dots, I_p such that $I_1 \cup \dots \cup I_p$ spans T and such that for any unstable arc J contained in some I_i ,
 - (a) $G(J)$ is finitely generated;
 - (b) $G(J)$ is not a proper subgroup of any conjugate of itself i. e. $\forall g \in G, G(J)^g \subset G(J) \Rightarrow G(J)^g = G(J)$.

Then either (G, \mathcal{H}) splits over the stabilizer of an unstable arc contained in some I_i , or T is piecewise-stable.

Proof. By enlarging some peripheral subgroups, we may assume that each $H \in \mathcal{H}$ is either finitely generated, or is not contained in a finitely generated elliptic subgroup of G .

By Proposition 1.31, consider $(G_k, \mathcal{H}_k) \curvearrowright T_k$ a sequence of actions on \mathbb{R} -trees converging strongly to T , and such that T_k splits as a graph of actions where each vertex action is either an indecomposable component, or an edge. Denote by $\varphi_k : G_k \rightarrow G_{k+1}$, $\Phi_k : G_k \rightarrow G$, $f_k : T_k \rightarrow T_{k+1}$, and $F_k : T_k \rightarrow T$ the morphisms of the corresponding direct system.

Without loss of generality, we can assume that there exists $\tilde{I}_1, \dots, \tilde{I}_p \subset T_0$ which map isometrically to I_1, \dots, I_p . By subdividing the arcs \tilde{I}_i and the edges occurring in the decomposition of T_k , we may assume that each \tilde{I}_i is either an edge or an indecomposable arc of T_0 . All the arcs of T_k we are going to consider embed isometrically into T . Say that an arc $I \subset T_k$ is *pre-stable* if its image in T (under F_k) is a stable arc. Otherwise, we say that I is *pre-unstable*. If I is contained in an indecomposable component, then it is pre-stable by Lemma 1.20 and Assertion 1 of Lemma 1.19.

To do some bookkeeping among pre-unstable edges, we shall construct inductively a combinatorial graph $\mathcal{T} = \mathcal{T}(\tilde{I}_1) \sqcup \dots \sqcup \mathcal{T}(\tilde{I}_p)$ as a disjoint union of rooted trees. We use standard terminology for rooted trees: the father of v is the neighbour of v closer to the root than v , a child of v is a neighbour of v which is not the father of v , an ancestor of v if a vertex on the segment joining v to the root. The level of a vertex is its distance to

the root. We denote by $\mathcal{T}_k(\tilde{I}_i)$ (resp. $\mathcal{T}_{\geq k}(\tilde{I}_i)$) the set of vertices of $\mathcal{T}(\tilde{I}_i)$ of level k (resp. at least k), and $\mathcal{T}_k = \mathcal{T}_k(\tilde{I}_1) \cup \dots \cup \mathcal{T}_k(\tilde{I}_p)$. (resp. $\mathcal{T}_{\geq k} = \mathcal{T}_{\geq k}(\tilde{I}_1) \cup \dots \cup \mathcal{T}_{\geq k}(\tilde{I}_p)$).

Each vertex v of level k of $\mathcal{T}(\tilde{I}_i)$ will be labeled by a pre-unstable edge J_v contained in $f_{0k}(\tilde{I}_i) \subset T_k$. We label the root of $\mathcal{T}(\tilde{I}_i)$ by \tilde{I}_i . Assume that \mathcal{T} has been constructed up to the level k . Subdivide the edge structure of T_{k+1} so that, for each $v \in \mathcal{T}_k$, $f_k(J_v)$ is a finite union of edges and of indecomposable arcs. We may also assume that the length of each edge is at most $1/2^k$. The indecomposable pieces of $f_k(J_v)$ are pre-stable, and we discard them. We also discard pre-stable edges of $f_k(J_v)$. For each pre-unstable edge J' contained in $f_k(J_v)$, we add new child of v labeled by J' .

If \mathcal{T} is finite, then each \tilde{I}_i is contained in a finite union of stable arcs. Since $\tilde{I}_1 \cup \dots \cup \tilde{I}_p$ spans T , T is piecewise stable. So we assume that \mathcal{T} is infinite.

To each vertex $v \in \mathcal{T}_k$, we attach two subgroups of G : $A_v = \Phi_k(G_k(J_v))$, and $B_v = G(F_k(J_v))$. Clearly, $A_v \subset B_v$. Given $u, v \in \mathcal{T}$, write $B_u < B_v$ if B_u is properly contained in some conjugate of B_v . Say that $v \in \mathcal{T}_k$ is *minimal* if there is no $u \in \mathcal{T}_k$ with $B_u < B_v$. Let $\mathcal{M} \subset \mathcal{T}$ be the set of minimal vertices, and $\mathcal{M}_k = \mathcal{M} \cap \mathcal{T}_k$. Since no B_v is a proper conjugate of itself, for each v , either $v \in \mathcal{M}_k$ or there exists $u \in \mathcal{M}_k$ with $B_u < B_v$.

Lemma 3.2. *There exists k_0 such that for all $k \geq k_0$, the following hold:*

1. *if $v \in \mathcal{T}_k \setminus \mathcal{M}_k$, then no child of v is minimal;*
2. *if $v \in \mathcal{M}_k$ then for any child $v' \in \mathcal{M}_{k+1}$ of v , $A_v = A_{v'}$;*

Proof. Say that v is a *clone* of v' if $B_v = B_{v'}$ and $A_v = A_{v'}$. A *genealogical line* is a sequence v_0, v_1, \dots of vertices of \mathcal{T} where v_{i+1} is a child of v_i . Say that $v \in \mathcal{T}$ is *immortal* if there is a genealogical line starting at v and consisting of clones of v .

We claim that any genealogical line v_0, v_1, \dots eventually consists of clones. By the ascending chain condition in T , there are at most finitely many indices i such that $B_{v_i} \subsetneq B_{v_{i+1}}$. So we can assume that $B_{v_i} = B_{v_0}$ for all i . Since J_{v_0} is pre-unstable, B_{v_0} is finitely generated. The strong convergence implies that for k large enough, every generator of B_{v_0} has a preimage in $G_k(f_{k_0k}(J_{v_0}))$ (where k_0 is the level of v_0). Since for each v_i of level at least k , $J_{v_i} \subset f_{k_0k}(J_{v_0})$, we get $B_{v_0} \subset A_{v_i} \subset B_{v_i} = B_{v_0}$. Therefore, all the vertices v_i of sufficiently large level are clones of each other. This proves the claim. In particular, this also proves that if v is immortal, then $A_v = B_v$.

Say that v is *post-immortal* if it is immortal or if it has an immortal ancestor. We claim that all but finitely many vertices of \mathcal{T} are post-immortal. Indeed, for each $i \in \{1, \dots, p\}$, the set of vertices of $\mathcal{T}(\tilde{I}_i)$ which are not post-immortal is a rooted subtree of $\mathcal{T}(\tilde{I}_i)$. If it is infinite, it contains a genealogical line. Since this line contains many immortal vertices, this is a contradiction.

Let k_0 be such that $\mathcal{T}_{\geq k_0}$ consists of post-immortal vertices. We shall use several times the following fact: any $v \in \mathcal{M}_{\geq k_0}$ is the clone of an immortal vertex. Indeed, let u' be an immortal ancestor of v , and let u be a clone of u' of same level as v ; then $A_{u'} = B_{u'} \subset A_v \subset B_v$, and since v is minimal, inclusions are equalities, and u' is a clone of v . It follows that for all $v \in \mathcal{M}_{\geq k_0}$, $A_v = B_v$.

If $v \in \mathcal{T}_k \setminus \mathcal{M}_k$, there exists $u \in \mathcal{M}_k$ with $B_u < B_v$. Since u is the clone of an immortal vertex, there exists u' of level $k+1$ with $B_u = B_{u'}$ and no child of v can be minimal. This proves assertion 1.

If $v \in \mathcal{M}_k$, and $v' \in \mathcal{M}_{k+1}$ is a child of v , then v has an immortal clone, so has an immortal clone u' of level $k+1$. We have $B_{u'} = A_{u'} = A_v \subset A_{v'} \subset B_{v'}$ and since v' is minimal, v' is a clone of v . Assertion 2 follows. \square

We always consider $k \geq k_0$. Let \mathcal{E}_k be the set of edges of our decomposition of T_k which are in the orbit of some J_v for some $v \in \mathcal{M}_k$. In our decomposition of T_k as a graph of actions, collapse all the vertex trees which are not in \mathcal{E}_k (see Definition 1.7). The resulting tree S_k is a graph of actions where each vertex action is an edge, so it is a simplicial tree.

The action $G_k \circ S_k$ is minimal by Lemma 1.9 and is therefore non-trivial. By assertion 1 of Lemma 3.2, f_k induces a natural map $S_k \rightarrow S_{k+1}$. Since any $H \in \mathcal{H}_k$ fixes a point in T_k , it also does in S_k , so the action on S_k is an action of the pair (G_k, \mathcal{H}_k) .

Let's check that the actions $G_k \circ S_k$ satisfy the hypotheses of Extended Scott's lemma. Consider an edge $e \in \mathcal{E}_k$. We may translate e so that e corresponds to an arc J_v , with $v \in \mathcal{M}_k$. Let $e' \in \mathcal{E}_{k+1}$ with $e' \subset f_k(e)$. Since $e' \in \mathcal{E}_{k+1}$, it is pre-unstable, so $e' = J_{v'}$ for some child $v' \in \mathcal{T}$ of v . Moreover, since $e' \in \mathcal{E}_{k+1}$, e' is in the orbit of some $J_{w'}$ with w' minimal, so $v' \in \mathcal{M}_{k+1}$. By assertion 2 of Lemma 3.2, $\Phi_{k+1}(G_{k+1}(e')) = A_{v'} = A_v = \Phi_k(G_k(e))$. Since Φ_k and Φ_{k+1} are one-to-one in restriction to arc stabilizers, we get $\varphi_k(G_k(e)) = G_{k+1}(e')$.

If $\mathcal{H} = \emptyset$, this is enough to apply Extended Scott's Lemma, so G splits over some group A_v . Since $A_v = B_v$ is the stabilizer of an unstable arc in T , the theorem is proved in the non-relative case.

If $\mathcal{H} \neq \emptyset$, we need to modify slightly the argument to ensure that the second hypothesis of the relative version of Extended Scott's Lemma holds. Let $\mathcal{H}' \subset \mathcal{H}$ be the subset consisting of subgroups which are not finitely generated and $\mathcal{H}'_k \subset \mathcal{H}_k$ the subset corresponding to \mathcal{H}' . The pairs (G, \mathcal{H}') and (G_k, \mathcal{H}'_k) are finitely generated. Recall that no $H \in \mathcal{H}'$ is contained in a finitely generated elliptic subgroup. In particular, for all $H \in \mathcal{H}'_k$, $\Phi_k(H)$ fixes no unstable arc of T . For each $e \in E(S_k)$, $\Phi_k(G_k(e))$ fixes an unstable arc in T so Theorem 2.4 applies. We thus get a non-trivial splitting $(G, \mathcal{H}') \circ S$.

In view of the moreover part of Theorem 2.4, since any $H \in \mathcal{H}_0$ fixes a point in S_0 , any $H \in \mathcal{H}$ fixes a point in S . Thus, S defines a non-trivial splitting of (G, \mathcal{H}) . \square

4 Piecewise stable actions

The goal of this section is the following result:

Theorem 4.1. *Let (G, \mathcal{H}) be a finitely generated pair having a piecewise stable action on an \mathbb{R} -tree T .*

Then, either (G, \mathcal{H}) splits over the stabilizer H of an infinite tripod (and the normalizer of H contains a non-abelian free group generated by two hyperbolic elements whose axes don't intersect), or T has a decomposition into a graph of actions where each vertex action is either

1. *simplicial: $G_v \circ Y_v$ is a simplicial action on a simplicial tree;*
2. *of Seifert type: the vertex action $G_v \circ Y_v$ has kernel N_v , and the faithful action $G_v/N_v \circ Y_v$ is dual to an arational measured foliation on a closed 2-orbifold with boundary;*
3. *axial: Y_v is a line, and the image of G_v in $\text{Isom}(Y_v)$ is a finitely generated group acting with dense orbits on Y_v .*

The proof relies on the following particular case of a result by Sela:

Theorem 4.2 ([Sel97]). *Consider a minimal action of a finitely generated group G on an \mathbb{R} -tree T with trivial arc stabilizers.*

Then, either G is freely decomposable, or T has a decomposition into a graph of actions where each vertex action is either

1. *of surface type: the vertex action $G_v \circ Y_v$ is dual to an arational measured foliation on a closed 2-orbifold with boundary;*
2. *axial: Y_v is a line, and the image of G_v in $\text{Isom}(Y_v)$ is a finitely generated group acting with dense orbits on Y_v .*

We shall give a proof of this result in Appendix A.

4.1 From piecewise stability to trivial arc stabilizers

Lemma 4.3. *Let G be a group having a piecewise stable action on an \mathbb{R} -tree T .*

Then T has a decomposition into a graph of actions such that, denoting by N_v the kernel of the vertex action $G_v \curvearrowright Y_v$, then $G_v/N_v \curvearrowright Y_v$ has trivial arc stabilizers.

Remark. The lemma doesn't assume any finite generation of G .

Proof. Remember that a subtree $Y \subset T$ is stable if the stabilizer of any arc $J \subset Y$ fixes Y . Given a stable arc $I \subset T$, consider Y_I the maximal stable subtree containing I . This is a well defined subtree because if two stable subtrees contain I , their union is still stable, and an increasing union of stable subtrees is stable. Moreover, Y_I is closed in T because the closure of a stable subtree is stable.

Let $(Y_v)_{v \in V}$ be the family of all maximal stable subtrees of T . By piecewise stability, any arc of T is contained in a finite union of them. By maximality, if $Y_u \cap Y_v$ is non-degenerate, then $Y_u = Y_v$. Therefore, the family $(Y_v)_{v \in V}$ is a transverse covering of T . Denote by G_v the global stabilizer of Y_v , and by N_v its pointwise stabilizer. Since Y_v is a stable subtree, $G_v/N_v \curvearrowright Y_v$ has trivial arc stabilizers. \square

4.2 Relative version of Sela's Theorem


Consider a decomposition of T as a graph of actions as in Lemma 4.3. Consider a vertex group G_v of the corresponding graph of groups Γ . Even if G is finitely generated, G_v may fail to be finitely generated. However, in this case, the peripheral structure (G_v, \mathcal{H}_v) of G_v in Γ is finitely generated (Lemma 1.15 and Lemma 1.11), and the action on T_v is an action of the pair (G_v, \mathcal{H}_v) . If we started with a finitely generated pair (G, \mathcal{H}) , G_v is finitely generated with respect to a set \mathcal{H}_v consisting of the peripheral structure of G_v in Γ together with some conjugates of elements of \mathcal{H} (Lemma 1.12). We will need a version of Sela's result applying in this context.

Proposition 4.4 (Relative version of Sela's result). *Consider a minimal action of a finitely generated pair (G, \mathcal{H}) on an \mathbb{R} -tree T with trivial arc stabilizers.*

Then, either (G, \mathcal{H}) is freely decomposable, or T has a decomposition into a graph of actions where each non-degenerate vertex action is either

1. *of surface type: the vertex action $G_v \curvearrowright Y_v$ is dual to an arational measured foliation on a closed 2-orbifold with boundary;*
2. *axial: Y_v is a line, and the image of G_v in $\text{Isom}(Y_v)$ is a finitely generated group acting with dense orbits on Y_v .*

Proof. We shall embed G into a finitely generated group \hat{G} and apply Sela's non-relative result. Let $\mathcal{H} = \{H_1, \dots, H_p\}$. We may assume that each H_i is non-trivial. For $i = 1, \dots, p$, choose some finitely generated group \hat{H}_i containing H_i . We may assume that \hat{H}_i is freely indecomposable by changing \hat{H}_i to $\hat{H}_i \times \mathbb{Z}/2\mathbb{Z}$. Consider the graph of groups Γ below.



We denote by u (resp. u_i) the vertex labeled by G (resp. \hat{H}_i). Clearly, the group $\hat{G} = \pi_1(\Gamma)$ is finitely generated.

We shall add a structure of a graph of actions on \mathbb{R} -trees on Γ . Let Y_u be a copy of T endowed with its natural action of G , and let Y_{u_i} be a point endowed with the trivial

action of \hat{H}_i . For each edge uu_i of Γ , we define its attaching point in Y_u as the (unique) point of T fixed by H_i . Let $\hat{G} \circ \hat{T}$ be the \mathbb{R} -tree dual to this graph of actions.

Since T is a union of axes of elements of G , and since \hat{T} is covered by translates of T , \hat{T} is a union of axes of elements of \hat{G} so \hat{T} is minimal.

This action has trivial arc stabilizers because $G \circ T$ does. Therefore, we can apply Sela's Theorem 4.2 to $\hat{G} \circ \hat{T}$.

Assume first that \hat{G} is freely decomposable, i. e. that \hat{G} acts non-trivially on a simplicial tree R with trivial edge stabilizers. Then \hat{H}_i is elliptic in R because it is freely indecomposable. If G fixes a point $x \in R$, then each H_i fixes x and cannot fix any other point because edge stabilizers are trivial. It follows that each \hat{H}_i fixes x and that \hat{G} fixes x , a contradiction. Thus, the action of G on R defines a non-trivial free decomposition of G relative to H_i .

Assume now that \hat{T} has a decomposition into a graph of actions where each vertex action is axial or of surface type. In particular, each vertex action is indecomposable. Let (Z_i) be the transverse covering of \hat{T} induced by this decomposition. Since Z_i is indecomposable, if $Z_i \cap T$ is non-degenerate, then $Z_i \subset T$ (Lemma 1.18). The family of subtrees Z_i contained in T is therefore a transverse covering of T . This gives a decomposition of $G \circ T$ as a graph of actions where each vertex action is axial or of surface type. \square

4.3 Proof of Theorem 4.1

Proof of Theorem 4.1. Consider a transverse covering $\mathcal{Y} = (Y_v)_{v \in V}$ of T as in Lemma 4.3. Denote by G_v the global stabilizer of Y_v , and by $N_v \triangleleft G_v$ the kernel of the vertex action $G_v \circ Y_v$. We denote by $\overline{G}_v = G_v/N_v$ the group acting with trivial arc stabilizers on Y_v .

We shall prove that either (G, \mathcal{H}) splits over a tripod stabilizer with the desired properties, or that each vertex action $G_v \circ Y_v$ has a decomposition into a graph of actions of the right kind. The theorem will follow.

If Y_v is a line, then $G_v \circ Y_v$ is either simplicial or of axial type, so we can assume that Y_v is not a line.

By Lemma 1.6, we can first decompose Y_v into a graph of actions where each vertex action has a global fixed point or a dense minimal subtree. Therefore, we can assume without loss of generality that for each v , \overline{G}_v has either a global fixed point or a dense minimal subtree in Y_v .

Lemma 4.5. *If \overline{G}_v fixes a point in Y_v , then Y_v is a simplicial tree.*

Proof. Using Lemma 1.16, consider a finite tree K spanning Y_v . Let $x_0 \in K$ be a \overline{G}_v -invariant point. Denote by K_1, \dots, K_n the closure of the connected components of $K \setminus \{x_0\}$. If K_i, K_j are such that there exists $g \in \overline{G}_v$ with $g.K_i \cap K_j$ non-degenerate, replace K_i and K_j by $g.K_i \cup K_j$. This way, one can assume that for all $i \neq j$ and all $g \in \overline{G}_v$, $g.K_i \cap K_j = \{x_0\}$. Since arc stabilizers are trivial, for all $g \in \overline{G}_v \setminus \{1\}$, $g.K_i \cap K_i = \{x_0\}$. Thus, Y_v is the union of translates of K_1, \dots, K_n , all glued along $\{x_0\}$. Since each K_i is a finite tree, Y_v is a simplicial tree. \square

Now we consider the case where $\overline{G}_v \circ Y_v$ has a dense minimal subtree. Since Y_v is not a line, Y_v contains an infinite tripod. This tripod is fixed by N_v . Moreover, consider $\overline{g}, \overline{h} \in \overline{G}_v$ two hyperbolic elements having distinct axes. In particular $[\overline{g}, \overline{h}] \neq 1$. Since arc stabilizers are trivial, the axes of \overline{g} and \overline{h} have compact intersection. Therefore, \overline{g} and $\overline{g}' = \overline{h}^k$ have disjoint axes for k large enough. Let g, g' be some preimages of $\overline{g}, \overline{g}'$ in G . Then $\langle g, g' \rangle$ is a free group generated by two elements having disjoint axes in T , and which normalizes N_v .

Lemma 4.6. *If Y_v is not minimal, then G splits over N_v .*

Proof. Let $G \circ S$ be the skeleton S of the transverse covering \mathcal{Y} . Since $G \circ T$ is minimal, $G \circ S$ is minimal (Lemma 1.15). Therefore, we need only to prove that there is an edge of S whose stabilizer is N_v .

Let $x \in Y_v \setminus \min(Y_v)$. Since $\min(Y_v)$ is dense in Y_v and does not contain x , $Y_v \setminus x$ is connected. If x does not lie in any other tree $Y \in \mathcal{Y}$, then $T \setminus \{x\}$ is convex, so $T \setminus G.x$ is a G -invariant subtree, contradicting the minimality of T .

Therefore, x is a vertex of S and (x, Y_v) is an edge of S (see section 1.3). Denote by $G(x, Y_v) = \{g \in G_v | g.x = x\}$ its stabilizer. Clearly, $N_v \subset G(x, Y_v)$. Conversely, consider $g \in G(x, Y_v)$ and $y \in Y_v \setminus \{x\}$. Since y and $g.y$ both lie in the convex set $Y_v \setminus \{x\}$, $[x, y] \cap [x, g.y]$ is a non-degenerate arc fixed by g . Therefore, $g \in N_v$. \square

There remains to analyse the case where Y_v is not a line and $G_v \circ Y_v$ is minimal without global fix point. Consider the graph of groups $\Gamma = S/G$. We identify v with a vertex of Γ . By Lemma 1.12 (or 1.11 if G is finitely generated), G_v is finitely generated relative to the incident edge groups together with at most one conjugate of each $H \in \mathcal{H}$. When $\mathcal{H} \neq \emptyset$, we make sure that for any $H \in \mathcal{H}$ having a conjugate in G_v , \mathcal{H}_v contains a conjugate of H . Let $\overline{\mathcal{H}}_v$ be the image of \mathcal{H}_v in $\overline{G}_v = G_v/N_v$. Then we can apply the relative version of Sela's Theorem (Proposition 4.4) to $(\overline{G}_v, \overline{\mathcal{H}}_v) \circ Y_v$.

Assume first that $(\overline{G}_v, \overline{\mathcal{H}}_v)$ is freely decomposable. Therefore, G_v splits over N_v relative to \mathcal{H}_v . Thus, we can refine Γ at v using this splitting, so G splits over N_v . This is really a splitting of (G, \mathcal{H}) because we made sure that for each $H \in \mathcal{H}$, either H has a conjugate in \mathcal{H}_v , or H is conjugate in some other vertex group of Γ .

Assume now that $\overline{G}_v \circ Y_v$ has a decomposition into a graph of actions where each vertex action is either axial or of surface type. Then clearly, the action $G_v \circ Y_v$ as a decomposition into a graph of actions where each vertex action is either axial or of Seifert type. \square

5 Proof of Main Theorem and corollaries

Recall that an action of the pair (G, \mathcal{H}) on a tree is an action of G where each $H \in \mathcal{H}$ fixes a point. In terms of graphs of groups, a splitting of the pair (G, \mathcal{H}) is an isomorphism of G with a graph of groups such that each H_i is contained in a conjugate of a vertex group (see section 4.2).

Theorem 5.1 (Main Theorem). *Consider a minimal action of finitely generated pair (G, \mathcal{H}) on an \mathbb{R} -tree T by isometries. Assume that*

1. T satisfies the ascending chain condition;
2. there exists a finite family of arcs I_1, \dots, I_p such that $I_1 \cup \dots \cup I_p$ spans T (see Definition 1.13) and such that for any unstable arc J contained in some I_i ,
 - (a) $G(J)$ is finitely generated;
 - (b) $G(J)$ is not a proper subgroup of any conjugate of itself i. e. $\forall g \in G, G(J)^g \subset G(J) \Rightarrow G(J)^g = G(J)$.

Then either (G, \mathcal{H}) splits over the stabilizer of an unstable arc, or over the stabilizer of an infinite tripod (whose normalizer contains a non-abelian free group generated by two elements having disjoint axes), or T has a decomposition into a graph of actions where each vertex action is either

1. simplicial: $G_v \circ Y_v$ is a simplicial action;
2. of Seifert type: the vertex action $G_v \circ Y_v$ has kernel N_v , and the faithful action $G_v/N_v \circ Y_v$ is dual to an arational measured foliation on a closed 2-orbifold with boundary;

3. *axial*: Y_v is a line, and the image of G_v in $\text{Isom}(Y_v)$ is a finitely generated group acting with dense orbits on Y_v .

Remark. The group over which G splits (i. e. the stabilizer of an unstable arc or of a tripod) is really its full pointwise stabilizer. This contrasts with [BF95, Theorem 9.5].

Proof of Main Theorem. Let $(G, \mathcal{H}) \circ T$ be as in Main Theorem. By Theorem 3.1, if (G, \mathcal{H}) does not split over the stabilizer of an unstable arc, T is piecewise stable. By Theorem 4.1, either (G, \mathcal{H}) splits over the stabilizer of an infinite tripod with the required properties, or T has a decomposition into a graph of actions of the desired type. \square

Corollary 5.2. *Under the hypotheses of Main Theorem, either T is a line or (G, \mathcal{H}) splits over a subgroup H which is an extension of a cyclic group (maybe finite or trivial) by an arc stabilizer.*

Proof. We can assume that T splits as a graph of actions \mathcal{G} where each vertex action $G_v \circ Y_v$ is either simplicial, of Seifert type, or axial. Let $(Y_v)_{v \in V}$ be the family of non-degenerate vertex trees. Let S be the skeleton of this transverse covering (see Lemma 1.5).

First, consider the case where S is reduced to a point v . This means that $T = Y_v$, and $G = G_v$, and that T is itself simplicial or of Seifert type (T is not a line so cannot be of axial type). If T is simplicial, the result is clear, so assume that T is of Seifert type. Let $N_v \triangleleft G_v$ be the kernel of this action and let Σ be a 2-orbifold with boundary, with cone singularities, such that $G_v/N_v = \pi_1(\Sigma)$ and holding an arational measured foliation to which $G_v/N_v \circ Y_v$ is dual. Consider a splitting of G_v/N_v corresponding to an essential simple closed curve (i. e. a curve which cannot be homotoped to a point, a cone point, or to the boundary). Such a curve exists because Σ holds an arational measured foliation. This defines a splitting of $\pi_1(\Sigma)$ over a cyclic group, and a splitting of $G = G_v$ over an extension of a cyclic group by N_v . Any subgroup $H \subset G$ elliptic in T corresponds to the fundamental group of a cone point or of a boundary component of Σ . Thus H is elliptic in this splitting and we get a splitting of the pair (G, \mathcal{H}) .

Now assume that S is not reduced to a point. By Lemma 1.15, $G \circ S$ is minimal and any $H \in \mathcal{H}$ is elliptic in S . In particular, given any edge e of S , the corresponding splitting of G over $G(e)$ is non-trivial. We shall prove that for some edge e of S , $G(e)$ is an extension of an arc stabilizer by a cyclic group, and the corollary will follow.

Assume that some action $G_v \circ Y_v \simeq \mathbb{R}$ is axial. Consider an edge $e = (x, Y_v)$ of S incident on v (see Lemma 1.5). Its stabilizer $G(e)$ is the stabilizer of x in G_v . Since the stabilizer of x in G_v/N_v is either trivial or $\mathbb{Z}/2\mathbb{Z}$, $G(e)$ is an extension of a cyclic group by N_v .

Assume that some action $G_v \circ Y_v$ is of Seifert type. Consider an edge $e = (x, Y_v)$ incident on v in S . Since $G(e)$ is the stabilizer of x in G_v , $G(e)$ is an extension by N_v of the stabilizer of a point in $\pi_1(\Sigma) \circ Y_v$ which is cyclic.

The only remaining case is when each $G_v \circ Y_v$ is simplicial. In this case $G \circ T$ is simplicial, and the result is clear. \square

Corollary 5.3. *Let G be a finitely generated group for which any small subgroup is finitely generated. Assume that G acts on an \mathbb{R} -tree T with small arc stabilizers.*

Then G splits over the stabilizer of an unstable arc, or over a tripod stabilizer, or T has a decomposition into a graph of actions as in Main Theorem. In particular, G splits over a small subgroup.

Proof. We prove that all hypotheses of the Main Theorem are satisfied. The set of small subgroups is closed under increasing union. Since small subgroups are finitely generated, any ascending chain of small subgroups is finite. The ascending chain condition follows.

Similarly, if $H \subsetneq H^g$ for some small subgroup $H \subset G$, then the set of small subgroup H^{g^n} is an infinite increasing chain, a contradiction. Then Main Theorem and corollary 5.2 apply. \square

In the following situation, stabilizers of unstable arcs can be controlled by tripod stabilizers.

Corollary 5.4. *Consider a finitely generated group G acting by isometries on an \mathbb{R} -tree T . Assume that*

1. *arc stabilizers have a nilpotent subgroup of bounded index (maybe not finitely generated);*
2. *tripod stabilizers are finitely generated (and virtually nilpotent);*
3. *no group fixing a tripod is a proper subgroup of any conjugate of itself;*
4. *any chain $H_1 \subset H_2 \dots$ of tripod stabilizers stabilizes.*

Then either G splits over a subgroup having a finite index subgroup fixing a tripod, or T has a decomposition as in the conclusion of Main Theorem.

Proof. Let k be a bound on the index of a nilpotent subgroup in an arc stabilizer. We first claim that the stabilizer of an unstable arc has a subgroup of bounded index fixing a tripod. Indeed, let A be the stabilizer of an unstable arc I , and let $B \supsetneq A$ be the stabilizer of a sub-arc J . If A has index at most k in B , then A contains a normal subgroup A' of index at most $k!$. Since for $g \in B \setminus A$, $g.I$ is an arc distinct from I and containing J , $B.I$ contains a tripod. This tripod is fixed by A' . Suppose that the index of A in B is larger than k . Let N_B be a nilpotent subgroup of index at most k in B and $N_A = A \cap N_B$. Because of indices, $N_A \subsetneq N_B$. Let N be the normalizer of N_A in N_B . It is an easy exercise to check that in a nilpotent group, no proper subgroup is its own normalizer so $N_A \subsetneq N$. Therefore, $N.I$ contains a tripod, and this tripod is fixed by N_A .

It follows that the stabilizer of an unstable arc is finitely generated. Since each tripod stabilizer is slender, any ascending chain of subgroups fixing tripods stabilizes. It follows that the stabilizer of an unstable arc cannot be properly conjugated into itself. Consider $A_1 \subsetneq \dots \subsetneq A_n \subsetneq \dots$ an ascending chain of arc stabilizers. Let N_n be a subgroup of bounded index of A_n fixing a tripod, and let $N_{i,n}$ be its intersection with A_i for $i \leq n$. Since $N_{i,n}$ has bounded index in A_i , it takes finitely many values for each i . By a diagonal argument, we get an ascending chain of subgroups fixing tripods, a contradiction. Thus, Main Theorem applies, and the corollary is proved. \square

6 An example

The goal of this section is to give an example of an action with trivial tripod stabilizers providing a counter-example to [RS94, Theorem 10.8] and [Sel97, Theorem 2.3 and 3.1]. Note however that the proof of Theorem 3.1 in [Sel97] is valid under the stronger assumption that T is *super-stable*, which is a natural hypothesis since it is satisfied in applications in [Sel97, Sel01a].

Theorem 6.1. *There exists a non-trivial minimal action of a group G on an \mathbb{R} -tree T such that*

- *G is finitely presented (and even word hyperbolic) and freely indecomposable*
- *tripod stabilizers are trivial*
- *T satisfies the ascending chain condition*
- *T has no decomposition into a graph of actions as in Main Theorem.*

Proof. Let A be a freely indecomposable hyperbolic group containing a free malnormal subgroup $M_1 = \langle a, b_1 \rangle$. For instance, one can take for A a surface group, and for M_1 , the fundamental group of a punctured torus contained in this surface. Let A' be another copy of the group A , $C = \langle a \rangle$, and let $G = A *_C A'$. By Bestvina-Feighn's Combination Theorem, G is hyperbolic [BF92].

For $i > 1$, define inductively $M_i \subset M_{i-1}$ by $M_i = \langle a, b_i \rangle$ where $b_i = b_{i-1} a b_{i-1}^2$. One can easily check that M_i is malnormal in M_{i-1} (one can also use the software Magnus to do so [Coo05]). Moreover, one also easily checks that $\bigcap_{i \geq 1} M_i = C$ since any reduced word w on $\{a, b_i\}$ defines a reduced word on $\{a, b_{i-1}\}$ by the obvious substitution, and its length is strictly larger if w is not a power of a .

Let Γ_k be the graph of groups and $G \circlearrowleft T_k$ be the corresponding Bass-Serre tree as shown in Figure 2.




Figure 2: The graph of groups Γ_k and the tree T_k

Let a_k, a'_k be the vertices of T_k fixed by A and A' respectively. Denote by $e_i^k \subset [a_k, a'_k]$ the edge of T_k such that $G(e_i^k) = M_i$, and by $\varepsilon^k \subset [a_k, a'_k]$ the edge such that $G(\varepsilon^k) = C$. Figure 2 shows a neighbourhood of $[a_k, a'_k]$ in T_k , and the action of the vertex stabilizers. The main feature is that from each vertex v with stabilizer M_i , the neighbourhood of v consists of one orbit of edges $M_i.e_{i+1}^k$, and one single edge e_i^k .

Assign length $1/2^i$ to the edge e_i^k , and length $1/2^k$ to ε^k . This way, $d(a_k, a'_k) = 1$ for all k . There is a natural morphism of \mathbb{R} -trees $f_k : T_k \rightarrow T_{k+1}$ sending e_i^k to e_i^{k+1} and ε^k to $e_{k+1}^{k+1} \cup \varepsilon^{k+1}$.

The length function of T_{k+1} is not larger than the length function of T_k , so T_k converges in the length function topology to an action on an \mathbb{R} -tree T . This action is non-trivial since for any $g \in A \setminus C$ and $g' \in A' \setminus C$, the translation length of gg' in every T_k is 2.

We shall prove that T_k converges strongly to T . Consider α, β two distinct edges of T_0 sharing a vertex v (T_0 is the tree dual to the amalgam $G = A *_C A'$). If v is in the orbit of a'_0 , then the image of α and β in T_k share only one point. Now assume without loss of generality that $v = a$, and $\alpha = \varepsilon^0$. Let $g \in A \setminus C$ be such that $g.\alpha = \beta$. Consider k_0 the smallest integer such that, $g \notin M_{k_0}$. Then for $k \geq k_0$, the image of α and β in T_k share exactly an initial segment of length $1 - 1/2^{k_0}$. In particular, f_k is one-to-one in restriction to the union of the images of α and β . Now if K is a finite subtree of T_0 , it follows that for k large enough, f_k is one-to-one in restriction to the image of K in T_k . This proves the strong convergence of T_k .

Say that an arc $I \subset T_k$ is *immersed in Γ_k* if the restriction to I of the quotient map $T_k \rightarrow \Gamma_k$ is an immersion. We claim that for any arc $I \subset T_k$ which is not immersed in Γ_k , $G(I)$ is trivial. It is sufficient to prove it for I of the form $I = \alpha \cup \beta$ where $\alpha, \beta \in E(T_k)$ are incident on a common vertex and are in the same orbit. The triviality of $G(I) = G(\alpha) \cap G(\beta)$ follows easily from the malnormality of M_1 in A , of M_{i+1} in M_i , and of C in M_k and A' . Since a tripod cannot be immersed, it follows that tripod stabilizers of T_k are trivial. Going to the limit, tripod stabilizers of T are trivial.

Now we study arc stabilizers of T . Denote by $F_k : T_k \rightarrow T$ the natural map, and by $a = F_k(a_k)$ and $a' = F_k(a'_k)$. Let $I = [u, v] \subset T$ be a non-degenerate arc with non-trivial stabilizer. By strong convergence, if $g \in G(I)$ then there exists a lift $I_k \subset T_k$ in restriction

to which F_k is isometric, and which is fixed by g . The argument above shows that the image of I_k in Γ_k is immersed. In particular, we can assume that $I_k \subset [a_k, a'_k]$. If I contains a' , then since $F_k^{-1}(a') = \{a'_k\}$, I_k intersects ε_k in a non-degenerate segment, so $G(I_k) = C$ and $G(I) = C$. If I does not contain a' , for k large enough, I_k does not intersect ε_k so $G(I_k) = M_i$ for some i independent of k . It follows that $G(I) = M_i$. The ascending chain condition for T follows immediately.

Since the restriction of F_k to the ball of radius $1 - 1/2^k$ around a_k is an isometry, the segment $[a, a'] \subset T$ contains an infinite number of branch points. In particular, T is not a simplicial tree.

We claim that no subtree Y of T is indecomposable. Since $[a, a']$ spans T , we may assume that $Y \cap [a, a']$ contains a non-degenerate arc I . By indecomposability, the orbit of any point of I is dense in I , a contradiction.

Finally, assume that T has a decomposition as in Main Theorem. Since there are no indecomposable subtrees in T , T is simplicial, a contradiction. \square

A Sela's Theorem

Theorem ([Sel97]). *Consider a minimal action of a finitely generated group G on an \mathbb{R} -tree T with trivial arc stabilizers.*

Then, either G is freely decomposable, or T has a decomposition into a graph of actions where each non-degenerate vertex action is either

1. *of surface type: the vertex action $G_v \curvearrowright Y_v$ is dual to an arational measured foliation on a closed 2-orbifold with boundary;*
2. *axial: Y_v is a line, and the image of G_v in $\text{Isom}(Y_v)$ is a finitely generated group acting with dense orbits on Y_v .*

A.1 Preliminaries

The following lemma is essentially contained in [LP97].

Lemma A.1. *Let $G \curvearrowright T$ be a minimal action with trivial arc stabilizers.*

There exists a sequence of \mathbb{R} -trees $G_k \curvearrowright T_k$ converging strongly to $G \curvearrowright T$ such that T_k is geometric, dual to a foliated 2-complex $G_k \curvearrowright X_k$ whose leaf space is Hausdorff, and such that $\Phi_k : G_k \rightarrow G$ is one-to-one in restriction to each point stabilizer.

Proof. Consider a sequence of actions $G_k \curvearrowright T_k$ dual to foliated 2-complexes X_k as in Theorem 1.30. Let N_k be the kernel of $\Phi_k : G_k \rightarrow G$. Let L_k be the subgroup of N_k generated by elements which preserve a leaf of X_k . This is a normal subgroup of G_k . Let $X'_k = X_k/L_k$ be the quotient foliated 2-complex, endowed with the natural action of $G'_k = G_k/L_k$. By construction, for each leaf l of X'_k , the global stabilizer of l embeds into G under the induced morphism $\Phi'_k : G'_k \rightarrow G$.

The natural map $X_k \rightarrow X'_k$ is a covering. Since $\pi_1(X_k)$ is generated by free homotopy classes of curves contained in leaves, and since L_k is generated by elements preserving a leaf, $\pi_1(X'_k)$ is generated by free homotopy classes of curves contained in leaves. Let T'_k be the leaf space made Hausdorff of X'_k . It is an \mathbb{R} -tree by Theorem 1.23. Since $L_k \subset N_k$, the map $f_k : T_k \rightarrow T$ factors through the natural map $T_k \rightarrow T'_k$. It follows that T'_k is geometric, dual to X'_k , and that $G'_k \curvearrowright T'_k$ converge strongly to $G \curvearrowright T$.

By Proposition 1.28, there exists a countable union of leaves \mathcal{S} on the complement of which two points of X'_k are identified in T'_k if and only if they lie on the same leaf. It follows that any element $g \in G'_k$ fixing an arc in T'_k preserves a leaf of X'_k . Since arc stabilizers of T are trivial, the image of g in G is trivial, so $g = 1$. Thus, arc stabilizers of $G'_k \curvearrowright T'_k$ are trivial.

By proposition A.3 below, the leaf space of X'_k is Hausdorff. In particular, a point stabilizer of T'_k coincides with a leaf stabilizer of X'_k , which embeds into G . The lemma follows. \square

Definition A.2. *We will say that $G \circledast T$ is nice if*

- *T is geometric, dual to a foliated 2-complex $G \circledast X$ such that $\pi_1(X)$ is generated by free homotopy classes of curves contained in leaves;*
- *arc stabilizers of T are trivial.*

The following result follows from the concatenation of Lemma 3.5 and 3.4 in [LP97]:

Proposition A.3 ([LP97]). *Assume that T is nice.*

Then the leaf space of X is Hausdorff: any two points of X identified in T are in the same leaf.

Lemma A.4. *Assume that $G \circledast T$ is nice. Let Y be an indecomposable component of T (as in Proposition 1.25), and H its global stabilizer.*

Then $H \circledast Y$ is nice.

Proof. Let $G \circledast X$ be a foliated 2-complex such that T_k is dual to X . Let $C \subset X/G$ be its cut locus (see Definition 1.26), and \tilde{C} its preimage in X . The tree Y is dual to the closure \bar{U} of a connected component U of $X \setminus \tilde{C}$. By Remark 1.29, $\pi_1(\bar{U})$ is generated by free homotopy classes of curves contained in leaves. This means that Y is nice. \square

Proposition A.5. *Assume that $G \circledast T$ is nice and let $H \circledast Y$ be an indecomposable component. Then one of the following holds:*

- *Axial type: Y is a line, and the image of H in $\text{Isom}(Y)$ is a finitely generated group acting with dense orbits on Y ;*
- *Surface type: H is the fundamental group of a 2-orbifold with boundary Σ holding an arational measured foliation and Y is dual to $\tilde{\Sigma}$;*
- *Exotic type: H has a non-trivial free decomposition $H \circledast S$ in which any subgroup of H fixing a point in Y fixes a point in S .*

Proof. This proposition is essentially well known: it is a way of describing the output of the Rips machine. This would follow from [BF95] if we knew that H is finitely presented (because of the finiteness hypothesis in [BF95, Definition 5.1]). But we can apply some arguments of [Gui98b] where finite presentation is not assumed.

We recall some vocabulary from [GLP94] or [Gui98b]. A closed multi-interval D is a finite union of compact intervals. A *partial isometry* of D is an isometry between closed sub-intervals of D . A *system of isometries* \mathcal{S} on D is a finite set of partial isometries of D . Its *suspension* is the foliated 2-complex obtained from D by gluing for each partial isometry $\varphi : I \rightarrow J \in \mathcal{S}$ by a foliated band $I \times [0, 1]$ on D joining I to J whose holonomy is given by φ . A *singleton* is a partial isometry $\varphi : I \rightarrow J$ where I is reduced to a point. We also call singleton the band corresponding to a singleton in the suspension of a system of isometries. The $\mathring{\mathcal{S}}$ -orbits are the equivalence classes for the equivalence relation generated by $x \sim y$ if $y = \varphi(x)$ for some non-singleton $\varphi : I \rightarrow J$ such that $x \in \mathring{I}$. A system of isometries \mathcal{S} is *minimal* if \mathcal{S} has no singleton and every $\mathring{\mathcal{S}}$ -orbit of any point in $D \setminus \partial D$ is dense. The system \mathcal{S} is *simplicial* if every \mathcal{S} -orbit is finite.

The proposition holds if $H \circledast Y$ is dual to a minimal system of isometries: this follows from [Gui98b] section 5 (axial case), proposition 7.2 (exotic case), and section 8 (surface case).

Now $H \circledast Y$ is dual to some foliated 2-complex $H \circledast X$, occurring as a Galois covering of the suspension of some system of isometries \mathcal{S} . Following [GLP94], [BF95] or [Gui98b, Prop 3.1], one can perform a sequence of Rips moves on \mathcal{S} so that \mathcal{S} becomes a disjoint union

finitely many systems of isometries which are either minimal or simplicial, together with a finite set of singletons joining them. This decomposition induces a decomposition of Y as a graph of actions as in Proposition 1.25. Since Y is indecomposable, this decomposition is trivial, and Y is dual to the suspension of a minimal component of \mathcal{S} . \square

A.2 Proof of the theorem

Proof of Sela's Theorem. Let $G_k \circ T_k$ be a sequence of geometric actions converging strongly to $G \circ T$ as in lemma A.1. Arc stabilizers of $G_k \circ T_k$ are trivial. By Proposition 1.25, T_k splits as a graph of actions \mathcal{G} , where each non-degenerate vertex action is either simplicial or indecomposable.

Assume that for all k , the simplicial part of T_k is non-empty. We shall prove that G is freely decomposable. Collapse the indecomposable components of T_k as in Definition 1.7. Let S_k be the obtained tree. Clearly, S_k is a simplicial tree, and edge stabilizers are trivial because arc stabilizer of T_k are trivial. The action $G_k \circ S_k$ is minimal by Lemma 1.9 and therefore non-trivial. The map $f_k : T_k \rightarrow T_{k+1}$ maps an indecomposable component of T_k into an indecomposable component of T_{k+1} so f_k induces a map $S_k \rightarrow S_{k+1}$. By Scott's Lemma, G is freely decomposable.

Now, we assume that for all k , T_k splits as a graph of indecomposable actions. Let S_k be the skeleton of the corresponding transverse covering of T_k . Recall that its vertex set $V(S)$ is $V_0(S) \cup V_1(S)$ where $V_1(S) = \mathcal{Y}$, and $V_0(S)$ is the set of points $x \in T$ lying in the intersection of two distinct trees of \mathcal{Y} . Since f_k maps an indecomposable tree into an indecomposable tree, f_k induces a map $V_1(S_k) \rightarrow V_1(S_{k+1})$. Moreover, f_k induces a map $V_0(S_k) \rightarrow V_0(S_{k+1}) \cup V_1(S_{k+1})$: for $x \in V_0(S_k)$, if $f_k(x)$ belongs to two distinct indecomposable components, we map x to $f_k(x) \in V_0(S_{k+1})$, otherwise, we map x to the only indecomposable component containing $f_k(x)$. This map extends to a map $g_k : S_k \rightarrow S_{k+1}$ sending an edge to an edge or a vertex. The number of orbits of edges of S_k is non-increasing so for k large enough, no edge of S_k is collapsed by g_k , and any pair of edges e_1, e_2 identified by $g_k(e_1)$ are in the same orbit. Moreover, using Scott's Lemma, we can assume that for k large enough, no edge of S_k has trivial stabilizer.

The following lemma will be proved in next sections.

Lemma A.6 (Stabilisation of indecomposable components). *For k large enough, the following holds. Consider an indecomposable component $H \circ Y$ of T_k , and let $H' \circ Y'$ be the indecomposable component of T_{k+1} containing $f_k(Y)$.*

Then $f_k|_Y$ is an isometry from Y onto Y' . Moreover, if Y is not a line, then $\varphi_k(H) = H'$.

Using this lemma, we shall prove that for k large enough, $f_k : T_k \rightarrow T_{k+1}$ is an isometry. If this is not the case, then there exists two arcs $J_1, J_2 \subset T_k$ with $J_1 \cap J_2 = \{x\}$ and $f_k(J_1) = f_k(J_2)$. By shortening them, we may assume that J_1 and J_2 lie in some indecomposable components Y_1, Y_2 . By Lemma A.6, $Y_1 \neq Y_2$. Therefore, the edges (x, Y_1) and (x, Y_2) of S_k are identified under g_k . By the assumption above, they lie in the same orbit. Consider $g \in G_k(x)$ such that $g.Y_1 = Y_2$. Let $Y' = g_k(Y_1) = g_k(Y_2)$.

First, assume that Y_1 is not a line. Since $\varphi_k(g)$ preserves Y' , there exists $\tilde{g} \in G_k(Y_1)$ with $\varphi_k(\tilde{g}) = \varphi_k(g)$ by Lemma A.6. Since f_k is isometric in restriction to Y_1 and since $\varphi_k(\tilde{g}) = \varphi_k(g)$ fixes $f_k(x)$, \tilde{g} fixes x . In particular, $g^{-1}\tilde{g}$ fixes x and lies in the kernel of φ_k . By Lemma A.1, φ_k is one-to-one in restriction to point stabilizers, so $g = \tilde{g}$. This is a contradiction because $g \notin G_k(Y_1)$.

Therefore, Y_1 is a line. Since φ_k is one-to-one in restriction to $G_k(x)$, $\varphi_k(g) \neq 1$. Since $\varphi_k(g)$ preserves the line Y' , and since arc stabilizers are trivial, $\varphi_k(g)$ acts on Y' as the reflection fixing $f_k(x)$. By the assumption above, edge stabilizers of S_k are non-trivial so consider $h \neq 1$ in the stabilizer of the edge (x, Y_1) . Since h fixes a point in T_k (namely, x),

$\varphi_k(h) \neq 1$. So $\varphi_k(h)$ acts as the same reflection as $\varphi_k(g)$ on Y' . Since arc stabilizers are trivial, $\varphi_k(g) = \varphi_k(h)$. Since g, h are contained in a point stabilizer, we get $g = h$. This is a contradiction since $g \notin G_k(Y_1)$.

This proves that for k large enough, f_k is an isometry. It follows that φ_k is an isomorphism because any element of $\ker \varphi_k$ must fix T_k pointwise.

Thus, Proposition A.5 gives us a decomposition of T . If there is an exotic type vertex action $H \odot Y$, the free splitting of H given in Proposition A.5 can be used to refine the decomposition of G induced by the skeleton S_k , thus giving a decomposition of G as a free product.

Finally, there remains prove the finite generation claimed in the axial case. Let $H \odot Y$ be an indecomposable component such that Y is a line. In particular, H occurs as a vertex group in the graph of groups given by the decomposition of T . By Lemma 1.11, H is finitely generated relative to the stabilizers of incident edges H_1, \dots, H_p . Denote by $\psi : H \rightarrow \text{Isom}(Y)$ the map induced by the action. Since each H_i fixes a point in Y , $\psi(H_i)$ is cyclic of order at most 2. Since $\psi(H)$ is finitely generated relative to $\psi(H_1), \dots, \psi(H_p)$, $\psi(H)$ is finitely generated. \square

A.3 Standard form for indecomposable components

We set up the material needed for the proof of Lemma A.6. Consider $H \odot Y$ a nice and indecomposable action. Given an arc $I \subset Y$ and a finite generating set $S \subset H$, we construct a foliated 2-complex $X(I, S)$ as a kind of thickened Cayley graph. Alternatively, one can view $X(I, S)$ as the covering space with deck group H of the suspension of a system of isometries on I . The point here is to start with an arbitrary arc I and to allow S to be large in order to ensure that Y is dual to $X(I, S)$.

We first define $X(I, S)$ for any arc $I \subset Y$, and any finite subset $S \subset H$. Start with $H \times I$, endowed with the action of H given by $g.(h, x) = (gh, x)$. For each $s \in S$, consider $K_s = (I \cap s^{-1}I)$, and for each $g \in H$, add a foliated band $K_s \times [0, 1]$ joining $\{g\} \times K_s$ to $\{gs\} \times s.K_s$ whose holonomy is given by the restriction of s to K_s . The map sending (g, x) to $g.x$ extends uniquely to a map $p : X(I, S) \rightarrow Y$ which is constant on each leaf.

Lemma A.7. *Consider $H \odot Y$ a nice indecomposable action. For every arc $I \subset Y$, there exists a finite generating set S of H such that*

- (i) *the leaf space of $X(I, S)$ is Hausdorff;*
- (ii) *Y is dual to $X(I, S)$.*

Before proving the lemma, we introduce some tools. A *holonomy band* in a foliated 2-complex Σ is a continuous map $b : I \times [0, 1] \rightarrow \Sigma$ such that $b|_{I \times \{0\}}$ (resp. $b|_{I \times \{1\}}$) is an isometric map to a subset of a transverse edge of Σ , and for each $x \in I$, $b(\{x\} \times [0, 1])$ is contained in a leaf segment, and this leaf segment is regular if $x \notin \partial I$. The following property is a restatement of Theorem 2.3 in [GLP95].

Proposition A.8 (Segment-closed property, [GLP95]). *Let Σ be a compact foliated 2-complex. Consider two arcs $I, J \subset \Sigma^{(1)}$ and an isometry $\varphi : I \rightarrow J$ such that for all but countably many $x \in I$, x and $\varphi(x)$ are in the same leaf.*

Then there exist a subdivision $I = I_1 \cup \dots \cup I_p$, and for each $i \in \{1, \dots, p\}$ a holonomy band b_i joining I_i to $\varphi(I_i)$ whose holonomy is the restriction of φ . \square

Corollary A.9. *Assume that $G \odot T$ is nice, dual to a foliated 2-complex $G \odot X$.*

Consider two arcs I, J contained in transverse edges of X and having the same image in T . Then there exist subdivisions $I = I_1 \cup \dots \cup I_p$, $J = J_1 \cup \dots \cup J_p$, such that for each $i \in \{1, \dots, p\}$ there exists a holonomy band joining I_i to J_i .

Proof. Let $\varphi : I \rightarrow J$ be an isometry such that for all $x \in I$, x and $\varphi(x)$ map to the same point in T . By Proposition 1.28, for all but countably many $x \in I$, x lies in the same leaf as $\varphi(x)$. Apply segment closed property to the images of I, J in the compact foliated 2-complex X/G . By lifting the obtained holonomy bands to X , we obtain a subdivision $I = I_1 \cup \dots \cup I_p$ and for each $i \in \{1, \dots, p\}$ a holonomy band b_i joining I_i to $g_i \cdot \varphi(I_i) \subset g_i \cdot J$ for some $g_i \in G$. Since x and $\varphi(x)$ map to the same point in T , g_i fixes the arc I_i , so $g_i = 1$. \square

Proof of Lemma A.7. Consider a foliated 2-complex $H \circlearrowleft X$ such that Y is dual to X . Let $\pi : X \rightarrow X/H$ be the covering map. Consider the cut locus $C \subset X/H$, \tilde{C} its preimage in X . The set $U = X \setminus \tilde{C}$ is connected: otherwise, Y would split as a graph of actions with at least two non-degenerate vertex trees by Proposition 1.25; this is impossible because Y is indecomposable (Lemma 1.18). Let $q : X \rightarrow Y$ the quotient map (Y is the leaf space of X).

Recall that $X^{(t)} \subset X^{(1)}$ denotes the union of all closed transverse edges of X . We aim to construct a collection of partial isometries Φ from $X^{(t)}$ to $H \times I$, and use Φ to transport the holonomy of the triangles of X to bands on $H \times I$.

Consider a finite set of arcs J_1, \dots, J_p contained in edges of $X^{(t)}$ such that $H \cdot (J_1 \cup \dots \cup J_p) = X^{(t)}$ and $q(J_1) \cup \dots \cup q(J_p) \supset I$. By subdividing each J_i , one may assume that either $q(J_i) \subset I$, or $q(J_i) \cap I$ is degenerate. Since Y is indecomposable, I spans Y , so up to further subdivision of each J_i , one may assume that for each $i \in \{1, \dots, p\}$ there exists $g_i \in H$ such that $q(g_i \cdot J_i) \subset I$. By replacing J_i by $g_i \cdot J_i$ for all i such that $q(J_i) \cap I$ is degenerate, we get $q(J_1) \cup \dots \cup q(J_p) = I$.

Let $\varphi_i : J_i \rightarrow q(J_i)$ be the partial isometry defined as the restriction of q to J_i . We view φ_i as a partial isometry whose domain of definition is $\text{dom } \varphi_i = J_i$ and whose range is $\text{Im } \varphi_i = \{1\} \times q(J_i) \subset X(I, S)$. For $g \in H$, we consider the g -conjugate of φ_i defined by

$$\varphi_i^g = g \circ \varphi_i \circ g^{-1} : g \cdot J_i \rightarrow \{g\} \times I.$$

Let $\Phi = \{\varphi_i^g | g \in H, i = 1, \dots, p\}$. By construction, $\bigcup_{\varphi \in \Phi} \text{dom } \varphi = X^{(t)}$, and $\bigcup_{\varphi \in \Phi} \text{Im } \varphi = H \times I$.

There is a kind of commutative diagram

$$\begin{array}{ccc} X^{(t)} & \xrightarrow{\Phi} & H \times I \\ & \searrow q & \swarrow p \\ & & Y \end{array}$$

meaning that for each $x \in X^{(t)}$, and any $\varphi \in \Phi$ defined on x , $q(x) = p(\varphi(x))$.

We now build a foliated complex X' by gluing foliated bands on $H \times I$. For each pair of arcs $g_1 \cdot J_{i_1}, g_2 \cdot J_{i_2} \subset X^{(t)}$ whose intersection K is non-empty, we add a foliated band joining $\varphi_{i_1}^{g_1}(K)$ to $\varphi_{i_2}^{g_2}(K)$ with holonomy $\varphi_{i_2}^{g_2} \circ (\varphi_{i_1}^{g_1})^{-1}$. For each triangle τ of X , and for each pair of arcs $g_1 \cdot J_{i_1}, g_2 \cdot J_{i_2} \subset \partial\tau$ such that the holonomy along the leaves of τ defines a partial isometry $\psi : K_1 \subset g_1 \cdot I_1 \rightarrow K_2 \subset g_2 \cdot I_2$, we add a foliated band joining $\varphi_{i_1}^{g_1}(K_1)$ to $\varphi_{i_2}^{g_2}(K_2)$, with holonomy $\varphi_{i_2}^{g_2} \circ \psi \circ (\varphi_{i_1}^{g_1})^{-1}$.

The important property is that Φ maps leaves to leaves in the following sense: if $x, y \in X^{(t)}$ are in the same leaf, and if $\varphi, \psi \in \Phi$ are defined on x and y respectively, then $\varphi(x)$ and $\psi(y)$ are in the same leaf of X' .

Since X is locally finite, so is X' . Therefore, the natural free, properly discontinuous action of H on X' is cocompact. Let $S \subset H$ be the finite set of elements $s \in H$ such that there is a band connecting $(1, I)$ to (s, I) . Then X' is naturally contained in $X(I, S)$. And Φ , viewed as a collection of maps $X^{(t)} \rightarrow X(I, S)$, still maps leaves to leaves.

Let $T(I, S)$ be the leaf space made Hausdorff of $X(I, S)$ and $\tilde{p} : X(I, S) \rightarrow T(I, S)$ be the quotient map. The map $p : H \times I \rightarrow Y$ being constant on any leaf of $X(I, S)$

(or rather on its intersection with $H \times I$), and isometric in restriction to any connected component of $H \times I$, it induces a natural distance decreasing map $f : T(I, S) \rightarrow Y$.

$$\begin{array}{ccc}
 X^{(t)} & \xrightarrow{\Phi} & H \times I \\
 q \downarrow & \nearrow p & \downarrow \tilde{p} \\
 Y & \xleftarrow{f} & T(I, S) \\
 & \searrow g & \\
 & &
 \end{array}$$

Since Φ maps leaves to leaves, and since Y is the set of leaves of $X^{(t)}$ (it is Hausdorff by Proposition A.3), Φ induces a map $g : Y \rightarrow T(I, S)$. This map is distance decreasing because any path in X written as a concatenation of leaf segments and of transverse arcs in $X^{(t)}$ defines a path having the same transverse measure in $X(I, S)$ as follows: subdivide its transverse pieces so that they can be mapped to $X(I, S)$ using Φ , and join the obtained paths by leaf segments of $X(I, S)$. Since f and g are distance decreasing, both are isometries. In particular Y is dual to $X(I, S)$.

We prove that the leaf space of $X(I, S)$ is Hausdorff. Assume that $x, y \in H \times I$ have the image under p . Consider $\tilde{x}, \tilde{y} \in X$ and $\varphi, \psi \in \Phi$ such that $\varphi(\tilde{x}) = x$ and $\psi(\tilde{y}) = y$. Then $q(\tilde{x}) = q(\tilde{y})$, so \tilde{x} and \tilde{y} are in the same leaf because the leaf space of X is Hausdorff. Therefore, x and y are in the same leaf of $X(I, S)$. \square

Remark. If the lemma holds for some $S \subset H$, it also holds for any S' containing S .

A.4 Pseudo-groups

Up to now we considered partial isometries between closed intervals. We now need to consider partial isometries between *open* intervals. We use notations like $\mathring{D}, \mathring{\varphi}$ to emphasize this point. An open *multi-interval* \mathring{D} is a finite union of copies of bounded open intervals of \mathbb{R} . The *pseudo-group of isometries* generated by some partial isometries $\mathring{\varphi}_1, \dots, \mathring{\varphi}_n$ between open intervals of D , is the set of partial isometries $\mathring{\varphi} : \mathring{I} \rightarrow \mathring{J}$ such that for any $x \in \mathring{D}$, there exists a composition $\mathring{\varphi}_{i_1}^{\pm 1} \circ \dots \circ \mathring{\varphi}_{i_k}^{\pm 1}$ which is defined on a neighbourhood of x , and which coincides with $\mathring{\varphi}$ on this neighbourhood. Two points $x, y \in \mathring{D}$ are in the same orbit if there exists $\mathring{\varphi} \in \Lambda$ such that $y = \mathring{\varphi}(x)$. A pseudo-group is *minimal* if its orbits are dense in \mathring{D} .

Consider an arc $I \subset Y$, where $H \circlearrowleft Y$ is an indecomposable nice action. Let $\mathring{I} = I \setminus \partial I$ and $\mathring{\varphi}_s$ be the restriction of s to the interior of $s^{-1} \cdot \mathring{I} \cap \mathring{I}$. Let Λ_0 be the pseudo-group of isometries on \mathring{I} generated $\{\mathring{\varphi}_s | s \in S\}$. Let $\mathring{D} \subset \mathring{I}$ be the set of points whose orbit under Λ_0 is infinite.

Let $\Lambda(I, S)$ be the restriction of Λ_0 to \mathring{D} . We claim that $\mathring{I} \setminus \mathring{D}$ is finite (in particular, \mathring{D} is an open multi-interval) and that $\Lambda(I, S)$ is minimal. Let C be the cut locus of $\Sigma = X(I, S)/H$. Since Y is indecomposable, $\Sigma \setminus C$ consists of only one minimal component, so every leaf l of $\Sigma \setminus C$ is dense in $\Sigma \setminus C$. Since for any such leaf l , $l \cap \mathring{I}$ is contained in a Λ_0 -orbit, the claim follows.

A pseudo-group of isometries Λ is *homogeneous* if for each $\mathring{\varphi} \in \Lambda$, any partial isometry $\mathring{\psi}$ extending $\mathring{\varphi}$ lies in Λ . When $\Lambda(I, S)$ is homogeneous, Y is a line (see for instance [Gui98b, section 5]).

We will use the following result, which is a slight generalisation of a result of [Lev93], which deals with the orientation preserving case.

Theorem A.10 ([Gui, Gui98a]). *Let Λ be a minimal finitely generated pseudo-group of isometries of an open multi-interval \mathring{D} . Then the set of non-homogeneous pseudo-groups of isometries containing Λ is finite.*

A.5 Stabilisation of indecomposable components

We are now ready to prove the stabilisation of indecomposable components.

Proof of Lemma A.6. Let $G_k \odot T_k$ be a sequence of geometric actions converging strongly to $G \odot T$ as in lemma A.1. Recall that $(\varphi_k, f_k) : G_k \odot T_k \rightarrow G_{k+1} \odot T_{k+1}$ and $(\Phi_k, F_k) : G_k \odot T_k \rightarrow G \odot T$ denote the maps of the corresponding direct system.

Let $H_{k_0} \odot Y_{k_0}$ be an indecomposable component of T_{k_0} . Consider $k \geq k_0$. Since $f_{k_0 k}(Y_{k_0})$ is indecomposable, it is contained in an indecomposable component Y_k of T_k (see Lemma 1.18 and 1.19). Let $I_{k_0} \subset Y_{k_0}$ be an arc which embeds into T under F_{k_0} , and let $I_k = f_{k_0 k}(I_{k_0})$.

By Lemma A.7, Y_k is dual to $X(I_k, S_k)$ for some finite set $S_k \subset H_k$. By enlarging each S_k , we can assume that for all k , $\varphi_k(S_k) \subset S_{k+1}$. Let $\Lambda(I_k, S_k)$ be the corresponding minimal pseudo-group of isometries on $\mathring{D}_k \subset \mathring{I}_k$. Under the natural identification between I_k and I_{k+1} , we get $\mathring{D}_k \subset \mathring{D}_{k+1}$, and $\Lambda(I_k, S_k) \subset \Lambda(I_{k+1}, S_{k+1})$. Since $I_k \setminus \mathring{D}_k$ is finite, for k large enough, $\mathring{D}_{k+1} = \mathring{D}_k$.

By Theorem A.10 above, for k large enough, either $\Lambda(I_k, S_k)$ is homogeneous, or $\Lambda(I_k, S_k) = \Lambda(I_{k+1}, S_{k+1})$. It is an exercise to show that if Λ is a minimal homogeneous pseudo-group of isometries, then any pseudo-group of isometries containing Λ is homogeneous. Thus, in the first case, $\Lambda(I_{k+1}, S_{k+1})$ is homogeneous. It follows that Y_k and Y_{k+1} are lines. Since H_k acts with dense orbits on Y_k , the morphism of \mathbb{R} -trees $f_k|_{Y_k} : Y_k \rightarrow Y_{k+1}$ is necessarily one-to-one: there exists an arc of Y_k which is embedded under f_k and using the action of H_k , we see that f_k is locally isometric. It follows that $f_k(Y_k) = Y_{k+1}$.

In the second case, the following result concludes the proof. \square

Lemma A.11 (The pseudo-group determines the action). *Assume that $(\varphi, f) : H \odot Y \rightarrow H' \odot Y'$ is a morphism between nice indecomposable actions. Consider $I \subset Y$ such that f is isometric in restriction to I and let $I' = f(I)$. Consider finite subsets $S \subset H$ and $S' \subset H'$ such that $\varphi(S) \subset S'$.*

Assume that Y is dual to $X(I, S)$, that Y' is dual to $X(I', S')$, and that $\Lambda(I, S) = \Lambda(I', S')$.

Then $f : Y \rightarrow Y'$ is a surjective isometry and $\varphi : H \rightarrow H'$ is an isomorphism.




Figure 3: The pseudo-group determines the action

Proof. Assume that f is not one-to-one. There exists two arcs $[x, a_1], [x, a_2] \subset Y$ such that $[x, a_1] \cap [x, a_2] = \{x\}$ and $f([x, a_1]) = f([x, a_2])$. Since I spans Y , one may shorten

these arcs and assume without loss of generality that $[x, a_1] \subset I$ and $[x, a_2] \subset h.I$ for some $h \in H$.

Consider $a' = f(a_1) = f(a_2)$, $x' = f(x)$ and $h' = \varphi(h)$ (see Figure 3). The two points $\tilde{a}' = h'^{-1}a'$ and $\tilde{x}' = h'^{-1}x'$ lie in I' . The subsets $\{1\} \times [x', a']$ and $\{h'\} \times [\tilde{x}', \tilde{a}']$ of $X(I', S')$ map to the same arc in Y' . By Corollary A.9, there exist arcs $[x', b'] \subset [x', a']$, $[\tilde{x}', \tilde{b}'] \subset [\tilde{x}', \tilde{a}']$, and a holonomy band in $X(I', S')$ joining $\{1\} \times [x', b']$ to $\{h'\} \times [\tilde{x}', \tilde{b}']$. This holonomy band defines a partial isometry $\psi' \in \Lambda(I', S')$ such that $\psi'((x', b')) = (\tilde{x}', \tilde{b}')$. By hypothesis, ψ' corresponds to an element of $\psi \in \Lambda(I, S)$, sending (x, b) to (\tilde{x}, \tilde{b}) where b, \tilde{x}, \tilde{b} are the points of I corresponding to $b', \tilde{x}', \tilde{b}'$ under the natural identification.

We now apply segment closed property to the partial isometry induced by ψ in $X(I, S)/H$. We get the existence of arcs $[x, c] \subset [x, b]$ and $[\tilde{x}, \tilde{c}] \subset [\tilde{x}, \tilde{b}]$ and of a holonomy band in $X(I, S)/H$ whose holonomy coincides with $\psi|_{(x,c)}$. Lifting this holonomy band to $X(I, S)$, we get a holonomy band joining $\{1\} \times [x, c]$ to $\{\tilde{h}\} \times [\tilde{x}, \tilde{c}]$ for some $\tilde{h} \in H$. Since $S' \supset \varphi(S)$, this holonomy band defines a holonomy band in $X(I', S')$ joining $\{1\} \times [x', c']$ to $\{\varphi(\tilde{h})\} \times [\tilde{x}', \tilde{c}']$ where $\tilde{c}' = f(\tilde{c})$. In particular, $\varphi(\tilde{h}h^{-1})$ fixes the arc $[x', \tilde{c}']$, so $\varphi(\tilde{h}h^{-1}) = 1$. But since $\{\tilde{h}\} \times \{\tilde{x}\}$ is in the same leaf as $\{h\} \times \{x\}$, $\tilde{h}h^{-1}$ fixes the point x in Y . Since φ is one-to-one in restriction to point stabilizers of Y , $h = \tilde{h}$. Since $\{h\} \times [\tilde{x}, \tilde{c}]$ maps into $[x, a_2]$ in Y , we get that $[x, a_2] \cap [x, a_1]$ contains $[x, c]$, a contradiction.

This proves that f is one-to-one. It follows that φ is one-to one since an element of its kernel must fix Y pointwise.

Let's prove that $\varphi(H) \supset H'$, the other inclusion being trivial. The fact that $f(Y) = Y'$ will follow because $H' \circlearrowleft Y'$ is minimal.

Since $H' \circlearrowleft Y$ is indecomposable, H' is generated by the set of elements $g \in H'$ such that $g.I' \cap I'$ is non-degenerate. Let $h' \in H'$ be such that $[x', a'] = h'.I' \cap I'$ is non-degenerate. Repeating the argument above, there exists an arc $[x', b'] \subset [x', a']$ and $\psi' \in \Lambda(I', S')$, $\psi' : (x', b') \rightarrow (\tilde{x}', \tilde{b}')$ where $\tilde{x}' = h^{-1}x'$, $\tilde{b}' = h^{-1}b'$. Let $\psi \in \Lambda(I, S)$ be the corresponding partial isometry. By segment closed property, there exist arcs $[x, c] \subset [x, b]$ and $[\tilde{x}, \tilde{c}] \subset [\tilde{x}, \tilde{b}]$ and a holonomy band in $X(I, S)/H$ joining them in $X(I, S)/H$, whose holonomy coincides with ψ . This band lifts in $X(I, S)$, to a band joining $\{1\} \times [x, c]$ to $\{\tilde{h}\} \times [\tilde{x}, \tilde{c}]$ for some $\tilde{h} \in H$. In $X(I', S')$, this bands joins the corresponding arcs $\{1\} \times [x, c]$ to $\{\tilde{h}\} \times [\tilde{x}, \tilde{c}]$. It follows that in Y , the actions of $\varphi(\tilde{h})$ and h coincide on $[x', c']$. Since arc stabilizers are trivial, $h = \varphi(\tilde{h})$. We conclude that $\varphi(H) = H'$. \square

References

- [Ali] Emina Alibegovic. Makanin-razborov diagrams for limit groups. math.GR/0410198.
- [BF] Mladen Bestvina and Mark Feighn. Outer limits. Preprint.
- [BF91] Mladen Bestvina and Mark Feighn. Bounding the complexity of simplicial group actions on trees. *Invent. Math.*, 103(3):449–469, 1991.
- [BF92] M. Bestvina and M. Feighn. A combination theorem for negatively curved groups. *J. Differential Geom.*, 35(1):85–101, 1992.
- [BF95] Mladen Bestvina and Mark Feighn. Stable actions of groups on real trees. *Invent. Math.*, 121(2):287–321, 1995.
- [BS] Igor Belegradek and Andrzej Szczepanski. Endomorphisms of relatively hyperbolic groups. math.GR/0501321.
- [Chi01] Ian Chiswell. *Introduction to Λ -trees*. World Scientific Publishing Co. Inc., River Edge, NJ, 2001.
- [CL95] Marshall M. Cohen and Martin Lustig. Very small group actions on \mathbb{R} -trees and Dehn twist automorphisms. *Topology*, 34(3):575–617, 1995.
- [CM87] Marc Culler and John W. Morgan. Group actions on \mathbb{R} -trees. *Proc. London Math. Soc.* (3), 55(3):571–604, 1987.

- [Coo05] Group Theory Cooperative. Magnus, computational package for exploring infinite groups, version 4.1.3 beta, June 2005. (G.Baumslag director).
- [Del99] Thomas Delzant. Sur l'accessibilité acylindrique des groupes de présentation finie. *Ann. Inst. Fourier (Grenoble)*, 49(4):1215–1224, 1999.
- [DS] Cornelia Druţu and Mark Sapir. Groups acting on tree-graded spaces and splittings of relatively hyperbolic group. math.GR/0601305.
- [Dun98] M. J. Dunwoody. Folding sequences. In *The Epstein birthday schrift*, pages 139–158 (electronic). *Geom. Topol.*, Coventry, 1998.
- [GLP94] D. Gaboriau, G. Levitt, and F. Paulin. Pseudogroups of isometries of \mathbb{R} and Rips' theorem on free actions on \mathbb{R} -trees. *Israel J. Math.*, 87(1-3):403–428, 1994.
- [GLP95] Damien Gaboriau, Gilbert Levitt, and Frédéric Paulin. Pseudogroups of isometries of \mathbb{R} : reconstruction of free actions on \mathbb{R} -trees. *Ergodic Theory Dynam. Systems*, 15(4):633–652, 1995.
- [Gro05] Daniel Groves. Limit groups for relatively hyperbolic groups. II. Makanin-Razborov diagrams. *Geom. Topol.*, 9:2319–2358 (electronic), 2005.
- [Gui] Vincent Guirardel. Non-homogeneous minimal pseudogroups are almost maximal. In preparation.
- [Gui98a] Vincent Guirardel. *Actions de groupes sur des arbres réels et dynamique dans la frontière de l'outre-espace*. PhD thesis, Université Toulouse III, January 1998.
- [Gui98b] Vincent Guirardel. Approximations of stable actions on \mathbb{R} -trees. *Comment. Math. Helv.*, 73(1):89–121, 1998.
- [Gui04] Vincent Guirardel. Limit groups and groups acting freely on \mathbb{R}^n -trees. *Geom. Topol.*, 8:1427–1470 (electronic), 2004.
- [Gui05] Vincent Guirardel. Cœur et nombre d'intersection pour les actions de groupes sur les arbres. *Ann. Sci. École Norm. Sup. (4)*, 38(6):847–888, 2005.
- [Ima79] Hideki Imanishi. On codimension one foliations defined by closed one-forms with singularities. *J. Math. Kyoto Univ.*, 19(2):285–291, 1979.
- [Lev93] Gilbert Levitt. La dynamique des pseudogroupes de rotations. *Invent. Math.*, 113(3):633–670, 1993.
- [Lev94] Gilbert Levitt. Graphs of actions on \mathbb{R} -trees. *Comment. Math. Helv.*, 69(1):28–38, 1994.
- [LP97] Gilbert Levitt and Frédéric Paulin. Geometric group actions on trees. *Amer. J. Math.*, 119(1):83–102, 1997.
- [Mor88] John W. Morgan. Ergodic theory and free actions of groups on \mathbb{R} -trees. *Invent. Math.*, 94(3):605–622, 1988.
- [MS84] John W. Morgan and Peter B. Shalen. Valuations, trees, and degenerations of hyperbolic structures. I. *Ann. of Math. (2)*, 120(3):401–476, 1984.
- [Pau88] Frédéric Paulin. Topologie de Gromov équivariante, structures hyperboliques et arbres réels. *Invent. Math.*, 94(1):53–80, 1988.
- [RS94] E. Rips and Z. Sela. Structure and rigidity in hyperbolic groups. I. *Geom. Funct. Anal.*, 4(3):337–371, 1994.
- [Sco73] G. P. Scott. Finitely generated 3-manifold groups are finitely presented. *J. London Math. Soc. (2)*, 6:437–440, 1973.
- [Sel] Zlil Sela. Diophantine geometry over groups viii: The elementary theory of a hyperbolic group. <http://www.ma.huji.ac.il/~zlil>.
- [Sel97] Z. Sela. Acylindrical accessibility for groups. *Invent. Math.*, 129(3):527–565, 1997.
- [Sel99] Z. Sela. Endomorphisms of hyperbolic groups. I. The Hopf property. *Topology*, 38(2):301–321, 1999.

- [Sel01a] Zlil Sela. Diophantine geometry over groups. I. Makanin-Razborov diagrams. *Publ. Math. Inst. Hautes Études Sci.*, 93:31–105, 2001.
- [Sel01b] Zlil Sela. Diophantine geometry over groups VI: The elementary theory of a free group. <http://www.ma.huji.ac.il/~zlil>, 2001.
- [Ser77] Jean-Pierre Serre. *Arbres, amalgames, SL_2* . Société Mathématique de France, Paris, 1977. Avec un sommaire anglais, Rédigé avec la collaboration de Hyman Bass, Astérisque, No. 46.
- [Sha91] Peter B. Shalen. Dendrology and its applications. In *Group theory from a geometrical viewpoint (Trieste, 1990)*, pages 543–616. World Sci. Publishing, River Edge, NJ, 1991.
- [Sta83] John R. Stallings. Topology of finite graphs. *Invent. Math.*, 71(3):551–565, 1983.
- [Swa] Gadde Swarup. Delzant’s variation on Scott complexity. arXiv:math.GR/0401308.

Vincent Guirardel
 Laboratoire Émile Picard, UMR 5580
 UFR MIG,
 Université Paul Sabatier
 31062 Toulouse cedex 9.
 France.
e-mail:guirardel@math.ups-tlse.fr