Geometric structure in the representation theory of p-adic groups

Anne-Marie Aubert, Paul Baum and Roger Plymen

Abstract

We conjecture the existence of a simple geometric structure underlying questions of reducibility of parabolically induced representations of reductive *p*-adic groups.

1 Introduction

In the representation theory of reductive p-adic groups, the issue of reducibility of induced representations is an issue of great intricacy: see, for example, the classic article by Bernstein-Zelevinsky [4] on GL(n) and the more recent article by Muić [7] on G_2 . It is our contention, expressed as a conjecture, that there exists a simple geometric structure underlying this intricate theory.

For the moment, our conjecture is *local*, in that it applies only to finite places. To explain our conjecture, we need to refine the usual concept of quotient space.

2 The extended quotient and the reduced quotient

Let Γ be a finite group and let X be a complex affine algebraic variety. Assume that Γ is acting on X as automorphisms of the affine algebraic variety X.

Definition 1. The quotient variety X/Γ is obtained by collapsing each orbit of X to a point.

If J is a finite group, c(J) denotes the set of conjugacy classes of J. If $x \in X$, Γ_x denotes the isotropy group of x:

$$\Gamma_x = \{ \gamma \in \Gamma \, : \, \gamma x = x \}.$$

The extended quotient, denoted $X//\Gamma$ is obtained from X by replacing each orbit $\{\gamma x : \gamma \in \Gamma\}$ by $c(\Gamma_x)$. To construct $X//\Gamma$, we proceed as follows. Let

$$\widetilde{X} := \{(\gamma, x) \in \Gamma \times X : \gamma x = x\}.$$

The group Γ acts on \widetilde{X} by:

$$\alpha(\gamma, x) = (\alpha \gamma \alpha^{-1}, \alpha x) \quad \text{with } (\gamma, x) \in \widetilde{X}, \, \alpha \in \Gamma.$$

Definition 2. The extended quotient, denoted $X//\Gamma$, is defined as

$$X/\!/\Gamma := \widetilde{X}/\Gamma$$

i.e. $X//\Gamma$ is the ordinary quotient (as in definition 1) for the action of Γ on \widetilde{X} .

The projection

$$\Gamma \times X \longrightarrow X, \quad (\gamma, x) \mapsto x$$

gives a map

$$\pi \colon X //\Gamma \longrightarrow X /\Gamma.$$

If p is an orbit in X, i.e. $p = \{\gamma x : \gamma \in \Gamma\}$, then the pre-image in $X//\Gamma$ of p is $c(\Gamma_x)$. Thus, in forming $X//\Gamma$, each orbit $\{\gamma x : \gamma \in \Gamma\}$ has been replaced by $c(\Gamma_x)$.

Definition 3. The map $\pi: X//\Gamma \longrightarrow X/\Gamma$ is the projection of the extended quotient on the ordinary quotient.

Lemma 1. The projection $\pi: X//\Gamma \to X/\Gamma$ is a finite morphism of algebraic varieties.

Let $X^{\gamma} := \{x \in X : \gamma x = x\}$ and denote by $Z(\gamma)$ the Γ -centralizer of γ . We have

$$X//\Gamma = \bigsqcup_{\gamma} X^{\gamma}/\mathbf{Z}(\gamma)$$

where one γ is chosen in each Γ -conjugacy class. Let $e \in \Gamma$ denote the identity element. Since $X^e/\mathbb{Z}(e)$ is the ordinary quotient X/Γ , we have $X/\Gamma \subset X//\Gamma$.

Definition 4. The reduced quotient is defined as

$$(X/\Gamma)_{\rho} := \pi(X//\Gamma - X/\Gamma).$$

Lemma 2. The reduced quotient is an algebraic variety.

Lemma 3. The extended quotient is multiplicative: if $\Gamma_1 \times X_1 \to X_1$ and $\Gamma_2 \times X_2 \to X_2$ are as above, then we have

$$(X_1 \times X_2) / (\Gamma_1 \times \Gamma_2) = (X_1 / / \Gamma_1) \times (X_2 / / \Gamma_2).$$

3 Application to the representation theory of *p*-adic groups

Let F be a local nonarchimedean field, let G be the group of F-rational points in a connected reductive algebraic group defined over F, and let Irr(G) be the set of irreducible smooth representations of G. We recall the data in the Bernstein programme: $\mathfrak{s} \in \mathfrak{B}(G)$, $\mathfrak{s} = [M, \sigma]_G$ is an inertial class in G(with M a Levi subgroup of G and σ a supercuspidal representation of M), $D^{\mathfrak{s}}$ is the $\Psi(M)$ -orbit of σ in Irr(M), with $\Psi(M)$ the group of unramified characters of M, $W^{\mathfrak{s}} = \{w \in N_G(M)/M : w \cdot \mathfrak{s} = \mathfrak{s}\}$ and $D^{\mathfrak{s}}/W^{\mathfrak{s}}$ is the quotient variety, a component of the Bernstein variety.

We will fix a point $\mathfrak{s} \in \mathfrak{B}(G)$ and write $D = D^{\mathfrak{s}}, W = W^{\mathfrak{s}}$. Let $\operatorname{Irr}(G)^{\mathfrak{s}}$ denote the \mathfrak{s} -component of $\operatorname{Irr}(G)$ in the Bernstein decomposition of $\operatorname{Irr}(G)$. We will equip the quotient variety D/W with the Zariski topology, and $\operatorname{Irr}(G)^{\mathfrak{s}}$ with the Jacobson topology coming from $\operatorname{Prim} \mathcal{H}(G)^{\mathfrak{s}}$. We note that irreducibility is an *open* condition, and so $(D/W)_{\mathrm{red}}$, the set of reducible points in D/W, is a sub-variety of D/W. Let q denote the cardinality of the residue field of F. Let $\mathcal{H}(G)$ denote the Hecke algebra of G, let $\mathcal{H}(G)^{\mathfrak{s}}$ denote the ideal of $\mathcal{H}(G)$ corresponding to \mathfrak{s} in the Bernstein decomposition of $\mathcal{H}(G)$, and let HP_* denote periodic cyclic homology.

Conjecture 1. There is an isomorphism

$$\operatorname{HP}_*(\mathcal{H}(G)^{\mathfrak{s}}) \cong \operatorname{H}^*(\mathcal{D}//\mathcal{W})$$

and a continuous bijection

$$\mu: D//W \to \operatorname{Irr}(G)^{\mathfrak{s}}$$

such that:

(1) There is an algebraic family $\pi_t \colon D//W \to D/W$ of finite morphisms of algebraic varieties, with $t \in \mathbb{C}^{\times}$, such that

$$\pi_1 = \pi, \qquad \pi_{\sqrt{q}} = (inf.ch.) \circ \mu.$$

If we let \mathfrak{X}_t be the image of π_t restricted to D/W - D/W then \mathfrak{X}_t is a flat family of algebraic varieties such that

$$\mathfrak{X}_1 = (D/W)_{\rho}, \qquad \mathfrak{X}_{\sqrt{q}} = (D/W)_{red}.$$

(2) For each irreducible component $\mathbf{c} \subset D//W$ there is a cocharacter $h_{\mathbf{c}} \colon \mathbb{C}^{\times} \to D$ such that

$$\pi_t(x) = \pi(h_{\mathbf{c}}(t) \cdot x)$$

for all $x \in \mathbf{c}$. If $\mathbf{c} = D/W$ then $h_{\mathbf{c}} = 1$.

Theorem 1. The conjecture is true for G = SL(2). If $\mathfrak{s} = [T, 1]_G$ then \mathfrak{X}_t is the 0-dimensional variety given by the polynomial $(x + 1)(x - t^2) = 0$.

4 The general linear group

Theorem 2. The conjecture is true for GL(n).

Proof. The proof uses Langlands parameters, together with some careful combinatorics. In effect, the *L*-parameters encode the extended quotient for GL(n). Let G = GL(n) = GL(n, F), n = mr, τ be an irreducible supercuspidal representation of GL(m, F),

$$\mathfrak{s} = [M, \sigma]_G = [\operatorname{GL}(m)^r, \tau^{\otimes r}]_G.$$

We have

$$D = D^{\mathfrak{s}} = (\mathbb{C}^{\times})^r, \quad W = W^{\mathfrak{s}} = S_r.$$

Let W_F be the Weil group of F, and let $\mathcal{L}_F = W_F \times \mathrm{SU}(2)$. Let $\mathrm{Hom}_{\mathrm{ss}}(\mathcal{L}_F, \mathrm{GL}(n, \mathbb{C}))$ denote the set of equivalence classes of Frobenius-semisimple smooth homomorphisms from \mathcal{L}_F to $\mathrm{GL}(n, \mathbb{C})$.

For each $n \ge 1$ we have the local Langlands correspondence [6]

$$\operatorname{rec}_F \colon \operatorname{Irr}(\operatorname{GL}(n, F)) \to \operatorname{Hom}_{\operatorname{ss}}(\mathcal{L}_F, \operatorname{GL}(n, \mathbb{C})).$$

We shall write

$$\Phi(G) = \operatorname{Hom}_{ss}(\mathcal{L}_F, \operatorname{GL}(n, \mathbb{C}))$$

We will denote by p the following partition of r:

$$a_1 + \dots + a_1 + \dots + a_l + \dots + a_l = r_1 a_1 + \dots + r_l a_l = r$$

where a_j is repeated r_j times. Let $\gamma \in S_r$ be the corresponding product of $r_1 + \cdots + r_l$ cycles. The fixed set D^{γ} is a complex torus of dimension $r_1 + \cdots + r_l$.

We recall that τ is supercuspidal representation of $\operatorname{GL}(m)$. Now let $\operatorname{rec}_F(\tau) = \eta \in \operatorname{Irr}_m(W_F)$. Denote by R(j) the *j*-dimensional irreducible complex representation of SU(2). Corresponding to the partition p we have the *L*-parameter

$$\phi = \eta \otimes R(a_1) \oplus \cdots \oplus \eta \otimes R(a_1) \oplus \cdots \oplus \eta \otimes R(a_l) \oplus \cdots \oplus \eta \otimes R(a_l)$$

where $\eta \otimes R(a_1)$ is repeated r_1 times, ..., $\eta \otimes R(a_l)$ is repeated r_l times.

Let $\Psi(W_F)$ denote the group of unramified quasicharacters of the Weil group W_F , and consider the complex torus $\Psi(W_F)^{r_1+\cdots+r_l}$. The *orbit* of ϕ in $\Phi(G)$, via the action of this complex torus, is

$$\mathcal{O}(\phi) = \operatorname{Sym}^{r_1} \mathbb{C}^{\times} \times \cdots \times \operatorname{Sym}^{r_l} \mathbb{C}^{\times} \subset \Phi(G).$$

Let $\psi_j \in \Psi(W_F)$ with $1 \leq j \leq r_1 + \cdots + r_l$. We will map each *L*-parameter in the orbit $\mathcal{O}(\phi)$ as follows:

$$\psi_1 \otimes \eta \otimes R(a_1) \oplus \cdots \oplus \psi_{r_1 + \dots + r_l} \otimes \eta \otimes R(a_l) \mapsto (\psi_1(\varpi), \dots, \psi_{r_1 + \dots + r_l}(\varpi)) \in D^{\gamma}$$

where ϖ is a uniformizer in F. This induces a *bijection*

$$\mathcal{O}(\phi) \cong D^{\gamma}/\mathbf{Z}(\gamma).$$

Let $\Phi(G)^{\mathfrak{s}}$ denote the \mathfrak{s} -component of $\Phi(G)$ in the Bernstein decomposition of $\Phi(G)$, so that

$$\Phi(G)^{\mathfrak{s}} = \operatorname{rec}_F(\operatorname{Irr}(G)^{\mathfrak{s}}).$$

We now take the disjoint union of the permutations γ , one chosen in each S_r -conjugacy class. This creates a *canonical* bijection

$$\Phi(G)^{\mathfrak{s}} \cong D/\!/W.$$

The reduced quotient $(D/W)_{\rho}$ is the hypersurface \mathfrak{X}_1 given by the single equation $\prod_{i\neq j}(z_i-z_j)=0$. The variety $(D/W)_{red}$ is the variety \mathfrak{X}_q given by the single equation $\prod_{i\neq j}(z_i-qz_j)=0$, according to a classical theorem [4, Theorem 4.2], [8]. The polynomial equation $\prod_{i\neq j}(z_i-tz_j)=0$ determines a flat family \mathfrak{X}_t of hypersurfaces. The hypersurface \mathfrak{X}_1 is the *flat limit* of the family \mathfrak{X}_t as $t \to 1$, as in [5, p.77].

Let **c** denote the irreducible variety $D^{\gamma}/Z(\gamma)$. The cocharacter $h_{\mathbf{c}}$ is given by

 $h_{\mathbf{c}}: t \mapsto (t^{a_1-1}, \dots, t^{1-a_1}, \dots, t^{a_r-1}, \dots, t^{1-a_r}) \in D.$

Finally, we have to use the multiplicativity of the extended quotient. \Box

5 The exceptional group G_2

We have chosen the exceptional group G_2 as an awkward example, requiring many delicate calculations, see [2]. Let $\mathfrak{s} = [T, \chi \otimes \chi]_G$ where $T \simeq F^{\times} \times F^{\times}$ is a maximal *F*-split torus of $G = G_2$ and χ is a ramified quadratic character of F^{\times} . Let $\{\alpha, \beta\}$ be a basis of a set of roots of *G* with α short and β long. The group $W^{\mathfrak{s}} \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ is generated by the fundamental reflections s_{α} and $s_{3\alpha+2\beta}$, and we have $D = D^{\mathfrak{s}} = \{\psi_1 \chi \otimes \psi_2 \chi : \psi_1, \psi_2 \in \Psi(F^{\times})\}$. We obtain

$$D^{\gamma}/\mathbf{Z}(\gamma) = \begin{cases} \{ \{(\lambda,\lambda), (\lambda^{-1},\lambda^{-1})\} : \lambda \in \mathbb{C}^{\times} \}, & \text{if } \gamma = s_{\alpha}, \\ \{ \{(\lambda,\lambda^{-1}), (\lambda^{-1},\lambda)\} : \lambda \in \mathbb{C}^{\times} \}, & \text{if } \gamma = s_{3\alpha+2\beta}, \\ (1,1) \sqcup (-1,-1) \sqcup (1,-1), & \text{if } \gamma = s_{\alpha}s_{3\alpha+2\beta}. \end{cases}$$

Denote by \mathfrak{C}_1 the line x - y = 0 and by \mathfrak{C}_2 the hyperbola xy - 1 = 0. Setting $\mathrm{pt}_1 := (1, 1), \, \mathrm{pt}_2 := (-1, -1), \, \mathrm{and} \, \mathrm{pt}_3 := (1, -1), \, \mathrm{we \ obtain}$

$$D/W = D/W \sqcup \mathfrak{C}_1 \sqcup \mathfrak{C}_2 \sqcup \mathrm{pt}_1 \sqcup \mathrm{pt}_2 \sqcup \mathrm{pt}_3, \qquad (D/W)_{\rho} = \mathfrak{C}_1 \cup \mathfrak{C}_2 \cup \mathrm{pt}_3.$$

The cocharacter $h_{\mathbf{c}}: \mathbb{C}^{\times} \to D$ is as follows:

$$t \mapsto (t, t^{-1})$$
 if $\mathbf{c} = \mathfrak{C}_1, (t^{-1}, t^{-1})$ if $\mathbf{c} = \mathfrak{C}_2, (1, t^{-2})$ if $\mathbf{c} = \mathrm{pt}_1$ or $\mathrm{pt}_2, (1, 1)$ if $\mathbf{c} = \mathrm{pt}_3$.

The variety \mathfrak{X}_t is the union of the line $x - t^2 y = 0$, the hyperbola $xy - t^{-2} = 0$ and the point pt₃. This is a flat family. The 2 curves admit 2 intersection points: $(1, t^{-2})$ and $(-1, -t^{-2})$. Now let $t = \sqrt{q}$. At each of the intersection points, the corresponding parabolically induced representation admits 4 irreducible inequivalent constituents. As for the point pt₃: the corresponding parabolically induced representation admits 2 irreducible inequivalent *tempered* constituents, see [7].

Theorem 3. The conjecture is true for the point $\mathfrak{s} = [T, \chi \otimes \chi]_G$.

Acknowledgements

The second author was partially supported by an NSF grant.

References

- A.-M. Aubert, P. Baum, R.J. Plymen, The Hecke algebra of a reductive p-adic group: a geometric conjecture, in: C. Consani, M. Marcolli (Eds.), Noncommutative geometry and number theory, Aspects of Mathematics E37, Vieweg Verlag 2006.
- [2] A.-M. Aubert, P. Baum, R.J. Plymen, Geometric structure in the representation theory of *p*-adic groups, preprint 2006.
- [3] J. Bernstein, Representations of *p*-adic groups, Notes by K.E. Rumelhart, Harvard University 1992.

- [4] I.N. Bernstein, A.V. Zelevinsky, Induced representations of reductive p-adic groups I, Ann. Sci. E.N.S. 4 (1977) 441–472.
- [5] D. Eisenbud, J. Harris, The geometry of schemes, Springer, 2001.
- [6] M. Harris, R. Taylor, The geometry and cohomology of some simple Shimura varieties, Annals of Math. Study 151, Princeton, 2001.
- [7] G. Muić, On the unitary dual of G_2 , Duke Math. J. 90 (1997) 465–493.
- [8] A.V. Zelevinsky. Induced representations of reductive *p*-adic groups II, Ann. Sci. Ec. Norm. Sup. 13 (1980) 154–210.

Anne-Marie Aubert, Institut de Mathématiques de Jussieu, U.M.R. 7586 du C.N.R.S., Paris, France. Email: aubert@math.jussieu.fr

Paul Baum, Pennsylvania State University, Mathematics Department, University Park, PA 16802, USA. Email: baum@math.psu.edu

Roger Plymen, School of Mathematics, Manchester University, Manchester M13 9PL, England. Email: plymen@manchester.ac.uk