# arXiv:math/0607440v1 [math.DS] 18 Jul 2006

# A NOTE ON SENSITIVITY OF SEMIGROUP ACTIONS

EDUARD KONTOROVICH AND MICHAEL MEGRELISHVILI

ABSTRACT. It is well known that for a transitive dynamical system (X, f) sensitivity to initial conditions follows from the assumption that the periodic points are dense. This was done by several authors: Banks, Brooks, Cairns, Davis and Stacey [2], Silverman [8] and Glasner and Weiss [6]. In the latter article Glasner and Weiss established a stronger result (for compact metric systems) which implies that a transitive non-minimal compact metric system (X, f) with dense set of almost periodic points is sensitive. This is true also for group actions as was proved in the book of Glasner [4].

Our aim is to generalize these results in the frame of a unified approach for a wide class of topological semigroup actions including one-parameter semigroup actions on Polish spaces.

# 1. INTRODUCTION

First we recall some well known closely related results regarding sensitivity of dynamical systems.

- **Theorem 1.1.** (1) (Banks, Brooks, Cairns, Davis and Stacey [2]; Silverman [8]<sup>1</sup>) Let X be an infinite metric space and  $f : X \to X$  be continuous. If f is topologically transitive and has dense periodic points then f has sensitive dependence on initial conditions.
  - (2) (Glasner and Weiss [6, Theorem 1.3]); see also Akin, Auslander and Berg [1]) Let X be a compact metric space and the system (X, f) is an M-system and not minimal. Then (X, f) is sensitive.
  - (3) (Glasner [4, Theorem 1.41]) Let X be a compact metric space. An almost equicontinuous M-system (G, X), where G is a group, is minimal and equicontinuous. Thus M-system which is not minimal equicontinuous is sensitive.

Topological transitivity of  $f: X \to X$  as usual, means that for every pair U and V of nonempty subsets of X there exists n > 0 such that  $f^n(U) \cap V$  is nonempty. Analogously can be defined general semigroup action version (see Definition 3.1.1).

If X is a compact metric space then (2) easily covers (1). In order to explain this recall that *M*-system means that the set of almost periodic points is dense in X (Bronstein condition) and, in addition, the system is topologically transitive. A very particular case of Bronstein condition is that X has dense periodic points (the so-called *P*-systems). If now X is infinite then it cannot be minimal.

Our aim is to provide a unified and generalized approach. We show that (2) and (3) remain true for a large class of *C*-semigroups (which contains: cascades, topological groups and one-parameter semigroups) and M-systems (see Definitions 2.1 and 4.1).

Date: July 18, 2006.

<sup>&</sup>lt;sup>1</sup>under different but very close assumptions

Our approach allows us also to drop the compactness assumption of X dealing with Polish phase spaces. A topological space is *Polish* means that it admits a separable complete metric.

We cover also (1) in the case of Polish phase spaces.

Here we formulate one of the main results (Theorem 5.7) of the present article.

**Main result:** Let (S, X) be a dynamical system where X is a Polish space and S is a C-semigroup. If X is an M-system which is not minimal or not equicontinuous. Then X is sensitive.

Acknowledgment. We thank E. Akin for helpful suggestions and S. Kolyada and L. Snoha for sending us their article [7].

# 2. Preliminaries

A dynamical system in the present article is a triple  $(S, X, \pi)$ , where S is a topological semigroup, X at least is a Hausdorff space and

$$\pi: S \times X \to X, \ (s, x) \mapsto sx$$

is a continuous action on X. Thus,  $s_1(s_2x) = (s_1s_2)x$  holds for every triple  $(s_1, s_2, x)$ in  $S \times S \times X$ . Sometimes we write the dynamical system as a pair (S, X) or even as X, when S is understood. The *orbit* of x is the set  $Sx := \{sx : s \in S\}$ . By  $\overline{A}$ we will denote the closure of a subset  $A \subset X$ . If (S, X) is a system and Y a closed S-invariant subset, then we say that (S, Y), the restricted action, is a *subsystem* of (S, X). For  $U \subset X$  and  $s \in S$  denote

$$s^{-1}U := \{ x \in X : sx \in U \}.$$

If  $S = \{f^n\}_{n \in \mathbb{N}}$  (with  $\mathbb{N} := \{1, 2, \dots\}$ ) and  $f : X \to X$  is a continuous function, then the classical dynamical system (S, X) is called a *cascade*. Notation: (X, f).

# **Definition 2.1.** Let S be a topological semigroup.

(1) We say that S is a (left) F-semigroup if for every  $s_0 \in S$  the subset  $S \setminus Ss_0$  is finite.

(2) We say that S is a C-semigroup if  $S \setminus Ss_0$  is relatively compact (that is, its closure is compact in S).

*Example 2.2.* (1) Standard one-parameter semigroup  $S := ([0, \infty), +)$  is a C-semigroup.

- (2) Every cyclic "positive" semigroup  $M := \{s^n : n \in \mathbb{N}\}$  is an F-semigroup. In particular, for every cascade (X, f) the corresponding semigroup  $S = \{f^n\}_{n \in \mathbb{N}}$  is an F-semigroup (and hence also a C-semigroup).
- (3) Every topological group is of course an F-semigroup.
- (4) Every compact semigroup is a C-semigroup.

**Definition 2.3.** Let (S, X) be a dynamical system where (X, d) is a metric space.

(1) A subset A of S acts equicontinuously at  $x_0 \in X$  if for every  $\epsilon > 0$  there exists  $\delta > 0$  such that  $d(x_0, x) < \delta$  implies  $d(ax_0, ax) < \epsilon$  for every  $a \in A$ .

- (2) A point  $x_0 \in X$  is called an *equicontinuity point* (notation:  $x_0 \in Eq(X)$ ) if A := S acts equicontinuously at  $x_0$ . If Eq(X) = X then (S, X) is *equicontinuously* at *x*<sub>0</sub>.
- (3) (S, X) is called *almost equicontinuous* (see [1, 4]) if the subset Eq(X) of equicontinuity points is a dense subset of X.

**Lemma 2.4.** Let (S, X) be a dynamical system where (X, d) is a metric space. Let  $A \subset S$  be a relatively compact subset. Then A acts equicontinuously on (X, d).

# 3. TRANSITIVITY CONDITIONS OF SEMIGROUP ACTIONS

**Definition 3.1.** The dynamical system (S, X) is called:

- (1) topologically transitive (in short: TT) if for every pair (U, V) of non-empty open sets U, V in X there exists  $s \in S$  with  $U \cap sV \neq \emptyset$ . Since  $s(s^{-1}U \cap V) = U \cap sV$ , it is equivalent to say that  $s^{-1}U \cap V \neq \emptyset$ .
- (2) point transitive (PT) if there exists a point x with dense orbit. Such a point is called *transitive point*. Notation:  $x_0 \in Trans(X)$ .
- (3) densely point transitive (DPT) if there exists a dense set  $Y \subset X$  of transitive points.

Of course always (DPT) implies (PT). In general, (TT) and (PT) are independent properties. For a detailed discussion of transitivity conditions (for cascades) see a review paper by Kolyada and Snoha [7].

As usual, X is *perfect* means that X is a space without isolated points. Assertions (1) and (2) in the following proposition are very close to Silverman's observation [8, Proposition 1.1] (for cascades).

# **Proposition 3.2.** (1) If X is a perfect topological space and S is an F-semigroup, then (PT) implies (TT).

- (2) If X is a Polish space then every (TT) system (S, X) is (DPT) (and hence also (PT)).
- (3) Every (DPT) system (S, X) is (TT).

Proof. (1) Let x be a transitive point with orbit Sx. Now, let U and V be nonempty open subsets of X. There exists  $s_1 \in S$  such that  $s_1x \in V$ . The subset  $S \setminus Ss_1$  is finite because S is an almost F-group. Since X is perfect, removing the finite subset  $(S \setminus Ss_1)x$  from the dense subset Sx we get again a dense subset. Therefore,  $Ss_1x$  is a dense subset of X. Then there exists  $s_2 \in S$  such that  $s_2s_1x \in U$ . Thus  $s_2^{-1}U \cap V \neq \emptyset$ . By Definition 3.1.1 this means that (S, X) is a (TT) dynamical system.

(2) If (S, X) is topologically transitive, then  $S^{-1}U$  is a dense subset of X for every open set U. We know that X is Polish. Then there exists a countable open base  $\mathcal{B}$ of the given topology. By the Baire theorem,  $\bigcap \{S^{-1}U : U \in \mathcal{B}\}$  is dense in X and every point of this set is a transitive point of the dynamical system X.

(3) Let U and V be nonempty open subsets in X. Since the set Y of point transitive points is dense in X, it intersects V. Therefore, we can choose a transitive point  $y \in V$ . Now by the transitivity of y there exists  $s \in S$  such that sy belongs to U. Hence, sy is a common point of U and sV.

**Lemma 3.3.** Let (X, d) be a metric S-system which is (TT). Then  $Eq(X) \subset Trans(X)$ .

Proof. Let  $x_0 \in Eq(X)$  and  $y \in X$ . We have to show that the orbit  $Sx_0$  intersects the  $\varepsilon$ -neighborhood  $B_{\varepsilon}(y) := \{x \in X : d(x, y) < \varepsilon\}$  of y for every given  $\varepsilon > 0$ . Since  $x_0 \in Eq(X)$  there exists a neighborhood U of  $x_0$  such that  $d(sx_0, sx) < \frac{\varepsilon}{2}$  for every  $(s, x) \in S \times U$ . Since X is (TT) we can choose  $s_0 \in S$  such that  $s_0U \cap B_{\frac{\varepsilon}{2}}(y) \neq \emptyset$ . This means that  $d(s_0x, y) < \frac{\varepsilon}{2}$  for some  $x \in U$ . Then  $d(s_0x_0, y) < \varepsilon$ .

# 4. MINIMALITY CONDITIONS

The following definitions are standard for compact X.

**Definition 4.1.** Let X be a not necessarily compact S-dynamical system.

- (1) X is called *minimal*, if  $\overline{Sx} = X$  for every  $x \in X$ . In other words, all points of X are transitive points.
- (2) A point x is called *minimal* if the subsystem  $\overline{Sx}$  is minimal.
- (3) A point x is called *almost periodic* if the subsystem  $\overline{Sx}$  is minimal and compact.
- (4) If the set of almost periodic points is dense in X, we say that (S, X) satisfies the *Bronstein condition*. If, in addition, the system (S, X) is (TT), we say that it is an *M*-system.
- (5) A point  $x \in X$  is a *periodic point*, if Sx is finite. If (S,X) is a (TT) dynamical system and the set of periodic points is dense in X, then we say that it is a *P*-system, [6].

If X is compact then a point in X is minimal iff it is almost periodic. Every periodic point is of course almost periodic. Therefore it is also obvious that every P-system is an M-system.

For a system (S, X) and a subset  $B \subset X$ , we use the following notation

$$N(x,B) = \{s \in S : sx \in B\}.$$

The following definition is also standard.

**Definition 4.2.** A subset  $P \subset S$  is (left) *syndetic*, if there exists a finite set  $F \subset S$  such that  $F^{-1}P = S$ .

The following lemma is a slightly generalized version of a well known criteria for almost periodic points (cf. Definition 4.1.3) in compact dynamical systems. In particular, it is valid for every semigroup S.

**Lemma 4.3.** Let (S, X) be a (not necessarily compact) dynamical system and  $x_0 \in X$ . Consider the following conditions:

- (1)  $x_0$  is an almost periodic point.
- (2) For every open neighborhood V of  $x_0$  in X there exists a finite set  $F \subset S$  such that  $F^{-1}V \supseteq Y := \overline{Sx_0}$ .
- (3) For every neighborhood V of  $x_0$  in X the set  $N(x_0, V)$  is syndetic.
- (4)  $x_0$  is a minimal point (i.e., the subsystem  $\overline{Sx_0}$  is minimal).

Then  $(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4)$ .

If X is compact then all four conditions are equivalent.

*Proof.* (1)  $\Rightarrow$  (2) : Suppose that (Y, S) is minimal and compact. Then for every open neighborhood V of  $x_0$  in X and for every  $y \in Y$  there exists  $s \in S$  such that  $sy \in V$ . Equivalently,  $y \in s^{-1}V$ . Therefore,  $\bigcup_{s \in S} s^{-1}V \supseteq Y$ . By compactness of Y we can choose a finite set  $F \subseteq S$  such that  $F^{-1}V \supseteq Y$ .

 $(2) \Rightarrow (3)$ : It suffices to show that  $F^{-1}N(x_0, V) = S$ , where F is a subset of S defined in (2). Assume otherwise, so that there exists  $s \in S$  such that  $s \notin F^{-1}N(x_0, V)$ . Then  $sx_0 \notin F^{-1}V$ . On the other hand clearly,  $sx_0 \in Y$ , contrary to our condition that  $F^{-1}V \supseteq Y$ .

 $(3) \Rightarrow (4) : Y = \overline{Sx_0}$  is non-empty, closed and invariant. It remains to show that if  $y \in Y$  then  $x_0 \in \overline{Sy}$ . Assume otherwise, so that  $x_0 \notin \overline{Sy}$ . Choose an open neighborhood V of  $x_0$  in X, such that  $\overline{V} \cap \overline{Sy} = \emptyset$ . By our assumption the set  $N(x_0, V)$  is syndetic. Therefore there is a finite set  $F := \{s_1, ..., s_n\}$  so that for each  $s \in S$  some  $s_i s x_0 \in V$ . That is each  $s x_0$  belongs to  $F^{-1}V = \bigcup_{i=1}^n s_i^{-1}V$  for every  $s \in S$ . Hence,  $Sx_0 \subseteq \bigcup_{i=1}^n s_i^{-1}V$ . Then

$$y \in \overline{Sx_0} \subset \overline{\bigcup_{i=1}^n s_i^{-1}V} = \bigcup_{i=1}^n \overline{s_i^{-1}V} \subset \bigcup_{i=1}^n s_i^{-1}\overline{V}.$$

But then  $Sy \cap \overline{V} \neq \emptyset$  contrary to our assumption.

If X is compact then by Definition 4.1 it follows that  $(4) \Rightarrow (1)$ .

### 5. Sensitivity and other conditions

**Proposition 5.1.** Let S be an C-semigroup. Assume that (X, d) is a point transitive (PT) S-system such that  $Eq(X) \neq \emptyset$ . Then every transitive point is an equicontinuity point. That is,  $Trans(X) \subset Eq(X)$  holds.

*Proof.* Let y be a transitive point and  $x \in Eq(X)$  be an equicontinuity point. We have to show that  $y \in Eq(X)$ . For a given  $\varepsilon > 0$  there exists a neighborhood O(x) of x such that

$$d(sx'', sx') < \epsilon \qquad \forall \ s \in S \qquad \forall \ x', x'' \in O(x).$$

Since y is a transitive point then there exists  $s_0 \in S$  such that  $s_0 y \in O(x)$ . Then  $O(y) := s_0^{-1}O(x)$  is a neighborhood of y. We have

$$d(ss_0y', ss_0y'') < \epsilon \qquad \forall \ s \in S \qquad \forall \ y', y'' \in O(y).$$

Since S is a C-semigroup the subset  $M := \overline{S \setminus Ss_0}$  is compact. Hence by Lemma 2.4 it acts equicontinuously on X. We can choose a neighborhood U(y) of y such that

$$d(ty', ty'') < \epsilon \qquad \forall \ t \in M \qquad \forall \ y', y'' \in U(y).$$

Then  $V := O(y) \cap U(y)$  is a neighborhood of y. Since  $S = M \cup Ss_0$  we obtain that  $d(sy', sy'') < \epsilon$  for every  $s \in S$  and  $y', y'' \in V$ . This proves that  $y \in Eq(X)$ .  $\Box$ 

**Proposition 5.2.** Let S be an C-semigroup. Assume that (X, d) is a metric S-system which is minimal and  $Eq(X) \neq \emptyset$ . Then X is equicontinuous.

*Proof.* If (S, X) is a minimal system then Trans(X) = X. Then if  $Eq(X) \neq \emptyset$  every point is an equicontinuity point by Proposition 5.1. Thus, Eq(X) = X.

**Proposition 5.3.** Let S be an C-semigroup. Assume that (X,d) is a Polish (TT) S-system. Then X is almost equicontinuous if and only if  $Eq(X) \neq \emptyset$ .

*Proof.* X is (DPT) by Proposition 3.2.2. That is, Trans(X) is dense in X. Assuming that  $Eq(X) \neq \emptyset$  we obtain by Proposition 5.1 that  $Trans(X) \subset Eq(X)$ . It follows that Eq(X) is also dense in X. Thus, X is almost equicontinuous. This proves "if" part. The remaining direction is trivial.

The following natural definition plays a fundamental role in many investigations about chaotic systems. The present form is a generalized version of existing definitions for cascades (see also [6, 5]).

**Definition 5.4.** (sensitive dependence on initial conditions) A metric S-system (X, d) is sensitive if it satisfies the following condition: there exists a (sensitivity constant) c > 0 such that for all  $x \in X$  and all  $\delta > 0$  there are some  $y \in B_{\delta}(x)$  and  $s \in S$  with d(sx, sy) > c.

We say that (S, X) is non-sensitive otherwise.

**Proposition 5.5.** Let S be an C-semigroup. Assume that (X,d) is a (TT) Polish S-system. Then the system is almost equicontinuous if and only if it is non-sensitive.

*Proof.* Clearly an almost equicontinuous system is always non-sensitive.

Conversely, the non-sensitivity means that for every  $n \in \mathbb{N}$  there exists a nonempty open subset  $V_n \subset X$  such that

$$diam(sV_n) < \frac{1}{n} \quad \forall \ (s,n) \in S \times \mathbb{N}.$$

Define

$$U_n := S^{-1} V_n \qquad R := \bigcap_{n \in \mathbb{N}} U_n.$$

Then every  $U_n$  is open. Moreover, since X is (TT), for every nonempty open subset  $O \subset X$  there exists  $s \in S$  such that  $O \cap s^{-1}U_n \neq \emptyset$ . This means that every  $U_n$  is dense in X. Consequently, by Baire theorem (making use that X is Polish), R is also dense. It is enough now to show that  $R \subset Eq(X)$ . Suppose  $x \in R$  and  $\epsilon > 0$ . Choose n so that  $\frac{1}{n} < \epsilon$ , then  $x \in U_n$  implies the existence of  $s_0 \in S$  such that  $s_0 x \in V_n$ . Put  $V = s_0^{-1}V_{\frac{1}{n}}$ . Therefore for  $y \in V$  and every  $s := s's_0 \in Ss_0$  we get

$$d(sx, sy) = d(s's_0x, s's_0y) < \frac{1}{n} < \epsilon.$$

But  $S \setminus Ss_0$  is relatively precompact set in S because S is a C-semigroup. Then by Lemma 2.4 the set  $S \setminus Ss_0$  acts on (X, d) equicontinuously. We have an open neighborhood O of x such that for all  $y \in O$  and for every  $s \in \overline{S \setminus Ss_0}$  holds  $d(sx, sy) < \epsilon$ . Define an open neighborhood  $M := O \cap V$  of x. Then  $d(sx, sy) < \epsilon$  for every  $s \in S$ and all  $y \in M$ . Thus,  $x \in Eq(X)$ .

**Theorem 5.6.** Let (X, d) be a Polish S-system where S is a C-semigroup. If X is an M-system and  $Eq(X) \neq \emptyset$  then X is minimal and equicontinuous.

*Proof.* Let  $x_0 \in X$  be an equicontinuity point. Since every M-system is (TT), by Lemma 3.3 we know that  $x_0 \in Trans(X)$ . Thus,  $\overline{Sx_0} = X$ . Therefore, for the minimality of X it is enough to show that  $x_0$  is a minimal point.

Since  $x_0 \in Eq(X)$ , given  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $0 < \delta < \frac{\epsilon}{2}$  and  $x \in B_{\delta}(x_0)$  implies  $d(sx_0, sx) < \frac{\epsilon}{2}$  for every  $s \in S$ . Since X is an M-system the set Y of all almost periodic points is dense. Choose  $y \in B_{\delta}(x_0) \cap Y$ . Then the set

$$N(y, B_{\delta}(x_0)) := \{ s \in S : sy \in B_{\delta}(x_0) \}$$

is a syndetic subset of S by Lemma 4.3. Clearly,  $N(y, B_{\delta}(x_0))$  is a subset of the set

$$N(x_0, B_{\epsilon}(x_0)) = \{ s \in S : d(sx_0, x_0) \le \varepsilon \}.$$

Then  $N_{\varepsilon} := N(x_0, B_{\epsilon}(x_0))$  is also syndetic (for every given  $\varepsilon > 0$ ). Using one more time Lemma 4.3 we conclude that  $x_0$  is a minimal point, as desired. Now the equicontinuity of X follows by Proposition 5.2.

**Theorem 5.7.** Let (X, d) be a Polish S-system where S is a C-semigroup. If X is an M-system which is not minimal or not equicontinuous. Then X is sensitive.

*Proof.* If X is non-sensitive then by Proposition 5.5 the system is almost equicontinuous. Theorem 5.6 implies that X is minimal and equicontinuous. This contradicts our assumption.  $\Box$ 

Now if the action is a cascade (X, f) or if S is a topological group (both are the case of C-semigroups, see Example 2.2) then we get, as a direct corollary, assertions (2) and (3) of Theorem 1.1. The assertion (1) is also covered in the case of Polish phase spaces X. Furthermore the main results are valid for a quite large class of actions including the actions of one-parameter semigroups on Polish spaces.

### References

- E. Akin, J. Auslander and K. Berg, When is a transitive map chaotic? in: Conference in Ergodyc Theory and Probability (V. Bergelson, K. March and J. Rosenblatt, eds.), de Gruyter, Berlin, 1996.
- J. Banks, J. Brooks, G. Cairns, G. Davis, P. Stacey, On Devaney's definition of chaos, Amer. Math. Monthly 99 (1992), 332-334.
- 3. R.L. Devaney, An introduction to chaotic dynamical systems, 2nd ed., Addison Wesley, 1989.
- 4. E. Glasner, Ergodic Theory via joinings, Math. Surveys and Monographs, AMS, 101, 2003.
- E. Glasner and M. Megrelishvili, Hereditarily non-sensitive dynamical systems and linear representations, Colloq. Math. 104 (2006), no. 2, 223-283.
- E. Glasner and B. Weiss, Sensitive dependence on initial conditions, Nonlinearity, 6 (1993), 1067-1075.
- S. Kolyada and L. Snoha, Some aspects of topological transitivity A survey, Grazer Math. Ber., Bericht Nr. 334 (1997), 3-35.
- S. Silverman, On maps with dense orbits and the definition of chaos, Rocky Mount. J. Math., 22 (1992), 353-375.

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, 52900 RAMAT-GAN, ISRAEL E-mail address: bizia@walla.co.il URL: http://www.math.biu.ac.il/~kantore

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, 52900 RAMAT-GAN, ISRAEL E-mail address: megereli@math.biu.ac.il URL: http://www.math.biu.ac.il/~megereli