# ASYMPTOTIC DIMENSION OF ONE RELATOR GROUPS

DMITRY MATSNEV

ABSTRACT. We show that one relator groups viewed as metric spaces with respect to the word-length metric have finite asymptotic dimension in the sense of Gromov and give an estimate of their asymptotic dimension in terms of the relator length.

#### 1. INTRODUCTION

A finitely generated group has asymptotic dimension not more than n if its underlying metric space (with respect to the length function corresponding to a given set of generators) has the following property: for any R > 0 there exists a uniformly bounded cover of the group such that any R-ball meets not more than (n+1) elements of the cover (see [Gr]). There is a way to extend this notion to all countable groups, not necessarily finitely generated.

Not all groups have finite asymptotic dimension  $(\mathbb{Z} \wr \mathbb{Z}$  is the standard example of a group which does not satisfy the definition above for any n), but for some classes of groups it is known that their asymptotic dimension is finite: for instance, Gromov noticed this for the case of hyperbolic groups in [Gr] (for precise proof, see [R]), Ji proved this for arithmetic groups in [Ji], and of course there are more examples.

The question whether a given group has finite asymptotic dimension has drawn more attention after results of Yu, who showed in [Yu] that such group satisfies the Novikov conjecture, and Higson and Roe, who deduced the exactness of such group in [HR].

The second part of the title concerns one relator groups. These are the ones which admit a presentation  $\langle S|r \rangle$ , where the generating set S is at most countable, and the only relator r is a word in S. Historically, one relator groups appeared as the fundamental groups of 2-manifolds, and one may regard them as a simplest class of groups which are close to free groups.

Date: July 20, 2006.

<sup>2000</sup> Mathematics Subject Classification. Primary 20F65; Secondary 20F05, 20F69.

Key words and phrases. asymptotic dimension, one relator groups.

### DMITRY MATSNEV

Within the scope of this work, the main motivation for the study of the asymptotic dimension of one relator groups were the results of Guentner (see [Gu]), showing that one relator groups are exact, and the ones of Beguin, Bettaieb, and Valette in [BBV], showing that the Baum-Connes conjecture (and therefore the Novikov conjecture as well) holds for such groups. A very natural question which arises immediately after comparing all the results mentioned above is: do one relator groups have finite asymptotic dimension? In this paper we are giving an affirmative answer to this question and estimate asymptotic dimension in terms of the relator length.

## 2. Asymptotic dimension of groups

First we outline a few basic definitions and some theorems on asymptotic dimension which shall be used later. For more detailed discussion and precise proofs, consult [R].

**Definition 2.1.** Given a metric space G, we say that its *asymptotic dimension* does not exceed n and write  $\operatorname{asdim} G \leq n$  if for any R > 0 there exists a uniformly bounded cover of G such that any R-ball in G meets not more than (n + 1) elements of the cover. The *asymptotic dimension* of G, asdim G, is then the minimal n satisfying this condition.

Any finitely generated group  $\Gamma$  with a generating set S can be endowed with a word-length metric

dist $(\gamma_1, \gamma_2)$  = length of a shortest word in  $S \cup S^{-1}$  representing  $\gamma_1^{-1} \gamma_2$ .

Since any two metrics for the same group, arising form different generating sets, are bi-Lipschitz equivalent, the notion of asymptotic dimension of a finitely generated group, viewed as a discrete metric space as stated above, is independent of the particular generating set.

**Definition 2.2.** For a (not necessarily finitely generated) group G, its asymptotic dimension, asdim G, is defined to be a supremum of asdim  $\Gamma$  over all finitely generated subgroups  $\Gamma$  of G.

This definition is consistent, for any inclusion of a subgroup into the over-group is a coarse embedding.

Now we collect some facts on the asymptotic dimension of groups, to be used in our discussion later on.

**Theorem 2.3.** For a subgroup H of a group G, asdim  $H \leq \operatorname{asdim} G$ .

**Theorem 2.4** (Bell and Dranishnikov, [BD]). For an HNN extension  $*_AG$  of a group G, asdim  $*_AG \leq 1 + \operatorname{asdim} G$ 

 $\mathbf{2}$ 

The original theorem in [BD] was formulated for a finitely generated base group G, but since any finitely generated subgroup of the HNN extension of G is a subgroup of an HNN extension of some finitely generated subgroup of G, one can run the original argument of Bell and Dranishnikov for every finitely generated subgroup of  $*_AG$  to obtain the theorem in the form we formulated here.

**Theorem 2.5** (Bell and Dranishnikov, [BD]). For a free product G \* H of two groups G and H, asdim  $G * H \leq \max\{\text{asdim } G, \text{asdim } H, 1\}$ .

Again, the assumption that both G and H are finitely generated is not crucial for the proof.

Finally, as a base for our inductive arguments in the next section, we state that the asymptotic dimension of a finite group is 0, and the asymptotic dimension of a free group is 1.

#### 3. One relator groups

Throughout this section, let G be a one relator group with a (possibly infinite) generating set S and relator r, that is,  $G = \langle S|r \rangle$  is a quotient of a free group on S by the minimal normal subgroup generated by r. We assume that r, a finite word in  $S \cup S^{-1}$ , is cyclically reduced as a word in the free group on S, and use |r| to denote its length in this free group. To omit the trivial cases, we assume that S contains at least two elements and |r| > 0.

For any real number x, we denote by  $\lceil x \rceil$  the minimal integer greater or equal to x.

# **Theorem 3.1.** In the notations above, asdim $G \leq \lceil \frac{|r|}{2} \rceil$ .

The rest of this section is devoted to the proof of this theorem.

First note that we can assume that G is finitely generated and every letter of S appears in r. Indeed, G is isomorphic to a free product of a finitely generated one relator group  $\Gamma$  with relator r and generating set consisting of letters which appear in r and the free group on all other letters. According to Theorem 2.5, asdim  $G \leq \max\{\operatorname{asdim} \Gamma, 1\}$ . If we can prove that  $\operatorname{asdim} \Gamma \leq \lceil \frac{|r|}{2} \rceil$ , the statement of the theorem for G will follow.

The argument is based on the induction on the length of r. For |r| = 1 the group G is isomorphic to a free group on all letters in S except the one which appears in r. Thus asdim  $G = 1 \le 1 = \lceil \frac{|r|}{2} \rceil$ .

For the inductive step suppose that the statement of the theorem has been proven for all one relator groups with relator length strictly less than |r|. Following the standard arguments of Magnus and Molchanovskii

#### DMITRY MATSNEV

(see [LS]), which we shall briefly describe in what follows, consider two cases:

Case 1. There exists a letter  $t \in S$  whose exponent sum in r is 0. To fix a notation, let  $S = \{t, b, c, d, ...\}$ , and, by means of a cyclic permutation or r, one may assume that the latter word begins with b or  $b^{-1}$ .

Let  $b_i$  denote  $t^i bt^{-i}$  for  $i \in \mathbb{Z}$ ,  $c_i$  denote  $t^i ct^{-i}$  for  $i \in \mathbb{Z}$ , and so on. Rewrite r scanning it from left to right and changing any occurrence of  $t^i x$  into  $t^i xt^{-i}t^i = x_it^i$  (here x represents any letter among  $b, c, d, \ldots$ , or their inverses), collecting the powers of adjacent t-letters together, and continuing with the leftmost occurrence of t or its inverse in the modified word. This way we do at least one cancellation of t and its inverse which happen to be next to each other, and the resulting word s, which represents r in terms of  $t, b_i, c_i, \ldots$ , and their inverses, has length not more than (|r| - 2).

Let m and M be the minimal and the maximal subscript of  $b_i$  occurring in s. Then

$$G \cong \langle t, b_m, \dots, b_M, c_i, d_i, \dots (i \in \mathbb{Z}) |$$
  
$$s, tb_i t^{-1} b_{i+1}^{-1} (i = m, \dots, M-1), tc_i t^{-1} c_{i+1}^{-1}, td_i t^{-1} d_{i+1}^{-1}, \dots (i \in \mathbb{Z}) > 0$$

Consider

$$H = \langle b_m, \ldots, b_M, c_i, d_i, \ldots (i \in \mathbb{Z}) | s \rangle.$$

According to our inductive assumption, asdim  $H \leq \lceil \frac{|s|}{2} \rceil \leq \lceil \frac{|r|}{2} \rceil - 1$ . Now  $G \cong *_{F < b_m, \dots, b_M, c_i, d_i, \dots, (i \in \mathbb{Z}) >} H$ , and, via Theorem 2.4, asdim  $G \leq 1 + \text{asdim } H \leq \lceil \frac{|r|}{2} \rceil$ .

Case 2. For  $al\bar{l}$  letters in S, their exponent sums in r are nonzero. Let  $S = \{u, v, c, d, ...\}$ , and assume that the exponent sums of u and v in r are  $\alpha$  and  $\beta$  respectively. Define the following homomorphism:

 $\Psi: u \mapsto bt^{-\beta}, v \mapsto t^{\alpha}, c \mapsto c, d \mapsto d, \dots$ 

Our group G embeds via  $\Psi$  into

$$C = < t, b, c, d, \dots | r(bt^{-\beta}, t^{\alpha}, c, d, \dots) >$$

and if p is the cyclically reduced  $r(bt^{-\beta}, t^{\alpha}, c, d, ...)$ , the exponent sum of t in p is 0, and b occurs in p.

Since G can be thought of as a subgroup of C, it would be enough to show that asdim  $C \leq \left\lceil \frac{|r|}{2} \right\rceil$ .

Now C is an HNN extension of some group H as in Case 1: assuming that p starts with b or  $b^{-1}$ , we introduce new variables  $b_i = t^i b t^{-i}$ ,  $c_i = t^i c t^{-i}$ , and so on for  $i \in \mathbb{Z}$ . Using these variables, we rewrite p (and therefore r as well) as a word s, eliminating all occurrences of t

4

and its inverse and substituting appropriate  $x_i$  for any other letter x among  $b, c, d, \ldots$  and their inverses. If p had at least two occurrences of t or  $t^{-1}$ , then  $|s| \leq |r| - 2$ , and, using our inductive assumption for s, asdim  $H \leq \lceil \frac{|r|-2}{2} \rceil$  as before. Invoking Theorem 2.4, we obtain the desired inequality asdim  $C \leq \lceil \frac{|r|}{2} \rceil$ .

If, however, p contains t or its inverse only at one place, p expresses t in terms of other generators, so that we can eliminate t from the generating set, and C is indeed a free group on remaining generators with asdim  $C = 1 \leq \left\lceil \frac{|r|}{2} \right\rceil$ .

Now the theorem is proven completely.

#### References

- [BBV] C. Beguin, H. Bettaieb, and A. Valette, K-theory for C\*-algebras of onerelator groups, K-Theory 16:277–298, 1999.
- [BD] G. Bell and A. Dranishnikov, A Hurewicz-type theorem for asymptotic dimension and applications to geometric group theory, to appear in Trans. Amer. Math. Soc.
- [Gr] M. Gromov, Asymptotic invariants of infinite groups, Cambridge University Press, Geometric Group Theory 2, 1993.
- [Gu] E. Guentner, Exactness of one relator groups, Proc. Amer. Math. Soc. 130(4): 1087–1093, 2002.
- [HR] N. Higson and J. Roe, Amenable group actions and the Novikov conjecture, J. Reine Angew. Math. 519:143–153, 2000.
- [Ji] L. Ji, Asymptotic dimension of arithmetic groups, preprint, 2003.
- [LS] R. Lyndon and P. Schupp, Combinatorial group theory, Springer-Verlag, 1977.
- [R] J. Roe, Lectures on coarse geometry, University Lecture Series 31 of the American Mathematical Society, 2003.
- [Yu] G. Yu, The Novikov Conjecture for groups with finite asymptotic dimension, Ann. of Math., 147(2):325–355, 1998.

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, U.S.A.

*Current address*: Departamento de Matemática, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 LISBOA, Portugal

E-mail address: matsnev@math.ist.utl.pt