Fixed Points of abelian actions

John Franks,* Michael Handel[†] and Kamlesh Parwani [‡]
October 27, 2019

Abstract

We prove that if \mathcal{F} is an abelian group of C^1 diffeomorphisms isotopic to the identity of a closed surface S of genus at least two then there is a common fixed point for all elements of \mathcal{F} .

1 Introduction and Notation

The Lefschetz theorem implies that if S is a closed surface of nonzero Euler characteristic and f is an element of the group $\operatorname{Diff}_0(S)$ of C^1 diffeomorphisms of S isotopic to the identity, then f has a fixed point. Several papers (see [1],[2], [9],[8] [6]) address the question of when an abelian subgroup \mathcal{F} of $\operatorname{Homeo}_0(S^2)$ or $\operatorname{Diff}_0(S^2)$ must have a global fixed point, meaning a point that is fixed by every element of the group. The $S = S^2$ case is settled in [8] where it is shown that there is always a subgroup \mathcal{F}_0 of index at most two with a global fixed point and that one can choose $\mathcal{F}_0 = \mathcal{F}$ if a certain \mathbb{Z}_2 -invariant vanishes. In this paper we therefore restrict our attention to those surfaces S with negative Euler characteristic.

The fixed point set $\operatorname{Fix}(f)$ is partitioned into Nielsen classes, each with its own Lefschetz index. In our current context, there is a preferred Nielsen class called the contractible Nielsen class whose Lefschetz index is nonzero. (Full definitions and details are provided in section 2.) The set of contractible fixed points of f will be denoted $\operatorname{Fix}_{\mathbf{c}}(f)$ and if \mathcal{F} is a subgroup of $\operatorname{Diff}_{\mathbf{0}}(S)$ then $\operatorname{Fix}_{\mathbf{c}}(\mathcal{F})$ will denote the points which are in $\operatorname{Fix}_{\mathbf{c}}(f)$ for all f in \mathcal{F} . We can now state our main theorem.

Theorem 1.1. Suppose S is a closed oriented surface of genus at least two and that \mathcal{F} is an abelian subgroup of $\mathrm{Diff}_0(S)$ Then $\mathrm{Fix}_c(\mathcal{F})$ is non-empty. In particular $\mathrm{Fix}(\mathcal{F})$ is non-empty.

^{*}Supported in part by NSF grant DMS0099640.

[†]Supported in part by NSF grant DMS0103435.

[‡]Supported in part by NSF grant DMS0244529.

There are two aspects of our proof that are of independent interest. In section 3 we develop a theory of Nielsen classes relative to certain infinite compact subsets K of S and more importantly of their Lefshetz indices. The key here is in isolating the property of K that is required for the standard definitions to work.

In section 4 we define reducing curves for a finite subset of \mathcal{F} relative to certain \mathcal{F} -invariant compact sets. These are analogous to the reducing curves that occur for Thurston canonical form, allowing us to decompose the surfce into simpler subsurfaces on which the group action is sufficiently well understood as to allow analysis. The methods of [8] are used in, but are not sufficient for, the construction of the set of reducing curves.

2 Contractible Fixed Points

Throughout this article S will denote a closed oriented surface of genus at least two.

A path α with endpoints in Fix(f) is an f-Nielsen path if $f(\alpha)$ is homotopic to α rel endpoints. Elements of Fix(f) are f-Nielsen equivalent if they are connected by an f-Nielsen path. As the notation indicates, this defines an equivalence relation on Fix(f) whose equivalence classes are referred to as f-Nielsen classes. When f is understood we refer to these simply as Nielsen paths and Nielsen classes.

Since S has arbitrarily small contractible neighborhoods, each Nielsen class is an open subset of Fix(f), and since the complement of a Nielsen class is a union of Nielsen classes, each Nielsen class is also closed. In particular, there are only finitely many Nielsen classes.

If $\tilde{f}: \tilde{S} \to \tilde{S}$ is a lift to the universal cover \tilde{S} of S then any path $\tilde{\alpha}$ with endpoints in $\operatorname{Fix}(\tilde{f})$ projects to a Nielsen path in S. Conversely, if α is a Nielsen path and \tilde{f} fixes one endpoint of a lift $\tilde{\alpha}$ of α then it also fixes the other endpoint of $\tilde{\alpha}$. Thus $\operatorname{Fix}(\tilde{f})$ is either empty or projects onto a single Nielsen class in $\operatorname{Fix}(f)$ that we say is the Nielsen class determined by \tilde{f} . If T is a covering translation of \tilde{S} then $\operatorname{Fix}(\tilde{f})$ and $\operatorname{Fix}(T\tilde{f}T^{-1}) = T(\operatorname{Fix}(\tilde{f}))$ project to the same Nielsen class of $\operatorname{Fix}(f)$. Conversely, if $\operatorname{Fix}(\tilde{f})$ and $\operatorname{Fix}(\tilde{f}')$ project to the same Nielsen class then there is a covering translation T such that \tilde{f} and $T\tilde{f}T^{-1}$ have a common fixed point and hence are equal.

If f is isotopic to the identity then there is a unique lift $\tilde{f}: \tilde{S} \to \tilde{S}$, called the identity lift of f, that commutes with all covering translations of \tilde{S} . Equivalently, if f_t is any isotopy from f_0 = identity to $f_1 = f$, and if \tilde{f}_t is the lift of f_t with \tilde{f}_0 = identity, then \tilde{f}_1 is the identity lift. The uniqueness of the identity lift follows from the fact that the group of covering translations has trivial center and so uses the fact that the Euler characteristic of S is negative. The identity lift has non-empty fixed point set. Indeed, if this fails then the unit tangent vector at \tilde{x} in the direction of the geodesic segment from \tilde{x} to $\tilde{f}(\tilde{x})$ determines a non-zero equivariant vector field on \tilde{S} and hence a non-zero vector field on S. The Nielsen class determined by the identity lift is called the contractible Nielsen class and is denoted Fix_c(f). Elements

of $\operatorname{Fix}_{c}(f)$ are said to be *contractible*. (The terminology comes from the fact that if f_{t} is an isotopy with $f_{0} = id$ and $f_{1} = f$ then the closed loop $f_{t}(p)$ is contractible in S.) Note that if $x \in \operatorname{Fix}_{c}(f)$ and \tilde{f} is the identity lift of f then \tilde{f} fixes every lift of x since it fixes some lift of x and commutes with every covering translation of \tilde{S} .

Suppose now that \mathcal{F} is a torsion free finitely generated abelian subgroup of $\operatorname{Homeo}_0(S)$. Because the identity lift \tilde{f} of each $f \in \mathcal{F}$ is unique, $f \to \tilde{f}$ defines a lift of \mathcal{F} to a subgroup $\tilde{\mathcal{F}}$ of $\operatorname{Homeo}(\tilde{S})$ called the *identity lift of* \mathcal{F} . We say that $x \in S$ is a *global fixed point* if $x \in \operatorname{Fix}(f)$ for all $f \in \mathcal{F}$ and is a *global contractible fixed point* if $x \in \operatorname{Fix}_c(f)$ for all $f \in \mathcal{F}$; these sets are denoted $\operatorname{Fix}(\mathcal{F})$ and $\operatorname{Fix}_c(\mathcal{F})$ respectively.

Lemma 2.1. If $\{f_1, \ldots, f_m\}$ generate an abelian subgroup \mathcal{F} of $\operatorname{Homeo}_0(S)$ then $\bigcap_{i=1}^m \operatorname{Fix}_c(f_i) \subset \operatorname{Fix}_c(\mathcal{F})$.

Proof. If $\tilde{x} \in \tilde{S}$ is a lift of $x \in \bigcap_{i=1}^m \operatorname{Fix_c}(f_i)$ then $\tilde{x} \in \operatorname{Fix}(\tilde{f}_i)$ for the identity lift \tilde{f}_i of each f_i . Since the \tilde{f}_i 's generate the identity lift $\tilde{\mathcal{F}}$ of \mathcal{F} , we have $x \in \operatorname{Fix_c}(\tilde{f})$ for each $\tilde{f} \in \tilde{\mathcal{F}}$. Thus $x \in \operatorname{Fix_c}(\mathcal{F})$.

Our interest in contractible fixed points comes from the following lemma.

Lemma 2.2. If there exists a global contractible fixed point for a finite index subgroup of \mathcal{F} then there exists a global contractible fixed point of \mathcal{F} .

Proof. If \mathcal{F}_0 has finite index in \mathcal{F} , if $x \in \operatorname{Fix}_c(\mathcal{F}_0)$ and if $\tilde{\mathcal{F}}_0$ is the identity lift of \mathcal{F}_0 then $\tilde{x} \in \operatorname{Fix}(\tilde{\mathcal{F}}_0)$ for any lift \tilde{x} of x. The orbit of \tilde{x} under $\tilde{\mathcal{F}}$ is therefore finite and it follows from Theorem 1.2 of [8] that $\operatorname{Fix}(\tilde{\mathcal{F}})$, and hence its projected image $\operatorname{Fix}_c(\mathcal{F})$, are non-empty.

By combining Lemma 2.2 with a result from [8] we obtain a criterion for proving the existence of global contractible fixed points.

Proposition 2.3. Suppose that \mathcal{F}_0 is a finite index subgroup of a finitely generated abelian subgroup \mathcal{F} of $\operatorname{Homeo}_0(S)$ and that K is a compact \mathcal{F}_0 -invariant set. Suppose also that $\gamma \subset S \setminus K$ is essential in $S \setminus K$ but inessential in S. Finally, suppose that there is a lift $\tilde{\gamma}$ of γ to the universal covering \tilde{S} that is preserved up to isotopy rel \tilde{K} , the complete pre-image of K, by each \tilde{f} in the identity lift \tilde{F}_0 of F_0 . Then $\operatorname{Fix}_{\mathbf{c}}(\mathcal{F}) \neq \emptyset$.

Proof. Let $\tilde{K}_{\tilde{\gamma}}$ be the intersection of \tilde{K} with the disk in \tilde{S} bounded by $\tilde{\gamma}$. Then $\tilde{K}_{\tilde{\gamma}}$ is compact and is $\tilde{\mathcal{F}}_0$ -invariant, meaning that it is \tilde{f} -invariant for every $\tilde{f} \in \tilde{\mathcal{F}}_0$. It follows from Theorem 1.2 of [8] that $\operatorname{Fix}(\tilde{\mathcal{F}}_0) \neq \emptyset$ and hence that $\operatorname{Fix}_{\mathbf{c}}(\mathcal{F}_0) \neq \emptyset$. By Lemma 2.2, $\operatorname{Fix}_{\mathbf{c}}(\mathcal{F}) \neq \emptyset$.

3 Relative Nielsen Classes

Some definitions in this section apply beyond the context of surface homeomorphisms so we consider homeomorphisms $h:M\to M$ of a manifold of unspecified dimension. For any compact h-invariant set K we can define isotopy and homotopy rel K and in particular can define Nielsen classes rel K. However, to make sense of the Lefschetz index of a Nielsen class we must place restrictions on K. In particular we want the relative Nielsen classes to be open and closed subsets of $\operatorname{Fix}(h)$. We make the following definitions.

Definition 3.1. If $\alpha(s)$ and $\beta(s)$ are paths with the same endpoints in a space M and K is a closed subset of M we will say that α and β are homotopic rel K provided there is a homotopy $\alpha_t(s)$, $t \in [0,1]$ satisfying

- 1. $\alpha_0 = \alpha$ and $\alpha_1 = \beta$.
- 2. If $\alpha_{t_0}(s) \in K$ for some s and some t_0 then $\alpha_t(s) = \alpha_{t_0}(s)$ for all $t \in [0,1]$.
- 3. $\alpha_t(0) = x$, $\alpha_t(1) = y$ for all $0 \le t \le 1$.

Equivalently $\alpha_0(s)$ and $\alpha_1(s)$ are homotopic rel K if they are homotopic as maps $[0,1] \to S$ relative to the subset $\alpha^{-1}(K) \cup \{0\} \cup \{1\}$ of [0,1].

Definition 3.2. If $h: M \to M$ is a homeomorphism with compact fixed point set and K is a compact invariant set we will say that an arc α joining points $x, y \in Fix(h)$ is a Nielsen arc for h rel K if there is a homotopy rel K from α to $h(\alpha)$.

Definition 3.3. If $x \in \text{Fix}(h)$ and if there are arbitrarily small neighborhoods $U_x \subset V_x$ of x with the property that for every $y \in U_x \cap \text{Fix}(h)$ there is a Nielsen arc α in U_x joining x and y such that the image of a homotopy rel K from α to $h(\alpha)$ lies in V_x then we say that h has locally constant Nielsen behavior rel K at x. If h has locally constant Nielsen behavior rel K at x for every $x \in \text{Fix}(h)$ then we say that h has locally constant Nielsen behavior rel K.

If $x \in \text{Fix}(h)$ is either not in K or is an isolated point in $\text{Fix}(h) \cap K$ then h has locally constant Nielsen behavior rel K at x. One can choose $U_x \subset V_x$ to be any contractible neighborhoods such that $f(U_x) \subset V_x$ and such that $K \cap V_x$ is either empty or $\{x\}$.

For general $x \in K$, h need not have locally constant Nielsen behavior at x as the following example illustrates. Let $h : \mathbb{R}^2 \to \mathbb{R}^2$ be given in polar co-ordinates by $h(r,\theta) = (r,\theta+f(r))$ where f(r) is a C^{∞} function with values between 0 and 1 which vanishes if and only if r=0 or $r=1/n, =1,2,3,\ldots$ Choose ν_n so that $1/(n+1) < \nu_n < 1/n$ and so $f(\nu_n)$ is a non-zero rational multiple of π . Then any point on the circle of radius ν_n is h-periodic with period > 1. Let K be a compact invariant set consisting of the origin together with a single periodic orbit in the circle of radius ν_n for each n. There is no Nielsen arc rel K joining the origin to any other fixed point (the other fixed points must lie on circles of radius 1/n).

Definition 3.4. Suppose h has locally constant Nielsen behavior rel K and let $x, y \in \text{Fix}(h)$. We say that x is Nielsen equivalent to y rel K if there is a Nielsen arc rel K connecting x to y. The Nielsen equivalence class of $x \in \text{Fix}(h)$ relative to K is denoted $\mathcal{N}(h, K, x)$.

It is immediate that if h has locally constant Nielsen behavior rel K then each Nielsen class is both open and closed in Fix(h). In particular, if Fix(h) is compact then there are only finitely many Nielsen classes rel K.

Having defined relative Nielsen classes in the appropriate context, we now define the Lefschetz index for a relative Nielsen class.

Definition 3.5. Suppose $h_t: M \to M$ is an isotopy rel K from h_0 to h_1 and suppose $H: M \times I \to M \times I$, given by $H(x,t) = (h_t(x),t)$, has locally constant Nielsen behavior rel $K \times I$. Let $\mathcal{N}_0 = \mathcal{N}(h_0, K, x_0)$ be a Nielsen class rel K for h_0 . We define \mathcal{N}_t , the Nielsen class rel K determined by \mathcal{N}_0 and h_t , to be $\mathcal{N}_t = \pi(\mathcal{N} \cap (M \times \{t\}))$ where $\mathcal{N} = \mathcal{N}(H, K \times I, (x_0, 0))$ and $\pi: M \times I \to M$ is projection.

The fact that H has locally constant Nielsen behavior rel $K \times I$ implies that h_t has locally constant Nielsen behavior rel K for each t. This is because if α is a Nielsen arc in $S \times I$ joining (x,t) to (y,t) then $\pi \circ \alpha$ is a Nielsen arc for $h_t : S \to S$. This same observation shows that \mathcal{N}_t is a Nielsen class rel K for h_t and in fact $\mathcal{N}_t = \mathcal{N}(h_t, K, x)$ for any $(x,t) \in \mathcal{N}$.

If X_0 is an open and closed subset of $\operatorname{Fix}(h)$ then the Lefschetz number $L(X_0,h)$ is well defined. We recall the definition of $L(X_0,h)$ following [4]. Choose compact X and open V such that $X_0 \subset X \subset V \subset S$ and such that $\operatorname{Fix}(h|V) = X_0$. We may embed M as a submanifold of \mathbb{R}^m for some m and consider a retraction $r:W \to M$ where W is a neighborhood of S. Let $W_V = r^{-1}(V)$ which is an open neighborhood of X in \mathbb{R}^m . There is a canonical "fundamental" homology class $[u] \in H_m(W_V, W_V \setminus X)$. This class is characterized by the property that for each $x \in X$ the inclusion $(W_V, W_V \setminus X) \to (W_V, W_V \setminus \{x\})$ takes u to the standard generator of $H_m(W_V, W_V \setminus \{x\})$. The preimage of 0 under $(id - h \circ r): W_V \to \mathbb{R}^m$ is X_0 so

$$(id - h \circ r) : (W_V, W_V \setminus X) \to (\mathbb{R}^m, \mathbb{R}^m \setminus \{0\}).$$

The Lefschetz fixed point index $L(X_0,h)$ of X_0 is the integer defined by

$$(id - h \circ r)_*([u]) = L(X_0, h)[o_m]$$

where $[o_m] \in H_m(\mathbb{R}^m, \mathbb{R}^m \setminus \{0\})$ is the standard generator. A proof that $L(X_0, h)$ depends only on X_0 and h and not on the choice of X and V can be found in [4].

Proposition 3.6. Suppose $h_t: M \to M$ is an isotopy rel K from h_0 to h_1 and suppose $H: M \times I \to M \times I$, given by $H(x,t) = (h_t(x),t)$, has locally constant Nielsen behavior rel $K \times I$. Let \mathcal{N}_0 be a Nielsen class rel K for h_0 and let \mathcal{N}_t be the Nielsen class determined by \mathcal{N}_0 and h_t . Then $L(\mathcal{N}_t, h_t)$ is independent of t.

Proof. Let Y be an open neighborhood in $M \times I$ of the compact Nielsen class $\mathcal{N} = \mathcal{N}(H, K \times I, (x_0, 0))$ where $x_0 \in \mathcal{N}_0$ and assume Y is chosen to be disjoint from any fixed points of H which are not in \mathcal{N} . We define $V_t = Y \cap (M \times \{t\})$. It follows from the fact that \mathcal{N} is compact that for each t_0 there is an $\epsilon > 0$, a compact set $X \subset M$, and an open set $V \subset M$ such that for each $t \in [t_0 - \epsilon, t_0 + \epsilon]$ we have $\mathcal{N}_t \subset X \subset V \subset V_t$.

It then follows from the definition of the Lefschetz index cited above that $L(N_t, h_t)$ can be defined in terms of X and V for $t \in [t_0 - \epsilon, t_0 + \epsilon]$. The defining equation becomes $(id - h_t \circ r)_*([u]) = L(\mathcal{N}_t, h_t)[o_m]$ which depends only on the homotopy class of h_t and is therefore constant on $[t_0 - \epsilon, t_0 + \epsilon]$. Therefore by the compactness of $[0, 1], L(N_t, h_t)$ is independent of t.

We now return to the surface case $f: S \to S$. The following lemma uses smoothness to establish a strong form of locally constant Nielsen behavior relative to Fix(f).

Lemma 3.7. Suppose that $f \in \text{Diff}_0(S)$, that x is an accumulation point in Fix(f) and that $U_{\epsilon}(x)$ is the open ϵ neighborhood of x in local coordinates around x. Then the following hold for all sufficiently small $\epsilon > 0$ and for $\alpha_{x,y}$ the straight line segment from x to y.

- 1. If $y \in \text{Fix}(f) \cap U_{\epsilon}(x)$ then $f(\alpha_{x,y}) \subset U_{\epsilon}(x)$.
- 2. There is a neighborhood W of f in $Diff_0(S)$ so that if $g \in W$, if $p \notin Fix(g)$ and if $p, g(p) \in U_{\epsilon}(x)$ then $\alpha_{p,g(p)} \cap Fix(g) = \emptyset$.

In particular, the straight line homotopy between $\alpha_{x,y}$ and $f(\alpha_{x,y})$ is a homotopy rel Fix(f) for all $y \in \text{Fix}(f) \cap U_{\epsilon}(x)$.

Proof. If $y_j \in \text{Fix}(f)$ and $y_j \to x$ then $(y_j - x)/\|y_j - x\|$ converges to a unit vector that is fixed by Df_x . If z_j is contained in the interior of α_{x,y_j} then $(z_j - x)/\|z_j - x\|$ and $(y_j - z_j)/\|y_j - z_j\|$ converge to this same unit vector. Thus, for all sufficiently large j, the angle between $\alpha_{x,f(z_j)}$ and α_{x,z_j} is less than one degree and similarly the angle between $\alpha_{y_j,f(z_j)}$ and α_{y_j,z_j} is less than one degree. In particular $f(\alpha_{x,y_j})$ is contained in the closure of U_{ϵ_j} where ϵ_j is the distance from x to y_j . This proves (1).

If (2) fails then there exist $f_j \to f$, $p_j \notin \text{Fix}(f_j)$ such that $p_j \to x$ and $q_j \in \text{Fix}(f_j) \cap \alpha_{p_j,f_j(p_j)}$. In this case $(p_j - q_j)/\|p_j - q_j\|$ converges to a eigenvector of Df_x with negative eigenvalue. This contradicts the fact that f preserves orientation and Df_x has an eigenvector with eigenvalue 1.

Motivated by Lemma 3.7 we focus on K satisfying the property that each $x \in K$ has a neighborhood that is either disjoint from Fix(f) or contained in Fix(f).

Lemma 3.8. Suppose that $f \in \text{Diff}_+(S)$ and that K is a compact invariant set such that $\text{Fix}(f) \cap K$ is an open (and closed) subset of K. Then f has locally constant Nielsen behavior rel K. Moreover, if f_t is an isotopy rel K and $F: S \times I \to S \times I$ is given by $F(x,t) = (f_t(x),t)$ then each f_t has locally constant Nielsen behavior rel K and F has locally constant Nielsen behavior rel K.

Proof. For the first part, we must show that f has locally constant Nielsen behavior rel K at x for each $x \in \text{Fix}(f)$ and for this there is no loss in assuming that x is an accumulation point in $\text{Fix}(f) \cap K$. We work in local coordinates around $x \in \text{Fix}(f)$, letting $U_{\epsilon}(x)$ be the open ϵ ball around x and letting $\alpha_{y,z}$ be the straight line segment connecting y to z. By Lemma 3.7 there exists $\epsilon > 0$ so that, setting $U = V = U_{\epsilon}(x)$, there is, for each $y \in \text{Fix}(f) \cap U$, a homotopy rel Fix(f) from $\alpha_{x,y}$ to $f(\alpha_{x,y})$ through paths in V. If ϵ is sufficiently small then $K \cap V \subset \text{Fix}(f) \cap V$ and so the homotopy is also rel K.

We proceed to the argument for F. Suppose that $0 \le t \le 1$ and that $(x,t) \in \operatorname{Fix}(F)$. Note that $K \cap \operatorname{Fix}(f_s)$ is independent of s since the isotopy is rel K. Thus each f_s has locally constant Nielsen behavior rel K. Moreover, there is no loss in assuming that x is an accumulation point in $\operatorname{Fix}(f_t) \cap K$. For $\gamma > 0$ let $J_{\gamma}(t)$ be the γ neighborhood of t in I. Choose ϵ_t as in the last sentence of the preceding paragraph with f replaced by f_t and choose δ_t so that $f_t(U_{\delta_t}(x)) \subset U_{\epsilon_t/2}(x)$. Denote $U_{\delta_t}(x) \times J_{\gamma}(t)$ by U and $U_{\epsilon_t}(x) \times J_{\gamma}(t)$ by V. Thus $U \subset V$ and if γ is sufficiently small then $F(U) \subset V$. If $(y,s) \in \operatorname{Fix}(F) \cap U$ let β be the path in U connecting x to y that is the concatenation of the "vertical" arc $\{x\} \times [t,s]$ with the "horizontal" arc $\alpha_{x,y} \subset U \times \{s\}$. To prove that $F(\beta)$ is homotopic to β rel $K \times I$ through paths in V it suffices to show that the straight line homotopy from $f_s(\alpha_{x,y})$ to $\alpha_{x,y}$ in $U_{\epsilon_t}(x)$ is a homotopy relative to K. This follows from Lemma 3.7(2) if γ is sufficiently small. \square

We conclude this section with a variant of Proposition 2.3.

Lemma 3.9. Suppose that $f \in \text{Homeo}_0(S)$, that K is a compact f-invariant set and $D \subset S$ is a disk whose boundary is disjoint from K and whose interior intersects K in a non-empty set that is disjoint from Fix(f). Suppose further that there is a lift $\tilde{f}: \tilde{S} \to \tilde{S}$, a component \tilde{D}_0 of the full pre-image of D in \tilde{S} and an isotopy rel \tilde{K} , the complete pre-image of K, from \tilde{f} to a homeomorphism that preserves \tilde{D}_0 . Then there exists $\tilde{x} \in \text{Fix}(\tilde{f})$ whose projected image $x \in \text{Fix}(f)$ is not Nielsen equivalent rel K to any element of K. If \tilde{f} is the identity lift then $x \in \text{Fix}_c(f)$.

Proof. The compact set $C = \tilde{K} \cap \tilde{D}$ is \tilde{f} -invariant. For any $\tilde{y} \in \tilde{K} \cap \operatorname{Fix}(\tilde{f})$, our hypotheses imply that \tilde{y} is disjoint from \tilde{D} and that if $R_{\tilde{y}} \subset \tilde{S}$ is a properly embedded ray $R_{\tilde{y}} \subset \tilde{S}$ initiating at \tilde{y} that is disjoint from \tilde{D} then $\tilde{f}(R_{\tilde{y}})$ is properly homotopic to $R_{\tilde{y}}$ rel C and rel endpoint. By Proposition 5.3 of [5], there exists $\tilde{x} \in \operatorname{Fix}(\tilde{f})$ for which there does not exist a properly embedded ray $R_{\tilde{x}} \subset \tilde{S}$ initiating at \tilde{x} such that $\tilde{f}(R_{\tilde{x}})$ is properly homotopic to $R_{\tilde{x}}$ rel C and rel endpoint. It follows that there does not exist an arc $\tilde{\alpha}$ connecting \tilde{x} to \tilde{y} such that $\tilde{f}(\alpha)$ is homotopic to α rel \tilde{C} and rel endpoints. The projected image $x \in \operatorname{Fix}(f)$ of \tilde{x} is therefore not Nielsen equivalent rel K to any $y \in K$.

4 Reducing Curves

We assume throughout this section that f_1, \ldots, f_m generate an abelian subgroup $\mathcal{F} \subset \mathrm{Diff}_0(S)$.

For each $x \in S$, define $A_x \subset \{1, \ldots, m\}$ by $i \in A_x$ if and only if $x \in Fix(f_i)$. The cardinality of A_x is called the *multiplicity* of x. A set K has *locally constant multiplicity* if it satisfies the equivalent conditions of the following lemma.

Lemma 4.1. The following are equivalent for any set K.

- 1. Each $x \in K$ has a neighborhood U such that the multiplicity of $y \in K \cap U$ is independent of y.
- 2. Each $x \in K$ has a neighborhood U such that A_y is independent of $y \in K \cap U$.
- 3. Fix $(f_i) \cap K$ is an open subset of K for all i,

Proof. Let \mathcal{P} denote the power set of $\{1, \ldots, m\}$. Property (2) asserts that the function $\mathcal{A}: K \to \mathcal{P}$ given by $\mathcal{A}(x) = A_x$ is locally constant. Likewise, property (1) asserts that the function $card_{\mathcal{A}}: K \to \mathbb{Z}$ is locally constant where $card_{\mathcal{A}}(x)$ is the cardinality of A_x . Finally (3) asserts that for each i the function $\psi_i: K \to \{0, 1\}$ is locally constant, where $\psi_i(x) = 1$ if $x \in \text{Fix}(f_i)$ and 0 otherwise.

To see that (2) implies (3) observe that $\psi_i = \phi_i \circ \mathcal{A}$ where $\phi_i : \mathcal{P} \to \{0, 1\}$ is the function defined by $\phi_i(A) = 1$ if $i \in A$ and 0 otherwise.

To see that (3) implies (1) observe that the function

$$card_{\mathcal{A}}(x) = \sum_{i=1}^{m} \psi_i(x).$$

To see that (1) implies (2) note the fact each $\operatorname{Fix}(f_i)$ is a closed set implies the function \mathcal{A} is semicontinuous in the sense that if $\lim x_n = x$ and $\mathcal{A}(x_n) = A_0$ is independent of n then $A_0 \subset A_x$. If in addition $\operatorname{card}_{\mathcal{A}}$ is locally constant it follows that the cardinality of A_0 equals the cardinality of A_x so $A_0 = A_x$. This implies that the function \mathcal{A} is continuous or equivalently locally constant.

Lemma 4.2. Suppose that Y and C are compact disjoint \mathcal{F} -invariant subsets of S and that $Y \subset \bigcup_{i=1}^m \operatorname{Fix}(f_i)$. Then there is a compact \mathcal{F} -invariant subset $K \subset Y$ with locally constant multiplicity such that for all i, each $y \in Y \cap \operatorname{Fix}(f_i)$ is f_i -Nielsen equivalent rel $K \cup C$ to some $x \in K$.

Proof. The proof is by induction on the maximal multiplicity M(Y) of an element of Y. If M(Y) = 1 then K = Y satisfies the conclusions of the lemma. We may therefore assume that the lemma holds for Y' with M(Y') < M(Y). Denote M(Y) by M and let K_M be the set of points in Y with multiplicity M.

If $x \in K_M$ is isolated in K_M , let U_x be an open neighborhood of x that intersects K_M in $\{x\}$. Otherwise, x is an accumulation point in K_M and hence an accumulation

point in $\operatorname{Fix}(f_i) \cap K$ for each $i \in A_x$. In this case we let U_x be an open neighborhood of x such that $U_x \cap \operatorname{Fix}(f_i) = \emptyset$ if $i \notin A_x$ and such that for each $i \in A_x$ and each $y \in U_x \cap \operatorname{Fix}(f_i)$ there is a path $\beta_{x,y}$ connecting x to y in U_x and a homotopy rel $\operatorname{Fix}(f_i)$ from $\beta_{x,y}$ to $f_i(\beta_{x,y})$ through paths in U_x . Such a neighborhood exists by Lemma 3.7.

The U_x 's cover K_M and so have a finite subcover $\{U_{x_1}, \ldots, U_{x_k}\}$. Define W to be the \mathcal{F} -orbit of $\bigcup_{j=1}^k U_{x_j}$, Y' to be the intersection of Y with the complement of W and C' to be $C \cup K_M$. The inductive hypothesis applies to Y' and C' to produce $K' \subset Y'$; let $K = K' \cup K_M$.

K has locally constant multiplicity because both K' and K_M do and are disjoint. It suffices to show that each $y \in Y \cap \operatorname{Fix}(f_i)$ is f_i -Nielsen equivalent rel $K \cup C$ to some $x \in K$. This follows from the inductive hypothesis if $y \in Y'$. We may therefore assume that $y \in W$. Suppose at first that $y \in U_{x_j}$ for some j. Since $\operatorname{Fix}(f_i) \cap U_{x_j} \supset K_M \cap U_{x_j} = (K \cup C) \cap U_{x_j}$ the homotopy from $\beta_{x_j,y}$ to $f_i(\beta_{x_j,y})$ is rel $K \cup C$. Thus y is f_i -Nielsen equivalent to x_j rel $K \cup C$. For the general case we must consider f(y) where $y \in U_{x_j}$ and $f \in \mathcal{F}$. Since K, C and $\operatorname{Fix}(f_i)$ are all f-invariant and since f commutes with f_i , the path $\gamma = f(\beta_{x_j,y})$ connecting $f(x_j)$ to f(y) is homotopic rel $K \cup C$ to $f_i(\gamma)$ showing that f(y) is f_i -Nielsen equivalent to $f(x_j)$ rel $K \cup C$.

Definition 4.3. A reducing set for $\{f_1, \ldots, f_m\}$ relative to a compact \mathcal{F} -invariant set K is a finite collection \mathcal{R} of disjoint simple closed curves in S with the following properties.

- 1. $\mathcal{R} \cap K = \emptyset$ and \mathcal{R} is \mathcal{F} -invariant up to isotopy rel K.
- 2. Each contractible component of $S \setminus \mathcal{R}$ contains at least two elements of K.
- 3. For i = 1, ..., m the following holds:
 - (3-i) If a component S_l of $S \setminus \mathcal{R}$ intersects K in an infinite set and if $S_l \cap K$ contains a fixed point of f_i , then f_i is isotopic rel K to a homeomorphism that is the identity on S_l .

Note that if there is a reducing set for $\{f_1, \ldots, f_m\}$ then each $K \cap \text{Fix}(f_i)$ is an open subset of K and so K has locally constant multiplicity.

Definition 4.4. If \mathcal{N}_i is a union of Nielsen classes for f_i then a compact set K is \mathcal{N}_i -complete if each element of \mathcal{N}_i is Nielsen equivalent rel K to an element of K.

Proposition 4.5. Suppose that \mathcal{N}_i is a union of Nielsen classes for f_i and that $K \subset \bigcup_{i=1}^m \mathcal{N}_i$ is compact, is \mathcal{F} -invariant, has locally constant multiplicity and is \mathcal{N}_i -complete for each $1 \leq i \leq m$. Then there is a reducing set \mathcal{R} for $\{f_1, \ldots, f_m\}$ rel K. Moreover if \mathcal{R}' is a finite set of disjoint simple closed curves satisfying (1) and (2) of Definition 4.3 then we may choose \mathcal{R} to contain \mathcal{R}' .

Proof. We show below that there is a finite collection \mathcal{R}_m of disjoint simple closed curves containing \mathcal{R}' and satisfying (1), (2) and (3-m) of Definition 4.3. This proves the proposition in the case that m = 1. Assume by induction on m that the proposition holds for $\{f_1, \ldots, f_{m-1}\}$. To produce the desired reducing set \mathcal{R} for $\{f_1, \ldots, f_m\}$ rel K, apply the inductive statement to $\{f_1, \ldots, f_{m-1}\}$ with R' replaced by \mathcal{R}_m .

We now prove the existence of \mathcal{R}_m , denoting f_m , \mathcal{R}_m and \mathcal{N}_m simply by f, \mathcal{R} and \mathcal{N} . The proof is based on arguments from section 2 of [8]. Both $B = \text{Fix}(f) \cap K$ and $C = K \setminus B$ are \mathcal{F} -invariant and are compact by the assumption that K has locally constant multiplicity. If B is finite then simply take $\mathcal{R} = \mathcal{R}'$. We assume now that B is infinite.

Define W to be the set of compact subsurfaces $W \subset S$ such that:

- (W-1) ∂W has finitely many components, each of which is disjoint from K and is either disjoint from \mathcal{R}' or is isotopic rel K to an element of \mathcal{R}' . Moreover, each contractible component of $S \setminus \partial W$ contains at least two points in K.
- (W-2) W contains all but finitely many points of B and every component of W intersects B in an infinite set.
- (W-3) f is isotopic rel K to a homeomorphism that is the identity on W.

Lemma 4.1 of [9] and the isotopy extension theorem (see the proof of Theorem 1.2 of [7] where W is denoted $W(\langle f \rangle, B, K \setminus B)$; see also Lemma 3.7 of this paper) imply that $W \neq \emptyset$. (This uses the fact that B is compact and is open in K.) There is a partial order on W defined by $W_1 < W_2$ if and only if W_1 is isotopic rel K to a subsurface of W_2 but is not isotopic rel K to W_2 .

The next step in the proof is to show that W has maximal elements. This is essentially Lemma 2.7 of [8]; the main difference is that instead of assuming that B contains each element of \mathcal{N} we are assuming that every element of \mathcal{N} is Nielsen equivalent rel K to an element of B. To compensate for this, we quote Lemma 3.9 instead of the Brouwer lemma to produce a fixed point in a certain universal cover. We make this precise as follows.

If $W_1 \in \mathcal{W}$ is not contained in a maximal element of \mathcal{W} then there is an infinite increasing sequence $W_1 \subset W_2 \subset \ldots$ of elements of \mathcal{W} that are non-isotopic rel K. We may assume that $W_l \cap B$ and the number of components of W_l are independent of l. The number of components of $S \setminus W_l$ is unbounded. We may therefore choose l so that some complementary component of W_l is a disk D that is disjoint from B.

There is a homeomorphism $\phi: S \to S$ that is isotopic to f rel K and such that $\phi|_{W_l}$ is the identity. Choose a component \tilde{W}_l^* of the full pre-image \tilde{W}_l of W_l in \tilde{S} and let $\tilde{\phi}$ be the lift of ϕ that restricts to the identity on \tilde{W}_l^* . The isotopy rel K from ϕ to f lifts to an isotopy rel K from $\tilde{\phi}$ to some lift \tilde{f} of f. Choose $g \in S \cap W_l$ and a lift $\tilde{g} \in \tilde{W}_l^*$. Then $\tilde{g} \in Fix(\tilde{\phi})$ and, since $\tilde{\phi}$ is isotopic to \tilde{f} rel K, $\tilde{g} \in Fix(\tilde{f})$. Thus every element of $Fix(\tilde{f})$ projects into $\mathcal{N} \subset K$.

On the other hand, some component \tilde{D}^* of the the full pre-image of D in \tilde{S} is a complementary component of \tilde{W}^*_l . Lemma 3.9 implies that there exists $\tilde{x} \in \text{Fix}(\tilde{f})$ whose projected image $x \in \text{Fix}(f)$ is not Nielsen equivalent rel K to an element of K. This contradiction completes the proof that W has maximal elements.

The hypotheses of Lemma 2.7 and Lemma 2.8 of [8] are slightly different than those of this proposition but their proofs carry over without change to this context. Together they imply that up to isotopy rel K there is a unique maximal element W of W and that one may choose W so that ∂W is disjoint from \mathcal{R}' . Let $\mathcal{R}_m = \partial W \cup \mathcal{R}'$. Properties 4.3(2) and 4.3(3-m) follow from the fact that $W \in W$. For 4.3(1), note that g(W) is also a maximal element of W for each $g \in \mathcal{F}$ because $gfg^{-1} = f$. The uniqueness of W up to isotopy rel K implies that W, and hence \mathcal{R}_m , is g-invariant up to isotopy rel K.

Corollary 4.6. Suppose that $\{f_1, \ldots, f_m\}$ generate an abelian subgroup of diffeomorphisms of S. Then there is a compact subset K of $\bigcup_{i=1}^m \operatorname{Fix_c}(f_i)$ that has locally constant Nielsen behavior for each f_i and that is $\operatorname{Fix_c}(f_i)$ -complete for each i and there is a reducing set \mathcal{R} for $\{f_1, \ldots, f_m\}$ relative to K.

Proof. Ley $Y = \bigcup_{i=1}^m \operatorname{Fix}_{\mathbf{c}}(f_i)$. The compact \mathcal{F} -invariant subset $K \subset Y$ produced by Lemma 4.2 applied to Y and $C = \emptyset$ has locally constant multiplicity and is $\operatorname{Fix}_{\mathbf{c}}(f_i)$ -complete for each i. The former implies by Lemma 3.8 that each f_i has locally constant Nielsen behavior rel K. The existence of a reducing set \mathcal{R} for $\{f_1, \ldots, f_m\}$ relative to K now follows from Proposition 4.5.

5 Thurston Normal Form

Our definition of a reducing set \mathcal{R} for $\{f_1, \ldots, f_m\}$ rel K places no restrictions on the components Y of $S \setminus \mathcal{R}$ that intersect K in a finite set. The natural restriction is that the isotopy class of Y relative to K determined by f_l should be irreducible in the sense of Thurston normal form (TNF) for each f_l . In this section we recall the appropriate definitions and a result about Thurston normal form for abelian subgroups. The actual restrictions will be made in the next section. Further details about Thurston normal form can be found, for example, in [11], [5], [10].

Let S be a finitely punctured surface of negative Euler characteristic, perhaps with boundary and let MCG(S) be the mapping class group of S.

Suppose that \mathcal{R}_T is a finite set of disjoint simple closed curves on S, all of whose complementary components S_i have negative Euler characteristic and that $f: S \to S$ is a homeomorphism that setwise preserves \mathcal{R}_T . For each S_i there exists a minimal $n_i \geq 1$ such that f^{n_i} preserves S_i . Let \bar{S}_i be the completion of S_i obtained by compactifying each end of S_i that corresponds to an element of \mathcal{R}_T with a boundary circle. Then $f^{n_i}|S_i$ determines an element $\alpha_i \in \mathrm{MCG}(\bar{S}_i)$ that depends only on the isotopy classes of the elements of \mathcal{R}_T and the isotopy class $\alpha \in \mathrm{MCG}(S)$ determined

by f. We say that \mathcal{R}_T is a TNF-reducing set for f (or α) if each α_i is either represented by a periodic homeomorphism or by a pseudo-Anosov homeomorphism.

Let \mathcal{S} be the set of isotopy classes of simple, closed, essential, non-peripheral curves on S. Then $\mathrm{MCG}(S)$ acts on \mathcal{S} and for each element α of $\mathrm{MCG}(S)$ we define $\mathrm{Per}(\alpha)$ to be the set of elements of \mathcal{S} on which α acts periodically. The intersection number $\gamma \cdot \delta$ of two elements $\gamma, \delta \in \mathcal{S}$ is the minimum, over all representatives c of γ and d of δ , of the cardinality of $c \cap d$. Following [10], we define $\mathrm{Per}_0(\alpha)$ to be those elements γ of $\mathrm{Per}(\alpha)$ such that $\gamma \cdot \delta = 0$ for all $\delta \in \mathrm{Per}(\alpha)$.

Lemma 5.1. $P_0(\alpha)$ is a TNF-reducing set for α .

Proof. We assume notation as above. By construction, $Per_0(\alpha_i) = \emptyset$ for all α_i . Lemma 2.2 of [10] (see also Lemma 2.12 of [8]) asserts that if α_i is not represented by a periodic homeomorphism then $Per(\alpha_i) = \emptyset$. It then follows from the Thurston normal form theorem that α_i is represented by a pseudo-Anosov homeomorphism.

Corollary 5.2. If A is an abelian subgroup of MCG(S) then $\bigcup_{\alpha \in A} Per_0(\alpha)$ is a finite A-invariant subset of S that is represented by disjoint simple closed curves and is a TNF-reducing set for each $\alpha \in A$.

Proof. If $\alpha, \beta \in A$ then $\beta(\operatorname{Per}_0(\alpha)) = \operatorname{Per}_0(\alpha)$ since the action of β on \mathcal{S} preserves $\operatorname{Per}(\alpha)$ and intersection numbers. It follows that $\operatorname{Per}_0(\alpha) \subset \operatorname{Per}(\beta)$ and hence that $\gamma \cdot \delta = 0$ for all $\gamma \in \operatorname{Per}_0(\alpha)$ and $\delta \in \operatorname{Per}_0(\beta)$. This proves that $\bigcup_{\alpha \in A} \operatorname{Per}_0(\alpha)$ is a finite subset of \mathcal{S} and since it contains a TNF-reducing set for each $\alpha \in A$ it is a TNF-reducing set for each $\alpha \in A$.

Remark 5.3. It is easy to see that $\operatorname{Per}_0(\alpha)$ is the minimal TNF-reducing set for α in the sense that every element of any TNF-reducing set for α is an element of $\operatorname{Per}_0(\alpha)$. It follows that $\bigcup_{\alpha \in A} \operatorname{Per}_0(\alpha)$ is a minimal subset of $\mathcal S$ that is a TNF-reducing set for every $\alpha \in A$. This minimal subset has a canonical characterization which we now describe. Define $\operatorname{Per}(A)$ to be those elements of $\mathcal S$ which are in $\operatorname{Per}(\alpha)$ for all $\alpha \in A$ and $\operatorname{Per}_0(A)$ to be those elements $\gamma \in \operatorname{Per}(A)$ such that $\gamma \cdot \delta = 0$ for all $\delta \in \operatorname{Per}(A)$. Since for all $\alpha, \beta \in A$ the set $\operatorname{Per}_0(\alpha)$ is finite and $\beta(\operatorname{Per}_0(\alpha)) = \operatorname{Per}_0(\alpha)$ it is clear that $\operatorname{Per}_0(\alpha) \subset \operatorname{Per}_0(A)$ for every $\alpha \in A$. Therefore $\operatorname{Per}_0(A)$ is a TNF-reducing set for every $\alpha \in A$. It is straightforward to show $\operatorname{Per}_0(A)$ is also minimal so $\bigcup_{\alpha \in A} \operatorname{Per}_0(\alpha) = \operatorname{Per}_0(A)$.

6 Proof of Theorem 1.1

We assume throughout this section that \mathcal{F} is an abelian subgroup of $\mathrm{Diff}^+(S)_0$ with rank k.

Definition 6.1. A subset Γ of \mathcal{F} will be called k-spanning if it has cardinality at least k and if every subset Γ_0 of cardinality k generates a finite index subgroup of \mathcal{F} .

Equivalently Γ is k-spanning if every subset Γ_0 of cardinality k is a basis of the vector space \mathbb{R}^k determined by the identification of \mathcal{F} with \mathbb{Z}^k .

Lemma 6.2. Given $m \geq k$ there is a subset $\Gamma \subset \mathbb{Z}^k$ of cardinality m such that Γ is k-spanning.

Proof. The desired property is equivalent to the requirement that Γ contains a set of generators and that any subset of $\Gamma_0 \subset \Gamma$ of cardinality k is linearly independent in \mathbb{R}^k . We induct on m. For m = k take Γ a set of generators of \mathbb{Z}^k . Given a Γ_m of size m with the desired property let $X \subset \mathbb{R}^k$ be the union of the spans of all subsets of Γ_m with cardinality k-1. Then X is a union of a finite number of (k-1)-dimensional subspaces. Let v be an element of \mathbb{Z}^k which is not in X and define $\Gamma_{m+1} := \Gamma_m \cup \{v\}$. If $\Gamma_0 = \{v, v_1, \dots v_{k-1}\}$ then v is not a linear combination of $\{v_1, \dots v_{k-1}\}$ so Γ_0 is linearly independent and Γ_{m+1} has the desired property.

We establish notation for the remainder of this section as follows. \mathcal{F}_m is a finite index subgroup of \mathcal{F} generated by a k-spanning set $\{f_1, \ldots, f_m\} \subset \mathcal{F}$. K is a compact set that has locally constant Nielsen behavior with respect to f_i for $1 \leq i \leq m$. \mathcal{R} is a reducing set for $\{f_1, \ldots, f_m\}$ relative to K.

Each $f \in \mathcal{F}_m$ is isotopic rel K to a homeomorphism F that preserves \mathcal{R} and so preserves each complementary component of \mathcal{R} . Suppose that Y is such a component, $K_Y = K \cap Y$ is finite and that \bar{Y} is the completion of Y obtained by compactifying each end of Y that corresponds to an element of \mathcal{R} with a boundary circle. The restriction F|Y determines an element $[f|\bar{Y}]$ of the mapping class group $\mathcal{G}_{\bar{Y}}$ rel K_Y that depends only on f and not the choice of F. The assignment $f \mapsto [f|\bar{Y}]$ defines a homomorphism $\phi_Y : \mathcal{F}_m \to \mathcal{G}_{\bar{Y}}$.

Definition 6.3. Assume notation as above. \mathcal{R} is TNF-complete for $\{f_1, \ldots, f_m\}$ if $[f_i|\bar{Y}]$ is either periodic or pseudo-Anosov for each f_i and for each component Y with finite K_Y . If \mathcal{R} is not TNF-complete then it is TNF-incomplete.

Lemma 6.4. Any reducing set \mathcal{R} for $\{f_1, \ldots, f_m\}$ relative to K extends to a TNF-complete reducing set for $\{f_1, \ldots, f_m\}$ relative to K.

Proof. Suppose that Y is a complementary component of \mathcal{R} and that K_Y is finite. Define $\phi_Y : \mathcal{F}_m \to \mathcal{G}_{\bar{Y}}$ as above. Since $\phi_Y(\mathcal{F}_m)$ is an abelian subgroup of $\mathcal{G}_{\bar{Y}}$, by Corollary 5.2 there is a finite set R_Y of disjoint non-parallel, non-peripheral, simple closed curves in $Y \setminus K_Y$ so that each element $\phi_Y(\mathcal{F}_m)$ is represented by a homeomorphism of Y whose restriction to each component of the complement of R_Y is either periodic or pseudo-Anosov. Taking the union of \mathcal{R} with R_Y for all such choices of Y produces the desired TNF-complete reducing set.

We assume throughout that \mathcal{R} is TNF-complete and that no proper subset of \mathcal{R} is a TNF-complete reducing set.

Let \mathcal{E} be the subset of \mathcal{R} consisting of those elements that are essential in S. Since $\chi(S) < 0$ equals the sum of the Euler characteristics of the components of the complement of \mathcal{E} in S there must be such a component with negative Euler characteristic. In fact if $\{N_i\}_{i=1}^r$ is the set of closures of those components of $S \setminus \mathcal{E}$ which have negative Euler characteristic then $1 \leq r \leq -\chi(S)$. The remaining components of $S \setminus \mathcal{E}$ are annuli and we let $\{A_j\}_{i=1}^s$ be those annuli that share a boundary component with some N_i . Three distinct A_j cannot be parallel in S so s is bounded by $2 - \chi(S)$, which is twice the genus of S and hence twice the maximum cardinality of a set of disjoint essential non-parallel curves in S. Let $S_- = (\bigcup_{i=1}^r N_i) \cup (\bigcup_{j=1}^s A_j)$, $S_0 = S \setminus S_-$, $K_- = K \cap S_-$ and $K_0 = K \cap S_0$. The number $|S_-|$ of subsurfaces N_i and A_j that compose S_- is bounded above by $2 - 2\chi(S)$.

Before beginning the formal proof of Theorem 1.1, we prove it in a pair of special cases.

Lemma 6.5. If K is $\operatorname{Fix}_{\mathsf{c}}(f_l)$ -complete then $\operatorname{Fix}_{\mathsf{c}}(f_l) \cap K_- \neq \emptyset$.

Proof. To simplify notation we write f_l simply as f. We assume that $\operatorname{Fix}_{\mathbf{c}}(f) \cap K_- = \emptyset$ and argue to a contradiction. Let $K_0 = K \setminus K_-$. The first step in the proof is to choose g isotopic to f and satisfying the following properties.

- 1. f agrees with g on the complement of $\bigcup A_j$.
- 2. The restriction of g to each A_i has a periodic orbit P_i .
- 3. Fix(f) = Fix(g) and $Fix_c(f) = Fix_c(g)$.
- 4. If $K' = K_0 \bigcup P_j$ then every point of $\operatorname{Fix}_{\mathbf{c}}(g)$ is Nielsen equivalent rel K' to a point of K'.

Choose an f recurrent point p_j in $K \cap \operatorname{int}(A_j)$. If p_j is periodic, let $P_j = \operatorname{orb}(p_j)$ and let $f|A_j = g|A_j$. Note that if p_j is fixed then it is not in $\operatorname{Fix}_c(g)$. Otherwise, perturb f in a neighborhood W_j of p_j in A_j that is disjoint from its f-image so that p_j becomes periodic with period greater than 1 and let $P_j = \operatorname{orb}(p_j)$. Properties (1) through (3) are satisfied.

The fact that we can achieve Property 4 requires justification. For each $x \in \operatorname{Fix_c}(f)$ choose neighborhoods $U_x \subset V_x$ as in the definition of locally constant Nielsen behavior rel K. Since $\operatorname{Fix_c}(f)$ is compact there exist x_1, \ldots, x_m such that $\operatorname{Fix_c}(f) \subset \bigcup_{l=1}^m U_{x_l}$. Let β_l be an f-Nielsen arc rel K connecting x_l to some element of $\operatorname{Fix_c}(f) \cap K$. The existence of β_l follows from our assumption that K is $\operatorname{Fix_c}(f_l)$ -complete. If the W_j 's are sufficiently small then $\bigcup W_j$ is disjoint from each V_{x_l} and the homotopy from $f(\beta_l)$ to β_l is rel $K \cup (\bigcup W_j)$. For $x \in U_{x_l} \cap \operatorname{Fix_c}(f)$, let β_x be an f-Nielsen arc that is the concatenation of an f-Nielsen arc in U_{x_l} connecting x to x_l with β_l . Then β_x is an f-Nielsen arc rel $K \cup (\bigcup W_j)$ and so is a g-Nielsen arc rel K'.

Let \mathcal{R}_T be a minimal set of reducing curves (with respect to Thurston normal form) for the element $[g|S_-]$ determined by g in the mapping class group of S_- relative to the finite set $\cup_j P_j$. After an isotopy we may assume that each essential element of \mathcal{R}_T

is a component of some ∂N_i . There is an isotopy rel K' from g to a homeomorphism G that preserves S_- , preserves \mathcal{R}_T and that is the identity on each N_i .

For each A_j , let X_j be the complement in A_j of the disks bounded by the inessential elements, if any, of \mathcal{R}_T that are contained in A_j .

We claim that $\bigcup_i \partial N_i \subset \mathcal{R}_T$. If this fails, then there exists adjacent N_i and X_j such that $G|(N_i \cup X_j)$ is the identity. If $A_j = X_j$ then $P_j = \{p_j\}$ is a single point $p_j \in \operatorname{Fix}_{\operatorname{c}}(G) \cap K_-$. Since G is isotopic to g rel P_j , $p_j \in \operatorname{Fix}_{\operatorname{c}}(g) = \operatorname{Fix}_{\operatorname{c}}(f)$ in contradiction to the assumption that $\operatorname{Fix}_{\operatorname{c}}(f) \cap K_- = \emptyset$. If $A_j \neq X_j$ then $A_j \setminus X_j$ is a single disk D_j that contains P_j contradicting property (4) above and Lemma 3.9 (applied with K' replacing K) which implies the existence of an element of $\operatorname{Fix}_{\operatorname{c}}(g)$ that is not g-Nielsen equivalent rel K' to any element of K'. This completes the proof that $\bigcup_i \partial N_i \subset \mathcal{R}_T$.

It is well known that N_i is a complete Nielsen class for $G|S_-$ relative to $\cup P_j$ and that the action of G on $\pi_1(X_j, \partial X_j)$ does not fix any element determined by an arc in X_j connecting distinct components of ∂X_j . It follows that N_i is a complete Nielsen class μ' for G relative to K'. Indeed, if there were a Nielsen path connecting $x \in N_i$ to $y \in S_0$ then its intersection with some X_j would contain an arc with endpoints on distinct components of A_j whose relative homotopy class would be fixed by the action of G.

On the other hand, $\mu' \subset \operatorname{Fix}_{\operatorname{c}}(G)$ has Lefschetz index equal to $\chi(N_i) \neq 0$. Since $\operatorname{Fix}(f) = \operatorname{Fix}(g)$ and since K' is the union of K_0 with a finite set, $K' \cap \operatorname{Fix}(g)$ is an open subset of K'. Lemma 3.8 and Proposition 3.6 imply that μ , the g-Nielsen class rel K' determined by μ' and the isotopy rel K' from g to G, is non-empty. Since $\mu' \subset \operatorname{Fix}_{\operatorname{c}}(G)$, $\mu \subset \operatorname{Fix}_{\operatorname{c}}(g)$. By Property 4, μ contains an element g that is contained in K' and so in not contained in K_i . Since $g \in \operatorname{Fix}(g) \cap K'$ and G is isotopic to g rel K', it follows that g is an element of g. This contradiction completes the proof. \square

For each $M = N_i$ or $M = A_j$ there is a component X_M of $S \setminus \mathcal{R}$ that is either equal to M or is obtained from M by removing disks whose boundaries are elements of $\mathcal{R} \setminus \mathcal{E}$.

Lemma 6.6. Suppose that $M = N_i$ or $M = A_j$ and that $\operatorname{Fix}_c(f_l) \cap K \cap X_M \neq \emptyset$ for at least k distinct values of l. Then $\operatorname{Fix}_c(\mathcal{F}) \neq \emptyset$.

Proof. For simplicity we denote X_M by X. Let \mathcal{F}' be the finite index subgroup of \mathcal{F} generated by the f_l 's satisfying $\operatorname{Fix}_{\operatorname{c}}(f_l) \cap K \cap X \neq \emptyset$. If each $f_l|X$ is isotopic rel K to a homeomorphism whose restriction to X is the identity then $K \cap X \subset \operatorname{Fix}_{\operatorname{c}}(f_l)$ for each l and $K \cap X$ is a non-empty subset of $\operatorname{Fix}_{\operatorname{c}}(\mathcal{F}')$. Lemma 2.2 then completes the proof.

We may therefore assume that some f_l , which we denote simply as f, is not isotopic rel K to a homeomorphism whose restriction to X is the identity. It follows that $X \cap K$ is finite. Since \mathcal{R} is TNF-complete, f|X determines an irreducible relative mapping class. Thus f is isotopic rel K to a homeomorphism F whose restriction

to X is either periodic with period greater than one or is pseudo-Anosov. At least two components of ∂X are contained in \mathcal{E} and so are F-invariant. There is also at least one F-invariant puncture in X. It follows that F|X is not periodic with period greater than one.

It remains to consider the case that F|X is pseudo-Anosov. Each element of \mathcal{F}' is isotopic to a homeomorphism that preserves X; restricting this homeomorphism to X defines a homomorphism $\phi: \mathcal{F}' \to \mathcal{G}$ where \mathcal{G} is the subgroup of the mapping class group of X rel $X \cap K$ consisting of elements that commute with the pseudo-Anosov mapping class $\phi(f)$. It is well known (see for example Lemma 2.10 of [8]) that \mathcal{G} is virtually cyclic. The subgroup $\mathcal{F}_0 = \phi^{-1}(\langle \phi(f) \rangle)$ has finite index in \mathcal{F} . Moreover, $K \cap \operatorname{Fix}_{\mathbf{c}}(f) \cap X \subset \operatorname{Fix}_{\mathbf{c}}(\mathcal{F}_0)$. Lemma 2.2 completes the proof.

We are now ready for the general case.

Proof of Theorem 1.1 Let $m = 3k(2 - 2\chi(S)) \geq 3k|S_-|$. Corollary 4.6 implies the existence of K and \mathcal{R} satisfying our standing hypothesis and with the additional feature that K is $\operatorname{Fix}_{\mathbf{c}}(f_l)$ -complete for all f_l . Lemma 6.5 implies that for each l there exists $M = N_i$ or $M = A_j$ such that $\operatorname{Fix}_{\mathbf{c}}(f_l) \cap K \cap M \neq \emptyset$. If there exists a single such M and at least k values of l such that $\operatorname{Fix}_{\mathbf{c}}(f_l) \cap M \cap K \neq \emptyset$ then Lemma 6.6 implies that $\operatorname{Fix}_{\mathbf{c}}(\mathcal{F}) \neq \emptyset$ and we are done. We may therefore assume that there are at least $2k|S_-|$ distinct values of l such that $\operatorname{Fix}_{\mathbf{c}}(f_l) \cap K \cap X_M = \emptyset$ for all $M = N_i$ and all $M = A_j$. After reordering the f_l 's we may assume that $\{f_1, \ldots, f_{m'}\}$ have this property where $m' = 2k|S_-|$.

Let K' be the subset of K obtained by removing $K \cap X_M$ for all M. Then K' has locally constant Nielsen behavior and is $\operatorname{Fix}_c(f_l)$ complete for $1 \leq l \leq m'$. Let \mathcal{R}' be a TNF-complete normal form for $\{f_l\}$ that contains \mathcal{R} . (Removing punctures may create new finitely punctured components of $S \setminus \mathcal{R}$ and so may cause \mathcal{R} to be TNF-incomplete with respect to K'.) Define subsurfaces N'_i , A'_j and S'_- using K' and \mathcal{R}' . Note that $S'_- \subset S_-$ and that each non-disk component of $S'_- \setminus \mathcal{R}'$ is contained in a non-disk component of $S_- \setminus \mathcal{R}$. It follows that $K' \cap X_{M'} = \emptyset$ for all $M' = N'_i$ and all $M' = A'_j$.

Since $m' = 2k|S_-|$ there exists a single M' and at least k distinct values of l such that $K \cap \operatorname{Fix}_{\mathbf{c}}(f_l) \cap D_l \neq \emptyset$ for some component D_l of $M \setminus X_M$. The fact that \mathcal{R}' is TNF-complete implies that each $f_l|X_M$ is isotopic rel K' to a homeomorphism F_l whose restriction to X_M is either the identity, is periodic with period greater than one or is pseudo-Anosov.

If each $F_l|X_M$ is the identity then $K' \cap M \cap \operatorname{Fix}(f_l) \subset \operatorname{Fix}_{\operatorname{c}}(f_l)$ for each l and Proposition 2.3 then completes the proof. We may therefore assume that some F_l , which we denote simply as F, is either periodic with period greater than one or is pseudo-Anosov. Every component of ∂M and every component of ∂X_M bounding a disk that intersects $K \cap \operatorname{Fix}(f)$ is F-invariant. Thus X_M has at least three invariant boundary components and $F|X_M$ is not periodic with period greater than one.

It remains to consider the case that $F|X_M$ is pseudo-Anosov. As in the proof of Lemma 6.6, there is a finite index subgroup \mathcal{F}_0 of \mathcal{F} whose image in the mapping

class group of X_M rel K' is the cyclic group generated by $[F|X_M]$. Thus each disk component of $M \setminus X_M$ that has non-trivial intersection with $K \cap \operatorname{Fix}_{\mathbf{c}}(f)$ satisfies the hypotheses of Proposition 2.3, which then completes the proof.

References

- [1] C. Bonatti, Un point fixe common pour des diffeomorphisms commutants de S² Ann. Math **129** (1989), 61–79
- [2] C. Bonatti, Diffomorphismes commutants des surfaces et stabilit des fibrations en tores. Topology 29 (1990), no.2 205–209
- [3] **R. F. Brown** The Lefschetz fixed point theorem. Scott, Foresman and Co., Glenview, Ill.-London (1971) vi+186 pp.
- [4] **A. Dold,** Fixed point index and fixed point theorem for Euclidean neighborhood retracts. Topology 4 (1965), 1–8
- [5] A. Fathi, F. Laudenbach and V. Poenaru, Travaux de Thurston sur les surfaces, Asterisque (1979) 66–67
- [6] S. Firmo A note on commuting diffeomorphisms of surfaces. Nonlinearity 18 (2005), n0.4 1511–1526
- [7] J. Franks and M. Handel, Periodic points of Hamiltonian surface diffeomorphisms, Geom. Topol. 7 (2003) 713–756
- [8] J. Franks, M. Handel and K. Parwani Fixed Points of abelian actions on S^2 . preprint
- [9] M. Handel, Commuting homeomorphisms of S^2 , Topology 31 (1992) 293–303
- [10] M. Handel and W. Thurston New Proofs of Some Results of Nielsen, Adv. in Math. 56 (1985) 173–191
- [11] **W. Thurston,** On the geometry and dynamics of diffeomorphisms of surfaces, Bull. Amer. Math. Soc. **19** (1988) 417–431