Character sums with Beatty sequences on Burgess-type intervals

WILLIAM D. BANKS Department of Mathematics University of Missouri Columbia, MO 65211 USA bbanks@math.missouri.edu

IGOR E. SHPARLINSKI Department of Computing Macquarie University Sydney, NSW 2109, Australia igor@ics.mq.edu.au

July 5, 2021

Abstract

We estimate multiplicative character sums taken on the values of a non-homogeneous Beatty sequence $\{\lfloor \alpha n + \beta \rfloor : n = 1, 2, ...\}$, where $\alpha, \beta \in \mathbb{R}$, and α is irrational. Our bounds are nontrivial over the same short intervals for which the classical character sum estimates of Burgess have been established.

2000 Mathematics Subject Classification: 11B50, 11L40, 11T24

1 Introduction

For fixed $\alpha, \beta \in \mathbb{R}$, the corresponding *non-homogeneous Beatty sequence* is the sequence of integers defined by

$$\mathcal{B}_{\alpha,\beta} = \left(\lfloor \alpha n + \beta \rfloor \right)_{n=1}^{\infty},$$

where $\lfloor x \rfloor$ denotes the greatest integer $\leq x$ for every $x \in \mathbb{R}$. Beatty sequences arise in a variety of apparently unrelated mathematical settings, and because of their versatility, the arithmetic properties of these sequences have been extensively explored in the literature; see, for example, [1, 6, 15, 16, 19, 22] and the references contained therein.

In this paper, we study character sums of the form

$$S_k(\alpha, \beta, \chi; N) = \sum_{n \leq N} \chi(\lfloor \alpha n + \beta \rfloor),$$

where α is *irrational*, and χ is a *non-principal character modulo* k. In the special case that k = p is a prime number, the sums $S_p(\alpha, \beta, \chi; N)$ have been previously studied and estimated nontrivially for $N \ge p^{1/3+\varepsilon}$, where $\varepsilon > 0$; see [2, 3].

Here, we show that the approach of [1] (see also [6, 18]), combined with a bound on sums of the form

$$U_k(t,\chi;M_0,M) = \sum_{M_0 < m \leqslant M} \chi(m) \mathbf{e}(tm) \qquad (t \in \mathbb{R}),$$
(1)

where $\mathbf{e}(x) = \exp(2\pi i x)$ for all $x \in \mathbb{R}$, yields a nontrivial bound on the sums $S_k(\alpha, \beta, \chi; N)$ for all sufficiently large N (see Theorem 4.1 below for a precise statement). In particular, in the case that k = p is prime, we obtain a nontrivial bound for all $N \ge p^{1/4+\varepsilon}$, which extends the results found in [2, 3].

It has recently been shown in [5] that for a prime p the least positive quadratic non-residue modulo p among the terms of a Beatty sequence is of size at most $p^{1/(4e^{1/2})+o(1)}$, a result which is complementary to ours. However, the underlying approach of [5] is very different and cannot be used to bound the sums $S_k(\alpha, \beta, \chi; N)$.

We remark that one can obtain similar results to ours by using bounds for double character sums, such as those given in [13]. The approach of this paper, however, which dates back to [1], seems to be more general and can be used to estimate similar sums with many other arithmetic functions f(m) provided that appropriate upper bounds for the sums

$$V(t, f; M_0, M) = \sum_{M_0 < m \leq M} f(m) \mathbf{e}(tm) \qquad (t \in \mathbb{R})$$

are known. Such estimates have been obtained for the characteristic functions of primes and of smooth numbers (see [11] and [12], respectively), as well as for many other functions. Thus, in principle one can obtain asymptotic formulas for the number of primes or smooth numbers in a segment of a Beatty sequence (in the case of smooth numbers, this has been done in [4] by a different method).

Acknowledgements. The authors would like to thank Moubariz Garaev for several fruitful discussions. This work began during a pleasant visit by W. B. to Macquarie University; the support and hospitality of this institution are gratefully acknowledged. During the preparation of this paper, I. S. was supported in part by ARC grant DP0556431.

2 Notation

Throughout the paper, the implied constants in the symbols O and \ll may depend on α and ε but are absolute otherwise. We recall that the notations U = O(V) and $U \ll V$ are equivalent to the assertion that the inequality $|U| \leq cV$ holds for some constant c > 0.

We also use the symbol o(1) to denote a function that tends to 0 and depends only on α and ε . It is important to note that our bounds are uniform with respect all of the involved parameters other than α and ε ; in particular, our bounds are uniform with respect to β . In particular, the latter means that our bounds also apply to the shifted sums of the form

$$\sum_{M+1\leqslant n\leqslant M+N}\chi(\lfloor\alpha n+\beta\rfloor)=\sum_{n\leqslant N}\chi(\lfloor\alpha n+\alpha M+\beta\rfloor),$$

and these bounds are uniform for all integers M.

In what follows, the letters m and n always denote non-negative integers unless indicated otherwise.

We use $\lfloor x \rfloor$ and $\{x\}$ to denote the greatest integer $\leq x$ and the fractional part of x, respectively.

Finally, recall that the *discrepancy* D(M) of a sequence of (not necessarily distinct) real numbers $a_1, \ldots, a_M \in [0, 1)$ is defined by

$$D(M) = \sup_{\mathcal{I} \subseteq [0,1)} \left| \frac{V(\mathcal{I}, M)}{M} - |\mathcal{I}| \right|,$$
(2)

where the supremum is taken all subintervals $\mathcal{I} = (c, d)$ of the interval [0, 1), $V(\mathcal{I}, M)$ is the number of positive integers $m \leq M$ such that $a_m \in \mathcal{I}$, and $|\mathcal{I}| = d - c$ is the length of \mathcal{I} .

3 Preliminaries

It is well known that for every irrational number α , the sequence of fractional parts $\{\alpha\}, \{2\alpha\}, \{3\alpha\}, \ldots$, is uniformly distributed modulo 1 (for instance, see [17, Example 2.1, Chapter 1]). More precisely, let $D_{\alpha,\beta}(M)$ denote the discrepancy of the sequence $(a_m)_{m=1}^M$, where

$$a_m = \{\alpha m + \beta\} \qquad (m = 1, 2, \dots, M).$$

Then, we have:

Lemma 3.1. Let α be a fixed irrational number. Then, for all $\beta \in \mathbb{R}$ we have

$$D_{\alpha,\beta}(M) \leq 2D_{\alpha,0}(M) = o(1) \qquad (M \to \infty),$$

where the function implied by o(1) depends only on α .

When more information about α is available, the bound of Lemma 3.1 can be made more explicit. For this, we need to recall some familiar notions from the theory of *Diophantine approximations*.

For an irrational number α , we define its *type* τ by the relation

$$\tau = \sup \Big\{ \vartheta \in \mathbb{R} : \liminf_{q \to \infty, \ q \in \mathbb{Z}^+} q^\vartheta \, \|\alpha q\| = 0 \Big\}.$$

Using Dirichlet's approximation theorem, it is easy to see that $\tau \ge 1$ for every irrational number α . The celebrated theorems of Khinchin [14] and of Roth [20] assert that $\tau = 1$ for almost all real numbers α (with respect to Lebesgue measure) and all algebraic irrational numbers α , respectively; see also [7, 21].

The following result is taken from [17, Theorem 3.2, Chapter 2]:

Lemma 3.2. Let α be a fixed irrational number of type $\tau < \infty$. Then, for all $\beta \in \mathbb{R}$ we have

 $D_{\alpha,\beta}(M) \leqslant M^{-1/\tau+o(1)} \qquad (M \to \infty),$

where the function implied by o(1) depends only on α .

Next, we record the following property of type:

Lemma 3.3. If α is an irrational number of type $\tau < \infty$ then so are α^{-1} and $a\alpha$ for any integer $a \ge 1$.

Finally, we need the following elementary result, which describes the set of values taken by the Beatty sequence $\mathcal{B}_{\alpha,\beta}$ in the case that $\alpha > 1$:

Lemma 3.4. Let $\alpha > 1$. An integer m has the form $m = \lfloor \alpha n + \beta \rfloor$ for some integer n if and only if

$$0 < \{\alpha^{-1}(m - \beta + 1)\} \leq \alpha^{-1}.$$

The value of n is determined uniquely by m.

Proof. It is easy to see that an integer m has the form $m = \lfloor \alpha n + \beta \rfloor$ for some integer n if and only if the inequalities

$$\frac{m-\beta}{\alpha} \leqslant n < \frac{m-\beta+1}{\alpha}$$

hold, and since $\alpha > 1$ the value of n is determined uniquely.

4 Character Sums

For every real number $\varepsilon > 0$ and integer $k \ge 1$, we put

$$B_{\varepsilon}(k) = \begin{cases} k^{1/4+\varepsilon} & \text{if } k \text{ is prime;} \\ k^{1/3+\varepsilon} & \text{if } k \text{ is a prime power;} \\ k^{3/8+\varepsilon} & \text{otherwise.} \end{cases}$$
(3)

Theorem 4.1. Let $\alpha > 0$ be a fixed irrational number, and let $\varepsilon > 0$ be fixed. Then, uniformly for all $\beta \in \mathbb{R}$, all non-principal multiplicative characters χ modulo k, and all integers $N \ge B_{\varepsilon}(k)$, we have

$$S_k(\alpha, \beta, \chi; N) = o(N) \qquad (k \to \infty),$$

where the function implied by o(N) depends only on α and ε .

Proof. We can assume that $\varepsilon < 1/10$, and this implies that $B_{\varepsilon}(k) \leq k^{2/5}$ in all cases. Observe that it suffices to prove the result in the case that $B_{\varepsilon}(k) \leq N \leq k^{1/2}$. Indeed, assuming this has been done, for any $N > k^{1/2}$ we put $N_0 = \lfloor k^{9/20} \rfloor$ and $t = \lfloor N/N_0 \rfloor$; then, since $B_{\varepsilon}(k) \leq N_0 \leq k^{1/2}$ we have

$$S_k(\alpha, \beta, \chi; N) = \sum_{j=0}^{t-1} \sum_{n \leq N_0} \chi(\lfloor \alpha(n+jN_0) + \beta \rfloor) + \sum_{tN_0 < n \leq N} \chi(\lfloor \alpha n + \beta \rfloor)$$
$$= \sum_{j=0}^{t-1} S_k(\alpha, \beta + \alpha j N_0, \chi; N_0) + O(N_0)$$
$$= o(tN_0) + O(Nk^{-1/20}) = o(N) \qquad (k \to \infty)$$

using the fact that our bounds are uniform with respect to β .

We first treat the case that $\alpha > 1$. Put $\gamma = \alpha^{-1}$, $\delta = \alpha^{-1}(1 - \beta)$, $M_0 = \lfloor \alpha + \beta - 1 \rfloor$, and $M = \lfloor \alpha N + \beta \rfloor$. From Lemma 3.4 we see that

$$S_k(\alpha, \beta, \chi; N) = \sum_{\substack{M_0 < m \leqslant M\\ 0 < \{\gamma m + \delta\} \leqslant \gamma}} \chi(m) = \sum_{\substack{M_0 < m \leqslant M}} \chi(m) \, \psi(\gamma m + \delta), \qquad (4)$$

where $\psi(x)$ is the periodic function with period one for which

$$\psi(x) = \begin{cases} 1 & \text{if } 0 < x \leqslant \gamma; \\ 0 & \text{if } \gamma < x \leqslant 1. \end{cases}$$

By a classical result of Vinogradov (see [23, Chapter 2, Lemma 2]) it is known that for any Δ such that

$$0 < \Delta < \frac{1}{8}$$
 and $\Delta \leqslant \frac{1}{2} \min\{\gamma, 1 - \gamma\},$

there is a real-valued function $\psi_{\Delta}(x)$ with the following properties:

- $\psi_{\Delta}(x)$ is periodic with period one;
- $0 \leq \psi_{\Delta}(x) \leq 1$ for all $x \in \mathbb{R}$;
- $\psi_{\Delta}(x) = \psi(x)$ if $\Delta \leq x \leq \gamma \Delta$ or $\gamma + \Delta \leq x \leq 1 \Delta$;

• $\psi_{\Delta}(x)$ can be represented as a Fourier series

$$\psi_{\Delta}(x) = \gamma + \sum_{j=1}^{\infty} \left(g_j \mathbf{e}(jx) + h_j \mathbf{e}(-jx) \right),$$

where the coefficients g_j, h_j satisfy the uniform bound

$$\max\{|g_j|, |h_j|\} \ll \min\{j^{-1}, j^{-2}\Delta^{-1}\} \qquad (j \ge 1).$$

Therefore, from (4) we derive that

$$S_k(\alpha, \beta, \chi; N) = \sum_{M_0 < m \leq M} \chi(m) \psi_\Delta(\gamma m + \delta) + O(V(\mathcal{I}, M_0, M)), \quad (5)$$

where $V(\mathcal{I}, M_0, M)$ denotes the number of integers $M_0 < m \leq M$ such that

$$\{\gamma m + \delta\} \in \mathcal{I} = [0, \Delta) \cup (\gamma - \Delta, \gamma + \Delta) \cup (1 - \Delta, 1)$$

Since $|\mathcal{I}| \ll \Delta$, it follows from Lemma 3.1 and the definition (2) that

$$V(\mathcal{I}, M_0, M) \ll \Delta N + o(N), \tag{6}$$

where the implied function o(N) depends only on α .

To estimate the sum in (5), we insert the Fourier expansion for $\psi_{\Delta}(\gamma m + \delta)$ and change the order of summation, obtaining

$$\sum_{M_0 < m \leq M} \chi(m) \psi_{\Delta}(\gamma m + \delta) = \gamma U_k(0, \chi; M_0, M)$$

+
$$\sum_{j=1}^{\infty} g_j \mathbf{e}(\delta j) U_k(\gamma j, \chi; M_0, M) + \sum_{j=1}^{\infty} h_j \mathbf{e}(-\delta j) U_k(-\gamma j, \chi; M_0, M),$$

where the sums $U_k(t, \chi; M_0, M)$ are defined by (1).

Since $M - M_0 \ll N$, using the well known results of Burgess [8, 9, 10] on bounds for partial Gauss sums, it follows that for any fixed $\varepsilon > 0$ there exists $\eta > 0$ such that

$$U_k(a/k,\chi;M_0,M) \ll N^{1-\eta} \tag{7}$$

holds uniformly for all $N \ge B_{\varepsilon}(k)$ and all integers *a*; clearly, we can assume that $\eta \le 1/10$.

Put $r = \lfloor \gamma k \rfloor$. Then, for any integer n, we have

$$\mathbf{e}(\gamma n) - \mathbf{e}(rn/k) \ll |\gamma n - rn/k| \leqslant |n|k^{-1},$$

which implies that

$$U_k(\gamma j, \chi; M_0, M) = U_k(rj/k, \chi; M_0, M) + O(N^2 k^{-1}|j|).$$

Using (7) in the case that $|j| \leq k N^{-1-\eta}$ we derive that

$$U_k(\gamma j, \chi; M_0, M) \ll N^{1-\eta},$$

and for $|j| > k N^{-1-\eta}$ we use the trivial bound

$$|U_k(\gamma j, \chi; M_0, M)| \ll N.$$

Consequently,

$$\sum_{M_0 < m \leq M} \chi(m) \psi_{\Delta}(\gamma m + \delta)$$

$$\ll N^{1-\eta} \sum_{j \leq kN^{-1-\eta}} (|g_j| + |h_j|) + N \sum_{j > kN^{-1-\eta}} (|g_j| + |h_j|)$$

$$\ll N^{1-\eta} \sum_{j \leq kN^{-1-\eta}} j^{-1} + N\Delta^{-1} \sum_{j > kN^{-1-\eta}} j^{-2}$$

$$\ll N^{1-\eta} \log k + N^{2+\eta} \Delta^{-1} k^{-1}.$$

Since $N^2 \leq k \leq N^4$, we see that

$$\sum_{M_0 < m \leq M} \chi(m) \psi_{\Delta}(\gamma m - \delta) \ll N^{1-\eta} \log N + N^{\eta} \Delta^{-1}.$$
 (8)

Inserting the bounds (6) and (8) into (5), choosing $\Delta = N^{(\eta-1)/2}$, and taking into account that $0 < \eta \leq 1/10$, we complete the proof in the case that $\alpha > 1$.

If $\alpha < 1$, put $a = \lceil \alpha^{-1} \rceil$ and write

$$S_k(\alpha, \beta, \chi; N) = \sum_{n \leq N} \chi(\lfloor \alpha n + \beta \rfloor)$$

=
$$\sum_{j=0}^{a-1} \sum_{m \leq (N-j)/a} \chi(\lfloor \alpha a m + \alpha j + \beta \rfloor)$$

=
$$\sum_{j=0}^{a-1} S_k(\alpha a, \alpha j + \beta, \chi; (N-j)/a).$$

Applying the preceding argument with the irrational number $\alpha a > 1$, we conclude the proof.

For an irrational number α of type $\tau < \infty$, we proceed as in the proof of Theorem 4.1, using Lemma 3.2 instead of Lemma 3.1, and also applying Lemma 3.3; this yields the following statement:

Theorem 4.2. Let $\alpha > 0$ be a fixed irrational number of type $\tau < \infty$. For every fixed $\varepsilon > 0$ there exists $\rho > 0$, which depends only on ε and τ , such that for all $\beta \in \mathbb{R}$, all non-principal multiplicative characters χ modulo k, and all integers $N \ge B_{\varepsilon}(k)$, we have

$$S_k(\alpha, \beta, \chi; N) \ll Nk^{-\rho}.$$

References

- A. G. Abercrombie, 'Beatty sequences and multiplicative number theory', Acta Arith. 70 (1995), 195–207.
- [2] W. Banks and I. E. Shparlinski, 'Non-residues and primitive roots in Beatty sequences', Bull. Aust. Math. Soc. (to appear).
- [3] W. Banks and I. E. Shparlinski, 'Short character sums with Beatty sequences', *Math. Res. Lett.* (to appear).
- [4] W. Banks and I. E. Shparlinski, 'Prime divisors in Beatty sequences', J. Number Theory (to appear).
- [5] W. Banks, M. Z. Garaev and I. E. Shparlinski, 'Density of non-residues in short intervals', *Preprint* 2006, 1-15.
- [6] A. V. Begunts, 'An analogue of the Dirichlet divisor problem', Moscow Univ. Math. Bull. 59 (2004), no. 6, 37–41.
- [7] Y. Bugeaud, Approximation by algebraic numbers, Cambridge Univ. Press, Cambridge, 2004.
- [8] D. A. Burgess, 'Partial Gaussian sums', Bull. London Math. Soc. 20 (1988), no. 6, 589–592.

- [9] D. A. Burgess, 'Partial Gaussian sums, II', Bull. London Math. Soc. 21 (1989), no. 2, 153–158.
- [10] D. A. Burgess, 'Partial Gaussian sums. III', Glasgow Math. J. 34 (1992), no. 2, 253–261.
- [11] H. Davenport, *Multiplicative number theory*, 2nd ed., Springer-Verlag, New York 1980.
- [12] E. Fouvry and G. Tenenbaum, 'Entiers sans grand facteur premier en progressions arithmétiques', London Math. Soc. 63 (1991), 449–494.
- [13] J. Friedlander and H. Iwaniec, 'Estimates for character sums', Proc. Amer. Math. Soc. 119 (1993), 365–372.
- [14] A. Y. Khinchin, 'Zur metrischen Theorie der diophantischen Approximationen', Math. Z. 24 (1926), no. 4, 706–714.
- [15] T. Komatsu, 'A certain power series associated with a Beatty sequence', Acta Arith. 76 (1996), 109–129.
- [16] T. Komatsu, 'The fractional part of $n\vartheta + \varphi$ and Beatty sequences', J. Théor. Nombres Bordeaux 7 (1995), 387–406.
- [17] L. Kuipers and H. Niederreiter, Uniform distribution of sequences, Wiley-Interscience, New York-London-Sydney, 1974.
- [18] G. S. Lü and W. G. Zhai, 'The divisor problem for the Beatty sequences', Acta Math. Sinica 47 (2004), 1213–1216 (in Chinese).
- [19] K. O'Bryant, 'A generating function technique for Beatty sequences and other step sequences', J. Number Theory 94 (2002), 299–319.
- [20] K. F. Roth, 'Rational approximations to algebraic numbers', Mathematika 2 (1955), 1–20 (see also 'Corrigendum', 2 (1955), 168).
- [21] W. M. Schmidt, *Diophantine Approximation*, Springer-Verlag, Berlin, 1980.
- [22] R. Tijdeman, 'Exact covers of balanced sequences and Fraenkel's conjecture', Algebraic number theory and Diophantine analysis (Graz, 1998), 467–483, de Gruyter, Berlin, 2000.

[23] I. M. Vinogradov, The method of trigonometrical sums in the theory of numbers, Dover Publ., NY, 2004.