#### hep-th/0608152 ITEP-TH-XX/06 HMI-06-XX TCD-MATH-06-XX

## Givental Integral Representation for Classical Groups<sup>1</sup>

A. Gerasimov

Institute for Theoretical and Experimental Physics, Moscow, 117259, Moscow, Russia School of Mathematics, Trinity College, Dublin 2, Ireland Hamilton Mathematics Institute, TCD, Dublin 2, Ireland,

D. Lebedev<sup>2</sup>

Institute for Theoretical and Experimental Physics, 117259, Moscow, Russia l'Institute des Hautes Études Scientifiques, 35 route de Chartres, Bures-sur-Yvette, France,

S. Oblezin<sup>3</sup>

Institute for Theoretical and Experimental Physics, 117259, Moscow, Russia Max-Planck-Institut für Mathematik, Vivatsgasse 7, D-53111 Bonn, Germany,

#### Abstract

We propose an integral representation for wave functions of open Toda chains corresponding to classical series of finite Lie algebras  $B_n$ ,  $C_n$ , and  $D_n$  thus generalizing Givental representation for  $A_n$ . Our approach is based on the connection between Baxter *Q*-operator for the closed Toda chain and Givental representation for a wave function uncovered previously. In this paper we also construct Baxter *Q*-operators for closed Toda chains corresponding to Lie algebras  $B_{\infty}$ ,  $C_{\infty}$ ,  $D_{\infty}$ , affine Lie algebras  $B_n^{(1)}$ ,  $C_n^{(1)}$ ,  $D_n^{(1)}$  and twisted forms  $A_{2n-1}^{(2)}$  and  $A_{2n}^{(2)}$ .

<sup>&</sup>lt;sup>1</sup>To be published in the Proceedings of the satellite ICM 2006 conference: "Integrable systems in Applied Mathematics", Colmenarejo (Madrid, Spain), 7-12 September 2006

<sup>&</sup>lt;sup>2</sup>E-mail: lebedev@ihes.fr

<sup>&</sup>lt;sup>3</sup>E-mail: Sergey.Oblezin@itep.ru

## 1 Introduction

The remarkable integral formula for the common eigenfunctions of  $A_n$  open Toda chain Hamiltonian operators was proposed in [Gi] (see also [JK]). It has many interesting properties. Its extremely simple form allows a graphical representation of the integrand, simple description of the integration contours and of the critical points of the integrand. It was shown in [GKLO] that this integral representation has a natural iterative structure allowing to connect  $A_{n-1}$  and  $A_n$  wave function by simple integral transformation. Moreover it was found that this integral transformation is given by the degenerate version of the Baxter Q-operator realizing quantum Bäcklund transformations in closed Toda chain [PG]. Q-operator was introduce by Baxter as a key tool in solving quantum integrable systems (see [B]). Thus it was demonstrated that Givental representation has a deep relation to the intricate elements of the theory of quantum integrable systems [F].

The puzzling property of the integral representation [Gi] is that this representation is based on a specific properties of the inclusion of the Lie algebras  $\mathfrak{sl}_{n+1} \subset \mathfrak{gl}_{n+1}$ . Its generalization to other classical series  $B_n$ ,  $C_n$ ,  $D_n$  is not obvious. Not surprisingly the generalization of the Baxter Q-operators to the corresponding affine Lie algebras is not known too. In this paper we solve both these problem and generalize the connection between these two subjects established in [GKLO] to all classical series. It turns out that there is an important difference between  $A_n$  case and the case of other classical Lie algebras. The kernel of the integral operator providing recursive construction of the integral representation for  $A_n$  case has simple form of the exponent of the sum of exponents in the natural coordinates. For other classical groups the same type of the iteration operator exists but it relates Toda chain wave functions for different classical series (i.e.  $D_n \to C_{n-1}$  and  $C_n \to D_n$ ). Therefore the integral operator connecting the wave functions for two neighborhood ranks in the same series (i.e.  $D_n \to D_{n-1}$ ) is a composition of the elementary integral operators. Hence the corresponding integral kernel does not have a simple form.

Let us note that in [GKLO] the interpretation of the Givental integral representation of wave function for  $A_n$  open Toda chain in terms representation theory of the corresponding Lie group SL(n + 1) was found. A degenerate version of the Baxter Q-operator also plays a crucial role in this interpretation leading to a particular factorized representation of the group element. In this note we restrict ourselves to the construction of the integral representations and integral kernels for the Baxter Q-operators for classical groups. The group theory interpretation generalizing that of [GKLO] will be given elsewhere [GLO]. To simplify a presentation we consider only the first non-trivial cases by constructing common eigenfunctions of open Toda chains for zero eigenvalues and explicitly check the eigenfunction property for quadratic Hamiltonians for all classical series. Our results in full generality will be published in the extended version of this note [GLO].

The plan of this paper is as follows. In Section 2 we summarize the results of [GKLO]. In Section 3 we construct kernels of the elementary integral operators intertwining Hamiltonian operators for open Toda chains corresponding to different classical finite Lie algebras. In Section 4 combining the results from Section 3 we give explicit integral representations for the wave functions of  $B_n$ ,  $C_n$  and  $D_n$  open Toda chains. In Section 5 we construct elementary integral operators intertwining Hamiltonian operators of closed Toda chains for (in general different) classical series of affine Lie algebras. In Section 6 using appropriate compositions of the elementary intertwining operators we construct integral kernels for Baxter Q-operators for all classical series of (twisted) affine Lie algebras. These results are compatible with the results obtained in Sections 3 and 4. We conclude in Section 7 with a short discussion of the directions of further research.

We were informed by E. Sklyanin that he also have a progress in the construction of Baxter Q-operators for Toda theories.

Acknowledgments: The authors are grateful to S. Kharchev for discussions at the initial stage of this project and to M. Kontsevich for his interest in this work. The research of A. Gerasimov was partly supported by the Enterprise Ireland Basic Research Grant. D. Lebedev is grateful to Institute des Hautes Études Scientifiques for a warm hospitality. S. Oblezin is grateful Max-Planck-Institut für Mathematik for excellent working conditions.

## 2 Recursive structure of Givental representation

In this section we recall a recursive construction of the Givental integral representation discussed in [GKLO].

The solution of a quantum integrable system starts with the finding of the full set of common eigenfunction of the quantum Hamiltonian operators. Thus for the open Toda chain corresponding to Lie algebra  $\mathfrak{gl}_n$  the first nontrivial quantum Hamiltonian is given by

$$H^{\mathfrak{gl}_n}(x) = -\frac{\hbar^2}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^{n-1} g_i e^{x_{i+1} - x_i} \,. \tag{2.1}$$

We shell assume  $g_i$  to be real and positive. In the following we will not distinguish  $\mathfrak{sl}_n$  and  $\mathfrak{gl}_n$ . Toda chains since the connection of all expression for these two cases is very simple.

In [Gi] the following remarkable representation for a common eigenfunction of quantum Hamiltonians of  $\mathfrak{gl}_n$  open Toda chain was proposed

$$\Psi(x_1,\ldots,x_n) = \int_{\Gamma} e^{-\frac{1}{\hbar}\mathcal{F}_n(x)} \bigwedge_{k=1}^{n-1} \bigwedge_{i=1}^k dx_{k,i},$$
(2.2)

where  $x_{n,i} := x_i$ , the function  $\mathcal{F}_n(x)$  is given by

$$\mathcal{F}_n(x) = \sum_{k=1}^{n-1} \sum_{i=1}^k \left( e^{x_{k,i} - x_{k+1,i}} + g_i e^{x_{k+1,i+1} - x_{k,i}} \right), \tag{2.3}$$

and the cycle  $\Gamma$  is a middle dimensional submanifold in the n(n-1)/2- dimensional complex torus with coordinates {exp  $x_{k,i}$ ,  $i = 1, \ldots, k$ ;  $k = 1, \ldots, n-1$ } such that the integral converges. The eigenfunction (2.2) corresponds to a particular eigenvalue

$$H^{\mathfrak{gl}_n}(x)\,\Psi(x_1,\cdots,x_n) = 0 \tag{2.4}$$

In the following we take  $\hbar = 1$  for convenience. Note that this representation can be easily generalized to the case of non-zero eigenvalues [JK].

In [GKLO] among other things the derivation of the integral representation (2.2) using the recursion over the rank n of the Lie algebra  $\mathfrak{gl}_n$  was given. It is easy to see that (2.2) can be represented in the following form

$$\Psi(x_1, \dots, x_n) = \int \bigwedge_{k=1}^{n-1} \bigwedge_{i=1}^k dx_{k,i} \prod_{k=1}^{n-1} Q_{k+1,k}(x_{k+1,1}, \dots, x_{k+1,k+1}; x_{k,1}, \dots, x_{k,k})$$
(2.5)

with the integral kernel

$$Q_{k+1,k}(x_{k+1,i}; x_{k,i}) = \exp\Big\{\sum_{i=1}^{k} e^{x_{k,i} - x_{k+1,i}} + g_i e^{x_{k+1,i+1} - x_{k,i}}\Big\},$$
(2.6)

Here we have  $x_i := x_{n,i}$ . The representation (2.2) then follows from a simple identity

$$H^{\mathfrak{gl}_{k+1}}(x_{k+1,i})Q_{k+1,k}(x_{k+1,i}, x_{k,i}) = Q_{k+1,k}(x_{k+1,i}, x_{k,i})H^{\mathfrak{gl}_k}(x_{k,i})$$
(2.7)

where

$$H^{\mathfrak{gl}_k}(x_i) = -\frac{1}{2} \sum_{i=1}^k \frac{\partial^2}{\partial x_{k,i}^2} + \sum_{i=1}^{k-1} g_i e^{x_{k,i+1} - x_{k,i}} \,.$$
(2.8)

Here and in the following we assume that in the relation similar to (2.7) the Hamiltonian operator on l.h.s. acts on right and the Hamiltonian on r.h.s. acts on the left. Thus the integral operator with the kernel  $Q_{k+1,k}$  intertwines Hamiltonian operators for  $A_{k+1}$  and  $A_k$  open Toda chains.

The integral operator defined by the kernel (2.6) is closely related with a Baxter Qoperator realizing Bäcklund transformations in a closed Toda chain corresponding to affine
Lie algebra  $\widehat{\mathfrak{gl}}_n$ . Baxter Q-operator can be written in the integral form with the kernel [PG]

$$Q^{\widehat{\mathfrak{gl}}_{n}}(x_{i}, y_{i}) = \exp\Big\{\sum_{i=1}^{n} e^{x_{i}-y_{i}} + g_{i}e^{y_{i+1}-x_{i}}\Big\},$$
(2.9)

where  $x_{i+n} = x_i$  and  $y_{i+n} = y_i$ . This operator commutes with the Hamiltonian operators of the closed Toda chain. Thus for example for the quadratic Hamiltonian we have

$$\mathcal{H}^{\widehat{\mathfrak{gl}}_n}(x_i)Q^{\widehat{\mathfrak{gl}}_n}(x_i, y_i) = Q^{\widehat{\mathfrak{gl}}_n}(x_i, y_i)\mathcal{H}^{\widehat{\mathfrak{gl}}_n}(y_i)$$
(2.10)

where quadratic Hamiltonian operator for  $\widehat{\mathfrak{gl}}_n$  closed Toda chain is given by

$$\mathcal{H}^{\hat{\mathfrak{gl}}_n} = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^n g_i e^{x_{i+1} - x_i} \,.$$
(2.11)

where  $x_{i+n} = x_i$ . The recursive operator (2.6) can be obtained from Baxter operator (2.9) in the limit  $g_n \to 0$ .

The main objective of this note is to generalize the representation (2.2), (2.3) to other classical series of finite Lie groups  $B_n$ ,  $C_n$  and  $D_n$ . Quite naturally to get recursive operators one would like to use a degeneration of the corresponding Baxter *Q*-operators of  $B_n^{(1)}$ ,  $C_n^{(1)}$ and  $D_n^{(1)}$  closed Toda chains. But unfortunately it seems that such operators are not known. So in the following we will solve both problems simultaneously by constructing analogs of recursive operators for open Toda chains and Baxter operators for closed Toda chains. Note that the first can be obtained from the second by a degeneration.

Before we give explicit expressions for recursive operators let us explain the main problem in the construction of the recursive operators for classical series different from  $A_n$ . As it was explained in [GKLO] the variables  $x_{k,i}$  in the integral representation have a clear meaning of linear coordinates on Cartan subalgebras of the intermediate Lie algebras entering recursion  $A_n \to A_{n-1} \to \cdots \to A_1$ . This is a consequence of the identity

$$\left(\dim\left(\mathfrak{gl}_{n}\right) - \operatorname{rk}\left(\mathfrak{gl}_{n}\right)\right) - \left(\dim\left(\mathfrak{gl}_{n-1}\right) - \operatorname{rk}\left(\mathfrak{gl}_{n-1}\right)\right) = 2\operatorname{rk}\mathfrak{gl}_{n-1}$$
(2.12)

However for other classical series there is no such simple relation and in general one should expect to find more integration variables in the integral representation then those arising as linear coordinates on intermediate Cartan subalgebras. The resolution of this problem is as follows. Elementary integral operators appear to intertwine Hamiltonians corresponding to Toda Chains for different Lie algebras. The recursive operators are then constructed as appropriate compositions of elementary intertwining operators. Let us stress that in the general case integral kernels of the recursive operators do not have a simple form of an exponent of a sum of exponents but are given by the integrals of such expressions. However the resulting integral representations for  $B_n$ ,  $C_n$  and  $D_n$  open Toda chain eigenfunctions are as close to that for  $A_n$  open Toda chain as one could imagine.

## **3** Elementary Intertwiners for open Toda chains

Let  $\mathfrak{g}$  be a simple Lie algebra,  $\mathfrak{h}$  be a Cartan subalgebra,  $n = \dim \mathfrak{h}$  be the rank of  $\mathfrak{g}$ ,  $R \subset \mathfrak{h}^*$ be the root system, W be the Weyl group. Let us fix a decomposition  $R = R_+ \cup R_-$  of the roots on positive and negative roots. Let  $\alpha_1, \ldots, \alpha_n$  be the bases of simple roots. Let (,) be a W-invariant bilinear symmetric form on  $\mathfrak{h}^*$  normalized so that  $(\alpha, \alpha) = 2$  for a long root. This form provides an identification of  $\mathfrak{h}$  with  $\mathfrak{h}^*$  and thus can be considered as a bilinear form on  $\mathfrak{h}$ . Choose an orthonormal basis  $e = \{e_1, \ldots, e_n\}$  in  $\mathfrak{h}$ . Then for any  $x \in \mathfrak{h}$  one has a decomposition  $x = \sum_{i=1}^n x_i e_i$ . One associates with these data an open Toda chain with the quadratic Hamiltonian

$$H^{R}(x_{i}) = -\frac{1}{2} \sum_{i=1}^{n} \frac{\partial^{2}}{\partial x_{i}^{2}} + \sum_{i=1}^{n} g_{i} e^{\alpha_{i}(x)}$$
(3.1)

with  $g_i$  are non-negative real numbers. For the standard facts on Toda theories corresponding to arbitrary root systems see e.g. [RSTS].

We start with explicit expressions for elementary intertwiner operators for open Toda chains.

### **3.1** $BC \leftrightarrow B$

Let  $e_i \ i = 1, \dots, n$  be an orthonormal basis in  $\mathbb{R}^n$ . Non-reduced root system of type  $BC_n$  can be defined as

$$\alpha_0 = 2e_1, \qquad \alpha_1 = e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad 1 \le i \le n-1$$
 (3.2)

and the corresponding Dynkin diagram is

$$\frac{\alpha_0}{\alpha_1} \iff \alpha_2 \longleftarrow \dots \longleftarrow \alpha_n \tag{3.3}$$

where the first vertex from the left is doubled vertex corresponding to non-reduced root  $\alpha_0 = 2\alpha_1$ .

Quadratic Hamiltonian operator of the corresponding open Toda chain is given by

$$H^{BC_n}(x_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \frac{g_1}{2} \left( e^{x_1} + e^{2x_1} \right) + \sum_{i=1}^{n-1} g_i e^{x_{i+1} - x_i}.$$
(3.4)

Let us stress that the same open Toda chain can be considered as a most general form of  $C_n$  open Toda chain (see e.g. [RSTS], Remark p.61). However in the following we will use the term  $BC_n$  open Toda chain to distinguish it from a more standard  $C_n$  open Toda chain that will be consider below.

A root system of type  $B_n$  can be defined as

$$\alpha_1 = e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad 1 \le i \le n - 1$$
(3.5)

and the corresponding Dynkin diagram is

$$\alpha_1 \longleftarrow \alpha_2 \longleftarrow \ldots \longleftarrow \alpha_n \,. \tag{3.6}$$

Quadratic Hamiltonian operator of the corresponding open Toda chain is given by

$$H^{B_n}(x_i) = -\frac{1}{2} \sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1} + \sum_{i=1}^{n-2} g_i e^{x_{i+1}-x_i}.$$
(3.7)

An elementary operator intertwining open Toda chain Hamiltonians for  $BC_n$  and  $B_{n-1}$  can be written in an integral form with a kernel

$$Q_{BC_n}^{B_{n-1}}(z_1,\ldots,z_n;x_1,\ldots,x_{n-1}) = \exp\left\{g_1e^{z_1} + \sum_{i=1}^{n-1}e^{x_i-z_i} + g_{i+1}e^{z_{i+1}-x_i}\right\}$$
(3.8)

satisfying the following relation

$$H^{BC_n}(z)Q^{B_{n-1}}_{BC_n}(z, x) = Q^{B_{n-1}}_{BC_n}(z, x)H^{B_{n-1}}(x).$$
(3.9)

Similarly an elementary operator intertwining  $B_n$  and  $BC_n$  Hamiltonians has an integral kernel

$$Q_{B_n}^{BC_n}(x_1,\ldots,x_n;z_1,\ldots,z_n) = \exp\Big\{g_1e^{z_1} + \sum_{i=1}^{n-1}\left(e^{x_i-z_i} + g_{i+1}e^{z_{i+1}-x_i}\right) + e^{x_n-z_n}\Big\}.$$
 (3.10)

## **3.2** $C \leftrightarrow D$

A root system of type  $C_n$  can be defined as

 $\alpha_1 = 2e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad 1 \le i \le n - 1,$ (3.11)

and the corresponding Dynkin diagram is

$$\alpha_1 \Longrightarrow \alpha_2 \longrightarrow \ldots \longrightarrow \alpha_n \,. \tag{3.12}$$

Quadratic Hamiltonian operator of the corresponding open Toda chain is given by

$$H^{C_n}(x_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^{n-1} g_i e^{x_{i+1} - x_i} + g_n e^{-2x_n},$$
(3.13)

A root system of type  $D_n$  is

$$\alpha_1 = e_1 + e_2, \qquad \alpha_{i+1} = e_{i+1} - e_i, \quad 1 \le i \le n$$
(3.14)

and the corresponding Dynkin diagram is

Quadratic Hamiltonian operator of the  $D_n$  open Toda chain is

$$H^{D_n}(x_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^{n-1} g_i e^{x_{i+1} - x_i} + g_{n-1} g_n e^{-x_n - x_{n-1}}.$$
(3.16)

An integral operator intertwining  $C_n$  and  $D_n$  Hamiltonians has the kernel

$$Q_{C_n}^{D_n}(z_1,\ldots,z_n;x_1,\ldots,x_n) = \exp\left\{\sum_{i=1}^{n-1} \left(e^{x_i-z_i} + g_{i+1}e^{z_{i+1}-x_i}\right) + e^{x_n-z_n} + g_n e^{-x_n-z_n}\right\}$$
(3.17)

Similarly an integral operator with the kernel

$$Q_{D_n}^{C_{n-1}}(x_1,\ldots,x_n;z_1,\ldots,z_{n-1}) = \exp\left\{\sum_{i=1}^{n-1} \left(e^{z_i-x_i} + g_i e^{x_{i+1}-z_i}\right) + g_n e^{-x_n-z_{n-1}}\right\}$$
(3.18)

intertwines the following  $D_n$  and  $C_{n-1}$  Hamiltonians

$$H^{D_n}(x_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^{n-1} g_i e^{x_{i+1} - x_i} + g_n e^{-x_n - x_{n-1}},$$
(3.19)

$$H^{C_{n-1}}(z_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_i^2} + \sum_{i=1}^{n-1} g_i e^{z_{i+1}-z_i} + 2g_{n-1}g_n e^{-2z_{n-1}}.$$
(3.20)

## 4 Givental representation for wave function

In the previous section explicit expressions for the kernels of elementary intertwining operators were presented. Now the construction of integral representations for eigenfunctions of open Toda chain Hamiltonians is a quite straightforward generalization of  $A_n$  case. Below we provide integral representations for all classical series.

#### **4.1** $B_n$

The eigenfunction for  $B_n$  open Toda chain is given by

$$\Psi^{B_n}(x_1,\ldots,x_n) = \int \bigwedge_{k=1}^{n-1} \bigwedge_{i=1}^k dx_{k,i} \prod_{k=1}^{n-1} Q^{B_k}_{B_{k+1}}(x_{k+1,1},\ldots,x_{k+1,k+1};x_{k,1},\ldots,x_{k,k})$$
(4.1)

where  $x_i := x_{n,i}$  and the kernel  $Q_{B_{k+1}}^{B_k}$  of the integral operator is given by the convolution of the kernels  $Q_{B_{k+1}}^{BC_{k+1}}$  and  $Q_{BC_{k+1}}^{B_k}$ 

$$Q_{B_{k+1}}^{B_k}(x_{k+1,i}; x_{k,j}) = \int \bigwedge_{i=1}^k dz_{k,i} Q_{B_{k+1}}^{BC_{k+1}}(x_{k+1,1}, \dots, x_{k+1,k+1}; z_{k+1,1}, \dots, z_{k+1,k+1}) \times (4.2) \times Q_{BC_{k+1}}^{B_k}(z_{k+1,1}, \dots, z_{k+1,k+1}; x_{k,1}, \dots, x_{k,k})$$

## **4.2** *C*<sub>*n*</sub>

The eigenfunction for  $C_n$  open Toda chain is given by

$$\Psi^{C_n}(z_1,\ldots,z_n) = \int \bigwedge_{k=1}^{n-1} \bigwedge_{i=1}^k dz_{k,i} \prod_{k=1}^{n-1} Q^{C_k}_{C_{k+1}}(z_{k+1,1},\ldots,z_{k+1,k+1};z_{k,1},\ldots,z_{k,k})$$
(4.3)

where  $z_i := z_{n,i}$  and the kernel  $Q_{C_{k+1}}^{C_k}$  of the integral operator is given by a convolution of the kernels  $Q_{C_{k+1}}^{D_{k+1}}$  and  $Q_{D_{k+1}}^{C_k}$ 

$$Q_{C_{k+1}}^{C_k}(z_{k+1,i}; z_{k,j}) = \int \bigwedge_{i=1}^k dx_{k,i} Q_{C_{k+1}}^{D_{k+1}}(z_{k+1,1}, \dots, z_{k+1,k+1}; x_{k+1,1}, \dots, x_{k+1,k}) \times$$

$$\times Q_{D_{k+1}}^{C_k}(x_{k+1,1}, \dots, x_{k+1,k+1}; z_{k,1}, \dots, z_{k,k})$$

$$(4.4)$$

#### **4.3** *D<sub>n</sub>*

The eigenfunction for  $D_n$  open Toda chain is given by

$$\Psi^{D_n}(x_1,\ldots,x_n) = \int \bigwedge_{k=1}^{n-1} \bigwedge_{i=1}^k dx_{k,i} \prod_{k=1}^{n-1} Q_{D_{k+1}}^{D_k}(x_{k+1,1},\ldots,x_{k+1,k+1};x_{k,1},\ldots,x_{k,k})$$
(4.5)

where  $x_i := x_{n,i}$  and the kernel  $Q_{D_{k+1}}^{D_k}$  of the integral operator is given by a convolution of the kernel  $Q_{D_{k+1}}^{C_{k+1}}$  and  $Q_{C_{k+1}}^{D_k}$ 

$$Q_{D_{k+1}}^{D_k}(x_{k+1,i}; x_{k,j}) = \int \bigwedge_{i=1}^k dz_{k,i} Q_{D_{k+1}}^{C_k}(x_{k+1,1}, \dots, x_{k+1,k+1}; z_{k+1,1}, \dots, z_{k+1,k+1}) \times (4.6) \times Q_{C_{k+1}}^{D_k}(z_{k+1,1}, \dots, z_{k+1,k+1}; x_{k,1}, \dots, x_{k,k})$$

#### 4.4 Givental diagrams

In this section we describe graphically the integrands of the integral representations of open Toda chain wave functions for classical series of Lie algebras. This is a generalization of Givental diagram for  $A_n$  introduced in [Gi]. Diagrams for other classical series can be obtained from those for  $A_n$  by a factorization. This factorization is closely related with a realization of classical series of Lie algebras as fixed point subalgebras of the involutions acting on an algebra  $\mathfrak{gl}_N$  for some N (for the explicit form of this relation that will be used below see [DS]). Let us remind the Givental diagram for  $A_N$ . The integrand (2.3) of the integral representation for  $A_{2n-1}$  can be concisely described by the following diagram



where N = 2n. We assign the variables  $x_{k,i}$  to the vertexes and the functions  $e^{y-x}$  to an arrows  $a_{x,y} = (x \longrightarrow y)$  of the diagram (4.7). The potential function  $\mathcal{F}(x_{k,i})$  (see (2.3)) is represented as a sum of the functions over all arrows.

Lie algebra  $D_n$  can be realized as a fixed point subalgebra of  $A_{2n-1}$  using the following involution

$$\iota : \quad X \longmapsto w_0^{-1} X^T w_0, \tag{4.8}$$

where  $w_0 \in \mathfrak{S}_{2n}$  is the longest element of Weyl group  $W(A_{2n-1})$  and  $X^T$  denotes the standard transposition. Correspondingly the diagram for  $D_n$  can be obtain from that for  $A_{2n-1}$  by identification of the variables assigned to the vertexes of  $A_{2n-1}$  diagram

$$w_0: \qquad x_{k,i} \longleftrightarrow -x_{k,k+1-i}. \tag{4.9}$$

The example of the resulting diagram for  $D_4$  is given by



To reconstruct the integrand we assign the term  $e^{z-x}$  to a single arrow  $a_{x,z} = (x \longrightarrow z)$ , and and  $e^{-z-x}$  to a double arrow  $a_{x,z} = (x = z)$ . Then the integrand for the Givental integral representation of the eigenfunction for  $D_4$  open Toda Chain is given by the exponent of the corresponding potential function  $\mathcal{F}(x, z)$  and for the wave function we have

$$\Psi^{D_4}(x_{41}, x_{42}, x_{43}, x_{44}) = \int \bigwedge_{k=1}^3 \bigwedge_{i=1}^k dz_{k,i} dx_{k,i} \cdot e^{\frac{i}{\hbar} \mathcal{F}(x_{k,i}, z_{k,i})}$$
(4.10)

It is easy to see that it is in complete accordance with (4.5).

Following the same reasoning one can readily obtain diagrams for  $C_n$  and  $B_n$ . In the case of  $C_n$  it is a quotient of  $A_{2n-1}$  diagram by the action (4.9) of involution (4.8). According to (4.4) it can also be obtained from the diagram for  $D_{n+1}$  (see (4.4)) by erasing the last row of vertexes and arrows on the right slope. For example, the diagram for  $C_3$  reads



Similarly, the diagram for  $B_n$  can be obtained by factorization of the diagram (4.7) with N = 2n + 1 over the involution (4.8) with the same action (4.9) of  $w_0 \in \mathfrak{S}_{2n+1}$ . For example

the diagram for  $B_3$  has the following form



Here one assigns a function  $e^x$  to an arrow  $a_{*,x} = (\longrightarrow x)$ .

## 5 Elementary intertwiners for closed Toda chains

In this section we generalize the construction of the elementary intertwiners to the classical series of affine Lie algebras. For the theory of affine Lie algebras see [K]. Let us recall the construction for the only known case of  $A_n^{(1)}$  closed Toda chain [PG]. The integral kernel of the Baxter intertwining Q-operator in this case reads

$$Q^{A_n^{(1)}}(x_1,\ldots,x_n;y_i,\ldots,y_n) = \exp\Big\{\sum_{i=1}^{n+1} e^{x_i - y_i} + g_{i+1}e^{y_{i+1} - x_i}\Big\}, \qquad y_{n+2} = y_1 \qquad (5.1)$$

It is easy to see that this operator intertwines the Hamiltonians for  $A_n^{(1)}$  closed Toda chain

$$\mathcal{H}^{A_n^{(1)}}(x_i) = -\frac{1}{2} \sum_{i=1}^{n+1} \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1 - x_{n+1}} + \sum_{i=1}^n g_{i+1} e^{x_{i+1} - x_i},$$
(5.2)

$$\mathcal{H}^{A_n^{(1)}}(y_i) = -\frac{1}{2} \sum_{i=1}^{n+1} \frac{\partial^2}{\partial y_i^2} + g_1 e^{y_1 - y_{n+1}} + \sum_{i=1}^n g_{i+1} e^{y_{i+1} - y_i} \,.$$
(5.3)

Below we give explicit expressions for the kernels of the integral operators intertwining Hamiltonians for other affine Lie algebras of classical type.

# **5.1** $A_{2n}^{(2)} \leftrightarrow BC_{n+1}^{(2)}$

Simple roots of the twisted affine root system  $A_{2n}^{(2)}$  are given in terms of the standard basis  $\{e_i\}$  as follows

$$\alpha_1 = e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad 1 \le i \le n-1 \qquad \alpha_{n+1} = -2e_n$$
(5.4)

and the corresponding Dynkin diagram is given by

 $\alpha_1 \Leftarrow \alpha_2 \leftarrow \ldots \leftarrow \alpha_{n-1} \Leftarrow \alpha_n.$ 

Simple roots of the twisted affine root system  $BC_n^{(2)}$  are as follows

$$\alpha_0 = 2e_1, \quad \alpha_1 = e_1, \quad \alpha_{i+1} = e_{i+1} - e_i, \quad 1 \le i \le n - 1, \quad \alpha_{n+1} = -e_n - e_{n-1} \tag{5.5}$$

and the corresponding Dynkin diagram is given by

The integral operator with the following kernel

$$Q_{A_{2n}^{(2)}}^{BC_{n+1}^{(2)}}(x_i, z_i) = \exp\left\{g_1 e^{z_1} + \sum_{i=1}^n \left(e^{x_i - z_i} + g_{i+1} e^{z_{i+1} - x_i}\right) + \frac{g_{2n+2}}{2} e^{-z_{n+1} - x_n}\right\}$$
(5.7)

intertwines Hamiltonian operators for  $A_{2n}^{\left(2\right)}$  and  $BC_{n+1}^{\left(2\right)}$ 

$$\mathcal{H}^{A_{2n}^{(2)}}(x_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{i+1}-x_i} + 2g_{n+1}g_{n+2} e^{-2x_n}, \qquad (5.8)$$

$$\mathcal{H}^{BC_{n+1}^{(2)}}(z_i) = -\frac{1}{2} \sum_{i=1}^{n+1} \frac{\partial^2}{\partial z_i^2} + \frac{g_1}{2} \left( e^{z_1} + e^{2z_1} \right) + \sum_{i=1}^n g_{i+1} e^{z_{i+1} - z_i} + g_{n+2} e^{-z_{n+1} - z_n}.$$
 (5.9)

**5.2** 
$$A_{2n-1}^{(2)} \leftrightarrow A_{2n-1}^{(2)}$$

Simple roots of the twisted affine root system  $A_{2n-1}^{(2)}$  are given by

$$\alpha_1 = 2e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad 1 \le i \le n-1 \qquad \alpha_{n+1} = -e_n - e_{n-1}, \qquad (5.10)$$
  
and corresponding Dynkin diagram is given by  
$$\alpha_1 \Longrightarrow \alpha_2 \longrightarrow \dots \longrightarrow \alpha_{n-1} \longrightarrow \alpha_n$$

$$\alpha_1 \Longrightarrow \alpha_2 \longrightarrow \dots \longrightarrow \alpha_{n-1} \longrightarrow \alpha_n$$

$$\downarrow$$

$$\alpha_{n+1}$$

The integral operator represented by the following kernel

$$Q_{A_{2n-1}^{(2)}}^{A_{2n-1}^{(2)}}(x_i, z_i) = \exp\left\{g_1 e^{x_1 + z_1} + \sum_{i=1}^{n-1} \left(e^{x_i - z_i} + g_{i+1} e^{z_{i+1} - x_i}\right) + e^{x_n - z_n} + g_{n+1} e^{-x_n - z_n}\right\}$$
(5.11)

intertwines Hamiltonian operators for  $A_{2n-1}^{(2)}$  closed Toda chains with different coupling constants

$$\mathcal{H}^{A_{2n-1}^{(2)}}(x_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + 2g_1 e^{2x_1} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{i+1}-x_i} + g_{n+1} e^{-x_n-x_{n-1}}, \quad (5.12)$$

$$\mathcal{H}^{A_{2n-1}^{(2)}}(z_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_i^2} + g_1 e^{z_1 + z_2} + \sum_{i=1}^{n-1} g_{i+1} e^{z_{i+1} - z_i} + 2g_{n+1} e^{-2z_n}.$$
 (5.13)

# **5.3** $B_n^{(1)} \leftrightarrow BC_{n+1}^{(1)}$

Simple roots of the affine root system  $B_n^{(1)}$  are

$$\alpha_1 = e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad 1 \le i \le n-1 \qquad \alpha_{n+1} = -e_n - e_{n-1}$$
(5.14)

and corresponding Dynkin diagram is given by

$$\alpha_1 \Leftarrow \alpha_2 \leftarrow \dots \leftarrow \alpha_{n-1} \leftarrow \alpha_n$$

$$\uparrow$$

$$\alpha_{n+1}$$

Simple roots of the affine root system  $BC_n^{(1)}$  are

$$\alpha_0 = 2e_1, \quad \alpha_1 = e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad 1 \le i \le n-1 \qquad \alpha_{n+1} = -2e_n \qquad (5.15)$$

and corresponding Dynkin diagram is given by

$$\alpha_0 \iff \alpha_2 \longleftarrow \ldots \longleftarrow \alpha_n \Longleftarrow \alpha_{n+1}$$

The integral operator represented by the following kernel

$$Q_{B_n^{(1)}}^{BC_{n+1}^{(1)}}(x_i, z_i) = \exp\left\{g_1 e^{z_1} + \sum_{i=1}^{n-1} \left(e^{x_i - z_i} + g_{i+1} e^{z_{i+1} - x_i}\right) + e^{x_n - z_n} + g_{n+1} e^{-x_n - z_n}\right\}$$
(5.16)

intertwines Hamiltonians of  $B_n^{(1)}$  and  $BC_{n+1}^{(1)}$  closed Toda chains

$$\mathcal{H}^{B_n^{(1)}}(x_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{i+1}-x_i} + g_{n+1} e^{-x_n-x_{n-1}}, \qquad (5.17)$$

$$\mathcal{H}^{BC_{n+1}^{(1)}}(z_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial z_i^2} + \frac{g_1}{2} \left( e^{z_1} + e^{2z_1} \right) + \sum_{i=1}^{n-1} g_{i+1} e^{z_{i+1}-z_i} - g_{n+1} e^{-2z_n}.$$
 (5.18)

## $\mathbf{5.4} \qquad C^{(1)} \leftrightarrow D^{(1)}$

Simple roots of the affine root system  $C_n^{(1)}$  are

$$\alpha_1 = 2e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad 1 \le i \le n - 1 \qquad \alpha_{n+1} = -2e_n \tag{5.19}$$

and corresponding Dynkin diagram is given by

$$\alpha_1 \Longrightarrow \alpha_2 \longrightarrow \ldots \longleftarrow \alpha_n \Longleftarrow \alpha_{n+1}$$

Simple roots of the affine root system  $D_n^{(1)}$  are

$$\alpha_1 = e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad 1 \le i \le n-1 \qquad \alpha_{n+1} = -e_n - e_{n-1}$$
(5.20)

and corresponding Dynkin diagram is given by



The integral operator with the following kernel

$$Q_{C_n^{(1)}}^{D_{n+1}^{(1)}}(x_i, z_i) = \exp\left\{g_1 e^{x_1 + z_1} + \sum_{i=1}^n \left(e^{x_i - z_i} + g_{i+1} e^{z_{i+1} - x_i}\right) + g_{n+2} e^{-z_{n+1} - x_n}\right\}$$
(5.21)

intertwines Hamiltonian operators for  $C_n^{(1)}$  and  $D_{n+1}^{(1)}$  closed Toda chains

$$\mathcal{H}^{C_n^{(1)}}(x_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + 2g_1 e^{2x_1} + \sum_{i=1}^{n-1} g_{i+1} e^{x_{i+1} - x_i} + 2g_{n+1} g_{n+2} e^{-2x_n}, \quad (5.22)$$

$$\mathcal{H}^{D_{n+1}^{(1)}}(z_i) = -\frac{1}{2} \sum_{i=1}^{n+1} \frac{\partial^2}{\partial z_i^2} + g_1 e^{z_1 + z_2} + \sum_{i=1}^n g_{i+1} e^{z_{i+1} - z_i} + g_{n+2} e^{-z_{n+1} - z_n}.$$
 (5.23)

On the other hand the integral operator with the kernel

$$Q_{D_n^{(1)}}^{C_{n-1}^{(1)}}(x_i, z_i) = \exp\left\{g_1 e^{x_1 + z_1} + \sum_{i=1}^{n-1} \left(e^{z_i - x_i} + g_{i+1} e^{x_{i+1} - z_i}\right) + g_{n+1} e^{-x_n - z_{n-1}}\right\}$$
(5.24)

intertwines Hamiltonian operators for  $D_n^{(1)}$  and  $C_{n-1}^{(1)}$  closed Toda chains

$$\mathcal{H}^{D_n^{(1)}}(x_i) = -\frac{1}{2} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1 + x_2} + \sum_{i=1}^n g_{i+1} e^{x_{i+1} - z_i} + g_{n+2} e^{-x_{n+1} - x_n}, \quad (5.25)$$

$$\mathcal{H}^{C_{n-1}^{(1)}}(z_i) = -\frac{1}{2} \sum_{i=1}^{n-1} \frac{\partial^2}{\partial z_i^2} + 2g_1 e^{2z_1} + \sum_{i=1}^{n-1} g_{i+1} e^{z_{i+1}-z_i} - 2g_{n+1}g_{n+2} e^{-2z_n}.$$
 (5.26)

## 6 Baxter operators

In this section we apply the results of the previous section for the construction Baxter integral Q-operators for all classical series of affine Lie groups. Let us stress that the results of this and the previous Section are in accordance with the results presented in Section 3 and Section 4. The elementary intertwining operators and recursive operators for finite Lie algebras can be obtained from the elementary intertwining operators and Baxter operators for affine Lie algebras by taking appropriate limits  $g_i \to 0$  in (5.7)-(5.24). This generalizes known relation between Baxter operator for  $A_n^{(1)}$  and recursive operators for  $A_n$ . Baxter Q-operator for  $A_{2n}^{(2)}$  can be defined by the integral kernel

$$Q^{A_{2n}^{(2)}}(x_1, \dots, x_n; y_1, \dots, y_n) = \int \bigwedge_{i=1}^{n+1} dz_i \, Q^{BC_{n+1}^{(2)}}_{A_{2n}^{(2)}}(x_1, \dots, x_n; z_1, \dots, z_{n+1}) \times$$

$$\times Q^{A_{2n}^{(2)}}_{BC_{n+1}^{(2)}}(z_1, \dots, z_{n+1}; y_1, \dots, y_n)$$
(6.1)

Baxter Q-operator for  $A_{2n-1}^{(2)}$  can be defined by the integral kernel

$$Q^{A_{2n-1}^{(2)}}(x_1, \dots, x_n; y_1, \dots, y_n) = \int \bigwedge_{i=1}^n dz_i \, Q^{A_{2n-1}^{(2)}}_{A_{2n-1}^{(2)}}(x_1, \dots, x_n; z_1, \dots, z_n) \times$$

$$\times Q^{A_{2n-1}^{(2)}}_{A_{2n-1}^{(2)}}(z_1, \dots, z_n; y_1, \dots, y_n)$$
(6.2)

Baxter Q-operator for  $B_n^{(1)}$  can be defined by the integral kernel

$$Q^{B_n^{(1)}}(x_1, \dots, x_n; y_1, \dots, y_n) = \int \bigwedge_{i=1}^{n+1} dz_i \, Q^{BC_{n+1}^{(1)}}_{B_n^{(1)}}(x_1, \dots, x_n; z_1, \dots, z_{n+1}) \times \qquad (6.3)$$
$$Q^{B_n^{(1)}}_{BC_{n+1}^{(1)}}(z_1, \dots, z_{n+1}; y_1, \dots, y_n)$$

Baxter Q-operator for  $C_n^{(1)}$  can be defined by the integral kernel

$$Q^{C_n^{(1)}}(x_1, \dots, x_n; y_1, \dots, y_n) = \int \bigwedge_{i=1}^n dz_i \, Q^{D_n^{(1)}}_{C_n^{(1)}}(x_1, \dots, x_n; z_1, \dots, z_n) \times \qquad (6.4)$$
$$Q^{C_n^{(1)}}_{D_n^{(1)}}(z_1, \dots, z_n; y_1, \dots, y_n)$$

Baxter Q-operator for  $D_n^{(1)}$  can be defined by the integral kernel

$$Q^{D_n^{(1)}}(x_1, \dots, x_n; y_1, \dots, y_n) = \int \bigwedge_{i=1}^n dz_i \, Q^{C_n^{(1)}}_{D_n^{(1)}}(x_1, \dots, x_n; z_1, \dots, z_n) \times \qquad (6.5)$$
$$Q^{D_n^{(1)}}_{C_n^{(1)}}(z_1, \dots, z_n; y_1, \dots, y_n)$$

## 7 Baxter operators for $B_{\infty}$ , $C_{\infty}$ and $D_{\infty}$

It is not difficult to find elementary intertwining and Baxter operators for the Toda chains connected with infinite root systems  $B_{\infty}$ ,  $C_{\infty}$  and  $D_{\infty}$ . These operators are quite similar to that for  $A_{\infty}$  (see [T] for a classical case) with an additional subtlety discussed above for affine root systems of classical type. By similar reason as for affine root systems we also need to include the infinite non-reduced root system  $BC_{\infty}$ .

Simple roots and Dynkin diagrams for infinite group  $A_{\infty}$ ,  $B_{\infty}$ ,  $C_{\infty}$ ,  $BC_{\infty}$  and  $D_{\infty}$  are as follows

$$B_{\infty}: \quad \alpha_1 = e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \quad i \in \mathbb{Z}_{>0},$$
 (7.1)

$$\alpha_1 \Leftarrow \alpha_2 \leftarrow \alpha_3 \leftarrow \dots \qquad (7.2)$$

$$C_{\infty}: \quad \alpha_1 = 2e_1, \qquad \alpha_{i+1} = e_{i+1} - e_i, \quad i \in \mathbb{Z}_{>0},$$
 (7.3)

$$\alpha_1 \Longrightarrow \alpha_2 \longrightarrow \alpha_3 \longrightarrow \dots \tag{7.4}$$

$$D_{\infty}: \quad \alpha_1 = e_1 + e_2, \qquad \alpha_{i+1} = e_{i+1} - e_i, \qquad i \in \mathbb{Z}_{>0}, \tag{7.5}$$

$$BC_{\infty}: \quad \alpha_1 = e_1, \qquad \alpha_2 = 2e_1, \qquad \alpha_i = e_{i+1} - e_i, \qquad i \in \mathbb{Z}_{>0},$$
 (7.7)

$$\frac{\alpha_0}{\alpha_1} \Longleftrightarrow \alpha_2 \longleftarrow \alpha_3 \longleftarrow \dots$$
(7.8)

Baxter Q-operator for  $A_{\infty}$  has the integral kernel

$$Q(x_i, y_i) = \exp\Big\{\sum_{i \in \mathbb{Z}} e^{x_i - y_i} + g_i e^{y_{i+1} - x_i}\Big\},$$
(7.9)

and intertwines  $A_\infty$  Toda chain Hamiltonians

$$\mathcal{H}^{A_{\infty}}(x_i) = -\frac{1}{2} \sum_{i \in \mathbb{Z}} \frac{\partial^2}{\partial x_i^2} + 2g_i e^{x_{i+1} - x_i}, \qquad (7.10)$$

$$\mathcal{H}^{A_{\infty}}(y_i) = -\frac{1}{2} \sum_{i \in \mathbb{Z}} \frac{\partial^2}{\partial y_i^2} + 2g_i e^{y_{i+1} - y_i}.$$
(7.11)

Its generalization to other classical series goes as follows. As in the previous cases we start with elementary intertwining operators. The integral operator with the kernel

$$Q_{B_{\infty}}^{BC_{\infty}}(x_i, z_i) = \exp\left\{g_1 e^{z_1} + \sum_{i>0} e^{x_i - y_i} + g_{i+1} e^{y_{i+1} - x_i}\right\}$$
(7.12)

intertwines  $B_{\infty}$  and  $BC_{\infty}$  Toda chain Hamiltonians

$$\mathcal{H}^{B_{\infty}}(x_i) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1} + \sum_{i=1}^{\infty} g_{i+1} e^{x_{i+1} - x_i},$$
(7.13)

$$\mathcal{H}^{BC_{\infty}}(z_i) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial z_i^2} + \frac{g_1}{2} \left( e^{z_1} + e^{2z_1} \right) + \sum_{i=1}^{\infty} g_{i+1} e^{z_{i+1} - z_i}$$
(7.14)

Baxter Q-operator for  $B_{\infty}$  then obtained by applying this intertwiner operators twice and has as a kernel

$$Q^{B_{\infty}}(x_i; y_i) = \int \bigwedge_{i=1}^{\infty} dz_i \cdot Q_{B_{\infty}}^{BC_{\infty}}(x_i, z_i) \cdot Q_{BC_{\infty}}^{B_{\infty}}(z_i, y_i)$$

Similarly the integral operator with the kernel

$$Q_{C_{\infty}}^{D_{\infty}}(x_i, z_i) = \exp\left\{g_1 e^{x_1 + z_1} + \sum_{i>0} e^{x_i - y_i} + g_i e^{y_{i+1} - x_i}\right\}$$
(7.15)

intertwines  $C_\infty$  and  $D_\infty$  Toda chain Hamiltonian operators

$$\mathcal{H}^{C_{\infty}}(x_i) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial x_i^2} + 2g_1 e^{2x_1} + \sum_{i=1}^{\infty} g_{i+1} e^{x_{i+1} - x_i},$$
(7.16)

$$\mathcal{H}^{D_{\infty}}(z_i) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial z_i^2} + g_1 e^{z_1 + z_2} + \sum_{i=1}^{\infty} g_{i+1} e^{z_{i+1} - z_i}.$$
(7.17)

and Baxter integral  $Q\text{-}\mathrm{operator}$  for  $C_\infty$  has as a kernel

$$Q^{C_{\infty}}(x_i; y_i) = \int \bigwedge_{i=1}^{\infty} dz_i \cdot Q_{C_{\infty}}^{D_{\infty}}(x_i, z_i) \cdot Q_{D_{\infty}}^{C_{\infty}}(z_i, y_i)$$

Finally the integral operator with the kernel

$$Q_{D_{\infty}}^{C_{\infty}}(x_i, z_i) = \exp\left\{g_1 e^{x_1 + z_1} + \sum_{i>0} e^{z_i - x_i} + g_{i+1} e^{x_{i+1} - z_i}\right\}$$
(7.18)

intertwines  $D_\infty$  and  $C_\infty$  Toda chain Hamiltonians

$$\mathcal{H}^{D_{\infty}}(x_i) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial x_i^2} + g_1 e^{x_1 + x_2} + \sum_{i=1}^{\infty} g_{i+1} e^{x_{i+1} - x_i},$$
(7.19)

$$\mathcal{H}^{C_{\infty}}(z_i) = -\frac{1}{2} \sum_{i=1}^{\infty} \frac{\partial^2}{\partial z_i^2} + 2g_1 e^{2z_1} + \sum_{i=1}^{\infty} g_{i+1} e^{z_{i+1}-z_i}.$$
(7.20)

Therefore Baxter integral Q-operator for  $D_{\infty}$  has the following kernel

$$Q^{D_{\infty}}(x_i; y_i) = \int \bigwedge_{i=1}^{\infty} dz_i \cdot Q^{C_{\infty}}_{D_{\infty}}(x_i, z_i) \cdot Q^{D_{\infty}}_{C_{\infty}}(z_i, y_i)$$

## 8 Conclusions

In this note we provide explicit expressions for recursive operators and Baxter operators connected with the classical series. This allows us to generalize Givental representation for the eigenfunctions of the open Toda chains quadratic Hamiltonians to the case of all classical series for zero eigenvalues. These representations can be straightforwardly generalized to the case of the common eigenvalue of the full set of the Hamiltonians with non-zero eiegenvalues [GLO]. The generalization relies on a connection between generating function for the Hamiltonians of Toda theories and (twisted) Yangians (see e.g. [MNO]). We would like to stress that the results presented in this note can be considered as a generalization to classical series of only a part of [GKLO]. The other part is connected with the interpretation of the elementary intertwining operators in representation theory framework and will be presented elsewhere. Taking into account the simplicity and a unified form of the arising constructions in the representation theory of classical groups one can expect important applications in the theory of L-functions and related subjects.

## References

- [B] R.J. Baxter, *Exactly solved models in statistical mechanics*, London: Academic Press, 1982.
- [DS] V.G. Drinfel'd, V.V. Sokolov, Lie algebras and equations of Korteveg-de Vries type, Translated from Itogi Nauki i Tekhniki, Seriya Sovremennye Problemy Matematiki,v. 24 (1984), 81–180.
- [F] L.D. Faddeev, Quantum completely integrable models in field theory, Sov. Sci. Rev., Sect. C (Math. Phys. Rev.) 1 (1980), 107-155.
- [GKLO] A. Gerasimov, S. Kharchev, D. Lebedev, S. Oblezin, On a Gauss-Givental representation for quantum Toda chain wave function, To be published in Int. Math. Res. Notices, (2006), [arXiv:math.RT/0505310].
- [GLO] A. Gerasimov, D. Lebedev, S. Oblezin, On a Gauss-Givental representation for classical groups, to appear.
- [Gi] A. Givental, Stationary Phase Integrals, Quantum Toda Lattices, Flag Manifolds and the Mirror Conjecture, AMS Trans. (2) 180 (1997), 103-115.
- [JK] D. Joe, B. Kim, Equivariant mirrors and the Virasoro conjecture for flag manifolds, Int. Math. Res. Notes 2003 No. 15 (2003), 859–882.
- [K] V. Kac, Infinite-dimensional Lie algebras, Third edition, Cambridge University Press, Cambridge 1990.
- [MNO] A. Molev, M. Nazarov, G. Olshanski, Yangians and classical Lie algebras, Russian Math. Surveys, 51, (1996), 205–282.

- [PG] V. Pasquier, M. Gaudin, The periodic Toda chain and a matrix generalization of the Bessel function recursion relation, J. Phys. A 25 (1992), 5243–5252.
- [RSTS] A.G. Reyman, M.A. Semenov-Tian-Shansky, *Integrable Systems. Group theory approach*, Modern Mathematics, Moscow-Igevsk: Institute Computer Sciences, 2003.
- [T] M. Toda, Theory of Nonlinear Lattices, Berlin, Springer-Verlag, 1981.