

# POISSON STRUCTURES ON THE HOMOLOGY OF THE SPACE OF KNOTS

KEIICHI SAKAI

**ABSTRACT.** In this article we study the Poisson algebra structure on the homology of the totalization of a fibrant cosimplicial space associated with an operad with multiplication. This structure is given as the Browder operation induced by the action of little disks operad, which was found by J. McClure and J. Smith. We show that the Browder operation coincides with the Poisson algebra structure on the Hochschild homology, which appears as the  $E^2$ -term of the homology spectral sequence constructed by A. Bousfield. In particular we consider a variant of the space of long knots in higher dimensional Euclidean space, and show that D. Sinha's homology spectral sequence computes the Poisson algebra structure of the homology of the space. We find a homology class which does not directly correspond to chord diagrams.

## 1. OVERVIEW

A *long knot* is an embedding  $\mathbb{R}^1 \hookrightarrow \mathbb{R}^n$ ,  $n \geq 3$ , which agrees with a fixed line outside a compact set. A long immersion is defined similarly.

Let  $\mathcal{K}_n$  and  $\text{Imm}_n$  be the spaces of long knots and long immersions in  $\mathbb{R}^n$  respectively. They are homotopy commutative Hopf spaces whose products are connecting sums. Consider the space

$$\mathcal{K}'_n := \text{the homotopy fiber of } \mathcal{K}_n \hookrightarrow \text{Imm}_n.$$

This is also a Hopf space, so its homology is a Hopf algebra. Moreover, when  $n > 3$ , there are two ways to define the Poisson algebra structure on the rational homology group of the space, as explained below. Our main theorem 1.4 shows that in both ways we obtain the same structure (up to sign).

D. Sinha [10] constructed a cosimplicial model for the space, using the method “functor calculus” due to T. Goodwillie [4, 5]. The readers are referred to [7] for the detailed introduction to operads.

**Theorem 1.1** ([10]). *There exists a topological operad  $X_n^\bullet$  with multiplication such that*

- (1) *there exist homotopy equivalences from  $\text{Conf}(\mathbb{R}^n, k) = (\mathbb{R}^n)^k \setminus \Delta$  to  $X^k$  for all  $k \geq 0$ ,*
- (2) *when  $n > 3$ ,  $\text{Tot} RX_n^\bullet$  is weakly equivalent to  $\mathcal{K}'_n$  where  $RX_n^\bullet$  is a fibrant replacement of the associated cosimplicial space.*  $\square$

The operad  $X_n^\bullet$  is called the *Kontsevich operad* in [10]. Since it is multiplicative, we can construct an associated cosimplicial space, which we also denote by  $X_n^\bullet$  in the above. Below its fibrant replacement will be also denoted by  $X_n^\bullet$ .

One of the remarkable features of such a cosimplicial space is the existence of an operad action on the totalization.

**Theorem 1.2** ([8]). *The totalization of a cosimplicial space associated with a topological operad  $\mathcal{O}$  with multiplication admits an action of the operad  $\mathcal{D}$  which is weakly equivalent to the little disks operad, that is, there exist maps*

$$\mathcal{D}(k) \times (\mathrm{Tot} \mathcal{O})^k \longrightarrow \mathrm{Tot} \mathcal{O}, \quad k \geq 0,$$

*with the associativity conditions.*  $\square$

This action induces an operation on the homology of the totalization, called the *Browder operation*,

$$\lambda : H_p(\mathrm{Tot} \mathcal{O}, \mathbb{Q}) \otimes H_q(\mathrm{Tot} \mathcal{O}, \mathbb{Q}) \longrightarrow H_{p+q+1}(\mathrm{Tot} \mathcal{O}, \mathbb{Q}),$$

which is determined by a generator of  $H_1(\mathcal{D}(2)) \cong \mathbb{Z}$ . This map can be regarded as a Poisson algebra structure on the homology [2].

On the other hand, since  $X_n^\bullet$  satisfies the conditions of Theorem 3.2 of [1], we can construct a homology spectral sequence  $E^r$  associated with a double complex  $S_*(X_n^\bullet)$ , the modules of singular chains of  $X_n^\bullet$ , and converging to  $H_*(\mathrm{Tot} X_n^\bullet)$ . Indeed Bousfield proved that (under the same conditions) the total complex  $TS(X_n^\bullet)$  is an algebraic model for  $\mathrm{Tot} X_n^\bullet$ , that is, there exists a quasi-isomorphism

$$\varphi : S_*(\mathrm{Tot} X_n^\bullet) \longrightarrow TS(X_n^\bullet).$$

By definition the  $E^1$ -term consists of the homology groups of  $X_n^\bullet$ . Moreover it is not hard to see that  $E^1$ -term is the *Hochschild complex* of the homology operad  $H_*(X_n^\bullet)$ , which V. Turchin called the *Poisson algebras operad* [11]. Since  $X_n^\bullet$  is multiplicative, there exist a natural product and a bracket on the complex that make the Hochschild homology  $HH_*(X_n^\bullet)$  a Poisson algebra [3, 11, 12].

In the case of  $X_n^\bullet$  we have the rational degeneracy of the spectral sequence.

**Theorem 1.3** ([6]). *When  $n > 3$ , the above homology spectral sequence degenerates at  $E^2$ -term rationally. Thus  $H_*(K'_n, \mathbb{Q})$  is the Hochschild homology of the Poisson algebras operad.*  $\square$

This theorem holds since the *formality* of the operad  $X_n^\bullet$  allows us to replace the vertical differentials of the spectral sequence by zero maps.

Define the natural filtrations:

$$F_p TS(X_n^\bullet)|_{\mathrm{deg}=q} = \prod_{l \geq p} S_{q+l}(X^l),$$

$$F_p H_q(TS(X_n^\bullet)) = \mathrm{im} \{H_q(F_p TS(X_n^\bullet)) \rightarrow H_q(TS(X_n^\bullet))\}$$

and denote its associated quotient by  $GH_*(TS(X_n^\bullet))$ :

$$GH_q(TS(X_n^\bullet)) = \bigoplus_p F_p H_q(TS(X_n^\bullet)) / F_{p+1} H_q(TS(X_n^\bullet)).$$

Thus we have two Poisson algebra structures on  $GH_*(TS(X_n^\bullet))$ : one induced via  $\varphi$  and the other defined on the Hochschild homology.

**Theorem 1.4.** *Suppose  $n > 3$ . Let  $X_n^\bullet$  be the Kontsevich operad, or (a fibrant replacement of) its associated cosimplicial space. The Browder operation induced on  $GH_*(TS(X_n^\bullet))$  via  $\varphi$  coincides up to sign with the Poisson structure defined on the Hochschild homology of Poisson algebras operad.*

**Remark 1.5.** *We will not use the specific properties of  $X_n^\bullet$ , except of the formality. Hence our construction would work in more general situations: for any topological operad  $\mathcal{O}$  with multiplication, the Browder operation defined on  $H_*(\text{Tot } R\mathcal{O})$  would be compatible with the Poisson structure on the  $E^2$ -term of the homology spectral sequence.*  $\square$

We should notice that P. Salvatore [9] has already known the result. His motivation was slightly different from ours. He proved that the space  $\mathcal{K}_n$  is also a double loop space, but the existence of certain homology class of degree  $3n - 8$  (discussed in the next section) shows that the first projection  $\mathcal{K}'_n \rightarrow \mathcal{K}_n$  is not a map of double loop spaces. Here we use the same result to prove that the Browder operation is not trivial and produces a homology class which does not correspond to any chord diagram (see below).

The proof will be given later. First we compute some homology groups of  $\mathcal{K}'_n$ .

## 2. SOME COMPUTATIONS

The subgroup  $\bigoplus_{k \geq 0} H_{(n-3)k}(\mathcal{K}_n)$  is known to be non-trivial, since it contains the subalgebra  $\mathcal{A}$  isomorphic to the algebra of chord diagrams (modulo 4-term relations). But few homology classes except of those in  $\mathcal{A}$  were known.

Theorem 1.4 implies that the spectral sequence computes the Poisson algebra structure on  $H_*(\mathcal{K}'_n, \mathbb{Q})$ . By computing the  $E^2$ -term explicitly (see [12]), we find that the Browder operation produces new homology classes.

To state the result, we need the following.

**Lemma 2.1** ([10]). *There exists a homotopy equivalence  $\mathcal{K}'_n \sim \mathcal{K}_n \times \Omega^2 S^{n-1}$ . This decomposition preserves the Hopf space structures.*  $\square$

The topology of  $\Omega^2 S^{n-1}$  is well known:

$$H_*(\Omega^2 S^{n-1}, \mathbb{Q}) \cong \begin{cases} \Lambda^*[ \iota ] \otimes P[\lambda(\iota, \iota)] & n \text{ is odd,} \\ \Lambda^*[ \iota ] & n \text{ is even,} \end{cases}$$

where  $\iota$  is the image of  $1 \in \pi_{n-3}(\Omega^2 S^{n-1})$  via the Hurewicz isomorphism, and  $\lambda$  is the Browder operation induced by the little disks action on  $\Omega^2 S^{n-1}$ . Using this, we can compute some low-degree homology groups of  $\mathcal{K}'_n$  and  $\mathcal{K}_n$ .

**Theorem 2.2.** *Suppose  $n > 4$ . Then*

$$H_{3n-8}(\mathcal{K}'_n, \mathbb{Q}) \cong \begin{cases} \mathbb{Q}\iota\lambda(\iota, \iota) \oplus \mathbb{Q}\lambda(\iota, v_2) & n \text{ is odd,} \\ \mathbb{Q}\lambda(\iota, v_2) & n \text{ is even} \end{cases}$$

where  $v_2 \in H_{2(n-3)}(\mathcal{K}_n, \mathbb{Q})$  is the class made from a chord diagram with two chords. By the Künneth formula

$$H_{3n-8}(\mathcal{K}_n, \mathbb{Q}) \cong \mathbb{Q}$$

and this group is generated by the homology class corresponding to  $\lambda(\iota, v_2)$ , which is the first example of a homology class of  $\mathcal{K}_n$  which does not directly correspond to chord diagrams.  $\square$

In other words, we can regard  $H_*(\Omega^2 S^{n-1}, \mathbb{Q})$  as acting on  $H_*(\mathcal{K}_n, \mathbb{Q})$  via the Browder operation, and this action is non-trivial.

The above computation suggests that we might be able to obtain more homology classes by applying  $\lambda(\iota, \cdot)$  to the elements of  $\mathcal{A}$ . It is possible in principle to compute them, but it would become more exhausting as the degrees raise (see [12]).

## 3. PROOF OF THE MAIN THEOREM

**3.1. algebraic model for  $\text{Tot } \mathcal{O}$ .** Let  $\mathcal{O}$  be a topological operad with multiplication and suppose the associated cosimplicial space is fibrant. Let  $TS(\mathcal{O})$  be the total complex of the double complex from the first section: its degree  $k$  part is

$$TS(\mathcal{O})_k = \prod_{l \geq 0} S_{k+l}(\mathcal{O}^l).$$

Bousfield [1] constructed a quasi-isomorphism

$$\varphi : S_*(\text{Tot } \mathcal{O}) \longrightarrow TS(\mathcal{O})$$

when  $\mathcal{O}$  satisfies some convergence conditions of [1]. This is defined as follows. We regard a chain  $f^\bullet = \sum a_i f_i^\bullet \in S_q(\text{Tot } \mathcal{O})$  as the sum of maps

$$f_i^l : \Delta^q \times \Delta^l \longrightarrow \mathcal{O}^l, \quad l \geq 0,$$

which satisfy the cosimplicial identities. We choose the Eilenberg-MacLane map  $EM \in S_{q+l}(\Delta^q \times \Delta^l)$ ,  $l, q \geq 0$ , which gives a chain equivalence

$$S_*(M) \otimes S_*(N) \xrightarrow{\cong} S_*(M \times N)$$

for any spaces  $M$  and  $N$ . Then the quasi-isomorphism  $\varphi$  is defined by

$$\varphi(f^\bullet) \in \prod_{l \geq 0} S_{q+l}(\mathcal{O}^l),$$

$$\varphi(f^\bullet)_l := \sum a_i (f_i^l \circ EM) \in S_{q+l}(\mathcal{O}^l).$$

Our main theorem states that this quasi-isomorphism preserves the Poisson algebra structures.

**3.2. disks action on  $\text{Tot } \mathcal{O}$ .** Studying the disks action of the form

$$\mathcal{D}(2) \times (\text{Tot } \mathcal{O})^2 \longrightarrow \text{Tot } \mathcal{O}$$

defined by McClure-Smith [8] in detail, we can describe the Browder operation on  $H_*(\text{Tot } \mathcal{O}, \mathbb{Q})$  as follows. We use the notation  $\circ_i$  for the operad structure:

$$\circ_i : \mathcal{O}(k) \times \mathcal{O}(l) \longrightarrow \mathcal{O}(k+l-1), \quad a \circ_i b = a \overbrace{(\text{id}, \dots, \text{id}, b, \text{id}, \dots, \text{id})}^{i-1}$$

for any operad  $\mathcal{O}$ .

**Theorem 3.1.** *Let  $g_1^\bullet \in S_q(\text{Tot } \mathcal{O})$  and  $g_2^\bullet \in S_s(\text{Tot } \mathcal{O})$  be cycles. Define the map*

$$\langle g_1^\bullet, g_2^\bullet \rangle^l : \Delta^q \times \Delta^s \times \Delta^1 \times \Delta^l \longrightarrow \mathcal{O}^l, \quad l \geq 0$$

by

$$\begin{aligned} \langle g_1^\bullet, g_2^\bullet \rangle^l(x, y, \tau, t) = & \\ & \begin{cases} g_1^{f_{\tau,t}^{-1}(1)}(x)(u^1(\tau, t)) \circ_{i_0(\tau,t)+1} g_2^{f_{\tau,t}^{-1}(2)}(y)(u^2(\tau, t)), & 0 \leq \tau \leq 1/2, \\ g_2^{\bar{f}_{\tau,t}^{-1}(2)}(y)(v^2(\tau, t)) \circ_{j_0(\tau,t)+1} g_1^{\bar{f}_{\tau,t}^{-1}(1)}(x)(v^1(\tau, t)), & 1/2 \leq \tau \leq 1 \end{cases} \end{aligned}$$

for some

- $f_{\tau,t}, \bar{f}_{\tau,t} : \{0, 1, \dots, l+2\} \rightarrow \{1, 2\}$ ,
- $i_0(\tau, t), j_0(\tau, t) \geq 0$ ,
- $u^i(\tau, t) \in \Delta^{f^{-1}(i)} := \Delta^{|f^{-1}(i)|-1}$  ( $i = 1, 2$ ) and

- $v^i(\tau, t) \in \Delta^{\bar{f}^{-1}(i)}$  ( $i = 1, 2$ )

(they are defined below). Then the map

$$\lambda : H_q(\text{Tot } \mathcal{O}) \otimes H_s(\text{Tot } \mathcal{O}) \rightarrow H_{q+s+1}(\text{Tot } \mathcal{O})$$

given by

$$\lambda(g_1^\bullet, g_2^\bullet) := (-1)^{q+1} \langle g_1^\bullet, g_2^\bullet \rangle \circ EM \in H_{q+s+1}(\text{Tot } \mathcal{O})$$

is the Browder operation, where  $EM$  is the Eilenberg-MacLane map.  $\square$

**Remark 3.2.** Though the symbols  $f_{\tau,t}$ ,  $i_0$ ,  $j_0$ ,  $u_i$  and  $v_i$  are as follows, their explicit forms are not important, so readers may skip here. For  $0 \leq \tau \leq 1/2$  and  $t = (t_0, \dots, t_l) \in \Delta^l$ , define:

- $i_0 := \min\{i; t_0 + \dots + t_i \geq \tau\}$ ,  $i_1 := \min\{i; t_0 + \dots + t_i \geq \tau + \frac{1}{2}\}$ ,
- $f = f_{\tau,t} : [l+2] := \{0, 1, \dots, l+2\} \rightarrow \bar{2} := \{1, 2\}$ ,  $f^{-1}(1) = \{0, \dots, i_0, i_0+1, \dots, l+2\}$ ,  $f^{-1}(2) = \{i_0+1, \dots, i_1+1\}$ ,
- $h = s^{i_1} s^{i_0} : [l+2] \rightarrow [l]$ ,
- $u = (u^1, u^2) = ((u_p)_{p \in f^{-1}(1)}; (u_q)_{q \in f^{-1}(2)}) \in \Delta^{f^{-1}(1)} \times \Delta^{f^{-1}(2)}$ , where

$$u_b = \begin{cases} 2t_b & b < i_0, \\ 2\tau - 2(t_0 + \dots + t_{i_0-1}) & b = i_0, \\ 2(t_0 + \dots + t_{i_0}) + 2\tau & b = i_0 + 1, \\ 2t_{b-1} & i_0 + 1 < b < i_1 + 1, \\ (2\tau + 1) - 2(t_0 + \dots + t_{i_1-1}) & b = i_1 + 1, \\ 2(t_0 + \dots + t_{i_1}) - (2\tau + 1) & b = i_1 + 2, \\ 2t_{b-2} & i_1 + 2 < b \leq l+2. \end{cases}$$

and for  $1/2 \leq \tau \leq 1$  and  $t = (t_0, \dots, t_l) \in \Delta^l$ , define:

- $j_0 := \min\{j; t_0 + \dots + t_j \geq 1 - \tau\}$ ,  $j_1 := \min\{j; t_0 + \dots + t_j \geq \frac{3}{2} - \tau\}$ ,
- $\bar{f} : [l+2] \rightarrow \bar{2}$ ,  $\bar{f}^{-1}(1) = \{j_0 + 1, \dots, j_1 + 1\}$ ,  $\bar{f}^{-1}(2) = \{0, \dots, j_0, j_1 + 2, \dots, l+2\}$ ,
- $\bar{h} = s^{j_1} s^{j_0} : [l+2] \rightarrow [l]$ ,
- $v = (v^1, v^2) = ((v_a)_{a \in \bar{f}^{-1}(1)}; (v_b)_{b \in \bar{f}^{-1}(2)}) \in \Delta^{\bar{f}^{-1}(1)} \times \Delta^{\bar{f}^{-1}(2)}$ , where

$$v_b = \begin{cases} 2t_b & b < j_0, \\ (2 - 2\tau) - 2(t_0 + \dots + t_{j_0-1}) & b = j_0, \\ 2(t_0 + \dots + t_{j_0}) - (2 - 2\tau) & b = j_0 + 1, \\ 2t_{b-1} & j_0 + 1 < b < j_1 + 1, \\ (3 - 2\tau) - 2(t_0 + \dots + t_{j_1-1}) & b = j_1 + 1, \\ 2(t_0 + \dots + t_{j_1}) - (3 - 2\tau) & b = j_1 + 2, \\ 2t_{b-2} & j_1 + 2 < b \leq l+2. \end{cases} \quad \square$$

Moreover the straightforward computation shows the following.

**Lemma 3.3.** For any given  $l$  and  $0 \leq i_0 \leq i_1 \leq l$ , define

$$\Delta(i_0, i_1) := \{(\tau, t) \in [0, 1/2] \times \Delta^l \mid i_\epsilon(\tau, t) = i_\epsilon \ (\epsilon = 0, 1)\}.$$

Then the correspondence

$$u = (u^1, u^2) : \Delta(i_0, i_1) \longrightarrow \Delta^{f_{\tau,t}^{-1}(1)} \times \Delta^{f_{\tau,t}^{-1}(2)},$$

defined in the above proof, is a homeomorphism. Similarly, if we define

$$\Delta'(j_0, j_1) := \{(\tau, t) \in [1/2, 1] \times \Delta^l \mid j_\epsilon(\tau, t) = j_\epsilon \ (\epsilon = 0, 1)\},$$

for any given  $l$  and  $0 \leq j_0 \leq j_1 \leq l$ , then

$$v = (v^1, v^2) : \Delta'(j_0, j_1) \longrightarrow \Delta^{\bar{f}_{\tau, t}^{-1}(1)} \times \Delta^{\bar{f}_{\tau, t}^{-1}(2)}$$

is a homeomorphism.  $\square$

**3.3. Browder operation in terms of  $TS(\mathcal{O})$ .** Via the quasi-isomorphism  $\varphi$ , the Browder operation will be interpreted as follows.

**Theorem 3.4.** *Given any  $h \in TS(\mathcal{O})_q$ , we will write  $h = (h_l)_{l \geq 0}$  with  $h_l \in S_{q+l}(\mathcal{O}^l)$ . Then for any  $g_1^\bullet \in H_q(\text{Tot } \mathcal{O})$  and  $g_2^\bullet \in H_s(\text{Tot } \mathcal{O})$ ,*

$$\begin{aligned} \varphi(\lambda(g_1^\bullet, g_2^\bullet))_l = & \sum_{p+r=l+1} (-1)^{(p+1)(r+1)+q+s} \left[ \sum_{1 \leq i \leq p} (-1)^{\epsilon_i} \varphi(g_1^\bullet)_p \circ_i \varphi(g_2^\bullet)_r \right. \\ & \left. - (-1)^{(q+1)(s+1)} \sum_{1 \leq j \leq r} (-1)^{\epsilon'_j} \varphi(g_2^\bullet)_r \circ_j \varphi(g_1^\bullet)_p \right] \end{aligned}$$

where

$$\epsilon_i = (q-1)(p-i) + (p-1)(r+s), \quad \epsilon'_j = (p-1)(r-j) + (p+q)(r-1).$$

*Proof.* The map

$$\Delta^q \times \Delta^s \times \Delta^1 \times \Delta^l \xrightarrow{\langle g_1, g_2 \rangle} X^l$$

pre-composed by  $EM$  represents  $\varphi(\lambda(g_1, g_2))_l$ . By definition  $\Delta^1 \times \Delta^l$  is triangulated by  $\Delta(i_0, i_1)$ 's and  $\Delta'(j_0, j_1)$ 's ( $0 \leq i_0 \leq i_1 \leq l$ ,  $0 \leq j_0 \leq j_1 \leq l$ ). By Theorem 3.1,  $\langle g_1, g_2 \rangle$  is  $g_1^p \circ_{i_0+1} g_2^r$  when restricted on  $\Delta^q \times \Delta^s \times \Delta(i_0, i_1)$ , where

$$p = |f_{\tau, t}^{-1}(1)| - 1 = l + i_0 - i_1 + 1, \quad r = |f_{\tau, t}^{-1}(2)| - 1 = i_1 - i_0,$$

and, when restricted on  $\Delta^q \times \Delta^s \times \Delta'(j_0, j_1)$ ,  $\langle g_1, g_2 \rangle$  is  $g_2^r \circ_{j_0+1} g_1^p$ , where  $p$  and  $r$  are determined similarly.

Thus  $\varphi(\lambda(g_1^\bullet, g_2^\bullet))$  should be a linear sum of  $\varphi(g_1^\bullet)_p \circ_i \varphi(g_2^\bullet)_r$ ,  $1 \leq i \leq p$ , and  $\varphi(g_2^\bullet)_r \circ_j \varphi(g_1^\bullet)_p$ ,  $1 \leq j \leq r$ . The coefficients are  $\pm 1$  because of Lemma 3.3. The signs  $\pm$  are those of the Jacobians of the maps

$$\begin{aligned} \Delta^q \times \Delta^s \times \Delta(i_0, i_1) &\xrightarrow{\sim} (\Delta^q \times \Delta^p) \times (\Delta^s \times \Delta^r), \\ \Delta^q \times \Delta^s \times \Delta'(j_0, j_1) &\xrightarrow{\sim} (\Delta^s \times \Delta^r) \times (\Delta^q \times \Delta^p). \end{aligned}$$

Explicit computations show that the signs are

$$(-1)^{1+i_0(i_1-i_0)(l-i_1)+ps} = (-1)^{(p+1)(r+1)+s+1} (-1)^{(r-1)(p-i_0-1)+(p-1)r+s},$$

and

$$(-1)^{1+j_0+(j_1-j_0)(l-j_1)+q(r+s)} = (-1)^{(p+1)(r+1)+qs+q+1} (-1)^{(p-1)(r-j_0-1)+(p+q)(r-1)}$$

respectively. They give the desired formula.  $\square$

**3.4. Poisson bracket on  $E^2$ .** Here let  $\mathcal{O}'$  be any operad of graded modules with multiplication (in our case  $\mathcal{O}' = H_*(X_n^\bullet)$ ). By unraveling the descriptions of [3, 11, 12] of the Hochschild complex  $(\mathcal{O}', \partial)$ , we can see the following.

**Theorem 3.5** ([3, 11, 12]). *The Poisson structure on  $H_*(\mathcal{O}')$  is induced by the (degree-preserving) map*

$$[-, -] : \mathcal{O}'(p) \times \mathcal{O}'(r) \longrightarrow \mathcal{O}'(p+r-1),$$

$$[x, y] := \sum_{1 \leq i \leq p} (-1)^{\epsilon_i} x \circ_i y - (-1)^{(q+1)(s+1)} \sum_{1 \leq j \leq r} (-1)^{\epsilon'_j} y \circ_j x$$

where  $q = \deg x$ ,  $s = \deg y$ , and  $\epsilon_i, \epsilon'_j$  are as in Theorem 3.4.  $\square$

Since the spectral sequence for  $\mathcal{O} = X_n^\bullet$  degenerates rationally at  $E^2$ , the natural map

$$\psi : E_{-p,q}^2 \otimes \mathbb{Q} \longrightarrow (F_p/F_{p+1})H_q(TS(X_n^\bullet), \mathbb{Q})$$

is an isomorphism, and via  $\psi$  a bracket is induced on  $GH_*(TS(X_n^\bullet))$ , where  $G$  denotes the associated quotient. Comparing Theorem 3.4 with Theorem 3.5, we can see that this bracket coincides on  $GH_*(TS(X_n^\bullet))$  with the Browder operation induced via  $\varphi$ .  $\square$

#### ACKNOWLEDGEMENT

The author expresses his great appreciation to Toshitake Kohno for his encouragement and advices. The author is also grateful to Victor Turchin and James McClure for reading the draft of the paper and giving him many suggestions, to Dev Sinha for answering his questions, and to Fred Cohen and Ryan Budney for teaching him about the little disks actions and so on.

This research is partially supported by the 21st century COE program at Graduate School of Mathematical Sciences, the University of Tokyo.

#### REFERENCES

- [1] A. Bousfield, *On the homology spectral sequence of a cosimplicial space*, Amer. J. Math., 109 (1987), no.2, 361-394.
- [2] F. Cohen, *The homology of  $C_{n+1}$ -space*,  $n \geq 0$ , Springer LNM 533, 207-351.
- [3] M. Gerstenhaber, A. Voronov, *Homotopy  $G$ -algebras and moduli space operad*, Internat. Math. Res. Notices (1995), no. 3, 141-153.
- [4] T. Goodwillie, *Calculus. II. Analytic functors*, K-Theory 5 (1991/92), no. 4, 295-332.
- [5] T. Goodwillie, *Calculus. III. Taylor series*, Geom. Topol. 7 (2003), 645-711.
- [6] P. Lambrechts, V. Turchin and I. Volić, *The rational homotopy type of spaces of knots in high codimension*, in preparation.
- [7] J. McClure, J. Smith, *Operads and cosimplicial objects: an introduction*, Axiomatic, Enriched and Motivic Homotopy Theory. Proc. NATO Adv. Study Inst. Edited by J.P.C. Greenlees. Kluwer (2004).
- [8] J. McClure, J. Smith, *Cosimplicial objects and little  $n$ -cubes I*, Amer. J. Math. 126 (2004), no.5, 1109-1153.
- [9] P. Salvatore, *Knots, operads and double loop spaces*, math.AT/0608490.
- [10] D. Sinha, *Operads and knot spaces*, J. Amer. Math. Soc. 19 (2006), no. 2, 461-486.
- [11] V. Tourtchine (Turchin), *On the homology of the spaces of long knots*, NATO Sci. Ser. II Math. Phys. Chem., 179, Kluwer (2004), 23-52.
- [12] V. Tourtchine, *On the other side of the bialgebra of chord diagrams*, math.QA/0411436.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCE, THE UNIVERSITY OF TOKYO  
 E-mail address: ksakai@ms.u-tokyo.ac.jp