CODIMENSION TWO PL EMBEDDINGS OF SPHERES WITH NONSTANDARD REGULAR NEIGHBORHOODS

M. CENCELJ, D. REPOVŠ AND A. SKOPENKOV

ABSTRACT. For a given polyhedron $K \subset M$ the notation $R_M(K)$ denotes a regular neighborhood of K in M. We study the following problem: find all pairs (m, k) such that if K is a compact k-polyhedron and M a PL m-manifold, then $R_M(fK) \cong R_M(gK)$, for each two homotopic PL embeddings $f, g: K \to M$. We prove that $R_{S^{k+2}}(S^k) \not\cong S^k \times D^2$ for each $k \geq 2$ and some PL sphere $S^k \subset S^{k+2}$ (even for any PL sphere $S^k \subset S^{k+2}$ having an isolated non-locally flat point with the singularity $S^{k-1} \subset S^{k+1}$ such that $\pi_1(S^{k+1} - S^{k-1}) \not\cong \mathbb{Z}$).

This paper is on the uniqueness of regular neighborhoods problem for distinct embeddings of one manifold into the other. Definitions of regular neighborhood and other notions of PL topology can be found in [RoSa72]. For any subpolyhedron $X \subset M$ of a PL manifold M we denote the regular neighborhood of X in M by $R_M(X)$. The following is the main result of the present paper:

Theorem 1. $R_{S^{k+2}}(S^k)$ is not homeomorphic to $S^k \times D^2$ for any $k \ge 2$ and (a) any PL sphere $S^k \subset S^{k+2}$ which is the suspension over a locally flat PL knot $S^{k-1} \subset S^{k+1}$ such that $G = \pi_1(S^{k+1} - S^{k-1}) \not\cong \mathbb{Z}$; or

(b) any PL sphere $S^k \subset S^{k+2}$ having an isolated non-locally flat point with the singularity $S^{k-1} \subset S^{k+1}$ such that $G = \pi_1(S^{k+1} - S^{k-1}) \not\cong \mathbb{Z}$.

For k = 2 in Theorem 1.b one can take *any* non-locally flat PL sphere $S^2 \subset S^4$.

Recall that for $x \in S^k \subset S^{k+2}$ the singularity at point x is the isotopy class of the submanifold $\partial D^{k+2} \cap S^k$ of ∂D^{k+2} . Here $(D^{k+2}, D^{k+2} \cap S^k)$ is any PL (k+2, k) ball pair which is a regular neighborhood of x in the pair (S^{k+2}, S^k) .

Of course, Theorem 1.a follows from Theorem 1.b, but the proof of the former is simpler (in particular for k = 2, because our argument uses the fundamental group whereas the proof of Theorem 1.b for that cases uses [Gab86, Gab87]).

We now give some motivation for Theorem 1.1. An important question in the topology of manifolds [Mas59] asks for which cases the normal vector bundle of a smooth k-manifold K is independent of the choice of the smooth embedding $K \to S^m$. This is the case when:

(a) m = k + 1; or

¹⁹⁹¹ Mathematics Subject Classification. Primary: 57M25, 57Q40, Secondary: 57M05, 57N40.

Key words and phrases. Embedding, regular neighborhood, Dehn surgery, fundamental group.

Skopenkov gratefully acknowledges the support of the Pierre Deligne fund based on his 2004 Balzan prize in mathematics, INTAS Grant No. YSF-2002-393, the Russian Foundation for Basic Research, Grants No 05-01-00993 and 04-01-00682, President of Russian Federation Grants MD-3938.2005.1 and NSH-1988.2003.1. Cencelj and Repovš were supported by Slovenian Research Agency grants BI-RU/05-07-04 and BI-RU/05-07-13.

(b) m = k + 2 and K is orientable [Ker65], [Rol76]; or

(c) K is a homotopy k-sphere and either m = k + 3 or $m \ge \frac{3k}{2} + 1$ [Ker59], [Mas59], [HaHi62]; or

(d) $K = \mathbb{R}P^2$ and m = 4 [Mas69].

On the other hand, there exist smooth embeddings $K \,\subset\, S^m$ with distinct normal bundles [HLS65; Haefliger's example in §2], [Mas69], [Roh70], [MiRe71]. For example, K can be the Klein bottle and $f_0, f_4 : K \to S^4$ can be any smooth embeddings with the normal Euler classes equal to 0 and 4, respectively. Then $\pi_1(R_{S^4}(f_0K)) \cong \mathbb{Z}^3$ while $\pi_1(R_{S^4}(f_4K))$ is not virtually abelian, i.e. it does not contain any abelian subgroups of finite index. This argument, due to Hillman, shows that for such embeddings even the spaces of normal bundles are non-homeomorphic.

Any locally flat PL embedding has a normal block bundle [RoSa68]. Thus the Massey question makes sense in the locally flat PL category. In this category the normal bundle does not depend on the embedding when $K = S^k$ (by the Zeeman Unknotting Balls Theorem) and in the cases (a), (b), (d) above [Wal67; Theorem 2], [RoSa68; Corollary 6.8]. In particular, $R_{S^4}(S^2) \cong S^2 \times D^2$ for any locally flat PL embedding $S^2 \subset S^4$ (cf. Theorem 1). Note that a PL sphere $S^n \subset \mathbb{R}^m$ has a PL standard regular neighborhood if either $m - n \neq 2$ or the embedding $S^n \subset \mathbb{R}^m$ is locally flat [Wal67, Zee63].

From now on we shall work in the PL (not locally flat PL) category, unless otherwise specified. As another polyhedral version of the Massey problem we can consider the question, whether for embeddings $f, g: K \to S^m$ the homeomorphism $g \circ f^{-1}: fK \to gK$ can be extended to a homeomorphism $R_M(fK) \cong R_M(gK)$ [LiSi69]. The analogues of Theorem 1 and conjectures (e), (f), (g) below for this problem are trivial. Theorem 1 was motivated by the following related problem, raised independently by Shtanko and Chernavski (private communication) and also in [Hor85, Hor90, Rep88]:

Find all pairs (m, k) such that if K is a compact k-polyhedron and M a PL m-manifold, then

(*) $R_M(fK) \cong R_M(gK)$ for each two homotopic PL embeddings $f, g: K \to M$.

The particular case of this problem for $M = S^m$ (then f and g are always homotopic) attracted special attention.

The property (*) holds for $m \ge 2k+2$ by general position. In general, many invariants of $R_M(fK)$ and $R_M(gK)$ coincide: the homotopy types, the intersection rings, the higher Massey products, the (classifying maps of) tangent bundles (and hence also all invariants deduced from the tangent bundle, e.g. characteristic classes and numbers). This implies that (*) also holds for m = 2k + 1 (because in this case an *m*-thickening is completely determined by its tangent bundle [LiSi69]) and for m = 2 (because a 2-manifold N with boundary is completely determined by $H_1(N, \mathbb{Z}_2)$ and its intersection form). Also,

$$R_M(fK) \times I^l \cong R_{M \times I^l}(fK) \cong R_{M \times I^l}(gK) \cong R_M(gK) \times I^l$$

for $l \geq 2k + 1 - m$. The boundary $\partial R_M(fK)$ is *l*-connected for $m \geq k + l + 2$ and *l*-connected K. All this suggests that distinguishing $R_M(fK)$ and $R_M(gK)$ is a nontrivial problem for $m \geq k + 3$.

The property (*) holds for $m = k + 1 \ge 3$ and a *fake surface* K [Cas65], [Cav92], [Mat73], [Rep88], [RBS99]. The property (*) fails for:

(a) $M = S^m, m = k + 1 \ge 3$ and $K = S^k \lor S^1 \lor S^1$ [Rep88]; or

(b) $M = S^3$, K a common spine of the granny knot and the square knot [MPR89], [CaHe94], [CLR97], [CHR98]; or

(c) $M = S^4$ and the Dunce Hat K [Zee64], [Hor85], [Hor90]; or

(d) $M = S^{k+2}, k \ge 2$ and $K = S^k$ by Theorem 1.

We conjecture that the property (*) also fails for:

(e) $M = S^{2k}$ and $K = \Sigma(S^{k-1} \sqcup S^{k-1})$ (one can take f and g to be the suspensions over the trivial link and the Hopf link, respectively); or

(f) $M = S^4$ and $K = \mathbb{R}P^2$ (f and g should be non-locally flat); or

(g) each $k+3 \leq m \leq 2k$ and some polyhedron or even manifold K.

The case (a) above was proved using the number of connected components of $\partial R_M(fK)$; the cases (c) and (d) was and will be proved using $\pi_1(\partial R_M(fK))$.

Proof of Theorem 1. If a PL embedding $S^2 \subset S^4$ is locally flat everywhere except at n points with singularities L_1, \ldots, L_n , then $\partial R_{S^4}(S^2)$ is obtained from ∂D^4 by the Dehn surgery over the knot $L_1 \# \dots \# L_n$ with certain framing (the details are analogous to the proof of Theorem 2.a below). Hence the case k = 2 of Theorem 1.b follows by a deep result of Gabai [Gab86, Gab87; Remark after Corollary 5], to the effect that the Dehn surgery along any non-trivial knot $S^1 \subset S^3$ can never yield $S^1 \times S^2$. (The argument of [Gab86, Gab87] uses foliation theory and cannot be generalized to obtain Theorem 1.b for k > 2.)

Theorem 1.b for k > 2 and Theorem 1.a are implied by the following result.

Theorem 2. For $k \geq 2$ and given embedding $S^k \subset S^{k+2}$ set $N = R_{S^{k+2}}(S^k)$. Then $G := \pi_1(S^{k+1} - S^{k-1})$ maps into $\pi_1(\partial N)$ in each of the following two cases:

(a) the embedding $S^k \subset S^{k+2}$ is the suspension over a locally flat PL knot $S^{k-1} \subset S^{k+1}$. (b) $k \geq 3$ and the embedding $S^k \subset S^{k+2}$ has an isolated non-locally flat point with the singularity $S^{k-1} \subset S^{k+1}$.



Figure 1.a

Proof of Theorem 2.a. Consider decomposition $S^{k+2} = D_+^{k+2} \bigcup_{\partial D_+^{k+2} = S^{k+1}} D_-^{k+2}$. Let

 $B^k_\pm:=D^{k+2}_\pm\cap S^k$ be the cone over S^{k-1} (Figure 1.a). Denote

$$N_{\pm} := R_{D_{\pm}^{k+2}}(B_{\pm}^k)$$
 so that $N_{\pm} \cap \partial D_{\pm}^{k+2} = R_{S^{k+1}}(S^{k-1})$ and

 $\partial_{\pm}N := \partial N_{\pm} - \operatorname{Int} R_{\partial N_{\pm}}(S^{k-1})$ so that $N = N_{+} \cup N_{-}$ and $\partial N = \partial_{+}N \cup \partial_{-}N$.

Since B_{\pm}^k is a cone, it follows that D_{\pm}^{k+2} collapses to B_{\pm}^k . Therefore D_{\pm}^{k+2} is also a regular neighborhood of B_{\pm}^k in D_{\pm}^{k+2} . Hence by the uniqueness of regular neighborhoods,

$$(\partial N_{\pm}, S^{k-1}) \cong (S^{k+1}, S^{k-1}), \text{ thus } \pi_1(\partial_{\pm}N) \cong G.$$

The composition $\partial_+ N \xrightarrow{i} \partial_+ N \cup \partial_- N \xrightarrow{r} \partial_+ N$ of the inclusion *i* and the 'symmetric' retraction *r* is a homeomorphism. So the induced composition $G \xrightarrow{i_*} \pi_1(\partial N) \xrightarrow{r_*} G$ is an isomorphism. Hence i_* is a monomorphism, and (a) is proved. \Box



Figure 1.b

Proof of Theorem 2.b. Let x be the isolated non-locally flat point. Then the singularity $S^{k-1} \subset S^{k+1}$ is locally flat. Consider the decomposition of S^{k+2} as above such that $x \in \mathring{D}^{k+2}_+$. Let $B^k_+ := D^{k+2}_+ \cap S^k$ be the cone over S^{k-1} with the vertex x (Figure 1.b). Let $B^k_- := D^{k+2}_- \cap S^k$ and $N_+ := D^{k+2}_+$. Define N_- and $\partial_{\pm}N$ as in the proof of (a). Then

 $N \cong N_+ \cup N_-$ and $\partial N = \partial_+ N \cup \partial_- N.$

We have

$$\pi_1(\partial_+ N) \cong G$$
 and $\pi_1(\partial_+ N \cap \partial_- N) \cong \pi_1(S^1 \times \partial D^k) \cong \mathbb{Z}$ for $k \ge 3$.

First consider the simple case when x is the only non-locally flat point (formally, the proof of the general case does not use the simpler case). Then B_{-}^{k} is locally flat. Since $N_{-} = R_{D_{-}^{k+2}}(B_{-}^{k})$, it follows that the pair (N_{-}, B_{-}^{k}) is standard [Zee63]. Therefore $\partial_{-}N \cong S^{1} \times D^{k}$. Hence $\pi_{1}(\partial N) \cong G$ by the van Kampen Theorem.

In the general case denote by a the generator of $\pi_1(\partial_+ N \cap \partial_- N) \cong H_1(\partial_+ N \cap \partial_- N; \mathbb{Z}) \cong \mathbb{Z}$. Observe that a is represented by a small circle in $\partial R_{\partial N_{\pm}}(S^{k-1})$, bounding a small 2disk in $R_{\partial N_{\pm}}(S^{k-1})$, transversal to S^{k-1} . By [Hud69; p. 57], $\partial N_- \cong S^{k+1}$. Hence the class of a is the generator of $H_1(\partial_{\pm}N,\mathbb{Z}) \cong \mathbb{Z}$. Thus the maps of $H_1(\cdot;\mathbb{Z})$ induced by the inclusions $\partial_+ N \cap \partial_- N \to \partial_{\pm} N$ are injective. Therefore the maps of π_1 unduced by the inclusions are also injective. Hence by the van Kampen Theorem [Rol76, pages 370-371] the inclusion-induced map $\pi_1(\partial_+N) \to \pi_1(\partial N)$ is also injective. \Box

We remark that under the assumptions of Theorem 2 the group $\pi_1(\partial N)$ maps onto $G := \pi_1(S^{k+1} - S^{k-1}).$

We remark that for $k \geq 3$ there is a PL embedding $S^k \subset S^{k+2}$ which is locally flat except at one point and such that $R_{S^{k+2}}(S^k) \not\cong S^k \times D^2$. Indeed, there exist slice knots $S^{k-1} \subset S^{k+1}$ having non-cyclic $\pi_1(S^{k+1} - S^{k-1})$ [Ker65], and for each slice knot $S^{k-1} \subset S^{k+1}$ there is an embedding $S^k \subset S^{k+2}$, having only one non locally-flat point with the singularity $S^{k-1} \subset S^{k+1}$ [FoMi66]. Now the remark follows by (the simple case of) Theorem 2.b. The remark also follows from Theorem 2.a because the regular neighborhood of the suspension over a knot L homeomorphic to that the regular neighborhood of certain knot with the only singular point (having singularity L#(-L)). Or else cf. [Gab86, Gab87; Remark after Corollary 5].

Theorem 1 was announced in [ORS01] and at the International Conference on Knots, Links and Manifold (Siegen, January 2001). We would like to acknowledge J. Hillman, A. A. Klyachko, W. B. R. Lickorish, E. Rafikov, M. A. Shtanko, J. Vrabec, and the referee for comments and suggestions.

References

- [Cas65] B. G. Casler, An embedding theorem for connected 3-manifolds with boundary, Proc. Amer. Math. Soc. 16 (1965), 559–556.
- [Cav92] A. Cavicchioli, Imbeddings of polyhedra in 3-manifolds, Ann. Mat. Pura Appl. 162 (1992), 157–177.
- [CaHe94] A. Cavicchioli and F. Hegenbarth, Knot manifolds with isomorphic spines, Fund. Math. 145 (1994), 79–89.
- [CHR98] A. Cavicchioli, F. Hegenbarth and D. Repovš, On manifold spines and cyclic presentation of groups, Knot theory, Banach Center Publ. 42 (PWN, Warsaw, 1998), 49-56.
- [CLR97] A. Cavicchioli, W. B. R. Lickorish and D. Repovš, On the equivalent spines problem, Boll. Unione Mat. Ital. A 11 (1997), 775–788.
- [FoMi66] R. H. Fox and J. Milnor, Singularities of 2-spheres in 4-space and equivalence of knots, Osaka J. Math. 3 (1966), 257–267.
- [Gab86] D. Gabai, Foliations and surgery on knots, Bull. Amer. Math. Soc. 15 (1986), 83–87.
- [Gab87] D. Gabai, Foliations and the topology of 3-manifolds. III., J. Diff. Geom. 26 (1987), 479–536.
- [HaHi62] A. Haefliger and M. W. Hirsch, Immersions in the stable range, Ann. of Math. (2) 75 (1962), 231–241.
- [Hor85] K. Horvatić, Regular neighborhoods of homeomorphic polyhedra, Berichte D. Math.-Stat. Sektion im Forschungszentrum Graz 251 (1985), 1–16.

- [Hor90] K. Horvatić, *Classical Problems of Geometric Topology*, Tehnička knjiga, Zagreb, 1990, (in Croatian).
- [HLS65] W. C. Hsiang, J. Levine and R. H. Sczarba, On the normal bundle of a homotopy sphere embedded in Euclidean space, Topology 3 (1965), 173–181.
- [Hud69] J. F. P. Hudson, *Piecewise-Linear Topology*, Benjamin, New York, Amsterdam, 1969.
- [Ker59] M. A. Kervaire, An interpretation of G. Whitehead's generalization of H. Hopf's invariant, Ann. of Math. (2) 62 (1959), 345–362.
- [Ker65] M. A. Kervaire, On higher dimensional knots, S. S. Cairns, Ed., Princeton Univ. Press, Princeton, 1965, 55–62.
- [LiSi69] W. B. R. Lickorish and L. C. Siebenmann, Regular neighborhoods and the stable range, Trans. Amer. Math. Soc. 139 (1969), 207–230.
- [Mas59] W. S. Massey, On the normal bundle of a sphere imbedded in Euclidean space, Proc. Amer. Math. Soc. 10 (1959), 959–964.
- [Mas69] W. S. Massey, Pontryagin squares in the Thom space of a bundle, Pacific J. Math. **31** (1969), 133–142.
- [Mat73] S. V. Matveev, Special skeletons of PL manifolds, Mat. Sbornik 92 (1973), 282–293 (in Russian).
- [MiRe71] R. J. Milgram and E. Rees, On the normal bundle to an embedding, Topology 10 (1971), 299–308.
- [MPR89] W. J. R. Mitchell, J. H. Przytycki and D. Repovš, On spines of knot spaces, Bull. Acad. Polon. Sci. 37:7–12 (1989), 563–565.
- [ORS01] A. Onischenko, D. Repovš and A. Skopenkov, Contractible resolutions of 2-polyhedra and embeddings into R⁴, Contemporary Math. 288 (2001), 396–400.
- [Rep88] D. Repovš, Regular neighborhoods of homotopically PL embedded compacta in 3-manifolds, Proc. Conf. Topol. Trieste 1986, Suppl. Rend. Circ. Mat. Palermo 18 (1988), 415–422.
- [RBS99] D. Repovš, N. Brodsky and A. B. Skopenkov, A classification of 3-thickenings of 2-polyhedra, Topol. Appl. 94 (1999), 307–314.
- [Roh70] V. A. Rohlin, On normal Euler numbers of the projective plane and the Klein bottle in the four-dimensional Euclidean space, Dokl. Akad. Nauk SSSR 191 (1970), 27–29 (in Russian).
- [Rol76] D. Rolfsen, Knots and Links, Publish or Perish, Berkeley, California, 1976.
- [RoSa68] C. P. Rourke and B. J. Sanderson, *Block bundles I*, Ann. of Math. (2) 87 (1968), 1–28.
- [RoSa72] C. P. Rourke and B. J. Sanderson, Introduction to Piecewise-Linear Topology, Ergebn. der Math. 69, Springer-Verlag, Berlin, 1972.
- [Wal67] C. T. C. Wall, Locally flat PL submanifolds with codimension two, Proc. Camb. Phil. Soc. 63 (1967), 5–8.
- [Zee63] E. C. Zeeman, Unknotting combinatorial balls, Ann. of Math. (2) 78 (1963), 501–526.
- [Zee64] E. C. Zeeman, On the dunce hat, Topology 2 (1964), 341–358.

INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, UNIVERSITY OF LJUBLJANA, P. O. BOX 2964, 1001 LJUBLJANA, SLOVENIA. E-MAIL: DUSAN.REPOVS@FMF.UNI-LJ.SI

INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, UNIVERSITY OF LJUBLJANA, P. O. BOX 2964, 1001 LJUBLJANA, SLOVENIA. E-MAIL: MATIJA.CENCELJ@FMF.UNI-LJ.SI

DEPARTMENT OF DIFFERENTIAL GEOMETRY, FACULTY OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY, MOSCOW, RUSSIA 119992. E-MAIL: SKOPENKO@MCCME.RU