

# On prime factors of class number of cyclotomic fields

Roland Quême

2006 nov 27

## Contents

<b>1</b>	<b>Some definitions</b>	<b>2</b>
<b>2</b>	<b>On Kummer and Stickelberger relation</b>	<b>2</b>
<b>3</b>	<b>Polynomial congruences mod <math>p</math> connected to the <math>p</math>-class group <math>C_p</math></b>	<b>5</b>
3.1	Stickelberger relation for prime ideals $\mathfrak{q}$ of inertial degree $f = 1$ . . . . .	5
3.2	Stickelberger relation for prime ideals $\mathfrak{q}$ of inertial degree $f > 1$ . . . . .	7
<b>4</b>	<b>On prime factors <math>h \neq p</math> of the class number of the <math>p</math>-cyclotomic field</b>	<b>10</b>
4.1	The general case . . . . .	11
4.2	The case of complex quadratic fields contained in $K$ . . . . .	16
4.3	The case of biquadratic fields contained in $K$ . . . . .	18

### Abstract

Let  $p$  be an odd prime. Let  $K = \mathbb{Q}(\zeta)$  be the  $p$ -cyclotomic number field. Let  $v$  be a primitive root mod  $p$  and  $\sigma : \zeta \rightarrow \zeta^v$  be a  $\mathbb{Q}$ -isomorphism of the extension  $K/\mathbb{Q}$  generating the Galois group  $G$  of  $K/\mathbb{Q}$ . Following the conventions of Ribenboim in [8], for  $n \in \mathbb{Z}$ , the notation  $v_n$  is understood by  $v_n = v^n \pmod p$  with  $1 \leq v_n \leq p - 1$ . Let  $P(\sigma) = \sum_{i=0}^{p-2} \sigma^i v_{-i} \in \mathbb{Z}[G]$  be the Stickelberger polynomial. We show that there exists a polynomial  $Q(\sigma) \in \mathbb{Z}[G]$  such that  $P(\sigma) \times (\sigma - v) = p \times Q(\sigma)$  and such that  $Q(\sigma)$  annihilates the class group  $\mathbf{C}$  of  $K$  ( $\mathfrak{a}^{Q(\sigma)}$  is a principal ideal for all ideals  $\mathfrak{a}$  of  $K$ ). This result brings some informations on the structure of the class group  $\mathbf{C}$  and gives:

1. some explicit congruences in  $\mathbb{Z}[v] \pmod p$  for the  $p$ -class group of  $K$  (the subgroup of exponent  $p$  of  $\mathbf{C}$ ),

2. some explicit congruences in  $\mathbb{Z}[v] \bmod h$  for the  $h$ -class group of  $K$  for the prime divisors  $h \neq p$  of the class number  $h(K)$ . We detail at the end the case of class number of quadratic and biquadratic fields contained in the cyclotomic field  $K$ .

This article is at elementary level.

*Remark:* we have not found in the literature some formulations corresponding to theorems 3.4 p. 8, 4.1 p. 11, 4.3 p. 13, 4.6 p. 17, 4.7 p. 17, and 4.8 p. 18.

## 1 Some definitions

In this section we give some definitions and notations on cyclotomic fields and  $p$ -class group used in this paper.

1. Let  $p$  be an odd prime. Let  $\mathbf{F}_p$  be the finite field of cardinal  $p$  and  $\mathbf{F}_p^*$  its multiplicative group. Let  $\zeta$  be a root of the polynomial equation  $X^{p-1} + X^{p-2} + \dots + X + 1 = 0$ . Let  $K$  be the  $p$ -cyclotomic field  $K = \mathbb{Q}(\zeta)$  and  $O_K$  its ring of integers. Let  $K^+$  be the maximal totally real subfield of  $K$ . Let  $v$  be a primitive root  $\bmod p$ . In this paper, following Ribenboim conventions in Ribenboim [8] for any  $n \in \mathbb{Z}$ , we note  $v_n = v^n \bmod p$  with  $1 \leq v_n \leq p - 1$ . Let  $G$  be the Galois group of the extension  $K/\mathbb{Q}$ . Let  $\sigma : \zeta \rightarrow \zeta^v$  be a  $\mathbb{Q}$ -isomorphism of the extension  $K/\mathbb{Q}$  generating  $G$ . Let  $\lambda = \zeta - 1$ . The prime ideal of  $K$  lying over  $p$  is  $\pi = \lambda O_K$ .
2. Let  $\mathbf{C}$  be the class group of  $K$ . Let  $C_p$  be the  $p$ -class of  $K$  (the subgroup of exponent  $p$  of  $\mathbf{C}$ ). Let  $C_p^+$  be the  $p$ -class group of  $K^+$ . Then  $C_p = C_p^+ \oplus C_p^-$  where  $C_p^-$  is called the relative  $p$ -class group. Let  $r^-$  be the rank of  $C_p^-$ .
3.  $C_p^-$  is the direct sum of  $r^-$  subgroups  $\Gamma_k$  of order  $p$  annihilated by  $\sigma - \mu_k \in \mathbf{F}_p[G]$  with  $\mu_k \in \mathbf{F}_p^*$ ,  $\mu_k = v_{2m_k+1}$  where  $m_k$  is a natural integer  $m_k$ ,  $1 \leq m_k \leq \frac{p-3}{2}$ ,

$$(1) \quad C_p^- = \bigoplus_{k=1}^{r^-} \Gamma_k.$$

## 2 On Kummer and Stickelberger relation

Stickelberger relation was already known by Kummer under the form of Jacobi resolvents for the cyclotomic field  $K$ , see for instance Ribenboim [8] (2.6) p. 119. In this section we derive some elementary properties from Stickelberger relation.

1. Let  $q \neq p$  be an odd prime. Let  $\zeta_q$  be a root of the minimal polynomial equation  $X^{q-1} + X^{q-2} + \dots + X + 1 = 0$ . Let  $K_q = \mathbb{Q}(\zeta_q)$  be the  $q$ -cyclotomic field. Let  $K_{pq} = \mathbb{Q}(\zeta_p, \zeta_q)$ . Then  $K_{pq}$  is the compositum  $KK_q$ . The ring of integers of  $K_{pq}$  is  $O_{K_{pq}}$ .

2. Let  $\mathfrak{q}$  be a prime ideal of  $O_K$  lying over the prime  $q$ . Let  $f$  be the order of  $q \bmod p$  and  $m = N_{K_p/\mathbb{Q}}(\mathfrak{q}) = q^f$ . If  $\psi(\alpha) = a$  is the image of  $\alpha \in O_K$  under the natural map  $\psi : O_K \rightarrow O_K/\mathfrak{q}$ , then for  $\psi(\alpha) = a \not\equiv 0$  define a character  $\chi_{\mathfrak{q}}^{(p)}$  on  $\mathbf{F}_m = O_K/\mathfrak{q}$  by

$$(2) \quad \chi_{\mathfrak{q}}^{(p)}(a) = \left\{ \frac{\alpha}{\mathfrak{q}} \right\}_p^{-1} = \overline{\left\{ \frac{\alpha}{\mathfrak{q}} \right\}_p},$$

where  $\left\{ \frac{\alpha}{\mathfrak{q}} \right\} = \zeta^c$  for some natural integer  $c$ , is the  $p^{\text{th}}$  power residue character mod  $\mathfrak{q}$ . We define the Gauss sum

$$(3) \quad g(\mathfrak{q}) = \sum_{x \in \mathbf{F}_m} (\chi_{\mathfrak{q}}^{(p)}(x) \times \zeta_q^{\text{Tr}_{\mathbf{F}_m/\mathbf{F}_q}(x)}) \in O_{K_{pq}}.$$

It follows that  $\mathbf{g}(\mathfrak{q}) \in O_{K_{pq}}$ . Moreover  $g(\mathfrak{q})^p \in O_K$ , see for instance Mollin [6] prop. 5.88 (c) p. 308.

3. The Stickelberger relation is classically:

$$(4) \quad \mathbf{g}(\mathfrak{q})^p O_K = \mathfrak{q}^S,$$

with  $S = \sum_{t=1}^{p-1} t \times \varpi_t^{-1}$ , where  $\varpi_t \in \text{Gal}(K/\mathbb{Q})$  is given by  $\varpi_t : \zeta \rightarrow \zeta^t$  (see for instance Mollin [6] thm. 5.109 p. 315).

The four following lemmas are derived in an elementary way from the Stickelberger relation.

**Lemma 2.1.** *If  $q \not\equiv 1 \pmod p$  then the Gauss sum  $g(\mathfrak{q}) \in \mathbb{Z}[\zeta]$ .*

*Proof.*

1. Let  $u$  be a primitive root mod  $q$ . Let  $\tau : \zeta_q \rightarrow \zeta_q^u$  be a  $\mathbb{Q}$ -isomorphism generating  $\text{Gal}(K_q/\mathbb{Q})$ . The  $\mathbb{Q}$ -isomorphism  $\tau$  is extended to a  $K_p$ -isomorphism of  $K_{pq}$  by  $\tau : \zeta_q \rightarrow \zeta_q^u, \zeta_p \rightarrow \zeta_p$ . Then  $g(\mathfrak{q})^p \in \mathbb{Z}[\zeta_p]$  and so

$$\tau(g(\mathfrak{q}))^p = g(\mathfrak{q})^p,$$

and it follows that there exists a natural integer  $\rho$  with  $\rho < p$  such that

$$\tau(g(\mathfrak{q})) = \zeta_p^\rho \times g(\mathfrak{q}).$$

Then  $N_{K_{pq}/K}(\tau(g(\mathfrak{q}))) = \zeta^{(q-1)\rho} \times N_{K_{pq}/K}(g(\mathfrak{q}))$  and so  $\zeta^{\rho(q-1)} = 1$ .

2. If  $q \not\equiv 1 \pmod p$ , it implies that  $\zeta^\rho = 1$  and so that  $\tau(g(\mathfrak{q})) = g(\mathfrak{q})$  and thus that  $g(\mathfrak{q}) \in O_K$ .

□

**Lemma 2.2.** Let  $S = \sum_{t=1}^{p-1} \varpi_t^{-1} \times t$  where  $\varpi_t$  is the  $\mathbb{Q}$ -isomorphism of the extension  $K/\mathbb{Q}$  given by  $\varpi_t : \zeta \rightarrow \zeta^t$  of  $K$ . Let  $P(\sigma) = \sum_{i=0}^{p-2} \sigma^i \times v_{-i} \in \mathbb{Z}[G]$ . Then  $S = P(\sigma)$ .

*Proof.* Let us consider one term  $\varpi_t^{-1} \times t$ . Then  $v_{-1} = v_{p-2}$  is a primitive root mod  $p$  and so there exists one and one  $i$  such that  $t = v_{-i}$ . Then  $\varpi_{v_{-i}} : \zeta \rightarrow \zeta^{v_{-i}}$  and so  $\varpi_{v_{-i}}^{-1} : \zeta \rightarrow \zeta^{v_i}$  and so  $\varpi_{v_{-i}}^{-1} = \sigma^i$  (observe that  $\sigma^{p-1} \times v_{-(p-1)} = 1$ ), which achieves the proof. □

**Lemma 2.3.**

$$(5) \quad P(\sigma) = \sum_{i=0}^{p-2} \sigma^i \times v_{-i} = v_{-(p-2)} \times \left\{ \prod_{k=0, k \neq 1}^{p-2} (\sigma - v_k) \right\} + p \times R(\sigma),$$

where  $R(\sigma) \in \mathbb{Z}[G]$  with  $\deg(R(\sigma)) < p - 2$ .

*Proof.* Let us consider the polynomial  $R_0(\sigma) = P(\sigma) - v_{-(p-2)} \times \left\{ \prod_{k=0, k \neq 1}^{p-2} (\sigma - v_k) \right\}$  in  $\mathbf{F}_p[G]$ . Then  $R_0(\sigma)$  is of degree smaller than  $p - 2$  and the two polynomials  $\sum_{i=0}^{p-2} \sigma^i v_{-i}$  and  $\prod_{k=0, k \neq 1}^{p-2} (\sigma - v_k)$  take a null value in  $\mathbf{F}_p[G]$  when  $\sigma$  takes the  $p - 2$  different values  $\sigma = v_k$  for  $k = 0, \dots, p - 2, \quad k \neq 1$ . Then  $R_0(\sigma) = 0$  in  $\mathbf{F}_p[G]$  which leads to the result in  $\mathbb{Z}[G]$ . □

**Lemma 2.4.**

$$(6) \quad P(\sigma) \times (\sigma - v) = p \times Q(\sigma),$$

where  $Q(\sigma) = \sum_{i=1}^{p-2} \delta_i \times \sigma^i \in \mathbb{Z}[G]$  is given by

$$(7) \quad \begin{aligned} \delta_{p-2} &= \frac{v_{-(p-3)} - v_{-(p-2)}v}{p}, \\ \delta_{p-3} &= \frac{v_{-(p-4)} - v_{-(p-3)}v}{p}, \\ &\vdots \\ \delta_i &= \frac{v_{-(i-1)} - v_{-i}v}{p}, \\ &\vdots \\ \delta_1 &= \frac{1 - v_{-1}v}{p}, \end{aligned}$$

with  $-p < \delta_i \leq 0$ .

*Proof.* We start of the relation in  $\mathbb{Z}[G]$

$$P(\sigma) \times (\sigma - v) = v_{-(p-2)} \times \prod_{k=0}^{p-2} (\sigma - v_k) + p \times R(\sigma) \times (\sigma - v) = p \times Q(\sigma),$$

with  $Q(\sigma) \in \mathbb{Z}[G]$  because  $\prod_{k=0}^{p-2} (\sigma - v_k) = 0$  in  $\mathbf{F}_p[G]$  and so  $\prod_{k=0}^{p-2} (\sigma - v_k) = p \times R_1(\sigma)$  in  $\mathbb{Z}[G]$ . Then we identify in  $\mathbb{Z}[G]$  the coefficients in the relation

$$(v_{-(p-2)}\sigma^{p-2} + v_{-(p-3)}\sigma^{p-3} + \cdots + v_{-1}\sigma + 1) \times (\sigma - v) = p \times (\delta_{p-2}\sigma^{p-2} + \delta_{p-3}\sigma^{p-3} + \cdots + \delta_1\sigma + \delta_0),$$

where  $\sigma^{p-1} = 1$ . □

**Remark:**

1. Observe that, with our notations,  $\delta_i \in \mathbb{Z}$ ,  $i = 1, \dots, p-2$ , but generally  $\delta_i \not\equiv 0 \pmod{p}$ .
2. We see also that  $-p < \delta_i \leq 0$ . Observe also that  $\delta_0 = \frac{v^{-(p-2)} - v}{p} = 0$ .

### 3 Polynomial congruences mod $p$ connected to the $p$ -class group $C_p$

We give some explicit polynomial congruences in  $\mathbb{Z}[v]$  mod  $p$  connected to the relative  $p$ -class group  $C_p^-$  of  $O_K$ . We apply successively the Stickelberger relation to prime ideals  $\mathfrak{q}$  of inertial degree  $f = 1$  and of inertial degree  $f > 1$ . As an application example we give an elementary straightforward proof that if  $\frac{p-1}{2}$  is odd then the Bernoulli Number  $B_{(p+1)/2} \not\equiv 0 \pmod{p}$ . We recall that  $r^-$  is the  $p$ -rank of the relative  $p$ -class group  $C_p^-$  of  $K$ .

#### 3.1 Stickelberger relation for prime ideals $\mathfrak{q}$ of inertial degree $f = 1$

**Theorem 3.1.**  $\prod_{k=1}^{r^-} (\sigma - v_{2m_k+1})$  divides  $Q(\sigma)$  in  $\mathbf{F}_p[G]$  and

$$(8) \quad Q(v_{2m_k+1}) = \sum_{i=1}^{p-2} v_{(2m_k+1) \times i} \times \left( \frac{v^{-(i-1)} - v^{-i} \times v}{p} \right) \equiv 0 \pmod{p},$$

where  $m_k$  is defined in relation (1) p. 2.

*Proof.* From Kummer, the group of ideal classes of  $K$  is generated by the classes of prime ideals of degree 1 (see for instance Ribenboim [8] (3A) p. 119). Let  $\mathfrak{q}$  a prime ideal of inertial degree 1 whose class  $Cl(\mathfrak{q}) \in C_p^-$  is annihilated by  $\sigma - \mu_k$  with  $\mu_k = v_{2m_k+1}$ . We start of  $g(\mathfrak{q})^p O_K = \mathfrak{q}^{P(\sigma)}$  and so  $g(\mathfrak{q})^{p(\sigma-v)} O_K = \mathfrak{q}^{P(\sigma)(\sigma-v)} = \mathfrak{q}^{pQ(\sigma)}$  and thus  $g(\mathfrak{q})^{(\sigma-v)} O_K = \mathfrak{q}^{Q(\sigma)}$ . It can be shown that  $g(\mathfrak{q})^{\sigma-v} \in K$ , see for instance Ribenboim [9] F. p. 440. Therefore  $Q(\sigma)$  annihilates the ideal class  $Cl(\mathfrak{q})$  and the congruence follows.  $\square$

**Remarks:**

1. Observe that  $\delta_i$  can also be written in the form  $\delta_i = -[\frac{v-i \times v}{p}]$  where  $[x]$  is the integer part of  $x$ , similar form also known in the literature.
2. Observe that it is possible to get other polynomials of  $\mathbb{Z}[G]$  annihilating the relative  $p$ -class group  $C_p^-$ : for instance from Kummer's formula on Jacobi cyclotomic functions we induce other polynomials  $Q_d(\sigma)$  annihilating the relative  $p$ -class group  $C_p^-$  of  $K$ : If  $1 \leq d \leq p-2$  define the set

$$I_d = \{i \mid 0 \leq i \leq p-2, \quad v_{(p-1)/2-i} + v_{(p-1)/2-i+ind_v(d)} > p\}$$

where the index  $ind_v(d)$  is the minimal integer  $s$  such that  $d = v_s$ . Then the polynomials  $Q_d(\sigma) = \sum_{i \in I_d} \sigma^i$  for  $d = 1, \dots, p-2$  annihilate the  $p$ -class  $C_p^-$  of  $K$ , see for instance Ribenboim [8] relations (2.4) and (2.5) p. 119.

3. See also in a more general context Washington, [10] corollary 10.15 p. 198.
4. It is easy to verify the consistency of relation (8) with the table of irregular primes and Bernoulli numbers in Washington, [10] p. 410.

An immediate consequence is an explicit criterium for  $p$  to be a regular prime:

**Corollary 3.2.** *Let  $p$  be an odd prime. Let  $v$  be a primitive root mod  $p$ . If the congruence*

$$(9) \quad \sum_{i=1}^{p-2} X^{i-1} \times \left( \frac{v_{-(i-1)} - v_{-i} \times v}{p} \right) \equiv 0 \pmod{p}$$

*has no solution  $X$  in  $\mathbb{Z}$  with  $X^{(p-1)/2} + 1 \equiv 0 \pmod{p}$  then the prime  $p$  is regular.*

We give, as an example, a straightforward proof of following lemma on Bernoulli Numbers (compare elementary nature of this proof with proof hinted by Washington in exercise 5.9 p. 85 using Siegel-Brauer theorem).

**Lemma 3.3.** *If  $2m+1 = \frac{p-1}{2}$  is odd then the Bernoulli Number  $B_{(p+1)/2} \not\equiv 0 \pmod{p}$ .*

*Proof.* Suppose that  $B_{(p+1)/2} \equiv 0 \pmod p$  and search for a contradiction. From previous corollary  $B_{(p+1)/2} \equiv 0 \pmod p$  implies that with  $\sigma - v_{p-(p+1)/2} \mid Q(\sigma)$ , hence that  $\sum_{i=1}^{p-2} v_{(p-1)/2}^i \times \delta_i \equiv 0 \pmod p$ . From  $v_{(p-1)/2} = p - 1$  we get

$$\sum_{i=1}^{p-2} (-1)^i \times \left( \frac{v_{-(i-1)} - v_{-i} \times v}{p} \right) \equiv 0 \pmod p.$$

Multiplying by  $p$

$$\sum_{i=1}^{p-2} (-1)^i \times (v_{-(i-1)} - v_{-i} \times v) \equiv 0 \pmod{p^2},$$

expanded to

$$(-1 + v_{-1} - v_{-2} + \cdots - v_{-(p-3)}) + (v_{-1}v - v_{-2}v + \cdots + v_{-(p-2)}v) \equiv 0 \pmod{p^2}$$

also

$$(-1 + v_{-1} - v_{-2} + \cdots - v_{-(p-3)}) + (v_{-1} - v_{-2} + \cdots + v_{-(p-2)})v \equiv 0 \pmod{p^2}.$$

Let us set  $V = -1 + v_{-1} - v_{-2} + \cdots - v_{-(p-3)} + v_{-(p-2)}$ . Then we get  $V - v_{-(p-2)} + v(V + 1) \equiv 0 \pmod{p^2}$ , and so  $V(1 + v) + v - v_{-(p-2)} \equiv 0 \pmod{p^2}$ . But  $v = v_{-(p-2)}$  and so  $V \equiv 0 \pmod{p^2}$ . But

$$\begin{aligned} -V &= 1 - v_{-1} + v_{-2} + \cdots + v_{-(p-3)} - v_{-(p-2)} = S_1 - S_2 \\ S_1 &= 1 + v_{-2} + \cdots + v_{-(p-3)}, \\ S_2 &= v_{-1} + v_{-3} + \cdots + v_{-(p-2)}. \end{aligned}$$

$v_{-1}$  is a primitive root mod  $p$  and so  $S_1 + S_2 = \frac{p(p-1)}{2}$ . Clearly  $S_1 \neq S_2$  because  $\frac{p(p-1)}{2}$  is odd and so  $-V = S_1 - S_2 \neq 0$  and  $-V \equiv 0 \pmod{p^2}$  with  $|-V| < \frac{p(p-1)}{2}$ , contradiction which achieves the proof.  $\square$

### 3.2 Stickelberger relation for prime ideals $\mathfrak{q}$ of inertial degree $f > 1$

Let  $\mathfrak{q}$  be a prime ideal of  $O_K$  with  $Cl(\mathfrak{q}) \in C_p$ . In this section we apply Stickelberger relation to the prime ideals  $\mathfrak{q}$  of inertial degree  $f > 1$  with the method used for the prime ideals of inertial degree 1 in section 3 p. 5. Observe, from lemma 2.1 p. 3, that  $f > 1$  implies that  $g(\mathfrak{q}) \in O_K$ , property used in this section (by opposite  $g(\mathfrak{q}) \notin K$  when  $f = 1$ ).

**A definition:** we say that the prime ideal  $\mathfrak{c}$  of a number field  $M$  is  $p$ -principal if the component of the class group  $\langle Cl(\mathfrak{c}) \rangle$  in  $p$ -class group  $D_p$  of  $M$  is trivial.

**Theorem 3.4.** *Let  $q$  be an odd prime with  $q \neq p$ . Let  $f$  be the order of  $q$  mod  $p$  and  $m = \frac{p-1}{f}$ . Let  $\mathfrak{q}$  be a prime ideal of  $O_K$  lying over  $q$ . If  $f > 1$  then*

1.  $g(\mathfrak{q}) \in O_K$  and  $g(\mathfrak{q})O_K = \mathfrak{q}^{P_1(\sigma)}$  where

$$(10) \quad P_1(\sigma) = \sum_{i=0}^{m-1} \left( \frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times \sigma^i \in \mathbb{Z}[G].$$

2. *There exists a natural integer  $n$  with  $1 \leq n \leq p-2$  such that  $\sigma - v_n$  divides  $P_1(\sigma)$  in  $\mathbf{F}_p[G]$  and  $\sigma - v_n$  annihilates  $Cl(\mathfrak{q})$ .*

*Proof.*

1.  $N_{K/\mathbb{Q}}(\mathfrak{q}) = q^f$  and  $\mathfrak{q} = \mathfrak{q}^{\sigma^m} = \dots = \mathfrak{q}^{\sigma^{(f-1)m}}$ . From Stickelberger relation  $g(\mathfrak{q})^p \mathbb{Z}[\zeta_p] = \mathfrak{q}^{P(\sigma)}$  where  $P(\sigma) = \sum_{i=0}^{m-1} \sum_{j=0}^{f-1} \sigma^{i+jm} v_{-(i+jm)}$ . Observe that, from hypothesis,  $\mathfrak{q} = \mathfrak{q}^{\sigma^m} = \dots = \mathfrak{q}^{\sigma^{(f-1)m}}$  so Stickelberger's relation implies that  $g(\mathfrak{q})^p O_K = \mathfrak{q}^{P(\sigma)}$  with

$$P(\sigma) = \sum_{i=0}^{m-1} \sum_{j=0}^{f-1} \sigma^i v_{-(i+jm)} = p \times \sum_{i=0}^{m-1} \left( \frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times \sigma^i,$$

where  $(\sum_{i=0}^{f-1} v_{-(i+jm)})/p \in \mathbb{Z}$  because  $v_{-m} - 1 \not\equiv 0 \pmod{p}$ .

2. Let  $P_1(\sigma) = \frac{P(\sigma)}{p}$ . Then from below  $P_1(\sigma) \in \mathbb{Z}[G]$ . From lemma 2.1 p. 3 we know that  $f > 1$  implies that  $g(\mathfrak{q}) \in O_K$ . Therefore

$$g(\mathfrak{q})^p O_K = \mathfrak{q}^{pP_1(\sigma)}, \quad g(\mathfrak{q}) \in O_K,$$

and so

$$g(\mathfrak{q})O_K = \mathfrak{q}^{P_1(\sigma)}, \quad g(\mathfrak{q}) \in O_K.$$

Then  $P_1(\sigma)$  annihilates  $Cl(\mathfrak{q})$  and thus there exists  $1 < n \leq p-2$  such that  $(\sigma - v_n) \mid P_1(\sigma)$ . □

## Remarks

1. For  $f = 2$  the value of polynomial  $P_1(\sigma)$  obtained from this lemma is  $P_1(\sigma) = \sum_{i=1}^{(p-3)/2} \sigma^i$ .
2. Let  $\mathfrak{q}$  be a prime not principal ideal of inertial degree  $f > 1$  with  $Cl(\mathfrak{q}) \in C_p$ . The **two** polynomials of  $\mathbb{Z}[G_p]$ ,  $Q(\sigma) = \sum_{i=0}^{p-2} \left( \frac{v_{-(i-1)} - v_{-i}}{p} \right) \times \sigma^i$  (see thm 3.1) and  $P_1(\sigma) = \sum_{i=0}^{m-1} \left( \frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times \sigma^i$  (see lemma 3.4) annihilate the ideal class  $Cl(\mathfrak{q})$ . When  $f > 1$  the lemma 3.4 p. 8 supplement the theorem 3.1 p. 5.
3. This result explains that, when  $f$  increases, the proportion of  $p$ -principal ideals  $\mathfrak{q}$  increases. For instance when  $f = \frac{p-1}{2}$  then all the prime ideals  $\mathfrak{q}$  are  $p$ -principal (see corollary 3.6 p. 10).

It is possible to derive some explicit congruences in  $\mathbb{Z}$  from this theorem.

**Corollary 3.5.** *Let  $q$  be an odd prime with  $q \neq p$ . Let  $f$  be the order of  $q \bmod p$  and let  $m = \frac{p-1}{f}$ . Let  $\mathfrak{q}$  be an prime ideal of  $O_K$  lying over  $q$ . Suppose that  $f > 1$ .*

1. *If  $\mathfrak{q}$  is not  $p$ -principal ideal there exists a natural integer  $l$ ,  $1 \leq l < m$  such that*

$$(11) \quad \sum_{i=0}^{m-1} \left( \frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times v^{lfi} \equiv 0 \pmod{p},$$

2. *If for all natural integers  $l$  such that  $1 \leq l < m$*

$$(12) \quad \sum_{i=0}^{m-1} \left( \frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times v^{lfi} \not\equiv 0 \pmod{p},$$

*then  $\mathfrak{q}$  is  $p$ -principal*

*Proof.*

1. Suppose that  $\mathfrak{q}$  is not  $p$ -principal. Observe at first that congruence (11) with  $l = m$  should imply that  $\sum_{i=0}^{m-1} (\sum_{j=0}^{f-1} v_{-(i+jm)})/p \equiv 0 \pmod{p}$  or

$$\sum_{i=0}^{m-1} \sum_{j=0}^{f-1} v_{-(i+jm)} \equiv 0 \pmod{p^2},$$

which is not possible because  $v_{-(i+jm)} = v_{-(i'+j'm)}$  implies that  $j = j'$  and  $i = i'$  and so that  $\sum_{i=0}^{m-1} \sum_{j=0}^{f-1} v_{-(i+jm)} = \frac{p(p-1)}{2}$ .

2. The polynomial  $P_1(\sigma)$  of lemma 3.4 annihilates the non  $p$ -principal ideal  $\mathfrak{q}$  in  $\mathbf{F}_p[G]$  only if there exists  $\sigma - v_n$  dividing  $P_1(\sigma)$  in  $\mathbf{F}_p[G]$ . From  $\mathfrak{q}^{\sigma^{m-1}} = 1$  it follows also that  $\sigma - v_n \mid \sigma^m - 1$ . But  $\sigma - v_n \mid \sigma^m - v_{nm}$  and so  $\sigma - v_n \mid v_{nm} - 1$ , thus  $nm \equiv 0 \pmod{p-1}$ , so  $n \equiv 0 \pmod{f}$  and  $n = lf$  for some  $l$ . Therefore if  $\mathfrak{q}$  is non  $p$ -principal there exists a natural integer  $l$ ,  $1 \leq l < m$  such that

$$(13) \quad \sum_{i=0}^{m-1} \left( \frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times v^{lfi} \equiv 0 \pmod{p},$$

3. The relation (12) is an immediate consequence of previous part of the proof.  $\square$

As an example we deal with the case  $f = \frac{p-1}{2}$ .

**Corollary 3.6.** *If  $p \equiv 3 \pmod{4}$  and if  $f = \frac{p-1}{2}$  then  $\mathfrak{q}$  is  $p$ -principal.*

*Proof.* We have  $f = \frac{p-1}{2}$ ,  $m = 2$  and  $l = 1$ . Then

$$(14) \quad \Sigma = \sum_{i=0}^{m-1} \left( \frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times v^{lfi} = \frac{\sum_{j=0}^{(p-3)/2} v_{-2j}}{p} - \frac{\sum_{j=0}^{(p-3)/2} v_{-(1+2j)}}{p}.$$

$\Sigma \equiv 0 \pmod{p}$  should imply that  $\sum_{j=0}^{(p-3)/2} v_{-2j} - \sum_{j=0}^{(p-3)/2} v_{-(1+2j)} \equiv 0 \pmod{p^2}$ . But  $\sum_{j=0}^{(p-3)/2} v_{-2j} + \sum_{j=0}^{(p-3)/2} v_{-(1+2j)} = \frac{p(p-1)}{2}$  is odd, which achieves the proof.  $\square$

**Remark:** this result can also be found directly because the quadratic subfield  $K_2$  of  $K$  is  $p$ -principal and that when  $f = \frac{p-1}{2}$  then  $\mathfrak{q} = \mathfrak{q}_2 O_K$  where  $\mathfrak{q}_2$  is a prime ideal of  $K_2$ .

## 4 On prime factors $h \neq p$ of the class number of the $p$ -cyclotomic field

In previous sections we considered the relative  $p$ -class group  $C_p^-$  of  $K$ . By opposite, in this section we apply Stickelberger relation to all the primes  $h \neq p$  dividing the class number  $h(K)$ . A first subsection is devoted to the general case of the relative class group  $\mathbf{C}^-$  of  $K$ , a second to the class group of the quadratic subfield of  $K$  and a third subsection to the class group at the biquadratic subfield of  $K$  when  $p \equiv 1 \pmod{4}$ .

## 4.1 The general case

1. The class group  $\mathbf{C}$  of  $K_p$  is the direct sum of the class group  $\mathbf{C}^+$  of the maximal totally real subfield  $K^+$  of  $K$  and of the relative class group  $\mathbf{C}^-$  of  $K$ .
2. Remind that  $v$  is a primitive root mod  $p$  and that  $v_n$  is to be understood as  $v^n \pmod{p}$  with  $1 \leq v_n \leq p-1$ . Let  $h(K)$  be the class number of  $K$ . Let  $h \neq p$  be a prime dividing  $h(K_p)$ , with  $v_h(h(K)) = \beta$ . Let  $d = \text{Gcd}(h-1, p-1)$ . Let  $\mathbf{C}(h)$  be the  $h$ -Sylow subgroup of the class group of  $K$  of order  $h^\beta$ . Then  $\mathbf{C}(h) = \bigoplus_{j=1}^{\rho} \mathbf{C}_j(h)$  where  $\rho$  is the  $h$ -rank of the abelian group  $\mathbf{C}(h)$  of order  $h^\beta$  and  $\mathbf{C}_j(h)$  are cyclic groups of order  $h^{\beta_j}$  where  $\beta = \sum_{i=1}^{\rho} \beta_j$ .
3. From Kummer (see for instance Ribenboim [8] (3A) p. 119), the prime ideals of  $O_K$  of inertial degree 1 generate the ideal class group. Therefore there exist in the subgroup of exponent  $h$  of  $\mathbf{C}(h)$  some prime ideals  $\mathfrak{q}$  of inertial degree 1 such that  $\langle Cl(\mathfrak{q}) \rangle = \bigoplus_{j=1}^{\rho} c_j$  where  $c_j$  is a cyclic group of order  $h$  and so that  $\langle Cl(\mathfrak{q}) \rangle$  is of order  $h^\rho$ .
4. Let  $P(\sigma) = \sum_{k=0}^{p-2} \sigma^k v_{-k}$  be the Stickelberger polynomial. From lemma 2.2 p. 4 Stickelberger relation is  $\mathfrak{q}^{P(\sigma)} = g(\mathfrak{q})^p O_K$  where  $g(\mathfrak{q})^p \in O_K$ . Therefore  $\mathfrak{q}^{P(\sigma)}$  is principal, a fortiori is  $\mathbf{C}(h)$ -principal (or  $Cl(\mathfrak{q})^{P(\sigma)}$  has a trivial component in  $\mathbf{C}(h)$ ). There exists a minimal polynomial  $V(X) \in \mathbf{F}_h(X)$  of degree  $\delta \leq \rho$  such that  $\mathfrak{q}^{V(\sigma)}$  is  $\mathbf{C}(h)$ -principal, if not the remainder of the division of  $P(X)$  by  $V(X)$  of degree smaller than  $\delta$  would annihilate also  $\mathfrak{q}$ . Therefore the irreducible polynomial  $V(X)$  divides  $P(X)$  in  $\mathbf{F}_h[X]$  for the indeterminate  $X$ .
5. If  $Cl(\mathfrak{q}) \in \mathbf{C}^-$  then  $\mathfrak{q}^{\sigma^{(p-1)/2+1}}$  is principal.
6. Let  $D(X) \in \mathbf{F}_h(X)$  defined by  $D(X) = \text{Gcd}(P(X), X^{(p-1)/2} + 1) \pmod{h}$ .

We obtain the following:

**Theorem 4.1.** *Suppose that the prime  $h \neq p$  divides the class number  $h(K)$ . Let  $D(X) = \text{Gcd}(P(X), X^{(p-1)/2} + 1) \pmod{h}$ . Then*

1.  $V(X)$  divides  $D(X)$  in  $\mathbf{F}_h[X]$ . The  $h$ -rank  $\rho$  of  $\mathbf{C}(h)$  is greater or equal to the degree  $\delta$  of  $V(X)$  and  $h^\rho \mid h(K)$ .
2. If  $h$  is coprime with  $p-1$  and with the class number  $h(E)$  of all intermediate fields  $\mathbb{Q} \subset E \subset K$ ,  $E \neq K$  then  $f \mid \rho$  where  $f$  is the order of  $h \pmod{p-1}$ .

*Proof.*

1. Reformulation of previous paragraph.
2. Immediate consequence of theorem 10.8 p. 187 in Washington [10] for the cyclic extension  $K/E$ .

□

**Remark:** we have taken the annihilation of  $\mathbf{C}$  by  $P(\sigma)$ . We have also, from Stickelberger relation,  $g(\mathbf{q})^{p(\sigma-v)}O_K = \mathbf{q}^{P(\sigma)(\sigma-v)} = \mathbf{q}^{p \times Q(\sigma)}$  where  $Q(\sigma) \in \mathbb{Z}[G]$  has been defined in lemma 2.4 p. 4, thus  $g(\mathbf{q})^{\sigma-v}O_K = \mathbf{q}^{Q(\sigma)}$ . We know that  $g(\mathbf{q})^{\sigma-v} \in K$  and so we see that  $Q(\sigma)$  annihilates also  $\mathbf{C}$ : we could take  $Q(\sigma)$  in place of  $P(\sigma)$  in previous theorem. When  $\mathbf{C}(h)$  is cyclic :

**Corollary 4.2.** *If  $\mathbf{C}(h)$  is cyclic then:*

1.  $V(X) = X - \nu$  with  $\nu \in \mathbf{F}_h^*$ .
2. In  $\mathbf{F}_h(X)$

$$(15) \quad \begin{aligned} V(X) &| X^d - 1, \quad d = \gcd(h-1, p-1), \\ V(X) &| \left\{ \sum_{i=0}^{d-1} X^i \times \frac{\sum_{k=0}^{(p-1)/d-1} v^{-(i+jd)}}{p} \right\}. \end{aligned}$$

3. If  $h$  is coprime with  $p-1$  and with the class number  $h(E)$  of all intermediate fields  $\mathbb{Q} \subset E \subset K$ ,  $E \neq K$  then  $h \equiv 1 \pmod{p-1}$ .
4. Let  $M$  be the smallest subfield of  $K$  such that  $h \mid h(M)$ . Let  $n = [M : \mathbb{Q}]$ . If  $h$  is coprime with  $p-1$  then  $h-1 \equiv 0 \pmod{n}$ .

*Proof.*

1.  $\rho = 1$  implies that  $V(\sigma) = \sigma - \nu$ .
2.  $X - \nu \mid X^{h-1} - \nu^{h-1}$  and so  $\sigma^{h-1} - \nu^{h-1}$  annihilates  $\mathbf{C}(h)$ . From  $\nu^{h-1} \equiv 1 \pmod{h}$  it follows that  $\sigma^{h-1} - 1$  annihilates  $\mathbf{C}(h)$ .  $\sigma^{p-1} - 1$  annihilates  $\mathbf{C}(h)$  and so  $\sigma^d - 1$  annihilates  $\mathbf{C}(h)$  and so  $X - \nu$  divides  $X^d - 1$  in  $\mathbf{F}_h[X]$ . Then apply theorem 4.1. Observe that  $\frac{\sum_{k=0}^{(p-1)/d-1} v^{-(i+jd)}}{p} \equiv 0 \pmod{p}$  and that we have assumed that the prime  $h \neq p$  in this section.
3. From theorem 4.1 with  $\rho = 1$  it follows that  $h \equiv 1 \pmod{p-1}$ .
4. From hypothesis we assume that  $h$  is coprime  $p-1$ . If there is no intermediate field  $E$ ,  $\mathbb{Q} \subset E \subset K$  with  $h \mid h(E)$  apply previous item. If there is an intermediate field  $E_1$ ,  $\mathbb{Q} \subset E \subset K$  with  $h \mid h(E_1)$  then  $E_1/\mathbb{Q}$  is cyclic and we can apply theorem 10.8 p. 187 in Washington [10] and we can pursue up to find the intermediate field  $E_n$  such that  $p \mid h(E_n)$  with  $E_n = M$  and no field between  $E_n$  and  $\mathbb{Q}$ . Then we apply theorem 10.8 p. 187 of Washington [10] to the cyclic extension  $M/\mathbb{Q}$ , thus  $h \equiv 1 \pmod{n}$  and  $n \mid p-1$ .

□

It is possible to enlarge previous results with another annihilation polynomial:

**Theorem 4.3.** *The polynomial  $\Pi(\sigma) = \sum_{i=0, v_{-i} \text{ even}}^{p-2} \sigma^i$  annihilates the non- $p$ -part of the class group  $\mathbf{C}$  of the  $p$ -cyclotomic field  $K$ .*

*Proof.*

1. Apply Stickelberger relation to field  $\mathbb{Q}(\zeta_{2p}) = \mathbb{Q}(\zeta_p)$ . Let  $\varpi_{2t+1} : \zeta_{2p} \rightarrow \zeta_{2p}^{2t+1}$ . The Stickelberger polynomial can be written

$$S_2 = \sum_{2t+1=1, t \neq p}^{2p-1} \varpi_{2t+1}^{-1} \times (2t+1).$$

2. Observe at first that  $\zeta_{2p} = -\zeta_p$ .

If  $t > (p-1)/2$

then  $\varpi_{2t+1} : \zeta_{2p} \rightarrow -\zeta_{2p}^{2t+1-p}$ , hence  $\varpi_{2t+1} : -\zeta_p \rightarrow -(-\zeta_p)^{2t+1-p}$ , hence  $\varpi_{2t+1} : -\zeta_p \rightarrow -(\zeta_p)^{2t+1-p}$  because  $2t+1-p$  is even, hence  $\varpi_{2t+1} : \zeta_p \rightarrow \zeta_p^{2t+1-p}$ , hence  $\varpi_{2t+1} = \varpi_{2t+1-p}$ . Then

$$\varpi_{2t+1}^{-1} \times (2t+1) = \varpi_{2t+1-p}^{-1} \times (2t+1-p) + p \times \varpi_{2t+1-p}^{-1}.$$

3. The Stickelberger polynomial is

$$S_2 = \sum_{t=0}^{(p-3)/2} \varpi_{2t+1}^{-1} \times (2t+1) + \sum_{t=(p+1)/2}^{(p-1)} \varpi_{2t+1}^{-1} \times (2t+1),$$

hence

$$S_2 = \sum_{t=0}^{(p-3)/2} \varpi_{2t+1}^{-1} \times (2t+1) + \sum_{t=(p+1)/2}^{(p-1)} \varpi_{2t+1-p}^{-1} \times (2t+1-p) + p \sum_{t=(p-1)/2}^{p-1} \varpi_{2t+1-p},$$

Then

$$S_2 = \sum_{t=1}^{p-1} \varpi_t^{-1} t + p \times \sum_{t=(p-1)/2}^{p-1} \varpi_{2t+1-p},$$

hence with  $P(\sigma)$  in lemma 2.2 p. 4

$$S_2 = P(\sigma) + p \sum_{t=(p-1)/2}^{p-1} \varpi_{2t+1-p},$$

Then with  $2t+1-p = v_{-i}$  we get

$$(16) \quad S_2 = P(\sigma) + p \sum_{i=0, v_{-i} \text{ even}}^{p-2} \sigma^i.$$

4. The polynomial  $P(\sigma)$  annihilates the class group  $\mathbf{C}$  of  $K$ . Therefore the polynomial  $p \sum_{i=0, v_{-i} \text{ even}}^{p-2} \sigma^i$  annihilates also  $\mathbf{C}$ . If  $h \neq p$  then  $\Pi(\sigma)$  annihilates  $\mathbf{C}(h)$ , which achieves the proof.

□

**Some examples:** Our results are consistent with examples taken in tables of relative class numbers of  $\mathbb{Q}(\zeta_n)$ ,  $3 \leq n \leq 1020$ , given in Washington [10], p. 412-420: verification made with a little MAPLE program.

1.  $p = 29$ ,  $p - 1 = 2^2 \cdot 7$ ,  $v = 2$ .  
 $h = 2$ ,  $2^3 \parallel h(K)$ ,  $d = 1$ ,  $V(\sigma) \mid \gcd(P(\sigma), \sigma^{14} + 1) \bmod 2 = (\sigma^3 + \sigma + 1)^2 \cdot (\sigma + 1) \cdot (\sigma^3 + \sigma^2 + 1)^2$ . Then  $\rho = 3$   $\mathbf{C}(h)$  is not cyclic and  $V(\sigma)$  is either the polynomial  $\sigma^3 + \sigma + 1$  or  $\sigma^3 + \sigma^2 + 1$ , without being able of say more.
2.  $p = 31$ ,  $p - 1 = 2 \cdot 3 \cdot 5$ ,  $v = 3$   
 $h = 3$ ,  $3^2 = h(K)$ ,  $\rho \leq 2$ ,  $d = 2$ ,  $V(\sigma) = \sigma + 1$ : 3 divides  $h(\mathbb{Q}(\sqrt{-p}))$ .
3.  $p = 131$ ,  $p - 1 = 2 \times 5 \times 13$ ,  $v = 2$ .
  - (a)  $h = 3$ ,  $3^3 \parallel h(K)$ ,  $\rho = 3$ ,  $d = 2$ ,  $V(\sigma) = \sigma^3 + 2\sigma^2 + 1$ :  $\mathbf{C}(h)$  is not cyclic.
  - (b)  $h = 5$ ,  $5^2 \parallel h(K)$ ,  $\rho \leq 2$ ,  $d = 2$ ,  $V(\sigma) = \sigma + 1$ ,  $h \mid h(\mathbb{Q}(\sqrt{-131}))$ .
  - (c)  $h = 53 \parallel h(K)$ ,  $\rho = 1$ ,  $d = 26$ ,  $V(\sigma) = \sigma + 46$ :  $\mathbf{C}(h)$  is cyclic.  $h \not\equiv 1 \pmod{p-1}$  and so there exists a field  $E$ ,  $\mathbb{Q} \subset E \subset K$  such that  $h \mid h(E)$ : it is  $E$  with  $[E : \mathbb{Q}] = 13$ .  $E/\mathbb{Q}$  is cyclic and so the theorem 4.2 item4 or the theorem 10.8 p. 187 in Washington can be applied to the extension  $E/\mathbb{Q}$  with  $f = 1$ ,  $h = 53$  and  $[E : \mathbb{Q}] = 13$  because  $53^1 \equiv 1 \pmod{13}$ .
  - (d)  $h = 1301 \parallel h(K)$ ,  $\rho = 1$ ,  $d = 130$ ,  $V(\sigma) = \sigma + 283$ :  $\mathbf{C}(h)$  is cyclic and  $h \equiv 1 \pmod{p-1}$ .
  - (e)  $h = 4673706701 \parallel h(K)$ ,  $\rho = 1$ ,  $d = 130$ ,  $V(\sigma) = \sigma + 3346914817$ :  $\mathbf{C}(h)$  is cyclic and  $h \equiv 1 \pmod{p-1}$ .
4.  $p = 137$ ,  $p - 1 = 2^3 \times 17$ ,  $v = 2$ .
  - (a)  $h = 17$ ,  $17^2 \parallel h(K)$ ,  $\rho \leq 2$ ,  $d = 8$ ,  $V(\sigma) = \sigma + 8$ .
  - (b)  $h = 47737 \parallel h(K)$ ,  $\rho = 1$ ,  $d = 136$ ,  $V(\sigma) = \sigma + 13288$ :  $\mathbf{C}(h)$  is cyclic and  $h \equiv 1 \pmod{p-1}$ .
  - (c)  $h = 46890540621121 \parallel h(K)$ ,  $\rho = 1$ ,  $d = 136$ ,  $V(\sigma) = \sigma + 14017446570735$ :  $\mathbf{C}(h)$  is cyclic and  $h \equiv 1 \pmod{p-1}$ .
5.  $p = 167$ ,  $p - 1 = 2 \times 83$ ,  $v = 5$ .
  - (a)  $h = 11 \parallel h(K)$ ,  $\rho = 1$ ,  $d = 2$ ,  $V(\sigma) = \sigma + 1$ :  $\mathbf{C}(h)$  is cyclic.  $h \mid h(\mathbb{Q}(\sqrt{-167}))$ .
  - (b)  $h = 499 \parallel h(K)$ ,  $\rho = 1$ ,  $d = 166$ ,  $V(\sigma) = \sigma + 491$ :  $\mathbf{C}(h)$  is cyclic and  $h \equiv 1 \pmod{p-1}$ .

- (c)  $h = 5123189985484229035947419 \parallel h(K)$ ,  $\rho = 1$ ,  $d = 166$ ,  
 $V(\sigma) = \sigma + 698130937752344432562779$ :  $\mathbf{C}(h)$  is cyclic and  $h \equiv 1 \pmod{p-1}$ .

**Lemma 4.4.** *Let  $E$  be a subfield of  $K$  with  $[K : \mathbb{Q}] = d$ . Let  $h$  be an odd prime number dividing  $h(L)$ . Then in  $\mathbf{F}_h[X]$*

$$(17) \quad \begin{aligned} V(X) &| \sum_{i=0}^{d-1} X^i, \\ V(X) &| \left\{ \sum_{i=0}^{d-1} X^i \times \frac{\sum_{k=0}^{(p-1)/d-1} v^{-(i+jd)}}{p} \right\}. \end{aligned}$$

*Proof.*  $\sigma^d - 1$  annihilates  $\mathbf{C}(h)$ . The Stickelberger polynomial

$$P(X) = \sum_{i=0}^{p-2} X^i v_{-i} = p \times \sum_{i=0}^{d-1} X^i \times \frac{\sum_{k=0}^{(p-1)/d-1} v^{-(i+jd)}}{p},$$

and from  $p \neq h$  it follows that

$$V(X) | \left\{ \sum_{i=0}^{d-1} X^i \times \frac{\sum_{k=0}^{(p-1)/d-1} v^{-(i+jd)}}{p} \right\}$$

in  $\mathbf{F}_p[G]$ . □

**Remark:** Compare lemma 4.4 for  $h \neq p$  with lemma 3.4 p. 8 proved when  $h = p$ .

### Some Remarks

1. In our propositions we apply the **simultaneous** annihilation of subgroup  $\mathbf{C}(h)$  of the class group by  $\sigma^{(p-1)/2} + 1$  and by the Stickelberger polynomial  $\sum_{k=0}^{p-2} \sigma^k v_{-k}$ , which improves the result obtained with the only norm polynomial (as in proof of theorem 10.8 p. 187 in Washington[10]) or with the only Stickelberger polynomial.
2. Let  $\delta$  be an integer  $1 \leq \delta \leq p-2$ . There exists an integer  $s$ ,  $0 \leq s \leq p-2$  such that  $\delta \equiv v^s \pmod{p}$ .  $s$  is called the index of  $\delta$  relative to  $v$  and denoted  $s = \text{ind}_v(\delta)$ . Let us define the polynomial

$$(18) \quad \begin{aligned} Q(X) &= \sum_{i \in I_\delta} X^i, \\ I_\delta &= \{i \mid 0 \leq i \leq p-2, v^{(p-1)/2-i} + v^{(p-1)/2-i+\text{ind}_v(\delta)} > p\}. \end{aligned}$$

From a result of Kummer on Jacobi cyclotomic functions, the polynomial  $Q(\sigma)$  annihilates the complete class group  $\mathbf{C}$  of  $K_p$  (see for instance Ribenboim [8] relation (2.5) p. 119). It follows that  $V(X) \mid Q(X)$  in  $\mathbf{F}_h[X]$  gives another criterium for the study of the structure of the group  $\mathbf{C}$ .

3. Observe that the theorem 4.3 is not trivial even when  $\mathbf{C}(h) \subset \mathbf{C}^+$  class group of  $K^+$ , by opposite to Stickelberger polynomial annihilating trivially  $\mathbf{C}(h) \subset \mathbf{C}^+$ .
4. The results could be generalized to the cyclotomic fields  $K_n = \mathbb{Q}(\zeta_n)$  where  $n$  is not prime.

## 4.2 The case of complex quadratic fields contained in $K$

In this paragraph we formulate directly previous result when  $h$  divides the class number of the complex quadratic field  $\mathbb{Q}(\sqrt{-p}) \subset K$ ,  $p \equiv 3 \pmod{4}$ ,  $p \neq 3$ .

**Theorem 4.5.** *Suppose that  $p \equiv 3 \pmod{4}$ ,  $p \neq 3$ . If  $h$  is an odd prime with  $h \mid h(\mathbb{Q}(\sqrt{-p}))$  then*

$$(19) \quad \sum_{i=0}^{p-2} (-1)^i v_{-i} \equiv 0 \pmod{h}.$$

*Proof.* Let  $\mathbf{Q}$  be the prime of  $\mathbb{Q}(\sqrt{-p})$  lying above  $\mathbf{q}$ . The ideals  $\mathbf{Q} \neq \sigma(\mathbf{Q})$  and so  $\mathbf{Q}^{\sigma+1}$  is principal because  $\mathbf{Q}^{\sigma^2} = \mathbf{Q}$ . Therefore  $\mathbf{Q}^{\sum_{i=0}^{p-2} (-1)^i v_{-i}}$  is principal and  $\sum_{i=0}^{p-2} (-1)^i v_{-i} \equiv 0 \pmod{h}$ .  $\square$

### Remarks:

1. The theorem 4.7 can also be obtained from Hilbert Theorem 145 see Hilbert [3] p. 119. See also Mollin, [6] theorem 5.119 p. 318.
2. From lemmas 2.4 p. 4 we could prove similarly:

Suppose that  $p \equiv 3 \pmod{4}$ ,  $p \neq 3$ . If  $h$  is an odd prime with  $h \mid h(\mathbb{Q}(\sqrt{-p}))$  then

$$(20) \quad 2 \times \sum_{i=0}^{(p-3)/2} (-1)^i v_{-i} - p \equiv 0 \pmod{h}.$$

3. Numerical evidences easily computable show more: If  $p \neq 3$  is prime with  $p \equiv 3 \pmod{4}$  then the class number  $h(\mathbb{Q}(\sqrt{-p}))$  verifies

$$(21) \quad h(\mathbb{Q}(\sqrt{-p})) = -\frac{\sum_{i=0}^{p-2} (-1)^i v_{-i}}{p}.$$

This result has been proved by Dirichlet by analytical number theory, see Mollin remark 5.124 p. 321. It is easy to verify this formula, for instance in tables of class numbers of complex quadratic fields in :

(a) H. Cohen [2] p. 502- 505, all the table for  $p \leq 503$ .

(b) in Wolfram table of quadratic class numbers [11] for large  $p$ .

4. When  $p \equiv 1 \pmod{4}$  this method cannot be applied to the quadratic field  $\mathbb{Q}(\sqrt{p}) \subset K$  because  $\sum_{i=0}^{p-2} (-1)^i v_{-i}$  is trivially null.

**Theorem 4.6.** *Suppose that  $p \equiv 3 \pmod{4}$ ,  $p \neq 3$ . If  $h$  is an odd prime with  $h \mid h(\mathbb{Q}(\sqrt{-p}))$  then*

$$(22) \quad \begin{aligned} \sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i &\neq 0, \\ \sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i &\equiv 0 \pmod{h}. \end{aligned}$$

*Proof.* We apply previous theorem 4.3 p. 13 observing that in that case  $\sigma^2(\mathbf{Q}) = \mathbf{Q}$  when  $\mathbf{Q}$  is a non-principal ideal of the quadratic field  $\mathbb{Q}(\sqrt{-p})$ , hence  $\sigma+1$  annihilates  $\mathbf{C}(h)$  and  $\sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i \equiv 0 \pmod{h}$ . This sum has  $\frac{p-1}{2}$  elements, thus of odd cardinal and cannot be null.  $\square$

**Theorem 4.7.** *Suppose that  $p \equiv 3 \pmod{4}$ ,  $p \neq 3$ . Let  $\delta$  be an integer  $1 \leq \delta \leq p-2$ . Let  $I_\delta$  be the set*

$$(23) \quad I_\delta = \{i \mid 0 \leq i \leq p-2, v_{(p-1)/2-i} + v_{(p-1)/2-i+ind_v(\delta)} > p\},$$

where, as seen above,  $ind_v(\delta)$  is the notation index of  $\delta$  relative to  $v$ . If  $h$  is an odd prime with  $h \mid h(\mathbb{Q}(\sqrt{-p}))$  then

$$(24) \quad \begin{aligned} \sum_{i \in I_\delta} (-1)^i &\neq 0, \\ \sum_{i \in I_\delta} (-1)^i &\equiv 0 \pmod{h}. \end{aligned}$$

*Proof.*  $I_\delta$  has an odd cardinal. Then see relation (23).  $\square$

**Remark:**

1. Observe that results of theorems 4.6 and 4.7 are consistent with existing tables of quadratic fields, for instance Arno, Robinson, Wheeler [1]. Numerical verifications seem to show more :

$$(25) \quad \sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i \equiv 0 \pmod{h(\mathbb{Q}(\sqrt{-p}))}.$$

2. Observe that if  $p \equiv 1 \pmod{4}$  then  $\sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i = 0$ .

### 4.3 The case of biquadratic fields contained in $K$

The following example is a generalization for the biquadratic fields  $L$  which are included in  $p$ -cyclotomic field  $K$  with  $p \equiv 1 \pmod{4}$ .

**Theorem 4.8.** *Let  $p$  be a prime with  $2^2 \parallel p - 1$ . Let*

$$(26) \quad S = \left( \sum_{i=0}^{(p-3)/2} (-1)^i v_{2i} \right)^2 + \left( \sum_{i=0}^{(p-3)/2} (-1)^i v_{2i+1} \right)^2.$$

*Let  $L$  be the field with  $\mathbb{Q}(\sqrt{p}) \subset L \subset K$ ,  $[L : \mathbb{Q}(\sqrt{p})] = 2$ . Let  $h$  be an odd prime number with  $h \mid h(L)$  and  $h \nmid h(\mathbb{Q}(\sqrt{p}))$ . Then  $S \neq 0$  and  $S \equiv 0 \pmod{h}$ .*

*Proof.*  $V(\sigma) \mid \sigma^4 - 1$ .  $h \nmid h(\mathbb{Q}(\sqrt{p}))$  and so  $V(\sigma) \mid \sigma^2 + 1$ .  $P(\sigma) = \sum_{i=0}^{(p-3)/2} \sigma^{2i} v^{-2i} + \sigma \times \sum_{i=0}^{(p-3)/2} \sigma^{2i} v^{2i+1}$ . Relation (26) follows.  $\square$

**Remarks:**

1.  $S$  does not depend of the primitive root  $v \pmod{p}$  chosen.
2. Numerical computations seem to show more :  $P(\sigma) \equiv 0 \pmod{p^2}$  and so

$$(27) \quad \frac{\left( \sum_{i=0}^{(p-3)/2} (-1)^i v_{2i} \right)^2 + \left( \sum_{i=0}^{(p-3)/2} (-1)^i v_{2i+1} \right)^2}{p^2} \equiv 0 \pmod{h}.$$

3. This result is a generalization for biquadratic fields of theorem 145 of Hilbert for quadratic fields.

## References

- [1] S. Arno, M.L. Robinson, F.S. Wheeler *Imaginary quadratic fields with small odd class number*, Acta Arith. 83, 1998, 295-300.
- [2] H. Cohen, *A course in computational number theory*, Springer-Verlag, 1993.
- [3] D. Hilbert, *The Theory of Algebraic Numbers*, Springer, 1998
- [4] K. Ireland, M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, 1982.
- [5] H. Koch, *Algebraic Number Theory*, Springer, 1997.
- [6] R.A. Mollin, *Algebraic Number Theory*, Chapman and Hall/CRC, 1999.
- [7] W. Narkiewicz, *Elementary and Analytic Theory of Numbers*, Springer-verlag, 1990.
- [8] P. Ribenboim, *13 Lectures on Fermat's Last Theorem*, Springer-Verlag, 1979.
- [9] P. Ribenboim, *Classical Theory of Algebraic Numbers*, Springer, 2001.
- [10] L.C. Washington, *Introduction to cyclotomic fields, second edition*, Springer, 1997.
- [11] *Class number of quadratic fields*,  
<http://www.mathworld.wolfram.com/ClassNumber.html>

Roland Quême

13 avenue du château d'eau

31490 Brax

France

mailto:roland.queme@wanadoo.fr

home page: <http://roland.queme.free.fr/>

\*\*\*\*\*

V10 - MSC Classification : 11R18; 11R29

\*\*\*\*\*