

ON PLOUFFE'S RAMANUJAN IDENTITIES

LINAS VEPSTAS

ABSTRACT. Recently, Simon Plouffe has discovered a number of identities for the Riemann zeta function at odd integer values. These identities are obtained numerically and are inspired by a prototypical series for Apéry's constant given by Ramanujan:

$$(3) = \frac{7^3}{180} \sum_{n=1}^{\infty} \frac{1}{n^3 (e^{2^n} - 1)}$$

This note provides an analytic derivation of these types of identities, providing insight into their structure, and giving a general layout in which to discuss them. The resemblance of these relations to the general structure of modular forms is briefly sketched.

1. INTRODUCTION

Inspired by an identity for (3) given in Ramanujan's notebooks [2, chapter 14, formulas 25.1 and 25.3],

$$(3) = \frac{7^3}{180} \sum_{n=1}^{\infty} \frac{1}{n^3 (e^{2^n} - 1)}$$

Plouffe describes a set of similar identities [4],[5] that were discovered numerically using arbitrary-precision software. For example, Plouffe gives an identity for (7):

$$(7) = \frac{19^7}{56700} \sum_{n=1}^{\infty} \frac{1}{n^7 (e^{2^n} - 1)}$$

This text provides an analytically derived formula for expressions of this type. The resulting general formula, valid for integer $m \geq 1$, is

$$(4m-1) = \sum_{n=1}^{\infty} \frac{1}{n^{4m-1} (e^{2^n} - 1)}$$

$$\frac{1}{2} (2)^{4m-1} \sum_{j=0}^{2m-1} (-1)^j \frac{B_{2j} B_{4m-2j}}{(2j)! (4m-2j)!}$$

where B_k is the k 'th Bernoulli number.

Similar sums have been explored before; for example, Apostol gives [1, see exercise 15 at end of chapter 1.]

$$\sum_{n=1, n \text{ odd}}^{\infty} \frac{n^{4k+1}}{1+e^n} = \frac{2^{4k+1}}{8k+4} B_{4k+2}$$

Similar sums involving double Bernoulli numbers also appear in the analysis of the Dedekind eta function; see for example, Sho Iseki's transformation formula, as described by Apostol [1, see theorem 3.5].

Date: 14 May 2006.

1991 Mathematics Subject Classification. 11M06.

Plouffe also notes similar relations for the other odd integers; for example,

$$(5) = \frac{5}{294} \sum_{n=1}^{\infty} \frac{72 X^{2n}}{n^5 (e^{2n} - 1)} - \frac{2}{35} \sum_{n=1}^{\infty} \frac{X^{2n}}{n^5 (e^{2n} + 1)}$$

The general form for this type of expression may be shown to be

$$\begin{aligned} 1 + \frac{X}{(4)^m} \sum_{n=1}^{\infty} \frac{2^{4m+1}}{(4m+1)} &= 2 \sum_{n=1}^{\infty} \frac{X^{2n}}{n^{4m+1} (e^{2n} - 1)} \\ &+ 2 \sum_{n=1}^{\infty} \frac{2^{4m+1}}{(4)^m} \frac{X^{2n}}{n^{4m+1} (e^{2n} + 1)} \\ &+ (2)^{4m+1} \sum_{j=0}^{\infty} \frac{X^{2n}}{(4)^{m+j}} \frac{B_{4m-4j+2}}{(4m-4j+2)!} \frac{B_{4j}}{(4j)!} \\ &+ \frac{1}{2} (2)^{4m+1} \sum_{j=0}^{\infty} \frac{2^{2j+1}}{(4)^j} \frac{B_{4m-2j+2}}{(4m-2j+2)!} \frac{B_{2j}}{(2j)!} \end{aligned}$$

A sum of the above expressions for $(4m - 1)$ and $(4m + 1)$ is noted by Borwein, *et al*[3, formula 60], who attribute it to Ramanujan[2, Ch. 14, entry 21(i), page 359].

The methods described in this text also allow for a large generalization of these types of sums. Defining

$$P_k (X) = \sum_{n=1}^{\infty} \frac{X^{kn}}{n^k (e^{2in} - 1)}$$

these generalizations follow from a modular equation relating $P_k (X)$ to $P_k (X^{-1})$ for odd integers k . In particular, interesting relationships result from considering $X \in \mathbb{Q}[\sqrt{-1}]$, the field of Gaussian rationals, as well as Diophantine roots of quadratics.

The rest of this paper is roughly laid out as follows: The second section gives a relationship between the sums and the polylogarithm, and thence to an integral on the complex plane. The third section examines the related contour integral, which is easily integrated via Cauchy's residue theorem to give a finite sum involving the Bernoulli numbers. The fourth section relates the contour integral to the polylogarithm integral, thus resulting in a functional equation for $P_k (X)$. The fifth section applies the functional equation, providing a variety of lemmas, many of which explain Plouffe's discoveries. The sixth section give a pair of relationships on the Bernoulli numbers that arise naturally in this context. The seventh section explores the modular nature of the relations on $P_k (X)$, followed by a conclusion. An appendix gives a derivation of an integral representation of the polylogarithm, that is central to the analysis.

2. THE POLYLOGARITHM

The recurring theme in Plouffe's identities is the sum

$$S_s (x) = \sum_{n=1}^{\infty} \frac{x^n}{n^s (e^{xn} - 1)}$$

with s usually an odd positive integer and $x = \frac{1}{m}$ or $x = 2$ or possibly other interesting values, such as $x = \frac{1}{m}$ for some integer m . This sum may be converted into a sum over polylogarithms, and subsequently into an integral. The integral may, after some difficulties, be converted into a contour integral, whereupon it may be evaluated by Cauchy's residue

theorem. The result is a finite sum whose general structure resembles those of Plouffe's and Ramanujan's identities. This section develops the first part of this analysis.

To find the polylogarithm, one expands

$$\begin{aligned}
 S_s(x) &= \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{x^{n(m+1)}}{n^s} \\
 &= \sum_{m=1}^{\infty} \text{Li}_s(x^m) e^{-xm}
 \end{aligned}$$

which is generally valid for $\langle x \rangle > 0$. The above is easily obtained by applying the expansion

$$\frac{1}{1-z} = \sum_{m=0}^{\infty} z^m$$

and the definition of the polylogarithm:

$$\text{Li}_s(z) = \sum_{n=1}^{\infty} \frac{z^n}{n^s}$$

The polylogarithm may be expressed in terms of an integral as

$$\text{Li}_s(e^{-u}) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma(z) (z+s)^{-s} u^{-z} dz$$

(See Appendix for derivation). Here, $\Gamma(z)$ is the Gamma function. The line of integration is taken to be to the right of all of the poles in the integrand, namely, $c > 1$. Using this in the summation, one obtains

$$\begin{aligned}
 S_s(x) &= \sum_{m=1}^{\infty} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma(z) (z+s)^{-s} (xm)^{-z} dz \\
 &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \Gamma(z) \frac{(z)}{x^z} (z+s)^{-s} (z) dz
 \end{aligned}$$

The exchange of the order of summation and integration is justified precisely when one has $c > 1$. The last integral has poles at $z = 1$ and $z + s = 1$ coming from the zeta functions and poles at all of the non-positive integers coming from the Gamma function.

If the integral can somehow be converted into a closed contour on the left, then it may be evaluated in a straight-forward way by means of Cauchy's residue theorem. Performing this closure is in fact harder than one might hope, as there are non-zero contributions to the contour from its closure. The next section evaluates the Cauchy integral, assuming that the contour can be closed. The section after that computes the contributions from closing the contour integral.

3. THE CONTOUR INTEGRAL

Define the contour integral as

$$I_s(x) = \frac{1}{2\pi i} \int \frac{\Gamma(z)}{x^z} (z+s)^{-s} (z) dz$$

where the contour encircles the poles at $z = 1$, $z + s = 1$ and $z = 0; -1; -2; \dots$ in the usual, right-handed fashion. Then, one uses Cauchy's theorem, which states that

$$f(a) = \frac{1}{2\pi i} \int \frac{f(z)}{z-a} dz$$

for simple poles, and that

$$f^0(a) = \frac{1}{2i} \int_{z=1-k}^1 \frac{f(z)}{(z-a)^2} dz$$

for double poles. For the pole at $z = 1$, one obtains the residue

$$\text{Res}(z = 1) = \frac{(s+1)}{x}$$

For the poles at $z = -n$, one obtains the residue

$$\text{Res}(z = -n) = \frac{(-x)^n}{n!} (s-n) \zeta(n)$$

and so one has

$$I_s(x) = \text{Res}(z = 1-s) + \frac{(s+1)}{x} + \sum_{n=0}^{\infty} \frac{(-x)^n}{n!} (s-n) \zeta(n)$$

For s not an integer, one has

$$\text{Res}(z = 1-s) = \frac{(1-s)}{x^{1-s}} \zeta(1-s)$$

However, the interesting case is for $s = k$ a positive integer. In this case, the pole overlays another pole from the Gamma, and one has a double pole. This is just a little trickier to evaluate:

$$\begin{aligned} \frac{1}{2i} \int_{z=1-k}^1 \frac{(z)}{x^z} (z+s) \zeta(z) dz &= \frac{1}{2i} \int_{z=1-k}^1 \frac{f(z)}{(z+k-1)^2} dz \\ &= \frac{d}{dz} \zeta(z+k-1)^2 (z) \zeta(z+k) \frac{(z)}{x^z} \Big|_{z=1-k} \end{aligned}$$

To perform the derivative, one will need to use the identities

$$\frac{d}{ds} \zeta(s-1) \zeta(s) \Big|_{s=1} =$$

where $\gamma = 0.577 \dots$ is the Euler-Mascheroni constant, and

$$\frac{d}{dz} \zeta(z+n) \zeta(z) \Big|_{z=-n} = (-1)^n \frac{(n+1)}{(n+1)} = (-1)^n \frac{H_n}{n!}$$

where $\zeta(z)$ is the digamma function, and H_n is the n 'th harmonic number. Putting these together, one obtains

$$\frac{1}{2i} \int_{z=1-k}^1 \frac{(z)}{x^z} (z+s) \zeta(z) dz = \frac{(-x)^{k-1}}{(k-1)!} [\zeta'(1-k) + (H_{k-1} \ln 2) \zeta(1-k)]$$

Adding this to the other contributions, one gets

$$\begin{aligned} I_k(x) &= \frac{(k+1)}{x} + \sum_{\substack{n=0 \\ n \neq k-1}}^{\infty} \frac{(-x)^n}{n!} (k-n) \zeta(n) \\ &\quad + \frac{(-x)^{k-1}}{(k-1)!} [\zeta'(1-k) + (H_{k-1} \ln 2) \zeta(1-k)] \end{aligned}$$

When k is an odd integer, the above simplifies in two ways. First, $\zeta(1-k)$ vanishes, because the zeta function vanishes at all negative even integers. Similarly, the infinite sum

becomes finite: when k is an odd integer, one has either $\binom{k-n}{n} = 0$ or $\binom{n}{n} = 0$ for all $n > k$. Thus, for k an odd integer, one has

$$I_k(x) = \frac{(k+1)}{x} + \sum_{\substack{n=0 \\ n \notin k-1}}^{x^k} \frac{(x)^n}{n!} \binom{k-n}{n} \binom{n}{n} + \frac{(x)^{k-1}}{(k-1)!} {}_0(1-k)$$

For the remainder of the paper, it is assumed that, in this context, $k > 1$ is an odd integer, unless explicitly stated otherwise. The above evaluation of the contour integral is the main result of this section. To see that this is a key result, one may substitute $k = 3$ and $x = 2$ to obtain

$$I_3(2) = (3) + \frac{7^3}{180}$$

which should be recognizable as a portion of Ramanujan's identity. For $k = 7$, one has

$$I_7(2) = (7) + \frac{19^7}{56700}$$

which resembles one of the results given by Plouffe. To complete the connection, one must relate the contour integral $I_k(x)$ to the sum $S_k(x)$. This is done in the next section.

First, however, to drive the point home, one must observe that most of the terms in the above expression are rational multiples of powers of x . This follows from the zeta function being related to the Bernoulli numbers B_n at even integers:

$$\zeta(2n) = (-1)^{n+1} \frac{(2\pi)^{2n} B_{2n}}{2(2n)!}$$

for integer $n \geq 0$. At the negative values, one has

$$\zeta(-n) = \frac{B_{n+1}}{n+1}$$

while the derivative is

$$\zeta'(-2n) = (-1)^n \frac{(2n)!}{2(2\pi)^{2n}} (2n+1)$$

for integer $n > 1$. Using these in the above expression for $I_k(x)$, and rearranging terms a bit, one gets

$$2I_k(x) = (k-1) \frac{x^{k-1}}{2-i} + \frac{1}{x(k+1)!} \sum_{j=0}^{(k-1)/2} \frac{x^{k+1-2j}}{2^j} (2-i)^{k+1-2j} B_{2j} B_{k+1-2j}$$

The above introduces the binomial coefficient

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Although the imaginary number $i = \sqrt{-1}$ appears in the above, it is always squared, and thus is just a sign-keeping device. Every term in the sum is purely real.

When the contour integral is written in this form, it may now be seen that for x being any rational multiple of π , that is, $x = p/q$ for any integers p, q that the coefficient of $\binom{k}{n}$ is a rational number, and that the second term is another rational times x^k .

A further curiosity in this regard is noted by Plouffe: if one takes $x = \frac{p}{p-q}$ for integers p and q , one also gets simple expressions: because k is odd, the coefficient of x^k is still a rational, and the coefficient of x^{k+1} is $\frac{p}{p-q}$ times some rational. For rational x , one still has that the sum is a rational polynomial in x^2 , and for $x = \frac{p}{p-q}$ one still has that the sum is a rational polynomial in x . The ocean-full of rationals here suggest that some sort of p -adic analysis might be interesting. The appearance of the square root suggests that there is a relation to complex multiplication, or that one may have interesting results on the field of Gaussian integers.

4. EVALUATING THE CONTOUR INTEGRAL

The goal of this section is to relate the sum $S_k(x)$ to the contour integral $I_k(x)$.

Theorem 4.1. *For odd integers k , one has that*

$$S_k(x) = I_k(x) + (-1)^{(k-1)/2} \frac{x^{k-1}}{2} S_k\left(\frac{4}{x}\right)$$

For the remainder of this text, this will be referred to as the "functional equation for S_k ".

Proof. To prove this result, consider evaluating the contour integral $I_s(x)$ for a tall rectangular contour surrounding the poles at $z = 1; 0; -1; \dots; -s$. Thus, write $I_s(x) = A + B + C + D$ with A being the integral from $c - ih$ to $c + ih$ for a constant $c > 1$ and the height h large, eventually taking the limit $h \rightarrow \infty$. That is, A forms the right hand side of the rectangular contour. In the limit of $h \rightarrow \infty$, one has by definition

$$A = S_s(x)$$

Let B and C be the top and bottom of the contour, so that for B , the integral runs from $c + ih$ to $ih - s$ leftwards. For C , the integral runs rightwards from $ih - s$ to $c - ih$; here we take $c > 0$. The integral D on the left hand side of the rectangle closes the contour, running downwards, from $ih - s$ to $c - ih$.

The integrals B and C will vanish in the limit of the height $h \rightarrow \infty$. This can be easily seen after a simple change of variable:

$$\begin{aligned} B &= \frac{1}{2i} \int_{c+ih}^{ih-s} \frac{z^s}{x^z} (s+z)(z) dz \\ &= \frac{1}{2i} \int_c^{ih-s} \frac{(u+ih)^s}{x^{u+ih}} (s+u+ih)(u+ih) du \end{aligned}$$

Much of the integrand is $O(1)$ in h , or polynomially thereabouts. The integrand is dominated by the Gamma function, which, from Stirling's approximation, may be seen to be

$$(u+ih) \Gamma(u+ih) \sim e^{-h} h^{u+ih-1/2}$$

The complex conjugate argument applies to C , and thus B and C vanish in the limit of $h \rightarrow \infty$.

To evaluate D , begin by writing

$$\begin{aligned} D &= \frac{1}{2i} \int_{ih-s}^{c-ih} \frac{z^s}{x^z} (s+z)(z) dz \\ &= \frac{1}{2i} x^{s+ih} \int_{ih-s}^{c-ih} x^{-u} (u-s)(u) (u-s) du \end{aligned}$$

after a change of variable $z = u + s$. One then applies the functional equations for Gamma:

$$\Gamma(z) \Gamma(1-z) = \frac{1}{\sin z}$$

and for zeta:

$$\zeta(s) = 2^{s-1} \sin \frac{s}{2} \Gamma(1-s) \Gamma(s)$$

to obtain

$$D = \frac{1}{2^{2i}} \frac{x}{2} \int_0^{2\pi} \frac{1}{2} \frac{R_{ih}}{ih} \frac{x}{4} \int_0^{2\pi} \frac{\sin \frac{u+s}{2} \sin \frac{u+s+}{2}}{\sin(u+s)} (1+u) (1+u) (1+s+u) du$$

Another change of variable, this time as $w = 1 + u + s$ take the integral to a slightly more recognizable form:

$$D = \frac{1}{i} \frac{x}{2} \int_0^{2\pi} \frac{1}{1+ih} \frac{x}{4} \int_0^{2\pi} \frac{\cos \frac{w}{2} \cos \frac{w+s}{2}}{\sin(w+s)} (w) (w) (w+s) dw$$

Next, by taking $s = k$ to be an odd integer, the trigonometric piece simplifies and loses its w dependence:

$$\frac{\cos \frac{w}{2} \cos \frac{w+k}{2}}{\sin(w+k)} = \frac{1}{2} (1)^{(k+1)/2}$$

and in the limit $h \rightarrow 1$, one finally reaches the claimed result:

$$D = (1)^{(k+1)/2} \frac{x}{2} \int_0^{2\pi} S_k \frac{4}{x}$$

that is,

$$I_k(x) = A + D = S_k(x) (1)^{(k+1)/2} \frac{x}{2} \int_0^{2\pi} S_k \frac{4}{x}$$

for odd integer k .

The next section will review the application of this and the preceding sections to specific, simple values of x , thus regaining many of Plouffe's sums.

5. LEMMAS AND APPLICATIONS

The first corollary demonstrates Plouffe's simplest sums for $(4m-1)$.

Corollary 5.1. For m integer, one has

$$I_{4m-1}(2) = 2S_{4m-1}(2):$$

Proof. Substitute $x = 2$ in the functional equation.

This corollary provides the first concrete result of this exposition, namely that

$$\sum_{n=1}^{\infty} \frac{1}{n^{4m-1} (e^{2n}-1)} = (4m-1) + \frac{(2)^{4m-1}}{(4m)!} \sum_{j=0}^{4m} \binom{4m}{2j} (1)^j B_{2j} B_{4m-2j}$$

which completely resolves one set of relationships given by Plouffe. The functional equation opens additional possibilities. By substituting $x = 2^p/q$, one obtains, for $k = 4m-1$, that

$$q^{k-1} \sum_{n=1}^{\infty} \frac{1}{n^k e^{2pn/q} - 1} + p^{k-1} \sum_{n=1}^{\infty} \frac{1}{n^k e^{2qn/p} - 1} = q^{k-1} I_k \frac{2^p}{q}$$

Thus, for example, by choosing $p = 2, q = 1$ and $k = 3$, one obtains

$$(3) = \frac{29^3}{900} - \frac{2}{5} \sum_{n=1}^{\infty} \frac{x^n}{n^3} \frac{1}{e^n - 1} + \frac{4}{e^{4n} - 1}$$

and similarly, for $k = 7$,

$$(7) = \frac{409^7}{94500} - \frac{2}{5} \sum_{n=1}^{\infty} \frac{x^n}{n^7} \frac{1}{e^n - 1} + \frac{4}{e^{4n} - 1}$$

and one may proceed in a similar manner.

The next corollary shows that something more is needed for (5), (9), and so on, since the most direct approach does not give any information for such sums.

Corollary 5.2. For k an odd integer, one has

$$I_k(x) = (-1)^{(k+1)/2} \frac{4^{k-1}}{x} I_k \left(\frac{4}{x} \right)$$

Proof. This may be proved by applying the functional equation twice in a row. That is, it may be proved by substituting $x \rightarrow 4/x$ in the functional equation and then employing the result.

Corollary 5.3. For m a positive integer, one has

$$I_{4m+1}(2) = 0:$$

Proof. Substitute $x = 2$ in the preceding corollary.

Following directly from the above is another corollary:

Corollary 5.4. For m a positive integer, one has

$$0 = \sum_{j=0}^{2m+1} (-1)^j \frac{B_{4m-2j+2}}{(4m-2j+2)!} \frac{B_{2j}}{(2j)!}$$

Proof. Write out the value of $I_{4m+1}(2)$ in detail.

From the above, it should be clear that the functional equation for S_k does not provide any statements about S_k when $k = 4m + 1$. To obtain results on sums involving $k = 4m + 1$, one must introduce

$$T_s(x) = \sum_{n=1}^{\infty} \frac{x^n}{n^s (e^{xn} + 1)}$$

Theorem 5.5. One has

$$T_s(x) = S_s(x) - 2S_s(2x)$$

Proof. This may be proved by re-writing in terms of the polylogarithm, along the lines of the earlier development:

$$T_s(x) = \sum_{m=1}^{\infty} (-1)^m L_s^m e^{-xm}$$

The even and odd terms are regrouped, as

$$T_s(x) = \sum_{m=1}^{\infty} L_s^m e^{-xm} - 2 \sum_{m=1}^{\infty} L_s^m e^{-2xm}$$

which is seen to be a sum of S_s 's.

The results for (5), etc. follow from a critical observation: that

$$S_s(x + 2i) = S_s(x)$$

is a periodic function. This periodicity is employed directly in the next theorem.

Theorem 5.6. For positive integer m , one has

$$S_{4m+1}(2) = I_{4m+1}(2(1+i)) + (1)^m \frac{T_{4m+1}(2)}{4^m} + 2^{-m} S_{4m+1}(2) - 4^m I_{4m+1}(2)$$

Proof. Using periodicity, one writes, for $k = 4m + 1$,

$$S_k(2 + 2i) = S_k(2) = I_k(2(1+i)) + 2^{(k-1)/2} e^{3i(k-1)/4} S_k(1-i)$$

The series $S_k(1-i)$ doesn't have an imaginary part; rather, it is an alternating series, which may be expanded and written as

$$S_k(1-i) = T_k(2) + \frac{T_k(2) + S_k(2)}{2^k}$$

The $T_k(2)$ term may be eliminated by writing

$$T_k(2) = S_k(2) - 2S_k(2)$$

and the $S_k(2)$ term may be eliminated by

$$\begin{aligned} S_k(2) &= I_k(2) + 2^{1-k} S_k(4) \\ &= I_k(2) + 2^{-k} [S_k(2) - T_k(2)] \end{aligned}$$

Performing the various substitutions suggested above proves the theorem.

As an example of the application of the above theorem, take $m = 1$, that is, $k = 5$. One easily finds that

$$I_5(2) = \frac{15}{32}(5) + \frac{5}{9 \cdot 64}$$

and that

$$I_5(2(1+i)) = \frac{5}{2}(5) + \frac{5}{9 \cdot 15}$$

Combining these, one gets

$$(5) = \frac{5}{294} - \frac{2}{35} [I_5(2) + 36S_5(2)]$$

which is given by Plouffe. The theorem may be used to generate similar expressions for all $(4m + 1)$.

Curiously, the theorem yields results for $m = 0$ as well. In this case, one finds

$$S_1(2) + T_1(2) = \frac{3}{6} \log 2$$

Many identities for $\zeta(3)$ are possible by taking two different expressions for a given zeta, and subtracting them, leaving behind a rational combination of the sums and $\log 2$. Thus, for example, the following theorem for Apéry's constant:

Theorem 5.7. A series expression for $\zeta(3)$ is given by

$$\zeta(3) = 720 \sum_{n=1}^{\infty} \frac{1}{n^3} - 900 \sum_{n=1}^{\infty} \frac{1}{(2n)^3} + 180 \sum_{n=1}^{\infty} \frac{1}{(4n)^3}$$

Proof. This follows by taking the general expression for $k = 4m - 1$:

$$\begin{aligned} 16^m S_{4m-1}(\cdot) + 4S_{4m-1}(4) &= 16^m I_{4m-1}(\cdot) \\ &= \frac{1}{2} (4m-1) [16^m + 4] \\ &= (2)^{4m-1} \sum_{j=0}^{2m} (4)^j \frac{B_{4m-2j} B_{2j}}{(4m-2j)! (2j)!} \end{aligned}$$

solving for $(4m-1)$ and then using

$$2S_{4m-1}(2) = I_{4m-1}(2) = (4m-1) + (2)^{4m-1} \sum_{j=0}^{2m} (1)^j \frac{B_{4m-2j} B_{2j}}{(4m-2j)! (2j)!}$$

to eliminate the appearance of the zeta. The resulting expression may then be solved for $(4m-1)$.

Relationships involving square roots also arise naturally.

Theorem 5.8. *One has*

$$(3) = \frac{5^3}{72} - \frac{1}{2} S_3 - 2^{\frac{p-3}{3}} - \frac{3}{2} S_3 - \frac{2^{\frac{p-3}{3}}}{3}$$

Proof. This follows from the general expression given earlier, that

$$S_k \frac{2^{\frac{p}{q}}}{q} + \frac{p}{q} S_k \frac{2^{\frac{q}{p}}}{p} = I_k \frac{2^{\frac{p}{q}}}{q}$$

Here, making the substitution $q = \frac{p}{\bar{p}}$ one obtains

$$\begin{aligned} S_k (2^{\frac{p}{\bar{p}}}) + p^{(k-1)=2} S_k \frac{2^{\frac{p}{\bar{p}}}}{p} &= \frac{1}{2} (k-1) (p)^{(k-1)=2} \\ &+ \frac{(1)^{(k-1)=2}}{2^{\frac{p}{\bar{p}}}} (2)^k \\ &+ \sum_{j=0}^{(k-1)=2} (p)^j \frac{B_{k+1-2j} B_{2j}}{(k+1-2j)! (2j)!} \end{aligned}$$

The specific result follows after choosing $k = 3$ and $p = 3$.

6. SOME BERNOULLI NUMBER IDENTITIES

In addition to the previously noted identity

$$0 = \sum_{j=0}^{2m+1} (1)^j \frac{B_{4m-2j+2} B_{2j}}{(4m-2j+2)! (2j)!}$$

there are several other identities on sums of Bernoulli numbers that result from the previous developments. These are briefly stated here.

Theorem 6.1. *For integer m , one has*

$$0 = \sum_{j=0}^{2m} (4)^j \frac{B_{4m-4j+2} B_{4j}}{(4m-4j+2)! (4j)!} + 2 \frac{B_{4m-4j} B_{4j+2}}{(4m-4j)! (4j+2)!}$$

Proof. Consider the sums resulting from $0 = I_{4m+1}(2(1+i)) - I_{4m+1}(2(1-i))$.

Theorem 6.2. For integer m , one has

$$\sum_{k=0}^{\infty} (-1)^k \frac{B_{4m-2k} B_{2k}}{(4m-2k)!(2k)!} = \sum_{j=0}^{\infty} (-4)^j \frac{B_{4m-4j} B_{4j}}{(4m-4j)!(4j)!} \\ = \sum_{j=0}^{\infty} (-4)^j \frac{B_{4m-4j-2} B_{4j+2}}{(4m-4j-2)!(4j+2)!}$$

Proof. Consider the sums resulting from the identity

$$I_{4m-1}(z) = I_{4m-1}(z(1+i)) + I_{4m-1}(z(1-i))$$

7. MODULAR RELATIONS

The various sums and quantities above can be seen to be quasi-modular by making a simple change of variable, namely by making the substitution $x = 2i$. By “quasi-modular”, it is meant that the various terms almost have simple behaviors under the Mobius transformation $z \mapsto (a+z)/(b+z)$ for integer $a; b; c$ and d . These relationships are brought to focus here.

First, define $K_k(z) = I_k(2i)$. This change of variable results in a definition which seems simpler than that for I_k . An expansion in z is often referred to as a “Fourier series” in the context of hyperbolic geometry:

$$K_k(z) = \frac{z^{-k-1}}{2} \sum_{j=0}^{\infty} \binom{k}{j} \frac{(2i)^k}{2} \sum_{j=0}^{\lfloor \frac{k-1}{2} \rfloor} \frac{B_{2j} B_{k+1-2j}}{(2j)!(k+1-2j)!}$$

This quantity is almost a modular form of weight $k-1$, in that

$$K_k\left(\frac{1}{z}\right) = z^{k-1} K_k(z)$$

Its only “almost” a modular form, because it is not periodic in z , that is

$$K_k(z+1) \notin K_k(z)$$

By contrast, $P_k(z) = S_k(2i)$ is periodic:

$$P_k(z+1) = P_k(z)$$

but is not quite modular under inversion:

$$P_k\left(\frac{1}{z}\right) = z^{k-1} P_k(z) + K_k\left(\frac{1}{z}\right)$$


One may define a simple variant of the sums that does have a simple transformation under inversion, namely

$$M_k(z) = P_k(z) - \frac{1}{2} K_k(z)$$

which transforms as

$$M_k\left(\frac{1}{z}\right) = z^{k-1} M_k(z)$$

However, M_k is then not periodic.

FIGURE 7.1. Phase of $P_{-1}(q)$ on the unit disk

This graphic shows the phase

$$\arg P_{-1}(q) = \arg \prod_{n=1}^{\infty} \frac{q^n}{q^n - 1}$$

where $\arg f(z) = \text{Im} \log f(z)$ is the usual arg of a function. The color scheme is such that black represents areas where $\arg > 0$ and red represents areas where $\arg < 0$. The absence of other colors indicates that the phase is rather closely confined to the vicinity of 0 for most all of the disk. Numerically, the absolute value of the phase is smaller than 10^{-3} for much of the disk. In particular, this indicates that there are no zeros at all in the interior of the disk, as a zero would be surrounded by a region where the phase wraps around by 2π .

The function $P_s(q)$ does have poles at $q = e^{2\pi i m/n}$ for all rationals m/n ; these are visible at the edges of the disk. The fractal nature of this image is the characteristic signature of a modular form of weight 2; the self-similar regions are just copies of the fundamental region of the modular group $SL(2; \mathbb{Z})$.

FIGURE 7.2. Phase of $P_{-5}(q)$ on the unit disk


This graphic shows the phase

$$\arg P_{-5}(q) = \arg \prod_{n=1}^{\infty} \frac{n^5}{q^n - 1}$$

The color scheme is as in the previous image, and similar conclusions apply: there are no zeros at all in the interior of the disk. The absolute value of the phase is tiny: numerically, it is within 10^{-9} of zero for much of the disk.

The fractal nature of this image is the characteristic signature of a modular form of weight 6; it should be compared to the image of the modular invariant \mathfrak{g}_3 shown in the next image.

The analytic structure of P_k is curious: it has a pole at $q = i1$ and, by the inversion formula and periodicity, at every rational value of q . This is clearly visible in the graphic

FIGURE 7.3. Graph of the modular invariant g_3 .

This figure shows the imaginary part of the modular invariant g_3 , one of the invariants of an elliptic curve. Specifically, it shows the imaginary part of

$$g_3(q) = \frac{8}{27} \left(1 - 504 \sum_{n=1}^{\infty} \frac{n^5 q^n}{1 - q^n} \right)$$

which is a modular form of weight 6. The colors are chosen such that black represents areas that are negative, blue and green represent areas with smaller values, and red those areas with the largest values.

below, which shows P_3 on the q -series or “punctured disk” coordinates $q = e^{2\pi i}$:

$$P_s(q) = \sum_{n=1}^{\infty} \frac{1}{n^s (q^n - 1)}$$

In this form, the relation to modular forms becomes the most immediate. Consider the Lambert series:

$$\sum_{n=1}^{\infty} \frac{1}{n^s} \frac{q^n}{1 - q^n} = (s) P_s(q) = \sum_{m=1}^{\infty} d_s(m) q^m$$

Here, $d_s(m)$ is the divisor function:

$$d_s(m) = \sum_{n \mid m} n^{-s}$$

with the notation $\sum_{n \mid m}$ denoting that the sum extends over all divisors n of m . This should be compared to the Eisenstein series [1, see section 3.10]

$$G_{2k}(q) = 2(2k) + \sum_{m=1}^{\infty} \frac{2(2k-1)2^k}{(2k-1)!} d_{2k-1}(m) q^m$$

etc. In essence, we are dealing with an extension of the Eisenstein series to general complex argument. XXX unfinished; this should be elaborated on a bit more. XXXX

8. CONCLUSIONS

Given any particular rational and set of sums, there is a rich variety of possible relationships that can be formed. It would be interesting to characterize the number of these relationships as a function of the number of sums involved. Thus, for example, given only $S_3(2)$, there seems to be only one relationship that can be formed to give (3). However, given $S_3(1)$, $S_3(2)$ and $S_3(4)$, there seem to be two distinct relationships, of which a linear combination can be taken: thus there is a one-dimensional linear space of solutions. It would be interesting to know what the higher order combinatorial possibilities may be, the dimensionality of the spaces they form, and how these depend on the selected series.

Another interesting avenue of research would be to strengthen the connections to modular forms, to see if additional insights can be gained by following that route. A different and fruitful avenue for understanding almost-modular structures is to pose them in terms of functions on a one-dimensional lattice (in the sense of the Ising or Potts model of physics), where translation is described by the shift operator. It is known that such lattice models are isomorphic to measure-preserving dynamical systems, such as the Baker’s map or more generally subshifts of finite type. In particular, it seems possible to re-express the functional equation for S_k , as given in section 4, in terms of (logarithms of) eigenfunctions of the Baker’s map.

Thanks to Simon Plouffe for generating interest in such sums.

9. APPENDIX: POLYLOG INTEGRAL

This appendix proves the following theorem:

Theorem 9.1. *The polylogarithm may be written as the integral*

$$Li_s^+(e^{-u}) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} (r) (r+s) u^{-r} dr$$

Proof. The proof below is cribbed from the Wikipedia article on polylogarithms[6]. One begins by writing the the Mellin transform of the polylog, as

$$M_s(r) = \int_0^1 \text{Li}_s(y) e^{-u} u^r \frac{du}{u}$$

Using an integral representation of the polylogarithm,

$$\text{Li}_s(w) = \frac{1}{(s)_0} \int_0^1 \frac{t^{s-1}}{w^{-1}e^t - 1} dt$$

and substituting, one obtains

$$M_s(r) = \frac{1}{(s)_0} \int_0^1 \int_0^1 \frac{u^{r-1} t^{s-1}}{y^{-1}e^{t+u} - 1} dt du$$

A change of variable $t = ab$ and $u = a(1-b)$ with $dt du = a da db$ gives

$$\begin{aligned} M_s(r) &= \frac{1}{(s)_0} \int_0^1 b^{s-1} (1-b)^{r-1} db \int_0^1 \frac{a^{r+s-1}}{y^{-1}e^a - 1} da \\ &= (r) \text{Li}_{r+s}(y) \end{aligned}$$

The inverse Mellin transform may now be employed to write

$$\text{Li}_s(y) e^{-u} = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} u^{-r} (r) \text{Li}_{r+s}(y) dr$$

By setting $y = 1$, one then uses $\text{Li}_{s+r}(1) = (s+r)$ to obtain the desired result.

REFERENCES

- [1] Tom M. Apostol. *Modular Functions and Dirichlet Series in Number Theory*. Springer, 2nd ed. edition, 1990.
- [2] Bruce C. Berndt. *Ramanujan's Notebooks*, volume II. Springer-Verlag, 1985.
- [3] Richard E. Crandall Jonathan M. Borwein, David M. Bradley. Computational strategies for the riemann zeta function. *J. Comp. App. Math.*, 21:11, 2000. <http://www.maths.ex.ac.uk/~mwatkins/zeta/borwein1.pdf>.
- [4] Simon Plouffe. Identities inspired from ramanujan notebooks ii. <http://www.lacim.uqam.ca/~plouffe/identities.html>, July 21 1998.
- [5] Simon Plouffe. Identities inspired by ramanujan notebooks (part 2). <http://www.lacim.uqam.ca/~plouffe/inspired2.pdf>, April 2006.
- [6] Wikipedia. Polylogarithm, 2006. <http://en.wikipedia.org/wiki/Polylogarithm>.

—

—

E-mail address: <linas@linas.org>