FIBERS OF CHARACTERS IN GELFAND-TSETLIN CATEGORIES

VYACHESLAV FUTORNY AND SERGE OVSIENKO

ABSTRACT. We solve the problem of extension of characters of commutative subalgebras in associative (noncommutative) algebras for a class of subrings (Galois rings) in skew group rings. These results can be viewed as a noncommutative analogue of liftings of prime ideals in the case of integral extensions of commutative rings. The proposed approach can be applied to the representation theory of many infinite dimensional algebras including universal enveloping algebras of reductive Lie algebras, Yangians and finite W-algebras. In particular, we advance in the representation theory of gl_n developing a theory of Gelfand-Tsetlin modules. Besides classification results we characterize their categories in the generic case.

Contents

1. Introduction]
2. Preliminaries	4
2.1. Notations	4
2.2. Integral extensions	ţ
2.3. Skew (semi)group rings	ţ
2.4. Galois rings	ţ
2.5. Galois orders	r
2.6. Harish-Chandra subalgebras	r
3. Gelfand-Tsetlin categories	8
3.1. Motivation	8
3.2. Gelfand-Tsetlin modules	(
3.3. A realization of Gelfand-Tsetlin modules	13
4. Generic representations	17
5. Representations of Galois orders	19
5.1. Extension of characters for Galois orders	19
5.2. Gelfand-Tsetlin modules over Galois orders	2.
5.3. Gelfand-Tsetlin modules for Galois orders	26
5.4. Gelfand-Tsetlin categories for Galois orders	26
5.5. Proof of Theorem B	28
6. Gelfand-Tsetlin modules for gl_n	28
7. Acknowledgment	29
References	29

1. Introduction

The functors of restriction onto subalgebras and induction from subalgebras are important tools in the representation theory. The effectiveness of these tools

depends upon the choice of a subalgebra. Denote by Specm A (Spec A) the space of maximal (prime) ideals in A, endowed with the Zarisky topology. In the classical commutative algebra setup, an integral extension $A \subset B$ of two commutative rings induces a map $\varphi: \operatorname{Spec} B \to \operatorname{Spec} A$, whose fibers are non-empty for every point of $\operatorname{Spec} A$ (e.g. $A = B^G$, where G is a finite subgroup of the automorphism group of B). In particular, every character of A can be extended to a character of its integral extension B. Moreover, if B is finite over A then all fibers $\varphi^{-1}(I)$, $I \in \operatorname{Spec} A$ are finite, and hence, the number of extensions of a character of A is finite. The Hilbert-Noether theorem provides an example of such situation with B being the symmetric algebra on a finite-dimensional vector space V and $A = B^G$, where G is a finite subgroup of GL(V).

The primary goal of this paper is to generalize these results to "semi-commutative" pairs $\Gamma \subset U$ where U is an associative (noncommutative) algebra over a base field \mathbbm{k} and Γ is an integral domain. The canonical embedding $\Gamma \subset U$ induces a functor from the category of U-modules which are direct sums of finite dimensional Γ -modules (Gelfand-Tsetlin modules with respect to Γ) to the category of torsion Γ -modules. This functor induces a "multivalued function" from Specm Γ associating to an ideal $\mathbbm{m} \in \operatorname{Specm} \Gamma$ the fiber $\Phi(\mathbbm{m})$ of left maximal ideals of U that contain \mathbbm{m} . Our goal is to find natural sufficient conditions for the fibers to be non-empty and finite for any point in $\operatorname{Specm} \Gamma$. On the other hand, for a maximal left ideal $I \subset U$ such that U/I is a Gelfand-Tsetlin module, it is interesting to investigate its support in $\operatorname{Specm} \Gamma$ (i.e. the set of $\mathbbm{m} \in \operatorname{Specm} \Gamma$, such that Γ/\mathbbm{m} is a subfactor of U/I as a Γ -module) and find the multiplicity of Γ/\mathbbm{m} in U/I.

A motivation for the study of such pairs (U, Γ) comes from the representation theory. The classical framework of *Harish-Chandra modules* ([D], Ch. 9) is related to a pair of a reductive Lie algebra \mathcal{F} and its reductive subalgebra \mathcal{F}' , where Uand Γ are their universal enveloping algebras respectively. A more general concept of Harish-Chandra modules (related to a pair (U, Γ)) was introduced in [DFO2].

The case when U is the universal enveloping algebra of a reductive finite dimensional Lie algebra and Γ is the universal enveloping algebra of a Cartan subalgebra leads to the theory of Harish-Chandra modules with respect to this Cartan subalgebra, commonly known as generalized weight modules. Classification of such simple modules is well known for gl_2 and for any simple finite-dimensional Lie algebra for modules with finite-dimensional weight spaces, due to Fernando [Fe] and Mathieu [Ma]. It remains an open problem in general. To approach this classification problem, the full subcategory of weight Gelfand-Tsetlin $U(\mathrm{gl}_n)$ -modules with respect to the Gelfand-Tsetlin subalgebra was introduced in [DFO1]. This class is based on natural properties of a Gelfand-Tsetlin basis for finite-dimensional representations of simple classical Lie algebras [GTs], [Zh], [M].

Gelfand-Tsetlin subalgebras were considered in [FM] in connection with the solutions of the Euler equation, in [Vi] in connection with subalgebras of maximal Gelfand-Kirillov dimension in the universal enveloping algebra of a simple Lie algebra, in [KW1], [KW2] in connection with classical mechanics, and also in [Gr] in connection with general hypergeometric functions on the Lie group $GL(n, \mathbb{C})$. A similar approach was used by Okunkov and Vershik in their study of the representations of the symmetric group S_n [OV], with U being the group algebra of S_n and Γ being the maximal commutative subalgebra generated by the Jucys-Murphy elements

$$(1i) + \ldots + (i-1i), \quad i = 1, \ldots, n.$$

In this case the elements of Specm Γ parametrize basis of irreducible representations of U. Recent advances in the representation theory of Yangians ([FMO]) and finite W-algebras ([FMO1]) are also based on similar techniques.

What is the intrinsic reason of the existence of Gelfand-Tsetlin formulae and of the successful study of Gelfand-Tsetlin representations of various classes of algebras? This question led us to the introduction in [FO1] of concepts of *Galois rings* and *Galois orders* in invariant skew (semi)group rings.

For the rest of the paper we assume that Γ is a commutative domain, K the field of fractions of Γ , $K \subset L$ a finite Galois extension, $\mathbb{M} \subset \operatorname{Aut} L$ a submonoid closed under conjugation by the elements of the Galois group G = G(L/K). Then the group G acts in a natural way on the skew group algebra $L * \mathbb{M}$. A Galois ring U over Γ is a finitely generated over Γ subring in the ring of G-invariants $\mathcal{K} = (L * \mathbb{M})^G \subset L * \mathbb{M}$ such that $KU = UK = \mathcal{K}$.

A Galois ring U over Γ is called right (respectively left) order if for any finite dimensional right (respectively left) K-subspace $W \subset \mathcal{K}$ (respectively $W \subset \mathcal{K}$), $W \cap U$ is a finitely generated right (respectively left) Γ -module. A Galois ring is order if it is both right and left order ([FO1]). Galois orders are natural versions of "noncommutative orders" in skew semigroup rings of invariants. In comparison with the classical notion of an order we note, that $\Gamma \subset U$ is not central, but maximal commutative subalgebra.

A class of Galois orders includes in particular the following subrings in the corresponding skew group rings ([FO1]): Generalized Weyl algebras over integral domains with infinite order automorphisms, e.g. n-th Weyl algebra A_n , quantum plane, q-deformed Heisenberg algebra, quantized Weyl algebras, Witten-Woronowicz algebra among the others ([Ba], [BavO]); the universal enveloping algebra of gl_n over the Gelfand-Tsetlin subalgebra ([DFO1], [DFO2]), associated shifted Yangians and finite W-algebras ([FMO], [FMO1]); certain rings of invariant differential operators on torus.

In this paper we develop a representation theory of Galois rings. We emphasize that the theory of Galois orders unifies the representation theories of universal enveloping algebras and generalized Weyl algebras. Our main result establishes sufficient conditions for the fiber $\Phi(m)$ to be nontrivial and finite. Let ℓ_m be any lifting of m to the integral closure of Γ in L, \mathcal{M}_m the stabilizer of ℓ_m in \mathcal{M} .

Theorem A. Let Γ be a commutative domain which is finitely generated as a k-algebra, U a Galois ring over Γ , $m \in \operatorname{Specm} \Gamma$. Suppose that \mathcal{M}_m is finite.

- If U is a right Galois order over Γ then the fiber $\Phi(m)$ is non-empty.
- If U is a Galois order over Γ then the fiber $\Phi(m)$ is finite.

This result guarantees that a class of Galois orders over Γ has a "nice" theory of Gelfand-Tsetlin modules (cf. Section 5.4), namely, for any $m \in \operatorname{Specm} \Gamma$ with finite \mathcal{M}_m there are finitely many isomorphism classes of simple Gelfand-Tsetlin modules whose support contains m (e.g. this number is limited by $1!2!\dots(n-1)!$ for gl_n , the same number bounds the dimensions of Γ -subspaces of these modules).

Note, that in the case $U=U(\operatorname{sl}_2)$ the structure of the category of Gelfand-Tsetlin modules is well known. This category splits into a direct sum of blocks. Each block is equivalent to the category of finite dimensional representations of complete algebras over $\Bbbk[c]_{\mathsf{m}}$, where $c\in U$ is the Casimir element and $\mathsf{m}\in\operatorname{Specm} \Bbbk[c]$ is a maximal ideal which acts on the corresponding block nilpotently. These algebras are presented by the quivers



The first quiver corresponds to the case of generic blocks, the second case corresponds to the blocks which contains Verma module, but does not contain a finite

dimensional module and the third quiver corresponds to the blocks, containing a finite dimensional sl_2 -module. Besides in this last case holds the relation ab = cd. In the situation of an arbitrary Galois order (even in the case of sl_n) we are far from such detailed description of the blocks of the category of Gelfand-Tsetlin modules. Nevertheless we are able to characterize the structure of generic blocks. We show that the block of generic Gelfand-Tsetlin modules looks exactly ass in the case of sl_2 , moreover there is enough of such representations to guarantee the faithful action of U on these modules. Namely, we have

Theorem B. Let Γ a commutative domain which is finitely generated and normal as a \mathbb{k} -algebra, U a Galois order over Γ with M being a group.

- (1) There exists a massive subset $W \subset \operatorname{Specm} \Gamma$ such that for any $m \in W$, $|\Phi(m)| = 1$ and hence there exists a unique simple U-module L_m whose support contains m. Moreover, the extension category generated by L_m contains all indecomposable modules whose support contains m and it is equivalent to the category $\Gamma_m \operatorname{mod}$
- (2) If M is a group, then for any nonzero $u \in U$ there exists a massive set of non-isomorphic simple Gelfand-Tsetlin U-modules on which u acts non-trivially.

Obtained results allow to study effectively a large category of modules over certain infinite-dimensional algebras. In particular, we are able to advance in the classification problem of irreducible weight gl_n -modules beyond the results of [Ma].

In Section 5 we establish sufficient conditions under which a block of a Gelfand-Tsetlin category contains a finite number of non-isomorphic simple modules (Corollary 5.4, Corollary 5.5).

In this paper we apply the theory only to Lie algebras of type A but we believe it can be extended to other types. Developed technique was used in [FMO1] to address the classification problem of irreducible Gelfand-Tsetlin modules for finite W-algebras and shifted Yangians associated with gl_n and to prove an analogue of the Gelfand-Kirillov conjecture for these algebras.

2. Preliminaries

2.1. Notations. All fields in the paper contain the base field k, which is algebraically closed of characteristic 0. All algebras in the paper are k-algebras. If A is an associative ring then by $A-\operatorname{mod}$ we denote the category of finitely generated left A-modules. Let $\mathcal C$ be a k-category, i.e. all $\operatorname{Hom}_{\mathcal C}$ -sets are endowed with a structure of a k-vector space and all the compositions maps are k-bilinear. The category of k-modules k-mod is defined as the category of k-linear functors k-mod, where k-mod is the category of k-vector spaces. The category of locally finitely generated k-modules we denote by k-mod.

For the rest of the paper we assume that Γ is a commutative domain, K the field of fractions of Γ , $K \subset L$ a finite Galois extension, $\mathcal{M} \subset \operatorname{Aut} L$ a submonoid closed under conjugation by the elements of the Galois group G = G(L/K). Then G acts on the skew semigroup ring $L*\mathcal{M}$ via the action $(lm)^g = l^g m^g$, where $m^g = g^{-1} mg$. We will always assume that if for any $m_1, m_2 \in \mathcal{M}$ from

$$m_1|_K = m_2|_K$$

follows $m_1 = m_2$. Such monoid \mathcal{M} is called *separating* with respect to K. A Galois ring U over Γ is a finitely generated over Γ subring of the ring $(L*\mathcal{M})^G$ of invariants such that $KU = UK = (L*\mathcal{M})^G$.

Let ℓ_m be any lifting of m to the integral closure of Γ in L, \mathcal{M}_m the stabilizer of ℓ_m in \mathcal{M} . Since m defines \mathcal{M}_{ℓ_m} up to conjugacy, by abuse of notations we denote it by \mathcal{M}_m .

2.2. Integral extensions. Let A be an integral domain, K its field of fractions and \hat{A} the integral closure of A in K. The ring A is called normal if $A = \hat{A}$. The following is standard

Corollary 2.1. (S], Ch. III, Prop. 11, Prop. 16) If \tilde{A} is noetherian, e.g. A is a finitely generated k-algebra, then \bar{A} is finite over \hat{A} . In this case \hat{A} is finite over A.

Let $i:A\hookrightarrow B$ be an integral extension. Then it induces a surjective map $\operatorname{Specm} B \to \operatorname{Specm} A$ ($\operatorname{Spec} B \longrightarrow \operatorname{Spec} A$). In particular, for any character χ : $A \to \mathbb{k}$ there exists a character $\tilde{\chi}: B \to \mathbb{k}$ such that $\tilde{\chi}|_A = \chi$. If, in addition, B is finite over A, i.e. finitely generated as an A-module, then the number of different characters of B which correspond to the same character of A, is finite. Hence we have in particular

Corollary 2.2. If A is a finitely generated k-algebra then for any character χ : $A \to \mathbb{k}$ there exists finitely many characters $\tilde{\chi} : \bar{A} \to \mathbb{k}$ such that $\tilde{\chi}|_A = \chi$.

2.3. Skew (semi)group rings. Let R be a ring, \mathcal{M} a submonoid of Aut R. The skew semigroup ring, R * M, is a free left R-module with a basis M and with the multiplication

$$(r_1m_1)\cdot (r_2m_2) = (r_1r_2^{m_1})(m_1m_2), \quad m_1, m_2 \in \mathcal{M}, \ r_1, r_2 \in R.$$

If $x \in R * M$ and $m \in M$ then denote by x_m the element in R such that $x = \sum_{m \in \mathcal{M}} x_m m$. Set

$$\operatorname{supp} x = \{ m \in \mathcal{M} | x_m \neq 0 \}.$$

If a finite group G acts by automorphisms on R and by conjugations on M then G acts on R * M. Denote by $(R * M)^G$ the invariants under this action. Then $x \in (R * \mathcal{M})^G$ if and only if $x_{m^g} = x_m^g$ for $m \in \mathcal{M}, g \in G$. For $\varphi \in \text{Aut } R$ and $a \in R$ set $H_{\varphi} = \{h \in G | \varphi^h = \varphi\}$ and

(2)
$$[a\varphi] := \sum_{g \in G/H_{\varphi}} a^g \varphi^g \in (R * \mathfrak{M})^G.^1$$

Then

(3)
$$R * \mathcal{M}^{G} = \bigoplus_{\varphi \in G \setminus \mathcal{M}} (R * \mathcal{M})_{\varphi}^{G}, \text{ where}$$
$$(R * \mathcal{M})_{\varphi}^{G} = \{ [a\varphi] \mid a \in R^{H_{\varphi}} \}.$$

(4)
$$\gamma \cdot [a\varphi] = [(a\gamma)\varphi], \ [a\varphi] \cdot \gamma = [(a\gamma^{\varphi})\varphi], \ \gamma \in R^G,$$
 and $[a\varphi] = [\varphi a^{\varphi^{-1}}].$

Remark 2.1. The formulae (4) means, that as left R-module $R[a\varphi]R$ is isomorphic to RR^{φ} and as a right R-module $R[a\varphi]R$ is isomorphic to RR^{φ}

2.4. Galois rings. Let R = L be a field, $K \subset L$ a finite Galois extension of fields, G = G(L/K) its Galois group and i the canonical embedding $K \hookrightarrow L$. Denote $K = L^{\hat{G}}$.

Let $S \subset \mathcal{M}$ be a finite G-invariant subset, $V \subset \mathcal{K}$ a Γ -subbimodule. Then we introduce the following Γ -subbimodule in V

(5)
$$V(S) = \{ x \in V \mid \operatorname{supp} x \subset S \}$$

¹The notation $g \in G/H_{\varphi}$ means that g runs over a set of representatives of cosets from G/Hand the result does not depend on a choice of these representatives.

We need the following simple fact.

Lemma 2.1. In the assumption above the right and the left dimensions of $\mathfrak{K}(S)$ coincides with |S/G|.

In particular, for any $\varphi \in \mathcal{M}$ holds

(6)
$$\dim_K \mathcal{K} = [L^{H_{\varphi}} : K] = [G : H_{\varphi}] = |G\varphi|.$$

Definition 1. (1) A monoid $\mathcal{M} \subset \operatorname{Aut} L$ is called separating (with respect to K) if for any $m_1, m_2 \in \mathcal{M}$ the equality $m_1|_K = m_2|_K$ implies $m_1 = m_2$.

(2) An automorphism $\varphi: L \longrightarrow L$ is called separating (with respect to K) if the monoid generated by $\{\varphi^g \mid g \in G\}$ in Aut L is separating.

Lemma 2.2. ([FO1], Proposition 2.3) Let M be a separating monoid with respect to K. Then

- $(1) \ \mathcal{M} \cap G = \{e\}.$
- (2) For any $m \in \mathcal{M}$, $m \neq e$ there exists $\gamma \in K$ such that $\gamma^m \neq \gamma$.
- (3) If $Gm_1G = Gm_2G$ for some $m_1, m_2 \in \mathcal{M}$, then there exists $g \in G$ such that $m_1 = m_2^g$.
- (4) If M is a group, then the statements (1), (2), (3) are equivalent and each of them implies that M is separating.

Recall from the introduction that an associative ring U containing Γ is called a Galois Γ -ring if it is finitely generated over Γ (Γ -subring) in \mathcal{K} and $KU = UK = \mathcal{K}$.

Remark 2.2. Since in the text below we will work with a fixed Galois ring, we will use the following agreement. We assume, that by definition every Galois U ring is endowed with the following data, presented in the definition of a Galois ring:

$$\Gamma \subset K, K \subset L, G = G(L/K), \mathcal{M} \subset \text{Aut } L \text{ is a separating monoid}$$

$$g\mathcal{M}g^{-1}=\mathcal{M},g\in G;\mathcal{K}=(L*\mathcal{M})^G,$$

and with a canonical embedding $i: U \hookrightarrow \mathcal{K}$, identical on Γ .

 $K, L, G, \mathcal{M}, \mathcal{K}, V(S)$, to explain the notation |S/G| - all ingredients! $I_L(\Gamma)$

Let U be a Galois ring over Γ in \mathcal{K} with a fixed embedding $i:\Gamma\to U$.

Lemma 2.3. (compare [FO1], Lemma 4.1) Let $u \in U$ be nonzero element,

$$T = \operatorname{supp} u, \ u = \sum_{m \in T} [a_m m].$$

Then

$$K(\Gamma u\Gamma) = (\Gamma u\Gamma)K = KuK = \bigoplus_{m \in T} V(a_m m),$$

where $V(a_m m) = K[a_m m]K$.

In particular it shows that for every $m \in \mathcal{M}$, U contains the elements $[b_1 m], \ldots, [b_k m]$ where b_1, \ldots, b_k is a K-basis in L^{H_m} . Let $e \in \mathcal{M}$ be the unit element and $U_e = U \cap Le$.

Theorem 2.1. (FO1], Theorem 4.1) Let U be a Galois ring over Γ . Then

- (1) $U_e \subset K$.
- (2) $U \cap K$ is a maximal commutative k-subalgebra in U.
- (3) The center Z(U) of U equals $U \cap K^{\mathfrak{M}}$.

2.5. Galois orders. In this section we recall the basic properties of Galois orders following [FO1]. For simplicity we only consider right Galois orders. Let M be a right Γ -submodule of a Galois order U. Set

$$\mathbb{D}_r(M) = \{ u \in U \mid \text{ there exists } \gamma \in \Gamma, \gamma \neq 0 \text{ such that } u \cdot \gamma \in M \}.$$

We have the following characterization of Galois orders.

Proposition 2.1. (FO1], Proposition 5.1) A Galois ring U over a noetherian Γ is a right order if and only if for every finitely generated right Γ -module $M \subset U$, the right Γ -module $\mathbb{D}_r(M)$ is finitely generated.

In particular, if U is right integral then $\Gamma \subset U_e$ is an integral extension and U_e is a normal ring.

2.6. Harish-Chandra subalgebras. Let U be a Galois ring over a commutative domain Γ . Assume U to be a \mathbb{k} -algebra.

Recall that Γ is called a Harish-Chandra subalgebra in U if $\Gamma u\Gamma$ is finitely generated both as a left and as a right Γ -module for any $u \in U$ [DFO2]. We will also say that Γ is a right (left) Harish-Chandra subalgebra if $\Gamma u\Gamma$ is finitely generated as a right (left) Γ -module for any $u \in U$. Harish-Chandra subalgebras of Galois rings have the following characterization.

Proposition 2.2. (compare [FO1], Corollary 5.3) If U is a Galois order over Γ and Γ is a noetherian \mathbb{k} -algebra then Γ is a Harish-Chandra subalgebra in U.

Let $S \subset \mathcal{M}$ be a finite G-invariant subset. Denote

$$U(S) = \{ u \in U \mid \text{supp } u \subset S \}.$$

For every $f \in \Gamma$ consider $f_S^r \subset \Gamma \otimes_{\mathbb{k}} K$ as follows

(7)
$$f_S^T = \prod_{s \in S} (f \otimes 1 - 1 \otimes f^{s^{-1}}) = \sum_{i=0}^{|S|} f^{|S|-i} \otimes T_i, \ (T_0 = 1).$$

Similarly we define $f_S^l = \prod_{s \in S} (f^s \otimes 1 - 1 \otimes f)$.

Lemma 2.4. ([FO1], Lemma 5.2) Let $S \subset M$ be a G-invariant subset, $m^{-1}(\Gamma) \subset \bar{\Gamma}$ $(m(\Gamma) \subset \bar{\Gamma} \text{ respectively}) \text{ for all } m \in \mathcal{M}. \text{ For any subset } X \subset \mathcal{M} \text{ set } f_X = f_X^r$ $(f_X = f_X^l \ respectively).$

- (1) $u \in U(S)$ if and only if $f_S \cdot u = 0$ for every $f \in \Gamma$. (2) If $T = \operatorname{supp} u \setminus S$ then $f_T \cdot u \in U(S)$ for every $f \in \Gamma$.
- (3) If $f_S = \sum_{i=1}^n f_i \otimes g_i$, $[am] \in L * \mathcal{M}$ then

$$f_S \cdot [am] = [(\sum_{i=1}^n f_i g_i^m a)m] = [\prod_{s \in S} (f - f^{ms^{-1}})am].$$

- (4) Let S be a G-orbit and T an G-invariant subset in M. The Γ -bimodule homomorphism $P_S^T(=P_S^T(f)): U(T) \longrightarrow U(S), \ u \mapsto f_{T \setminus S} \cdot u, f \in \Gamma$ is either zero or $\operatorname{Ker} P_S^T = U(T \setminus S)$.
- (5) Let $S = S_1 \sqcup \cdots \sqcup S_n$ be the decomposition of S in G-orbits and $P_{S_s}^S$: $U(S) \longrightarrow U(S_i), i = 1, ..., n$ are defined in (4) nonzero homomorphisms. Then the homomorphism

(8)
$$P^S: U(S) \longrightarrow \bigoplus_{i=1}^n U(S_i), \ P^S = (P_{S_1}^S, \dots, P_{S_n}^S),$$

is a monomorphism.

Theorem 2.2. (cf. [FO1], Theorem 5.1) Let U be a Galois ring over a noetherian Γ and assume that Γ is a right (left) Harish-Chandra \mathbb{k} -subalgebra of U. Then the following statements are equivalent.

- (1) U is right (respectively left) Galois order.
- (2) U(S) is finitely generated right (respectively left) Γ -module for any finite G-invariant $S \subset \mathcal{M}$.
- (3) $U(G \cdot m)$ is finitely generated right (respectively left) Γ -module for any $m \in \mathcal{M}$.

Theorem 2.3. (cf. [FO1], Theorem 5.2) Let $U \subset \mathcal{K}$ be a Galois ring over a noetherian Γ and \mathcal{M} a subgroup of Aut L.

- (1) If U_e is integral extension of Γ and $m^{-1}(\Gamma) \subset \bar{\Gamma}$ (respectively $m(\Gamma) \subset \bar{\Gamma}$), then U is a right (respectively left) Galois order.
- (2) If U_e is an integral extension of Γ and Γ is a Harish-Chandra \mathbb{k} -subalgebra in U, then U is Galois order over Γ .

3. Gelfand-Tsetlin categories

3.1. **Motivation.** The constructions of this section are the main tools we will use to investigate the class of Gelfand-Tsetlin U-modules. First such constructions appeared in [DFO2], but to make our paper more readable, we consider here a special case of a commutative subalgebra Γ and present it in more details. Though we will work with Galois orders, in this section we just assume that Γ is a commutative Harish-Chandra subalgebra in a finitely generated associative algebra U.

Before going into details we give some motivation of the constructions below. Let U be a finite dimensional associative algebra over \mathbb{k} , $R \subset U$ its Levi subalgebra, $\Gamma \subset U$ the center of R. Then $\Gamma = \bigoplus_{i=1}^n \mathbb{k} e_i$, where $\{e_1, \ldots, e_n\}$ is the complete (i.e. $e_1 + \cdots + e_n = 1$) family of mutually orthogonal idempotents. Obviously, $\Gamma \subset U$ is a Harish-Chandra subalgebra.

This data allows to present U as a ring of the $n \times n$ matrices of the form

$$(9) (e_j U e_i)_{i,j=1,\dots,n}, \quad u \longmapsto \begin{pmatrix} e_1 u e_1 & \dots & e_1 u e_n \\ \vdots & \ddots & \vdots \\ e_n u e_1 & \dots & e_n u e_n \end{pmatrix}$$

with the standard multiplication. This presentation is called the two-sided Pierce decomposition of U. Besides, we can associate with U a k-linear category

$$\mathcal{A} = \mathcal{A}(U; \Gamma)$$
 where $\mathrm{Ob}\,\mathcal{A} = \{1, \dots, n\}, \, \mathcal{A}(i, j) = e_j U e_i,$

and the composition of morphism is defined by the multiplication in U. One easy but important remark is the existence of canonical isomorphism

(10)
$$U - \operatorname{Mod} \simeq A - \operatorname{Mod}.$$

If $\Gamma = R$ (equivalently, U is a basic or Morita reduced algebra), then the category \mathcal{A} is presented usually as a quiver with relations. This presentation is the key feature in the investigation of finite dimensional representations of U (see e.g. [DK], [GR] for details).

The definition of \mathcal{A} can be rewritten as follows. As objects of \mathcal{A} one can consider Specm $\Gamma = \{\mathsf{m}_1, \ldots, \mathsf{m}_n\}$, where m_i is the kernel of the projection of Γ onto $\Bbbk e_i$. Besides set $\mathcal{A}(\mathsf{m}_i, \mathsf{m}_j) = \Gamma/\mathsf{m}_j \otimes_{\Gamma} U \otimes_{\Gamma} \Gamma/\mathsf{m}_i$ with the composition of morphisms induced by the product in U. The constructions below can be considered as a generalization of the two sided Pierce decomposition for the case of a commutative Harish-Chandra subalgebra $\Gamma \subset U$, where U is not necessary finite dimensional. As above, we associate with the pair $\Gamma \subset U$ a category \mathcal{A} with $\mathrm{Ob}\,\mathcal{A} = \mathrm{Specm}\,\Gamma$

(see Definition 3 for details). Unfortunately, there is no equivalence of categories of U-modules and A-modules. Instead we have a weaker result for the full subcategory of Gelfand-Tsetlin U-modules (see Theorem 3.2).

3.2. **Gelfand-Tsetlin modules.** In the next two subsection we recall and develop results from [DFO2]. We assume U an algebra over k and $\Gamma \subset U$ is a commutative finitely generated (in particular, noetherian) subalgebra. The following is the key notion of the paper.

Definition 2. A finitely generated U-module M is called Gelfand-Tsetlin module (with respect to Γ) provided that the restriction $M|_{\Gamma}$ is a direct sum of Γ -modules

(11)
$$M|_{\Gamma} = \bigoplus_{\mathsf{m} \in \operatorname{Specm} \Gamma} M(\mathsf{m}),$$

where

$$M(\mathsf{m}) = \{ v \in M | \mathsf{m}^k v = 0 \text{ for some } k \ge 0 \}.$$

When for all $m \in \operatorname{Specm} \Gamma$ and all $x \in M(m)$ holds mx = 0 such Gelfand-Tsetlin module M is called weight module (with respect to Γ). More generally, for a left (right) Γ -module X and $m \in \operatorname{Specm} \Gamma$ we call an element $x \in X$ left (right) m-nilpotent, provided that $m^k x = 0$ (xm^k = 0) for some $k \geq 1$.

All Gelfand-Tsetlin modules form a full abelian and closed with respect to extensions subcategory $\mathbb{H}(U,\Gamma)$ in U-mod. A full subcategory of $\mathbb{H}(U,\Gamma)$ consisting of weight Gelfand-Tsetlin we denote by $\mathbb{H}W(U,\Gamma)$.

The support of a Gelfand-Tsetlin module M is a set

$$\operatorname{supp} M = \{ \mathsf{m} \in \operatorname{Specm} \Gamma \mid M(\mathsf{m}) \neq 0 \}.$$

For $D \subset \operatorname{Specm} \Gamma$ denote by $\mathbb{H}(U,\Gamma,D)$ the full subcategory in $\mathbb{H}(U,\Gamma)$ formed by M such that $\operatorname{supp} M \subset D$. For a given $\mathsf{m} \in \operatorname{Specm} \Gamma$ we denote by $\chi_{\mathsf{m}} : \Gamma \longrightarrow \Gamma/\mathsf{m}$ the corresponding character of Γ . Conversely, for a character $\chi : \Gamma \longrightarrow \mathbb{k}$ denote $\mathsf{m}_{\chi} = \operatorname{Ker} \chi$, so we will identify the set of all characters of Γ with $\operatorname{Specm} \Gamma$. If there exists a simple Gelfand-Tsetlin module M with $M(\mathsf{m}) \neq 0$ then we say that the character χ_{m} lifts to M.

For any pair $(m, n) \in \operatorname{Specm} \Gamma \times \operatorname{Specm} \Gamma$, $a \in U$ and $m, n \geq 0$ we introduce the left Γ -module $L_{a,m^m} = \Gamma a \Gamma / \Gamma a m^m$ and the right Γ -module $R_{a,n^n} = \Gamma a \Gamma / n^n a \Gamma$.

We assume that Γ is a Harish-Chandra subalgebra in U. Then has a natural structure of Γ -bimodule, equivalently, $\Gamma \otimes \Gamma$ -module. For a Γ -bimodule V we will use the following notations:

$$_{\mathbf{n}^n}V = V/\mathbf{n}^nV, \quad V_{\mathbf{m}^m} = V/V\mathbf{m}^m, \quad _{\mathbf{n}^n}V_{\mathbf{m}^m} = V/(\mathbf{n}^nV + V\mathbf{m}^m).$$

Lemma 3.1. In the assumption above holds the following.

- (1) The modules $L_{a,m}$, $R_{a,n}$ are finite dimensional.
- (2) Any from the following three conditions
 - (a) Γ/n is the subquotient of $\Gamma a\Gamma/\Gamma am$ as a left Γ -module.
 - (b) Γ/m is the subquotient of $\Gamma a \Gamma/na \Gamma$ as a right Γ -module.
 - (c) $\Gamma/\mathsf{n} \otimes_{\Gamma} \Gamma a \Gamma \otimes_{\Gamma} \Gamma/\mathsf{m} \neq 0$.

defines the same set X_a of pairs $(m, n) \in \operatorname{Specm} \Gamma \times \operatorname{Specm} \Gamma$.

(3) For $m, n \in \operatorname{Specm} \Gamma$ define the set

$$X_a(\mathsf{m}) = \{\mathsf{n} \in \operatorname{Specm} \Gamma \mid (\mathsf{m}, \mathsf{n}) \in X_a\}$$

and the set

$$X^a(\mathsf{n}) = \{\mathsf{m} \in \operatorname{Specm} \Gamma \mid (\mathsf{m}, \mathsf{n}) \in X_a\}.$$

Then both $X_a(\mathbf{m})$ and $X^a(\mathbf{n})$ are finite. Besides, the kernels of simple subquotient of all L_{a,\mathbf{m}^m} (respectively R_{a,\mathbf{n}^n}) belong to X_a (respectively X^a).

(4) Let M be a Gelfand-Tsetlin U-module, $m \in \operatorname{Specm} \Gamma$. Then

(12)
$$aM(\mathsf{m}) \subset \sum_{\mathsf{n} \in X_a(\mathsf{m})} M(\mathsf{n}).$$

(5) Let M be a Gelfand-Tsetlin U-module, $\pi_n : M \longrightarrow M(n)$, $n \in \operatorname{Specm} \Gamma$ the canonical projection and $m \notin X^a(n)$. Then

(13)
$$\pi_{\mathsf{n}}(aM(\mathsf{m})) = 0.$$

(6) If X is a finite-dimensional Γ -module then $U \otimes_{\Gamma} X$ is a Gelfand-Tsetlin module.

Proof. We will prove the statements for $L_{a,m}$. The case of $R_{a,m}$ is analogous.

Since Γ is finitely generated, $\dim_k \Gamma/\mathsf{m}^m < \infty$. Then $L_{a,\mathsf{m}^m} \simeq \Gamma a \Gamma \otimes_{\Gamma} \Gamma/\mathsf{m}^m$ is finite dimensional, since $\Gamma a \Gamma$ is finitely generated as right Γ -module. That proves (1). It follows that if $L_{a,\mathsf{m}} = \bigoplus_{k=1}^t L_k$ is a decomposition in the direct sum of indecomposable left Γ -modules, then for every $k = 1, \ldots, t$ there exists $\mathsf{n}_k \in \operatorname{Specm} \Gamma$ and $n_k \geq 1$, such that $\mathsf{n}_k^{n_k} L_k = 0$. In particular, the subquotients L_k are isomorphic to Γ/n_k . On the other hand $\Gamma/\mathsf{n} \otimes_{\Gamma} L_k \simeq L_k/\mathsf{n} L_k$ is nonzero if and only if $\mathsf{n} = \mathsf{n}_k$, that, together with (1), proves (2) and (3).

To prove (4) consider any $x \in M(m)$. Then there exists m > 1, such that $m^m x = 0$. It follows that the left Γ -submodule $\Gamma a \Gamma x \subset M$ is the factor of L_{a,m^m} . Then the statement follows from (3). The statement (5) is proved analogously.

To show (6) it is enough to consider the case $\dim_{\mathbb{k}} V = 1$. But this follows from (1).

Denote by Δ the minimal equivalence on Specm Γ containing all X_a , $a \in U$ and by $\Delta(U, \Gamma)$ the set of the Δ -equivalence classes on Specm Γ .

Lemma 3.2. Let X, X' be Gelfand-Tsetlin modules, supp $X \subset D$, supp $X' \subset D'$, where D and D' are different classes of Δ -equivalence. Then

$$\text{Hom}_U(X, X') = 0, \ \text{Ext}_U^1(X, X') = 0.$$

Proof. Obviously, $\operatorname{Hom}_{\Gamma}(X, X') = 0$ all the more $\operatorname{Hom}_{U}(X, X') = 0$. It is enough to prove that every exact sequence in $U - \operatorname{mod}$

$$0 \longrightarrow X' \xrightarrow{\sigma} Y \xrightarrow{\pi} X \longrightarrow 0$$

splits. Since $D \cap D' = \emptyset$, this sequence splits uniquely as a sequence of Γ -modules, thus we can assume

$$Y(\mathsf{m}) = X'(\mathsf{m}) \oplus X(\mathsf{m}), \mathsf{m} \in \operatorname{supp} Y.$$

Since $D \cap D' = \emptyset$, in this sum either $X'(\mathsf{m}) = 0$ or $X(\mathsf{m}) = 0$. For $a \in U$ holds $aX'(\mathsf{m}) \subset X'$ and $aX(\mathsf{m}) \subset X$, by Lemma 3.1, (4). Hence X' and X are U-submodules in Y.

Immediately from Lemma 3.1, (4), (5) and Lemma 3.2 we obtain the following

Corollary 3.1.

$$\mathbb{H}(U,\Gamma) = \bigoplus_{D \in \Delta(U,\Gamma)} \mathbb{H}(U,\Gamma,D).$$

The subcategory $\mathbb{H}(U,\Gamma,D)$, where $D\in\Delta(U,\Gamma)$ will be called a *block* of $\mathbb{H}(U,\Gamma)$.

3.3. A realization of Gelfand-Tsetlin modules. For a Γ -bimodule V denote by ${}_{\mathsf{n}}\hat{V}_{\mathsf{m}}$ the I-adic completion of $\Gamma \otimes_{\mathbb{k}} \Gamma$ -module V, where $I \subset \Gamma \otimes \Gamma$ is a maximal ideal $I = \mathsf{n} \otimes \Gamma + \Gamma \otimes \mathsf{m}$, in other words

$$_{\mathsf{n}}\hat{V}_{\mathsf{m}} = \varprojlim_{n,m} {}_{\mathsf{n}^n}V_{\mathsf{m}^m}.$$

Let B be a Γ -bimodule, satisfying the Harish-Chandra condition, namely every finite generated bimodule in B is finitely generated both from the left and from the right. Denote by F(B) the set of finitely generated Γ -subbimodules in B. Note, that if $V, W \in F(B)$ and $V \subset W$, then the canonical embedding induces a monomorphism ${}_{\mathsf{n}}V_{\mathsf{m}} \hookrightarrow_{\mathsf{n}} W_{\mathsf{m}}$. It allows us to give the following definition.

The finitary completion ${}_{\mathsf{n}}B_{\mathsf{m}}$ of the bimodule B by the ideal $I = \mathsf{n} \otimes \Gamma + \Gamma \otimes \mathsf{m}$ is a $(\widehat{\Gamma \otimes \Gamma})_I$ module

(14)
$${}_{\mathsf{n}}\widetilde{B}_{\mathsf{m}} = \varinjlim_{V \in F(B)} {}_{\mathsf{n}}\widehat{V}_{\mathsf{m}}.$$

For any $V \in F(B)$, $\mathsf{m}, \mathsf{n} \in \operatorname{Specm} \Gamma$, m, n > 0 we have a family of Γ -bimodule morphisms

$${}_{\mathsf{n}}\hat{V}_{\mathsf{m}} \longrightarrow_{\mathsf{n}^n} V_{\mathsf{m}^m} \longrightarrow_{\mathsf{n}^n} B_{\mathsf{m}^m},$$

where the first is the canonical map from the projective limit and the second is induced by the embedding $V \subset B$. This family defines a homomorphism

$$\Psi:_{\mathbf{n}} \widetilde{B}_{\mathbf{m}} = \varinjlim_{V \in F(B)} {}_{\mathbf{n}} \hat{V}_{\mathbf{m}} \longrightarrow \varprojlim_{n,m} {}_{\mathbf{n}} {}_{\mathbf{m}} B_{\mathbf{m}^m} = \widehat{{}_{\mathbf{n}} B}_{\mathbf{m}}$$

If B is finitely generated as Γ -bimodule then Ψ is an isomorphism.

Definition 3. Define a category $A = A_{U,\Gamma}$ with objects $Ob A = Specm \Gamma$. The space of morphisms from m to n is defined as the completion of U, where i.e.

$$\mathcal{A}(\mathsf{m},\mathsf{n}) \ = \ {}_{\mathsf{n}}\widetilde{U}_{\mathsf{m}} \ \big(= \varinjlim_{V \in F(U)} \varprojlim_{n,m} {}_{\mathsf{n}^n}V_{\mathsf{m}^m}. \big)$$

The spaces $\mathcal{A}(m,n)$ are endowed with the standard topology defined by the limits. Besides, any $\mathcal{A}(m,n)$ is endowed with a canonical structure of a completed $\Gamma_n - \Gamma_m$ -bimodule, i.e. a complete $\Gamma_n \hat{\otimes} \Gamma_m$ -module, where $\hat{\otimes}$ is the completed tensor product. Set

$${}^{\mathsf{I}}V_{\mathsf{m}^m} = \{ \bar{a} \in V_{\mathsf{m}^m} \mid {}^{\mathsf{I}}\bar{a} = 0 \text{ for some } l \ge 1 \},$$

 ${}_{\mathsf{n}^n}V^{\mathsf{I}} = \{ \bar{b} \in {}_{\mathsf{n}^n}V \mid \bar{b}{}^{\mathsf{I}} = 0 \text{ for some } l \ge 1 \},$

For $V \subset U$ consider V_{m^m} as a left Γ -module. By Lemma 3.1, (3) for every $a \in V$ the exists a finite set $X_a(\mathsf{m}) = \{\mathsf{n}_1, \ldots, \mathsf{n}_k\}$ and $N \geq 1$, $N = N(a, \mathsf{m})$, such that $\mathsf{n}_1^N \ldots \mathsf{n}_k^N$ annihilates the class \bar{a} of a in $V/V\mathsf{m}^m$. Hence by the Chinese remainder theorem

$$V_{\mathsf{m}^m} = \bigoplus_{\mathsf{I} \in X_\mathsf{m}} {}^\mathsf{I}V_{\mathsf{m}^m}, \text{ where } X_\mathsf{m} = \bigcup_{a \in V} X_a(\mathsf{m})$$

(15) and

$${}_{\mathsf{n}^n}V = igoplus_{\mathsf{I} \in X^\mathsf{n}} {}_{\mathsf{n}^n} \, V^\mathsf{I} \,, \text{ where } X^\mathsf{n} = igcup_{a \in V} X^a(\mathsf{n}).$$

Lemma 3.3. (1) Let $D, D' \subset \operatorname{Specm} \Gamma$ be two different classes of Δ -equivalence, $m \in D$, $n \in D'$. Then $\mathcal{A}(m, n) = 0$.

(2) We have a decomposition of A into a direct sum of its full subcategories,

$$\mathcal{A} = \bigoplus_{D \in \Delta(U,\Gamma)} \mathcal{A}_D,$$

where A_D is the restriction of A on D.

- (3) If for $a \in U$ holds $n \notin X_a(m)$, then the class of a in A(m, n) equals 0.
- (4) If for $a \in U$ holds $m \notin X^a(n)$, then the class of a in A(m,n) equals 0.

Proof. Let $m \in \operatorname{Specm} \Gamma$ and $m \geq 1$. Then

$${}_{\mathsf{n}^n}U_{\mathsf{m}^m}=\Gamma/\mathsf{n}^n\otimes_{\Gamma}U_{\mathsf{m}^m}=\Gamma/\mathsf{n}^n\otimes_{\Gamma}\bigoplus_{l\in X_m}{}^{\mathrm{I}}U_{\mathsf{m}^m}=\bigoplus_{l\in X_m}\Gamma/\mathsf{n}^n\otimes_{\Gamma}{}^{\mathrm{I}}U_{\mathsf{m}^m}.$$

If m and n belong to different classes of Δ -equivalence, then $n \neq l$ and all the summands in the last sum equal 0. It proves (1). The statement (2) follows immediately from (1).

The statements (3) and (4) are proved analogously to (1).

3.3.1. The composition of morphisms. Let V, W be finitely generated Γ -subbimodules in U. Then the bimodule $T \subset U$, spanned on all products vw, $v \in V$, $w \in W$ is finitely generated subbimodule in U, since Γ is a Harish-Chandra subalgebra in U. By $\mu: V \otimes_{\Gamma} W \longrightarrow T$ we denote the map $\mu(v \otimes w) = vw$, $v \in V, w \in W$.

For a Γ -bimodule B denote by ${}_{\mathsf{n}^n}\mathbb{1}_{\mathsf{m}^m} (={}_{\mathsf{n}^n}\mathbb{1}_{\mathsf{m}^m}(B))$ the canonical epimorphism

$$_{\mathsf{n}^n} \mathbb{1}_{\mathsf{m}^m} : B \longrightarrow B/(\mathsf{n}^n B + B \mathsf{m}^m).$$

Lemma 3.4. Let $V, W \subset U$ be finitely generated Γ -bimodules, T = VW, $\mathsf{m}, \mathsf{n} \in \operatorname{Specm} \Gamma, m, n \geq 0$.

- (1) For $p, p' \in \operatorname{Specm} \Gamma$, $p \neq p'$ holds $n^n V^p \otimes_{\Gamma} p' W_{m^n} = 0$.
- (2) The induced by the decomposition (15) homomorphism

$$\Phi: \bigoplus_{\mathsf{p} \in X^{\mathsf{n}} \cap X_{\mathsf{m}}} (\mathsf{n}^n \ V^{\mathsf{p}} \otimes_{\Gamma} \ \mathsf{^p} W_{\mathsf{m}^n}) \longrightarrow \mathsf{n}^n V \otimes_{\Gamma} W_{\mathsf{m}^m}$$

is an isomorphism of Γ -bimodules.

(3) There exists P > 0, (= P(m, n, m, n, V, W)), such that for any $p \ge P$ the canonical projections

$${}_{\mathsf{n}^n}\pi_{\mathsf{p}^p}:{}_{\mathsf{n}^n}V\longrightarrow{}_{\mathsf{n}^n}V_{\mathsf{p}^p},\,{}_{\mathsf{p}^p}\pi_{\mathsf{m}^m}:W_{\mathsf{m}^m}\longrightarrow{}_{\mathsf{p}^p}W_{\mathsf{m}^m}.$$


 $induce\ isomorphisms$

$$_{\mathsf{n}^n}\pi^{\mathsf{p}^p}:_{\mathsf{n}^n}V^{\mathsf{p}}\longrightarrow_{\mathsf{n}^n}V_{\mathsf{p}^p},\ ^{\mathsf{p}^p}\pi_{\mathsf{m}^m}:{}^{\mathsf{p}}W_{\mathsf{m}^m}\longrightarrow_{\mathsf{p}^p}W_{\mathsf{m}^m}.$$

(4) There exists P > 0, such that for $p \ge P$ there exists a unique homomorphism

$$\widehat{\mu}(\mathsf{m}^m,\mathsf{n}^n):\bigoplus_{\mathsf{p}\in X^\mathsf{n}\cap X_\mathsf{m}}(\mathsf{n}^nV_\mathsf{p^p}\otimes_{\widehat{\Gamma}_\mathsf{p}}\mathsf{p^p}W_\mathsf{m^n})\longrightarrow \mathsf{n}^nT_\mathsf{m^m},$$

which makes the diagram below



commutative. Here $n^n \mu_{m^m}$ is induced by μ , Φ is defined following (15) and

$$c_{\mathsf{p}}: {_{\mathsf{n}^n}}\,V^{\mathsf{p}}\,\otimes_{\Gamma}\,{^{\mathsf{p}}\!W_{\mathsf{m}^n}} \xrightarrow{\quad {_{\mathsf{n}^n}}\,\pi^{\mathsf{p}^p}\,\otimes\,{^{\mathsf{p}^p}}\,\pi_{\mathsf{m}^m}} \to {_{\mathsf{n}^n}}V_{\mathsf{p}^p}\otimes_{\Gamma\mathsf{p}^p}W_{\mathsf{m}^n} \longrightarrow {_{\mathsf{n}^n}}V_{\mathsf{p}^p}\otimes_{\widehat{\Gamma}_{\mathsf{n}}}{_{\mathsf{p}^p}}W_{\mathsf{m}^n},$$

where the second arrow is induced by the canonical homomorphism $\Gamma \to \widehat{\Gamma}_{p}$. The homomorphism $\widehat{\mu}(\mathbf{m}^{m}, \mathbf{n}^{n})$ does not depend on $p \geq P$. (5) Denote

$$\widehat{\mu}_{\mathsf{p}}(\mathsf{m}^m,\mathsf{n}^n):{}_{\mathsf{n}^n}V_{\mathsf{p}^p}\otimes_{\widehat{\Gamma}_{\mathsf{n}}\mathsf{p}^p}W_{\mathsf{m}^n}\longrightarrow{}_{\mathsf{n}^n}T_{\mathsf{m}^m}$$

the restriction of $\widehat{\mu}_{p}(m^{m}, n^{n})$. Then

$$\widehat{\mu}(\mathsf{m}^m,\mathsf{n}^n) = \sum_{\mathsf{p} \in X^{\mathsf{n}} \cap X_{\mathsf{m}}} \widehat{\mu}_{\mathsf{p}}(\mathsf{m}^m,\mathsf{n}^n).$$

Besides, in the assumption of (4) there exists $S \geq 1$, such that for every $p \in X^n \cap X_m$, $v \in V$, $w \in W$ and its classes $\bar{v} \in {}_{n^n}V_{p^p}$, $w \in {}_{p^p}W_{m^n}$, $s \geq S$ and

(16) $\gamma \in \Gamma$, such that $\gamma \equiv 1 \mod p^s$, $\gamma \equiv 0 \mod q^s$, $q \in X^n \cap X_m$, $q \neq p$ holds

$$\widehat{\mu}_{p}(\mathsf{m}^{m},\mathsf{n}^{n})(\bar{v}\otimes\bar{w})=\overline{v\gamma w},$$

where $\overline{v\gamma w}$ is the class of $v\gamma w$ in $n^n T_{m^m}$.

(6) Let $m' \leq m$, $n' \leq n$ be integral. Then for large enough p satisfying the conditions of (3) there exists the commutative diagram of $\hat{\Gamma}_n - \hat{\Gamma}_m$ homomorphisms

where the horizontal arrows are induced by the canonical projections and the vertical arrows are of corresponding homomorphisms $\widehat{\mu}_p(\mathsf{m}^m,\mathsf{n}^n)$.

(7) Let $V \subset V'$, $W \subset W'$ be a finitely generated Γ -subbimodules in U, T' = V'W'. Then for large enough p satisfying the conditions of (3) there exists the commutative diagram of $\hat{\Gamma}_n - \hat{\Gamma}_m$ homomorphisms

where the horizontal arrows are induced by the canonical embeddings and the vertical arrows are corresponding homomorphisms $\widehat{\mu}_{p}(\mathbf{m}^{m}, \mathbf{n}^{n})$.

Proof. The statement (1) obviously follows from the Chinese remainder theorem. The statement (2) follows from the decomposition (15) and from the statement (1) above.

To prove the statement (3) note first that the sequence of finite dimensional modules $_{\mathsf{n}^n}V_{\mathsf{p}^p}$ stabilizes beginning from some P hence $_{\mathsf{n}^n}V^{\mathsf{p}}$ is a factor of $_{\mathsf{n}^n}V_{\mathsf{p}^p}$. On other hand every $_{\mathsf{n}^n}\pi_{\mathsf{p}^p}$ factorizes through $_{\mathsf{n}^n}V^{\mathsf{p}}$, which prove (3) for V. The case of W is considered analogously.

Note that for big enough p all c_p are isomorphisms. The first map in the definition of c_p is isomorphism due to (3) and the second map is isomorphism, since both multipliers in the tensor product are finite dimensional p-torsion modules. Hence the third vertical arrow in the diagram is an isomorphism. Besides, Φ is an isomorphism by (2), that proves (4). Independence on p is obvious.

Let $\bar{v} = \sum_{l} \bar{v}_{l}$, $\bar{w} = \sum_{l} \bar{w}$, be the decompositions from (15). Then for $\gamma', \gamma'' \in \Gamma$, satisfying (16) for large enough p hold

$$\bar{v}\gamma' = \bar{v}_{\mathrm{p}} + \sum_{\mathrm{I} \not\in X^{\mathrm{n}} \cap X_{\mathrm{m}}} \bar{v}_{\mathrm{I}}\gamma', \quad \gamma''\bar{w} = {}_{\mathrm{p}}\bar{w} + \sum_{\mathrm{I} \not\in X^{\mathrm{n}} \cap X_{\mathrm{m}}} \gamma''_{\mathrm{I}}\bar{w}.$$

Then following (1) $v\gamma' \otimes \gamma''w = v_{\mathsf{p}} \otimes_{\mathsf{p}} w$, which competed the proof of (5).

The statements (6) and (7) follow from (5). In these diagrams both images of $\bar{v} \otimes \bar{w}$ from the left upper corner of the diagram for convenient $\gamma \in \Gamma$ equal of the class $\overline{v\gamma w}$ in the right lower corner.

Then a composition in the category A is defined as follows. Since direct limits commute with tensor product, we can write down

$$\mathcal{A}(\mathsf{I},\mathsf{n}) \otimes_{\widehat{\Gamma}_\mathsf{I}} \mathcal{A}(\mathsf{m},\mathsf{I}) \simeq \varinjlim_{V \in F(U)} \varinjlim_{W \in F(U)} {}_\mathsf{n} V_\mathsf{I} \otimes_{\widehat{\Gamma}_\mathsf{I}} {}_\mathsf{I} W_\mathsf{m}$$

Then we have the following composition

$${}_{\mathbf{n}}V_{\mathbf{l}}\otimes_{\widehat{\Gamma}_{\mathbf{l}}}{}_{\mathbf{l}}W_{\mathbf{m}}\xrightarrow{\widehat{m}_{\mathbf{l}}(\mathbf{m}^{m},\mathbf{n}^{n})}{}_{\mathbf{n}^{n}}T_{\mathbf{m}^{m}}\xrightarrow{i_{\mathbf{n},\mathbf{m}}}\mathcal{A}(\mathbf{m},\mathbf{n}),$$

where the first homomorphism is constructed above and the second is the canonical map in the direct limit. These maps enter in corresponding commutative diagrams, i.e. Lemma 3.4, (6), hence they define correctly a map

$$\mathcal{A}(p,n) \times \mathcal{A}(m,p) \longrightarrow \mathcal{A}(m,n).$$

- 3.3.2. Generalized Pierce decomposition. We start from the following categorical statement. Assume \mathcal{C} is a \mathbb{k} -category with sums and products and $\{C_i \mid i \in \mathcal{I}\}$ be a family of objects of \mathcal{C} . Denote by (*) and (**) the following properties of this family:
 - (*) for every $j \in \mathcal{I}$ there exist finitely many $i \in \mathcal{I}$, such that $\mathcal{C}(C_i, C_j)$ is nonzero. Consider the vector space

$$\Pi_{\mathcal{I}} = \prod_{i \in \mathcal{I}, j \in \mathcal{I}} \mathcal{C}(C_j, C_i),$$

written as $\mathcal{I} \times \mathcal{I}$ matrices, provided that j's correspond to columns and i's corresponds to rows. In general the standard "column-by-row" product of such matrices isn't correctly defined. By $M_{\mathcal{I}}$ denote the subspaces of $\Pi_{\mathcal{I}}$, formed by the matrices with finitely many nonzero elements in any column and in any row. Then "column-by-row" product turns it into k-algebras.

Lemma 3.5. Assume the family $\{C_i \mid i \in \mathcal{I}\}\$ of objects of the k-category \mathcal{C} satisfies the property (**).

(1) There exists a canonical isomorphism of k-algebras

$$\mathrm{M}_{\mathfrak{I}} \simeq \mathrm{End}_{\mathfrak{C}}(\underset{i \in \mathfrak{I}}{\oplus} C_i),$$

where End_C denotes the endomorphisms ring in the category C.

(2) Let $C_{\mathfrak{I}}$ be the restriction the category C on \mathfrak{I} , $M_{\mathfrak{I}}-\mathrm{mod}_{r}$ a full subcategory in $M_{\mathfrak{I}}-\mathrm{mod}$ consisting on modules M, such that $M=\bigoplus_{i\in \mathfrak{I}}e_{ii}M$, where e_{ii} where e_{ii} is a matrix unit corresponding $i\in \mathfrak{I}$. Then there exists the canonical equivalence

$$F: \mathcal{A}_{\mathcal{I}} - \text{Mod} \simeq M_{\mathcal{I}} - \text{mod}_r,$$

where for
$$N \in \mathcal{A}_{\mathfrak{I}}$$
 – Mod holds $F(N) = \bigoplus_{i \in \mathfrak{I}} N(i)$.

Proof. Every element $(f_{ij} \mid i, j \in \mathcal{I}) \in \Pi_{\mathcal{I}}$ defines canonically a homomorphism $f: \underset{i \in \mathcal{I}}{\oplus} C_i \longrightarrow \prod_{i \in \mathcal{I}} C_i$. By the condition (*) the image of f belongs to $\underset{i \in \mathcal{I}}{\oplus} C_i \subset \prod_{i \in \mathcal{I}} C_i$. The second statement is standard.

The following statement is an analogue of two-sided Pierce decomposition for the pair $\Gamma \subset U$.

Theorem 3.1. Denote for $u \in U$ by [u] the matrix from M_A , such that $[u]_{m,n} = 0$ $(u_{n,m}), m, n \in \operatorname{Specm} \Gamma, \text{ where } u_{n,m} \text{ is the image of } u \text{ in } \mathcal{A}(m,n).$

- (1) The mapping $i_{\mathcal{A}} = []: U \longrightarrow M_{\mathcal{A}}$, which sends $u \in U$ to [u] is a homomorphism of \mathbb{k} -algebras.
- (2) Endow Π_A with the topology of direct product and $M_D \subset \Pi_A$ with the induced topology. Then the image of $\pi_D i_A$ is dense in M_D .

Proof. Following Lemma 3.3, (3), (4), the matrix [u] has finitely many nonzero elements in any row and any column, hence $[u] \in \mathcal{M}_{\mathcal{A}}$.

Fix $u, v \in U$, $m, n \in \operatorname{Specm} \Gamma$, $m, n \geq 1$. Also fix V, W, T and P > 0, satisfying the conditions of Lemma 3.4. Then from

$$\widehat{\mu}(\mathsf{m}^m,\mathsf{n}^n) = \sum_{\mathsf{p} \in X^\mathsf{n} \cap X_\mathsf{m}} \widehat{\mu}_\mathsf{p}(\mathsf{m}^m,\mathsf{n}^n)$$

follows (as in the proof of Lemma 3.4, (5))

$$\overline{uv} = \sum_{\mathbf{p} \in X^{\mathbf{n}} \cap X_{\mathbf{m}}} \widehat{\mu}_{\mathbf{p}}(\mathbf{m}^m, \mathbf{n}^n) (\bar{v}_{\mathbf{p}} \otimes_{\mathbf{p}} \bar{w}),$$

where \overline{uv} is a class of $uv \in {\bf n}^n T_{{\bf m}^m}$. Taking the limits we obtain that

$$[uv]_{\mathsf{m},\mathsf{n}} = \sum_{\mathsf{p} \in X^\mathsf{n} \cap X_\mathsf{m}} [u]_{\mathsf{n},\mathsf{p}}[v]_{\mathsf{p},\mathsf{m}},$$

that proves the first statement.

To prove the second statement it is enough to show that if $\mathcal{A}(m,n) \subset M_D$, then $\mathcal{A}(\mathsf{m},\mathsf{n}) \subset \overline{\mathrm{Im}\,\pi_D i_{\mathcal{A}}}$. First note that the image of U is dense in the image of $\mathcal{A}(\mathsf{m},\mathsf{n}) \subset \mathrm{M}_D$, i.e. matrices from M_D , which are zero in all positions except (n, m). Let $u_i \in U \mid i = 1, \ldots$ be a sequence in U, which converges to an element f from $\mathcal{A}(m,n)$. Note, that any D is by definition at most countable. Consider an increasing sequence of finite subsets $S_i \subset D$, $i = 1, \ldots$ such that $\bigcup_{i=1}^{\infty} S_i = D$, strictly increasing sequence of integers $k_i > 0$ and the elements $\mu_i, \nu_i \in \Gamma$, such that

$$\left\{ \begin{array}{ll} \mu_i & \equiv & 1 \mod & \mathsf{m}^{k_i} \\ \mu_i & \equiv & 0 \mod & \mathsf{m}'^{k_i} \mod & \mathsf{m}' \in S_i, \mathsf{m}' \neq \mathsf{m}. \\ \\ \left\{ \begin{array}{ll} \nu_i & \equiv & 1 \mod & \mathsf{n}^{k_i} \\ \nu_i & \equiv & 0 \mod & \mathsf{n}'^{k_i} \mod & \mathsf{n}' \in S_i, \mathsf{n}' \neq \mathsf{n}. \end{array} \right.$$

and

$$\begin{cases} \nu_i \equiv 1 \mod & \mathsf{n}^{k_i} \\ \nu_i \equiv 0 \mod & \mathsf{n}'^{k_i} \mod \\ \end{cases}$$

Then $i_D(\mu_i)$ (correspondingly $i_D(\nu_i)$) tends in M_D to the diagonal matrix unit in the position m (correspondingly n),hence the sequence $i_D(\nu_i u_i \mu_i) = i_D(u_i) i_D(\mu_i)$ converges to f since it tends to 0 in all positions except (n, m) and tends to f in the position (n, m).

3.3.3. Category equivalence for Gelfand-Tsetlin modules. We consider the category of \mathbb{k} – Mod endowed with the discrete topology and consider the category \mathcal{A} – Mod_d of continuous functors $M: \mathcal{A} \longrightarrow \mathbb{k} - \text{Mod}$, which in [DFO2] are called discrete modules. It means, for every $m \in \operatorname{Specm} \Gamma$ there exists $m = m(m) \geq 0$, such that $\mathsf{m}^m M(\mathsf{m}) = 0$. For a discrete \mathcal{A} -module M define a Gelfand-Tsetlin U-module

 $\mathbb{F}(M)$ by setting

(18)
$$\mathbb{F}(M) = \bigoplus_{\mathsf{m} \in \operatorname{Specm} \Gamma} M(\mathsf{m}) \text{ and for } x \in M(\mathsf{m}), a \in U$$

$$\operatorname{set} \, ax = \sum_{\mathsf{n} \in \operatorname{Specm} \Gamma} a_{\mathsf{m},\mathsf{n}} x,$$

where $a_{\mathsf{m},\mathsf{n}}$ is the image of a in $\mathcal{A}(\mathsf{m},\mathsf{n})$. If $f: M \longrightarrow N$ is a morphism in $\mathcal{A} - \mathrm{mod}_d$ then define $\mathbb{F}(f) = \bigoplus_{\mathsf{m} \in \mathrm{Specm}\,\Gamma} f(\mathsf{m})$.

Theorem 3.2. (DFO2), Theorem 17) The defined above \mathbb{F} is an equivalence

$$\mathbb{F}: \mathcal{A} - \text{mod}_d \longrightarrow \mathbb{H}(U, \Gamma),$$

which induces a functorial isomorphism

$$\operatorname{Ext}^1_U(X,Y) \simeq \operatorname{Ext}^1_{\mathcal{A}}(\mathbb{F}(X),\mathbb{F}(Y)), \ X,Y \in \mathbb{H}W(U,\Gamma).$$

Proof. Following Lemma 3.5, (2) there exists a canonical equivalence

$$\mathcal{A} - \text{mod}_d \simeq M_{\mathcal{A}} - \text{mod}_r$$

that together with the functor, induced by $i_{\mathcal{A}}: U \hookrightarrow \mathcal{M}_{\mathcal{A}}$ (Theorem 3.1, (1)) gives us the functor $\mathcal{A} - \text{mod}_d \longrightarrow U - \text{mod}$, besides the image of this functor belongs to $\mathbb{H}(U,\Gamma)$.

Let $u \in U$, $m, n \in \operatorname{Specm} \Gamma$, $u_{m,n}$ the image of u in $\mathcal{A}(m, n)$. Let

$$M = \sum_{\mathsf{m} \in \operatorname{Specm} \Gamma} M(\mathsf{m})$$

be a Gelfand-Tsetlin module. The action of $u_{\mathsf{m},\mathsf{n}}$ on M is described in the next lemma.

Lemma 3.6. Let $m_u: M(\mathsf{m}) \longrightarrow M$ be the map $x \longmapsto ux, \ x \in M(\mathsf{m})$. Then $u_{\mathsf{m,n}}: M(\mathsf{m}) \to M(\mathsf{n}), \ u_{\mathsf{m,n}} = \pi_{\mathsf{n}} m_u, \ where \ \pi_{\mathsf{n}} \ is \ the \ canonical \ projection \ of \ M$ onto M_{n} . Besides, if $\mathsf{n} \not\in X_u$, then $u_{\mathsf{m,n}} = 0$.

Proof. Condition $n \notin X_u(m)$ holds if and only if $\Gamma u\Gamma = nu\Gamma + \Gamma um$, equivalently, $u \in nu\Gamma + \Gamma um$. But then $u \in n^n u\Gamma + \Gamma um^m$ for every m, n > 0 and hence $u_{m,n} = 0$. On other hand in this case $\pi_n m_u = 0$ by (4). By the Chinese remainder theorem, there exists a sequence $\gamma_n \in \Gamma, n \geq 1$, such that $\gamma_n \equiv 1 \pmod{n^n}$, $\gamma_n \equiv 0 \pmod{l^n}$ for any $l \in X_u(m), l \neq n$. Then in A(m, n) holds $\lim_{n \to \infty} \gamma_n u = u_{m,n}$. On the other hand the operator $\bar{m}_u = \lim_{n \to \infty} m_{\gamma_n u} : M(m) \to M$ is well defined and $\pi_n m_u = \pi_m \bar{m}_u$ and $\pi_{n'} m_u = 0$ for any $n' \neq n$. Then (18) completes the proof. \square

We will identify a discrete \mathcal{A} -module N with the corresponding Gelfand-Tsetlin module $\mathbb{F}(N)$. Let $\Gamma_{\mathsf{m}} = \lim_{\leftarrow m} \Gamma/\mathsf{m}^m$ be the completion of Γ by $\mathsf{m} \in \operatorname{Specm} \Gamma$. For $\mathsf{m} \in \operatorname{Specm} \Gamma$ denote by $\hat{\mathsf{m}}$ a completion of m . Consider a two-sided ideal $I \subset \mathcal{A}$ generated by $\hat{\mathsf{m}}$ for all $\mathsf{m} \in \operatorname{Specm} \Gamma$ and set $\mathcal{A}(W) = \mathcal{A}/I$. Then Theorem 3.2 implies the following statement.

Corollary 3.2. The categories $\mathbb{H}W(U,\Gamma)$ and $\mathcal{A}(W) - \text{mod}_d$ are equivalent.

The following standard statement shows usefulness of the category \mathcal{A} for calculation simple Gelfand-Tsetlin modules over U.

Lemma 3.7. Let in assumption above m, M simple A module, $M(m) \neq 0$, N simple A(m,m)-module. Then the correspondences

$$M \longmapsto M|_{\mathcal{A}(\mathsf{m},\mathsf{m})} \ and \ N \longmapsto \mathcal{A} \otimes_{\mathcal{A}(\mathsf{m},\mathsf{m})} N/J,$$

where J is a unique maximal A-submodule in the induced module, realizes a bijection between the sets of isoclasses of simple A(m,m)-modules and simple A-modules M such that $M(m) \neq 0$.

The subalgebra Γ is called big in $m \in \operatorname{Specm} \Gamma$ if $\mathcal{A}(m,m)$ is finitely generated as a right Γ_m —module. The importance of this concept is described in the following statement.

Lemma 3.8. ([DFO2], Corollary 19) If Γ is big in $m \in \operatorname{Specm} \Gamma$ then there exists finitely many non-isomorphic irreducible Gelfand-Tsetlin $U-modules\ M$ such that $M(m) \neq 0$. For any such module M holds $\dim_{\mathbb{R}} M(m) < \infty$.

4. Generic representations

Assume \mathbbm{k} is uncountable. A non-empty set $X \subset \operatorname{Specm} \Gamma$ is called *massive*, provided that X contains an intersection of countable many dense open subsets. In this case X is dense in $\operatorname{Specm} \Gamma$.

Lemma 4.1. Let M be a finitely generated right Γ -module. Then the set of $\mathsf{m} \in \operatorname{Specm} \Gamma$ such that $\operatorname{Tor}_1^{\Gamma}(M, \Gamma/\mathsf{m}) = 0$ contains an open dense subset $X \subset \operatorname{Specm} \Gamma$.

Proof. Let $R^{\bullet}: \ldots \xrightarrow{d^2} \Gamma^{n_2} \xrightarrow{d^1} \Gamma^{n_1} \xrightarrow{d^0} \Gamma^{n_0} \longrightarrow 0 \ldots$ be a free resolution of M. It induces the resolution $R^{\bullet} \otimes_{\Gamma} K$ of $M \otimes_{\Gamma} K$. Denote $r = \dim_K \operatorname{Im}(d^1 \otimes \operatorname{id}_K)$. Let $d^i(\mathsf{m})$ be the specialization of d^i in $\mathsf{m} \in \operatorname{Specm} \Gamma$. Then m satisfying the conditions $\dim \operatorname{Im} d^1(\mathsf{m}) = n_1 - r$ and $\dim \operatorname{Im} d^2(\mathsf{m}) = r$ form an open dense set X. Then for $\mathsf{m} \in X$ the first cohomology of $R^{\bullet} \otimes_{\Gamma} \Gamma/\mathsf{m}$ equals 0 that completes the proof.

Proposition 4.1. For a nonzero $u \in U, u \neq 0$ there exists a massive set $\Omega_u \subset \operatorname{Specm} \Gamma$ such that for every $m \in \Omega_u$ the image \bar{u} of u in U/Um is nonzero.

Proof. Let $m \in \operatorname{Specm} \Gamma$, $N = u\Gamma$. Then $\bar{u} = 0$ if and only if $u = \sum_{i=1}^{n} u_i m_i$, for some

 $u_i \in U, m_i \in \mathsf{m}, \ i = 1, \ldots, n.$ Denote $S = \bigcup_{i=1}^n \operatorname{supp} u_i$ and M = U(S). If $\bar{u} = 0$, then the exact sequence of right Γ -modules

$$0 \longrightarrow N \longrightarrow M \longrightarrow M/N \longrightarrow 0$$

becomes non-exact after tensoring with Γ/m , i.e. $\operatorname{Tor}_1^{\Gamma}(M/N, \Gamma/m) \neq 0$. Denote

$$X(u,S) = \{\mathsf{m} \in \operatorname{Specm} \Gamma \mid \operatorname{Tor}_1^{\Gamma}(M/N,\Gamma/\mathsf{m}) = 0.$$

Following Lemma 4.1 we can set
$$\Omega_u = \bigcap_{S \subset \mathcal{M}} X(u, S)$$
.

Here we show, that for any Galois ring there exists a "large" set of $m \in \operatorname{Specm} \Gamma$, such that the character $\chi = \chi_m$ extends to a simple U-module.

Let $U_{\chi} = U/(U\mathsf{m}_{\chi})$. We call U_{χ} the universal module generated by a χ -eigenvector. If Γ is a Harish-Chandra subalgebra then $U_{\chi} \in \mathbb{H}(U,\Gamma)$. In general U_{χ} could be zero.

Let Γ_0 be the algebra generated by $\bigcup_{m\in\mathcal{M}} m(\Gamma)$ and $\bar{\Gamma}_0$ its integral closure in L (if $\Gamma_0 \subset \bar{\Gamma}$ then U is a right Galois order), $\mathcal{L} = \operatorname{Specm} \bar{\Gamma}_0$, $\Omega = \operatorname{Specm} \Gamma$. The embedding $\Gamma \hookrightarrow \bar{\Gamma}_0$ induces the mapping $\pi_0 : \mathcal{L} \longrightarrow \Omega$. The group G acts on \mathcal{L} and the orbits of this action are in the canonical bijection with $\operatorname{Specm} \Gamma_S$ for some

multiplicative set S. Since $\Gamma \subset \overline{\Gamma}_0$ we can write $\gamma(\ell)$ instead of $\gamma(\pi_0(\ell))$ for $\gamma \in \Gamma$ and $\ell \in \mathcal{L}$. If $\varphi \in \operatorname{Aut} \overline{\Gamma}_0$ and $\gamma \in \Gamma$, then $\gamma^{\varphi}(\ell) = \gamma(\varphi^{-1} \cdot \ell)$.

Let $A_{\mathcal{M}}$ be the set of all a^m , where $m \in \mathcal{M}$ and $[a\varphi]$ runs all standard generators of U. Denote by Λ_1 an algebra generated over $\bar{\Gamma}_0$ by $A_{\mathcal{M}}$, and let Λ_2 be an algebra generated over Λ_1 by all a^{-1} , $a \in A_{\mathcal{M}}$. Denote $\mathcal{L}_i = \operatorname{Specm} \Lambda_i$, $\Omega_i = \pi_0(\mathcal{L}_i)$, i = 1, 2. Then

$$\mathcal{L}_2 \subset \mathcal{L}_1 \subset \mathcal{L}, \quad \Omega_2 \subset \Omega_1 \subset \Omega.$$

By $\mathcal{L}_r \subset \mathcal{L}$ denote the set of ℓ , such that \mathcal{M} acts on ℓ freely and $\mathcal{M} \cdot \ell \cap G \cdot \ell$ consists just of ℓ (in terms of 5.1 for $\mathsf{m} = \pi_0(\ell) \in \operatorname{Specm} \Gamma$ holds $S(\mathsf{m}, \mathsf{m}) = \{e\}$) and $\Omega_r = \pi_0(\mathcal{L}_r)$).

The following standard fact is obvious.

Lemma 4.2. Let $\pi: \mathcal{L} \to \Omega$ be the canonical projection. If $X \subset \mathcal{L}$ is a massive subset then $\pi_0(X)$ is massive in Ω .

If $\mathcal{M} \cdot \Gamma \subset \overline{\Gamma}$ then any $m \in \mathcal{M}$ defines an automorphism of $\overline{\Gamma}_0 = \overline{\Gamma}$ and, hence induces an automorphism of \mathcal{L} . In particular, this holds when Γ is a Harish-Chandra subalgebra.

Lemma 4.3. Assume \mathbb{k} is uncountable. Then the sets $\mathcal{L}_i \subset \mathcal{L}$ and $\Omega_i \subset \operatorname{Specm} \Gamma$, i = 1, 2, are massive. Moreover, if $\mathbb{M} \cdot \Gamma \subset \overline{\Gamma}$ then \mathcal{L}_r and Ω_r are massive.

Proof. Every element $a \in L$ defines a rational function on \mathcal{L} with a non-empty open domain of definition $X(a) \subset \mathcal{L}$. Then $\mathcal{L}_i, i = 1, 2$ by definition are intersections of countably many sets of the form X(a), hence are massive.

For any $m \in \mathcal{M}, m \neq e$, denote $\mathfrak{X}_m = \{\ell \in \mathcal{L} \mid m \cdot \ell \in G \cdot \ell\}$ and for $m \in \mathcal{M}, g \in G$ define a closed in \mathcal{L} subset $\mathcal{L}(m,g) = \{\ell \in \mathcal{L} \mid m \cdot \ell = g \cdot \ell\}$. Then $\mathfrak{X}_m = \bigcup_{g \in G} \mathcal{L}(m,g)$ is closed subset in \mathcal{L} , since G is finite. Assume that $\mathcal{L} = \mathfrak{X}_m$ for some $m \neq e$. Since \mathcal{L} is irreducible, we conclude that $\mathcal{L}(m,g) = \mathcal{L}$ for some $g \in G$, and hence m = g. This is a contradiction since \mathcal{M} is separating. But $\bigcup_{m \in \mathcal{M}, m \neq e} \mathfrak{X}_m$ is the complement of Ω_r in \mathcal{L} and \mathcal{L}_r is massive. The sets Ω_i , i = 1, 2, r, are massive by Lemma 4.2.

The following useful fact is obvious.

Lemma 4.4. Let $\bar{\Gamma}$ be the integral closure on Γ in L, $\ell_{\mathsf{m}_k}, \ldots, \ell_{\mathsf{m}_k} \in \operatorname{Specm} \bar{\Gamma}$ belong to different orbits of G. Then there exists $\gamma \in \Gamma$, such that $\gamma(\ell_{\mathsf{m}_1}), \ldots, \gamma(\ell_{\mathsf{m}_k})$ in other words Γ distinguishes the orbits of G.

Theorem 4.1. Suppose that the field k is uncountable.

- (1) There exists a massive subset $X_1 \subset \operatorname{Specm} \Gamma$, such that for every $\chi \in X_1$, U_{χ} is nonzero Gelfand-Tsetlin module and $\sup U_{\chi} \subset \mathbb{O}_{\mathsf{m}}$, where $\chi = \chi_{\mathsf{m}}$ and $\mathbb{O}_{\mathsf{m}} = \mathbb{M} \cdot \mathsf{m}$.
- (2) If M is a group, then there exists a massive set $X_2 \subset X_1$, such that for any $\chi \in X_2$ the module U_{χ} is a unique $\mathfrak{O}_{\mathsf{m}}$ -graded irreducible U-module generated by a χ -eigenvector and $\sup U_{\chi} = \mathfrak{O}_{\mathsf{m}}$.
- (3) If M is a group and $M \cdot \Gamma \subset \overline{\Gamma}$, then there exists a massive set $X_r \subset X_2$, such that for any $\chi \in X_r$, U_{χ} is irreducible weight U-module with 1-dimensional components. In this case there is a canonical isomorphism of \mathbb{R} -vector spaces $\mathbb{R}M \simeq U_{\chi}$.

Proof. Note that a skew semigroup algebra $\Lambda_1 * \mathcal{M} \subset L * \mathcal{M}$ contains the Galois ring U. A character $\tilde{\chi}: \Lambda_1 \to \mathbb{k}$ induces a $\Lambda_1 * \mathcal{M}$ -module $M(\tilde{\chi})$ which is isomorphic to $\underset{\varphi \in \mathcal{M}}{\oplus} \Lambda_1/(\tilde{\mathsf{m}}^{\varphi})$, as a \mathbb{k} -vector space, where $\tilde{\mathsf{m}} = \operatorname{Ker} \tilde{\chi}$. Let $X_1 = \Omega_1$. Since the restriction of $M(\tilde{\chi})$ on U is nontrivial then for every $\chi \in X_1$, $U_{\chi} \neq 0$. Clearly,

 U_{χ} is a Gelfand-Tsetlin module. Moreover, since U_{χ} is Specm \mathcal{L} -graded, it has an irreducible quotient with a nonzero χ -eigenvector. This implies (1). Assume now that \mathcal{M} is a group and set $X_2 = \Omega_2$. Let $\chi \in X_2$, $\mathsf{m} = \operatorname{Ker} \chi$ and $\sum_{h \in \mathcal{M}} [x_h h]$ a

generator of U. By assumption, for every $n \in \mathcal{M} \cdot m$ holds $x_h(n) \neq 0$, hence every graded component of U_χ generates the whole U_χ . Therefore, U_χ is irreducible as \mathcal{O}_m -graded U-module. Moreover, since U_χ is the universal module generated by a χ -eigenvector, it is a unique such graded irreducible module, implying statement (2). Note that if \mathcal{M} acts on m with a nontrivial stabilizer then U_χ is not irreducible. Suppose now that $\mathcal{M} \cdot \Gamma \subset \bar{\Gamma}$. Then Ω_r is massive by Lemma 4.3. Consider a subset $X_r = X_2 \cap \Omega_r$. Since Γ distinguishes the components by Lemma 4.4, it implies the irreducibility of U_χ for any $\chi \in X_r$. The basis elements of U_χ in this case are labeled by the elements of $\mathbb{k}\mathcal{M}$ which completes the proof of (3).

5. Representations of Galois orders

Given $\mathsf{m} \in \operatorname{Specm} \Gamma$ fix $\ell_{\mathsf{m}} \in \operatorname{Specm} \overline{\Gamma}$ and denote $\mathfrak{M}_{\ell_{\mathsf{m}}} \subset \mathfrak{M}$ $(G_{\ell_{\mathsf{m}}} \subset G)$ the stabilizer of ℓ_{m} in \mathfrak{M} (in G). The ideal m defines $\mathfrak{M}_{\ell_{\mathsf{m}}}$ and $G_{\ell_{\mathsf{m}}}$ uniquely up to G-conjugation. It allows us to use the notation $\mathfrak{M}_{\mathsf{m}}$ instead of $\mathfrak{M}_{\ell_{\mathsf{m}}}$ and G_{m} instead of $G_{\ell_{\mathsf{m}}}$.

5.1. Extension of characters for Galois orders. The property of the Galois ring U to be a Galois order has the following immediate impact on the representation theory of U. We will consider right Galois orders. The case of left orders is analogous.

Let U be a Galoir ring over Γ , Γ a noetherian algebra which is a right Harish-Chandra subalgebra of U, $\mathsf{m} \in \operatorname{Specm} \Gamma$. Denote $S = \mathcal{M}_{\mathsf{m}}$.

Lemma 5.1. Assume $U = U \mathsf{m}$ holds. Then for every $k \geq 1$ there exist $\gamma_k \in \Gamma \setminus \mathsf{m}$, $v_j \in U$, $\nu_j \in \mathsf{m}^k$, $j = 1, \ldots, N$ such that

(19)
$$\gamma_k = \sum_{j=1}^N v_j \nu_j$$

and supp $v_j \in S$, j = 1, ..., n.

Proof. The condition $U = U \mathbf{m}$ is equivalent to the condition $1 \in U \mathbf{m}$, i.e. holds the equality

(20)
$$1 = \sum_{i=1}^{n} u_i \mu_i, \text{ where } u_i \in U, \mu_i \in \mathsf{m}.$$

We use induction in k to prove the statement of the lemma without the condition supp $v_i \in S$, i = 1, ..., n. The base of induction k = 1 follows immediately from (20). The induction step is obtained by substitution of the right side of (20) in (19): if

(21)
$$1 = \sum_{j=1}^{N} w_j \nu_j, \text{ where } w_j \in U, \nu_j \in \mathsf{m}^k, j = 1, \dots, N,$$

then

$$1 = \sum_{j=1}^{N} w_j \nu_j = \sum_{j=1}^{N} w_j \cdot 1 \cdot \nu_j =$$
$$\sum_{j=1}^{N} w_j \left(\sum_{i=1}^{n} u_i \mu_i \right) \nu_j = \sum_{j=1}^{N} \sum_{i=1}^{n} w_j u_i (\mu_i \nu_j).$$

It proves the induction step, since all $\mu_i \nu_j \in \mathsf{m}^{k+1}$. Denote

$$T = \bigcup_{i=1}^{N} \operatorname{supp} w_i \setminus S.$$

Since $T \cap \mathcal{M}_{\mathsf{m}} = \emptyset$, the ideals ℓ_{m}^t and ℓ_{m} for every $t \in T$ belong to different G-orbits. By Lemma 4.4 there exists $f \in \Gamma$ such that $f(\ell_{\mathsf{m}}) \neq f(\ell_{\mathsf{m}}^t)$ for every $t \in T$. Without loss of generality we can assume that (Lemma 2.4, (3))

$$f_T(\mathsf{m},\mathsf{m}) = \prod_{t \in T} (f(\ell_\mathsf{m}) - f^{t^{-1}}(\ell_\mathsf{m})) = \prod_{t \in T} (f(\ell_\mathsf{m}) - f(\ell_\mathsf{m}^t)) = 1.$$

In particular, it implies that $f_T \in 1 + m \otimes \Gamma + \Gamma \otimes m$.

Then by Lemma 2.4, (3), $\gamma_k = f_T \cdot 1 \in 1 + m$. Besides, by Lemma 2.4, (2), $v_i = f_T \cdot w_i$ satisfies $v_i \in U(S)$. Applying f_T to the equality (21) we obtain (19),

$$\gamma_k = \sum_{i=1}^N v_i \nu_i$$
, where $\gamma_k \in \Gamma \setminus \mathsf{m}, \, v_i \in U(S), \nu_i \in \mathsf{m}^k$.

Corollary 5.1. Let Γ be a noetherian algebra, U a right Galois order over Γ , $m \in \operatorname{Specm} \Gamma$, such that $|\mathcal{M}_m| < \infty$. Then $Um \neq U$. In particular, there exists a left simple U-module M, generated by $x \in M$, such that $m \cdot x = 0$.

Proof. The set $S = \mathcal{M}_m$ is finite and the algebra U is an Galois order, hence by Theorem 2.2, (2), U(S) is finitely generated as a right Γ-module. Then applying the Artin-Rees Lemma (Theorem 8.5. [Mat]) for a right Γ- module U(S) and its submodule Γ we conclude that there exists $c \geq 0$ such that for every $k \geq c$

$$U(s)\mathsf{m}^k\cap\Gamma=(U(S)\mathsf{m}^c\cap\Gamma)\mathsf{m}^{k-c}.$$

But for every k > c there exists (by Lemma 5.1) $\gamma_k \in U(S) \mathfrak{m}^k \cap \Gamma$, such that $\gamma_k \not\in \mathfrak{m}$, hence $\gamma_k \not\in (U(S) \mathfrak{m}^c \cap \Gamma) \mathfrak{m}^{k-c}$, since k-c > 0. The obtained contradiction shows that $U \neq U\mathfrak{m}$.

Denote by v the image of 1 in $U/U m \neq 0$. Then mv = 0 which defines a structure of Gelfand-Tsetlin module on U/U m. Any nonzero simple quotient of U/U m satisfies the statement. Therefore, as the module M we can choose any simple quotient of U/U m and set x = v.

Corollary 5.1 implies **Theorem A**,(1) stated in the Introduction. Of course, the property of a Galois ring to be right order is not a necessary condition to guarantee an extension of an arbitrary character of Γ to a U-module.

The following statement is probably well known but we include the proof for the convenience of the reader. Note that we consider Proposition 5.1 as a motivation for introducing the notion of a Galois order.

Proposition 5.1. Let $i: A \hookrightarrow B$ be an embedding of integral domains with a regular A. Assume the induced morphism of varieties $i^*: \operatorname{Specm} B \to \operatorname{Specm} A$ is surjective (e.g. $A \subset B$ is an integral extension). If $b \in B$ and $ab \in A$ for some nonzero $a \in A$ then $b \in A$.

Proof. Under the assumption the map i induces an epimorphism of Spec B onto Spec A. Fix $m \in \operatorname{Specm} A$ and let $ab = a' \in A$. Since the localization A_m is a unique factorization domain, we can assume that a and a' are coprime in A_m . If a is invertible in A_m then $b \in A_m$. If a is not invertible in A_m then there exists $P \in \operatorname{Spec} A_m$ such that $a \in P$ and $a' \notin P$. It shows that P does not lift to Spec B_m . Hence $b \in A_m$ for every $m \in \operatorname{Specm} A$, holds $b \in A$ ([Mat], Theorem 4.7).

Proposition 5.1 gives the converse statement to Theorem A.

Lemma 5.2. Let $U \subset L*M$ be a Galois ring over a noetherian Γ . If every character $\chi: \Gamma \longrightarrow \mathbb{k}$ extends to a representation of U then $U_e \subset \overline{\Gamma} \cap K$. If in addition M is a group and Γ is a Harish-Chandra subalgebra then U is a Galois order.

Proof. If χ extends to a representation of U, then it extends to a representation of $U_e \subset K$ in particular. Proposition 5.1 implies that U_e belongs to the integral closure of Γ in K. The second statement follows immediately from Theorem 2.3.

Hence we obtain the following module-theoretic characterization of left and right Galois orders.

Corollary 5.2. Let U be a Galois ring with respect to a noetherian algebra Γ , \mathbb{M} a group and $m^{-1}(\Gamma) \subset \overline{\Gamma}$ ($m(\Gamma) \subset \overline{\Gamma}$) for any $m \in \mathbb{M}$. Then every character $\chi : \Gamma \to \mathbb{k}$ extends to a simple left (right) U-module if and only if U is right (left) Galois order.

- 5.2. Gelfand-Tsetlin modules over Galois orders. In this subsection we assume that U is a Galois order over Γ where Γ is normal noetherian over \Bbbk . In particular, $\Gamma = \tilde{\Gamma} = U_e$ and $\bar{\Gamma}$ is finite over Γ by Corollary 2.1. Also Γ is a Harish-Chandra subalgebra by Proposition 2.1. If $\ell \in \operatorname{Specm} \bar{\Gamma}$ projects to $\mathsf{m} \in \operatorname{Specm} \Gamma$ then we will write $\ell = \ell_\mathsf{m}$ and say that ℓ_m is lying over m . Let ℓ_m and ℓ_n be some maximal ideals of $\bar{\Gamma}$ lying over m and n respectively. Note that given $\mathsf{m} \in \operatorname{Specm} \Gamma$ the number of different ℓ_m is finite due to Corollary 2.2.
- 5.2.1. Finiteness statements for the category \mathcal{A} . Denote by $S(\mathsf{m},\mathsf{n})$ the following G-invariant subset in \mathcal{M}

(22)
$$S(\mathbf{m}, \mathbf{n}) = \{ m \in \mathcal{M} \mid \ell_{\mathbf{n}} \in GmG \cdot \ell_{\mathbf{m}} \} = \{ m \in \mathcal{M} \mid g_{2}\ell_{\mathbf{n}}$$
$$= mq_{1}\ell_{\mathbf{m}} \text{ for some } q_{1}, q_{2} \in G \}.$$

Obviously this definition does not depend on the choice of ℓ_m and ℓ_n .

Lemma 5.3. Let $m \in \operatorname{Specm} \Gamma$ and M is a group. Then

(1) $|\mathcal{M}_m| < \infty$ if and only if for any $n \in \operatorname{Specm} \Gamma$ both $\mathit{S}(m,n)$ and $\mathit{S}(n,m)$ are finite. In this case

$$|S(\mathsf{m},\mathsf{n})| = |S(\mathsf{n},\mathsf{m})|.$$

(2) If $|\mathcal{M}_{\mathsf{m}}| < \infty$ and $S(\mathsf{m},\mathsf{n}) \neq \emptyset$, $S(\mathsf{n},\mathsf{m}) \neq \emptyset$ are nonempty, then $|\mathcal{M}_{\mathsf{n}}| < \infty$

$$|S(\mathsf{m},\mathsf{n})| \le \frac{|G|^2 |\mathcal{M}_\mathsf{m}|}{|G_\mathsf{m}||G_\mathsf{n}|}.$$

Proof. $S(\mathsf{m},\mathsf{n})$ is infinite if and only if there exists infinitely many $m \in \mathcal{M}$, such that $m^{-1}g\ell_{\mathsf{n}} = \ell_{\mathsf{m}}$ for some $g \in G$. Since the group \mathcal{M}_{m} is infinite. To prove the equality note, that $m \in S(\mathsf{m},\mathsf{n})$ if and only if $m^{-1} \in S(\mathsf{n},\mathsf{m})$. It proves (1).

If $|\mathcal{M}_{\mathsf{m}}| < \infty$ then there exists at most $|\mathcal{M}_{\mathsf{m}}|$ elements $\varphi \in \mathcal{M}$, such that $\ell_{\mathsf{n}} = \varphi \ell_{\mathsf{m}}$. Considering the G-orbits of ℓ_{m} and ℓ_{n} , which have the lengths $\frac{|G|}{|G_{\mathsf{m}}|}$ and $\frac{|G|}{|G_{\mathsf{n}}|}$ respectively, we obtain the statement (2).

Lemma 5.4. Let $\mathsf{m}, \mathsf{n} \in \operatorname{Specm} \Gamma$, $S = S(\mathsf{m}, \mathsf{n})$, $m, n \geq 0$. Then $U = U(S) + \mathsf{n}^n U + U \mathsf{m}^m$. Moreover, for every $u \in U, k \geq 0$ there exists $u_k \in U(S)$, such that $u \in u_k + \mathsf{n}^{\lfloor k/2 \rfloor} u \Gamma + \Gamma u \mathsf{m}^{\lfloor k/2 \rfloor}$.

Proof. Fix $u \in U$ and denote $T = \text{supp } u \setminus S$. If $T = \emptyset$ then $u \in U(S)$. Let $T \neq \emptyset$. We show by induction in k, that there exists $u_k \in U(S)$, such that

(24)
$$u \in u_k + \sum_{i=0}^k \mathsf{n}^{k-i} u \mathsf{m}^i, \ u_k \in U(S) \ (\text{ hence } u_{k+1} - u_k \in \sum_{i=0}^k \mathsf{n}^{k-i} u \mathsf{m}^i).$$

Since ℓ_{m}^t and ℓ_{n} belong to different G-orbits if $t \notin S$, then by Lemma 4.4 there exists $f \in \Gamma$ such that $f(\ell_{\mathsf{n}}) \neq f(\ell_{\mathsf{m}}^t)$ for every $t \in T$. Without loss of generality we can assume that $f_T(\mathsf{n},\mathsf{m}) = \prod_{t \in T} (f(\ell_{\mathsf{n}}) - f^{t^{-1}}(\ell_{\mathsf{m}})) = 1$, which implies $f_T \in \mathcal{T}$

 $1+\mathsf{n}\otimes\Gamma+\Gamma\otimes\mathsf{m}$. Set $u_1=f_T\cdot u$. Then u_1 belongs to $u+\mathsf{n}u\Gamma+\Gamma u\mathsf{m}$ and, hence, $u\in u_1+\mathsf{n}u\Gamma+\Gamma u\mathsf{m}$. Moreover, $u_1\in U(S)$ by Lemma 2.4, (2). We prove the induction step $k\Rightarrow k+1$. Writing in (24) the expression for u in the right hand side we obtain

$$u \in u_k + \sum_{i=0}^k \mathsf{n}^{k-i} (u_k + \sum_{j=0}^k \mathsf{n}^{k-j} u \mathsf{m}^j) \mathsf{m}^i \subset u_k + \sum_{i=0}^k \mathsf{n}^{k-i} u_k \mathsf{m}^i + \sum_{i=0}^{k+1} \mathsf{n}^{k+1-i} u \mathsf{m}^i,$$

that proofs of the induction step, since $u_k + \sum_{i=0}^k \mathsf{n}^{k-i} u_k \mathsf{m}^i \subset U(S)$.

Remark 5.1. We use notations from Lemma 3.1. If $\mathcal{A}(m,n)$ is finitely generated as Γ -bimodule then the category \mathcal{A} can be equivalently defined as follows ([DFO2], 1.4). Let $I, m, n \in \operatorname{Specm} \Gamma$. To define the composition

$$\mathcal{A}(n, I) \times \mathcal{A}(m, n) \longrightarrow \mathcal{A}(m, I)$$

considered pairs $a, b \in U$ and m, n > 0, the left Γ -module $M = L_{a,m^m}$ and the right Γ -module $N = R_{b,l^l}$. By the Harish-Chandra property both modules are finite dimensional over k. As above M and N as Γ -modules decompose

(25)
$$M = M_0 \oplus M_1$$
, where $n^n M_0 = 0$, $nM_1 = M_1$,

(26)
$$N = N_0 \oplus N_1$$
, where $N_0 n^n = 0, N_1 n = N_1$

for some $n \geq 0$. These decompositions induce (not uniquely!) decompositions in U

(27)
$$a = a_0 + a_1, \mathsf{n}^n a_0 \in U\mathsf{m}^m, a_1 \in \mathsf{n}U + U\mathsf{m}^m, b = b_0 + b_1, b_0 \mathsf{n}^n \in \mathsf{l}^l U, a_1 \in U\mathsf{n} + \mathsf{l}^l U.$$

Then the class of b_0a_0 in ${}_{l}U_{m^m}$ depends only on the classes of a in ${}_{n^n}U_{m^m}$ and the class of b in ${}_{l}U_{n^n}$. Taking the limit as $l,m \longrightarrow \infty$ we obtain the product we need. Obviously, this definition is equivalent to the one given above.

Theorem 5.1. For any $m, n \in \operatorname{Specm} \Gamma$ such that S = S(m, n) is finite, the completed $\Gamma_n - \Gamma_m$ -bimodule $\mathcal{A}(m, n)$ (see (3)) is finitely generated. Moreover, $\mathcal{A}(m, n)$ coincides with the image of U(S) in \mathcal{A} . Besides, $\mathcal{A}(m, n)$ is finitely generated both as a right Γ_m -module and as a left Γ_n -module as well.

Proof. In view of Lemma 5.4 and formula (3) we have an embedding

$$\mathcal{A}(\mathsf{m},\mathsf{n}) \ \subset \lim_{\leftarrow n,m} U(S)/\left((\mathsf{n}^n U + U\mathsf{m}^m) \cap U(S)\right).$$

Moreover, by Remark 5.1, this embedding is in fact an isomorphism.

Since U(S) is a noetherian Γ -bimodule by Theorem 2.2, the generators of U(S) as a Γ -bimodule generate any $U(S)/((\mathsf{n}^n U + U\mathsf{m}^m) \cap U(S))$ as a Γ -bimodule, and hence generate $\mathcal{A}(\mathsf{m},\mathsf{n})$ as a complete $\Gamma_\mathsf{n} - \Gamma_\mathsf{m}$ -bimodule ([Mat], Theorem 8.7). The statement, that $\mathcal{A}(\mathsf{m},\mathsf{n})$ is finitely generated both from the left and from the right,

follows from Theorem 2.2, (2) and from Theorem 8.7, [Mat]. This completes the proof. $\hfill\Box$

Here we prove an analogue of Corollary 5.1.

Theorem 5.2. Let U be a Galois order over a noetherian algebra, $m, n \in \operatorname{Specm} \Gamma$ such that S(m,n), $|\mathcal{M}_m|$, $|\mathcal{M}_n|$ are finite and $S(m,n) \neq \emptyset$. Then the image of U(S) in $\mathcal{A}(m,n)$ (3) is nonzero.

Proof. Consider U as a $\Gamma \otimes \Gamma$ -module and denote by I the ideal $\mathsf{n} \otimes \Gamma + \Gamma \otimes \mathsf{n}$ in $\Lambda \otimes \Lambda$. Then the class of $u \in U$ in $\mathcal{A}(\mathsf{m},\mathsf{n})$ is 0 if for any $N \geq 0$ holds $u \in I^N U$. We prove the following statement: if for some $n \geq 0$ holds $U(S) \subset I^N U$, then $U(S) \subset I^N U(S)$.

Assume $U(S) \subset I^N U$, equivalently, if v_1, \ldots, v_k are the generators of U(S) as Γ -bimodule, then

(28)
$$v_i = \sum_{j=1}^k \nu_{ij} u_{ij}$$
, for some $\nu_{ij} \in I^N \subset \Gamma \otimes \Gamma, u_{ij} \in U, i = 1, \dots, k$.

Set $T = \bigcup_{i=1}^{n} \operatorname{supp} u_i \setminus S$. As in the proof of Lemma 5.4 construct the element

$$f_T = 1 - F \in 1 + \mathsf{n} \otimes \Gamma + \Gamma \otimes \mathsf{m}$$

such that for alls $v_{ij} = f_T \cdot u_{ij}$ holds supp $v_{ij} \subset S$. Applying f_T to the equality 28 we obtain

(29)
$$v_i = F \cdot v_i + \sum_{j=1}^k \nu_{ij} v_{ij}, \ \nu_{ij} \in I^N, v_{ij} \in U(S), i = 1, \dots, k.$$

Substituting the value v_i in the right side of (29) we obtain

$$v_i = F^2 \cdot v_i + (F+1) \sum_{j=1}^k \nu_{ij} v_{ij},$$

and iterated this procedure N-1 times we obtain

$$v_i = F^N \cdot v_i + (F^{N-1} + \dots + F + 1) \cdot \sum_{j=1}^k \nu_{ij} v_{ij},$$

for some
$$\nu_{ij} \in I^N$$
, $v_{ij} \in U(S)$, $i = 1, ..., k$,

which shows $v_i \in I^N U(S)$, since v_i itself and all v_{ij} belongs to U(S). In particular, it means that

$$U(S) = \bigcap_{n=1}^{\infty} I^N U(S).$$

But then Krull Theorem (Theorem 8.9, [Mat]), there exists $a \in 1 + I$, such that

$$a \cdot U(S) = 0.$$

Then the action of a is defined on $V = U(S) \otimes_{\Gamma} K \subset \mathcal{K}$ and $a \cdot V = 0$ as well.

But, following Lemma 2.3, all irreducible summand s V as K-bimodule has a form $V(\varphi)$ for some $\varphi \in \mathcal{M}$, and since supp $V = S(\mathsf{m},\mathsf{n})$, there exist coimages ℓ_{m} and ℓ_{n} , such that $\ell_{\mathsf{n}} = \ell_{\mathsf{m}}^{\varphi}$. Note, that the K-bimodule $V(\varphi)$ is isomorphic to the $L^{H_{\varphi}}$, endowed with the structure of K-bimodule.

The $a \in 1 + I \subset \Gamma \otimes \Gamma$ can be written in the form

$$a = 1 + \sum_{i=1}^{m} \nu_i \otimes \alpha_i + \sum_{j=1}^{n} \beta_j \otimes \mu_j, \alpha_i, \beta_j \in \Gamma, \mu_i \in \mathsf{m}, \nu_j \in \mathsf{n}.$$

Write down the action of a on $1 \in L^{H_{\varphi}}$.

$$0 = a \cdot 1 = \left(1 + \sum_{i=1}^{m} \nu_i \otimes \alpha_i + \sum_{j=1}^{n} \beta_j \otimes \mu_j\right) \cdot 1 =$$

$$1 + \sum_{i=1}^{m} \nu_i^{\varphi} \alpha_i + \sum_{j=1}^{n} \beta_j^{\varphi} \mu_j \in 1 + \ell_n,$$

because all the elements in the formulas above belongs to $\overline{\Gamma}$, $\nu_i \in \ell_n$ and $\mu_j^{\varphi} \in \ell_n$, since $m^{\varphi} \subset \ell_m^{\varphi} = \ell_n$. But $0 \notin 1 + \ell_n$, that completes the proof.

Note that this theorem in the case m=n together with Theorem 3.2 gives another proof of Corollary 5.1.

5.2.2. Finiteness statements for Gelfand-Tsetlin modules. Denote by $r(\mathsf{m},\mathsf{n})$ the minimal number of generators of $U(S(\mathsf{m},\mathsf{n}))$ as a right Γ -module. Since Γ is a Harish-Chandra subalgebra from $|S(\mathsf{m},\mathsf{m})|<\infty$ follows $r(\mathsf{m},\mathsf{n})<\infty$. In this case $\mathcal{A}(\mathsf{m},\mathsf{m})$ is finitely generated as a right Γ_{m} -module (i.e. Γ is big in m). In particular there exists finitely many non-isomorphic simple \mathcal{A} -modules M such that $M(\mathsf{m})\neq 0$, besides $M(\mathsf{m})$ is finite dimensional (Lemma 3.8).

Lemma 5.5. Let $\mathsf{m}, \mathsf{n} \in \operatorname{Specm} \Gamma$, $S = S(\mathsf{m}, \mathsf{n})$, M a Gelfand-Tsetlin module $U \otimes_{\Gamma} \Gamma/\mathsf{m}$, $x = 1 \otimes (1 + \mathsf{m}) \in M(\mathsf{m})$, $\pi_{\mathsf{n}} : M \to M(\mathsf{n})$ the canonical projection. Then

$$\mathcal{A}(\mathsf{m},\mathsf{n})\cdot x = \pi_\mathsf{n}(Ux) = \pi_\mathsf{n}(U(S)x).$$

In particular if M is generated by a vector in M(m) annihilated by m, then

$$\dim_{\mathbb{k}} M(\mathsf{n}) \leq r(\mathsf{m},\mathsf{n}).$$

Proof. The first equality follows from Lemma 3.6. To prove the second equality consider some $u \in U$. Then by the Chinese remainder theorem there exists $\gamma \in \Gamma$ such that $\gamma ux = \pi_{\mathsf{n}}(ux)$ and we change u to γu . Let k = 2t, where $\mathsf{n}^t ux = 0$. Then from Lemma 5.4, $ux = u_k x$, where $u_k \in U(S)$.

Theorem 5.3. Let U be a Galois order over Γ where Γ is a normal noetherian \mathbb{k} -algebra, M is a U-module.

- (1) If for some $m \in D$, \mathcal{M}_m is finite and $M \in \mathbb{H}(U,\Gamma,D)$ is simple then M(m) is finite dimensional. Both $\dim_{\mathbb{K}} M(m)$ and the number of isomorphism classes of simple modules N, such that $N(m) \neq 0$, are bounded by r(m,m).
- (2) If in addition M is a group then for any $n \in D$

$$\dim_{\mathbb{k}} M(\mathsf{n}) \leq r(\mathsf{m},\mathsf{n}) < \infty.$$

(3) Let U be free as a right Γ -module and any projective module over Γ is free. If M is generated over U by $x \in M$, mx = 0, $m \in \operatorname{Specm} \Gamma$, then

(30)
$$r(\mathsf{m},\mathsf{n}) \leq |S(\mathsf{m},\mathsf{n})/G|$$
 If $M = U \otimes_{\Gamma} \Gamma/\mathsf{m}$ then in (30) holds the equality.

Proof. The statement (1) follows from Lemma 5.5 and Lemma 5.3. The statement (2) follows from Lemma 5.3, (1). To prove (3) note that U(S) is a direct summand of U for any G-invariant subset $S \subset \mathcal{M}$. Hence U(S) is free as a right Γ -module and its rank coincides with the K-dimension of $KU(S) \subset U$. But, $\dim_K U(S) = |S/G|$ (Lemma 2.1). Applying this argument for $S = S(\mathfrak{m},\mathfrak{n})$ we obtain the inequality $r(\mathfrak{m},\mathfrak{n}) \leq |S(\mathfrak{m},\mathfrak{n})/G|$. The statement about the cyclic module is obvious.

5.2.3. Finiteness statements for blocks of Gelfand-Tsetlin modules. Let D be an equivalence class in Δ , $m, n \in D$ and $\varphi \in S(m, n)$. If there exists a pair of elements of U, $[a_-\varphi^{-1}]$ and $[a_+\varphi]$, such that $a_-a_+^{\varphi^{-1}}$ and $a_-^{\varphi}a_+$ do not belong to $\{g\ell_m, g \in G\}$ then we say that m and n are connected by φ . Endow D with a structure of a non-oriented graph as follows. The vertices are the elements of D and the edges between m and n are in the bijection with those φ that connect m and n.

Lemma 5.6. If m and n in D are connected by $\varphi \in \mathcal{M}$, then $m \simeq n$ in \mathcal{A}_D .

Proof. As in Lemma 4.4, for every $n \geq 0$ there exists a function $f_n \in \Gamma$, such that

$$f_n \equiv 1 \pmod{\ell_n^n}$$
 and $f(\varphi^g \cdot \ell_m) = 0$ for $g \in G$ such that $\ell_n \neq \varphi^g \cdot \ell_m$.

Consider the element $x_n = [a_-\varphi^{-1}]f_n[a_+\varphi]$. By the assumption, a_- and $a_+^{\varphi^{-1}}$ are nonzero in m and n and hence in $\mathcal A$ there exist $g:m\longrightarrow n$ and $f:n\longrightarrow m$ such that $fg=1_m$. Analogously one constructs f',g' such that $g'f'=1_n$.

Immediately from Lemma 5.6 follows

Corollary 5.3. If m and n belong to the same connected component of the graph D then they are isomorphic in A_D .

Theorem 5.4. Let D be a class of Δ -equivalence. Suppose M is a group, D has a finite number of connected components and for some $m \in D$ the group M_m is finite. Then the module U/Um is of finite length.

Proof. By Corollary 5.3 the skeleton \mathcal{B}_D of the category \mathcal{A}_D contains a finite number of objects, $\operatorname{Ob}\mathcal{B}_D = \{\mathsf{n}_1,\ldots,\mathsf{n}_k\}$. Consider $U/U\mathsf{m}$ as an element in $\mathcal{A}-\operatorname{mod}_d$ and denote the \mathcal{B}_D -module $M=(U/U\mathsf{m})|_{\mathcal{B}_D}$. Then by Theorem 5.3, (2), $\dim_k M(\mathsf{n}_i) \leq r(\mathsf{m},\mathsf{n}_i)$ for any $i=1,\ldots,k$. Since the categories $\mathcal{A}-\operatorname{mod}_d$ and $\mathcal{B}_D-\operatorname{mod}_d$ are equivalent, it completes the proof.

From the proof above we obtain

Corollary 5.4. In the notation of Theorem 5.4 the length of U/Um and the number of simples in the block $\mathbb{H}(U,\Gamma,D)$ does not exceed $\sum_{i=1}^{k} r(m,n_i)$, where n_i runs a set of representatives of connected components of D.

Applying Theorem 5.3, (1) and (3) and Lemma 5.3, (23) we obtain also.

Corollary 5.5. Let U and Γ satisfy the conditions of Theorem 5.3, (3), D a class of Δ -equivalence such that \mathcal{M}_m is finite for some $m \in D$. If the graph D contains $c(D) < \infty$ connected components, then for the number s(D) of isomorphism classes of simple U-modules in the block $\mathbb{H}(U,\Gamma,D)$ holds

$$1 \le s(D) \le |G||\mathcal{M}_{\mathsf{m}}|c(D).$$

The same holds for the length of the module U/Um.

5.2.4. Proof of Theorem A.. We will show that under some conditions for Galois orders there exists (up to isomorphism) finitely many simple Gelfand-Tsetlin modules extending a given character of Γ (hence we will prove **Theorem A**).

Let U be a Galois order over Γ , $\mathbf{m} \in \operatorname{Specm} \Gamma$. Assume that Γ is noetherian over \mathbb{k} and \mathcal{M}_{m} is finite. Then $S(\mathsf{m},\mathsf{m})$ is finite by Lemma 5.3. Consider $\chi:\Gamma \longrightarrow \mathbb{k}$ such that $\mathsf{m} = \operatorname{Ker} \chi$. If Γ is not normal then $\tilde{\Gamma}$ is a finite Γ -module and χ admits finitely many extensions to $\tilde{\Gamma}$, by Corollary 2.2. Hence, it is enough to prove the statement in the case $\tilde{\Gamma} = \Gamma$. But then Theorem 5.1 implies that Γ is big in m . By Lemma 3.8 there exists only finitely many non-isomorphic extensions of χ to simple U-modules, which completes the proof of Theorem A.

5.3. Gelfand-Tsetlin modules for Galois orders. In this subsection we assume that U is a Galois order over noetherian Γ . Let $\bar{\Gamma}$ be the integral closure of Γ in L, $\varphi \in \mathcal{M}$ and $i : \Gamma \to \bar{\Gamma}$, $i_{\varphi} : \varphi(\Gamma) \to \bar{\Gamma}$ the canonical embeddings, π and π_{φ} the induced mappings of the maximal spectra, $\mathbf{m} \in \operatorname{Specm} \Gamma$, $\pi^{-1}(\mathbf{m}) = \{\ell_1, \ldots, \ell_k\}$. The following lemma describes the sets $X_a, a \in U$ (see Lemma 3.1).

Lemma 5.7. Let $a \in L^{H_{\varphi}}$ and $V = \Gamma[a\varphi]\Gamma$. Then the set of simple factors of the right Γ -module $V \otimes_{\Gamma} \Gamma/m$ coincides with the set of simples of the form Γ/n , $n \in \pi(\varphi(\pi^{-1}(m))) = \{\pi(\ell_1^{\varphi}), \ldots, \pi(\ell_k^{\varphi})\}$. Besides, for $u \in U$ and $m \in \operatorname{Specm} \Gamma$ holds $X_u(m) = \pi(\sup u \cdot \pi^{-1}(m))$.

Proof. Write $V = [a\Gamma\Gamma^{\varphi}\varphi]$ and $V\mathsf{m} = [a\Gamma\Gamma^{\varphi}\mathsf{m}^{\varphi}\varphi]$. Since $\Gamma^{\varphi} \subset \Gamma\Gamma^{\varphi}$ is a finite integral extension, then $V \neq V\mathsf{m}$ and $V/V\mathsf{m} \simeq \Gamma\Gamma^{\varphi}/\Gamma\mathsf{m}^{\varphi}$ (Remark 2.1). The homomorphisms $p: \bar{\Gamma} \to \mathbb{k}$ extending $p^{\varphi}: \Gamma^{\varphi} \to \Gamma^{\varphi}/\mathsf{m}^{\varphi} \simeq \mathbb{k}$ form a G^{φ} - orbit $\{\pi(\ell_{k}^{\varphi}), \ldots, \pi(\ell_{k}^{\varphi})\}$, whose restrictions on Γ give all characters of Γ . This proves the first statement.

Let $V = \Gamma u \Gamma$. Then Lemma 2.4, (5) reduces the second statement to the case of Γ -bimodule V generated by elements of the form $[a_1\varphi], \ldots, [a_k\varphi]$. Hence the second statement follows from the first one.

5.4. Gelfand-Tsetlin categories for Galois orders. In this subsection we study in more details the category of Gelfand-Tsetlin modules over Galois order U over Γ . We assume that Γ is normal noetherian \Bbbk -algebra and Ω_2 and Ω_r are as in Section 4. For $\mathsf{m} \in \operatorname{Specm} \Gamma$ denote by $D(\mathsf{m})$ denote the class of Δ -equivalence, containing m .

Theorem 5.5. (1) Let $S(\mathsf{m},\mathsf{n}) = \emptyset$ for some $\mathsf{m},\mathsf{n} \in \operatorname{Specm} \Gamma$. Then $\mathcal{A}(\mathsf{m},\mathsf{n}) = \emptyset$

(2) Let
$$a \in U$$
, $a = \sum_{i=1}^{n} [a_i m_i]$, $m_i \in M$, $a_i \in L^*$ and

$$\mathfrak{X}(a) = \{ (m_i \cdot \ell^g, \ell) \mid \ell \in \mathcal{L}, i = 1, \dots, n, g \in G \}.$$

Then $X_a \subset (\pi \times \pi)(\mathfrak{X}(a))$. In particular, if $m, n \in \operatorname{Specm} \Gamma$ and $u \in U$, then the image of u in $\mathcal{A}(m, n)$ is nonzero, only if $u \in S(m, n)$.

- (3) If $m \in \Omega_r$, then $\mathcal{A}(m, m)$ is a homomorphic image of Γ_m . In particular, there exists a unique up to isomorphism simple U-module M, extending the character $\chi : \Gamma \longrightarrow \Gamma/m$.
- (4) Let $m \in \Omega_r$, D = D(m), $M_m = \mathcal{A}_D/\mathcal{A}_D\hat{m}$, where $\hat{m} \subset \Gamma_m$ is the completed ideal. Then U/Um is canonically isomorphic to $\mathbb{F}(M_m)$.
- (5) Let \mathcal{M} be a group, $\mathbf{m} \in \Omega_r \cap \Omega_2$. Then for every $\mathbf{n} \in D(\mathbf{m})$,

$$\mathcal{A}(\mathsf{n},\mathsf{n}) \simeq \Gamma_\mathsf{n}$$

and all objects of A_D are isomorphic.

Proof. The statement (1) follows from Lemma 5.4 and (3). To prove (2) note, that the Γ-bimodule $\Gamma a\Gamma$ is a factor of $\bigoplus_{i=1}^n \Gamma[am_i]\Gamma$, so it is enough to prove, that if for $\mathsf{m}, \mathsf{n} \in \operatorname{Specm}\Gamma$, Γ/n is a subfactor of $\Gamma[am]\Gamma/\mathsf{m}$, then $(\mathsf{m}, \mathsf{m}) \in (\pi \times \pi)(\mathfrak{X}(a))$.

 $\mathsf{m}, \mathsf{n} \in \operatorname{Specm} \Gamma$, Γ/n is a subfactor of $\Gamma[am]\Gamma/\mathsf{m}$, then $(\mathsf{m}, \mathsf{m}) \in (\pi \times \pi)(\mathcal{X}(a))$. Since Ext^1 -spaces between non-isomorphic simples in commutative case are zero, we can consider just factors instead of subfactors.

Consider the canonical embedding of Γ -subbimodules in $L * \mathcal{M}$

$$i: \Gamma[am]\Gamma \hookrightarrow \bar{\Gamma} \otimes_{\Gamma} \Gamma[am]\Gamma \otimes_{\Gamma} \bar{\Gamma} \simeq \bar{\Gamma}[am]\bar{\Gamma}.$$

Then, since $\bar{\Gamma}$ is a faithfully flat (i.e. $\bar{\Gamma}$ is flat and $\bar{\Gamma} \otimes_{\Gamma} M \neq 0$ for any nonzero Γ -module M, [Mat], Theorem 7.2) Γ -module, every $\chi : \Gamma[am]\Gamma \longrightarrow \mathbb{k}$ lifts to

$$\bar{\chi}: \bar{\Gamma} \otimes_{\Gamma} \Gamma[am]\Gamma \otimes_{\Gamma} \bar{\Gamma} \longrightarrow \mathbb{k}.$$

Hence, every simple factor of $\Gamma[am]\Gamma$ as Γ -bimodule factorizes through i. Consider a homomorphism of Γ -bimodules

$$\bar{\Gamma} \otimes_{\mathbb{k}} \bar{\Gamma} \xrightarrow{\pi} \bar{\Gamma}[am] \bar{\Gamma} \xrightarrow{\bar{\chi}} \mathbb{k}.$$

Then the composition is just the pair $(m \cdot \ell^g, \ell)$, where $\ell \in \mathcal{L}$ and $g \in G$. It proves (2). To prove (3) we note that, by Lemma 5.4 and by (3), $\mathcal{A}(\mathsf{m}, \mathsf{m})$ is generated as Γ_{m} -bimodule by the central element \bar{e} , the class of $e \in U$. On other hand, there exists the canonical complete algebra homomorphism $i: \Gamma_{\mathsf{m}} \longrightarrow \mathcal{A}(\mathsf{m}, \mathsf{m})$, $i(1) = \bar{e}$, which is clearly surjective. Obviously, $\mathcal{A}_D - \mathrm{mod}_d$ is equivalent to the full subcategory $\mathbb{F}(\mathcal{A}_D - \mathrm{mod}_d) \subset U - \mathrm{mod}$. For $\mathsf{m} \in \mathrm{Specm}\,\Gamma$ consider the functor $W_{\mathsf{m}}: U - \mathrm{mod} \longrightarrow \mathbb{k} - \mathrm{Mod}$, which sends $M \in U - \mathrm{mod}$ to

$$W_{\mathsf{m}}(M) = \{ x \in M \mid \mathsf{m} \cdot x = 0 \}.$$

This is obviously a representable functor, namely

$$W_{\mathsf{m}} \simeq \operatorname{Hom}_U(U/U\mathsf{m}, -).$$

On the other hand, for an indecomposable N, which does not belong to the image of \mathbb{F} , holds

$$\operatorname{Hom}_{U}(\mathbb{F}(M_{\mathsf{m}}), N) = 0,$$

and for $N = \mathbb{F}(N')$ we have

$$\operatorname{Hom}_{U}(\mathbb{F}(M_{\mathsf{m}}), \mathbb{F}(N')) \simeq \operatorname{Hom}_{\mathcal{A}}(M_{\mathsf{m}}, N') \simeq W_{\mathsf{m}}(\mathbb{F}(N')) = W_{\mathsf{m}}(N),$$

where all isomorphisms are functorial, i.e, $U/Um \simeq \mathbb{F}(M_m)$. It implies (4). Let us prove (5). As shown, $\mathcal{A}(m, m)$ is a homomorphic image of Γ_m .

Consider in $\mathbb{F}(\mathcal{A}_{D(m)} - \text{mod}_d)$ a *U*-module:

$$M_{\chi,n} = U \otimes_{\Gamma} \Gamma/\mathsf{m}^n \simeq \bigoplus_{\mathsf{n} \in \mathfrak{O}_\mathsf{m}} \Gamma/\mathsf{n}^n.$$

Any nonzero element from Γ_{m} acts nontrivially on $M_{\chi,n}$ for any n. Thus $\mathcal{A}(\mathsf{m},\mathsf{m}) \simeq \Gamma_{\mathsf{m}}$ by the Krull intersection theorem ([Mat], Theorem 8.10, (II)).

Corollary 5.6. Let M be a group, $D = D(m) \subset \operatorname{Specm} \Gamma$ a $\Delta(U, \Gamma)$ -equivalence class of a maximal ideal $m \in \Omega_r \cap \Omega_2$. Then the category $\mathbb{H}(U, \Gamma, D)$ is equivalent to the category $\Gamma_m - \operatorname{mod}$.

Proof. Since all objects in \mathcal{A}_D are isomorphic by Theorem 5.5, (5) the categories $\mathcal{A}_D - \text{mod}$ and $\mathcal{A}(\mathsf{m},\mathsf{m}) - \text{mod}$ are equivalent. Note that the functors of restriction $res: \mathbb{H}(U,\Gamma,D) \to \mathcal{A}(\mathsf{m},\mathsf{m}) - \text{mod}$ and of induction $ind: \mathcal{A}(\mathsf{m},\mathsf{m}) \to \mathbb{H}(U,\Gamma,D)$ are quasi-inverse.

Remark 5.2. Recall that if m is nonsingular point of Specm Γ , then Γ_m is isomorphic to the algebra of formal power series in $GKdim \Gamma$ variables.

Remark 5.3. In the assumption of Theorem 5.5, (2) a nonzero element $u \in S(m, n)$ can turn zero in A(m, n).

5.5. **Proof of Theorem B.** Theorem 4.1, Theorem 5.5 and Corollary 5.6 immediately imply the first part of **Theorem B**.

The second part of Theorem B is an analogue of the Harish-Chandra theorem for the universal enveloping algebras ([D]). In particular it shows that the subcategories in U-mod, described in Corollary 5.6, contain enough modules. Suppose that conditions of Theorem B are satisfied. Consider the massive set Ω_u constructed in Proposition 4.1 and set

$$\Omega'_u = \Omega_u \cap \Omega_2 \cap \Omega_r$$
.

Then for any $m \in \Omega'_u$ the element u acts nontrivially on U/Um which is simple by Theorem 5.5. This completes the proof of Theorem B.

6. Gelfand-Tsetlin modules for gl_n

Consider the general lineal Lie algebra gl_n with the standard basis $e_{ij}, i, j = 1, \ldots, n$. Set $U = U(\mathrm{gl}_n), \ U_m = U(\mathrm{gl}_m), \ 1 \leq m \leq n$. Let Z_m be the center of U_m . We identify gl_m for $m \leq n$ with a Lie subalgebra of gl_n spanned by $\{e_{ij} \mid i, j = 1, \ldots, m\}$, so that we have the following chain of inclusions

$$\mathrm{gl}_1 \subset \mathrm{gl}_2 \subset \ldots \subset \mathrm{gl}_n$$
.

It induces the inclusions $U_1 \subset U_2 \subset \ldots \subset U_n$ of the universal enveloping algebras. The Gelfand-Tsetlin subalgebra ([DFO1]) Γ in U is generated by $\{Z_m \mid m = 1, \ldots, n\}$, where Z_m is a polynomial algebra in m variables $\{c_{mk} \mid k = 1, \ldots, m\}$,

(31)
$$c_{mk} = \sum_{(i_1,\dots,i_k)\in\{1,\dots,m\}^k} e_{i_1i_2}e_{i_2i_3}\dots e_{i_ki_1}.$$

Hence Γ is a polynomial algebra in $\frac{n(n+1)}{2}$ variables $\{c_{ij} \mid 1 \leqslant j \leqslant i \leqslant n\}$ ([Zh]).

Denote by K be the field of fractions of Γ . Let $\mathcal{M} \subset \mathcal{L}$, $\mathcal{M} \simeq \mathbb{Z}^{\frac{n(n-1)}{2}}$ be a free abelian group generated by δ^{ij} , $1 \leq j \leq i \leq n-1$, $(\delta^{ij})_{kl} = 1$ if i = k, j = l and 0 otherwise. For $i = 1, \ldots, n$ denote by S_i the i-th symmetric group and set $G = S_1 \times \ldots \times S_n$. The group G acts on \mathfrak{l}_i : $(s \cdot \ell)_{ij} = \ell_{s_j(i)j}$ for $\ell = (\ell_{ij}) \in \mathcal{L}$ and $s = (s_i) \in G$. Also the group \mathcal{M} acts on \mathcal{L} as follows $\delta^{ij} \cdot \ell = \ell + \delta^{ij}$, $\delta^{ij} \in \mathcal{M}$.

Let Λ be a polynomial algebra in variables $\{\lambda_{ij} \mid 1 \leq j \leq i \leq n\}$ and L be the fraction field of Λ . Let $i: \Gamma \longrightarrow \Lambda$ be an embedding such that

$$i(c_{mk}) = \sum_{i=1}^{m} (\lambda_{mi} + m)^k \prod_{j \neq i} (1 - \frac{1}{\lambda_{mi} - \lambda_{mj}}).$$

The image of i coincides with the subalgebra of G—invariant polynomials in Λ ([Zh]) which we identify with Γ . The homomorphism i can be extended to the embedding of the fields $K \subset L$, where $L^G = K$ and G = G(L/K) is the Galois group. An action of the group G by conjugations on M induces its action on L * M.

Let e be the identity element of the group \mathcal{M} . Consider a linear map $\mathbf{t}:U\longmapsto\mathcal{K}$ such that

(32)
$$t(e_{mm}) = ee_{mm}, \ t(e_{m\,m+1}) = \sum_{i=1}^{m} A_{mi}^{+} \delta^{mi}, \ t(e_{m+1\,m}) = \sum_{i=1}^{m} A_{mi}^{-} (\delta^{mi})^{-1},$$

where

$$A_{mi}^{\pm} = \mp \frac{\prod_{j} (\lambda_{m\pm 1,j} - \lambda_{mi})}{\prod_{j\neq i} (\lambda_{mj} - \lambda_{mi})}.$$

The map t is an algebra homomorphism by the Harish-Chandra theorem. Moreover, we have

Proposition 6.1. (see [FO1], Proposition 7.2) Let $S = \Gamma \setminus \{0\}$. Then

- t is an embedding;
- $UK = KU \simeq (L * \mathbb{Z}^m)^G$, m = n(n-1)/2;
- U is a Galois order over Γ .

Since \mathcal{M} acts freely on Specm Γ then \mathcal{M}_{m} is trivial. Hence $|G|^2$ provides a bound for the number of isomorphism classes of simple U-modules V, such that $V(\mathsf{m}) \neq 0$, and for the $\dim_{\mathbb{K}} V(\mathsf{m})$. But this bound can be improved in this case. Set $Q_n = \prod_{i=1}^{n-1} i!$.

Corollary 6.1. Let $U=U(\operatorname{gl}_n),\ \Gamma\subset U$ is the Gelfand-Tsetlin subalgebra, D a $\Delta\text{-class},\ \mathsf{m}\in D.$ Then

(1) For a simple U-module $M(m) \neq 0$ holds

$$\dim_{\mathbb{K}} M(\mathsf{m}) \leq Q_n$$
.

- (2) There exist at most Q_n isoclasses of simple U-modules N, such that $N(\mathsf{m}) \neq 0$.
- (3) Denote by c(D) the number of connected components in D. Then the number of isomorphism classes of simple modules in $\mathbb{H}(U,\Gamma,D)$ does not exceed $c(D)Q_n$. The same number bounds the length of U/U m.

Proof. In any Δ -equivalence class D the central character is fixed and hence $D = \pi(\ell_{\mathsf{m}} + \mathbb{Z}^{n(n-1)/2})$ for some $\mathsf{m} \in D$. Then for $\mathsf{n} \in D$ and some fixed ℓ_{n} there exists finitely many (bounded by Q_n) $s \in \mathbb{Z}^{n(n-1)/2}$ such that ℓ_{n} and $\ell_{\mathsf{n}} + s$ differ by the action of G. It immediately implies (2). Note that we obtain a better bound than given by Theorem 5.3. Since U is free as a left (right) Γ -module ([O]) the statement (1) follows from Lemma 5.3 and Theorem 5.3,(3). Finally (3) follows from (2) and Corollary 5.5.

Remark 6.1. We believe that the bound Q_n can not be improved. It is known to be exact for n = 2 and n = 3 [DFO1].

Remark 6.2. Note that U/Um is holonomic and hence it has a finite length [Be].

Remark 6.3. Due to Corollary 6.1 the category of Gelfand-Tsetlin gl_n -modules splits into a direct sum of full subcategories each of which is equivalent to a module category of some associative algebra (quiver with relations).

7. Acknowledgment

The first author is supported in part by the CNPq grant (processo 301743/2007-0) and by the Fapesp grant (processo 2005/60337-2). The second author is grateful to Fapesp for the financial support (processos 2004/02850-2 and 2006/60763-4) and to the University of São Paulo for the hospitality during his visits.

References

- [Ba] Bavula V., Generalized Weyl algebras and their representations, Algebra i Analiz 4 (1992), 75–97. (English translation: St. Petersburg Math. J. 4 (1993), 71–92.
- [BavO] Bavula V., Oystaeyen F., Simple Modules of the Witten-Woronowicz algebra, Journal of Algebra 271 (2004), 827–845.
- [Be] Bernstein I.N., Modules over a ring of differential operators. An investigation of the fundamental solutions of equations with constant coefficients, Funkt. Anal. i Prilozehn. 5 (1971), no 2, 1-16.
- [D] Dixmier J., Algèbres Enveloppantes. Paris: Gauthier-Villars, 1974.

- [Dr] Drozd, Yu. A., On representations of the Lie algebra sl₂, in Ukrainian, Bulletin of the Kiev State University, Math, and Mech., 25 (1983), 70-77.
- [DK] Drozd Yu. A., Kirichenko V.V., Finite Dimensional Algebras, with Appendix by Vlastimil Dlab. Berlin Springer Verlag, 1994. - 249p.
- [DFO1] Drozd Yu.A., Ovsienko S.A., Futorny V.M. On Gelfand-Zetlin modules, Suppl. Rend. Circ. Mat. Palermo, 26 (1991), 143-147.
- [DFO2] Drozd Yu., Ovsienko S., Futorny V., Harish Chandra subalgebras and Gelfand Zetlin modules, in: "Finite dimensional algebras and related topics", NATO ASI Ser. C., Math. and Phys. Sci., 424, (1994), 79-93.
- [Fe] Fernando S., Lie algebra modules with finite-dimensional weight spaces I, Trans.Amer.Math.Soc. 322 (1990), 757-781.
- [FM] Fomenko T., Mischenko A., Euler equation on finite-dimensional Lie groups, Izv. Akad. Nauk SSSR, Ser. Mat. 42 (1978), 396-415.
- [FMO] Futorny V., Molev A., Ovsienko S., Harish-Chandra modules for Yangians, Representation Theory, AMS, 9 (2005), 426-454.
- [FMO1] Futorny V., Molev A., Ovsienko S., Gelfand-Kirillov Conjecture and Harish-Chandra modules for finite W-algebras, arXiv:0801.0903v2.
- [FO1] Futorny V., Ovsienko S., Galois orders, arXiv:math/0610069v2.
- [FO2] Futorny V., Ovsienko S., Galois algebras II: Representation theory, preprint RT-MAT 2006-14, 28p.
- [FO3] Futorny V., Ovsienko S. Kostant's theorem for special filtered algebras, Bull. London Math. Soc. 37 (2005), 1-13.
- [GR] Gabriel P., Roiter A.V. Representations of finite-dimensional algebras, Encyclopedia of the Mathematical Sciences, Springer, Algebra VIII, 1992. Vol. 73. 182 p.
- [GTs] Gelfand I.M., Tsetlin M.S., Finite dimensional representations of the group of unimodular matrices, Doklady Akad. Nauk SSSR, 71 (1950), 1017-1020.
- [Gr] Graev M.I., Infinite-dimensional representation sof the Lie algebra gl(n, C) related to complex analogs of the Gelfand-Tsetlin patterns and general hupergeometric functions on the Lie group GL(n, C), Acta Appl. Mathematicae 81 (2004), 93-120.
- [KW1] Kostant B., Wallach N.: Gelfand-Zeitlin theory from the perspective of classical mechanics I. In Studies in Lie Theory Dedicated to A. Joseph on his Sixtieth Birthday, Progress in Mathematics, Vol. 243, (2006), 319-364.
- [KW2] Kostant B., Wallach N.: Gelfand-Zeitlin theory from the perspective of classical mechanics II. In The Unity of Mathematics In Honor of the Ninetieth Birthday of I.M. Gelfand, Progress in Mathematics , Vol. 244, (2006), 387–420.
- [Mat] Matsumura H., Commutative ring theory, Cambridge Studies in Advanced Mathematics, 8, Cambridge University Press, 1997.
- [Ma] Mathieu O., Classification of irreducible weight modules, Ann. Inst. Fourier 50 (2000), 537-592.
- [M] Molev A. I., Gelfand-Tsetlin bases for classical Lie algebras, in "Handbook of Algebra", (M. Hazewinkel, Ed.), Elsevier, to appear.
- [OV] Okunkov A., Vershik A., A new approach to representation theory of symmetric groups, Selecta Math. New Series, 2 (1996), 581-605.
- [O] Ovsienko S., Strongly nilpotent matrices and Gelfand-Tsetlin modules, Linear Algebra and Its Appl., 365 (2003), 349-367.
- [S] Serre J. P., Local algebra, Springer Monographs in Mathematics, Springer, 2000.
- [Vi] Vinberg E., On certain commutative subalgebras of a universal enveloping algebra, Math. USSR Izvestiya 36 (1991), 1-22.
- [Zh] Zhelobenko D.P. Compact Lie groups and their representations, Nauka, Moscow, 1970 (Translations of mathematical monographs, 40, AMS, Providence, Rhode Island, 1973).

Instituto de Matemática e Estatística, Universidade de São Paulo, Caixa Postal 66281, São Paulo, CEP 05315-970, Brasil

E-mail address: futorny@ime.usp.br

Faculty of Mechanics and Mathematics, Kiev Taras Shevchenko University, Vladimirskaya 64, 00133, Kiev, Ukraine

 $E\text{-}mail\ address: \verb|ovsienko@zeos.net|$