# COMPOSITION OPERATORS WITHIN SINGLY GENERATED COMPOSITION C\*-ALGEBRAS

THOMAS L. KRIETE, BARBARA D. MACCLUER, AND JENNIFER L. MOORHOUSE

ABSTRACT. Let  $\varphi$  be a linear-fractional self-map of the open unit disk  $\mathbb{D}$ , not an automorphism, such that  $\varphi(\zeta) = \eta$  for two distinct points  $\zeta, \eta$  in the unit circle  $\partial \mathbb{D}$ . We consider the question of which composition operators lie in  $C^*(C_{\varphi}, \mathcal{K})$ , the unital  $C^*$ -algebra generated by the composition operator  $C_{\varphi}$  and the ideal  $\mathcal{K}$  of compact operators, acting on the Hardy space  $H^2$ . This necessitates a companion study of the unital  $C^*$ -algebra generated by the composition operators induced by all parabolic non-automorphisms with common fixed point on the unit circle.

# 1. INTRODUCTION

Given any analytic self-map  $\varphi$  of the unit disk  $\mathbb{D}$  in the complex plane, one can form the composition operator  $C_{\varphi} : f \to f \circ \varphi$ , which acts as a bounded operator on the Hardy space  $H^2$ . This paper is the second in a series of three investigating spectral theory in  $C^*$ -algebras generated by certain composition and Toeplitz operators. In the first article [15], we studied  $C^*(T_z, C_{\varphi})$ , the unital  $C^*$ algebra generated by the unilateral shift  $T_z$  on  $H^2$  and a single composition operator  $C_{\varphi}$  with  $\varphi$  satisfying

(1)

 $\begin{cases} \varphi \text{ is a linear-fractional self-map of } \mathbb{D} \text{ which is not an automorphism, and} \\ \varphi(\zeta) = \eta \text{ for distinct points } \zeta, \eta \text{ in the unit circle } \partial \mathbb{D}. \end{cases}$ 

Throughout the current paper,  $\varphi$  will always have this meaning. The algebra  $C^*(T_z, C_{\varphi})$  necessarily contains the ideal  $\mathcal{K}$  of compact operators on  $H^2$ . The main result of [15] identifies  $C^*(T_z, C_{\varphi})/\mathcal{K}$  with a certain  $C^*$ -algebra of  $2 \times 2$  matrix valued functions; see Theorem 4.12 of [15]. The case where  $\varphi$  is replaced by an automorphism of  $\mathbb{D}$ , or even a discrete group of automorphisms, has been studied by Jury [12], [13], and has a rather different character.

The shift  $T_z$  does not appear to play a role in the questions we consider in this paper; accordingly we omit it and study  $C^*(C_{\varphi}, \mathcal{K})$ , the unital  $C^*$ -algebra generated by  $C_{\varphi}$ , for  $\varphi$  as described above, and the compact operators. The composition  $\varphi \circ \varphi$  has sup-norm strictly less than 1, so that  $C_{\varphi}^2 = C_{\varphi \circ \varphi}$  is compact and nonzero. Since  $\varphi$  has no boundary fixed point, a theorem of Guyker [10] shows that  $C_{\varphi}$  is irreducible if and only if  $\varphi(0) \neq 0$ . It follows that when  $\varphi(0) \neq 0$  the unital  $C^*$ -algebra  $C^*(C_{\varphi})$  generated by  $C_{\varphi}$  alone contains  $\mathcal{K}$ ; see [5], p.74. We want our  $C^*$ -algebras to always contain  $\mathcal{K}$ , and we indicate this by continuing to write  $C^*(C_{\varphi}, \mathcal{K})$  if the irreducibility criterion  $\varphi(0) \neq 0$  is in doubt.

Date: July 10, 2008.

<sup>2000</sup> Mathematics Subject Classification. Primary 47B33.

Let  $\mathcal{P}$  denote the dense subalgebra of  $C^*(C_{\varphi}, \mathcal{K})$  consisting of all finite linear combinations of the identity I, all words in  $C_{\varphi}$  and  $C_{\varphi}^*$ , and all compact operators. An element B in  $\mathcal{P}$  has a unique representation of the form

(2) 
$$B = cI + f(C_{\varphi}^*C_{\varphi}) + g(C_{\varphi}C_{\varphi}^*) + C_{\varphi}p(C_{\varphi}^*C_{\varphi}) + C_{\varphi}^*q(C_{\varphi}C_{\varphi}^*) + K$$

where f, g, p and q are polynomials, f(0) = 0 = g(0), c is complex, and K is compact. Let  $s = 1/|\varphi'(\zeta)|$  and write  $\mathcal{D}$  for the  $C^*$ -algebra of continuous  $2 \times 2$ matrix-valued functions F on [0, s] with F(0) a scalar multiple of the identity, equipped with the supremum operator norm. It was shown in [15] that there is a unique \*-homomorphism  $\Psi$  of  $C^*(C_{\varphi}, \mathcal{K})$  onto  $\mathcal{D}$  with Ker  $\Psi = \mathcal{K}$  and such that

(3) 
$$\Psi(B) = \begin{bmatrix} c+g & rp \\ rq & c+f \end{bmatrix}$$

where B is given by Equation (2) and  $r(t) = \sqrt{t}$ . Equivalently, we have a short exact sequence of  $C^*$ -algebras

$$0 \to \mathcal{K} \xrightarrow{i} C^*(C_{\varphi}, \mathcal{K}) \xrightarrow{\Psi} \mathcal{D} \to 0$$

where *i* is inclusion. For any operator *T* on  $H^2$  we write  $||T||_e$  for the essential norm of *T*, that is, the distance from *T* to the ideal  $\mathcal{K}$ . We note that if *T* is in  $C^*(C_{\varphi}, \mathcal{K})$ , then  $||T||_e = ||\Psi(T)||$ .

For bounded operators A and B on  $H^2$ , let us write  $A \equiv B \pmod{\mathcal{K}}$  if there exists a compact operator K with A = B + K. In [15] the authors used C. Cowen's well-known adjoint formula [7] to show that if

$$\psi(z) = \frac{az+b}{cz+d}$$

is a linear-fractional self map of  $\mathbb{D}$ , not an automorphism but satisfying  $|\psi(z_0)| = 1$  for some  $z_0 \in \partial \mathbb{D}$ , then

(4) 
$$C_{\psi}^* \equiv \frac{1}{|\psi'(z_0)|} C_{\sigma_{\psi}} \pmod{\mathcal{K}}$$

where  $\sigma_{\psi}$  is the so-called "Krein adjoint" of  $\psi$ ,

$$\sigma_{\psi}(z) = \frac{\overline{a}z - \overline{c}}{-\overline{b}z + \overline{d}}$$

We denote the Krein adjoint of  $\varphi$  itself simply by  $\sigma$ , which the reader can distinguish by context from the spectrum  $\sigma(T)$  or essential spectrum  $\sigma_e(T)$  of an operator T.

Now consider an operator B in the dense subalgebra  $\mathcal{P}$ , as expressed in Equation (2). Since  $I = C_z, C_{\varphi}^* C_{\varphi} \equiv s C_{\varphi \circ \sigma} \pmod{\mathcal{K}}$  and  $C_{\varphi} C_{\varphi}^* \equiv s C_{\sigma \circ \varphi} \pmod{\mathcal{K}}$ , where  $s = 1/|\varphi'(\zeta)|$ , we can rewrite Equation (2) as

(5) 
$$B = cC_z + A_1 + A_2 + A_3 + A_4 + K'$$

where K' is compact and  $A_1, A_2, A_3$ , and  $A_4$  are finite linear combinations of composition operators whose associated self-maps are chosen, respectively, from the four lists

(6) 
$$\{(\varphi \circ \sigma)_{n_1}\}, \{(\sigma \circ \varphi)_{n_2}\}, \{(\varphi \circ \sigma)_{n_3} \circ \varphi\}, \{(\sigma \circ \varphi)_{n_4} \circ \sigma\}.$$

Here we write  $(\psi)_n$  for the  $n^{th}$  iterate of a self-map  $\psi$  of  $\mathbb{D}$ , and let  $n_k$  range over the positive integers for k = 1, 2, and over the non-negative integers for k = 3, 4. See [15] for further details. Thus  $C^*(C_{\varphi}, \mathcal{K})$  is spanned, modulo the compacts, by actual composition operators. This leads to our main question: for which analytic self-maps  $\psi$  of  $\mathbb{D}$  does  $C_{\psi}$  lie in  $C^*(C_{\varphi}, \mathcal{K})$ ?

In particular we will describe explicitly, in both function-theoretic and operatortheoretic terms, all linear-fractional composition operators lying in  $C^*(C_{\varphi}, \mathcal{K})$ . This description plays a role in the third paper of our series [16] which is devoted to spectral theory in Toeplitz-composition algebras with several generators. Along the way here we show that if  $C^*(\mathbb{P}_{\gamma})$  denotes the unital  $C^*$ -algebra generated by composition operators induced by the parabolic non-automorphisms fixing  $\gamma$  in the unit circle, then there is a short exact sequence

$$0 \to \mathcal{K} \to C^*(\mathbb{P}_{\gamma}) \to C([0,1]) \to 0$$

of  $C^*$ -algebras, where C([0, 1]) denotes the algebra of continuous functions on the unit interval.

# 2. Necessary conditions

If  $\psi : \mathbb{D} \to \mathbb{D}$  is analytic we write  $F(\psi)$  for the set of points  $\alpha$  in  $\partial \mathbb{D}$  at which  $\psi$  has a finite angular derivative in the sense of Caratheodory; see [8],[21]. In particular, if  $\alpha$  is in  $F(\psi)$ , the nontangential limit  $\psi(\alpha)$  necessarily exists and has modulus one. We write  $\psi'(\alpha)$  for the angular derivative of  $\psi$  at  $\alpha$ . Recall that this is the ordinary derivative if  $\psi$  extends analytically to a neighborhood of  $\alpha$  and  $|\psi(\alpha)| = 1$ .

It is well known that if  $C_{\psi}$  is compact on  $H^2$ , then  $F(\psi)$  is empty [20]. When  $C_{\psi}$  is considered as acting on the Bergman space in  $\mathbb{D}$ , the converse assertion is true [17]. On the space  $H^2$  considered here, however, " $C_{\psi}$  is compact" is a strictly stronger requirement than " $F(\psi)$  is empty"; see, for example, [8],[21]. Our first goal is to show that when  $C_{\psi}$  lies in  $C^*(C_{\varphi}, \mathcal{K})$  these two conditions are equivalent.

First we recall that a linear-fractional self-map  $\rho$  of  $\mathbb{D}$  is *parabolic* if  $\rho$  fixes one point  $\gamma$  in  $\partial \mathbb{D}$  and is conjugate, via the map  $(\gamma + z)/(\gamma - z)$ , to translation in the right half-plane  $\Omega = \{w : \operatorname{Re} w > 0\}$  by a complex number a with non-negative real part. We denote this parabolic map by  $\rho_a$ , or by  $\rho_{\gamma,a}$  if the fixed point  $\gamma$  is not clear from the context. A linear-fractional map  $\rho$  with fixed point  $\gamma$  in  $\partial \mathbb{D}$  is parabolic provided  $\rho'(\gamma) = 1$ . Another representation of  $\rho_a$  will prove useful. The map  $\tau_{\gamma}(z) = i(\gamma - z)/(\gamma + z)$  carries  $\mathbb{D}$  onto the upper half-plane  $\{w : \operatorname{Im} w > 0\}$ and takes  $\gamma$  to 0, rather than infinity. We write u for the conjugate of  $\rho_a$  by  $\tau_{\gamma}$ :  $u = \tau_{\gamma} \circ \rho_a \circ \tau_{\gamma}^{-1}$ . One readily computes that u(w) = iw/(i + wa), and so

$$(7) u''(0) = 2ia$$

Also important for us will be several lower bounds for the essential norm of a linear combination of composition operators. Given an analytic self-map  $\psi$  of  $\mathbb{D}$  and  $\alpha$  in  $F(\psi)$ , we call  $D_1(\psi, \alpha) \equiv (\psi(\alpha), \psi'(\alpha))$  the first order data vector for  $\psi$  at  $\alpha$ . If we have a finite collection of maps  $\psi_1, \psi_2, \dots, \psi_n$  and  $\alpha$  lies in the union of the finite angular derivative sets  $F(\psi_1), F(\psi_2), \dots F(\psi_n)$ , we define

$$\mathcal{D}_1(\alpha) = \{ D_1(\psi_j, \alpha) : 1 \le j \le n \text{ and } \alpha \in F(\psi_j) \},\$$

the set of possible first order data vectors at  $\alpha$  coming from that collection of maps. Theorem 5.2 in [14] states that if  $\mathcal{D}_1(\alpha)$  is non-empty, then for any complex numbers  $c_1, \dots, c_n$ ,

(8) 
$$||c_1 C_{\psi_1} + \dots + c_n C_{\psi_n}||_e^2 \ge \sum_{\mathbf{d} \in \mathcal{D}_1(\alpha)} \left| \sum_{\substack{\alpha \in F(\psi_j) \\ D_1(\psi_j, \alpha) = \mathbf{d}}} c_j \right|^2 \frac{1}{|d_1|},$$

where **d** =  $(d_0, d_1)$ .

There is a higher order version of this lower bound which works provided that for the specific  $\alpha$  in  $F(\psi)$ ,  $\psi$  is analytically continuable across  $\partial \mathbb{D}$  at  $\alpha$  and  $|\psi(e^{i\theta})| < 1$ for  $e^{i\theta}$  near to, but not equal to,  $\alpha$ . More detail can be found in [14], where a somewhat larger class of maps is considered. For such  $\alpha$ , the curve  $\psi(e^{i\theta})$ ,  $e^{i\theta}$  near  $\alpha$ , automatically has positive and even order of contact 2m with  $\partial \mathbb{D}$  when  $e^{i\theta} = \alpha$ ; that is,

$$\frac{1-|\psi(e^{i\theta})|^2}{|\psi(\alpha)-\psi(e^{i\theta})|^{2m}}$$

is bounded above and away from 0 for  $e^{i\theta}$  near  $\alpha.$  For  $k\geq 1$  the  $k^{th}$  order data vector

$$D_k(\psi, \alpha) \equiv (\psi(\alpha), \psi'(\alpha), \cdots, \psi^{(k)}(\alpha))$$

makes sense. Given a finite collection  $\psi_1 \cdots, \psi_n$  of such maps and  $k \ge 2$ , we write  $\mathcal{M}_k(\alpha)$  for the set of integers  $j, 1 \le j \le n$ , for which  $F(\psi_j)$  contains  $\alpha$  and the order of contact of  $\psi_j$  at  $\alpha$  is at least k. Further, put

$$\mathcal{D}_k(\alpha) = \{D_k(\psi_j, \alpha) : j \in \mathcal{M}_k(\alpha)\},\$$

the set of possible  $k^{th}$  order data vectors at  $\alpha$  for associated orders of contact at least k. With this notation, Theorem 5.7 in [14] states that for any  $k \geq 2$  and complex constants  $c_1 \cdots, c_n$ ,

(9) 
$$||c_1 C_{\psi_1} + \dots + c_n C_{\psi_n}||_e^2 \ge \sum_{\mathbf{d} \in \mathcal{D}_{k-1}(\alpha)} \left| \sum_{\substack{j \in \mathcal{M}_k(\alpha) \\ D_{k-1}(\psi_j, \alpha) = \mathbf{d}}} c_j \right|^2 \frac{1}{|d_1|},$$

where  $\mathbf{d} = (d_0, d_1, \cdots, d_{k-1}).$ 

For the case k = 2, we need a calculation which appears in the proof of a more delicate version of the inequality (9); see Lemma 5.9 in [14]. Given  $\psi$  as above, with  $\alpha$  in  $F(\psi)$ , convert it into a self-map u of the upper half-plane fixing the origin by  $u = \tau_{\psi(\alpha)} \circ \psi \circ \tau_{\alpha}^{-1}$ . Given our finite collection  $\psi_1 \cdots , \psi_n$ , associate  $u_j$ to  $\psi_j$  in this manner. For D > 0 we write  $\Gamma_{\alpha,D}$  for the locus of the equation  $(1 - |z|^2)/(|\alpha - z|^2) = 4D$ , a circle internally tangent to  $\partial \mathbb{D}$  at  $\alpha$ . We have

(10) 
$$\lim_{\Gamma_{\alpha,D}} \left\| (\overline{c_1}C_{\psi_1}^* + \dots + \overline{c_n}C_{\psi_n}^*) \frac{k_z}{\|k_z\|} \right\|^2 = \sum_{\mathbf{d}\in\mathcal{D}_1(\alpha)} \left\| \sum_{\substack{j\in\mathcal{M}_2(\alpha)\\D_1(\psi_j,\alpha)=\mathbf{d}}} \overline{c_j}k_{w_j}^+ \right\|_{H_2^2}^2$$

where  $k_z$  is the Szego kernel for the Hardy space  $H^2$  in the disk,  $H^2_+$  is the Hardy space of the right half-plane  $\Omega$ ,  $k_w^+(z) = 1/(\overline{w} + z)$  is its reproducing kernel,  $w_j = u'_j(0)/2 - iDu''_j(0)$ , and the limit is taken as  $z \to \alpha$  along  $\Gamma_{\alpha,D}$ . Since  $u''_j(0)$ necessarily has non-negative imaginary part,  $w_j$  is a complex number automatically lying in  $\Omega$ . For further discussion of this circle of ideas, see [14]. We note for future reference that a non-automorphism linear-fractional self-map  $\psi$  of  $\mathbb{D}$  has order of contact two at the unique point in  $F(\psi)$ .

Finally, we need a variant of a result of Berkson [4] and Shapiro and Sundberg [19], which states that if  $\psi_1, \dots, \psi_n$  are distinct analytic self-maps of  $\mathbb{D}$  and  $J(\psi_i) = \{e^{i\theta} : |\psi_i(e^{i\theta})| = 1\}$ , then for any complex constants  $c_1, \dots, c_n$ ,

(11) 
$$\|c_1 C_{\psi_1} + \dots + c_n C_{\psi_n}\|_e^2 \ge \frac{1}{2\pi} \sum_{j=1}^n |c_j|^2 |J(\psi_j)|,$$

where |J| denotes the Lebesgue measure of J; see Exercise 9.3.2 in [8].

**Theorem 1.** Let  $\psi$  be an analytic self-map of  $\mathbb{D}$  such that  $C_{\psi}$  lies in  $C^*(C_{\varphi}, \mathcal{K})$ , where  $\varphi$  is as in (1). If  $F(\psi)$  is empty, then  $C_{\psi}$  is compact.

*Proof.* Suppose that  $C_{\psi}$  lies in  $C^*(C_{\varphi}, \mathcal{K})$  and  $F(\psi)$  is empty. We want to show that  $C_{\psi}$  is compact, or equivalently, that the matrix function

$$\Psi(C_{\psi}) = \left[ \begin{array}{cc} f_2 & f_3 \\ f_4 & f_1 \end{array} \right]$$

is identically zero on [0, s]. Given a small  $\epsilon > 0$  (size to be specified later), there exists B in  $\mathcal{P}$  given by Equation (2) and equivalently by Equation (5), such that  $||C_{\psi} - B|| < \epsilon$ . If we write

$$Y_1 = f(C_{\varphi}^* C_{\varphi}), Y_2 = g(C_{\varphi} C_{\varphi}^*), Y_3 = C_{\varphi} p(C_{\varphi}^* C_{\varphi}), Y_4 = C_{\varphi}^* q(C_{\varphi} C_{\varphi}^*),$$

and  $Y = Y_1 + Y_2 + Y_3 + Y_4$ , it is clear that  $A_k \equiv Y_k \pmod{\mathcal{K}}$  for each *i*, and  $\Psi(Y) = \Psi(A)$ , where  $A = A_1 + A_2 + A_3 + A_4$ . Now using the representation (5) for *B*, we have

$$||C_{\psi} - cC_z - A||_e < \epsilon.$$

Since A is a finite linear combination of composition operators, we see from the inequality (11) that

$$\epsilon^2 > ||C_{\psi} - cC_z - A||_e^2 \ge \frac{|J(\psi)|}{2\pi} + |c|^2.$$

From this we find that  $|c| < \epsilon$ , and since  $\epsilon > 0$  is arbitrary,  $|J(\psi)| = 0$ . In particular we have  $||C_{\psi} - A||_{e} < 2\epsilon$ , hence

$$\left\| \begin{bmatrix} f_2 - g & f_3 - rp \\ f_4 - rq & f_1 - f \end{bmatrix} \right\| = \|\Psi(C_{\psi} - Y)\| = \|C_{\psi} - A\|_e < 2\epsilon.$$

It follows that  $|f_3(t) - \sqrt{t}p(t)| < 2\epsilon$  for  $0 \le t \le s$ , and similarly for the other three matrix entries.

We will show that  $f_3$  vanishes identically on [0, s]. Suppose not, so that its supremum norm  $||f_3||_{\infty}$  is positive. Without loss of generality we may assume  $8\epsilon < ||f_3||_{\infty}$ . It follows that there is a non-degenerate closed subinterval I of [0, s], depending only on  $f_3$  and not containing zero, with  $\sqrt{t}|p(t)| \ge ||f_3||_{\infty}/2$  for t in I. Thus

(12) 
$$\int_{I} |p(t)|^2 dt \ge \frac{\|f_3\|_{\infty}^2 |I|}{4s}$$

We return to this inequality below.

Now we want to apply Equation (10) to the linear combination A, which we write as

(13) 
$$A = c_1 C_{\psi_1} + \cdots + c_m C_{\psi_m}.$$

Recall that the normalized Szego kernel functions  $k_z/||k_z||$  tend to zero weakly as  $|z| \rightarrow 1$ , and so

$$||T||_e = ||T^*||_e \ge \limsup_{|z| \to 1} \left\| T^* \left( \frac{k_z}{||k_z||} \right) \right\|$$

for any bounded operator T on  $H^2$ . The linear-fractional maps  $\psi_1, \dots, \psi_m$  in Equation (13) are taken from the four lists in (6). The maps in each of these lists have a common angular derivative set (a singleton) and a single common first order data vector. For example, the maps  $\psi_i$  from the first list all have  $F(\psi_i) = \{\eta\}$  and first order data vector  $D_1(\psi_i, \eta) = (\eta, 1)$ , which we call  $\mathbf{d}_1$ . The following table summarizes the corresponding information for each of the four lists:

TABLE I			
$\psi_i$ chosen from	$F(\psi_i)$	unique first-order data vector	
$\{(\varphi \circ \sigma)_{n_1}: n_1 \ge 1\}$	$\{\eta\}$	$\mathbf{d}_1 = (\eta, 1)$	
$\{(\sigma \circ \varphi)_{n_2}: n_2 \ge 1\}$	$\{\zeta\}$	$\mathbf{d}_2 = (\zeta, 1)$	
$\{(\varphi \circ \sigma)_{n_3} \circ \varphi: n_3 \ge 0\}$	$\{\zeta\}$	$\mathbf{d}_3 = (\eta, \varphi'(\zeta))$	
$\{(\sigma \circ \varphi)_{n_4} \circ \sigma : n_4 \ge 0\}$	$\{\eta\}$	$\mathbf{d}_4 = (\zeta, \sigma'(\eta))$	

Since  $C^*_{\psi}(k_z) = k_{\psi(z)}$ , our hypothesis that  $F(\psi)$  is empty says exactly that

$$\lim_{|z| \to 1} \left\| C_{\psi}^* \frac{k_z}{\|k_z\|} \right\|^2 = \lim_{|z| \to 1} \frac{1 - |z|^2}{1 - |\psi(z)|^2} = 0$$

(see [8], p.132) and thus

$$4\epsilon^{2} > \|C_{\psi} - A\|_{e}^{2} \geq \limsup_{|z| \to 1} \left\| (C_{\psi}^{*} - A^{*}) \frac{k_{z}}{\|k_{z}\|} \right\|^{2}$$
$$\geq \lim_{\Gamma_{\zeta,D}} \left\| (\overline{c_{1}}C_{\psi_{1}}^{*} + \dots + \overline{c_{m}}C_{\psi_{m}}^{*}) \frac{k_{z}}{\|k_{z}\|} \right\|^{2}.$$

We evaluate the limit on the right via Equation (10) with  $\alpha = \zeta$ . Note that  $\mathcal{D}_1(\zeta) = \{\mathbf{d}_2, \mathbf{d}_3\}$ . Discarding the (necessarily non-negative)  $\mathbf{d}_2$  term from the right-hand side of Equation (10) yields

(14) 
$$4\epsilon^{2} > \lim_{\Gamma_{\zeta,D}} \left\| (\overline{c_{1}}C_{\psi_{1}}^{*} + \dots + \overline{c_{m}}C_{\psi_{m}}^{*}) \frac{k_{z}}{\|k_{z}\|} \right\|^{2} \ge \left\| \sum_{\substack{\zeta \in F(\psi_{i}) \\ D_{1}(\psi_{i},\zeta) = \mathbf{d}_{3}}} \overline{c_{i}}k_{w_{i}}^{+} \right\|_{H^{2}_{+}}^{2}$$

We can relabel  $\psi_1, \dots, \psi_m$  so that the relevant  $\psi'_i s$  occur at the beginning, starting with i = 0. Then the right-hand side of (14) becomes

(15) 
$$\left\|\sum_{k=0}^{n} \overline{c_k} k_{w_k}^{+}\right\|_{H^2_{+}}^{2} = \sum_{i,j=0}^{n} \overline{c_i} c_j \frac{1}{\overline{w_i} + w_j} = \sum_{i,j=0}^{n} \overline{c_i} c_j \int_0^1 t^{\overline{w_i} + w_j - 1} dt$$
$$= \int_0^1 \left|\sum_{k=0}^{n} c_k t^{w_k}\right|^2 t^{-1} dt$$

for appropriate  $n \leq m - 1$ .

Let us evaluate  $c_k$  in terms of the polynomial p occuring in the upper right entry of the matrix function  $\Psi(A) = \Psi(Y)$ . If

$$p(z) = \sum_{k=0}^{n} b_k z^k,$$

we have

$$\begin{split} Y_3 &= C_{\varphi} p(C_{\varphi}^* C_{\varphi}) &\equiv C_{\varphi} p(s C_{\sigma} C_{\varphi}) (\text{mod } \mathcal{K}) \\ &= C_{\varphi} p(s C_{\varphi \circ \sigma}) \\ &= \sum_{k=0}^n b_k s^k C_{(\varphi \circ \sigma)_k \circ \varphi}, \\ &= A_3, \end{split}$$

so that, relabeling if necessary,  $\psi_k = (\varphi \circ \sigma)_k \circ \varphi$  and  $c_k = b_k s^k$  for  $k = 0, 1, \cdots, n$ .

Next we compute  $w_k$  for  $k = 0, 1, \dots, n$ . Let us convert  $\psi_k$  into a self-map  $U_k$ of the upper half-plane fixing the origin as described prior to Equation (10):  $U_k = \tau_\eta \circ \psi_k \circ \tau_\zeta^{-1}$ . We can do the same for the composition factors of  $\psi_k = (\varphi \circ \sigma)_k \circ \varphi$ . The map  $\varphi \circ \sigma$  is a positive parabolic non-automorphism with fixed point  $\eta$ . Let a > 0be its translation number, so that  $\varphi \circ \sigma = \rho_a$ . Thus for  $k \ge 1$ ,  $(\varphi \circ \sigma)_k = \rho_{ka}$ , and its half-plane transplant  $u_k = \tau_\eta \circ \rho_{ka} \circ \tau_\eta^{-1}$  satisfies  $u_k''(0) = 2ika$  by Equation (7). We write v for the half-plane version of  $\varphi : v = \tau_\eta \circ \varphi \circ \tau_\zeta^{-1}$ . We have

$$U_k = \tau_\eta \circ \psi_k \circ \tau_\zeta^{-1} = (\tau_\eta \circ \rho_{ka} \circ \tau_\eta^{-1}) \circ (\tau_\eta \circ \varphi \circ \tau_\zeta^{-1}) = u_k \circ v.$$

Now  $u'_k(0) = \rho'_{ka}(\eta) = 1$  and  $v'(0) = |\varphi'(\zeta)| = \frac{1}{s}$ , and we find

$$U'_k(0) = \frac{1}{s}$$
 and  $U''_k(0) = v''(0) + \frac{2ika}{s^2}$ .

From the discussion following Equation (10) we see that

$$w_k = \frac{1}{2s} - iDv''(0) + k\left(\frac{2aD}{s^2}\right).$$

To this point D has been an arbitrary positive number. Let us choose D so that  $2aD/s^2 = 1$  and put  $\mu = \frac{1}{2s} - iDv''(0)$ , a complex number with positive real part. Thus  $w_k = \mu + k$  and we can write the right hand side of Equation (15) as

$$\int_{0}^{1} \left| \sum_{k=0}^{n} b_{k} s^{k} t^{k} \right|^{2} t^{2\operatorname{Re}\mu-1} dt = \int_{0}^{1} |p(st)|^{2} t^{2\operatorname{Re}\mu-1} dt$$
$$= \frac{1}{s^{2\operatorname{Re}\mu}} \int_{0}^{s} |p(t)|^{2} t^{2\operatorname{Re}\mu-1} dt$$

We consider two cases: if  $2\text{Re}\mu - 1 \ge 0$  then this last integral is at least

$$\frac{t_0^{2\operatorname{Re}\mu-1}}{s^{2\operatorname{Re}\mu}} \int_I |p(t)|^2 dt$$

where  $t_0 > 0$  is the left-hand endpoint of I, and if  $2\text{Re}\mu - 1 < 0$  it is at least

$$\frac{1}{s} \int_{I} |p(t)|^2 dt.$$

For small enough  $\epsilon > 0$ , either case of this inequality, combined with the inequality (14) and Equation (15), is incompatible with the inequality (12), yielding the desired contradiction. It follows that  $f_3 \equiv 0$  on [0, s]. Entirely similar arguments show that  $f_1, f_2$  and  $f_4$  vanish identically on [0, s], hence  $\Psi(C_{\psi}) = 0$ .

With Theorem 1 in hand, we can present our necessary conditions for membership in  $C^*(C_{\omega}, \mathcal{K})$ .

**Theorem 2.** Let  $\varphi$  be as in (1). Suppose  $\psi$  is an analytic self-map of  $\mathbb{D}$  with  $C_{\psi}$  lying in  $C^*(C_{\varphi}, \mathcal{K})$  and  $C_{\psi}$  not compact. Then either  $\psi(z) = z$  or one of the following holds:

(a)  $F(\psi) = \{\zeta\}, \psi(\zeta) = \eta \text{ and } \psi'(\zeta) = \varphi'(\zeta).$ (b)  $F(\psi) = \{\zeta\}, \psi(\zeta) = \zeta \text{ and } \psi'(\zeta) = 1.$ (c)  $F(\psi) = \{\eta\}, \psi(\eta) = \zeta \text{ and } \psi'(\eta) = 1/\varphi'(\zeta).$ (d)  $F(\psi) = \{\eta\}, \psi(\eta) = \eta \text{ and } \psi'(\eta) = 1.$ (e)  $F(\psi) = \{\zeta, \eta\} \text{ with } \psi(\zeta) = \eta, \psi'(\zeta) = \varphi'(\zeta), \psi(\eta) = \eta \text{ and } \psi'(\eta) = 1.$ (f)  $F(\psi) = \{\zeta, \eta\} \text{ with } \psi(\eta) = \zeta, \psi'(\eta) = 1/\varphi'(\zeta), \psi(\zeta) = \zeta \text{ and } \psi'(\zeta) = 1.$ 

*Proof.* If  $\psi$  has no finite angular derivative, then Theorem 1 guarantees that  $C_{\psi}$  is compact. Thus we may assume  $F(\psi)$  is non-empty. We also assume  $\psi$  is not the identity, else there is nothing to prove. If  $C_{\psi}$  is in  $C^*(C_{\varphi}, \mathcal{K})$ , the density of the polynomial subalgebra  $\mathcal{P}$  says that given  $\epsilon$ , we may find a scalar c and a finite linear combination A of composition operators with associated maps from the lists (6) so that

$$\|C_{\psi} - A - cC_z\|_e < \epsilon.$$

As in the beginning of the proof of Theorem 1, we may then conclude that  $|\psi(e^{i\theta})| < 1$  a.e.,  $|c| < \epsilon$ , and that

$$(16) ||C_{\psi} - A|| < 2\epsilon.$$

The self-maps of  $\mathbb{D}$  which define the composition operators in the linear combination A appear among those in Table I above, along with their angular derivative sets (all singletons) and first order data vectors. Suppose that  $\lambda$  is in  $F(\psi)$  and  $D_1(\psi, \lambda) = \mathbf{d}$ . If  $\lambda$  is not in  $\{\zeta, \eta\}$ , then the inequality (8) gives

(17) 
$$||C_{\psi} - A||_{e}^{2} \ge \frac{1}{|\psi'(\lambda)|}$$

contradicting (16). Similarly, if  $\lambda = \zeta$  and **d** is neither **d**<sub>2</sub> nor **d**<sub>3</sub> from Table I, or if  $\lambda = \eta$  and **d** is neither **d**<sub>1</sub> nor **d**<sub>4</sub>, the inequality (8) and Table I again imply (17). It follows that if  $F(\psi)$  is a singleton, one of conditions (a)-(d) must hold.

The remainder of the proof considers the possibility that  $F(\psi) = \{\zeta, \eta\}$ . The Julia-Caratheodory theory says a non-identity analytic self-map of  $\mathbb{D}$  cannot have fixed points at distinct points  $\zeta, \eta$  in  $\partial \mathbb{D}$  with derivative 1 at each. If we have both  $\psi(\zeta) = \eta, \psi'(\zeta) = \varphi'(\zeta)$  and  $\psi(\eta) = \zeta, \psi'(\eta) = 1/\varphi'(\zeta)$ , then  $\psi \circ \psi$  fixes both  $\zeta$  and  $\eta$  with derivative 1 at each, so that  $\psi \circ \psi$  is the identity map, contradicting the observation above that  $|\psi(e^{i\theta})| < 1$  almost everywhere. Thus if  $F(\psi) = \{\zeta, \eta\}$ , either (e) or (f) must hold, completing the proof.

We will see in Section 5 that there are indeed maps  $\psi$  of types (e) and (f) in Theorem 2 for which  $C_{\psi}$  belongs to  $C^*(C_{\varphi}, \mathcal{K})$ .

### 3. The $C^*$ -algebra induced by parabolic non-automorphisms

Let us write  $\mathcal{B}(H^2)$  for the algebra of bounded operators on  $H^2$ . A bounded operator T on  $H^2$  is essentially normal if T and  $T^*$  commute modulo  $\mathcal{K}$ ; normal operators and compact operators give trivial examples of essentially normal operators. The only normal composition operators  $C_{\psi}$  are those of the form  $\psi(z) = az, |a| \leq 1$ . Bourdon, Levi, Narayan, and Shapiro [1] showed that if  $\psi$  is linear-fractional with  $\|\psi\|_{\infty} = 1$  and not a rotation, then  $C_{\psi}$  is essentially normal exactly when  $\psi$  is a parabolic non-automorphism. Let us select a point  $\gamma$  in  $\partial \mathbb{D}$  and consider the set  $\{\rho_a : \text{Re } a > 0\}$  of all parabolic non-automorphisms fixing  $\gamma$ . Here, as earlier, a is the translation number for  $\rho_a$ . Any two of the maps  $\rho_a$  commute under composition and in fact  $\rho_a \circ \rho_b = \rho_{a+b}$ , so  $C_{\rho_a}$  and  $C_{\rho_b}$  commute. One can easily check that the Krein adjoint of  $\rho_a$  is  $\rho_{\overline{a}}$ . Since  $\rho'_a(\gamma) = 1$ , it follows from Equation (4) that  $C^*_{\rho_a} = C_{\rho_{\overline{a}}} + K$  for some compact operator K. A recent theorem of Montes-Rodríguez, Ponce-Escudero and Shkarin [18] shows that  $C_{\rho_a}$  is irreducible. Moreover  $C^*(C_{\rho_a})$ , the unital  $C^*$ -algebra generated by  $C_{\rho_a}$ , contains the commutator of  $C_{\rho_a}$  and  $C^*_{\rho_a}$  which we know is compact but non-zero. Thus  $C^*(C_{\rho_a})$  contains  $\mathcal{K}$  and  $C^*(C_{\rho_a})/\mathcal{\tilde{K}}$  is commutative. Now let  $\mathbb{P}_{\gamma}$  denote the set of all composition operators  $C_{\rho}$ , where  $\rho$ , fixing  $\gamma$ , ranges over  $\{\rho_a : \text{Re } a > 0\}$ . We write  $C^*(\mathbb{P}_{\gamma})$  for the unital C<sup>\*</sup>-algebra generated by the operators in  $\mathbb{P}_{\gamma}$ . Clearly  $C^*(\mathbb{P}_{\gamma})$  contains  $\mathcal{K}$ , and, by the above remarks,  $C^*(\mathbb{P}_{\gamma})/\mathcal{K}$  is also commutative. In this section we compute and apply the Gelfand representation of this quotient algebra.

We begin with two lemmas.

**Lemma 1.** For a > 0 there is an operator  $A \ge 0$  and a compact operator K with  $C_{\rho_a} = A + K$ .

Proof. Then

$$C_{\rho_{a/2}} = \frac{1}{2} (C_{\rho_{a/2}} + C^*_{\rho_{a/2}}) + \frac{1}{2} (C_{\rho_{a/2}} - C^*_{\rho_{a/2}})$$
  
$$\equiv B + J$$

where B is self-adjoint and J is compact. Thus  $C_{\rho_a} = C_{\rho_{a/2}}C_{\rho_{a/2}} = (B+J)^2 = B^2 + (BJ + JB + J^2)$ . Since  $B^2$  is positive and  $BJ + JB + J^2$  is compact, we are done.

**Lemma 2.** Let a, b be positive with b/a = m/n, with m and n positive integers. Suppose  $0 < \lambda \leq 1$  and there is a sequence  $f_k$  of unit vectors in  $H^2$  converging weakly to zero such that

$$\|(C_{\rho_a} - \lambda)f_k\| \to 0.$$

Then

$$\|(C_{\rho_b} - \lambda^{m/n})f_k\| \to 0.$$

*Proof.* First observe that

$$C_{\rho_a}^m - \lambda^m = [C_{\rho_a}^{m-1} + \lambda C_{\rho_a}^{m-2} + \dots + \lambda^{m-2} C_{\rho_a} + \lambda^{m-1}][C_{\rho_a} - \lambda]$$

$$(C_{\rho_a}^m - \lambda^m) f \parallel \lambda 0 \text{ or } h \to \infty \text{ Since } C_{\rho_a}^m - C_{\rho_a}^n$$

In particular,  $\|(C^m_{\rho_a} - \lambda^m)f_k\| \to 0$  as  $k \to \infty$ . Since  $C^m_{\rho_a} = C^n_{\rho_b}$ ,  $\|(C^n_{\rho_b} - \lambda^m)f_k\| \to 0.$ 

Also note that we may factor  $C_{\rho_b}^n - \lambda^m = C_{\rho_b}^n - (\lambda^{m/n})^n$  as

$$[C_{\rho_b}^{n-1} + \lambda^{m/n} C_{\rho_b}^{n-2} + \dots + (\lambda^{m/n})^{n-2} C_{\rho_b} + (\lambda^{m/n})^{n-1}][C_{\rho_b} - \lambda^{m/n}]$$

Apply Lemma 1 to  $C_{\rho_b}$  to write

 $C_{\rho_b}^{n-1} + \lambda^{m/n} C_{\rho_b}^{n-2} + \dots + (\lambda^{m/n})^{n-2} C_{\rho_b} + (\lambda^{m/n})^{n-1} I = T + (\lambda^{m/n})^{n-1} I + K$ for some positive T and compact K. We have

(18) 
$$\| (C_{\rho_b}^n - \lambda^m) f_k \| = \| (T + (\lambda^{m/n})^{n-1} + K) (C_{\rho_b} - \lambda^{m/n}) f_k \|$$

Since K is compact,  $||K(C_{\rho_b} - \lambda^{m/n})f_k|| \to 0$  as  $k \to \infty$ . Since the left-hand side of Equation (18) goes to 0 as  $k \to \infty$ , we see, writing  $c = (\lambda^{m/n})^{n-1}$  that

$$\|(T+cI)(C_{\rho_b}-\lambda^{m/n})f_k\|\to 0.$$

But  $||(T+cI)(C_{\rho_b}-\lambda^{m/n})f_k||^2$  is equal to

$$||T(C_{\rho_b} - \lambda^{m/n})f_k||^2 + 2c\langle T(C_{\rho_b} - \lambda^{m/n})f_k, (C_{\rho_b} - \lambda^{m/n})f_k \rangle + c^2 ||(C_{\rho_b} - \lambda^{m/n})f_k||^2 \ge c^2 ||(C_{\rho_b} - \lambda^{m/n})f_k||^2$$

where the last inequality uses the positivity of T, so that  $\|(C_{\rho_b} - \lambda^{m/n})f_k\| \to 0$ .  $\Box$ 

The essential spectrum  $\sigma_e(T)$  of a bounded operator on  $H^2$  is by definition the spectrum of the coset [T] in  $\mathcal{B}(H^2)/\mathcal{K}$ . We recall from [15] that if a > 0,  $\sigma_e(C_{\rho_a}) = [0, 1]$ . We will need the notion of joint essential spectrum, which is treated by Dash in [9]. If Re a > 0, the coset  $[C_{\rho_a}]$  of  $C_{\rho_a}$  modulo  $\mathcal{K}$  will also be denoted by  $x_a$ . By either Lemma 1 or the discussion preceeding it,  $x_a$  is self-adjoint. Given a and b, the joint essential spectrum  $\sigma_e(C_{\rho_a}, C_{\rho_b})$  is defined to be the joint spectrum  $\sigma(x_a, x_b)$  of the pair  $x_a, x_b$  in the Calkin algebra  $\mathcal{B}(H^2)/\mathcal{K}$ . This set coincides with the joint spectrum in the commutative unital subalgebra  $C^*(x_a, x_b)$  generated by  $x_a$  and  $x_b$ . If  $\mathcal{M}$  is the maximal ideal space of this algebra, and  $\widehat{}$  denotes the Gelfand transform, then the map  $\ell \mapsto (\widehat{x_a}(\ell), \widehat{x_b}(\ell))$  is a homeomorphism of  $\mathcal{M}$  onto  $\sigma(x_a, x_b)$ . Let us assume that a and b are positive. A theorem of Dash [9] states, in this context, using  $C^*_{\rho_a} \equiv C_{\rho_a} \pmod{\mathcal{K}}$  and similarly for  $C_{\rho_b}$ , that  $(\lambda, \mu)$  lies in  $\sigma_e(C_{\rho_a}, C_{\rho_b})$  if and only if there exists a sequence  $\{f_k\}$  of unit vectors in  $H^2$ , converging weakly to zero, such that  $\|(C_{\rho_a} - \lambda)f_k\|$  and  $\|(C_{\rho_b} - \mu)f_k\|$  both tend to zero as  $k \to \infty$ .

# **Corollary 1.** Suppose that a, b are positive and b/a is rational. Then

$$\sigma_e(C_{\rho_a}, C_{\rho_b}) = \{ (t^a, t^b) : 0 \le t \le 1 \}.$$

*Proof.* We know that

 $\sigma_e(C_{\rho_a}, C_{\rho_b}) \subset \sigma_e(C_{\rho_a}) \times \sigma_e(C_{\rho_b}) = [0, 1] \times [0, 1].$ 

Let  $0 < \lambda \leq 1$ . Since  $\lambda \in \sigma_e(C_{\rho_a})$ , we may find unit vectors  $f_k$  with  $f_k \to 0$ weakly and  $||(C_{\rho_a} - \lambda)f_k|| \to 0$ . By Lemma 2,  $||(C_{\rho_b} - \lambda^{b/a})f_k|| \to 0$  as well, so that  $(\lambda, \lambda^{b/a})$  is in  $\sigma_e(C_{\rho_a}, C_{\rho_b})$  by Dash's theorem. Setting  $t^a = \lambda$  we have  $\{(t^a, t^b) : 0 < t \leq 1\} \subset \sigma_e(C_{\rho_a}, C_{\rho_b})$ . The set on the right is compact in  $\mathbb{R}^2$ , so it contains (0,0) as well, giving  $\{(t^a, t^b) : 0 \leq t \leq 1\} \subset \sigma_e(C_{\rho_a}, C_{\rho_b})$ . Conversely, if  $(\lambda, \mu) \in \sigma_e(C_{\rho_a}, C_{\rho_b})$ , Dash's theorem gives the existence of a sequence of unit vectors  $f_k$  converging weakly to 0 with both  $||(C_{\rho_a} - \lambda)f_k||$  and  $||(C_{\rho_b} - \mu)f_k||$  tending to zero as  $k \to \infty$ . If  $\lambda > 0$ ,  $||(C_{\rho_b} - \lambda^{b/a})f_k|| \to 0$  by Lemma 2. Thus  $\mu = \lambda^{b/a}$  and  $(\lambda, \mu) = (\lambda, \lambda^{b/a}) = (t^a, t^b)$  for some  $t, 0 \leq t \leq 1$ . If  $\mu > 0$ , the symmetric result says  $\lambda = \mu^{a/b} > 0$ , and, putting  $\mu = t^b$ ,  $(\lambda, \mu) = (\mu^{a/b}, \mu) = (t^a, t^b)$ , again of the desired form. As for (0, 0), we already know it lies in  $\sigma_e(C_{\rho_a}, C_{\rho_b})$ , and of course it has the form  $(0^a, 0^b)$ . We will need the fact that on the domain  $\Omega = \{a : \text{Re } a > 0\}$  the map  $a \mapsto C_{\rho_a}$  is a holomorphic function of a in the operator norm topology; see for example the discussion in the proof of Theorem 6.1 in [6]. We continue to denote the coset of  $C_{\rho_a}$  by  $x_a$  and to keep in mind that when a > 0,  $\sigma(x_a) = [0, 1]$ .

**Theorem 3.** There is a unique \*-isomorphism  $\Gamma : C([0,1]) \to C^*(\mathbb{P}_{\gamma})/\mathcal{K}$  such that  $\Gamma(t^a) = [C_{\rho_a}]$  for all  $a \in \Omega$ .

Proof. First consider a = 1 and  $x_1 = [C_{\rho_1}]$ . Since  $\sigma(x_1) = [0, 1]$  we may define a \*-isomorphism  $\Gamma : C([0, 1]) \mapsto C^*(x_1)$  by sending p to  $p(x_1)$  for any polynomial p. Fix any rational number r > 0. By Corollary 1,

$$\sigma(x_1, x_r) = \sigma_e(C_{\rho_1}, C_{\rho_r}) = \{(t, t^r) : 0 \le t \le 1\}.$$

The map  $p(x_1, x_r) \mapsto p(z_1, z_2)$ , where p is a two-variable polynomial, extends to a unique \*-isomorphism of  $C^*(x_1, x_r)$  onto  $C(\sigma(x_1, x_r))$ . Since  $\sigma(x_1, x_r)$  is homeomorphic to [0, 1] via the map  $t \mapsto (t, t^r)$ , we see that  $p(x_1, x_r) \mapsto p(t, t^r)$  defines a \*-isomorphism of  $C^*(x_1, x_r)$  onto C([0, 1]). Let  $\tilde{\Gamma}$  denote the inverse of this map, that is  $\tilde{\Gamma} : p(t, t^r) \mapsto p(x_1, x_r)$ . Since polynomials in t span C([0, 1]),  $x_1$  generates the  $C^*$ -algebra  $C^*(x_1, x_r)$  and  $C^*(x_1) = C^*(x_1, x_r)$ . It follows that  $\tilde{\Gamma} = \Gamma$ . Since ris arbitrary in the set  $\mathbb{Q}_+$  of positive rationals, we have shown that

$$C^*(\{x_r : r \in \mathbb{Q}_+\}) = C^*(x_1).$$

Moreover,  $\Gamma(t^r) = x_r, r \in \mathbb{Q}_+$ . It is easy to see that the map  $a \mapsto t^a$  is a normholomorphic map of the right half plane into C([0,1]) and thus that  $a \mapsto \Gamma(t^a)$  is norm-holomorphic from the right half-lane to  $\mathcal{B}(H^2)/\mathcal{K}$ , as is the function  $a \mapsto x_a$ . We have seen that these functions agree on  $\mathbb{Q}_+$ , hence they must agree on the right half-plane  $\Omega$ .

We record three immediate consequences.

**Corollary 2.** If  $a_1, \dots, a_n$  lie in the right half-plane  $\Omega$ , then

$$\sigma_e(C_{\rho_{a_1}}, \cdots, C_{\rho_{a_n}}) = \{(t^{a_1}, \cdots, t^{a_n}) : 0 \le t \le 1\}.$$

**Corollary 3.** If  $\rho$  is a parabolic non-automorphism fixing  $\gamma$ , then  $C^*(C_{\rho}) = C^*(\mathbb{P}_{\gamma})$ .

**Corollary 4.** If  $\varphi$  is as in (1), then  $\mathbb{P}_{\zeta}$  and  $\mathbb{P}_{\eta}$  are both subsets of  $C^*(C_{\varphi}, \mathcal{K})$ .

# 4. LINEAR-FRACTIONAL MAPS

The goal of this section is to find all linear-fractional  $\psi$  with  $C_{\psi}$  in  $C^*(C_{\varphi}, \mathcal{K})$ , where  $\varphi$  satisfies the conditions of (1). Since  $C_{\psi}$  is compact if  $\|\psi\|_{\infty} < 1$ , our interest is in the case  $\|\psi\|_{\infty} = 1$ .

**Lemma 3.** If  $\varphi$  is as in (1),  $C^*(C_{\varphi}, \mathcal{K})$  contains  $C_{\psi}$  for all linear-fractional  $\psi$ :  $\mathbb{D} \to \mathbb{D}$  with  $\psi(\zeta) = \eta$ ,  $\psi'(\zeta) = \varphi'(\zeta)$  and  $\psi(\mathbb{D})$  properly contained in  $\varphi(\mathbb{D})$ .

Proof. Set  $\tau = \varphi^{-1} \circ \psi$ , noting that the hypothesis  $\psi(D) \subset \varphi(\mathbb{D})$  means that  $\tau$  is well-defined. Since this containment is proper, and  $\tau'(\zeta) = 1$ ,  $\tau$  is a parabolic nonautomorphism with fixed point  $\zeta$ . Since  $\varphi \circ \tau = \psi$ ,  $C_{\psi} = C_{\tau}C_{\varphi}$ . By Corollary 4,  $C_{\tau} \in C^*(C_{\varphi}, \mathcal{K})$ , from which the conclusion follows.  $\Box$ 

Now consider the parabolic non-automorphism  $\rho = \varphi \circ \sigma$ . The unique fixed point for  $\rho$  and its iterates  $(\rho)_n$  is  $\eta$ . Fix an integer  $n \geq 1$  and let  $\varphi_1 = (\rho)_n \circ \varphi$ . Clearly  $\varphi_1(\mathbb{D})$  is properly contained in  $\varphi(\mathbb{D})$ . Note that  $\varphi_1(\zeta) = \eta$  and, since  $\rho'(\eta) = 1, \varphi'_1(\zeta) = \varphi'(\zeta)$ . It follows from Lemma 3 that  $C^*(C_{\varphi_1}, \mathcal{K})$  is contained in  $C^*(C_{\varphi}, \mathcal{K})$ . Let *i* denote the corresponding inclusion map. Since  $i(\mathcal{K}) = \mathcal{K}$ , *i* induces a \*-homomorphism

$$\hat{i}: C^*(C_{\varphi_1}, \mathcal{K})/\mathcal{K} \to C^*(C_{\varphi}, \mathcal{K})/\mathcal{K}$$

given by  $\hat{i}([T]) = [T]$ , where [T] denotes the coset, modulo  $\mathcal{K}$  of the operator T. Note that  $\hat{i}$  is itself an inclusion. Also observe that the map  $\Psi : C^*(C_{\varphi}, \mathcal{K}) \to \mathcal{D}$ induces a \*-isomorphism  $\Phi : C^*(C_{\varphi}, \mathcal{K})/\mathcal{K} \to \mathcal{D}$  given by  $\Phi([T]) = \Psi(T)$ . Let  $\Phi_1$ denote the corresponding \*-isomorphism  $\Phi_1 : C^*(C_{\varphi_1}, \mathcal{K})/\mathcal{K} \to \mathcal{D}$ . Keep in mind that  $\Phi_1$  should be defined by  $\Phi_1([T]) = \Psi_1(T)$ , where  $\Psi_1 : C^*(C_{\varphi_1}, \mathcal{K}) \to \mathcal{D}$  is associated to  $\varphi_1$  as  $\Psi$  is associated to  $\varphi$ . Thus if B in  $C^*(C_{\varphi_1}, \mathcal{K})$  is given by (2), but with  $\varphi$  replaced by  $\varphi_1$ , then  $\Psi_1(B)$  is given by (3). We have a commutative diagram

where  $\Lambda = \Phi \circ \hat{i} \circ \Phi_1^{-1}$ . We seek to identify  $\Lambda$  explicitly.

**Lemma 4.** For any element F in  $\mathcal{D}$ ,

(19) 
$$(\Lambda F)(t) = F(t^{2n+1}/s^{2n}), \ 0 \le t \le s.$$

*Proof.* For the purposes of the proof, we use  $\Lambda$  to denote the map given by formula (19), and then show, with this redefinition, that it coincides with  $\Phi \circ \hat{i} \circ \Phi_1^{-1}$ , that is, that  $\Lambda \circ \Phi_1 = \Phi \circ \hat{i}$ . Recall that  $C_{\varphi}^* \equiv sC_{\sigma} \pmod{\mathcal{K}}$  so that

$$C_{\varphi_1} = C_{\varphi} C_{(\rho)_n} = C_{\varphi} (C_{\sigma} C_{\varphi})^n$$
$$\equiv \frac{1}{s^n} C_{\varphi} (C_{\varphi}^* C_{\varphi})^n \pmod{\mathcal{K}},$$

and, taking adjoints,  $C_{\varphi_1}^* \equiv \frac{1}{s^n} C_{\varphi}^* (C_{\varphi} C_{\varphi}^*)^n$  modulo the compacts. Calculations using these two facts show that if we write  $y = [C_{\varphi_1}]$  and  $x = [C_{\varphi}]$ , we have, for each non-negative integer m,

$$(y^*y)^m = \frac{1}{s^{2nm}} (x^*x)^{(2n+1)m},$$
  

$$(yy^*)^m = \frac{1}{s^{2nm}} (xx^*)^{(2n+1)m}.$$
  

$$y(y^*y)^m = \frac{1}{s^{(2m+1)n}} x(x^*x)^{(2n+1)m+n}.$$

and

$$y^*(yy^*)^m = \frac{1}{s^{(2m+1)n}} x^*(xx^*)^{(2n+1)m+n}$$

The left-hand sides in these four equations are elements in  $C^*(C_{\varphi_1}, \mathcal{K})/\mathcal{K}$ , while the right-hand sides represent the same objects as elements of  $C^*(C_{\varphi}, \mathcal{K})/\mathcal{K}$ . We first act on  $y(y^*y)^m$  by  $\hat{i}$ , followed by  $\Phi$ . We then act on  $y(y^*y)^m$  by  $\Phi_1$ , followed by  $\Lambda$  (as defined by Equation (19)). As the reader can see from the following picture, we end up with a common result, the matrix function in the lower right-hand corner.

$$\begin{array}{cccc} y(y^*y)^m & & \stackrel{\hat{i}}{\longrightarrow} & \frac{1}{s^{(2m+1)n}} x(x^*x)^{(2n+1)m+n} \\ & \Phi_1 \\ & & & & \downarrow \Phi \\ \begin{bmatrix} 0 & \sqrt{t} \ t^m \\ 0 & 0 \end{bmatrix} & \stackrel{\Lambda}{\longrightarrow} & \begin{bmatrix} 0 & \frac{\sqrt{t} \ t^{(2n+1)m+n}}{s^{(2m+1)n}} \\ 0 & 0 \end{bmatrix} \end{array}$$

One can check that we also arrive at common values when  $\Lambda \circ \Phi_1$  and  $\Phi \circ i$  act on  $(y^*y)^m$ , and similarly for  $(yy^*)^m$  and  $y^*(yy^*)^m$ . Since elements of the form  $(y^*y)^m$ ,  $(yy^*)^m$ ,  $y(y^*y)^m$  and  $y^*(yy^*)^m$ , together with the identity, span  $C^*(C_{\varphi_1}, \mathcal{K})/\mathcal{K}$ , we have  $\Lambda \circ \Phi_1 = \Phi \circ \hat{i}$  as desired

It is clear from (19) that  $\Lambda$  is an automorphism of  $\mathcal{D}$ . It follows that i is an isomorphism and thus that i has range equal to all of  $C^*(C_{\varphi}, \mathcal{K})$ , that is,

(20) 
$$C^*(C_{\varphi_1},\mathcal{K}) = C^*(C_{\varphi},\mathcal{K}).$$

More generally, we have the following result.

**Theorem 4.** Let  $\psi$  be a linear-fractional map of  $\mathbb{D}$ , not an automorphism, with  $\psi(\zeta) = \varphi(\zeta)$  and  $\psi'(\zeta) = \varphi'(\zeta)$ , where  $\varphi$  is as in (1). Then  $C^*(C_{\psi}, \mathcal{K}) = C^*(C_{\varphi}, \mathcal{K})$ .

*Proof.* The circles  $\varphi(\partial \mathbb{D})$  and  $\psi(\partial \mathbb{D})$  are both internally tangent to  $\partial \mathbb{D}$  at  $\eta$ . If  $\varphi(\mathbb{D})$  is a proper subset of  $\psi(\mathbb{D})$ , then

(21) 
$$C^*(C_{\varphi}, \mathcal{K}) \subset C^*(C_{\psi}, \mathcal{K})$$

by Lemma 3. Suppose on the other hand that  $\psi(\mathbb{D}) \subset \varphi(\mathbb{D})$ . If *a* is the (necessarily positive) translation number of the parabolic map  $\varphi \circ \sigma$  (so that  $\varphi \circ \sigma = \rho_a$  in the terminology of Section 2), then  $(\varphi \circ \sigma)_n = \rho_{na}$ , and the radius of the disk  $\rho_{na}(\mathbb{D})$  shrinks to zero as  $n \to \infty$ . Thus there exists *n* with  $\rho_{na}(\mathbb{D})$  properly contained in  $\psi(D)$ . If  $\varphi_1 = (\varphi \circ \sigma)_n \circ \varphi = \rho_{na} \circ \varphi$ , then  $\varphi_1(\mathbb{D})$  is also properly contained in  $\psi(\mathbb{D})$ . Since  $\varphi_1(\zeta) = \eta = \psi(\zeta)$  and  $\varphi'_1(\zeta) = \rho'_{na}(\eta)\varphi'(\zeta) = \varphi'(\zeta) = \psi'(\zeta)$ , Lemma 3 implies that  $C^*(C_{\psi}, \mathcal{K})$  contains  $C^*(C_{\varphi_1}, \mathcal{K})$ , which by (20) coincides with  $C^*(C_{\varphi}, \mathcal{K})$ . The result is that (21) holds, whatever the relationship between the disks  $\varphi(\mathbb{D})$  and  $\psi(\mathbb{D})$ . The statement of the theorem is symmetric in  $\varphi$  and  $\psi$ , so symmetry implies that the containment reverse to that in (21) also holds, completing the proof.  $\Box$ 

**Theorem 5.** Suppose  $\varphi$  is as in (1). Let  $\psi$ , not the identity, be any linear-fractional self-map of  $\mathbb{D}$  with  $\|\psi\|_{\infty} = 1$ . Then  $C_{\psi}$  is in  $C^*(C_{\varphi}, \mathcal{K})$  if and only if  $\psi$  is not an automorphism and one of the following conditions holds:

- (a)  $\psi(\zeta) = \eta$  and  $\psi'(\zeta) = \varphi'(\zeta)$ .
- (b)  $\psi(\zeta) = \zeta$  and  $\psi'(\zeta) = 1$ .
- (c)  $\psi(\eta) = \zeta$  and  $\psi'(\eta) = 1/\varphi'(\zeta)$ .
- (d)  $\psi(\eta) = \eta$  and  $\psi'(\eta) = 1$ .

Proof. The "only if" statement follows immediately from Theorem 2 and the hypothesis that  $\psi$  is linear-fractional. Conversely, let  $\psi$  be a linear-fractional map which is not an automorphism. If  $\psi$  is parabolic with fixed point at either  $\zeta$  or  $\eta$ , the result follows from Corollary 4; this handles the cases (b) and (d). If  $\psi$  is as in (a), then we have  $C_{\psi} \in C^*(C_{\varphi}, \mathcal{K})$  by Theorem 4. Finally, if  $\psi$  satisfies condition (c), then its Krein adjoint  $\sigma_{\psi}$  is a linear-fractional self-map of  $\mathbb{D}$ , not an automorphism, which satisfies condition (a), so that  $C_{\sigma_{\psi}} \in C^*(C_{\varphi}, \mathcal{K})$ . Since  $C^*_{\psi} \equiv sC_{\sigma_{\psi}}$  modulo the compacts, this completes the argument.

The maps satisfying (a)-(d) in Theorem 5 can be described more explicitly. Given a point  $\gamma$  on  $\partial \mathbb{D}$ , let us write  $\rho_{\gamma,a}$  for the unique parabolic map fixing  $\gamma$  with translation number a. This will be a self-map of  $\mathbb{D}$  when Re  $a \geq 0$ , but when Re a < 0,  $\rho_{\gamma,a}$  takes  $\mathbb{D}$  onto a larger disk, whose boundary is externally tangent to  $\partial \mathbb{D}$  at  $\gamma$ . Clearly, the linear-fractional non-automorphisms of  $\mathbb{D}$  satisfying (b) or (d) are, respectively,  $\rho_{\zeta,a}$  or  $\rho_{\eta,a}$  with Re a > 0. Imaginary a gives an automorphism of  $\mathbb{D}$ , but in this case  $C^*(C_{\varphi}, \mathcal{K})$  does not contain the corresponding composition operator.

Consider now a linear-fractional non-automorphism  $\psi$  of  $\mathbb{D}$  satisfying (a). If  $\psi(\mathbb{D}) \subset \varphi(\mathbb{D})$ , we can define  $\rho = \psi \circ \varphi^{-1}$  which fixes  $\eta$  and carries  $\mathbb{D}$  to  $\mathbb{D}$ . Moreover,  $\rho'(\eta) = \psi'(\zeta)/\varphi'(\zeta) = 1$ , so  $\rho$  is parabolic; say  $\rho = \rho_{\eta,a}$  where Re  $a \ge 0$ , and we find  $\psi = \rho_{\eta,a} \circ \varphi$ . On the other hand, if  $\psi$  satisfies (a) and  $\varphi(\mathbb{D})$  is a proper subset of  $\psi(\mathbb{D})$ , we put  $\rho = \varphi \circ \psi^{-1}$  which is again a parabolic self-map of  $\mathbb{D}$ ; this time a non-automorphism. Thus  $\rho = \rho_{\eta,a}$  with Re a > 0 and we find  $\psi = \rho_{\eta,a}^{-1} \circ \varphi = \rho_{\eta,-a} \circ \varphi$ . If b is the unique positive number with  $\rho_{\eta,b}(\mathbb{D}) = \varphi(\mathbb{D})$ , then  $\rho_{\eta,-a} \circ \varphi$  is a non-automorphism self-map of  $\mathbb{D}$  exactly when Re a < b. Rephrasing and summarizing, we conclude that the non-automorphisms  $\psi$  of  $\mathbb{D}$  satisfying (a) are precisely the maps of the form  $\rho_{\eta,a} \circ \varphi$ , Re a > -b. Similarly, if c is the unique positive number with  $\rho_{\zeta,c}(\mathbb{D}) = \sigma(\mathbb{D})$ , then the non-automorphisms  $\psi$  satisfying (c) are exactly the maps of the form  $\psi = \rho_{\zeta,a} \circ \sigma$  with Re a > -c. The next result shows that the positive translation numbers b and c are nicely related to each other, and to the translation numbers of the positive parabolic non-automorphisms  $\varphi \circ \sigma$  and  $\sigma \circ \varphi$ .

**Theorem 6.** Let b and c be the unique positive numbers with  $\rho_{\eta,b}(\mathbb{D}) = \varphi(\mathbb{D})$  and  $\rho_{\zeta,c}(\mathbb{D}) = \sigma(\mathbb{D})$ , respectively. We have  $c = |\varphi'(\zeta)|b$ , and moreover,  $\varphi \circ \sigma = \rho_{\eta,2b}$  and  $\sigma \circ \varphi = \rho_{\zeta,2c}$ .

*Proof.* Clearly there is no loss of generality in assuming that  $\zeta = 1$ . The non-affine linear-fractional self-maps of  $\mathbb{D}$  which send 1 to  $\eta \in \partial \mathbb{D}$  can be written in the form

$$\varphi(z) = \eta \ \frac{(1+s+sd)z + (d-s-sd)}{z+d}$$

where  $s = |\varphi'(1)|$  and Re  $\frac{d-1}{d+1} \ge s$  (see [3]). A computation shows that  $\varphi'(1) = \eta s$ and  $\varphi''(1) = -2\eta s/(1+d)$ . The image of the unit circle under  $\varphi$  is a circle with curvature  $\kappa_1 = |\varphi'(1)|^{-1} \text{Re } [1 + \varphi''(1)/\varphi'(1)] = \frac{1}{s} \text{Re } [1 - \frac{2}{1+d}]$ . Since

$$\sigma(z) = \frac{\overline{\eta}(1+s+s\overline{d})z-1}{-\overline{\eta}(\overline{d}-s-s\overline{d})z+\overline{d}}$$

we find that  $\sigma'(\eta) = \overline{\eta}/s$  and

$$\sigma''(\eta) = \frac{2\overline{\eta}^2(\overline{d} - s - s\overline{d})}{s^2(1 + \overline{d})},$$

so that the image of the unit circle under  $\sigma$  is a circle with curvature

$$\kappa_2 = s \operatorname{Re} \left\{ 1 + \eta \; \frac{2\overline{\eta}^2 (\overline{d} - s - s\overline{d})}{s^2 (1 + \overline{d})} \cdot \frac{s}{\overline{\eta}} \right\} = 2 - 2 \operatorname{Re} \; \frac{1}{1 + d} - s$$

A positive parabolic non-automorphism fixing 1 and corresponding to translation by a has the form ((2 + d)z - 1)/(z + d) where a = -2/(d + 1); by the above calculations the image of the unit circle under this map has curvature 1 + Re a. Thus, if  $\varphi(1) = \eta$  and  $|\varphi'(1)| = s$ , the unique positive value b such that the curvature of  $\varphi(\partial \mathbb{D})$  is equal to the curvature of the image of  $\partial \mathbb{D}$  under the positive parabolic map which corresponds to translation by b > 0 satisfies

$$1 + b = \frac{1}{s} \operatorname{Re} \left( 1 - \frac{2}{1+d} \right);$$

that is,

$$b = \frac{1}{s} \operatorname{Re}\left(1 - \frac{2}{1+d}\right) - 1.$$

Similarly, the curvature of the circle  $\sigma(\partial \mathbb{D})$  is equal to the curvature of the circle which is the image of the unit circle under the positive parabolic non-automorphism corresponding to translation by c precisely when  $c = 1 - s - 2\operatorname{Re}\frac{1}{1+d}$ . Thus c = sb. This conclusion also holds when  $\varphi$  is an affine map,  $\varphi(z) = \eta(sz + 1 - s)$ , where the computations are easier.

For the final statement, let  $\psi = \rho_{\eta,-b} \circ \varphi$ , so that  $\psi$  is an automorphism of  $\mathbb{D}$ and  $\varphi = \rho_{\eta,b} \circ \psi$ . Since the Krein adjoint of an automorphism is its inverse, we have

$$\sigma = \sigma_{\varphi} = \sigma_{\psi} \circ \sigma_{\rho_{\eta,b}} = \psi^{-1} \circ \rho_{\eta,\overline{b}} = \psi^{-1} \circ \rho_{\eta,b}$$

and thus  $\varphi \circ \sigma = \rho_{\eta,b} \circ \psi \circ \psi^{-1} \circ \rho_{\eta,b} = \rho_{\eta,2b}$ . Similarly,  $\sigma \circ \varphi = \rho_{1,2c} = \rho_{\zeta,2c}$ .  $\Box$ 

The remarks preceeding Theorem 6 express the linear-fractional maps  $\psi$  with  $C_{\psi}$  belonging to  $C^*(C_{\varphi}, \mathcal{K})$  in terms of  $\varphi, \sigma, \rho_{\eta,a}$  and  $\rho_{\zeta,a}$  for appropriate ranges of the translation numbers a. We describe below a corresponding operator-theoretic description of  $C_{\psi}$  modulo  $\mathcal{K}$ , in terms of the polar factors of  $C_{\varphi}$  and  $C_{\varphi}^*$ . In [15] it was shown that every operator B in  $C^*(C_{\varphi}, \mathcal{K})$  has a representation generalizing Equation (2) and having the form B = T + K with K compact and

(22) 
$$T = cI + f(C_{\varphi}^*C_{\varphi}) + g(C_{\varphi}C_{\varphi}^*) + Uh(C_{\varphi}^*C_{\varphi}) + U^*k(C_{\varphi}C_{\varphi}^*),$$

where f and h are continuous on  $\sigma(C_{\varphi}^*C_{\varphi})$ , g and k are continuous on  $\sigma(C_{\varphi}C_{\varphi}^*)$ , all four functions vanish at zero, and U is the partial isometry polar factor (which in this case is unitary) of  $C_{\varphi}$ . The restrictions of f, g, h and k to the interval [0, s], which coincides with both of the essential spectra  $\sigma_e(C_{\varphi}^*C_{\varphi})$  and  $\sigma_e(C_{\varphi}C_{\varphi}^*)$ , are uniquely determined by B. We call T a *distinguished representative* of the coset [B], and recall from [15] that

$$\Psi(B) = \Psi(T) = \begin{bmatrix} c+g & h \\ k & c+f \end{bmatrix}$$

We start with the operator  $(C_{\varphi}^*C_{\varphi})^a$ , defined by the self-adjoint functional calculus, where Re a > 0. Note that

$$C^*_{\varphi}C_{\varphi} \equiv sC_{\sigma}C_{\varphi} \pmod{\mathcal{K}} = sC_{\varphi\circ\sigma} = sC_{\rho_{\eta,2b}}$$

where the last equality follows from Theorem 6. In Theorem 3, take  $\gamma = \eta$  and consider the \*-isomorphism  $\Gamma$ , here called  $\Gamma_{\eta}$  to emphasize the fixed point  $\eta$ . We have

$$[(C^*_{\varphi}C_{\varphi})^a] = [C^*_{\varphi}C_{\varphi}]^a = s^a [C_{\rho_{\eta,2b}}]^a = s^a \Gamma_{\eta}(t^{2b})^a = s^a \Gamma_{\eta}(t^{2ba}) = s^a [C_{\rho_{\eta,2ba}}].$$

The first and fourth equalities follow, respectively, from the facts that the coset map  $B \mapsto [B]$  and  $\Gamma_{\eta}$  are each \*-homomorphisms. Relabeling 2ba as a, we see that  $s^{-\frac{a}{2b}}(C_{\varphi}^*C_{\varphi})^{\frac{a}{2b}}$  is a distinguished representative of  $[C_{\rho_{\eta,a}}]$  for Re a > 0. A similar argument shows that the coset  $[C_{\rho_{\zeta,a}}]$  has distinguished representative  $s^{-\frac{a}{2c}}(C_{\varphi}C_{\varphi}^*)^{\frac{a}{2c}}$  for Re a > 0.

Now consider  $\rho_{\eta,a} \circ \varphi$ , which we know to be a self-map of  $\mathbb{D}$ , but not an automorphism, when Re a > -b. First we look at the case Re a > 0. We have

$$C_{\rho_{\eta,a} \circ \varphi} = C_{\varphi} C_{\rho_{\eta,a}} = U(C_{\varphi}^* C_{\varphi})^{\frac{1}{2}} C_{\rho_{\eta,a}} \equiv s^{-\frac{a}{2b}} U(C_{\varphi}^* C_{\varphi})^{\frac{1}{2} + \frac{a}{2b}} \pmod{\mathcal{K}}$$

by our above discussion. By the spectral theorem,  $(C^*_{\varphi}C_{\varphi})^z$  is holomorphic for Re z > 0 in the weak operator topology, and therefore in the operator norm topology; see [11], Theorem 3.10.1. Thus the cosets  $[C_{\rho_{\eta,a}\circ\varphi}]$  and  $[s^{-\frac{a}{2b}}U(C^*_{\varphi}C_{\varphi})^{\frac{1}{2}+\frac{a}{2b}}]$  are both holomorphic  $\mathcal{B}(H^2)/\mathcal{K}$ -valued functions of a, Re a > -b, which agree on the subset  $\{a : \text{Re } a > 0\}$ . Hence they agree on all of  $\{a : \text{Re } a > -b\}$ , showing that  $s^{-\frac{a}{2b}}U(C^*_{\varphi}C_{\varphi})^{\frac{1}{2}+\frac{a}{2b}}$  is a distinguished representative of  $[C_{\rho_{\eta,a}\circ\varphi}]$  when Re a > -b. An analogous statement holds for  $[C_{\rho_{\zeta,a}\circ\sigma}]$  with Re a > -c. The following table summarizes these conclusions.

TABLE II				
Linear-fractional $\psi$ with $C_{\psi}$ in $C^*(C_{\varphi}, \mathcal{K})$				
Condition on $\psi$ in Theorem 5	$\psi$	Distinguished representative of $[C_{\psi}]$	Matrix function $\Psi(C_{\psi})(t), \ 0 \le t \le s$	
(d)	$\rho_{\eta,a},$ Re $a > 0$	$s^{-\frac{a}{2b}}(C^*_{\varphi}C_{\varphi})^{\frac{a}{2b}}$	$\left[\begin{array}{cc} \left(\frac{t}{s}\right)^{\frac{a}{2b}} & 0\\ 0 & 0 \end{array}\right]$	
(b)	$\begin{aligned} \rho_{\zeta,a}, \\ \text{Re } a > 0 \end{aligned}$	$s^{-\frac{a}{2c}} (C_{\varphi} C_{\varphi}^*)^{\frac{a}{2c}}$	$\left[\begin{array}{cc} 0 & 0 \\ 0 & \left(\frac{t}{s}\right)^{\frac{a}{2c}} \end{array}\right]$	
(a)	$\rho_{\eta,a} \circ \varphi,$ Re $a > -b$	$s^{-\frac{a}{2b}}U(C_{\varphi}^*C_{\varphi})^{\frac{1}{2}+\frac{a}{2b}}$	$\left[\begin{array}{cc} 0 & \sqrt{t} \left(\frac{t}{s}\right)^{\frac{a}{2b}} \\ 0 & 0 \end{array}\right]$	
(c)	$\rho_{\zeta,a} \circ \sigma,$ Re $a > -c$	$s^{-\frac{a}{2c}-1}U^*(C_{\varphi}C_{\varphi}^*)^{\frac{1}{2}+\frac{a}{2c}}$	$\left[\begin{array}{cc} 0 & 0\\ \frac{\sqrt{t}}{s} \left(\frac{t}{s}\right)^{\frac{a}{2c}} & 0 \end{array}\right]$	

Given an operator B in  $C^*(C_{\varphi}, \mathcal{K})$ ,  $\sigma_e(B)$  and  $\|B\|_e$  coincide with  $\sigma(\Psi(B))$  and  $\|\Psi(B)\|$ , respectively. Thus, if B is a finite linear combination of composition operators  $C_{\psi}$  with  $\psi$ 's chosen from Column 2 in Table II, one can calculate  $\Psi(B)$  from Column 4 and in principle read off  $\sigma_e(B)$  and  $\|B\|_e$ ; see Theorem 4.17 in [15].

It is known [14] that the collection of linear-fractional composition operators  $C_{\psi}$  with  $\psi$  a non-automorphism having  $\|\psi\|_{\infty} = 1$  is linearly independent modulo  $\mathcal{K}$ . The following result shows that this remains true when  $\mathcal{K}$  is replaced by the larger subspace  $C^*(C_{\varphi}, \mathcal{K})$  of  $\mathcal{B}(H^2)$ . **Theorem 7.** Let  $\varphi$  be as in (1). Suppose that  $\beta_1, \dots, \beta_q$  are distinct linearfractional self-maps of  $\mathbb{D}$  and that  $a_1 \dots, a_q$  are non-zero complex numbers. If the linear combination  $a_1C_{\beta_1} + \dots + a_qC_{\beta_q}$  lies in  $C^*(C_{\varphi}, \mathcal{K})$ , then so do the individual operators  $C_{\beta_1}, \dots, C_{\beta_q}$ .

*Proof.* Let us discard those  $C_{\beta_i}$ 's which lie in  $C^*(C_{\varphi}, \mathcal{K})$  and assume for the purpose of obtaining a contradiction that there are some left over. Relabel these as  $C_{\beta_1}, \dots, C_{\beta_r}$ , let  $a_1, \dots, a_r$  be the corresponding constants, and put  $T = a_1 C_{\beta_1} + \dots + a_r C_{\beta_r}$ , which lies in  $C^*(C_{\varphi}, \mathcal{K})$ . Here, none of the summands are compact, so  $\|\beta_i\|_{\infty} = 1$  for  $i = 1, \dots, r$ . Now we proceed almost as in the proof of Theorem 1, with T playing the role of  $C_{\psi}$ . Given  $\epsilon > 0$  there exists A as in that proof and a complex c such that  $\|T - cC_z - A\|_e < \epsilon$ . By the inequality (11)

$$||T - cC_z - A||_e^2 \ge |c|^2 + \frac{1}{2\pi} \sum_{i=1}^r |a_i|^2 |J(\beta_i)|$$

so that  $|c| < \epsilon$ , since each  $|J(\beta_i)|$  must be zero, so  $\beta_i$  is a non-automorphism. As earlier, we have  $||T - A||_e < 2\epsilon$ .

According to Corollary 5.17 in [14], the cosets  $[C_{\beta_1}], \cdots, [C_{\beta_r}]$  are linearly independent in  $\mathcal{B}(H^2)/\mathcal{K}$ . It follows that T is not compact, so the matrix function

(23) 
$$\Psi(T) = \begin{bmatrix} f_2 & f_3 \\ f_4 & f_1 \end{bmatrix}$$

is not identically zero on [0, s]. As in Theorem 1, we focus on  $f_3$  and aim for a contradiction by assuming that  $||f_3||_{\infty} > 0$ . An appropriate choice of  $\epsilon$  again yields the inequality (12), where p has the same meaning as there. Again we write A in the form (13), and thus have

$$4\epsilon^{2} > \|T - A\|_{e}^{2} \geq \limsup_{|z| \to 1} \left\| (T^{*} - A^{*}) \frac{k_{z}}{\|k_{z}\|} \right\|^{2}$$
$$\geq \lim_{\Gamma_{\zeta,D}} \left\| \left( \sum_{i=1}^{r} \overline{a_{i}} C_{\beta_{i}}^{*} - \sum_{i=1}^{m} \overline{c_{i}} C_{\psi_{i}}^{*} \right) \frac{k_{z}}{\|k_{z}\|} \right\|^{2}$$
$$\geq \left\| \sum_{\substack{\zeta \in F(\psi_{i}) \\ D_{1}(\psi_{i}, \zeta) = \mathbf{d}_{3}} \overline{c_{i}} k_{w_{i}}^{+} \right\|_{H_{+}^{2}}^{2}$$

The rest of the argument follows that of Theorem 1 exactly, reaching the same contradiction.  $\hfill \Box$ 

#### 5. Non linear-fractional maps

In this section we explore maps  $\psi$ , satisfying either condition (e) or (f) of Theorem 2, for which  $C_{\psi}$  lies in  $C^*(C_{\varphi}, \mathcal{K})$ ; our main result shows that such maps exist. We begin with a lemma about finite Blaschke products.

**Lemma 5.** Given  $\zeta$ ,  $\eta$  distinct points on  $\partial \mathbb{D}$ , and positive numbers  $t_1, t_2$ , there exists a finite Blaschke product B with the properties  $B(\eta) = \eta$ ,  $B(\zeta) = \eta$ ,  $B'(\eta) = t_1$  and  $|B'(\zeta)| = t_2$ . Moreover,  $B'(\zeta) = \eta \overline{\zeta} t_2$ .

*Proof.* Clearly there is no loss of generality in taking  $\eta = 1$ . Initially we will also suppose that  $\zeta = -1$ ; this condition will be removed at the end. A finite Blaschke product  $B(z) = \prod \frac{|a_n|}{a_n} (a_n - z)/(1 - \overline{a_n}z)$  will meet the conditions B(1) = 1, B(-1) = 1 if both

$$\prod \frac{|a_n|}{a_n} \frac{a_n - 1}{1 - \overline{a_n}} = 1$$

and

$$\prod \frac{|a_n|}{a_n} \frac{a_n + 1}{1 + \overline{a_n}} = 1$$

It is easy to see that both of these conditions will be met if the zeros of B are chosen to be a collection of conjugate pairs  $\{a, \overline{a}\}$ . The conditions  $B'(1) = t_1, |B'(-1)| = t_2$  are satisfied if

(24) 
$$\sum \frac{1 - |a_n|^2}{|1 - a_n|^2} = t_1$$

and

(25) 
$$\sum \frac{1 - |a_n|^2}{|1 + a_n|^2} = t_2$$

respectively (see [2]).

Next observe that for any t > 0,  $\{z : 1 - |z|^2 = t|1 - z|^2\}$  is a circle centered at (t/(t+1), 0) with radius 1/(t+1) and  $\{z : 1 - |z|^2 = t|1+z|^2\}$  is a circle centered at (-t/(t+1), 0) with radius 1/(1+t). As  $t \to 0$ , the centers of these circles approach 0 and the radii tend to 1. Thus given  $t_1, t_2$  arbitrary positive numbers we may choose m a positive integer sufficiently large so that the circles  $\{z : 1 - |z|^2 = \frac{t_1}{2^m}|1-z|^2\}$  and  $\{z : 1 - |z|^2 = \frac{t_2}{2^m}|1+z|^2\}$  intersect in a conjugate pair of points  $a, \overline{a}$ . Consider the Blaschke product B(z) with a zero of order  $2^{m-1}$  at a and a zero of order  $2^{m-1}$  at  $\overline{a}$ . Since the zeros occur at conjugate pairs, B(1) = 1 and B(-1) = 1. By construction

$$(1 - |a|^2)/(|1 - a|^2) = (1 - |\overline{a}|^2)/(|1 - \overline{a}|^2) = \frac{t_1}{2^m},$$

and

$$(1-|a|^2)/(|1+a|^2) = (1-|\overline{a}|^2)/(|1+\overline{a}|^2) = \frac{t_2}{2^m},$$

so that the zeros of B satisfy Equations (24) and (25) as desired, and  $B'(1) = t_1$ ,  $|B'(-1)| = t_2$ .

Now suppose  $\zeta \in \partial \mathbb{D}$  is not equal to -1. Find a parabolic automorphism  $\tau$  fixing 1, with derivative 1 there, and taking  $\zeta$  to -1; a unique such  $\tau$  exists since (purely imaginary) translations act transitively on the boundary of the right half-plane. Then for B as constructed above,  $B \circ \tau$  is a finite Blaschke product fixing 1, with derivative  $t_1$  at 1, sending  $\zeta$  to 1 and having derivative  $|(B \circ \tau)'(\zeta)| = |B'(-1)||\tau'(\zeta)|$ ; since |B'(-1)| can be arbitrarity prescribed and  $\tau$  depends only on the value of  $\zeta$ , this means  $|(B \circ \tau)'(\zeta)|$  can be chosen to be an arbitrary positive number. Finally observe that if B is a Blaschke product with  $B(\zeta) = \eta$  and  $|B'(\zeta)| = s$ , then we must have  $B'(\zeta) = \eta \overline{\zeta}s$ , since  $\overline{\eta}\zeta B(z)$  fixes  $\zeta$ , and hence has positive derivative there.

**Theorem 8.** Suppose that  $\varphi$  is as in (1). There exist analytic self-maps  $\psi_1$  and  $\psi_2$  of  $\mathbb{D}$ , satifying conditions (e) and (f) of Theorem 2 respectively, such that  $C_{\psi_1}$  and

 $C_{\psi_2}$  lie in  $C^*(C_{\varphi}, \mathcal{K})$ . Moreover,  $\psi_1$  and  $\psi_2$  can be taken to extend continuously to the closed disk  $\overline{\mathbb{D}}$ .

Proof. We shall indicate the construction for  $\psi_1$  with the normalization  $\eta = 1$ . First consider a simply connected domain  $\Omega$  in  $\mathbb{D}$ , whose boundary is a smooth Jordan curve which at 1 and at -1 includes an arc of an internally tangent circle, such that  $\overline{\Omega} \cap \partial \mathbb{D} = \{-1, 1\}$ . It is easy to construct a conformal map  $\rho : \mathbb{D}$  onto  $\Omega$ , extending to a homeomorphism of  $\overline{D}$  onto  $\overline{\Omega}$ , with the properties  $\rho(1) = 1, \rho(-1) = -1$  and  $\rho'(1) = \frac{1}{2}$ . The map  $\rho$  will be analytic in a neighborhood of both 1 and -1. Let  $\tau$  be the unique parabolic automorphism of  $\mathbb{D}$  fixing 1, having derivative 1 there, and mapping  $\zeta$  to -1. By the uniqueness statement,  $\tau'(\zeta)$  is determined by  $\zeta$ . Using Lemma 5 construct a finite Blaschke product B with B(1) = 1, B(-1) = 1, B'(1) = 2 and |B'(-1)| chosen so that  $|B'(-1)||\rho'(-1)||\tau'(\zeta)| = s$ , where  $s = |\varphi'(\zeta)|$ . The map  $\psi_1 \equiv B \circ \rho \circ \tau$  is a self-map of  $\mathbb{D}$ , with finite angular derivative set  $F(\psi_1) = \{\zeta, 1\}$  and satisfying  $\psi_1(\zeta) = 1, \psi_1(1) = 1, \psi_1'(1) = 1$  and  $\psi_1'(\zeta) = \overline{\zeta}s$ , this last condition following from  $|\psi_1'(\zeta)| = s$  and  $\psi_1(\zeta) = 1$ . Clearly  $\psi_1$  has order of contact two at 1 and  $\zeta$ .

Any linear-fractional map  $\beta$  is uniquely determined by its second order data vector  $D_2(\beta, z_0) = (\beta(z_0), \beta'(z_0), \beta''(z_0))$  at any point  $z_0$  of analyticity. The curvature of the curve  $\psi_1(\partial \mathbb{D})$  at the points  $\psi_1(\zeta)$  and  $\psi_1(1)$  is determined by  $D_2(\psi_1, \zeta)$ and  $D_2(\psi_1, 1)$ , respectively. By construction of  $\psi_1$ , these curvatures exceed unity. There exists unique linear-fractional maps  $\beta_1, \beta_2$  of  $\mathbb{D}$  with the second order data vectors

$$D_2(\beta_1, 1) = D_2(\psi_1, 1) = (1, 1, \psi_1''(1)) \text{ and } D_2(\beta_2, \zeta) = D_2(\psi_1, \zeta) = (1, \overline{\zeta}s, \psi_1''(\zeta)).$$

The curvature of  $\{\psi_1(e^{i\theta}): e^{i\theta} \in \partial \mathbb{D}\}$  matches that of  $\beta_1(\partial \mathbb{D})$  at  $e^{i\theta} = 1$  and that of  $\beta_2(\partial \mathbb{D})$  at  $e^{i\theta} = \zeta$ . Thus  $\beta_1$  and  $\beta_2$  are non-automorphism self-maps of  $\mathbb{D}$ . Since  $\varphi'(\zeta) = \psi'_1(\zeta) = \beta'_2(\zeta)$ , Theorem 5 shows that  $C_{\beta_1}$  and  $C_{\beta_2}$  are in  $C^*(C_{\varphi}, \mathcal{K})$ , and hence so is  $C_{\beta_1} + C_{\beta_2}$ . By Corollary 5.16 of [14],  $C_{\psi_1} \equiv C_{\beta_1} + C_{\beta_2} \pmod{\mathcal{K}}$  and thus  $C_{\psi_1}$  is in  $C^*(C_{\varphi}, \mathcal{K})$ .

For  $\psi_2$ , note that the Krein adjoint  $\sigma$  of  $\varphi$  satisfies  $\sigma(\eta) = \zeta$  and  $\sigma'(\eta) = 1/\varphi'(\zeta)$ . We apply the first part of the proof, with  $\varphi$  replaced by  $\sigma$ , to find a self map  $\psi_2$  of  $\mathbb{D}$  with  $F(\psi_2) = \{\zeta, \eta\}, \ \psi_2(\eta) = \zeta, \ \psi'_2(\eta) = \sigma'(\eta) = 1/\varphi'(\zeta), \ \psi_2(\zeta) = \zeta$ , and  $\psi'_2(\zeta) = 1$  such that  $C_{\psi_2} \in C^*(C_{\sigma}, \mathcal{K}) = C^*(C_{\varphi}, \mathcal{K})$  as desired.  $\Box$ 

Our last theorem shows that for sufficiently nice  $\psi$ , Theorem 2 is almost the whole story.

**Theorem 9.** Let  $\psi$  be an analytic self-map of  $\mathbb{D}$  such that  $F(\psi)$  is a finite set,  $\psi$  extends analytically to a neighborhood of each point in  $F(\psi)$ , and for any open set U of  $\partial \mathbb{D}$  containing  $F(\psi)$ ,  $\|\chi_{\partial \mathbb{D} \setminus U}\psi\|_{\infty} < 1$ . If  $\varphi$  is as in (1), then  $C_{\psi}$  lies in  $C^*(C_{\varphi}, \mathcal{K})$  if and only if

- (i) one of the conditions (a)-(f) of Theorem 2 holds, and
- (ii) the map  $\psi$  has order of contact two at each point of  $F(\psi)$ .

*Proof.* Suppose  $\psi$  is as described and  $C_{\psi}$  is in  $C^*(C_{\varphi}, \mathcal{K})$ . By Theorem 2, one of (a)-(f) holds. Suppose that  $\gamma$  is in  $F(\psi)$  and  $\psi$  has order of contact exceeding two at  $\gamma$ . Following Theorem 1, we let  $\epsilon > 0$  and find a finite linear combination A of composition operators whose self-maps are chosen from the lists (6) such that  $\|C_{\psi} - A\| < \epsilon$ . At the same time we apply the inequality (9) to the linear

combination  $C_{\psi} - A$  at the point  $\alpha = \gamma$ . The maps in the lists (6), being linearfractional non-automorphisms, all have order of contact two at the unique points in their angular derivative sets. Taking the left side of (9) to be  $||C_{\psi} - A||_e^2$  and k = 3 on the right side, the sum on the right-hand side has only one term, namely  $1/|\psi'(\gamma)|$ , giving

$$\epsilon^2 > ||C_{\psi} - A||_e^2 \ge \frac{1}{|\psi'(\gamma)|},$$

a contradiction. Thus  $\psi$  must have order of contact two at  $\gamma$ .

Conversely, suppose  $\psi$  satisfies (i) and (ii). If  $F(\psi) = \{\zeta\}$ , let  $\beta$  be the unique linear-fractional map with  $D_2(\beta, \zeta) = D_2(\psi, \zeta)$ . Since  $\psi$  has order of contact two at  $\zeta$ , the curvature of  $\{\psi(e^{i\theta}) : e^{i\theta} \in \partial \mathbb{D}\}$  at  $e^{i\theta} = \zeta$  exceeds unity, so that  $\beta(\partial \mathbb{D})$ , having the same curvature, is internally tangent to  $\partial \mathbb{D}$  at  $\psi(\zeta)$  and bounds a proper subdisk of  $\mathbb{D}$ . Thus  $\beta$  is a non-automorphism of  $\mathbb{D}$  satisfying (a) or (b) of Theorem 5, and  $C_\beta$  lies in  $C^*(C_{\varphi}, \mathcal{K})$ . The same argument covers the case  $F(\psi) = \{\eta\}$ .

If  $F(\psi) = \{\zeta, \eta\}$  we proceed as in the proof of Theorem 8 to produce linearfractional non-automorphisms  $\beta_1$  and  $\beta_2$  of  $\mathbb{D}$  with  $C^*(C_{\varphi}, \mathcal{K})$  containing  $C_{\beta_1}$  and  $C_{\beta_2}$  and  $C_{\psi} \equiv C_{\beta_1} + C_{\beta_2} \pmod{\mathcal{K}}$ .

# References

- P. Bourdon, D. Levi, S. Narayan and J. Shapiro, Which linear fractional composition operators are essentially normal?, J. Math. Anal. Appl. 280 (2003), 30–53.
- [2] P. Ahern and D. Clark, On inner functions with H<sup>p</sup>-derivative, Michigan Math. J. 21 (1974), 115–127.
- [3] E. Basor and D. Retsek, Extremal non-compactness of composition operators with linear fractional symbol, J. Math. Anal. Appl. 322 (2006), 749–763.
- [4] E. Berkson, Composition operators isolated in the uniform operator topology, Proc. Amer. Math. Soc. 81 (1981), 230–232.
- [5] J. Conway, A Course in Operator Theory, Amer. Math. Soc., Providence, 2000.
- [6] C. Cowen, Composition operators on H<sup>2</sup>, J. Operator Theory 9 (1983), 77-106.
- [7] C. Cowen, Linear-fractional composition operators on H<sup>2</sup>, Integral Equations Operator Theory 11 (1988), 151–160.
- [8] C. Cowen and B. MacCluer, Composition Operators on Spaces of Analytic Functions, CRC Press, Boca Raton, 1995.
- [9] A. Dash, Joint essential spectra, Pacific J. Math., 64 (1976), 119–128.
- [10] J. Guyker, On reducing subspaces of composition operators, Acta Sci. Math. (Szeged) 53 (1989), 369–376.
- [11] E. Hille and R. Phillips, Functional Analysis and Semi-Groups, Amer. Math. Soc., Providence, 1957.
- [12] M. Jury, C\*-algebras generated by groups of composition operators, Indiana Univ. Math. J. 56 (2007), 3171–3192.
- [13] M. Jury, The Fredholm index for elements of Toeplitz-composition C\*-algebras, Integral Equations Operator Theory, 58 (2007), 341–362.
- [14] T. Kriete and J. Moorhouse, Linear relations in the Calkin algebra for composition operators, Trans. Amer. Math. Soc. 359 (2007), 2915–2944.
- [15] T. Kriete, B. MacCluer, and J. Moorhouse, Toeplitz-composition C\*-algebras, J. Operator Theory 58 (2007), 135–156.
- [16] T. Kriete, B. MacCluer, and J. Moorhouse, Spectral theory for algebraic combinations of Toeplitz and composition operators, preprint.
- [17] B. MacCluer and J. Shapiro, Angular derivatives and compact composition operators on the Hardy and Bergman spaces, Canadian J. Math. 38 (1986), 878–906.
- [18] A. Montes-Rodríguez, M. Ponce-Escudero, and S. Shkarin, Invariant subspaces of parabolic self-maps in the Hardy space, preprint.
- [19] J. Shapiro and C. Sundberg, Isolation amongst the composition operators, Pacific J. Math. 145 (1990), 117–152.

- [20] J. Shapiro and P. Taylor, Compact, nuclear, and Hilbert-Schmidt composition operators on  $H^2$ , Indiana Univ. Math. J. 23 (1973), 471–496.
- [21] J. Shapiro, Composition Operators and Classical Function Theory, Springer-Verlag, New York, 1993.

Department of Mathematics, University of Virginia, Charlottesville, VA 22904  $E\text{-}mail\ address: \texttt{tlk8q@virginia.edu}$ 

Department of Mathematics, University of Virginia, Charlottesville, VA 22904 E-mail address: bdm3f@virginia.edu

Department of Mathematics, Colgate University, Hamilton, NY 13356 E-mail address: jmoorhouse@mail.colgate.edu