THE ORNSTEIN UHLENBECK BRIDGE AND APPLICATIONS TO MARKOV SEMIGROUPS

B. GOLDYS AND B. MASLOWSKI

ABSTRACT. For an arbitrary Hilbert space-valued Ornstein-Uhlenbeck process we construct the Ornstein-Uhlenbeck Bridge connecting a starting point x and an endpoint y that belongs to a certain linear subspace of full measure. We derive also a stochastic evolution equation satisfied by the OU Bridge and study its basic properties. The OU Bridge is then used to investigate the Markov transition semigroup associated to a nonlinear stochastic evolution equation with additive noise. We provide an explicit formula for the transition density and study its regularity. Given the Strong Feller property and the existence of an invariant measure we show that the transition semigroup maps L^p functions into continuous functions. We also show that transition operators are q-summing for some q > p > 1, in particular of Hilbert-Schmidt type.

Contents

1. Introduction	1
2. Preliminaries on OU Processes and Bridges	3
2.1. Measurable Linear Mappings	3
2.2. Conditional Distributions	4
2.3. Some Properties of the Ornstein-Uhlenbeck Process	Ę
2.4. Fundamentals on OU Bridge	8
3. SDE associated to the OU Bridge	11
4. Applications to Semilinear Equations	18
References	27

1. Introduction

Let (Z_t^x) be an Ornstein-Uhlenbeck process on a separable Hilbert space H. By this we mean that (Z_t^x) is a solution to a linear stochastic evolution equation

$$\begin{cases} dZ_t^x = AZ_t^x dt + \sqrt{Q} dW_t, \\ Z_0^x = x \in H. \end{cases}$$
 (1.1)

¹⁹⁹¹ Mathematics Subject Classification. 60H15, 60H10, 60J60, 60J35, 35R60.

Key words and phrases. Ornstein-Uhlenbeck process, pinned process, measurable linear mapping, stochastic semilinear equation, transition density.

This work was partially supported by the UNSW Faculty Research Grant and GAČR grant 201/04/0750.

In the above equation (W_t) is a standard cylindrical Wiener process defined on a certain stochastic basis $(\Omega, \mathscr{F}, (\mathscr{F}_t), \mathbb{P})$ and $Q = Q^* \geqslant 0$ is a bounded operator on H. We assume that the operator (A, dom(A)) is a generator of a C_0 -semigroup (S_t) on H. Under the assumptions given below the solution to (1.1) is defined by the formula

$$Z_t^x = S_t x + \int_0^t S_{t-s} \sqrt{Q} dW_s. \tag{1.2}$$

The aim of this paper is to study the basic properties of the Ornstein-Uhlenbeck Bridge (sometimes called a Pinned Ornstein-Uhlenbeck process) $(\hat{Z}_t^{x,y})$ associated to the Ornstein-Uhlenbeck process (Z_t^x) and its applications. Let us recall informally, that this process is defined via the formula

$$\mathbb{P}\left(\left.Z_{t}^{x} \in B\right| Z_{T}^{x} = y\right) = \mathbb{P}\left(\hat{Z}_{t}^{x,y} \in B\right), \quad t < T,$$

where $x, y \in H$ and $B \subset H$ is a Borel set. Intuitively, it is an Ornstein-Uhlenbeck process "conditioned to go from x at time t = 0 to y at time t = T" (a rigorous definition is given in Section 2, cf. Def. 2.15). The importance of various types of bridge processes in the theory of finite dimensional diffusions is well recognised, see for example [20]. In infinite dimensional framework this concept was developed in [17] in order to study regularity of transition semigroup of certain linear and nonlinear diffusions on Hilbert space. In [15] and [16] an Ornstein-Uhlenbeck Bridge is introduced in order to obtain lower estimates on the transition kernel of some semilinear stochastic evolution equations. Those estimates provide a powerful tool to study exponential ergodicity and V-uniform ergodicity for such equations. In particular, they allowed us to obtain in [11] explicit estimates of the rate of exponential convergence to the invariant measure.

In the present paper the OU Bridge is studied under much more general conditions and in more detail. We provide also further applications of the OU Bridge to the analysis of transition densities and the regularity of associated Markov semigroups. Regularity of Strongly Feller transition semigroups was studied in [8] (see also references therein). We use methods completely different from [8] and obtain stronger results but for bounded drifts only while the aforementioned paper allows linearly growing drifts. Closely related results for semigroups that are not strongly Feller may be found in [3]. For the regularity of strongly Feller semigroups associated to the OU process we refer to [5].

Let us describe the contents of this paper. In Section 2 we provide, for the reader's convenience, some relevant facts about linear measurable mappings and conditional distributions of Hilbert space valued Gaussian random vectors. Then we give a definition of the OU Bridge and some basic results on OU processes and OU Bridges. Some of the technical results from [11] that are needed in the sequel are stated without proof and others (Lemma 2.8, Proposition 2.11 and Lemma 3.3) are reproved under more general conditions. In Section 3, a stochastic equation for the OU Bridge is derived. A new Brownian Motion adapted to the filtration of the Ornstein Uhlenbeck Bridge is obtained and then it is shown that the Bridge process is a unique mild (and weak) solution of a linear

nonhomogeneous stochastic evolution equation with singular coefficients. Section 4 is devoted to applications of the previous results to semilinear stochastic equations; continuity of Markov transition densities (with respect to the Gaussian invariant measure ν that is an invariant measure with respect to the OU process) is proved (Theorem 4.5 and Remark 4.9), the Markov semigroup is shown to map the space $L^p(H,\nu)$, p>1, into the space of continuous functions on H (Theorem 4.6) and is also shown to be Hilbert-Schmidt on $L^2(H,\nu)$ and q-summing (in particular, compact) as a mapping $L^p(H,\nu) \to L^q(H,\nu)$ even if q>p provided the gap between q and p is not too large (Theorem 4.7). At the end of the section the results are illustrated in the case of one-dimensional semilinear stochastic parabolic equation (Example 4.10) in which case all conditions imposed in the paper are verified or specified.

ACKNOWLEDGEMENT. The authors are grateful to Jan Seidler for his valuable comments and suggestions.

2. Preliminaries on OU Processes and Bridges

In this section we collect, for the reader convenience, some properties of infinite-dimensional OU processes and Gaussian random variables which will be useful in the paper. We also define the OU Bridge and recall some known results that will be useful in the sequel.

2.1. Measurable Linear Mappings. Let H be a real separable Hilbert space and let $\mu = N(0, C)$ be a centered Gaussian measure on H with the covariance operator C such that $\overline{\operatorname{im}(C)} = H$. The space $H_C = \operatorname{im}(C^{1/2})$ endowed with the norm $|x|_C = |C^{-1/2}x|$ can be identified as the Reproducing Kernel Hilbert Space of the measure μ . In the sequel we will denote by $\{e_n : n \ge 1\}$ the eigenbasis of C and by $\{c_n : n \ge 1\}$ the corresponding set of eigenvalues:

$$Ce_n = c_n e_n, \quad n \geqslant 1.$$

For any $h \in H$ we define

$$\phi_n(x) = \sum_{k=1}^n \frac{1}{\sqrt{c_k}} \langle h, e_k \rangle \langle x, e_k \rangle, \quad x \in H.$$

The following two lemmas are well known (see e.g. [11]):

Lemma 2.1. The sequence (ϕ_n) converges in $L^2(H,\mu)$ to a limit ϕ and

$$\int_{H} |\phi(x)|^2 \, \mu(dx) = |h|^2.$$

Moreover, there exists a measurable linear space $\mathcal{M}_h \subset H$, such that $\mu(\mathcal{M}_h) = 1$, ϕ is linear on \mathcal{M}_h and

$$\phi(x) = \lim_{n \to \infty} \phi_n(x), \quad x \in \mathcal{M}_h. \tag{2.1}$$

We will use the notation $\phi(x) = \langle h, C^{-1/2}x \rangle$.

Let H_1 be another real separable Hilbert space and let $T: H \to H_1$ be a bounded operator. The Hilbert-Schmidt norm of T will be denoted by $||T||_{HS}$. Let

$$\tilde{T}_n x = \sum_{k=1}^n \frac{1}{\sqrt{c_k}} \langle x, e_k \rangle T e_k, \quad x \in H.$$

Lemma 2.2. Let $T: H \to H_1$ be a Hilbert-Schmidt operator. Then the sequence (\tilde{T}_n) converges in $L^2(H, \mu; H_1)$ to a limit \tilde{T} and

$$\int_{H} \left| \tilde{T}(x) \right|_{H_{1}}^{2} \mu(dx) = \|T\|_{HS}^{2}.$$

Moreover, there exists a measurable linear space $\mathcal{M}_T \subset H$, such that $\mu(\mathcal{M}_T) = 1$, \tilde{T} is linear on \mathcal{M}_T and

$$\tilde{T}(x) = \lim_{n \to \infty} \tilde{T}_n x, \quad x \in \mathcal{M}_T.$$
 (2.2)

We will use the notation $TC^{-1/2}x := \tilde{T}(x)$.

The above procedure is specified in the following Lemma (the proof of which may be found in [11]) to operator-valued functions:

Lemma 2.3. Let $K(t,s): H \to H$ be an operator-valued, strongly measurable function, such that for each $a \in (0,T)$

$$\int_0^a \int_0^a \|K(t,s)\|_{HS}^2 \, ds dt < \infty. \tag{2.3}$$

Then the following holds.

- (a) There exists a Borel set $B \subset [0,T]^2$ of full Lebesgue measure such that the measurable linear mapping $K(t,s)C^{-1/2}$ is well defined for all $(s,t) \in B$.
- (b) There exists a measurable mapping $f:[0,T)^2 \times H \to H$ and a measurable linear space $\mathcal{M} \subset H$ of full measure such that $f(t,s,y) = K(t,s)C^{-1/2}y$ for $y \in \mathcal{M}$ and for each a < T

$$\int_0^a |f(t,s,y)| \, ds < \infty$$

for almost all $t \in [0,T]$. We will use the notation $K(t,s)C^{-1/2}y := f(t,s,y)$.

2.2. Conditional Distributions. Let H_1 and H_2 be two real separable Hilbert spaces and let $(X, Y) \in H_1 \times H_2$ be a Gaussian vector with mean values

$$m_X = \mathbb{E}X$$
, and $m_Y = \mathbb{E}Y$.

The covariance operator of X is determined by the equation

$$\mathbb{E}\langle X - m_X, h \rangle \langle X - m_X, k \rangle = \langle C_X h, k \rangle, \quad h, k \in H_1, \tag{2.4}$$

and a similar condition determines the covariance C_Y of Y. The covariance operator $C_{XY}: H_1 \to H_2$ is defined by the condition

$$\langle C_{XY}h, k \rangle = \mathbb{E} \langle X - m_X, h \rangle \langle Y - m_Y, k \rangle, \quad h \in H_1, k \in H_2,$$

and then $C_{XY}^* = C_{YX}$.

For a linear closable operator G on H the closure of G will be denoted by \overline{G} . The next theorem is well known, see for example [14]

Theorem 2.4. Assume that C_X is injective. Then the following holds.

(a) We have

$$\operatorname{im}\left(C_{YX}\right) \subset \operatorname{im}\left(C_X^{1/2}\right),$$
 (2.5)

the operator $T = C_X^{-1/2}C_{YX}$ is of Hilbert-Schmidt type on H and $T^* = \overline{C_{XY}C_X^{-1/2}}$.

(b) We have

$$\mathbb{E}\left(Y|X\right) = m_Y + T^* C_X^{-1/2} \left(X - m_X\right), \quad \mathbb{P}^X - a.s.$$

(c) The conditional distribution of Y given X is Gaussian $N\left(\mathbb{E}\left(Y|X\right), C_{Y|X}\right)$, where

$$C_{Y|X} = C_Y - T^*T.$$

Moreover, the random variables $T^*C_X^{-1/2}X$ and $\left(Y-T^*C_X^{-1/2}X\right)$ are independent.

2.3. Some Properties of the Ornstein-Uhlenbeck Process. The following hypothesis is a standing assumption for the rest of the paper.

Hypothesis 2.5. For every t > 0

$$\int_{0}^{t} \left\| S_{s} Q^{1/2} \right\|_{HS}^{2} ds < \infty, \tag{2.6}$$

and

$$\overline{\operatorname{im}(Q_t)} = H, (2.7)$$

where, in view of (2.6)

$$Q_t = \int_0^t S_s Q S_s^* ds. \tag{2.8}$$

is a well defined trace class operator.

It is well known that if Hypothesis 2.5 holds then the process (1.2) is a well defined H-valued, Gaussian and Markov process, see [7].

Let μ denote the probability law of the process $\{Z_t^0: t \in [0,1]\}$ that is concentrated on $L^2(0,T;H)$ and let $\mathcal{L}: L^2(0,T;H) \to C(0,T;H)$ be defined by the formula

$$\mathcal{L}u(t) = \int_0^t S_{t-s}Q^{1/2}u(s)ds.$$
 (2.9)

Note that, cf. [7], im(\mathscr{L}) = $RKHS(\mu)$ (the Reproducing Kernel Hilbert Space of the measure μ). We will use the notation μ_t^x for the Gaussian measure $N(S_tx, Q_t)$ and μ_t for μ_t^0 . By the properties of Gaussian distribution μ_t^x is the probability distribution of a random variable Z_t^x and we set $Z_t = Z_t^0$. In the rest of this subsection we give several statements on properties of the family of covariance operators $\{Q_t : t \leq T\}$ that will be useful later.

The definition of Q_t given in (2.8) yields immediately a simple identity that will be frequently used:

$$Q_T = Q_t + S_t Q_{T-t} S_t^*, \quad t \leqslant T. \tag{2.10}$$

Lemma 2.6. We have

$$\operatorname{im}\left(Q_t^{1/2}\right) \subset \operatorname{im}\left(Q_T^{1/2}\right), \quad t \leqslant T,$$

hence the operator $U_t = Q_T^{-1/2}Q_t^{1/2}$ is bounded on H for every $t \leqslant T$ and $||U_t|| \leqslant 1$. Moreover, $U_t^* = \overline{Q_t^{1/2}Q_T^{-1/2}}$, the closure of the operator $Q_t^{1/2}Q_T^{-1/2}$ defined on the domain $(Q_T^{1/2})$.

Proof. From the definition of the covariance operators Q_t it follows that $|Q_t x|^2 \leq |Q_T x|^2$ for each $x \in H$ and $0 \leq t \leq T$ and the conclusion easily follows.

Lemma 2.7. (a) The operator $V_t = Q_T^{-1/2} S_{T-t} Q_t^{1/2}$ is well defined and bounded on H and

$$||V_t|| \le 1, \quad t \in (0, T).$$
 (2.11)

Moreover,

$$\lim_{t \to T} V_t^* x = \lim_{t \to T} V_t x = x, \quad x \in H.$$
(2.12)

(b) For any $t \in [0, T]$

$$Q_{T-t} = Q_T^{1/2} \left(I - V_t V_t^* \right) Q_T^{1/2}. \tag{2.13}$$

Proof. The inequality (2.11) has been proved in [18], the convergence (2.12) in [11]. Part (b) follows immediately from (2.10).

Under a slightly stronger condition we show that the inequality (2.11) is sharp, more precisely, we have

Lemma 2.8. The following conditions are equivalent:

(a) For any $t \in (0,T]$

$$\operatorname{im}\left(Q_t^{1/2}\right) = \operatorname{im}\left(Q_T^{1/2}\right). \tag{2.14}$$

- (b) im (U_t) is dense in H for each $t \in (0,T)$.
- (c) We have

$$||V_t|| < 1, \quad t \in (0, T).$$
 (2.15)

Proof. Obviously (a) implies (b).

To prove that (b) implies (c) note first that (2.10) yields

$$\left| Q_{T-t}^{1/2} x \right|^2 = \left| Q_T^{1/2} x \right|^2 - \left| V_t^* Q_T^{1/2} x \right|^2,$$

hence putting $y = Q_T^{1/2}x$ we obtain

$$\left| Q_{T-t}^{1/2} Q_T^{-1/2} y \right|^2 = |y|^2 - |V_t^* y|^2.$$

Assume that $||V_t^*|| = 1$ for a certain $t \in (0,T)$. Since $\operatorname{im}\left(Q_T^{1/2}\right)$ is dense in H, there exists a sequence $y_n \in \operatorname{im}\left(Q_T^{1/2}\right)$, such that $|y_n| = 1$ and $|V_t^*y_n| \to 1$. Therefore,

$$\lim_{n \to \infty} \left| Q_{T-t}^{1/2} Q_T^{-1/2} y_n \right|^2 = \lim_{n \to \infty} \left(1 - |V_t^* y_n|^2 \right) = 0. \tag{2.16}$$

Let y_{n_k} be a subsequence converging weakly to $y \in H$. Since

$$\operatorname{im}\left(Q_{T-t}^{1/2}\right) \subset \operatorname{im}\left(Q_T^{1/2}\right), \quad t \leqslant T,$$

and

$$\left(Q_T^{-1/2}Q_{T-t}^{1/2}\right)^* = \overline{Q_{T-t}^{1/2}Q_T^{-1/2}},$$

we find that

$$Q_{T-t}^{1/2}Q_T^{-1/2}y_{n_k} \to \overline{Q_{T-t}^{1/2}Q_T^{-1/2}}y$$
, weakly,

and by (2.16) we obtain $\overline{Q_{T-t}^{1/2}Q_T^{-1/2}}y = 0$ and since $|V_t^*y| = 1$ we obtain $y \neq 0$. It follows that the range of the operator $Q_T^{-1/2}Q_{T-t}^{1/2}$ is not dense in H, which shows that (b) implies (c).

Finally, assume that (c) holds. Then (2.13) and Proposition B1 in [7] yield

$$\operatorname{im}\left(Q_{T-t}^{1/2}\right) = \operatorname{im}\left(Q_T^{1/2}\left(I - V_t V_t^*\right)^{1/2}\right).$$

Since $||V_t|| < 1$, the operator $I - V_t V_t^* : H \to H$ is an isomorphism, hence

$$\operatorname{im}\left(Q_{T-t}^{1/2}\right) = \operatorname{im}\left(Q_T^{1/2}\right), \quad t < T,$$

and (a) follows.

Remark 2.9. Necessary and sufficient conditions for (2.14) to hold are not known but it was proved to be satisfied in the following cases.
(a) If

$$\operatorname{im}(S_t) \subset \operatorname{im}\left(Q_t^{1/2}\right), \quad t > 0,$$

then (2.14) holds. It is known that the above condition is equivalent to the strong Feller property of the OU transition semigroup $R_t\phi(x) = \mathbb{E}\phi(Z_t^x)$, see [7] for details.

(b) Assume that the process (Z_t^x) admits a nondegenerate invariant measure ν and $\operatorname{im}(Q)$ is dense in H. Let $H_Q = \operatorname{im}(Q^{1/2})$ be endowed with the norm $|x|_Q = |Q^{-1/2}x|$. Assume

that H_Q is invariant for the semigroup (S_t) and its restriction to H_Q is a C_0 -semigroup in H_Q . Then (2.14) holds, see [10]. These assumptions are satisfied for any process (Z_t^x) with the transition semigroup analytic in $L^2(H,\nu)$, in particular they are satisfied for any reversible OU process.

We define the operator $B: Q_T^{1/2}(H) \to L^2(0,T;H)$,

$$Bx(t) = Q^{1/2} S_{T-t}^* Q_T^{-1/2} x, \quad t \in [0, T], \quad x \in Q_T^{1/2}(H).$$

The following simple Lemma has been proved in [11]:

Lemma 2.10. (a) The operator B with the domain $dom(B) = Q_T^{1/2}(H)$ extends to a bounded operator (still denoted by B) $B: H \to L^2(0,T;H)$. Moreover,

$$|Bx|_{L^2(0,T;H)} = |x|_H, \quad x \in H.$$

(b) Setting

$$H \ni x \to \mathcal{K}x(t) = K_t x \in L^2(0, T; H), \tag{2.17}$$

where

$$K_t = Q_t^{1/2} V_t^*, (2.18)$$

we have $\mathscr{K} = \mathscr{L}B$. In particular the operator $\mathscr{K} : H \to C(0,T;H)$ is bounded.

2.4. **Fundamentals on OU Bridge.** In the present subsection we give the definition and some basic properties of the OU Bridge.

Since $V_t^* = \overline{Q_t^{1/2} S_{T-t}^* Q_T^{-1/2}}$ is bounded, the operator K_t is of Hilbert-Schmidt type on H for each $t \in [0, T)$. Also, $\mathcal{K} : H \to L^2(0, T; H)$ is Hilbert-Schmidt.

Note that if K_t is defined by (2.18) then, in view of Lemma 2.2, the measurable function $K_tQ_T^{-1/2}$ is well defined for each $t \in [0, T]$. We will start from the definition of the process (\hat{Z}_t) ,

$$\hat{Z}_t = Z_t - K_t Q_T^{-1/2} Z_T, \quad t \in [0, 1), \quad \text{and} \quad \hat{Z}_1 = 0.$$

Proposition 2.11. (a) An H-valued Gaussian process (\hat{Z}_t) is independent of Z_T .

(b) The covariance operator \hat{Q}_t of \hat{Z}_t is given by

$$\hat{Q}_t = Q_t^{1/2} \left(I - V_t^* V_t \right) Q_t^{1/2}. \tag{2.19}$$

- (c) The process (\hat{Z}_t) is mean-square continuous on [0,T].
- (d) If, moreover, one of the equivalent conditions (a)-(c) of Lemma 2.8 holds then

$$\operatorname{im}\left(\hat{Q}_{t}^{1/2}\right) = \operatorname{im}\left(Q_{t}^{1/2}\right), \quad t \in (0, T). \tag{2.20}$$

Proof. Theorem 2.4 yields immediately (a) since $\hat{Z}_t = Z_t - \mathbb{E}(Z_t | Z_T)$. Invoking (c) of Theorem 2.4 with $C_X = Q_T$, $C_Y = Q_t$ and $T^* = K_t$ and (2.18) we obtain

$$\hat{Q}_t = Q_t - K_t K_t^* = Q_t^{1/2} (I - V_t^* V_t) Q_t^{1/2}, \quad t < T.$$

Using (2.11) we find easily that

$$\lim_{t \to 0} \operatorname{tr}\left(\hat{Q}_t\right) = 0. \tag{2.21}$$

To prove that

$$\lim_{t \to T} \operatorname{tr}\left(\hat{Q}_t\right) = 0,\tag{2.22}$$

we note first that

$$\operatorname{tr}\left(\hat{Q}_{t}\right) = \operatorname{tr}\left(\left(I - V_{t}^{*}V_{t}\right)\left(Q_{t} - Q_{T}\right)\right) + \operatorname{tr}\left(\left(I - V_{t}^{*}V_{t}\right)Q_{T}\right).$$

Next, it is easy to see that

$$0 \leqslant \lim_{t \to T} \operatorname{tr} \left((I - V_t^* V_t) (Q_T - Q_t) \right) \leqslant \lim_{t \to T} \operatorname{tr} \left(Q_T - Q_t \right) = 0.$$
 (2.23)

Finally,

$$\operatorname{tr}\left(\left(I - V_t^* V_t\right) Q_T\right) = \operatorname{tr}\left(Q_T\right) - \operatorname{tr}\left(V_t Q_T V_t^*\right)$$
$$= \operatorname{tr}\left(Q_T\right) - \sum_{k=0}^{\infty} \left|Q_T^{1/2} V_t^* e_k\right|^2,$$

where $\{e_k : k \ge 1\}$ is a CONS in H. Therefore,

$$\lim_{t \to T} \operatorname{tr} \left((I - V_t^* V_t) \, Q_T \right) = 0 \tag{2.24}$$

by Lemma 2.7 and the Dominated Convergence Theorem. Combining (2.23) and (2.24) we obtain (2.22) and, consequently, (c). Part (d) follows immediately from Lemma 2.8 and (2.19).

Proposition 2.12. The conditional distribution of the process (Z_t^x) in the space $H_2 = L^2(0,T;H)$ given Z_T^x is $N(\lambda,\overline{Q})$, where

$$\lambda(t) = S_t x + K_t Q_T^{-1/2} Z_T, \tag{2.25}$$

$$\overline{Q} = \tilde{Q} - \mathcal{K}\mathcal{K}^*, \tag{2.26}$$

where \tilde{Q} is the covariance operator of the process (Z_t^x) in H_2 , $\tilde{Q}: H \to H_2$,

$$[\tilde{Q}y](t) = \int_0^t R(t,s)y(s)ds, \quad y \in H_2,$$

and

$$R(t,s)z = \int_0^s S_{t-r}QS_{s-r}^*zdr, \quad z \in H, \quad 0 \leqslant s \leqslant t \leqslant T,$$

and $\mathcal{K}: H \to H_1$ is defined in (2.17).

Proof. We use Theorem 2.4 with $H_1 = H$, $H_2 = L^2(0, T; H)$, $X = Z_t^x$, $Y = (Z_t^x)$, $C_X = Q_T$, and $C_Y = \tilde{Q}$. By the definition of the covariance C_{XY} ,

$$\langle C_{XY}k, h \rangle_{L^2(0,T;H)} = \mathbf{E} \langle Z_T^x, k \rangle \langle Z^x, h \rangle_{L^2(0,T;H)}, \quad k \in H_1, h \in H_2,$$

it is easy to compute $[C_{XY}k](t) = Q_t S_{T-t}^* k$, $t \in [0,T]$. Hence we have $T^* = \overline{C_{XY}C_X^{-1/2}} = \mathcal{K}$ and $T: H_2 \to H_1$, $Ty = \mathcal{K}^* y = \int_0^T K_t^* y(t) dt$. By Theorem 2.4 we have that

$$\overline{Q} = C_Y - T^*T = \tilde{Q} - \mathcal{K}\mathcal{K}^*,$$

and

$$\lambda(t) = \mathbb{E}(Z_t^x | Z_T^x) = \mathbb{E}(S_t x + Z_t | Z_T^x) = \mathbb{E}(S_t x + \hat{Z}_t + K_t Q_t^{-1/2} Z_T | Z_T^x)$$

which yields $\lambda(t) = S_t x + K_t Q_T^{-1/2} Z_T$, because \hat{Z}_t and Z_T^x are stochastically independent, hence (2.25) and (2.26) hold true.

Recall that μ_T denotes the probability law of Z_T on H.

Proposition 2.13. There exists a Borel subspace $\mathcal{M} \subset H$ such that $\mu_T(\mathcal{M}) = 1$ and for all $x \in H$ and $y \in S_T x + \mathcal{M}$ the H-valued Gaussian process

$$\hat{Z}_{t}^{x,y} = Z_{t}^{x} - \mathcal{K}Q_{T}^{-1/2} \left(Z_{T}^{x} - y\right), \tag{2.27}$$

is well defined with paths in $L^2(0,T;H)$ and

$$\hat{Z}_{t}^{x,y} = S_{t}x - \mathcal{K}Q_{T}^{-1/2}(S_{T}x - y) + \hat{Z}_{t}, \quad \mathbb{P} - a.s.$$
 (2.28)

Proof. By Lemma 2.2 we can choose a measurable linear space \mathscr{M} such that $\mathscr{K}Q_T^{-1/2}$ is linear on \mathscr{M} and $\mu_T(\mathscr{M})=1$. Therefore, $\mathscr{K}Q_T^{-1/2}(Z_T^x-y)$ is well defined for any $y\in S_Tx+\mathscr{M}$ and (2.28) holds.

Theorem 2.14. Let $\Phi: L^2(0,T;H) \to \mathbb{R}$ be a Borel mapping such that

$$\mathbb{E}\left|\Phi\left(Z^{x}\right)\right|<\infty.$$

Then

$$\mathbb{E}\left(\Phi\left(Z^{x}\right)\middle|Z_{T}^{x}=y\right)=\mathbb{E}\Phi\left(\hat{Z}^{x,y}\right),\quad\mu_{T}^{x}-a.e.\tag{2.29}$$

where the left-hand side of (2.29) is defined as a function $g_{\Phi} = g_{\Phi}(y) \in L^1(H, \mu_T^x)$ such that $\mathbb{E}(\Phi(Z^x)|Z_T^x) = g_{\Phi}(Z_T^x)$ \mathbb{P} -a.s.

Proof. We have to show that

$$\mathbb{E}(\Phi(Z^x)|Z_T^x) = \mathbb{E}(\Phi(\hat{Z}^{x,y}))|_{Z_T^x = y} \quad \mathbb{P} - a.s.$$

By Proposition 2.12 we have

$$\mathbb{E}(\Phi(Z^x)|Z_T^x) = \int_H \Phi(z)N(\lambda, \overline{Q})(dz) \quad \mathbb{P} - a.s., \tag{2.30}$$

where λ and \overline{Q} are defined by (2.25) and (2.26), respectively. On the other hand, the covariance operator \hat{Q} of the process $\hat{Z}_t^{x,y}$ in H_2 is by (2.28) the same as the one of \hat{Z}_t .

Since $Z_t = \hat{Z}_t + K_t Q_T^{-1/2} Z_T$ and the summands on the right-hand side are independent random variables, we obtain $\tilde{Q} = \hat{Q} + \mathcal{K} \mathcal{K}^*$, that is, $\hat{Q} = \overline{Q}$. Also, we have

$$\mathbb{E}\hat{Z}_t^{x,y} = S_t x - \mathcal{K}Q_T^{-1/2}(S_T x - y),$$

and therefore

$$\mathbb{E}(\Phi(\hat{Z}^{x,y}))|_{Z_T^x=y} = \int_H \Phi(z) N(S_t x - K_t Q_T^{-1/2}(S_T x - y), \overline{Q})(dz)|_{Z_T^x=y} = \int_H \Phi(z) N(\lambda, \overline{Q})(dz)$$

 $\mathbb{P} - a.s.$, which together with (2.30) concludes the proof.

Definition 2.15. Given $x, y \in H$ and an H-valued OU process (Z_t^x) , a process $(\hat{Z}_t^{x,y})$ satisfying (2.29) is called an Ornstein-Uhlenbeck Bridge (connecting points x at time t = 0 and y at time t = T). The probability law of the process $(\hat{Z}_t^{x,y})$ in the space $L^2(0,T;H)$ will be denoted by $\hat{\mu}^{x,y}$.

Thus we have shown that the OU Bridge may be written in the form (2.27) or (2.28) and its probability law $\hat{\mu}^{x,y}$ is $N(\gamma, \overline{Q})$ where $\gamma(t) = \mathbb{E}[\lambda(t)|Z_T^x = y] = S_t x - K_t Q_T^{-1/2}(S_T x - y)$ $\mu_T^x - a.e.$

The following Theorem has been proved in [11]:

Theorem 2.16. Let \mathscr{E} be a Banach space such that $\mu(\mathscr{E}) = 1$. Then $\hat{\mu}^{0,y}(\mathscr{E}) = 1$ for $y \in \mathscr{M}$.

3. SDE ASSOCIATED TO THE OU BRIDGE

In the sequel we will need the following

Hypothesis 3.1. For any t > 0

$$\operatorname{im}\left(S_{t}Q^{1/2}\right) \subset \operatorname{im}\left(Q_{t}^{1/2}\right). \tag{3.1}$$

Remark 3.2. Condition (3.1) is satisfied in some important cases.

- (a) If the process (Z_t^x) is strong Feller then im $(S_t) \subset \operatorname{im}\left(Q_t^{1/2}\right)$ and therefore (3.1) holds.
- (b) Let $H_Q = Q^{1/2}(H)$ be endowed with the norm $|x|_Q = |Q^{-1/2}x|$, where Q is assumed to be nondegenerate. Assume that $S_tH_Q \subset H_Q$ for all $t \ge 0$ and (S_t) restricted to H_Q is a C_0 -semigroup. It was proved in [10] that in this case $S_t(H) \subset Q_t^{1/2}(H)$ for all t > 0 and there exists c > 0 such that

$$\left\| Q_t^{-1/2} S_t Q^{1/2} \right\| \leqslant \frac{c}{\sqrt{t}}, \quad t > 0.$$

Assume additionally that the process (Z_t^x) admits a Gaussian invariant measure ν . Then, cf. [10], (S_t) is a C_0 -semigroup on H_Q if the transition semigroup of the process (Z_t^x) is analytic on $L^2(H,\nu)$, in particular this holds for a symmetric Ornstein-Uhlenbeck process. Explicit conditions for the analyticity and symmetry of the transition semigroup of the process (Z_t^x) in $L^2(H,\nu)$ may be found in [10] and [6].

Lemma 3.3. Assume that Hypothesis 3.1 holds. Then the function

$$t \to \left| Q_t^{-1/2} S_t Q^{1/2} h \right|,$$

is nonincreasing on $(0, \infty)$ for each $h \in H$.

Proof. By Lemma 2.7 we have

$$\left\| Q_{t+s}^{-1/2} S_t Q_s^{1/2} \right\| \leqslant 1. \tag{3.2}$$

By assumption the operator $Q_{t+s}^{-1/2}S_{t+s}Q^{1/2}$ is well defined and bounded and $S_sQ^{1/2}h \in \operatorname{im}\left(Q_s^{1/2}\right)$. Therefore, by (3.2)

$$\left| Q_{t+s}^{-1/2} S_{s+t} Q^{1/2} h \right| = \left| Q_{t+s}^{-1/2} S_t Q_s^{1/2} Q_s^{-1/2} S_s Q^{1/2} h \right|$$

$$\leq \left| Q_s^{-1/2} S_s Q^{1/2} h \right|,$$

and (b) follows.

Let

$$Y_u = \int_u^T S_{T-s} Q^{1/2} dW_s, \quad u \leqslant T.$$

Since the operator-valued function $t \to Q_t$ is continuous in the weak operator topology and all the operators Q_t are compact for t > 0, there exists a measurable choice of eigenvectors $\{e_k(t) : k \ge 1\}$ and eigenvalues $\{\lambda_k(t) : k \ge 1\}$. For each $n \ge 1$ we define a process

$$X_u^n = \sum_{k=1}^n \frac{1}{\sqrt{\lambda_k(T-u)}} \langle Y_u, e_k(T-u) \rangle F_u^* e_k(T-u),$$

where $F_u = Q_{T-u}^{-1/2} S_{T-u} Q^{1/2}$.

Lemma 3.4. There exists a measurable stochastic process (X_u) defined on [0,T) such that for each a < T

$$\lim_{n \to \infty} \mathbb{E} \int_0^a |X_u^n - X_u|^2 \, du = 0. \tag{3.3}$$

and for each $h \in H$ and a < T the series

$$\langle X_u, h \rangle = \sum_{k=1}^{\infty} \frac{1}{\sqrt{\lambda_k(T-u)}} \langle Y_u, e_k(T-u) \rangle \langle e_k(T-u), F_u h \rangle$$
 (3.4)

converges in $L^2(0,a)$ in mean square. Moreover, if $0 \le u \le v < T$ then for all $h, k \in H$

$$\mathbb{E}\langle X_u, h \rangle \langle X_v, k \rangle = \left\langle F_u h, Q_{T-u}^{-1/2} Q_{T-v}^{1/2} F_v k \right\rangle, \tag{3.5}$$

where the operator $Q_{T-u}^{-1/2}Q_{T-v}^{1/2}$ is bounded.

Proof. For $u \leqslant v \leqslant T$

$$\mathbb{E}\langle Y_u, h \rangle \langle Y_v, k \rangle = \langle Q_{T-v}h, k \rangle, \quad h, k \in H.$$
(3.6)

Therefore

$$\mathbb{E} \langle X_u^n - X_u^m, h \rangle^2 = \sum_{j=m+1}^n \frac{1}{\lambda_k(T-u)} \mathbb{E} \langle Y_u, e_k(T-u) \rangle^2 \langle e_k(T-u), F_u h \rangle^2$$

$$= \sum_{j=m+1}^n \langle e_k(T-u), F_u h \rangle^2 \underset{n,m \to \infty}{\longrightarrow} 0, \tag{3.7}$$

hence the process

$$\langle X_u, h \rangle = \sum_{k=1}^{\infty} \frac{1}{\sqrt{\lambda_k(T-u)}} \langle Y_u, e_k(T-u) \rangle \langle e_k(T-u), F_u h \rangle = \left\langle Q_{T-u}^{-1/2} Y_u, F_u h \right\rangle$$

is well defined for each $h \in H$ and u < T. For u, v such that $0 < u \le v < T$ we have

$$\operatorname{im}\left(Q_{T-v}^{1/2}\right) \subset \operatorname{im}\left(Q_{T-u}^{1/2}\right). \tag{3.8}$$

Let P_n is an orthogonal projection on $\lim \{e_k(T-v): k \leq n\}$ and $F_u^n = P_n F_u$. Then $Q_{T-u}^{-1/2} F_u^n$ is bounded on H. Let

$$X_u^n = \left(Q_{T-u}^{-1/2} F_u^n \right)^* Y_u.$$

By (3.6)

$$\mathbb{E} \langle X_u^n, h \rangle \langle X_v^n k \rangle = \left\langle Q_{T-v} Q_{T-u}^{-1/2} F_u^n h, Q_{T-v}^{-1/2} F_v^n k \right\rangle$$
$$= \left\langle F_u^n h, Q_{T-u}^{-1/2} Q_{T-v}^{1/2} F_v^n k \right\rangle.$$

By (3.8) the operator $Q_{T-u}^{-1/2}Q_{T-v}^{1/2}$ is bounded and therefore

$$\mathbb{E}\left\langle Q_{T-u}^{-1/2}Y_u, F_u h \right\rangle \left\langle Q_{T-v}^{-1/2}Y_v, F_v k \right\rangle = \lim_{n \to \infty} \mathbb{E}\left\langle X_u^n, h \right\rangle \left\langle X_v^n, k \right\rangle$$
$$= \left\langle F_u h, Q_{T-u}^{-1/2} Q_{T-v}^{1/2} F_v k \right\rangle.$$

It follows from (3.5) that

$$\mathbb{E}\left\langle X_{u},h\right\rangle ^{2}=\left|F_{u}h\right|^{2},$$

and by Lemma 3.3 we obtain for $u \leq a$

$$\mathbb{E} \left\langle X_u^n, h \right\rangle^2 \leqslant \mathbb{E} \left\langle X_u, h \right\rangle^2 \leqslant |h|^2 \left\| F_{T-a} \right\|^2.$$

Then (3.7) and the Dominated Convergence Theorem yield

$$\lim_{n,m\to\infty} \int_0^a \sup_{|h|\le 1} \mathbb{E} \left\langle X_u^n - X_u^m, h \right\rangle^2 du = 0.$$

As a consequence we find that (3.3) holds for any $a \in (0, T)$.

By Lemma 3.3 a cylindrical process

$$I_t = \int_0^t F_u^* Q_{T-u}^{-1/2} Y_u du$$

is well defined, that is for any $h \in H$ the real-valued process

$$\langle I_t, h \rangle = \int_0^t \left\langle Q_{T-u}^{-1/2} Y_u, F_u h \right\rangle du$$

is well defined for all t < T.

Lemma 3.5. The cylindrical process

$$\zeta_t = W_t - \int_0^t F_u^* Q_{T-u}^{-1/2} Y_u du, \quad t \leqslant T,$$

is a standard cylindrical Wiener process on H.

The proof of this Lemma is omitted; it is a word by word repetition of the proof of Lemma 4.7 in [11] if we use Lemmas 3.3 and 3.4 above.

Theorem 3.6. For all t < T

$$\mathbb{E} \int_{0}^{t} \left| S_{t-s} Q^{1/2} F_{s}^{*} Q_{T-s}^{-1/2} S_{T-s} \hat{Z}_{s} \right|^{2} ds < \infty, \tag{3.9}$$

and

$$\hat{Z}_t = -\int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} S_{T-s} \hat{Z}_s ds + \int_0^t S_{t-s} Q^{1/2} d\zeta_s, \quad \mathbb{P} - a.s.$$
 (3.10)

Proof. We will show first that the operator $Q_{T-s}^{-1/2}S_{T-s}\hat{Q}_sS_{T-s}^*Q_{T-s}^{-1/2}$ is bounded. Let $h,k\in H$. Then by Proposition 2.11 and (2.10) we obtain

$$\left\langle S_{T-s} \hat{Q}_{s} S_{T-s}^{*} h, k \right\rangle = \left\langle S_{T-s} Q_{s} S_{T-s}^{*} h, k \right\rangle - \left\langle S_{T-s} Q_{s}^{1/2} V_{s}^{*} V_{s} Q_{s}^{1/2} S_{T-s}^{*} h, k \right\rangle$$

$$= \left\langle \left(Q_{T} - Q_{T-s} \right) h, k \right\rangle - \left\langle Q_{T}^{-1/2} S_{T-s} Q_{s} S_{T-s}^{*} h, Q_{T}^{-1/2} S_{T-s} Q_{s} S_{T-s}^{*} k \right\rangle$$

$$= \left\langle \left(Q_{T} - Q_{T-s} \right) h, k \right\rangle - \left\langle Q_{T}^{-1/2} \left(Q_{T} - Q_{T-s} \right) h, Q_{T}^{-1/2} \left(Q_{T} - Q_{T-s} \right) k \right\rangle$$

$$= \left\langle \left(Q_{T} - Q_{T-s} \right) h, k \right\rangle - \left\langle \left(Q_{T} - Q_{T-s} \right) Q_{T}^{-1} \left(Q_{T} - Q_{T-s} \right) h, k \right\rangle$$

$$= \left\langle \left(Q_{T} - Q_{T-s} \right) h, k \right\rangle - \left\langle \left(Q_{T} - Q_{T-s} \right) \left(I - Q_{T}^{-1} Q_{T-s} \right) h, k \right\rangle$$

$$= \left\langle \left(Q_{T-s} - Q_{T-s} Q_{T}^{-1} Q_{T-s} \right) h, k \right\rangle = \left\langle Q_{T-s}^{1/2} \left(I - Q_{T-s}^{1/2} Q_{T}^{-1} Q_{T-s}^{1/2} \right) Q_{T-s}^{1/2} h, k \right\rangle .$$

Since the operator $Q_{T-s}^{1/2}Q_T^{-1}Q_{T-s}^{1/2}$ is bounded for s < T we find that the operator

$$T_s = Q_{T-s}^{-1/2} S_{T-s} \hat{Q}_s S_{T-s}^* Q_{T-s}^{-1/2} = I - Q_{T-s}^{1/2} Q_T^{-1} Q_{T-s}^{1/2}$$
(3.11)

is bounded as well. Therefore, for $s \leq T - \epsilon$ Lemma 3.3 and (3.11) yield

$$\mathbb{E} \left| S_{t-s} Q^{1/2} F_s^* \left(Q_{T-s}^{-1/2} S_{T-s} \hat{Z}_s \right) \right|^2$$

$$= \left\| S_{t-s} Q^{1/2} F_s^* \left(Q_{T-s}^{-1/2} S_{T-s} \hat{Q}_s^{1/2} \right) \right\|_{HS}^2 \leqslant \left\| S_{t-s} Q^{1/2} \right\|_{HS}^2 \left\| F_s \right\|^2 \left\| T_s^{1/2} \right\|$$

$$\leqslant \left\| S_{t-s} Q^{1/2} \right\|_{HS}^2 \left\| F_{T-\epsilon} \right\|^2,$$

which completes the proof of (3.9). As a byproduct of the argument given above we proved also that the process $Q_{T-s}^{-1/2}S_{T-s}\hat{Z}_s$ is well defined for all $s \leq T$. Now, we are ready to prove (3.10). By Lemma 3.5 we have

$$\hat{Z}_t = Z_t - K_t Q_T^{-1/2} Z_T$$

$$= \int_0^t S_{t-s} Q^{1/2} d\zeta_s + \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{1-s}^{-1/2} Y_s ds - K_t Q_T^{-1/2} Z_T,$$

and since

$$Y_s = Z_T - S_{T-s}Z_s = Z_T - S_{T-s}K_sQ_T^{-1/2}Z_T - S_{T-s}\hat{Z}_s,$$

we find that

$$\hat{Z}_t = \int_0^t S_{t-s} Q^{1/2} d\zeta_s - \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} S_{T-s} \hat{Z}_s ds$$

$$+ \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} \left(Z_T - S_{T-s} K_s Q_T^{-1/2} Z_T \right) ds - K_t Q_T^{-1/2} Z_T.$$

It remains to show that

$$\int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} \left(Z_T - S_{T-s} K_s Q_T^{-1/2} Z_T \right) ds - K_t Q_T^{-1/2} Z_T = 0.$$
 (3.12)

To this end note first that

$$K_t Q_T^{-1/2} Z_T = \left(\int_0^t S_{t-s} Q^{1/2} F_s^* ds \right) Q_T^{-1/2} Z_T, \tag{3.13}$$

and

$$S_{T-t}K_tQ_T^{-1/2}Z_T = \left(\int_0^t S_{T-s}Q^{1/2}F_s^*ds\right)Q_T^{-1/2}Z_T$$
$$= (Q_T - Q_{T-t})Q_T^{-1}Z_T = Z_T - Q_{T-t}Q_T^{-1}Z_T,$$

and thereby

$$Z_T - S_{T-t}K_tQ_T^{-1/2}Z_T = Q_{T-t}Q_T^{-1}Z_T. (3.14)$$

Finally, (3.14) and the definition of F_s^* give

$$\int_{0}^{t} S_{t-s} Q^{1/2} F_{s}^{*} Q_{T-s}^{-1/2} \left(Z_{T} - S_{T-s} K_{s} Q_{T}^{-1/2} Z_{T} \right) ds$$

$$= \int_{0}^{t} S_{t-s} Q^{1/2} F_{s}^{*} Q_{T-s}^{-1/2} Q_{T-s} Q_{T}^{-1} Z_{T} ds = \left(\int_{0}^{t} S_{t-s} Q^{1/2} F_{s}^{*} ds \right) Q_{T}^{-1/2} Z_{T},$$
and (3.12) follows from (3.13).

We will consider now the general case of the bridge $(\hat{Z}_t^{x,y})$ connecting points $x \in H$ and y. We will impose the stronger condition (2.14) which is now formulated as a separate hypothesis:

Hypothesis 3.7. For any $t \in (0,T]$

$$\operatorname{im}\left(Q_t^{1/2}\right) = \operatorname{im}\left(Q_T^{1/2}\right).$$

For $y \in H_1 := \operatorname{im}(Q_T^{1/2})$ we define

$$Ny(t) = \int_0^t S_{t-s}Q^{1/2}F_s^*Q_{T-s}^{-1/2}yds, \quad t \leqslant T - \epsilon.$$

Theorem 3.8. Assume that Hypotheses 3.1 and 3.7 hold. Then the following holds.

- (a) The operator $N: H_1 \to L^2(0, T \epsilon; H)$ is Hilbert-Schmidt.
- (b) For any $x \in H$ and $y \in \mathcal{M}$

$$\hat{Z}_{t}^{x,y} = S_{t}x - \int_{0}^{t} S_{t-s}Q^{1/2}F_{s}^{*}Q_{T-s}^{-1/2}S_{T-s}\hat{Z}_{s}^{x,y}ds + \int_{0}^{t} S_{t-s}Q^{1/2}d\zeta_{s} + \int_{0}^{t} S_{t-s}Q^{1/2}F_{s}^{*}Q_{T-s}^{-1/2}yds.$$
(3.15)

Proof. Recall that by Lemma 2.10 (b) we have $\mathcal{K} = \mathcal{L}B$, hence for $z \in \mathcal{M}$

$$K_t Q_T^{-1/2} z = \int_0^t S_{t-s} Q^{1/2} B_s Q_T^{-1/2} z ds.$$
 (3.16)

Next, for $s \leq T - \epsilon$

$$\sup_{s \le T - \epsilon} \left\| Q_{T-s}^{-1/2} Q_T^{1/2} \right\| = \left\| Q_{\epsilon}^{-1/2} Q_T^{1/2} \right\| < \infty,$$

and invoking Lemma 3.3 we find that

$$\|NQ_{T}^{1/2}\|_{HS}^{2} \leqslant \int_{0}^{T-\epsilon} \|\int_{0}^{t} S_{t-s}Q^{1/2}F_{s}^{*}Q_{T-s}^{-1/2}Q_{T}^{1/2}ds\|_{HS}^{2} dt$$

$$\leqslant \left(\int_{0}^{T} \|S_{s}Q^{1/2}\|_{HS}^{2} ds\right) \left(\int_{0}^{T-\epsilon} \|F_{s}^{*}Q_{T-s}^{-1/2}Q_{T}^{1/2}\|^{2} ds\right)$$

$$\leqslant \|F_{T-\epsilon}\|^{2} \|Q_{\epsilon}^{-1/2}Q_{T}^{1/2}\|^{2} \left(\int_{0}^{T} \|S_{s}Q^{1/2}\|_{HS}^{2} ds\right) < \infty.$$

Therefore, the measurable function

$$y \to \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} y ds,$$

is well defined. We are ready now for the proof of (3.15). Let $x, y \in \text{im}\left(Q_T^{1/2}\right)$. Then Hypothesis 3.7 yields $S_T x \in \text{im}\left(Q_T^{1/2}\right)$, hence $y \in \mathcal{M}$. By (2.28) we have

$$\hat{Z}_{t}^{x,y} = \hat{Z}_{t} + S_{t}x - K_{t}Q_{T}^{-1/2} (S_{T}x - y),$$

and Theorem 3.6 yields

$$\hat{Z}_t^{x,y} = S_t x - K_t Q_T^{-1/2} S_T x + K_t Q_T^{-1/2} y$$

$$-\int_{0}^{t} S_{t-s}Q^{1/2}F_{s}^{*}Q_{T-s}^{-1/2}S_{T-s}\hat{Z}_{s}ds + \int_{0}^{t} S_{t-s}Q^{1/2}d\zeta_{s}$$

$$= S_{t}x - K_{t}Q_{T}^{-1/2}S_{T}x + K_{t}Q_{T}^{-1/2}y$$

$$-\int_{0}^{t} S_{t-s}Q^{1/2}F_{s}^{*}Q_{T-s}^{-1/2}S_{T-s}\left(\hat{Z}_{s}^{x,y} - S_{s}x + K_{s}Q_{T}^{-1/2}S_{T}x - K_{s}Q_{T}^{-1/2}y\right)ds + \int_{0}^{t} S_{t-s}Q^{1/2}d\zeta_{s}$$

$$= -K_{t}Q_{T}^{-1/2}S_{T}x + \int_{0}^{t} S_{t-s}Q^{1/2}F_{s}^{*}Q_{T-s}^{-1/2}S_{T-s}\left(S_{s} - K_{s}Q_{T}^{-1/2}S_{T}\right)xds$$

$$+K_{t}Q_{T}^{-1/2}y + \int_{0}^{t} S_{t-s}Q^{1/2}F_{s}^{*}Q_{T-s}^{-1/2}S_{T-s}K_{s}Q_{T}^{-1/2}yds$$

$$+S_{t}x - \int_{0}^{t} S_{t-s}Q^{1/2}F_{s}^{*}Q_{T-s}^{-1/2}S_{T-s}\hat{Z}_{s}^{x,y}ds + \int_{0}^{t} S_{t-s}Q^{1/2}d\zeta_{s}$$

$$=: H_{t}x + G_{t}y + S_{t}x - \int_{0}^{t} S_{t-s}Q^{1/2}F_{s}^{*}Q_{T-s}^{-1/2}S_{T-s}\hat{Z}_{s}^{x,y}ds + \int_{0}^{t} S_{t-s}Q^{1/2}d\zeta_{s}. \tag{3.17}$$

We will show first that

$$G_t y = \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} y ds.$$
 (3.18)

For $y \in \operatorname{im}\left(Q_T^{1/2}\right)$

$$S_{T-t}K_t y = \int_0^t S_{t-s} Q S_{T-s}^* Q_T^{-1/2} y ds = \int_0^t S_{T-s} Q S_{T-s}^* Q_T^{-1/2} y ds$$
$$= (Q_T - Q_{T-t}) Q_T^{-1/2} y, \tag{3.19}$$

and therefore

$$\begin{split} F_s^*Q_{T-s}^{-1/2}S_{T-s}K_sy &= F_s^*Q_{T-s}^{-1/2}Q_T^{1/2}y - F_s^*Q_{T-s}^{1/2}Q_T^{-1/2}y \\ &= F_s^*Q_{T-s}^{-1/2}Q_T^{1/2}y - Q^{1/2}S_{T-s}^*Q_T^{-1/2}y. \end{split}$$

Hence, taking Lemma 2.10 (b) into account we find that

$$G_t y = K_t Q_T^{-1/2} y + \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} S_{T-s} K_s Q_T^{-1/2} y ds$$
$$= K_t Q_T^{-1/2} y + \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} y ds - K_t Q_T^{-1/2} y,$$

and (3.18) follows. Next, we claim that for $x \in \text{im}\left(Q_T^{1/2}\right)$

$$H_t x = 0. (3.20)$$

Indeed, using (3.19) we obtain

$$H_t x = -K_t Q_T^{-1/2} S_T x + \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} S_{T-s} \left(S_s - K_s Q_T^{-1/2} S_T \right) x ds$$

$$= -K_t Q_T^{-1/2} x + \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} S_T x ds - \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} S_{T-s} K_s Q_T^{-1/2} S_T x ds$$

$$= -K_t Q_T^{-1/2} x + \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} S_T x ds$$

$$- \int_0^t S_{t-s} Q^{1/2} F_s^* Q_{T-s}^{-1/2} \left(Q_T - Q_{T-t} \right) Q_T^{-1/2} S_T x ds = 0,$$

which yields (3.20) for $x \in \text{im}\left(Q_T^{1/2}\right)$ and therefore for all $x \in H$. Finally, combining (3.17), (3.18) and (3.20) we obtain (3.15).

Corollary 3.9. Assume Hypotheses 3.1 and 3.7. Then for each t < T, and $h \in \text{dom}(A^*)$ and all $x \in H$ and $y \in \mathcal{M}$

$$\left\langle \hat{Z}_{t}^{x,y}, h \right\rangle = \left\langle x, h \right\rangle + \int_{0}^{t} \left\langle \hat{Z}_{s}^{x,y}, A^{*}h \right\rangle ds - \int_{0}^{t} \left\langle F_{s}^{*} Q_{T-s}^{-1/2} S_{T-s} \hat{Z}_{s}^{x,y}, Q^{1/2}h \right\rangle ds + \int_{0}^{t} \left\langle F_{s}^{*} Q_{T-s}^{-1/2} y, Q^{1/2}h \right\rangle ds + \left\langle \zeta_{t}, Q^{1/2}h \right\rangle.$$

Proof. On any interval $[0, T_0]$ with $T_0 < T$ and for any $y \in \mathcal{M}$ the functions

$$s \to Q^{1/2} F_s^* Q_{T-s}^{-1/2} S_{T-s} \hat{Z}_s^{x,y}$$
 and $s \to Q^{1/2} F_s^* Q_{T-s}^{-1/2} y$

are \mathbb{P} -a.s. Bochner integrable by Theorem 3.8 and therefore standard results about the equivalence of weak and strong solutions of deterministic and stochastic evolution equations can be applied to prove the corollary, see for example [1] for deterministic and [2], [19] for stochastic versions.

4. Applications to Semilinear Equations

In this Section, transition densities and Markov semigroups defined by semilinear stochastic equations are studied using the OU Bridge. Throughout the Section we assume (beside (2.5)) that the OU process (Z_t^x) is strongly Feller, that is, the condition

$$\operatorname{im}(S_t) \subset \operatorname{im}(Q_t^{1/2}), \quad t \in (0, T), \tag{4.1}$$

is satisfied. Note that (4.1) trivially implies the preceding Hypotheses 3.1 and 3.7 (or (2.14)). Let $(\mathscr{P}, \|.\|_{var})$ denote the space of probability measures on the Borel sets of H endowed with the metric of total variation. We start from a simple proposition where some continuity properties of the OU Bridge are given.

Proposition 4.1. (a) For each $t \in (0,T)$, $y \in \mathcal{M}$, where \mathcal{M} has been defined in Proposition 2.13, the mappings

$$x \mapsto \hat{Z}_t^{x,y}(\omega), \quad H \to H,$$
 (4.2)

$$x \mapsto \hat{Z}^{x,y}(\omega), \quad H \to L^2(0,T;H),$$
 (4.3)

are continuous for \mathbb{P} -almost all $\omega \in \Omega$, and the mapping

$$x \mapsto \hat{\mu}_t^{x,y}, \quad H \to (\mathscr{P}, \|.\|_{var}),$$
 (4.4)

is continuous.

(b) If, moreover, for each $t \in (0,T)$ we have $K_tQ_T^{-1/2} \in \mathcal{L}(\hat{H},H)$, where \hat{H} is a separable Banach space continuously embedded into H, then the mapping $y \mapsto \hat{Z}_t^{x,y}(\omega)$ is $\hat{H} \to H$ \mathbb{P} -a.s. continuous. Similarly, if

$$\overline{\mathcal{K}Q_T^{-1/2}} \in \mathcal{L}(\hat{H}, L^2(0, T; H)) \tag{4.5}$$

then $\mathcal{M} \supset \hat{H}$ and the mapping $y \mapsto \hat{Z}^{x,y}(\omega)$ is \mathbb{P} -a.s. $\hat{H} \to L^2(0,T;H)$ continuous.

Proof. (a) By (4.1) we have that $S_Tx \in \operatorname{im}(Q_T^{1/2})$ for each $x \in H$ and hence $S_Tx \in \mathcal{M}$ by construction of \mathcal{M} , hence $y \in \mathcal{M}$. Furthermore, (4.1) implies that the mappings $\mathcal{K}Q_T^{-1/2}S_T$ and $K_tQ_T^{-1/2}S_T$, $t \in (0,T]$, are in $\mathcal{L}(H,L^2(0,T;H))$ and $\mathcal{L}(H)$, respectively, and (4.2) and (4.3) follow by (2.28). To show (4.4) we recall Proposition 2.11 and Lemma 2.8, by which we have $\operatorname{im}(\hat{Q}_t^{1/2}) = \operatorname{im}(Q_t^{1/2})$. Hence the measures $(\hat{\mu}_T^{x,y}), x \in H$, are equivalent and

$$\psi^{y}(t,x,z) = \frac{d\mu_{t}^{x,y}}{d\mu_{t}^{0,y}}(z) = \exp\left(-\frac{1}{2}\left|Q_{t}^{-1/2}S_{t}x\right|^{2} + \frac{1}{2}\left|Q_{T}^{-1/2}S_{T}x\right|^{2} + \left\langle Q_{t}^{-1/2}z, Q_{t}^{-1/2}S_{t}x\right\rangle\right). \tag{4.6}$$

Indeed, by the Cameron-Martin formula we have

$$\psi^{y}(t, x, z) = \exp\left(-\frac{1}{2} \left| \hat{Q}_{t}^{-1/2} m \right|^{2} + \left\langle \hat{Q}_{t}^{-1/2} z, \hat{Q}_{t}^{-1/2} m \right\rangle\right),$$

where $m = Q_t^{1/2} (I - V_t^* V_t) Q_t^{-1/2} S_t x$. Then using (2.19) we get (4.6) and the assertion easily follows.

The proof of part (b) is completely analogous.

Remark 4.2. (a) The equivalent form of the density (4.6) is

$$\psi^{y}(t,x,z) = \exp\left(-\frac{1}{2}\left| (I - V_{t}^{*}V_{t})^{1/2} Q_{t}^{-1/2} S_{t} x \right|^{2} + \left\langle Q_{t}^{-1/2} z, Q_{t}^{-1/2} S_{t} x \right\rangle \right).$$

(b) Note that the OU Bridge $(\hat{Z}_t^{x,y})$ satisfies the SDE (3.15) which defines an (inhomogeneous) Markov process on the interval (0,T). By (4.4) this process is strongly Feller.

Now consider a stochastic semilinear evolution equation of the form

$$dX_t = AX_t dt + F(X_t) dt + \sqrt{Q} dW_t, \quad X_0 = x \in H$$
(4.7)

where A, W_t and Q are as before and $F: H \to H$ is a nonlinear continuous mapping. Suppose that $\operatorname{im}(F) \subset \operatorname{im}(Q^{1/2})$ and set $G:=Q^{-1/2}F$.

Hypothesis 4.3. The mapping $G: H \to H$ is bounded and continuous.

Now we formulate technical assumptions on the linear part of the equation. For simplicity of presentation, it is stated in the form that is verifiable in examples and includes all assumptions made previously in the paper.

Hypothesis 4.4. Assume either

- (i) $\dim H < \infty$ or
- (ii) There exist $\alpha \in (0,1)$ and $\beta < \frac{1+\alpha}{2}$ such that

$$\int_0^{T_0} t^{-\alpha} ||S_t Q^{1/2}||_{HS}^2 dt < \infty \quad and$$

$$||Q_t^{-1/2}S_t|| \leqslant \frac{c}{t^{\beta}}, \quad t \in (0, T_0),$$

for some c > 0 and $T_0 > 0$.

Conditions from (ii) are often used in the theory of stochastic equations and have been widely studied (cf.[7] or [11], see also the Example below). Note that Hypothesis 4.4 (ii) implies all previous assumptions made in the paper on the linear part of the equation (4.7) (i.e., all except for Hypothesis 4.3).

It is well known (see e.g. [19]) that under Hypotheses 4.3 and 4.4 the equation (4.7) defines an H-valued Markov process induced by the mild formula

$$X_{t} = S_{t}x + \int_{0}^{t} S_{t-r}F(X_{r})dr + \int_{0}^{t} S_{t-r}\sqrt{Q}d\widetilde{W}_{r}, \quad t \geqslant 0,$$
(4.8)

where \widetilde{W}_t is a standard cylindrical Wiener process on H defined on a suitable probability space.

Finally, we assume that the OU process defined by the linear equation (1.1) has an invariant measure ν that will be used as a reference measure. This is equivalent to the condition

$$\sup_{t>0} tr(Q_t) < \infty. \tag{4.9}$$

If (4.9) holds then ν is a centered Gaussian measure with the covariance operator

$$Q_{\infty} = \int_0^{\infty} S_t Q S_t^* dt.$$

Moreover, it has been shown in [4] that $S_tQ_{\infty}^{1/2}(H)\subset Q_{\infty}^{1/2}(H)$ and the family of operators

$$S_0(t) = Q_{\infty}^{-1/2} S_t Q_{\infty}^{1/2}, \quad t \geqslant 0,$$

defines a C_0 -semigroup of contractions on H. Moreover, if part (ii) of Hypothesis 4.4 holds then $||S_0(t)|| < 1$ for all t > 0.

Denote by (P_t) the transition Markov semigroup defined by the equation (4.7) and set

$$P(t, x, \Gamma) = P_t 1_{\Gamma}(x), x \in H, t > 0$$

and Γ Borel sets in H, and

$$d(t, x, y) = \frac{P(t, x, dy)}{\nu(dy)}.$$

It is standard to see that the density d exists, because Girsanov Theorem may be used to show the equivalence of measures $P(t, x, dy) \sim \mu_t^x$, and $\mu_t^x \sim \nu$ by (4.1) (see e.g. [11]).

Theorem 4.5. Let Hypotheses 4.3, 4.4 and (4.9) be satisfied and let T > 0 be fixed. Then for ν -almost all $y \in H$ the mapping $x \mapsto d(T, x, y)$ is continuous on H.

Theorem 4.6. Let Hypotheses 4.3, 4.4 and (4.9) be satisfied. Then for p > 1, T > 0, we have

$$P_T(L^p(H,\nu)) \subset \mathscr{C}(H),$$

that is, the semigroup (P_t) maps the space $L^p(H,\nu)$ into the space of continuous functions on H.

For p, q > 1 we introduce the notation

$$||P_t||_{p,q} = \left(\int_H \left(\int_H d^{p'}(t,x,y)\nu(dy)\right)^{q/p'} \nu(dx)\right)^{1/q},$$

where $p' = \frac{p}{p-1}$. Note that $||P_t||_{2,2}$ is a Hilbert-Schmidt norm of P_t . Moreover, if $||P_t||_{p,q} < \infty$ then the operator $P_t: L^p(H,\nu) \to L^q(H,\nu)$ is compact. Under assumptions more general than ours necessary and sufficient conditions were given in [3] for boundedness of the operator $P_t: L^p(H,\nu) \to L^q(H,\nu)$. In the theorem below we use different arguments based on the formula for transition densities to show that a stronger property holds: $||P_t||_{p,q} < \infty$.

Theorem 4.7. Let Hypotheses 4.3, 4.4 and (4.9) be satisfied. Then for any fixed T > 0 and q > 0 satisfying

$$q < 1 + \frac{p-1}{\|S_0(T)\|^2}$$

we have $||P_T||_{p,q} < \infty$. In particular, the operator $P_T : L^p(H,\nu) \to L^q(H,\nu)$ is q-summing and P_T is Hilbert-Schmidt in the space $L^2(H,\nu)$.

By the above mentioned equivalence of probabilities we may write

$$d(T, x, y) = \frac{P(T, x, dy)}{\mu_T^x(dy)} \cdot \frac{\mu_T^x(dy)}{\mu_T^0(dy)} \cdot \frac{\mu_T^0(dy)}{\nu(dy)}$$
(4.10)

$$=: h(T, x, y) \cdot g(T, x, y) \cdot k(T, y), \tag{4.11}$$

where k does not depend on x, g is given by the Cameron-Martin formula

$$g(T, x, y) = \exp\{\left\langle x, \overline{S_T^* Q_T^{-1/2}} Q_T^{-1/2} y \right\rangle - \frac{1}{2} |Q_T^{-1/2} S_T x|^2\}$$
(4.12)

for ν -almost all $y \in H$, and h may be expressed by means of the OU Bridge $(\hat{Z}_t^{x,y})$,

$$h(T, x, y) = \mathbb{E} \exp\{\rho(\hat{Z}^{x,y}) - \int_0^T \left\langle G(\hat{Z}_s^{x,y}), B_1(s)\hat{Z}_s + B_2(s)x - B_3(s)y \right\rangle ds\}$$
(4.13)

(cf.[11], Theorem 5.2), where

$$\rho(\hat{Z}^{x,y}) = \int_0^T \left\langle G(\hat{Z}_s^{x,y}), dW_s \right\rangle - \frac{1}{2} \int_0^T |G(\hat{Z}_s^{x,y})|^2 ds$$

and

$$B_1(s) = (Q_{T-s}^{-1/2} S_{T-s} Q^{1/2})^* Q_{T-s}^{-1/2} S_{T-s},$$

$$B_2(s) = (Q_T^{-1/2} S_{T-s} Q^{1/2})^* Q_T^{-1/2} S_T,$$

$$B_3(s) y = (Q_T^{-1/2} S_{T-s} Q^{1/2})^* Q_T^{-1/2} y, \quad y \in \operatorname{im} \left(Q_T^{1/2} \right).$$

From Lemma 2.10 it follows that

$$\int_0^T |B_2(s)x|^2 ds = |Q_T^{-1/2} S_T x|^2, \quad x \in H,$$
(4.14)

and by [11], Proposition 4.9, we have that

$$\mathbb{E} \int_0^T |B_1(s)\hat{Z}_t| ds < \infty \tag{4.15}$$

and

$$\int_0^T |B_3(s)y| ds < \infty \tag{4.16}$$

for ν - almost all $y \in \mathcal{M}$ (with no loss of generality we may assume that (4.16) holds for all $y \in \mathcal{M}$, $\nu(\mathcal{M}) = 1$). The proofs of Theorems 4.5, 4.6 and 4.7 are based on the following technical lemma:

Lemma 4.8. Given T > 0 and $q \in [0, \infty)$, there exists a constant $k_q > 0$ such that

$$h_{q}(T, x, y) := \mathbb{E} \exp\{q(\rho(\hat{Z}^{x,y}) - \int_{0}^{T} \left\langle G(\hat{Z}_{s}^{x,y}), B_{1}(s)\hat{Z}_{s} + B_{2}(s)x - B_{3}(s)y \right\rangle ds)\}$$

$$\leqslant k_{q} \exp\{k_{q}(|x| + \int_{0}^{T} |B_{3}(s)y|ds)\}$$
(4.17)

for all $x \in H$ and $y \in \mathcal{M}$, in particular,

$$h(t, x, y) \le k_1 \exp\{k_1(|x| + \int_0^T |B_3(s)y|ds)\}.$$

Proof. By the Cauchy inequality we have

$$h_{q}(T, x, y) \leq (\mathbb{E} \exp\{2q\rho(\hat{Z}^{x,y}\})^{1/2}$$

$$\times (\mathbb{E} \exp\{2q(\int_{0}^{T} |\langle G(\hat{Z}_{s}^{x,y}), B_{1}(s)\hat{Z}_{s} + B_{2}(s)x - B_{3}(s)y\rangle |ds)\})^{1/2}$$

$$(4.18)$$

and since the process $s \mapsto G(\hat{Z}_s^{x,y})$ is bounded the first expectation on the right-hand side of (4.18) is bounded (uniformly w.r.t. x and y). By (4.14) and (4.16) we thus have

$$h_q(T, x, y) \le C_q(\mathbb{E}\exp\{C_q \int_0^T (|B_1(s)\hat{Z}_s| + |B_2(s)x| + |B_3(s)y|)ds\})^{1/2}$$
 (4.19)

$$\leq \tilde{C}_q \exp{\{\tilde{C}_q(|Q_t^{-1/2}S_Tx| + \int_0^T |B_3(s)y|ds)\}} (\mathbb{E}\exp{\{\tilde{C}_q\int_0^T |B_1(s)\hat{Z}_s|ds\}})^{1/2}$$

for some C_q , \tilde{C}_q , and (4.17) follows by (4.15) and the Fernique inequality.

Proof of Theorem 4.5. Without loss of generality (dropping, if necessary, a set of ν -measure zero) we may suppose that g(T, x, y) and k(T, y) are defined for all $y \in \mathcal{M}$. By (4.12) we have that the mapping $x \mapsto g(T, x, y)k(T, y)$ is continuous, so we only have to prove continuity of the mapping $x \mapsto h(T, x, y), y \in \mathcal{M}, T > 0$. Let $x_n \to x_0$ in H. First we show (possibly, for a subsequence) that

$$\lim_{n \to \infty} \exp\{\rho(\hat{Z}^{x_n, y}) - \int_0^T \left\langle G(\hat{Z}_s^{x_n, y}), B_1(s)\hat{Z}_s + B_2(s)x_n - B_3(s)y \right\rangle ds\}$$

$$= \exp\{\rho(\hat{Z}^{x_0, y}) - \int_0^T \left\langle G(\hat{Z}_s^{x_0, y}), B_1(s)\hat{Z}_s + B_2(s)x_0 - B_3(s)y \right\rangle ds\}$$
(4.20)

 \mathbb{P} -a.s. We have

$$\int_{0}^{T} \left| \left\langle G(\hat{Z}_{s}^{x_{n},y}), B_{1}(s) \hat{Z}_{s} + B_{2}(s) x_{n} - B_{3}(s) y \right\rangle - \int_{0}^{T} \left\langle G(\hat{Z}_{s}^{x_{0},y}), B_{1}(s) \hat{Z}_{s} + B_{2}(s) x_{0} - B_{3}(s) y \right\rangle \right| ds$$

$$\leq \int_{0}^{T} \left| G(\hat{Z}_{s}^{x_{n},y}) - G(\hat{Z}_{s}^{x_{0},y}) \right| (|B_{1}(s) \hat{Z}_{s}| + |B_{2}(s) x_{0}| + |B_{3}(s) y|) ds$$

$$+ \int_{0}^{T} \left| G(\hat{Z}_{s}^{x_{0},y}) \right| \cdot |B_{2}(s) (x_{n} - x_{0}) ds, \tag{4.21}$$

which tends to zero by continuity and boundedness of G, (4.14) and Dominated Convergence Theorem. Also, we have

$$\mathbb{E}|\rho(\hat{Z}^{x_n,y}) - \rho(\hat{Z}^{x_0,y})| \leqslant C\left(\left(\mathbb{E}\int_0^T |G(\hat{Z}_s^{x_n,y}) - G(\hat{Z}_s^{x_0,y})|^2 ds\right)^{1/2} + \mathbb{E}\int_0^T |G(\hat{Z}_s^{x_n,y}) - G(\hat{Z}_s^{x_0,y})|^2 ds\right),$$

which again tends to zero by Dominated Convergence Theorem, so there is a subsequence converging \mathbb{P} -a.s. Taking into account (4.21) we obtain (4.20). By (4.17) (used, for instance, with q=2) the random variables on the left-hand side of (4.20) are integrable uniformly in n, hence the convergence in (4.20) holds also in the space $L^1(\Omega)$ and, consequently, we obtain $h(T, x_n, y) \to h(T, x_0, y)$. Since we may choose a subsequence with this property from an arbitrary sequence $x_n \to x_0$, the convergence takes place for the whole sequence.

Proof of Theorem 4.6. Let T > 0, $\phi \in L^p(H, \nu)$ and $x_n \to x_0$ in H. Then

$$|P_T\phi(x_n) - P_T\phi(x_0)| \leqslant \int_H |\phi(y)| |d(T, x_n) - d(T, x_0, y)| \nu(dy)$$

$$\leqslant (\int_H |\phi|^p d\nu)^{1/p} (\int_H |d(T, x_n, y) - d(T, x_0, y)|^{p'} \nu(dy))^{1/p'},$$

so by Theorem 4.5 it suffices to show that

$$\int_{H} (d(T, x_n, y))^q \nu(dy) < c_q, \quad q \in (1, \infty),$$
(4.22)

where c_q does not depend on n. The same property (uniform boundedness in arbitrary $L^q(H,\nu)$) has been shown for Gaussian densities $g(T,x_n,\cdot)$ and $k(T,\cdot)$ in [5], so we only have to show (4.22) where $d(T,x_n,y)$ is replaced by $h(T,x_n,y)$. However, by Lemma 4.8 and Hölder inequality we have

$$\int_{H} (h(T, x_{n}, y))^{q} \nu(dy) \leqslant \int_{H} h_{q}(T, x_{n}, y) \nu(dy)$$

$$\leqslant k_{q} \exp\{k_{q}|x_{n}|\} \int_{H} \exp\{\int_{0}^{T} |B_{3}(s)y|ds\} \nu(dy) < c_{q}$$
(4.23)

where c_q does not depend on n, since the sequence x_n is obviously bounded and

$$\int_{H} \exp\{\int_{0}^{T} |B_{3}(s)y|ds\}\nu(dy) < \infty$$

by (4.16), (4.1) and the Fernique inequality.

Proof of Theorem 4.7. We can rewrite (4.10) in the form

$$d(T, x, Y) = h(T, x, y)H(T, x, y),$$

where

$$H(T, x, y) = \frac{\mu_T^x(dy)}{\nu(dy)}.$$

Invoking the Hölder inequality we obtain

$$||P_{T}\phi||_{p,q}^{q} = \int_{H} \left(\int_{H} h H \phi \nu(dy) \right)^{q} \nu(dx)$$

$$\leq \int_{H} \left(\left(\int_{H} h^{p'} H^{p'} \nu(dy) \right)^{1/p'} \left(\int_{H} |\phi|^{p} \nu(dy) \right)^{1/p} \right)^{q} \nu(dx)$$

$$= ||\phi||_{p}^{q} \int_{H} \left(\int_{H} h^{p'} H^{p'} \nu(dy) \right)^{q/p'} \nu(dx).$$
(4.24)

It remains to show that

$$K = \int_{H} \left(\int_{H} h^{p'} H^{p'} \nu(dy) \right)^{q/p'} \nu(dx) < \infty. \tag{4.25}$$

Indeed, using successively the Hölder equality we obtain for any r > 1

$$K \leqslant \int_{H} \left(\int_{H} h^{p'r'} \nu(dy) \right)^{q/p'r'} \left(\int_{H} H^{p'r} \nu(dy) \right)^{q/p'r} \nu(dx)$$

$$\leqslant \left(\int_{H} \left(\int_{H} h^{p'r'} \nu(dy) \right)^{q/p'} \nu(dx) \right)^{1/r'} \left(\int_{H} \left(H^{p'r} \nu(dy) \right)^{q/p'} \nu(dx) \right)^{1/r}. \tag{4.26}$$

It was shown in [5] that

$$\int_{H} \left(H^{a'} \nu(dy) \right)^{b/a'} \nu(dx) < \infty, \tag{4.27}$$

for any $a, b \ge 1$, such that

$$b \leqslant 1 + \frac{a-1}{\|S_0(T)\|^2}. (4.28)$$

Putting

$$a = \frac{p'r}{p'r - 1}$$
 and $b = qr$,

we find that there exists r > 1 such that (4.28) holds. Therefore, for such an r

$$\int_{H} \left(H^{p'r} \nu(dy) \right)^{q/p'} \nu(dx) = \int_{H} \left(H^{a'} \nu(dy) \right)^{b/a'} \nu(dx) < \infty. \tag{4.29}$$

Next, we need to show that

$$\int_{H} \left(\int_{H} h^{p'r'} \nu(dy) \right)^{q/p'} \nu(dx) < \infty. \tag{4.30}$$

To prove (4.30) we note that if $\frac{q}{n'} \ge 1$ then

$$\int_{H} \left(\int_{H} h^{p'r'} \nu(dy) \right)^{q/p'} \nu(dx) \leqslant \int_{H} \int_{H} h^{r'q} \nu(dy) \nu(dx)$$

However, using Lemma 4.8 for $\tilde{q} = r'q$ we have

$$\int_{H} \int_{H} (h(T, x, y))^{\tilde{q}} \nu(dx) \nu(dy) \leqslant \int_{H} \int_{H} h_{\tilde{q}}(T, x, y) \nu(dx) \nu(dy)$$

$$\leqslant \int_{H} \int_{H} k_{\tilde{q}} \exp\{k_{\tilde{q}}(|x| + \int_{0}^{T} |B_{3}(s)y|ds)\} \nu(dx) \nu(dy)$$

$$\leqslant k_{\tilde{q}} \int_{H} \exp\{k_{\tilde{q}}|x|\} \nu(dx) \int_{H} \exp\{k_{\tilde{q}} \int_{0}^{T} |B_{3}(s)y|ds)\} \nu(dy)$$

$$= k_{\tilde{q}} \mathbb{E} e^{k_{\tilde{q}}|\tilde{Z}|} \cdot \mathbb{E} \exp\{k_{\tilde{q}} \int_{0}^{T} |B_{3}(s)\tilde{Z}|ds\}$$

where \tilde{Z} is an arbitrary random variable with probability distribution ν . By (4.16), (4.1) and the Fernique inequality we conclude that (4.30) holds true. The proof of (4.30) for the case when $\frac{q}{n'} < 1$ is even simpler and is omitted.

Remark 4.9. There is a natural question whether the transition density is regular (continuous) "in y", that is, whether the mapping $y \mapsto d(T, x, y)$ is continuous, at least on a certain subspace $\hat{H} \subset H$) of full measure. In the Gaussian case the formulas for the density may be used to conclude that if $\overline{S_T^*Q_T^{-1}} \in \mathcal{L}(\hat{H}, H)$ then $y \to g(T, x, y)$ is continuous on \hat{H} for all T > 0 and $x \in H$ (cf. the Cameron-Martin formula (4.12)). A similar well-known

formula for k(T,y) (see e.g. [5]) yields $\hat{H} \to H$ continuity of the mapping $y \mapsto k(T,y)$ provided

$$C(T) := \overline{Q_{\infty}^{-1/2}(I - S_0(T)S_0^*(T))^{-1}S_0(T)S_0^*(T)Q_{\infty}^{-1/2}} \in \mathcal{L}(\hat{H}, H)$$
(4.31)

where $S_0(T) = Q_{\infty}^{-1/2} S_T Q_{\infty}^{1/2}$. Following the proof of Theorem 4.5 we can easily see that the remaining factor, the function h(T, x, y) is continuous in $y \in \hat{H}$ if the mapping $y \to \hat{Z}_t^{x,y}$ is $\hat{H} \to H$ a.s. continuous (which by Proposition 4.1 (b) happens if $K_t Q_T^{-1/2} \in \mathcal{L}(\hat{H}, H)$) and

$$B_3 \in \mathcal{L}(\hat{H}, L^1(0, T; H)).$$
 (4.32)

We are able to verify these additional conditions in some important cases (supposing that the standing assumptions of this Section (4.3), (4.4) and (4.9) are satisfied).

- (a) All three conditions are satisfied if $\dim H < \infty$.
- (b) In the commutative case the first two conditions are satisfied with $\hat{H} = H$. However, condition (4.32) is not satisfied with $\hat{H} = H$ even in simple infinite dimensional situations and a smaller space \hat{H} must be considered (cf. Example 4.10 below for details).

Example 4.10. Consider the semilinear stochastic heat equation

$$\frac{\partial u}{\partial t}(t,\xi) = \frac{\partial^2 u}{\partial \xi^2}(t,\xi) + f(u(t,\xi)) + \eta(t,\xi), \quad (t,\xi) \in \mathbb{R}_+ \times (0,1), \tag{4.33}$$

with an initial condition and Dirichlet boundary conditions

$$u(0,\xi) = x(\xi), \quad u(t,0) = u(t,1) = 0, \quad t \geqslant 0, \, \xi \in (0,1)$$
 (4.34)

where $f: \mathbb{R} \to \mathbb{R}$ is bounded and continuous and η denoted formally a space-dependent white noise. As well known (see e.g. [7] for fundamentals on the theory of stochastic evolution equations) the system (4.33) - (4.34) may be understood as an equation of the form (4.7) in the space $H = L^2(0,1)$ where $A = \frac{\partial^2}{\partial \xi^2}$, $\operatorname{dom}(A) = H_0^1(0,1) \cap H^2(0,1)$, $F: H \to H$, $F(y)(\xi) := f(y(\xi))$, $y \in H$, $\xi \in (0,1)$, and \sqrt{Q} is a bounded operator on $H = L^2(0,1)$. We assume that the operator Q is boundedly invertible on H, (i.e., the noise is nondegenerate). Then Hypothesis 4.3 is obviously satisfied and Hypothesis 4.4 (ii) is satisfied with $\beta = \frac{1}{2}$ and arbitrary $\alpha \in (0, \frac{1}{2})$ (cf.[11], Example 9.2 and references therein). Thus the conclusions of Theorems 4.5, 4.7 and 4.6 hold true in the present example.

As far as continuity of the transition density "in the variable y" is concerned (cf. Remark 4.9), the problem is more difficult and we only can verify our conditions in the diagonal (commutative) case. Denote by (e_n) and (α_n) the orthonormal basis in H consisting of eigenvectors of the operator -A and its corresponding eigenvalues (so we have $\alpha_n > 0$, $\alpha_n \sim n^2$), and assume that Q commutes with A, that is,

$$Qe_n = \lambda_n e_n, \quad 0 < \inf \lambda_n \leqslant \sup \lambda_n < \infty.$$

Then it is easy to compute eigenvalue expansions of all operators that are needed in Remark 4.9. We have

$$K_t Q_T^{-1} e_n = \frac{1 - e^{-2\alpha_n t}}{1 - e^{-2\alpha_n T}} e^{-\alpha_n (T - t)} e_n, \tag{4.35}$$

$$Q_T^{-1} S_T^* e_n = 2e^{-\alpha_n T} \frac{\alpha_n}{\lambda_n} (1 - e^{-\alpha_n T})^{-1} e_n, \tag{4.36}$$

$$C(T)e_n = 2e^{-2\alpha_n T} \frac{\alpha_n}{\lambda_n} (1 - e^{-2\alpha_n T})^{-1} e_n,$$
(4.37)

$$B_3(s)e_n = 2e^{-\alpha_n(T-s)} \frac{\alpha_n}{\sqrt{\lambda_n}} (1 - e^{-\alpha_n T})^{-1} e_n.$$
 (4.38)

Obviously, all operators given in (4.35) and (4.37) are in $\mathcal{L}(H)$, but it is easy to see that $||B_3(s)|| \sim \frac{1}{T-s}$, so B_3 is not an element of $\mathcal{L}(H, L^1(0,T;H))$ and we do not obtain the continuity in y in the norm of H. However, taking $\hat{H} = \text{dom}((-A)^{\delta})$ endowed with the graph norm for any $\delta > 0$ (which coincides with a suitable Sobolev-Slobodetskii space) we may easily check that the condition (4.32) is satisfied and we may conclude that the mapping $y \mapsto d(T, x, y)$ is $\hat{H} \to H$ continuous.

In the present case it is also easy to write equation (3.15) for the OU Bridge that splits into a sequence of independent one-dimensional equations for particular coordinates $\hat{z}_n^{x,y}(t) := \langle \hat{Z}_t^{x,y}, e_n \rangle$. We obtain

$$d\hat{z}_n^{x,y}(t) = \left[-\alpha_n \hat{z}_n^{x,y}(t) - 2\alpha_n e^{-\alpha_n(T-t)} (1 - e^{-2\alpha_n(T-t)})^{-1} (e^{-\alpha_n(T-t)} \hat{z}_n^{x,y}(t) - y_n) \right] dt + \sqrt{\lambda_n} d\zeta_n(t)$$

for $t \in (0,T)$ with the initial condition

$$\hat{z}_n^{x,y}(0) = x_n,$$

where $x_n = \langle x, e_n \rangle$, $y_n = \langle y, e_n \rangle$ and $\zeta_n(t) = \langle \zeta_t, e_n \rangle$. Here we do not have to assume that the eigenvalues α_n are all negative, only $\alpha_n \neq 0$. If $\alpha_n = 0$ for some n the corresponding equation takes the form

$$d\hat{z}_n^{x,y}(t) = \frac{y_n - \hat{z}_n^{x,y}(t)}{T - t}dt + \sqrt{\lambda_n}d\zeta_n(t), \quad t \in (0, T),$$

which is a well-known equation for a one-dimensional Brownian Bridge.

References

- [1] Ball J. M.: Strongly continuous semigroups, weak solutions, and the variation of constants formula *Proc. Amer. Math. Soc.* 63 (1977), 370-373
- [2] Chojnowska-Michalik A.: Stochastic differential equations in Hilbert spaces, in: Probability theory (Papers, VIIth Semester, Stefan Banach Internat. Math. Center, Warsaw, 1976), pp. 53-74, Banach Center Publ., 5, PWN, Warsaw, 1979
- [3] Chojnowska-Michalik A.: Transition semigroups for stochastic semilinear equations on Hilbert spaces Dissertationes Math. 396 (2001)
- [4] Chojnowska-Michalik A. and Goldys B.: Nonsymmetric Ornstein-Uhlenbeck semigroup as second quantized operator. J. Math. Kyoto Univ. 36 (1996), 481-498

- [5] Chojnowska-Michalik A. and Goldys B.: On regularity properties of nonsymmetric Ornstein-Uhlenbeck semigroup in L^p spaces, *Stochastics and Stochastics Rep.* 59 (1996), 183-209
- [6] Chojnowska-Michalik A. and Goldys B.: Symmetric Ornstein-Uhlenbeck Semigroups and their Generators, *Probab. Theory and Related Fields* 124 (2002), 459-486
- [7] Da Prato G. and Zabczyk J.: Stochastic Equations in Infinite Dimensions, Cambridge University Press 1992
- [8] Fuhrman M.: Regularity properties of transition probabilities in infinite dimensions *Stochastics and Stochastics Rep.* 69 (2000), 31-65
- [9] Goldys B.: On analyticity of Ornstein-Uhlenbeck semigroups, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. 10 (1999), 131-140
- [10] Goldys B. and van Neerven J.M.A.M.: Transition semigroups of Banach space valued Ornstein-Uhlenbeck processes, *Acta Appl. Math.* 76 (2003), 283-330
- [11] Goldys B. and Maslowski B.: Lower estimates of transition densities and bounds on exponential ergodicity for stochastic PDE's, Ann. Probab. 34 (2006), 1451-1496
- [12] Lyons T. J and Zheng W.A.: On Conditional Diffusion Processes, *Proc. Royal Soc. Edinburgh* 115A (1990), 243-255
- [13] Ma Zhi Ming and Röckner M.: Introduction to the theory of (nonsymmetric) Dirichlet forms, Springer-Verlag, 1992
- [14] Mandelbaum A.: Linear estimators and measurable linear transformations on a Hilbert space, Z. Wahrsch. Verw. Gebiete 65 (1984), 385-397
- [15] Maslowski B. and Simão I.: Asymptotic properties of stochastic semilinear equations by the method of lower measures, *Colloquium Math.* 72 (1997), 147-171
- [16] Maslowski B. and Simão I.: Long time behaviour of non-autonomous SPDE's, Stochastic Processes and Applications 95 (2001), 285-309
- [17] Simão I.: Pinned Ornstein-Uhlenbeck processes on an infinite-dimensional space, Stochastic Analysis and Applications (Powys, 1995), World Sci. Publishing, River Edge, NJ, 1996.
- [18] van Neerven J.M.A.M.: Nonsymmetric Ornstein-Uhlenbeck Semigroups in Banach Spaces, J. Funct. Anal. 155 (1998), 495-535
- [19] Ondreját M.: Brownian representations of cylindrical martingales, martingale problem and strong Markov property of weak solutions of SPDEs in Banach spaces, *Czechoslovak Math. J.* 55 (2005), 1003-1039
- [20] Yor M.: Some Aspects of Brownian Motion, Birkhäuser 1992

School of Mathematics, The University of New South Wales, Sydney 2052, Australia $E\text{-}mail\ address$: B.Goldys@unsw.edu.au

Institute of Mathematics, Academy of Sciences of Czech Republic, Žitná 25, 11567 Praha 1, Czech Republic

E-mail address: maslow@math.cas.cz