## 2000]54H11, 22A05

## A CLASS OF GROUPS IN WHICH ALL UNCONDITIONALLY CLOSED SETS ARE ALGEBRAIC

OL'GA V. SIPACHEVA

ABSTRACT. It is proved that, in any subgroup of a direct product of countable groups, the property of being an unconditionally closed set coincides with that of being an algebraic set.

Markov [2] called a subset A of a group G unconditionally closed in G if it is closed in any Hausdorff group topology on G. Clearly, all solution sets of equations in G, as well as their finite unions and arbitrary intersections, are unconditionally closed.

**Definition** (Markov [2]). A subset A of a group G with identity element 1 is said to be *elementary algebraic* in G if there exists a word w = w(x) in the alphabet  $G \cup \{x^{\pm 1}\}$  (x is a variable) such that  $A = \{x \in G : w(x) = 1\}$ . Finite unions of elementary algebraic sets are *additively algebraic* sets. An arbitrary intersection of additively algebraic sets is called *algebraic*. Thus, the algebraic sets in G are the solution sets of arbitrary conjunctions of finite disjunctions of equations.

In his 1945 paper [2], Markov showed that any algebraic set is unconditionally closed and posed the problem of whether the converse is true. In [3] (see also [1]), he solved this problem for countable groups by proving that any unconditionally closed set in a countable group is algebraic. In this paper, we show that the answer is also positive for subgroups of direct products of countable groups (in particular, for all Abelian groups); the proof is based on Markov's ideas. In the general case, the answer is negative (at least, under the continuum hypothesis) [4].

**Definition** (Markov [3]). Let *m* be a positive integer. By a multiplicative function of *m* arguments we mean an arbitrary word on the alphabet  $\{(1, \pm 1), \ldots, (m, \pm 1)\}$ . The *length* of the function is equal to the length of this word. Suppose that *G* is a group and  $x_1, \ldots, x_m \in G$ . The value of a multiplicative function  $\Phi = (j_1, \varepsilon_1) \ldots (j_n, \varepsilon_n)$  (of *m* arguments) at  $x_1, \ldots, x_n$  in *G* is defined as

$$\Phi_G(x_1,\ldots,x_n) = \prod_{i=1}^n x_{j_i}^{\varepsilon_i}.$$

**Definition.** For a subset A of a group G,  $\widetilde{A}^G$  (or simply  $\widetilde{A}$ ) denotes the *algebraic* closure of A in G, i.e., the least algebraic set in G containing A.

In what follows, we consider algebraic closures only in H.

**Theorem 1.** If H is a subgroup of a direct product of countable groups and  $A \subset H$ , then A is unconditionally closed in H if and only if it is algebraic in H.

*Proof.* Let G be the direct product of countable groups  $G_{\alpha}$ , where  $\alpha \in I$ , and let H be a subgroup of G. By 1 and  $1_{\alpha}$  we denote the identity elements of G and  $G_{\alpha}$ , respectively.

<sup>1991</sup> Mathematics Subject Classification. [.

This work was financially supported by the Russian Foundation for Basic Research (project no. 06-01-00761).

Suppose that  $A \subset H$ ,  $1 \in \tilde{A}$ , and  $1 \notin A$  (in particular, A is not algebraic in H). Let us show that A is not unconditionally closed in H. We set  $a_0 = 1$ and take an arbitrary element  $a_1 \in H \setminus \{1\}$ . This element has only finitely many nonzero coordinates; we denote their indices by  $\beta_1, \ldots, \beta_{k_1}$ . Note that the set of all multiplicative functions of length < 6 with 2 arguments is finite. Let  $\mathfrak{M}_1$  be the (finite) subset of this set consisting of all functions  $\Phi$  for which

$$\Phi(a_1,1) \neq 1$$

We set

$$A_{\Phi} = \{ x \in H : \Phi(a_1, x) = 1 \} \quad \text{for} \quad \Phi \in \mathfrak{M}_1$$

and

$$B_1 = \bigcup_{\Phi \in \mathfrak{M}_1} A_{\Phi}.$$

The set  $B_1$  is additively algebraic. We have  $1 \notin B_1$ . On the other hand,  $1 \in A$ . Since  $\widetilde{A}$  is the algebraic closure of A, we have  $A \setminus B_1 \neq \emptyset$ . Take  $x_1 \in A \setminus B_1$ . Note that  $x_1 \neq a_0, a_1$ . Indeed,  $x_1 \neq a_0 = 1$  because  $x_1 \in A \ni 1$ , and  $x \neq a_1$  because  $\Phi = (1, 1)(1, -1)$  is a multiplicative function of length 2 with two arguments for which  $\Phi(a_1, a_1) = a_1 a_1^{-1} = 1$  but  $\Phi(a_1, 1) = a_1 \neq 1$ .

Let  $\{\alpha_1, \ldots, \alpha_{n_1}\}$  be a finite subset of the index set I such that  $\{\beta_1, \ldots, \beta_{k_1}\} \subset \{\alpha_1, \ldots, \alpha_{n_1}\}$  and the indices of all nonidentity coordinates of the element  $x_1 \in G = \prod_{\alpha \in I} G_\alpha$  are contained in  $\{\alpha_1, \ldots, \alpha_{n_1}\}$  (here and in what follows,  $\prod$  denotes direct product). Let us number all elements a of the at most countable set  $H \cap \prod_{\alpha \in I} G'_\alpha \setminus \{a_0, a_1\}$ , where  $G'_\alpha = G_\alpha$  for  $\alpha \in \{\alpha_1, \ldots, \alpha_{n_1}\}$  and  $G'_\alpha = \{1_\alpha\}$  for  $\alpha \notin \{\alpha_1, \ldots, \alpha_{n_1}\}$ , by positive integers  $\geq 2$  so that infinitely many numbers remain unoccupied but the number 2 is occupied, i.e.,  $a_2$  is defined (e.g., we can put  $a_2 = x_1$  and use only even numbers for numbering; in this case, we have  $H \cap \prod_{\alpha \in I} G'_\alpha = \{a_0, a_1, a_2, a_4, a_6, \ldots\}$ . The set  $H \cap \prod_{\alpha \in I} G'_\alpha$  may be finite, but it necessarily contains  $a_0, a_1$ , and  $a_2$ ).

Take an element  $a_2$ . Arguing as above, we choose  $x_2 \in A$  so that if  $\Phi$  is a multiplicative function of length < 9 with 4 arguments and  $\Phi(a_1, a_2, x_1, x_2) = 1$ , then  $\Phi(a_1, a_2, x_1, 1) = 1$ , and show that  $x_2 \neq a_0, a_1, a_2$ . Let  $n_2 > n_1$ , and let  $\{\alpha_1, \ldots, \alpha_{n_2}\} \subset I$  be a set of indices which contains the indices of all nonidentity coordinates of  $x_2$ . Let us number all unnumbered elements a of the at most countable set  $H \cap \prod_{\alpha \in I} G''_{\alpha}$ , where  $G''_{\alpha} = G_{\alpha}$  for  $\alpha \in \{\alpha_1, \ldots, \alpha_{n_2}\}$  and  $G''_{\alpha} = \{1_{\alpha}\}$  for  $\alpha \notin \{\alpha_1, \ldots, \alpha_{n_2}\}$ , by positive integers that have not been used at the preceding step so that infinitely many numbers remain unoccupied but the number 3 is occupied (e.g., we can put  $a_3 = x_2$  and use only odd numbers divisible by 3 for numbering).

At the *j*th step, the situation is as follows. Positive integers  $n_1 < \cdots < n_{j-1}$  and indices  $\alpha_1, \ldots, \alpha_{n_{j-1}} \in I$  are defined; elements  $x_1, \ldots, x_{j-1} \in A$  and  $a_1, \ldots, a_j \in$  $H \cap \prod_{\alpha \in I} G_{\alpha}^{\prime\prime\prime}$ , where  $G_{\alpha}^{\prime\prime\prime} = G_{\alpha}$  for  $\alpha \in \{\alpha_1, \ldots, \alpha_{n_{j-1}}\}$  and  $G_{\alpha}^{\prime\prime} = \{1_{\alpha}\}$  for  $\alpha \notin \{\alpha_1, \ldots, \alpha_{n_{j-1}}\}$ , are defined; all elements *a* of the at most countable set  $H \cap$  $\prod_{\alpha \in I} G_{\alpha}^{\prime\prime\prime}$  are numbered by positive integers, and infinitely many positive integers remain unoccupied (however, the numbers  $1, \ldots, j$  are occupied by the elements  $a_1, \ldots, a_j$ ). Moreover, for any multiplicative function  $\Phi$  of length < 3j with 2(j - 1) arguments satisfying the condition  $\Phi(a_1, \ldots, a_{j-1}, x_1, \ldots, x_{j-1}) = 1$ , we have  $\Phi(a_1, \ldots, a_{j-1}, x_1, \ldots, x_{j-2}, 1) = 1$ . Let us define  $x_j$ .

The set of all multiplicative functions of length < 3(j+1) with 2j arguments is finite. Let  $\mathfrak{M}_j$  be its subset consisting of all functions  $\Phi$  for which

$$\Phi(a_1,\ldots,a_j,x_1,\ldots,x_{j-1},1) \neq 1.$$

We set

$$A_{\Phi} = \{x : \Phi(a_1, \dots, a_j, x_1, \dots, x_{j-1}, x) = 1\} \quad \text{for} \quad \Phi \in \mathfrak{M}_j$$

and

$$B_j = \bigcup_{\Phi \in \mathfrak{M}_j} A_\Phi.$$

The set  $B_j$  is additively algebraic, and  $1 \notin B_j$ . On the other hand,  $1 \in \widetilde{A}$ . Since  $\widetilde{A}$  is the algebraic closure of A, we have  $A \setminus B_j \neq \emptyset$ . Choose any  $x_j \in A \setminus B_j$ . Clearly,  $x_j \neq a_0 = 1, a_1, \ldots, a_j$ , because if  $x_j = a_i$ , then  $\Phi = (i, 1)(2j, -1)$  is a multiplicative function of length < 3(j+1) with 2j arguments for which  $\Phi(a_1, \ldots, a_j, x_1, \ldots, x_j) = a_i x_j^{-1} = 1$  but  $\Phi(a_1, \ldots, a_j, x_1, \ldots, x_{j-1}, 1) \neq 1$ . Take sufficiently large number  $n_j > n_{j-1}$  and set  $\{\alpha_{n_{j-1}+1}, \ldots, \alpha_{n_j}\} \subset I$  for which all coordinates of the element  $x_j$  with indices not belonging to  $\{\alpha_1, \ldots, \alpha_{n_j}\}$  are the identity elements of the corresponding factors. Let us number all elements a of the at most countable set  $H \cap \prod_{\alpha \in I} G_{\alpha}^{\prime\prime\prime\prime}$  (where  $G_{\alpha}^{\prime\prime\prime\prime} = G_{\alpha}$  for  $\alpha \in \{\alpha_1, \ldots, \alpha_{n_j}\}$  and  $G_{\alpha}^{\prime\prime\prime\prime} = \{1_{\alpha}\}$  for  $\alpha \notin \{\alpha_1, \ldots, \alpha_{n_j}\}$ ) which have not been numbered at the preceding steps by positive integers not occupied at the preceding steps so that infinitely many numbers remain unoccupied. If the element  $a_{j+1} = x_j$ .

As a result, we obtain a countable set  $I^* = \{\alpha_i : i \in \mathbb{N}\} \subset I\}$ ; all elements of the set  $G^* = H \cap \prod_{\alpha \in I} G^*_{\alpha}$ , where  $G^*_{\alpha} = G_{\alpha}$  for  $\alpha \in I^*$  and  $G^*_{\alpha} = \{1_{\alpha}\}$  for  $\alpha \notin I^*$ , are numbered by nonnegative integers:  $G^* = \{a_0, a_1, a_2, \ldots\}$ , where and  $a_0 = 1$ . We also have a sequence  $\{x_1, x_2, \ldots\} \subset A$  such that if  $\Phi$  is a multiplicative function of length  $\langle 3(j+1) \rangle$  with 2j arguments and  $\Phi(a_1, \ldots, a_j, x_1, \ldots, x_j) = 1$ , then  $\Phi(a_1, \ldots, a_j, x_1, \ldots, x_{j-1}, 1) = 1$ . We set  $A^* = A \cap G^*$ .

Note that  $G^*$  is a countable normal subgroup of H and  $A^* \subset G^*$ . Thus, the situation is quite similar to that considered by Markov in [3]. The role of the group G from [3] is played by  $G^*$ , and the role of A is played by  $A^*$ . In [3, Sections 3–5], Markov defined sequences  $\{b_i\}_{i=1}^{\infty}$ ,  $\{a_i\}_{i=1}^{\infty}$ , and  $\{x_i\}_{i=1}^{\infty}$  with certain properties (the properties of the first sequence are labeled by 2.31 and 2.32 in his paper, the properties of the second are labeled by 2.41–2.44, and the properties of the third are labeled by 2.51 and 2.52). We set  $b_i = 1$  for all i; the sequences  $\{a_i\}_{i=1}^{\infty}$  and  $\{x_i\}_{i=1}^{\infty}$  are already constructed. These sequences have all of the properties mentioned above except 2.32. However, 2.32 is not used in Sections 6–12 of [3]; therefore, the argument from these sections applies to the case under consideration without any changes (except that G should be replaced by  $G^*$  and A by  $A^*$ ); in particular, the functions  $f_j$  on the sets

$$A_j = \{1, a_j\} \cup \{a_k^{-1} x_i a_k : i \ge j, k \le i\}$$

given by

$$f_j(1) = 0,$$
  $f_j(a_j) = 1,$   $f_j(a_k^{-1}x_ia_k) = \frac{1}{i}$ 

are well defined and can be extended to seminorms  $N_j$  on the group  $G^*$ . Following Markov [3, Section 11], for any positive integer n and sets of integers  $\{p_j\}_{j=1}^n$  and  $\{q_j\}_{j=1}^n$ , where  $p_j \ge 0$  and  $q_j > 0$  for  $j = 1, \ldots, n$ , we define a seminorm  $N_{q_1,\ldots,q_n}^{p_1,\ldots,p_n}$  on  $G^*$  by

$$N_{q_1,\dots,q_n}^{p_1,\dots,p_n}(x) = \sum_{j=1}^n N_{q_j}(a_{p_j}^{-1}xa_{p_j});$$

we have  $N_q^0 = N_q$  for q = 1, 2, ..., because  $a_0 = 1$ . Markov showed that these seminorms determine a group topology  $\mathcal{T}^*$  on  $G^*$ ; its neighborhood base at the identity consists of all sets of the form

$$U_N = \{ x \in G^* : N(x) < 1 \},\$$

where  $N = N_{q_1,...,q_n}^{p_1,...,p_n}$  for some  $p_1,...,p_n \ge 0$  and  $q_1,...,q_n > 0$ .

Let us prove that 1 belongs to the closure of  $A^*$  in this topology. It is sufficient to show that  $U_N$  intersects  $A^*$  for any N.

Let  $N = N_{q_1,\ldots,q_n}^{p_1,\ldots,p_n}$ . Take a positive integer s such that  $s > n, s > p_j$ , and  $s > q_j$  for  $j = 1, \ldots, n$ . By definition, we have

$$N(x_s) = \sum_{j=1}^{n} N_{q_j}(a_{p_j}^{-1}x_s a_{p_j}) = \frac{n}{s} < 1.$$

Therefore,  $x_s \in U_N$ . On the other hand,  $x_s \in A^*$  by construction.

Thus, the set  $A^*$  is not closed in the group  $G^*$  with the topology  $\mathcal{T}^*$ . Since  $G^*$  is a normal subgroup of H, the neighborhoods of the identity in the topology  $\mathcal{T}^*$  form a neighborhood base of the identity for some group topology  $\mathcal{T}$  on H. Since any neighborhood of the identity in  $\mathcal{T}^*$  intersects  $A^*$  and  $A^* = A \cap G^* \subset A$ , it follows that any neighborhood of the identity in  $\mathcal{T}$  intersects A. Thus, A is not unconditionally closed in H.

Now, let *B* be an arbitrary nonalgebraic set in *H*; then  $\widetilde{B} \neq B$ . Take  $b \in \widetilde{B} \setminus B$ . We have  $1 \notin b^{-1}B$ . On the other hand,  $1 \in b^{-1}\widetilde{B}$ . According to Lemma 12 from [3],  $b^{-1}\widetilde{B} = \widetilde{b^{-1}B}$ ; hence  $1 \in \widetilde{b^{-1}B} \setminus b^{-1}B$ . As was shown above, this implies the nonclosedness of the set  $b^{-1}B$  (and, therefore, of *B* itself) in some group topology on *H*. Thus, if the set *B* is not algebraic in *H*, then it cannot be unconditionally closed in *H*. On the other hand, any algebraic set in the group *H* is unconditionally closed in *H* [3, Theorem 1].

**Corollary.** If G is an Abelian group and  $A \subset G$ , then A is unconditionally closed in G if and only if it is algebraic in G.

This assertion follows immediately from the fact that any Abelian group can be embedded in a direct product of countable groups as a subgroup (see, e.g., [5]).

## References

- A. A. Markov. "On unconditionally closed sets," Dokl. Akad. Nauk SSSR 44 (5), 196–197, 1944; English translation: A. Markoff, "On unconditionally closed sets," C. R. (Doklady) Acad. Sci. URSS (N.S.) 44, 180–181 (1944).
- [2] A. A. Markov, "On free topological groups," Izv. Akad. Nauk SSSR, Ser. Mat. 9 (1), 3– 64, 1945; English translation: "Three papers on topological groups: I. On the existence of periodic connected topological groups. II. On free topological groups. III. On unconditionally closed sets," Amer. Math. Soc. Transl. 30 (1950).
- [3] A. A. Markov, "On unconditionally closed sets," Mat. Sb. 18 (1), 3–26, 1946; English translation: "Three papers on topological groups: I. On the existence of periodic connected topological groups. II. On free topological groups. III. On unconditionally closed sets," Amer. Math. Soc. Transl. 30 (1950).
- [4] O. V. Sipacheva, "Consistent solution of Markov's problem about algebraic sets," ArXiv:math.GR/0605558.
- [5] I. Kaplansky, Infinite Abelian Groups (Univ. of Michigan Press, Ann Arbor, 1954).

DEPARTMENT OF GENERAL TOPOLOGY AND GEOMETRY, MECHANICS AND MATHEMATICS FAC-ULTY, MOSCOW STATE UNIVERSITY, LENINSKIE GORY, MOSCOW, 119992 RUSSIA

*E-mail address*: o-sipa@yandex.ru

4