# MORDELL-WEIL GROUPS OF A HYPERKÄHLER MANIFOLD -A QUESTION OF F. CAMPANA

### KEIJI OGUISO

ABSTRACT. Mordell-Weil groups of different abelian fibrations of a hyperkähler manifold may have non-trivial relation even among elements of infinite order, but have essentially no relation, as its birational transformation. Precise definition of the terms "essentially no relation" will be given in Introduction.

#### 1. INTRODUCTION - BACKGROUND AND THE STATEMENT OF MAIN RESULT

Unless stated otherwise, varieties are assumed to be normal, projective and defined over  $\mathbf{C}$ . Main results are Theorems (1.5) and (1.6).

Let M be a hyperkähler manifold, i.e. a simply-connected projective manifold admitting an everywhere non-degenerate global holomorphic 2-form  $\sigma_M$  s.t.  $H^0(M, \Omega_M^2) = \mathbb{C}\sigma_M$  ([GHJ, Part III]). We denote by Bir M the group of birational transformations of M. A morphism  $\varphi : M \longrightarrow B$  onto a variety B is called an *abelian fibration* if its generic fiber  $M_\eta(\ni O)$  is a positive dimensional abelian variety defined over  $\mathbb{C}(B)$ . The group  $M_\eta(\mathbb{C}(B))$ , called the Mordell-Weil group of  $\varphi$  and denoted from now by MW( $\varphi$ ), can be naturally regarded as an abelian subgroup of Bir M. In [Og3](see also [Og4]), we have shown the following:

**Theorem 1.1.** (1) Let G < Bir M. Then either  $G > \mathbb{Z} * \mathbb{Z}$  or G is an almost abelian group of rank at most  $\text{Max}(1, \rho(M) - 2)$ , the latter of which is finitely generated.

(2) Assume that M admits 2 different abelian fibrations  $\varphi_i : M \longrightarrow B_i$  s.t. rank  $MW(\varphi_i) > 0$  (i = 1, 2). Let  $f_i \in MW(\varphi_i)$  s.t. ord  $f_i = \infty$ . Then the subgroup  $\langle f_1, f_2 \rangle$  of Bir M contains  $\mathbf{Z} * \mathbf{Z}$  as its subgroup.

Unfortunately, the proof in [Og3], which is based on Tits' alternative [Ti] together with properties of Salem polynomial (cf. [Mc]), tells us no explicit way to find  $\mathbf{Z} * \mathbf{Z}$ in  $\langle f_1, f_2 \rangle$ . Then F. Campana asked the following:

Question 1.2. Under the same condition as in Theorem (1.1)(2),

$$\langle f_1, f_2 \rangle = \langle f_1 \rangle * \langle f_2 \rangle \simeq \mathbf{Z} * \mathbf{Z}$$

for suitably chosen  $f_i \in MW(\varphi_i)$  (i = 1, 2)?

In the view of an observation of Cantat [Ca], one might ask an even stronger:

Question 1.3. Under the same condition as in Theorem (1.1)(2),

$$\langle f_1, f_2 \rangle = \langle f_1 \rangle * \langle f_2 \rangle \simeq \mathbf{Z} * \mathbf{Z}$$

for any  $f_i \in MW(\varphi_i)$  s.t. ord  $(f_i) = \infty$  (i = 1, 2) at least when dim M = 2?

<sup>2000</sup> Mathematics Subject Classification. 14J50, 14J28.

The aim of this note is to give an affirmative answer to the first question in a slightly more general form and a negative answer to the second question. We note that there are infinitely many K3 surfaces and hyperkähler manifolds admitting 2 or more different abelian fibrations of positive Mordell-Weil rank (see eg. [Og3]).

**Definition 1.4.** Let G be a group and let  $G_i$   $(i = 1, 2, \dots, s)$  be subgroups of G. We say that the groups  $G_i$   $(i = 1, 2, \dots, s)$  have essentially no relation if there are finite index subgroups  $H_i < G_i$  s.t.

$$\langle H_1, H_2, \cdots, H_s \rangle = H_1 * H_2 * \cdots * H_s$$
.

Our main result is the following:

**Theorem 1.5.** (1) Let  $\varphi_i : M \longrightarrow B_i$   $(i = 1, 2, \dots, s)$  be mutually different abelian fibrations on a hyperkähler manifold M s.t. rank  $MW(\varphi_i) =: r_i > 0$ . Then, in Bir M, the Mordell-Weil groups  $MW(\varphi_i)$   $(i = 1, 2, \dots, s)$  have essentially no relation, i.e. there are finite index subgroups  $H_i < MW(\varphi_i)$  s.t.

$$\langle H_1, H_2, \cdots, H_s \rangle = H_1 * H_2 * \cdots * H_s \simeq \mathbf{Z}^{r_1} * \mathbf{Z}^{r_2} * \cdots * \mathbf{Z}^{r_s}$$
.

In particular, for given  $g_i \in MW(\varphi_i)$  with  $\operatorname{ord} g_i = \infty$ , there are positive integers  $m_i$   $(i = 1, 2, \dots, s)$  s.t.

$$\langle g_1^{m_1}, g_2^{m_2}, \cdots, g_s^{m_s} \rangle = \langle g_1^{m_1} \rangle * \langle g_2^{m_2} \rangle * \cdots * \langle g_s^{m_s} \rangle \simeq \mathbf{Z} * \mathbf{Z} * \cdots * \mathbf{Z}$$
(s-factors).

(2) There are a K3 surface S admitting 2 different Jacobian fibrations  $\varphi_i : S \longrightarrow \mathbf{P}^1$  and elements  $f_i \in \mathrm{MW}(\varphi_i)$  with  $\operatorname{ord} f_i = \infty$  (i = 1, 2) s.t. in  $\operatorname{Aut} S$ ,

$$\langle f_1, f_2 \rangle \simeq \mathbf{Z}/2 * \mathbf{Z}/3$$
, whence,  $\langle f_1, f_2 \rangle \neq \langle f_1 \rangle * \langle f_2 \rangle (\simeq \mathbf{Z} * \mathbf{Z})$ ,

but, for each integer  $n \geq 2$ ,

$$\langle f_1^n, f_2^n \rangle = \langle f_1^n \rangle * \langle f_2^n \rangle \simeq \mathbf{Z} * \mathbf{Z}$$
.

Recall that a K3 surface is a 2-dimensional hyperkähler manifold, and an abelian fibration on a surface is usually called a Jacobian fibration. We also note that Aut S = Bir S for a K3 surface (or more generally for a minimal surface). We shall prove Theorem (1.5)(1) in Section 2 and Theorem (1.5)(2) in Section 3. Our proof here is much inspired by an elementary hyperbolic geometry explained in the book [Th] of Thurston and an account [Ha] of de la Harpe, whose applicability is suggested by S. Cantat in connection with [Og3].

As a more concrete example, we shall also show, in Section 4, the following:

**Theorem 1.6.** There is a K3 surface S s.t. for each positive integer s,

Aut 
$$S > \mathbf{Z}^{18} * \mathbf{Z}^{18} * \dots * \mathbf{Z}^{18}$$
 (s-factors).

Note that if  $\mathbf{Z}^r < \operatorname{Aut} S$  for some K3 surface S, then  $r \leq 18$  by Theorem (1.1)(2).

Acknowledgement. I would like to express my deep thank to Professors F. Campana, S. Cantat, J.M. Hwang, M. Paun, and J. Winkelman for their valuable discussions, mostly in Johns Hopkins University (Recent Developments in Higher Dimensional Algebraic Geometry, March 2006), Luminy (Dynamics and Complex Geometry, June 2006) and Oberwolfach (Komplexe Analysis, August 2006).

 $\mathbf{2}$ 

### 2. PROOF OF THEOREM (1.5)(1)

1. The next Theorem and its proof are taken from Tits' article [Ti]. This simple theorem is very useful in our proof of Theorem (1.5).

**Theorem 2.1.** Let s be an integer s.t.  $s \ge 2$  and let G be a group which acts faithfully on a non-empty set S. Let  $G_i$   $(i = 1, 2, \dots, s)$  be subgroups of G, let  $S_i$  $(i = 1, 2, \dots, s)$  be subsets of S s.t.  $S \setminus \bigcup_{i=1}^s S_i \ne \emptyset$ , and let p be an element of  $S \setminus \bigcup_{i=1}^s S_i$ . Put  $G_i^0 := G_i \setminus \{id\}$  and  $\tilde{S}_i := S_i \cup \{p\}$ . Suppose that  $G_i^0(\tilde{S}_j) \subset S_i$  for all pairs  $i \ne j$ . Then

$$\langle G_1, G_2, \cdots, G_s \rangle = G_1 * G_2 * \cdots * G_s$$
.

Proof. Consider  $g_1g_2\cdots g_m$ , where  $g_k \in G_{i_k}^0$  and  $i_1 \neq i_2 \neq \cdots \neq i_m$ . Using  $s \geq 2$ , one can choose an integer i s.t.  $i \neq i_m$  and  $1 \leq i \leq s$ . Then  $G_{i_m}^0(\tilde{S}_i) \subset S_{i_m}, G_{i_{m-1}}^0 G_{i_m}^0(\tilde{S}_i) \subset S_{i_{m-1}}$ , and finally  $G_{i_1}^0 G_{i_2}^0 \cdots G_{i_{m-1}}^0 G_{i_m}^0(\tilde{S}_i) \subset S_{i_1}$ . Thus  $g_1g_2\cdots g_m(p) \subset S_{i_1}$ . Hence  $g_1g_2\cdots g_m(p) \neq p$  by  $p \notin S_{i_1}$ , and therefore  $g_1g_2\cdots g_m \neq id$ .

**2.** Let M be a hyperkähler manifold. As it is mentioned in Introduction, M is assumed to be projective. In this paragraph, we recall some basic facts about M and fix some notations.

(2.1) The Néron-Severi group NS(M) is a non-degenerate lattice of singature  $(1, \rho(M) - 1)$  w.r.t. the Beauville form (\*, \*\*). (See eg. [GHJ, Part III]; the Beauville form here is called the Beauville-Bogomolov form there). Here  $\rho(M) := \operatorname{rank} NS(M)$  is the Picard number of M. Note that  $c_1 : \operatorname{Pic} M \longrightarrow NS(M)$  is an isomorphism for M.

Let O(NS(M)) be the group of isometries of the lattice NS(M). Then, there is a natural group homomorphism

$$r_{NS}$$
: Bir  $M \longrightarrow O(NS(M))$ ;  $g \mapsto (g^{-1})^* | NS(M)$ 

(see eg. [GHJ, Part III]), and  $|\text{Ker} r_{NS}| < \infty$  (see eg. [Hu], [Og2]). For the last statement, projectivity of M is essential ([Og2]).

(2.2) In what follows, we denote the scalar extension  $NS(M) \otimes K$  (K is a field) simply by  $NS(M)_K$ . Whenever we discuss topology, we regard  $NS(M)_{\mathbf{R}} \simeq \mathbf{R}^{\rho(M)}$  as the Euclidean space by a suitable Euclidean norm ||\*||.

Let  $\mathcal{C}$  be the *positive cone* of NS(M), that is, the connected component of  $\{x \in NS(M)_{\mathbf{R}} | (x, x) > 0\}$  containing ample classes. We denote by  $\overline{\mathcal{C}}$  the closure of  $\mathcal{C}$  in  $NS(M)_{\mathbf{R}}$ , and by  $\partial \overline{\mathcal{C}}$ , the boundary of  $\mathcal{C}$ . We define

$$S := \{ x \in \partial \mathcal{C} \mid ||x|| = 1 \} .$$

This space S can be identified with the set of the rays  $\{\mathbf{R}_+ x \mid x \in \partial \mathcal{C} \setminus \{0\}\}$ , by the natural bijection  $\mathbf{R}_+ x \leftrightarrow x/||x||$ , and is also homeomorphic to the sphere  $S^{\rho(M)-2}$ .

(2.3) Let  $\varphi: M \longrightarrow B$  be an abelian fibration on M. Then, by a fundamental result due to Matsushita [Ma], B is **Q**-factorial and of Picard number 1. Let v be the (uniquely determined) primitive element in  $\mathbf{R}_+\varphi^*h_B \cap NS(M)$ . Here  $h_B$  is an ample class of Pic B. We call this v the primitive vector corresponding to  $\varphi$ . Note that  $v \in (\partial \overline{C} \cap NS(M)) \setminus \{0\}$ . We denote  $v/||v|| \in S$  by  $\overline{v}$  and call  $\overline{v}$  the unit vector corresponding to  $\varphi$ .

Let  $\hat{G}$  be one of the maximal free abelian subgroups of

$$\operatorname{Im}(r_{NS}|\mathrm{MW}(\varphi):\mathrm{MW}(\varphi)\longrightarrow \mathrm{O}(NS(M)))$$
.

Then  $\tilde{G} \simeq \mathbf{Z}^r$   $(r = \operatorname{rank} \operatorname{MW}(\varphi))$  by the paragraph (2.1). Note that the primitive vector v is stable under  $\tilde{G}$ , i.e.  $\tilde{G}(v) = v$ . This is because  $\rho(B) = 1$ .

Given another abelian fibration  $\varphi' : M \longrightarrow B'$ , we write the primitive (resp. unit) element of  $\varphi'$  by v' (resp. by  $\overline{v'}$ ). Again by  $\rho(B) = \rho(B') = 1$ , we have  $\varphi = \varphi'$  iff v = v', or equivalently, iff  $\overline{v} = \overline{v'}$ .

(2.4) Let  $\varphi: M \longrightarrow B, v, \overline{v}, \tilde{G}$  be the same as in (2.3). Following [Th, Problem 2.5.24], we consider the spaces:

$$\begin{split} \tilde{L}^{\mathbf{Q}} &:= \left\{ x \in NS(M)_{\mathbf{Q}} | (x, v) = 1 \right\} , \ \tilde{L} := \left\{ x \in NS(M)_{\mathbf{R}} | (x, v) = 1 \right\} ; \\ L^{\mathbf{Q}} &:= \tilde{L}^{\mathbf{Q}} / \mathbf{Q} v \ , \ L := \tilde{L} / \mathbf{R} v \ ; \\ P &:= \left\{ x \in \partial \mathcal{C} \mid (x, v) = 1 \right\} \subset \tilde{L} \ . \end{split}$$

Here, since  $v \in NS(M)$  and (v, v) = 0, the mapping  $av : x \mapsto x + av$ ,  $a \in \mathbf{Q}$  (resp.  $a \in \mathbf{R}$ ) defines a faithful action of  $\mathbf{Q}v$  (resp. of  $\mathbf{R}v$ ) on  $\tilde{L}^{\mathbf{Q}}$  (resp. on  $\tilde{L}$ ). We note that these spaces depend on (the choice of) an abelian fibration  $\varphi : M \longrightarrow B$ .

Remark 2.2. Though the rational spaces  $\tilde{L}^{\mathbf{Q}}$  and  $L^{\mathbf{Q}}$  are not considered in [Th, Problem 2.5.14], it is important to consider these spaces for our proof of Theorem (1.5).

By definition, the space  $L^{\mathbf{Q}}$  (resp. L) is a  $(\rho(M) - 2)$ -dimensional rational (resp. real) affine space. Or more explicitly, by taking a point in  $u \in L^{\mathbf{Q}}$  as 0 and by taking a rational basis of the tangent space at u, we can identify  $L^{\mathbf{Q}}$  (resp. L) with the vector space  $\mathbf{Q}^{\rho(M)-2}$  (resp.  $\mathbf{R}^{\rho(M)-2}$ ). Under these identifications, we have also  $L = (L^{\mathbf{Q}})_{\mathbf{R}}$ . We denote by p the natural quotient maps :

$$p: \tilde{L}^{\mathbf{Q}} \longrightarrow L^{\mathbf{Q}} , \ p: \tilde{L} \longrightarrow L .$$

Finally, we define the map  $\pi$  by

$$\pi: P \longrightarrow S \; ; \; x \mapsto x/||x|| \; .$$

Since  $\tilde{G}v = v$  and (v, v) = 0, the group  $\tilde{G}$  acts on  $\tilde{L}^{\mathbf{Q}}$ ,  $\tilde{L}$ ,  $L^{\mathbf{Q}}$ , L, P and S. Here the action of  $\tilde{G}$  on S is defined by  $x \mapsto g(x)/||g(x)||$ . By definiton, the actions of  $\tilde{G}$  on these spaces are equivariant under p and  $\pi$ .

**3.** In this paragraph, again closely following [Th, Problem 2.5.14], we shall prove three propositions, which are crucial in our proof of Theorem (1.5)(1). Let  $\varphi: M \longrightarrow B$  be the same as in the paragraph 2. We use the same notation as in the paragraphs (2.3) and (2.4).

**Proposition 2.3.** The map  $p|P: P \longrightarrow L$  is a homeomorphism and the map

$$\iota := \pi \circ (p|P)^{-1} : L \longrightarrow S$$

is a homeomorphism onto  $S \setminus \{\overline{v}\}$ , i.e. S is a one point compactification of the real affine space L through  $\iota$ . Moreover, the actions of  $\tilde{G}$  on L and S are equivariant under  $\iota$ .

*Proof.* The last assertion follows from the last statement in the paragraph (2.4). Let us show the first two assertions. For a given  $x \in \tilde{L}$ , there is a unique  $\alpha$  s.t.  $(x + \alpha v, x + \alpha v) = 0$ , namely  $\alpha = -(x, x)/2$ . Since  $(x + \alpha v, v) = 1 > 0$ , we have  $x + \alpha v \in \partial C$ , and therefore  $x + \alpha v \in P$ . By the uniqueness of  $\alpha$ , the map  $x + \mathbf{R}v \mapsto x + \alpha v$  is well-defined and gives the inverse  $q: L \longrightarrow P$  of p|P.

By (v, v) = 0, we have  $\pi(P) \subset S \setminus \{\overline{v}\}$ . Let  $x \in \partial \mathcal{C} \setminus \{0\}$ . By the Schwartz inequality, we have (x, v) > 0 unless  $x \in \mathbf{R}v$ . Thus, for  $x \in S \setminus \{\overline{v}\}$ , we have

4

 $x/(x,v) \in P$  and  $\pi(x/(x,v)) = x$ . Thus,  $x \mapsto x/(x,v)$  gives the inverse of  $\pi : P \longrightarrow S \setminus \{\overline{v}\}$ . Hence  $\iota = \pi \circ q$  satisfies the requirement.  $\Box$ 

**Proposition 2.4.** There is a finite index subgroup G of  $\tilde{G}$  s.t. G acts on the real affine space L as parallel transformations. Moreover, the action of G on L is faithful and discrete.

Remark 2.5. For both statements, it seems quite essential that the action  $\hat{G}$  is defined over the *lattice* NS(M), or in other words, over  $\mathbb{Z}$ . Being parallel transformations is false even for a cyclic orthogonal subgroup of  $NS(M)_{\mathbb{Q}}$  fixing some non-zero rational vector v with (v, v) = 0, as there is an infinite-order  $3 \times 3$  rotation matrix of rational entries. An account of [Ha, Page 135] seems to miss this point. See remark (2.7) for discreteness.

*Proof.* Since v is a primitive vector with (v, v) = 0, one can choose an integral basis of  $v_{NS(M)}^{\perp}$  as  $\langle v, w_1, \dots, w_n \rangle$ .

Choose  $u \in NS(M)_{\mathbf{Q}}$  s.t. (v, u) = 1. One can choose such u, because NS(M) is non-degenerate. Then

$$\langle v, w_1, \cdots, w_n, u \rangle$$

forms **Q**-basis of  $NS(M)_{\mathbf{Q}}$ . Note that, in this notation,  $\rho(M) = n + 2$ .

Let  $g \in \tilde{G}$ . Since g(v) = v and  $g(v_{NS(M)}^{\perp}) = v_{NS(M)}^{\perp}$ , the matrix representation M(g) of g w.r.t. the basis above is of the form:

$$M(g) := \begin{pmatrix} 1 & {}^{t}\mathbf{a}(g) & c(g) \\ \mathbf{o} & A(g) & \mathbf{b}(g) \\ 0 & {}^{t}\mathbf{o} & d(g) \end{pmatrix}$$

Here A(g) is the matrix representation of the action of g on the *lattice*  $N := v_{NS(M)}^{\perp}/\mathbb{Z}v$  w.r.t. its *integral* basis  $\langle [w_i] \rangle_{i=1}^n$ . Thus A(g) is an *integral* matrix, while  $\mathbf{b}(g) \in \mathbf{Q}^n$  is, in general, a rational vector.

Since N is of negative definite, its orthogonal transformation A(g) is diagonalizable, and the eigenvalues of A(g) are of absolute value 1. On the other hand, since A(g) is an integral matrix, its eigenvalues are algebraic integers. Thus, by Kronecker's theorem, the eigenvalues of A(g) are roots of 1. Since A(g) is diagonalizable, A(g) is then of finite order. Let us denote by m(g) the order of A(g).

Let  $\langle \tilde{g}_i \rangle_{i=1}^r$  be a generator of the free abelian group  $\tilde{G}$ . Put  $g_i := \tilde{g}_i^{2m(\tilde{g}_i)}$  and define the subgroup G of  $\tilde{G}$  by

$$G := \langle g_1, g_2, \cdots, g_r \rangle$$
.

Since  $\hat{G}$  is a finitely generated abelian group, G is a finite index subgroup of  $\hat{G}$ . Moreover, for each  $g \in G$ , we have  $A(g) = I_n$  (the identity matrix) and d(g) = 1, i.e.

$$M(g) := \begin{pmatrix} 1 & {}^{t}\mathbf{a}(g) & c(g) \\ \mathbf{o} & I_n & \mathbf{b}(g) \\ 0 & {}^{t}\mathbf{o} & 1 \end{pmatrix} .$$

For d(g) = 1, observe that det  $M(\tilde{g}_i) = \pm 1$ , whence det  $M(\tilde{g}_i^{2m(\tilde{g}_i)}) = 1$ , and therefore det M(g) = 1 for  $g \in G$ .

Let  $Q \in L$ . Then Q is uniquely expressed in the form:

$$Q = [u] + \sum_{i=1}^{n} x_i(Q)[w_i] ,$$

and the vector valued function  $\mathbf{x} := (x_i)_{i=1}^n$  gives an affine coordinate of both L and  $L^{\mathbf{Q}}$ . As it is remarked in the paragraph (2.4), under this coordinate, one can identify

$$L^{\mathbf{Q}} = \mathbf{Q}^n, \, L = (\mathbf{Q}^n) \otimes \mathbf{R} = \mathbf{R}^n$$

By the shape of M(g), the action of  $g \in G$  on  $L = \mathbb{R}^n$  is expressed by the parallel transformation by  $\mathbf{b}(g) \in \mathbb{Q}^n$ . This shows the first statement.

For the discreteness and faithfulness of the action of G on  $L = \mathbf{R}^n$ , it suffices to show Lemma (2.6) below. Indeed,  $\langle \rho(g_i) \rangle_{i=1}^r$  then forms a part of real basis of  $L = (\mathbf{Q}^n) \otimes \mathbf{R}$ .

Lemma 2.6. The map

$$\rho: G \simeq \mathbf{Z}^r \longrightarrow \mathbf{Q}^n \; ; \; g \mapsto \mathbf{b}(g)$$

is an injective group homomorphism.

Remark 2.7. Note that the map  $(a, b) \mapsto a + b\sqrt{2}$  defines an injective group homomorphism from  $\mathbb{Z}^2$  to the group  $\mathbb{R}$  of the parallel transformations of  $\mathbb{R}$ , but the image  $\{a + b\sqrt{2} | a, b \in \mathbb{Z}\}$  is not discrete in  $\mathbb{R}$ . So, the fact that G acts on the underlying rational space  $L^{\mathbb{Q}}$  seems crucial for the discreteness.

*Proof.* Assume that  $\mathbf{b}(g) = \mathbf{o}$ . Then, by the shape of M(g), we have g(v) = v,  $g(w_i) = w_i + a_i(g), g(u) = u + c(g)v$ . From

$$(u, u) = (g(u), g(u)) = (u, u) + 2c(g)(u, v) = (u, u) + 2c(g) ,$$

we have c(g) = 1. Then, from

$$(u, w_i) = (g(u), g(w_i)) = (u, w_i) + a_i(g)(u, v) = (u, w_i) + a_i(g) ,$$

we have  $a_i(g) = 0$ . Hence  $M(g) = I_{n+2}$ , i.e. g = id.

This completes the proof of Proposition (2.4).

**Proposition 2.8.** Let  $U \subset S$  be a compact neighborhood of the unit vector  $\overline{v}$  and let V be a non-empty compact subset of  $S \setminus \{\overline{v}\}$ . (See paragraphs (2.3) and (2.4) for the definition of S and  $\overline{v}$ .) Then, there is a finite index subgroup H of  $\tilde{G}$  s.t.  $(H \setminus \{id\})(V) \subset U$ .

*Proof.* Identifying L with  $S \setminus \{\overline{v}\}$  by  $\iota$  (see Proposition (2.3)), we may assume that

$$V \subset L$$
 and  $B := \{x \in L | ||x|| > r\} \subset U \setminus \{\overline{v}\} \subset L$ 

for some r > 0. Here || \* || is some Euclidean norm of L (w.r.t. some fixed origin). Let  $G \simeq \mathbf{Z}^r$  be a finite index subgroup of  $\tilde{G}$  found in Proposition (2.4). Then,

*G* acts on *L* by parallel transformations, say by  $\{t(g) \in \mathbf{R}^n | g \in G\}$ .

Since the action of G is discrete by Proposition (2.4), we have

$$\inf \left\{ ||t(g)|| \mid g \in G \setminus \{id\} \right\} > 0 .$$

Thus, there is some positive integer c s.t.  $g^c(V) \subset B$  for all  $g \in G \setminus \{id\}$ . Set  $H := \{g^c | g \in G\}$ . Since G is a finitely generated abelian group, this set H is a finite index subgroup of G, and therefore, is a finite index subgroup of  $\tilde{G}$ . This H satisfies the requirement.

4. Now we shall complete the proof of Theorem (1.5)(1).

Let  $\varphi_i : M \longrightarrow B_i$   $(i = 1, \dots, s)$  be mutually different abelian fibrations. For each *i*, we denote by  $\overline{v}_i \in S$  the unit element corresponding to  $\varphi_i$ . Note that  $\overline{v}_i \neq \overline{v}_j$ if  $i \neq j$ .

Put  $r_i := \operatorname{rank} \operatorname{MW}(\varphi_i)$  and choose one of the maximal free abelian subgroups  $\tilde{G}_i \simeq \mathbf{Z}^{r_i}$  of  $\operatorname{Im}(r_{NS} : \operatorname{MW}(\varphi_i) \longrightarrow \operatorname{O}(NS(M)))$  for each *i*.

Let  $U_i \subset S$  be a compact neighborhood of  $\overline{v}_i$ . We can take  $U_i \cap U_j = \emptyset$  if  $i \neq j$ and  $S \setminus \bigcup_{i=1}^s U_i \neq \emptyset$ . Choose  $p \in S \setminus \bigcup_{i=1}^s U_i$  and put  $\overline{U}_i := U_i \cup \{p\}$ . Then, by Proposition (2.8), there are finite index subgroups  $H'_i$  of  $G_i$  s.t.

$$(H'_i \setminus \{id\})(\overline{U}_j) \subset U_i$$

for all pairs  $i \neq j$  in  $\{1, 2, \dots, s\}$ . Then, by Theorem (2.1), we obtain

$$\langle H'_1, H'_2, \cdots, H'_s \rangle = H'_1 * H'_2 * \cdots * H'_s \simeq \mathbf{Z}^{r_1} * \mathbf{Z}^{r_2} * \cdots * \mathbf{Z}^{r_s}$$
.

One can choose free abelian subgroups  $H_i$  of  $MW(\varphi_i)$  s.t.  $r_{NS}(H_i) = H'_i$ . These  $H_i$  satisfy the requirement.

## 3. Proof of Theorem (1.5)(2)

Let *E* be an elliptic curve. Let us consider the product abelian surface  $A := E \times E$ , and its associated Kummer K3 surface S := Km A. By definition, *S* is the minimal resolution of the quotient surface  $A/\iota$  ( $\iota$  is defined below).

Let  $\tilde{p}_i : A \longrightarrow E$  be the projection to the *i*-th factor, and let  $p_i : S \longrightarrow \mathbf{P}^1$  be the Jacobian fibration on S, induced by  $\tilde{p}_i$ .

Let

$$\tilde{f}_2 := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} , \quad \tilde{f}_1 := \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} , \quad \iota := \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$$

and put  $\tilde{G} := \langle \tilde{f}_1, \tilde{f}_2 \rangle$  in SL (2, **Z**). Then  $\tilde{G} = \text{SL}(2, \mathbf{Z})$  (see eg. [Kn, Page 80 Exercise (3.2)]), and SL(2, **Z**) acts faithfully on A by

$$f_2(x,y) = (x+y,y)$$
,  $f_1(x,y) = (x,x+y)$ .

By the shape of  $\tilde{f}_i$ , we see that  $\tilde{f}_i \in MW(\tilde{p}_i)$  and  $\operatorname{ord} \tilde{f}_i = \infty$ . Since  $\tilde{f}_i \iota = \iota \tilde{f}_i$ , we also see that  $\tilde{f}_i$  descends to  $f_i \in MW(p_i)$  and  $\operatorname{ord} f_i = \infty$ .

Set  $G := \langle f_1, f_2 \rangle$  in Aut S. Then,  $G \simeq \mathrm{SL}(2, \mathbf{Z}) / \langle \iota \rangle = \mathrm{PSL}(2, \mathbf{Z}).$ 

We shall show that  $(S, f_1, f_2)$  satisfies the requirement.

It is well-known that  $PSL(2, \mathbb{Z}) \simeq \mathbb{Z}/2 * \mathbb{Z}/3$  (see eg. [Kn, Page 147]). In particular,  $\langle f_1, f_2 \rangle \neq \langle f_1 \rangle * \langle f_2 \rangle$ ; if otherwise,  $\mathbb{Z}/2 * \mathbb{Z}/3 \simeq \mathbb{Z} * \mathbb{Z}$ , a contradiction.

It remains to show that  $\langle f_1^n, f_2^n \rangle = \langle f_1^n \rangle * \langle f_2^n \rangle$  for each  $n \ge 2$ . Our proof of this fact closely follows [Ha, Example 1]. Note that  $PSL(2, \mathbb{Z}) < PGL(2, \mathbb{Z})$ .

In  $PGL(2, \mathbf{Z})$ , we put

$$g_2 := f_2^n = \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$$
,  $g_1 := f_1^n := \begin{pmatrix} 1 & 0 \\ n & 1 \end{pmatrix}$ ,  $j := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .

Then  $j^2 = id$  and  $g_1 = jg_2j$  in PGL(2, **Z**). So, if there would be a non-trivial relation among  $g_1$  and  $g_2$ , say,  $h(g_1, g_2) = id$ , then substituting  $g_1^l = jg_2^l j$  into the relation  $h(g_1, g_2) = id$ , we would have a non-trivial relation among  $g_2$  and j. Thus, we suffice to show that  $\langle g_2, j \rangle = \langle g_2 \rangle * \langle j \rangle$  in PGL(2, **Z**).

To prove this claim, we consider the natural fractional linear action of  $PGL(2, \mathbb{Z})$ on  $\mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$ , and the following subsets and a point of  $\mathbb{P}^1$ :

$$U_1 := \{ z \in \mathbf{C} \mid |z| < 1 \} , \ U_2 := \{ z \in \mathbf{C} \mid |\operatorname{Re} z| > 1 \} \cup \{ \infty \} , \ P := 2\sqrt{-1} \notin U_1 \cup U_2$$

Then  $j(U_2 \cup \{P\}) \subset U_1$  and  $g_2^k(U_1 \cup \{P\}) \subset U_2$  for each  $k \neq 0$  (by  $n \geq 2$ ). Thus, by Theorem (2.1), we have  $\langle g_2, j \rangle = \langle g_2 \rangle * \langle j \rangle$ , and we are done.

## 4. Proof of Theorem (1.6)

Let  $\varphi : S \longrightarrow \mathbf{P}^1$  be a Jacobian K3 surface s.t. rank MW( $\varphi$ ) = 18 and  $\rho(S) = 20$ . Such a Jacobian K3 surface exists by [Co] and [Ni]. It is also known that, given a Jacobian K3 surface M, one can find such a Jacobian K3 surface arbitrarily close to M in the period domain ([Og1]).

We shall show that this S satisfies the requirement.

Let  $e \in \operatorname{Pic} S \simeq NS(S)$  be the fiber class of  $\varphi$ . Since  $\varphi$  has at least three singular fibers (see eg. [Ca]), the subgroup

$$\operatorname{Aut} \varphi := \{ f \in \operatorname{Aut} S \,|\, f^*(e) = e \}$$

of  $\operatorname{Aut} S$  is an almost abelian group (of rank 18).

On the other hand, since  $\rho(S) = 20$ , we have  $\mathbf{Z} * \mathbf{Z} \subset \operatorname{Aut} S$  by [Og3]. In particular, Aut S is not an almost abelian group.

Hence the coset Aut  $S/\text{Aut }\varphi$  is an infinite set. Thus, the Aut S-orbit of e is an infinite set as well. Therefore, S admits infinitely many different Jacobian fibrations with Mordell-Weil rank 18. Now the result follows from Theorem (1.5)(1).

#### References

- [Ca] S. Cantat, Sur la dynamique du groupe d'automorphismes des surfaces K3, Transform. Groups 6 (2001) 201-214.
- [Co] D. A. Cox, Mordell-Weil groups of elliptic curves over  $\mathbf{C}(t)$  with  $p_g = 0$  or 1, Duke Math. J. **49** (1982) 677 689.
- [GHJ] M. Gross, D. Huybrechts and D. Joyce, Calabi-Yau manifolds and related geometries, Springer-Verlag (2003).
- [Ha] P. de la Harpe, Free groups in linear groups, L'Enseignement Math. 29 (1983) 129–144.
- [Hu] D. Huybrechts, Compact hyperkähler manifolds: basic results, Invent. Math. 135 (1999)
  63–113. Erratum: "Compact hyperkähler manifolds: basic results" Invent. Math. 152 (2003) 209–212.
- [Kn] T. Kondo, Group Theory, Iwanami Shoten (1991) (in Japanese).
- [Ma] D. Matsushita, On fibre space structures of a projective irreducible symplectic manifold, Topology 38 (1999) 79–83. Addendum: On fibre space structures of a projective irreducible symplectic manifold, Topology 40 (2001) 431–432.
- [Mc] C. T. McMullen, Dynamics on K3 surfaces: Salem numbers and Siegel disks, J. Reine Angew. Math. 545 (2002) 201–233.
- [Ny] K. Nishiyama, Examples of Jacobian fibrations on some K3 surfaces whose Mordell-Weil lattices have the maximal rank 18, Comment. Math. Univ. St. Paul. 44 (1995) 219–223.
- [Og1] K. Oguiso, Local families of K3 surfaces and applications, J. Algebraic Geom. 12 (2003) 405–433.
- [Og2] K. Oguiso, Bimeromorphic automorphism groups of non-projective hyperkähler manifolds
   a note inspired by C. T. McMullen, revised.
- [Og3] K. Oguiso, Tits alternative in hypekähler manifolds, Math. Research. Letters 13 (2006) 307–316.
- [Og4] K. Oguiso, Salem polynimials and bimeromprphic automorphisms of hyperkähler manifolds, revised in Sugaku Exposition (in Japanese).
- [Th] W. P. Thurston, *Three-dimensional geometry and topology*, Princeton University Press (1997).

8

[Ti] J. Tits, Free subgroups in linear groups, J. Algebra 20 (1972) 250–270.

DEPARTMENT OF ECONOMY, KEIO UNIVERSITY, HIYOSHI HOKUKU-KU YOKOHAMA, JAPAN, AND KOREA INSTITUTE FOR ADVANCED STUDY, 207-43 CHEONRYANGNI-2DONG, DONGDAEMUN-GU, SEOUL 130-722, KOREA

E-mail address: oguiso@hc.cc.keio.ac.jp