On Lusztig's conjecture for connected and disconnected exceptional groups

Olivier BRUNAT brunat@math.rwth-aachen.de

October 29, 2018

ABSTRACT

Lusztig conjectured that the almost characters of a finite reductive group are up to a scalar the same as the characteristic functions of the rational character sheaves defined on the corresponding algebraic group. We propose in this paper to verify this conjecture for the uniform almost characters of the Suzuki and the Ree groups and to show that a conjecture of the same type holds for the disconnected case of type B_2 . We then state a conjecture on the possible values of the uniform almost characters on unipotent elements in the disconnected case of type F_4 .

1 Introduction

Let \mathbf{G}_0 be a connected reductive group defined over the finite field with q elements and $F_0: \mathbf{G}_0 \to \mathbf{G}_0$ such that F_0^{δ} is a Frobenius map for some positive integer δ . We denote by $\mathbf{G}_0^{F_0}$ and $\mathbf{G}_0^{F_0^{\delta}}$ the finite groups of fixed points under F_0 and F_0^{δ} respectively. The group $\mathbf{G}_0^{F_0^{\delta}}$ is F_0 -stable and the restriction of F_0 to $\mathbf{G}_0^{F_0^{\delta}}$ (denoted by the same symbol) is an automorphism of $\mathbf{G}_0^{F_0^{\delta}}$. We are interested in the irreducible characters of the extension $\tilde{G}_0 = \mathbf{G}_0^{F_0^{\delta}} \rtimes \langle F_0 \rangle$. A relevant example in which this situation occurs is when \mathbf{G}_0 is a simple algebraic group of type B_2 , G_2 or F_4 and F_0 is the isogeny that defines the Suzuki and the Ree groups; in these cases, F_0^2 is a Frobenius map and $\mathbf{G}_0^{F_0^2}$ is a simple group (except for the first value of the parameter). In [2] and [3], we compute the character table of \tilde{G}_0 in the cases of type B_2 and G_2 respectively. But the employed methods are essentially *ad hoc* and cannot be used for type F_4 .

When the algebraic group is connected, Lusztig has shown that the computation of the irreducible characters of $\mathbf{G}_0^{F_0}$ is reduced to the determination of the *almost characters* of $\mathbf{G}_0^{F_0}$ (this is a family of class functions on $\mathbf{G}_0^{F_0}$ defined using the generalized characters of Deligne-Lusztig) [12]. To obtain the values of these functions, he develops the theory of character sheaves on \mathbf{G}_0 : to every F_0 stable character sheaf on \mathbf{G}_0 we can associate in a natural way a class function on $\mathbf{G}_0^{F_0}$, the so-called characteristic function of the sheaf [14]. He conjectures that the characteristic functions of character sheaves coincide (up to a scalar) with the almost characters [14, V]. With some assumptions on the characteristic and on the center of \mathbf{G}_0 , Shoji showed that this conjecture holds when F_0 is a Frobenius map [23]. Recently, Bonnafé proved that Lusztig's conjecture holds in type A_n when the center of \mathbf{G}_0 is not connected [1].

The advantage of this approach is that the characteristic functions of F_0 stable character sheaves can be computed using a uniform algorithm described by Lusztig in [14, V.§24]; the input of this algorithm is the generalized Springer correspondence introduced by Lusztig [13].

Recently, Lusztig started a series of papers in which he develops a theory of character sheaves for disconnected groups and defines a generalized Springer correspondence in this case [15]; we do not know at present whether Lusztig's algorithm is valid in this situation.

In this note, we propose to apply this theory to the groups \tilde{G}_0 interpreted as a fixed-point subgroup of a disconnected reductive group $\tilde{\mathbf{G}} = \mathbf{G} \rtimes \langle \tau \rangle$ (here \mathbf{G} is a connected reductive group constructed from \mathbf{G}_0 and τ is an automorphism of finite order of \mathbf{G}) under a generalized Frobenius F, using a method developed by Digne in [6]. In the case that \mathbf{G}_0 is of type B_2 or F_4 , the element τ (viewed as an element of $\tilde{\mathbf{G}}$) is unipotent and we thus can determine the generalized Springer correspondence on the coset $\mathbf{G}\tau$. We can then apply Lusztig's algorithm *a priori* as in the connected case; we show that Lusztig's conjecture holds for the disconnected type B_2 (using the explicit table in [2]) and we give the possible values of the uniform almost characters on unipotent elements for the type F_4 .

The paper is organized as follows: in Section 2 we determine the classes of $\tilde{\mathbf{G}}$ from those of \mathbf{G}_0 . In Section 3, we apply Lusztig's algorithm to \mathbf{G}_0 (in the case of types B_2 , G_2 and F_4) with the twisted map F_0 that defines the Suzuki and Ree groups and verify that Lusztig's conjecture holds in those cases for the uniform almost characters on the unipotents elements. In Section 4, we verify Lusztig's conjecture in the disconnected case of type B_2 and give a conjecture for the case F_4 .

2 The group \widetilde{G}_0

In this section, \mathbf{G}_0 is a connected reductive group defined over \mathbb{F}_q (where q is a power of 2) and F_0 is a map on \mathbf{G}_0 such that F_0^2 is a Frobenius map.

2.1 Restriction of scalars

Now we will interpret the group $\widetilde{G}_0 = \mathbf{G}_0^{F_0^2} \rtimes \langle F_0 \rangle$ as a fixed-point subgroup under an endomorphism F of an algebraic group $\widetilde{\mathbf{G}}$; we use the construction developed in [6]. We set $\mathbf{G} = \mathbf{G}_0 \times \mathbf{G}_0$ and $\widetilde{\mathbf{G}} = \mathbf{G} \rtimes \langle \tau \rangle$, where τ is the algebraic automorphism of \mathbf{G} defined by $(g_1, g_2)^{\tau} = (g_2, g_1)$. The group $\tilde{\mathbf{G}}$ is a non-connected reductive group with identity component \mathbf{G} . Let $F: \tilde{\mathbf{G}} \to \tilde{\mathbf{G}}$ be the homomorphism defined by $F(g_1, g_2) = (F_0(g_2), F_0(g_1))$ and $F(\tau) = \tau$. Let \mathbf{T}_0 be a maximal rational torus of \mathbf{G}_0 contained in a rational Borel subgroup of \mathbf{G}_0 . The group $\mathbf{B} = \mathbf{B}_0 \times \mathbf{B}_0$ is an *F*-rational Borel subgroup of \mathbf{G} containing $\mathbf{T} = \mathbf{T}_0 \times \mathbf{T}_0$ as maximal *F*-rational torus. The groups \mathbf{T} and \mathbf{B} are τ -stable, thus τ is quasi semisimple. It is not difficult to show that τ is in fact quasi-central [8, 1.15]. We put $\tilde{\mathbf{B}} = \mathbf{B} \rtimes \langle \tau \rangle$ and $\tilde{\mathbf{T}} = \mathbf{T} \rtimes \langle \tau \rangle$; we have $\tilde{\mathbf{B}} = N_{\tilde{\mathbf{G}}}(\mathbf{B})$ and $\tilde{\mathbf{T}} = N_{\tilde{\mathbf{G}}}(\mathbf{T}, \mathbf{B})$. Thus $\tilde{\mathbf{T}}$ and $\tilde{\mathbf{B}}$ are respectively rational "torus" and rational "Borel" in the sense of Digne and Michel [8, 1.2].

Proposition 2.1 We have the following isomorphisms:

$$\begin{array}{lll} \mathbf{G}^{\tau} &\simeq & \mathbf{G}_{0}, \\ \left(\mathbf{G}^{\tau}\right)^{F} &\simeq & \mathbf{G}_{0}^{F_{0}}, \\ \mathbf{G}^{F} &\simeq & \mathbf{G}_{0}^{F_{0}^{2}}, \\ \mathbf{\widetilde{G}}^{F} &\simeq & \mathbf{G}_{0}^{F_{0}^{2}} \rtimes \left\langle F_{0} \right\rangle \end{array}$$

Proof — The isomorphisms are given by $\varphi : \mathbf{G}_0 \to \mathbf{G}^{\tau}, \ g \mapsto (g,g)$ and $\varphi' : \mathbf{G}_0^{F_0^2} \to \mathbf{G}^F, \ g \mapsto (g, F_0(g))$. Moreover we have $\varphi' \circ F_0 = \tau \circ \varphi'$, it follows that $F|_{\mathbf{G}^F}$ acts on \mathbf{G}^F as $F_0|_{\mathbf{G}_0^{F_0^2}}$ on $\mathbf{G}_0^{F_0^2}$ and the result holds.

2.2 Conjugacy classes

We will now describe the conjugacy classes of the groups appearing in §2.1. We are interested in the conjugacy classes of elements belonging to the coset $\mathbf{G}\tau$.

Lemma 2.1 Two elements $(g_1, g_2)\tau$ and $(g'_1, g'_2)\tau$ of $\mathbf{G}\tau$ are conjugate in $\widetilde{\mathbf{G}}$ if and only if g_1g_2 and $g'_1g'_2$ are conjugate in \mathbf{G}_0 .

Proof — First recall that $(g_1, g_2)\tau$ and $(g'_1, g'_2)\tau$ are conjugate in \mathbf{G} if and only if they are conjugate by an element in \mathbf{G} . Then there exists $h = (h_1, h_2) \in \mathbf{G}$ such that $h_1g_1 = g'_1h_2$ and $h_2g_2 = g'_2h_1$. It follows that g_1g_2 and $g'_1g'_2$ are conjugate. Conversely if $g_1g_2 = hg'_1g'_2h^{-1}$ for some $h \in \mathbf{G}_0$, we have:

$$(g_1, 1)(g_2, g_1)\tau(g_1, 1)^{-1} = (g_1g_2, 1)\tau = (hg'_1g'_2h^{-1}, 1)\tau = (hg'_1, h)(g'_2, g'_1)\tau(hg'_1, h)^{-1}$$

Since $\tau(g_1, g_2)\tau\tau^{-1} = (g_2, g_1)\tau$ the result follows.

In the following, if H is a subgroup of \mathbf{G}_0 , we denote by $\mu(H) = \{(h,h) \in \mathbf{G} \mid h \in H\} \subseteq \mathbf{G}$.

Proposition 2.2 The map $f : \mathbf{G}_0 \mapsto \mathbf{G}\tau$, $x \mapsto (x, 1)\tau$ induces a bijection from the conjugacy classes of \mathbf{G}_0 to the classes in the coset $\mathbf{G}\tau$ and we have

$$C_{\widetilde{\mathbf{G}}}(f(x)) = \mu(C_{\mathbf{G}_0}(x)).\langle (x, 1)\tau \rangle.$$

Moreover f induces a bijection between the unipotent classes of \mathbf{G}_0 and the unipotent classes in the coset \mathbf{G}_{τ} .

Proof — We first remark that every conjugacy class in the coset $\mathbf{G}\tau$ has a representative of the form f(g) for some $g \in \mathbf{G}_0$ because $(x_1, x_2)\tau$ is conjugate to $f(x_1x_2)$. The bijection between the classes is now a consequence of Lemma 2.1. Moreover we have:

$$((x_1, x_2)\tau)^{2m} = ((x_1x_2)^m, (x_2x_1)^m).$$

We denote by d the order of $(x_1, x_2)\tau$ and by d' the one of x_1x_2 . Since x_1x_2 and x_2x_1 are conjugate, they have the same order. Thus we have $((x_1, x_2)\tau)^{2d'} =$ 1 and it follows that d divides 2d'. Moreover 2 divides d, then there exists a positive integer k such that d = 2k. Thus k divides d'. Conversely we have 1 = $((x_1, x_2)\tau)^{2k}$; it follows that $(x_1x_2)^k = 1$ and d' divides k. We have k = d' and we have to prove that

$$|\langle ((x_1, x_2), \tau) \rangle| = 2 |\langle x_1 x_2 \rangle|.$$

Recall that the characteristic of the considered groups is 2, it follows that the bijection induced by f is compatible with the unipotent classes.

Let $x \in \mathbf{G}_0$, we then have $\mu(\mathbf{C}_{\mathbf{G}_0}(x)).\langle (x,1)\tau \rangle \subseteq \mathbf{C}_{\mathbf{G}}(f(x))$. Conversely if $(g_1,g_2)(x,1)\tau = (x,1)\tau(g_1,g_2)$ we have $(g_1x,g_2)\tau = (xg_2,g_1)\tau$ and $g_1 \in \mathbf{C}_{\mathbf{G}_0}(x)$. If $(g_1,g_2)\tau(x,1)\tau = (x,1)\tau(g_1,g_2)\tau$, then $(g_1,g_2x) = (xg_2,g_1)$ and $g_2 \in \mathbf{C}_{\mathbf{G}_0}(x)$ as required.

We now discuss the *F*-rational classes of $\tilde{\mathbf{G}}$ in the coset $\mathbf{G}\tau$. We first recall a relevant result which is a direct consequence of Lang's theorem:

Theorem 2.1 Let **G** be a connected reductive group defined over \mathbb{F}_q and F a generalized Frobenius map on **G**. Suppose that **G** acts transitively on a set X, suppose that there exists $F' : X \to X$ such that F'(gx) = F(g)F'(x) for every $g \in \mathbf{G}$ and $x \in X$ and suppose finally that $\operatorname{Stab}_{\mathbf{G}}(x)$ is a closed subgroup of **G**. Then $X^{F'}$ is non-empty and if $x_0 \in X^{F'}$, we set

$$A(x_0) = \operatorname{Stab}_{\mathbf{G}}(x_0) / \operatorname{Stab}_{\mathbf{G}}(x_0)^{\circ}$$

and by π the canonical projection of $\operatorname{Stab}_{\mathbf{G}}(x_0)$ onto $A(x_0)$. Then the map F naturally induces an automorphism on $A(x_0)$, denoted by the same symbol, and the \mathbf{G}^F -orbits of $X^{F'}$ are in bijection with the F-classes of $A(x_0)$ (i.e., the orbits on $A(x_0)$ for the operation $g.h = g^{-1}hF(g)$). Moreover the correspondence is such that if $x = gx_0$ then we associate to the \mathbf{G}^F -class of x the F-class of $\pi(g^{-1}F(g))$.

Let $x\tau \in \widetilde{\mathbf{G}}$, we denote by $A(x\tau) = C_{\mathbf{G}}(x\tau)/C_{\mathbf{G}}(x\tau)^{\circ}$. If $x\tau$ is rational, we denote by F the automorphism of $A(x\tau)$ induced by F. We have:

Proposition 2.3 Every rational class on $\mathbf{G}\tau$ has an F-stable representative. Moreover the map f defined in Proposition 2.2 gives a bijection between the F_0 -rational classes of \mathbf{G}_0 and the F-classes in the coset $\mathbf{G}\tau$. Finally, the conjugacy classes of $\mathbf{G}_0^{F_0}$ are in bijection with the conjugacy classes of $\mathbf{G}_0^{F_0}$ are in bijection with the conjugacy classes of $\mathbf{G}_0^{F_0}$ and if \tilde{g} is a representative of a class of $\mathbf{G}_0^{F_0}$ and if \tilde{g} is a representative of the corresponding class on $\mathbf{G}^F\tau$ then we have

$$\begin{array}{lll} \mathbf{C}_{\widetilde{\mathbf{G}}^{F}}(\widetilde{g}) &\simeq & \mu(\mathbf{C}_{\mathbf{G}_{0}^{F}}(g)).\langle (g,1)\tau \rangle, \\ |\mathbf{C}_{\widetilde{\mathbf{G}}^{F}}(\widetilde{g})| &= & 2|\mathbf{C}_{\mathbf{G}_{0}^{F}}(g)|. \end{array}$$

Proof — The map f gives a bijection between the F_0 -rational classes of \mathbf{G}_0 and the F-classes in the coset \mathbf{G}_{τ} because $F((g, 1)\tau)$ is conjugate to $(g, 1)\tau$ if and only if $F_0(g)$ and g are conjugate. Let $g_0 \in \mathbf{G}_0^{F_0}$, then we have

$$F((g_0, 1)\tau) = (g_0, 1)^{-1}(g_0, 1)\tau(g_0, 1).$$

Since $(g_0, 1) \in \mathbf{G}$, we can find $u \in \mathbf{G}$ such that $(g_0, 1) = u^{-1}F(u)$ (we apply Lang's property in the connected group \mathbf{G} with the generalized Frobenius F). The element $u(g_0, 1)\tau u^{-1}$ is rational and we have

$$u \operatorname{C}_{\mathbf{G}}((g_0, 1)\tau)u^{-1} = \operatorname{C}_{\mathbf{G}}(u(g_0, 1)\tau u^{-1}).$$

Let $g, h \in C_{\mathbf{G}}((g_0, 1)\tau)$, we then have

$$uh^{-1}u^{-1}ugu^{-1}F(u)F(h)F(u^{-1}) = uh^{-1}g(g_0,1)F(h)(g_0,1)^{-1}u^{-1}.$$

Consequently the *F*-classes of $C_{\mathbf{G}}(u(g_0, 1)\tau u^{-1})$ are in bijection with the $^{(g_0,1)}F$ classes of $C_{\mathbf{G}}((g_0, 1)\tau)$. We can apply Theorem 2.1 with $X = \operatorname{Cl}((g_0, 1)\tau)$ and F'being the restriction of *F* to this set; we take $u(g_0, 1)\tau u^{-1}$ as rational element. It follows that the $\widetilde{\mathbf{G}}^F$ -classes of $\operatorname{Cl}(f(x_0))^F$ are in bijection with the $^{(g_0,1)}F$ classes of $A(f(x_0))$. Using Proposition 2.2, we have $C_{\mathbf{G}}(f(g_0)) = \mu(C_{\mathbf{G}_0}(g_0))$. For every $(h, h) \in C_{\mathbf{G}}(f(g_0))$, we then have

$$(g_0,1)F(h,h) = (g_0F_0(h),F_0(h)),$$

but since F_0 acts on $C_{\mathbf{G}_0}(g_0)$ we have $F_0(h) \in C_{\mathbf{G}_0}(g_0)$ and it follows that the operation of ${}^{(g_0,1)}F$ on $C_{\mathbf{G}}(f(g_0))$ is the same as the one of F on $C_{\mathbf{G}}(f(g_0))$. Since μ is an isomorphism of algebraic groups, it follows that $A(f(g_0))$ is isomorphic to $A_{\mathbf{G}_0}(g_0) = C_{\mathbf{G}_0}(g_0)/C_{\mathbf{G}_0}(g_0)^{\circ}$ and we have

$$F \circ \mu = \mu \circ F_0,$$

thus the F_0 -classes of $A_{\mathbf{G}_0}(g_0)$ are in bijection with the *F*-classes of $A(f(g_0))$. This proves that the classes of \mathbf{G}_0^F are in correspondence with the classes of $\widetilde{\mathbf{G}}^F$ in the coset $\mathbf{G}\tau$. We set π (resp. π_0) to be the canonical projection of $C_{\mathbf{G}}(f(g_0))$ onto $A(f(g_0))$ (resp. $C_{\mathbf{G}_0}(g_0)$ onto $A_{\mathbf{G}_0}(g_0)$). Let $\pi_0(a)$ be a representative of a F_0 -class of $A_{\mathbf{G}_0}(g_0)$ and let \tilde{a} be a corresponding class representative of $\mathbf{G}_0^{F_0}$ given in Theorem 2.1. We then have

$$C_{\mathbf{G}_0^{F_0}}(\widetilde{a}) \simeq \{h \in C_{\mathbf{G}_0}(g_0) \mid F(h) = a^{-1}ha\}.$$

Moreover the element $\pi(a, a) = (\pi_0(a), \pi_0(a))$ is a representative of the $(g_0, 1)F$ class of $A(f(g_0))$. Thus

$$C_{\mathbf{G}^F}(\widetilde{a,a}) \simeq \{h \in C_{\mathbf{G}}(f(g_0)) \mid F(h) = a^{-1}ha\},\$$

where $(\widetilde{a,a})$ is a representative of the class of $\widetilde{\mathbf{G}}^F$ corresponding to $\pi(a,a)$. Hence $C_{\mathbf{G}^F}(\widetilde{a,a})$ is isomorphic to $\mu(C_{\mathbf{G}^F_o}(\widetilde{a}))$ and the result is proven.

Remark 2.1 The element u in the preceding proof can be explicitly chosen as follows: since g_0 belongs to the connected group \mathbf{G}_0 , using Lang's property in this group, we can find $x \in \mathbf{G}_0$ such that $g_0 = x^{-1}F_0^2(x)$. We then have

$$(g_0, 1) = (x, F_0(x))^{-1} F(x, F_0(x)).$$

It follows that $y = (F_0(x)x^{-1}, F_0(F_0(x)x^{-1}))\tau$ is a rational element in the rational class of $(g_0, 1)\tau$. In [7] Digne and Michel define the Shintani correspondence: this is a bijection N_{F/F^2} between the conjugacy classes of $\mathbf{G}_0^{F_0}$ and the conjugacy classes of $\mathbf{G}_0^{F_0^2} \rtimes \langle F_0 \rangle$ in the coset $\mathbf{G}_0^{F_0^2} F_0$. We then remark that

$$y = \varphi' \left((N_{F/F^2}(g_0))^{-1} \right)$$

where φ' is the explicit isomorphism between $\mathbf{G}_0^{F_0^2} \rtimes \langle F_0 \rangle$ and $\widetilde{\mathbf{G}}^F$ constructed in the proof of Proposition 2.1.

3 Lusztig's conjecture for the Suzuki and the Ree groups

In this section, we will prove that the simple groups of types B_2 , G_2 and F_4 with the Frobenius map defining the Suzuki and Ree groups satisfy Lusztig's conjecture. For the explicit calculations in this section, we will use programs developed in GAP [9] by the author. We will first recall some generalities.

3.1 Generalized Springer correspondence

Let \mathbf{G}_0 be a connected reductive group defined over the finite field of q elements and F_0 be the corresponding Frobenius map. We set \mathcal{N} to be the \mathbf{G}_0 -classes of pairs (u, ϕ) , where u is a unipotent element of \mathbf{G}_0 and ϕ an irreducible character of the component group $A_{\mathbf{G}_0}(u)$ defined in the proof of Proposition 2.3. Two pairs (u, ϕ) and (v, ψ) are in the same \mathbf{G}_0 -class if there existe $x \in \mathbf{G}_0$ such that $v = x^{-1}ux$ and $\psi(t) = \phi(xtx^{-1})$ for every $t \in A_{\mathbf{G}_0}(v)$ (note that it is independent of the choice of x). In [13], Lusztig associates to every pair (u, ϕ) a 4-tuple $(\mathbf{L}, v, \psi, \rho)$, where \mathbf{L} is a Levi subgroup of $\mathbf{G}_0, v \in \mathbf{L}$ is unipotent, $\psi \in \operatorname{Irr}(A_{\mathbf{L}}(v))$ and $\rho \in \operatorname{Irr}(N_{\mathbf{G}_0}(\mathbf{L})/\mathbf{L})$. If $\mathbf{L} = \mathbf{G}_0$ then the pair is said *cuspidal*. Lusztig's construction defines a bijection between \mathcal{N} and the set of 4-tuples as above, such that (v, ψ) is cuspidal for \mathbf{L} . We write $\rho_{u,\phi}$ for ρ if (u, ϕ) corresponds to $(\mathbf{L}, v, \psi, \rho)$. The map $(u, \phi) \mapsto \rho_{u,\phi}$ is the generalized Springer correspondence.

3.2 Lusztig's conjecture

We use the same notation as in §3.1. We denote by \mathbf{T}_0 a maximal rational torus of \mathbf{G}_0 contained in a rational Borel \mathbf{B}_0 of \mathbf{G}_0 . We define the Weyl group $W = \mathrm{N}(\mathbf{T}_0)/\mathbf{T}_0$ of \mathbf{G}_0 . Since \mathbf{T}_0 is rational, it follows that F_0 acts as an automorphism on W; we denote it with the same symbol. Let $\rho \in \mathrm{Irr}(W)$ such that $\rho^{F_0} = \rho$; using Clifford theory we can extend ρ to $\widetilde{W} = W \rtimes \langle F_0 \rangle$. We choose such an extension $\widetilde{\rho}$ and we define the *uniform almost characters of* $\mathbf{G}_0^{F_0}$ as

$$R_{\widetilde{\rho}} = \frac{1}{|W|} \sum_{w \in W} \widetilde{\rho}(wF_0) R_w,$$

where R_w is the generalized Deligne-Lusztig character $R_{\mathbf{T}_w}^{\mathbf{G}_0}(\mathbf{1}_{T_w}^{F_0})$ associated to the rational torus \mathbf{T}_w of \mathbf{G}_0 corresponding to w. For definition and properties of these characters we refer for example to [5, §7].

Let C be a rational unipotent class in \mathbf{G}_0 and \mathcal{E} be an F_0 -stable irreducible local system on \mathbf{G}_0 which is \mathbf{G}_0 -equivariant for the conjugation action of \mathbf{G}_0 . In [14], Lusztig associates to (C, \mathcal{E}) an irreducible F_0 -stable perverse sheaf on \mathbf{G}_0 . We can attach to it its characteristic function $X_{C,\mathcal{E}}$ on $\mathbf{G}_0^{F_0}$ ([14, II.8.4] for the definition), defined up to a scalar.

Following [22, 3.5], there is a correspondence between the pairs (C, \mathcal{E}) defined previously and the pairs (u, ϕ) defined in §3.1 such that u is rational and ϕ is F_0 stable (the set of such pairs is denoted by \mathcal{N}^{F_0} in the following); we denote by $X_{u,\phi}$ the corresponding characteristic function in this correspondence. We denote by \mathcal{N}_0 (resp. $\mathcal{N}_0^{F_0}$) the pairs of \mathcal{N} (resp. \mathcal{N}^{F_0}) such that the associated 4tuple has the form $(\mathbf{T}_0, 1, 1, \rho_{u,\phi})$. Note that the restriction of the map $(u, \phi) \mapsto$ $\rho_{u,\phi}$ to \mathcal{N}_0 is the original Springer correspondence.

Lusztig conjectures that the uniform almost characters $R_{\tilde{\rho}_{u,\phi}}$ and the characteristic functions $X_{u,\phi}$ with $(u,\phi) \in \mathcal{N}_0^{F_0}$ coincide up to a scalar. We propose to verify that this conjecture holds on unipotent elements.

Remark 3.1 Lusztig's conjecture is more general. Lusztig defines in [12, 13.6] the almost characters. Lusztig's conjecture asserts that these class functions coincide up to a scalar with the characteristic functions of every F_0 -stable character sheaf on \mathbf{G}_0 . To compute the values of these functions, it is sufficient to

compute the values of the characteristic functions $X_{C,\mathcal{E}}$ on the unipotent elements; see [14, II.8.5].

3.3 Lusztig's algorithm

We keep the notation of the preceding sections. Our aim is to compute the values of the characteristic functions $X_{u,\phi}$ for $(u,\phi) \in \mathcal{N}_0^{F_0}$ on the unipotent elements of $\mathbf{G}_0^{F_0}$ and to compare them with the uniform almost characters of $\mathbf{G}_0^{F_0}$. Let $(u,\phi) \in \mathcal{N}_0^{F_0}$; we define $\varphi_{u,\phi} : \operatorname{Cl}_{\mathbf{G}_0}(u)^{F_0} \to \overline{\mathbb{Q}}_{\ell}$ as follows: using The-

Let $(u, \phi) \in \mathcal{N}_0^{F_0}$; we define $\varphi_{u,\phi} : \operatorname{Cl}_{\mathbf{G}_0}(u)^{F_0} \to \overline{\mathbb{Q}}_{\ell}$ as follows: using Theorem 2.1, if a is a representative of an F_0 -class of $A_{\mathbf{G}_0}(u)$, we denote by g_a the corresponding class representative of $\mathbf{G}_0^{F_0}$. We set $\varphi_{u,\phi}(g) = \widetilde{\phi}(aF_0)$ if gis conjugate to g_a in $\mathbf{G}_0^{F_0}$ and $\varphi_{u,\phi}(g) = 0$ if $g \notin \operatorname{Cl}_{\mathbf{G}_0}(u)^{F_0}$; it depends on the choice of an extension $\widetilde{\phi}$ of the F_0 -stable character ϕ to $A_{\mathbf{G}_0}(u) \rtimes \langle F_0 \rangle$ and hence it is only well-defined up to a non-zero scalar multiple. Note that the class function $\varphi_{u,\phi}$ is up to a scalar the class function $Y_{u,\phi}$ defined in [14, V.24.2.3].

We now recall Lusztig's algorithm. We first choose an order on $\mathcal{N}_0^{F_0}$ extending the natural order on unipotent classes of $\mathbf{G}_0^{F_0}$ defined by $u \leq v$ when $\operatorname{Cl}_{\mathbf{G}_0}(u)$ is contained in the Zariski closure of $\operatorname{Cl}_{\mathbf{G}_0}(v)$. Following Lusztig, we write $(u, \phi) \sim$ (v, ψ) when u is conjugate to v. If u is a unipotent element of \mathbf{G}_0 , we denote by \mathcal{B}_u the variety of Borel subgroups of \mathbf{G}_0 containing u. We set $d_u = \dim(\mathcal{B}_u)$. We remark that if $u \leq v$ then $d_v \leq d_u$. In [14], Lusztig shows that the funtions $Y_{u,\phi}$ (with $(u,\phi) \in \mathcal{N}_0^{F_0}$) form a basis of the uniform functions on the set of unipotent elements of $\mathbf{G}_0^{F_0}$. Note that the function $X_{u,\phi}$ defined in [14, V.24.2.8] is the characteristic function of the sheaf divided by $q^{-(a_0+r)/2}$; in our case, we have $a_0 + r = \dim(\mathbf{C}_{\mathbf{G}_0}(u)) - \operatorname{rk}(\mathbf{G}_0) = 2d_u$; see [25, 2.7]. With our notation, we have

$$q^{-d_u} X_{u,\phi} = \sum_{(v,\psi)\in\mathcal{N}_0^{F_0}} p_{(v,\psi),(u,\phi)} Y_{v,\psi}.$$
 (1)

We set P to be the matrix with coefficients $p_{(v,\psi),(u,\phi)}$. In [14, V.24], Lusztig shows that the coefficients of P are rational, that the diagonal elements of Pare 1 and that P is an upper triangular matrix. Moreover, if $(u, \phi) \sim (v, \psi)$ but $(u, \phi) \neq (v, \psi)$ then $p_{(v,\psi),(u,\phi)} = 0$. We give a way to compute the matrix P. We introduce the nonsingular bilinear form

$$(f, f') = \sum_{g \in X} f(g) \widetilde{f'}(g),$$

where X denotes the set of unipotent elements of $\mathbf{G}_0^{F_0}$ and $\tilde{f'}$ is defined in [14, V.24.2.12]. We set

$$w_{(u,\phi),(v,\psi)} = \frac{|\mathbf{G}_{0}^{F_{0}}|}{|W|} \sum_{w \in W} \frac{\widetilde{\rho}_{u,\phi}(wF_{0})\widetilde{\rho}_{v,\psi}(wF_{0})}{|\mathbf{T}_{w}^{F_{0}}|q^{d_{u}+d_{v}}},$$

where $\tilde{\rho}_{u,\phi}$ is an extension of the F_0 -stable character $\rho_{u,\phi}$ given in §3.1 to the group $W \rtimes \langle F_0 \rangle$ and \mathbf{T}_w is the rational torus of \mathbf{G}_0 corresponding to w.

u	$A_{\mathbf{G}_0}(u)$	ϕ	$ ho_{u,\phi}$	d_u
u_1	1	1	ε	4
u_2	1	1	φ_1	2
u_3	1	1	φ_2	2
u_4	1	1	χ	1
u_5	$\mathbb{Z}/2\mathbb{Z}$	1	1_W	0

Table 1: Generalized Springer correspondence for type B_2 .

We set Ω to be the matrix with coefficients $w_{(u,\phi),(v,\psi)}$. Lusztig proves that the matrix P defined as above is the unique solution of

$${}^{t}P\Lambda P = \Omega,$$

where Λ is a block diagonal matrix, which is uniquely determined from the equation when we choose the dimension of the blocks compatible with the relation \sim .

3.4 The Suzuki groups

Let \mathbf{G}_0 be a simple algebraic group of type B_2 defined over the algebraic closure of the field of 2 elements. Let n be a non-negative integer; we set $q = 2^n \sqrt{2}$ and $F_0 = \sigma \circ F_{2^n}$, where F_{2^n} is the standard Frobenius map on \mathbf{G}_0 over \mathbb{F}_{2^n} and σ is the bijective endomorphism arising from the symmetry of the Dynkin diagram, see [4, §12.3]. The Suzuki group of parameter q^2 is defined as $\mathbf{G}_0^{F_0}$; it has order $q^4(q^2 - 1)(q^4 + 1)$. The Weyl group of \mathbf{G}_0 is the dihedral group of order 8. We denote by w_a and w_b the two generators of W corresponding to the roots a and b respectively. We denote by $x_r(t)$ (for a root r and $t \in \mathbb{F}_2$) the Chevalley generators of \mathbf{G}_0 and we set $x_a = x_a(1)$ and $x_b = x_b(1)$. By convention, we choose b for the long root. The group W has 5 irreducible characters. We denote by ε the sign character of W and by χ the irreducible character of degree 2. The two other linear characters are denoted by φ_1 and φ_2 , the choice is such that $\varphi_1(w_a) = -1$.

In Table 1, we recall the generalized Springer correspondence for a simple group of type B_2 in characteristic 2; see [25, IV.1]. We set $u_1 = 1$, $u_2 = x_b$, $u_3 = x_a$, $u_4 = x_{a+b}x_{2a+b}$ and $u_5 = x_ax_bx_{a+b}$ as representatives for the unipotent classes of \mathbf{G}_0 . The F_0 -stable classes of \mathbf{G}_0 have representatives u_1 , u_4 and u_5 . The class $\operatorname{Cl}(u_5)^{F_0}$ splits into two classes of $\mathbf{G}_0^{F_0}$; representatives of these classes are $x_ax_bx_{a+b}$ and $x_ax_bx_{2a+b}$ and are denoted by ρ and ρ^{-1} respectively in [27].

3.4.1 Uniform almost characters

The Weyl group W of \mathbf{G}_0 has three F_0 -stable characters $\mathbf{1}_W$, ε and χ . We choose extensions $\widetilde{\mathbf{1}}_W$, $\widetilde{\varepsilon}$ and $\widetilde{\chi}$ to $W \rtimes \langle F_0 \rangle$. The values of these characters appear in Table 2. The decomposition into irreducible components of the uniform almost characters (computed using the definition of uniform almost characters in §3.2)

	F_0	$w_a F_0$	$w_a w_b w_a F_0$
$\widetilde{\varepsilon}$	1	-1	-1
$\widetilde{\chi}$	0	$-\sqrt{2}$	$\sqrt{2}$
$\widetilde{1}_W$	1	1	1

Table 2: Values of extensions on WF_0 .

	u_1	u_4	ρ	$ ho^{-1}$
$R_{\widetilde{\varepsilon}}$	q^4	0	0	0
$R_{\widetilde{\chi}}$	$q(q^2-1)$	-q	0	0
$R_{\widetilde{1}_W}$	1	1	1	1

Table 3: Values of the uniform almost characters of the Suzuki group.

of the Suzuki groups are given in [10, Table 1]. We have $R_{\tilde{1}_W} = 1_{\mathbf{G}_0^{F_0}}$, $R_{\tilde{\varepsilon}} = \operatorname{St}_{\mathbf{G}_0^{F_0}}$ and $R_{\tilde{\chi}} = \frac{1}{\sqrt{2}}(W_1 + W_2)$, where W_1 and W_2 are the two cuspidal unipotent characters of the Suzuki group (see [27]); note that the results there are correct, however, one has to use our choices for $\tilde{\chi}$ given in Table 2 contrary to what they claim in [10, Table 1]. Thus, using the character table of the Suzuki group given in [27], we deduce

Proposition 3.1 The values of the uniform almost characters of the Suzuki groups on the unipotent elements of $\mathbf{G}_{0}^{F_{0}}$ are given in Table 3.

3.4.2 Lusztig's algorithm for the Suzuki groups

The order on $\mathcal{N}_0^{F_0}$ is $(u_1, 1) \leq (u_4, 1) \leq (u_5, 1)$. For the construction of the functions $\varphi_{u,\phi}$, we need to fix an extension of ϕ to $A_{\mathbf{G}_0}(u) \rtimes \langle F_0 \rangle$; we choose the trivial character for $A_{\mathbf{G}_0}(u_1) \rtimes \langle F_0 \rangle$ and for $A_{\mathbf{G}_0}(u_3) \rtimes \langle F_0 \rangle$, and the non-trivial character for $A_{\mathbf{G}_0}(u_4) \rtimes \langle F_0 \rangle$. We then give the resulting functions in Table 4. Recall that $|\mathbf{T}_1^{F_0}| = q^2 - 1$, $|\mathbf{T}_{w_a}^{F_0}| = q^2 - \sqrt{2}q + 1$ and $|\mathbf{T}_{w_a w_b w w_a}^{F_0}| = q^2 + \sqrt{2}q + 1$.

Theorem 3.1 When $\mathbf{G}_0^{F_0}$ is the Suzuki group of parameter q^2 , then the ma-

	u_1	u_4	ρ	$ ho^{-1}$
$\varphi_{u_1,1}$	1	0	0	0
$\varphi_{u_4,1}$	0	-1	0	0
$\varphi_{u_5,1}$	0	0	1	1

Table 4: Values of the functions $\varphi_{u,\phi}$ for the Suzuki group.

trix Ω defined in §3.3 is

$$\Omega = \begin{bmatrix} 1 & q^2 - 1 & 1 \\ q^2 - 1 & q^6 - q^2 & -q^6 + q^4 \\ 1 & -q^6 + q^4 & q^8 \end{bmatrix}$$

The matrices P and Λ resulting from Lusztig's algorithm described in §3.3 are

$$P = \begin{bmatrix} 1 & q^2 - 1 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} \quad and \quad \Lambda = \begin{bmatrix} 1 & 0 & 0 \\ 0 & q^6 - q^4 + q^2 - 1 & 0 \\ 0 & 0 & q^8 - q^6 + q^4 - q^2 \end{bmatrix}$$

Moreover the characteristic functions $X_{u,\phi}$ with $(u,\phi) \in \mathcal{N}_0^{F_0}$ computed using Relation (1) in §3.3 coincide with the uniform almost characters $R_{\tilde{\rho}_{u,\phi}}$ on the unipotent elements of $\mathbf{G}_0^{F_0}$.

3.5 The Ree groups of type G_2

Let \mathbf{G}_0 be a simple algebraic group of type G_2 defined over $\overline{\mathbb{F}}_3$; we set $q = 3^n \sqrt{3}$ for some non-negative integer n and $F_0 = \sigma \circ F_{3^n}$, where σ is described in [4, §12.4] and F_{3^n} is the standard Frobenius map over \mathbb{F}_{3^n} . The finite group $\mathbf{G}_0^{F_0}$ is the Ree group of type G_2 with parameter q^2 which has order $q^6(q^2-1)(q^6+1)$. The Weyl group of \mathbf{G}_0 is the dihedral group of order 12. We denote by w_a and w_b the reflections of W corresponding to the roots a and b respectively (here b is the long root). As before, we denote by $x_r(t)$ the Chevalley generators of \mathbf{G}_0 . The group W has 6 irreducible characters. We denote by $\mathbf{1}_W$ the trivial character and by ε the sign of W. There are two other linear characters; we denote by ϵ_a (resp. ϵ_b) the one which satisfies $\epsilon_a(w_a) = -1$ (resp. $\epsilon_b(w_b) = -1$). We denote by θ' and θ'' the two characters of degree 2 such that $\theta'(w_a w_b) = 1$.

In Table 5, we recall the generalized Springer correspondence for a simple group of type G_2 in characteristic 3; see [26]. To obtain the correspondence for θ' and θ'' with our notations, we use that $b_{\theta'} = 1$ and $b_{\theta''} = 2$, where b_{χ} denotes the smallest positive integer k such that χ occurs in the representation of W on the space of homogeneous polynomials of degree k on the natural complex reflection representation of W. The representatives of unipotent classes of \mathbf{G}_0 are $u_1 = 1$, $u_2 = x_{2a+b}(1)$, $u_3 = x_{3a+2b}(1)$, $u_4 = x_{2a+b}(1)x_{3a+2b}(1)$, $u_5 = x_{a+b}(1)x_{3a+b}(1)$ and $u_6 = x_a(1)x_b(1)$. The F₀-stable classes of \mathbf{G}_0 have representatives u_1, u_4, u_5 and u_6 . The sets $\mathrm{Cl}(u_5)^{F_0}$ and $\mathrm{Cl}(u_6)^{F_0}$ split into two and three classes of $\mathbf{G}_0^{F_0}$ respectively. The representatives are denoted by T, T^{-1} and Y, YT, YT^{-1} following the notation of [28].

3.5.1 Uniform almost characters

The F_0 -stable characters of W are 1_W , ε , θ' and θ'' . In Table 6, we give the values of an extension of one of their extensions on the coset WF_0 . To simplify the notation, we set $w_1 = w_a$, $w_2 = w_a w_b w_a$ and $w_3 = w_a w_b w_a w_b w_a$. In [10, Table 1], the decomposition of the uniform almost characters of the Ree groups

u	$A_{\mathbf{G}_0}(u)$	ϕ	$ ho_{u,\phi}$	d_u
u_1	1	1	ε	6
u_2	1	1	ϵ_a	3
u_3	1	1	ϵ_b	3
u_4	1	1	$\theta^{\prime\prime}$	2
u_5	$\mathbb{Z}/2\mathbb{Z}$	1	θ'	1
u_6	$\mathbb{Z}/3\mathbb{Z}$	1	1_W	0

Table 5: Generalized Springer correspondence for type G_2 .

	F_0	w_1F_0	w_2F_0	w_3F_0
$\widetilde{\varepsilon}$	1	-1	$^{-1}$	-1
$\widetilde{\theta}^{\prime\prime}$	0	1	-2	1
$\widetilde{\theta}'$	0	$-\sqrt{3}$	0	$\sqrt{3}$
$\widetilde{1}_W$	1	1	1	1

extensions	on	WF_0 .
	extensions	extensions on

are given. We use the notation of [28] for the unipotent characters of ${\bf G}_0^{F_0}.$ We have:

$$\begin{aligned} R_{\tilde{1}_W} &= 1_{\mathbf{G}_0^{F_0}}, \\ R_{\tilde{\varepsilon}} &= \mathrm{St}_{\mathbf{G}_0^{F_0}}, \\ R_{\tilde{\theta}'} &= \frac{1}{2\sqrt{3}} (\xi_5 + \xi_6 + \xi_7 + \xi_8 + 2\xi_9 + 2\xi_{10}), \\ R_{\tilde{\theta}''} &= \frac{1}{2} (\xi_5 - \xi_6 + \xi_7 - \xi_8). \end{aligned}$$

Using the character table of $\mathbf{G}_{0}^{F_{0}}$ given in [28], we deduce

Proposition 3.2 The values of the uniform almost characters on the unipotent elements of $\mathbf{G}_0^{F_0}$ are given in Table 7.

	u_1	u_4	T	T^{-1}	Y	YT	YT^{-1}
$R_{\widetilde{\varepsilon}}$	$ \begin{array}{c} $	0	0	0	0	0	0
$R_{\widetilde{\theta}^{\prime\prime}}$	$q^2(q^2-1)$	$-q^2$	0	0	0	0	0
$R_{\widetilde{\theta}'}$	$q(q^4 - 1)$	-q	-q	-q	0	0	0
$R_{\tilde{1}_W}$	1	1	1	1	1	1	1

Table 7: Values of the uniform almost characters of the Ree groups of type G_2 .

_	u_1	u_4	T	T^{-1}	Y	YT	YT^{-1}
$\varphi_{u_1,1}$	1	0	0	0	0	0	0
$\varphi_{u_4,1}$	0	1	0	0	0	0	0
$\varphi_{u_5,1}$	0	0	-1	-1	0	0	0
$\begin{array}{c} \varphi_{u_1,1} \\ \varphi_{u_4,1} \\ \varphi_{u_5,1} \\ \varphi_{u_6,1} \end{array}$	0	0	0	0	1	1	1

Table 8: Values of the functions $\varphi_{u,\phi}$ for the Ree groups of type G_2 .

3.5.2 Lusztig's algorithm for the Ree groups of type G_2

We first recall the orders of the finite tori of $\mathbf{G}_{0}^{F_{0}}$ corresponding to $w \in W$; we have $|\mathbf{T}_{1}^{F_{0}}| = q^{2}-1$, $|\mathbf{T}_{w_{1}}^{F_{0}}| = q^{2}-\sqrt{3}q+1$, $|\mathbf{T}_{w_{2}}^{F_{0}}| = q^{2}+1$ and $|\mathbf{T}_{w_{3}}^{F_{0}}| = q^{2}+\sqrt{3}q+1$. We choose the order $(u_{1},1) \leq (u_{4},1) \leq (u_{5},1) \leq (u_{6},1)$ on $\mathcal{N}_{0}^{F_{0}}$. In order to define the $\varphi_{u,\phi}$ we choose as extensions the trivial character of $A_{\mathbf{G}_{0}}(u_{1}) \rtimes \langle F_{0} \rangle$, of $A_{\mathbf{G}_{0}}(u_{4}) \rtimes \langle F_{0} \rangle$ and of $A_{\mathbf{G}_{0}}(u_{6}) \rtimes \langle F_{0} \rangle$. For $A_{\mathbf{G}_{0}}(u_{5}) \rtimes \langle F_{0} \rangle$, we choose the extension of the trivial character of $A_{\mathbf{G}_{0}}(u_{5}) \rtimes \langle F_{0} \rangle$. In Table 8, we give the values of the corresponding functions.

Theorem 3.2 When $\mathbf{G}_0^{F_0}$ is the Ree group of type G_2 with parameter q^2 , then the matrix Ω defined in §3.3 is

$$\Omega = \begin{bmatrix} 1 & 1-q^2 & q^4-1 & 1\\ 1-q^2 & q^8-q^6-q^4-q^2 & -q^8+q^4 & q^8-q^6\\ q^4-1 & -q^8+q^4 & q^{10}+q^8-q^6-q^4 & -q^{10}+q^6\\ 1 & q^8-q^6 & -q^{10}+q^6 & q^{12} \end{bmatrix}$$

The matrices P and Λ resulting from Lusztig's algorithm described in §3.3 are

$$P = \begin{bmatrix} 1 & 1 - q^2 & q^4 - 1 & 1 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad and \quad \Lambda = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & p_1(q) & 0 & 0 \\ 0 & 0 & q^2 p_1(q) & 0 \\ 0 & 0 & 0 & q^4 p_1(q) \end{bmatrix},$$

where $p_1(q) = (q^4 - 1)(q^4 - q^2 + 1)$. Moreover for every $(u, \phi) \in \mathcal{N}_0^{F_0}$ the resulting characteristic function $X_{u,\phi}$ coincides with the uniform almost character $R_{\tilde{\rho}_{u,\phi}}$ on the unipotent elements of $\mathbf{G}_0^{F_0}$.

3.6 The Ree groups of type F_4

Let \mathbf{G}_0 be a simple algebraic group of type F_4 defined over $\overline{\mathbb{F}}_2$; we set $q = 2^n \sqrt{2}$ for some non-negative integer n and $F_0 = \sigma \circ F_{2^n}$, where σ is described in [4, §12.3]. The finite group $\mathbf{G}_0^{F_0}$ is the Ree group of type F_4 with parameter q^2 which has order $q^{24}(q^2 - 1)(q^6 + 1)(q^8 - 1)(q^{12} + 1)$. The Weyl group of \mathbf{G}_0 has 1152 elements and has 25 irreducible characters. The character table of Wwas computed by Kondo in [11]; we use his notation.

u	$A_{\mathbf{G}_0}(u)$	ϕ	0	d_u						
	~		$ ho_{u,\phi}$		•	u	$A_{\mathbf{G}_0}(u)$	ϕ	$ ho_{u,\phi}$	d_u
x_0	1	1	$\chi_{1,4}$	24		x_{14}	1	1	$\chi_{9,3}$	6
x_1	1	1	$\chi_{2,4}$	16		x_{15}	1	1	$\chi_{6,1}$	6
x_2	1	1	$\chi_{2,2}$	16						
x_3	1	1	$\chi_{4,5}$	13		x_{16}	1	1	$\chi_{16,1}$	5
x_4	1	1	$\chi_{9,4}$	10		x_{17}	\mathfrak{S}_3	1	$\chi_{12,1}$	4
								θ	$\chi_{6,2}$	
x_5	$\mathbb{Z}/2\mathbb{Z}$	1	$\chi_{8,4}$	9		x_{20}	$\mathbb{Z}/2\mathbb{Z}$	1	$\chi_{8,3}$	3
		ϵ	$\chi_{1,2}$			x_{22}	$\mathbb{Z}/2\mathbb{Z}$	1	$\chi_{8,1}$	3
x_7	$\mathbb{Z}/2\mathbb{Z}$	1	$\chi_{8,2}$	9			,			
		ϵ	$\chi_{1,3}$			x_{24}	D_8	1	$\chi_{9,1}$	2
x_9	$\mathbb{Z}/2\mathbb{Z}$	1		8				ϵ'	$\chi_{2,1}$	
-			$\chi_{4,1}$					$\epsilon^{\prime\prime}$	$\chi_{2,3}$	
x_{11}	1	1	$\chi_{4,3}$	7		x_{29}	$\mathbb{Z}/2\mathbb{Z}$	1	$\chi_{4,2}$	1
x_{12}	1	1	$\chi_{4,4}$	7		x_{31}	$\mathbb{Z}/4\mathbb{Z}$	1		0
x_{13}	1	1	$\chi_{9,2}$	6		<i>w</i> 31	<i>ш</i> /4 <i>Ш</i>	1	$\chi_{1,1}$	0

Table 9: Springer correspondence for type F_4 .

The generalized Springer correspondence of a simple algebraic group of type F_4 in characteristic 2 is determined in [26, p.330]. Remark that the characters denoted by χ_4 , $\chi_{4,1}$, $\chi_{4,2}$, $\chi_{4,3}$, $\chi_{4,4}$, χ_{12} and χ_{16} in [26] are those denoted by $\chi_{4,1}, \chi_{4,2}, \chi_{4,3}, \chi_{4,4}, \chi_{4,5}, \chi_{12,1}$ and $\chi_{16,1}$ in [11] respectively. The 20 unipotent classes of \mathbf{G}_0 were classified by Shinoda and we use the notation of [21, Table I]. The representatives x_i are defined as products of Chevalley generators in [20, p.139]. We recall in Table 9 the Springer correspondence of \mathbf{G}_0 . We denote by ϵ the non-trivial character of $\mathbb{Z}/2\mathbb{Z}$, by θ the irreducible character of \mathfrak{S}_3 of degree 2, and by ϵ' and ϵ'' the two linear characters of D_8 distinct from the trivial and the sign characters. There are 10 unipotent classes of \mathbf{G}_0 which are F_0 -stable. They have as representatives the elements x_0, x_3, x_4, x_9 , x_{15} , x_{16} , x_{17} , x_{24} , x_{29} and x_{31} . In [21], Shinoda shows that F_0 acts trivially on $A_{\mathbf{G}_0}(x_9)$, $A_{\mathbf{G}_0}(x_{29})$, $A_{\mathbf{G}_0}(x_{17})$ and $A_{\mathbf{G}_0}(x_{31})$. The action of F_0 on $A_{\mathbf{G}_0}(x_{24})$ is not trivial; this group has three F_0 -classes. Note that $\epsilon'^{F_0} = \epsilon''$. This permits to parametrize the conjugacy classes of $\mathbf{G}_0^{F_0}$. We denote by u_i $(0 \le i \le 18)$ a system of representatives as in [21, Table 2].

3.6.1 Uniform almost characters

The group W has eleven F_0 -stable characters: $\chi_{1,1}, \chi_{1,4}, \chi_{4,1}, \chi_{4,2}, \chi_{4,5}, \chi_{6,1}, \chi_{6,2}, \chi_{9,1}, \chi_{9,4}, \chi_{12,1}$ and $\chi_{16,1}$. For each of these characters, we choose an extension to $W \rtimes \langle F_0 \rangle$. We give the values of these extensions on the coset WF_0 in Table 10. We take the elements w_i $(1 \leq i \leq 11)$ defined in [21, p.8] for a system of representatives of the F_0 -classes of W.

The unipotent characters of $\mathbf{G}_0^{F_0}$ were computed by Malle in [16]; in the following, we will use his notation. The decomposition of the uniform almost

	w_1F_0	w_2F_0	w_3F_0	w_4F_0	w_5F_0	w_6F_0	$w_7 F_0$	w_8F_0	w_9F_0	$w_{10}F_{0}$	$w_{11}F_0$
$\widetilde{\chi}_{1,4}$	1	-1	-1	$^{-1}$	1	1	1	1	1	1	1
$\widetilde{\chi}_{4,5}$	0	0	$-\sqrt{2}$	$\sqrt{2}$	0	$2\sqrt{2}$	$-2\sqrt{2}$	0	0	$\sqrt{2}$	$-\sqrt{2}$
$\widetilde{\chi}_{9,4}$	1	1	$^{-1}$	$^{-1}$	$^{-1}$	3	3	$^{-3}$	0	0	0
$\widetilde{\chi}_{4,1}$	2	0	0	0	2	2	2	2	$^{-1}$	-1	-1
$\widetilde{\chi}_{6,1}$	0	0	0	0	2	$^{-2}$	$^{-2}$	-4	$^{-1}$	1	1
$\widetilde{\chi}_{16,1}$	0	0	0	0	0	$4\sqrt{2}$	$-4\sqrt{2}$	0	0	$-\sqrt{2}$	$\sqrt{2}$
$\widetilde{\chi}_{6,2}$	$^{-2}$	0	0	0	0	4	4	2	$^{-1}$	1	1
$\widetilde{\chi}_{12,1}$	2	0	0	0	$^{-2}$	$^{-2}$	$^{-2}$	2	$^{-1}$	1	1
$\widetilde{\chi}_{9,1}$	1	$^{-1}$	1	1	$^{-1}$	3	3	$^{-3}$	0	0	0
$\widetilde{\chi}_{4,2}$	0	0	$-\sqrt{2}$	$\sqrt{2}$	0	$-2\sqrt{2}$	$2\sqrt{2}$	0	0	$-\sqrt{2}$	$\sqrt{2}$
$\widetilde{\chi}_{4,1}$	1	1	1	1	1	1	1	1	1	1	1

Table 10: Values of extensions on WF_0 .

characters of $\mathbf{G}_0^{F_0}$ in unipotent components is given in [10, Table 1]. When $\chi_{i,j}$ is an F_0 -stable irreducible character of W, we write $R_{i,j}$ for $R_{\tilde{\chi}_{i,j}}$ to simplify. We recall:

$$\begin{array}{rcl} R_{1,4} &=& \chi_4, \\ R_{4,5} &=& \frac{1}{\sqrt{2}}(\chi_7 + \chi_8), \\ R_{9,4} &=& \chi_3, \\ R_{4,1} &=& \frac{1}{4}(\chi_9 + \chi_{10} + 2\chi_{11} - \chi_{12} - \chi_{13} - 2\chi_{14} - \chi_{15} - \chi_{16} - \chi_{17} - \chi_{18}), \\ R_{6,1} &=& \frac{1}{3}(\chi_{12} + \chi_{13} - \chi_{14} + \chi_{19} + \chi_{20} - 2\chi_{21}), \\ R_{16,1} &=& \frac{1}{2\sqrt{2}}(\chi_9 - \chi_{10} - \chi_{12} + \chi_{13} + \chi_{15} + \chi_{16} - \chi_{17} - \chi_{18}), \\ R_{6,2} &=& \frac{1}{6}(-3\chi_9 - 3\chi_{10} - \chi_{12} - \chi_{13} - 2\chi_{14} + 2\chi_{19} + 2\chi_{20} + 2\chi_{21}), \\ R_{12,1} &=& \frac{1}{12}(3\chi_9 + 3\chi_{10} + 6\chi_{11} + \chi_{12} + \chi_{13} + 2\chi_{14} + 3\chi_{15} + 3\chi_{16} + 3\chi_{17} \\ &\quad + 3\chi_{18} + 4\chi_{19} + 4\chi_{20} + 4\chi_{21}), \\ R_{9,1} &=& \chi_2, \\ R_{4,2} &=& \frac{1}{\sqrt{2}}(\chi_5 + \chi_6), \\ R_{4,1} &=& \chi_1. \end{array}$$

Using the table in [16], we deduce

Proposition 3.3 The values on the unipotent elements of the uniform almost characters of Ree groups of type F_4 are given in Table 11.

3.6.2 Lusztig's algorithm for the Ree groups of type F_4

The order on $\mathcal{N}_0^{F_0}$ is such that

$$(1,4) \le (4,5) \le (9,4) \le (4,1) \le (6,1) \le (16,1) \le (6,2) \le (12,1) \le (9,1) \le (4,2) \le (1,1).$$

	u_0	u_1	u_2	u_3	u_4	u_5
$R_{1,4}$	q^{24}	0	0	0	0	0
$R_{4,5}$	$q^{23} - q^{19} + q^{17} - q^{13}$	$-q^{13}$	0	0	0	0
$R_{9,4}$	$q^{22} - q^{20} + q^{16} - q^{12} + q^{10}$	$-q^{12} + q^{10}$	q^{10}	0	0	0
$R_{4,1}$	$q^{16} + q^8$	q^8	q^8	q^8	q^8	0
$R_{6,1}$	$q^{18} - q^{16} - q^8 + q^6$	$-q^{8} + q^{6}$	$-q^{8} + q^{6}$	0	0	q^6
$R_{16,1}$	$-q^{19} + 2q^{15} - q^{13} - q^{11} + 2q^9 - q^5$	$-q^{11} + 2q^9 - q^5$	$q^{9} - q^{5}$	$q^7 - q^5$	$q^7 - q^5$	$-q^{5}$
$R_{6,2}$	$-q^{18} + q^{16} - 2q^{12} + q^8 - q^6$	$-q^{12} + q^8 - q^6$	$-q^{6}$	$-q^4$	$-q^4$	0
$R_{12,1}$	$q^{20} + q^4$	q^4	q^4	q^4	q^4	q^4
$R_{9,1}$	$q^{14} - q^{12} + q^8 - q^4 + q^2$	$q^8 - q^4 + q^2$	$-q^{4} + q^{2}$	$q^6 - q^4 + q^2$	$q^6 - q^4 + q^2$	$-q^{4} + q^{2}$
$R_{4,2}$	$q^{11} - q^7 + q^5 - q$	$-q^{7} + q^{5} - q$	$q^5 - q$	$q^3 - q$	$q^3 - q$	-q
$R_{4,1}$	1	1	1	1	1	1

u_6	u_7	u_8	u_9	u_{10}	u_{11}	u_{12}	u_{13}	u_{14}	u_{15}	u_{16}	u_{17}	u_{18}
$R_{1,4} = 0$	0	0	0	0	0	0	0	0	0	0	0	0
$R_{4,5} = 0$	0	0	0	0	0	0	0	0	0	0	0	0
$R_{9,4}$ 0	0	0	0	0	0	0	0	0	0	0	0	0
$R_{4,1} = 0$	0	0	0	0	0	0	0	0	0	0	0	0
$R_{6,1} = 0$	0	0	0	0	0	0	0	0	0	0	0	0
$R_{16,1}$ $-q^5$	0	0	0	0	0	0	0	0	0	0	0	0
$R_{6,2} - q^4$	$-2q^4$	0	q^4	0	0	0	0	0	0	0	0	0
$R_{12,1}$ q^4	q^4	q^4	q^4	0	0	0	0	0	0	0	0	0
$R_{9,1} - q^4$	$+q^2$ q^2	q^2	q^2	q^2	q^2	q^2	0	0	0	0	0	0
$R_{4,2} = q^3 -$	$q \qquad 2q^3 - q$	-q	$-q^3 - q$	-q	-q	-q	-q	-q	0	0	0	0
$R_{4,1}$ 1	1	1	1	1	1	1	1	1	1	1	1	1

Table 11: Values on unipotent elements of the uniform almost characters of Ree groups of type F_4 .

The orders of the finite tori corresponding to elements $w \in W$ are

w	$ \mathbf{T}_w^{F_0} $		$ \mathbf{T}_w^{F_0} $
w_1	$(q^{2} - 1)^{2}$ $q^{4} - 1$ $(q^{2} - 1)(q^{2} - \sqrt{2}q + 1)$ $(q^{2} - 1)(q^{2} + \sqrt{2}q + 1)$		
wo	$a^4 - 1$	w_7	$(q^2 + \sqrt{2}q + 1)^2$ $(q^2 + 1)^2$
~ <u>~</u>	$(a^2 + 1)(a^2 + \sqrt{2}a + 1)$	w_8	$(q^2+1)^2$
w_3	$(q - 1)(q - \sqrt{2}q + 1)$	w_9	$q^{4} - q^{2} + 1$ $q^{4} - \sqrt{2}q^{3} + q^{2} - \sqrt{2}q + 1$ $q^{4} + \sqrt{2}q^{3} + q^{2} + \sqrt{2}q + 1$
w_4	$(q^2 - 1)(q^2 + \sqrt{2q} + 1)$	w_{10}	$a^4 - \sqrt{2}a^3 + a^2 - \sqrt{2}a + 1$
w_5	$q^4 + 1$		$a^{4} + \sqrt{2}a^{3} + a^{2} + \sqrt{2}a + 1$
w_6	$q^4 + 1 (q^2 - \sqrt{2}q + 1)^2$	w_{11}	$q + \sqrt{2}q + q + \sqrt{2}q + 1$

Using Table 10, we can compute the matrix Ω defined in §3.3. We now give the selected extension of ϕ to $A_{\mathbf{G}_0^{F_0}}(u) \rtimes \langle F_0 \rangle$ for $(u, \phi) \in \mathcal{N}_0^{F_0}$. We choose the trivial character of $A_{\mathbf{G}_0^{F_0}}(u) \rtimes \langle F_0 \rangle$ for $(x_0, 1)$, $(x_4, 1)$, $(x_9, 1)$, $(x_{15}, 1)$, $(x_{17}, 1)$, $(x_{24}, 1)$ and $(x_{31}, 1)$. We choose the extension of the trivial character which is not the trivial character for $(x_3, 1)$, $(x_{16}, 1)$ and $(x_{29}, 1)$. The group $A_{\mathbf{G}_0}(x_{17})$ is isomorphic to \mathfrak{S}_3 and F_0 acts trivially on it. Thus $A_{\mathbf{G}_0}(x_{17}) \rtimes \langle F_0 \rangle$ is a direct product; we choose for $\tilde{\theta}$ the extension of θ such that $\tilde{\theta}(F_0) = -2$. We then can compute the corresponding functions $\varphi_{u,\phi}$.

Theorem 3.3 When $\mathbf{G}_0^{F_0}$ is the Ree group of type F_4 with parameter q^2 , then the matrix P resulting from Lusztig's algorithm is

	1	$p_1(q)$	$p_2(q)$	$p_3(q)$	$p_4(q)$	$p_5(q)$	$p_6(q)$	$p_7(q)$	$p_8(q)$	$p_1(q)$	1	
	0	1	$p_9(q)$	-1	$p_9(q)$	$p_{10}(q)$	$p_{11}(q)$	-1	$p_{12}(q)$	$p_{13}(q)$	-1	
	0	0	1	1	$-p_{9}(q)$	$p_{14}(q)$	$-q^2$	1	$-p_9(q)$	$p_{14}(q)$	1	
	0	0	0	1	0	$p_9(q)$	$^{-1}$	1	$p_{15}(q)$	$p_9(q)$	1	
	0	0	0	0	1	-1	0	1	$-p_{9}(q)$	$^{-1}$	1	
P =	0	0	0	0	0	1	1	-1	$p_9(q)$	$-p_{9}(q)$	-1	,
	0	0	0	0	0	0	1	0	0	$-q^2$	0	
	0	0	0	0	0	0	0	1	1	-1	1	
	0	0	0	0	0	0	0	0	1	-1	1	
	0	0	0	0	0	0	0	0	0	1	-1	
	0	0	0	0	0	0	0	0	0	0	1	

where

$$\begin{array}{rcl} p_1(q) &=& (q^2-1)(q^2+1)^2(q^4-q^+1)\\ p_2(q) &=& (q^8-q^4+1)(q^4-q^2+1)\\ p_3(q) &=& q^8+1\\ p_4(q) &=& (q^2-1)^2(q^4-q^3+q^2-q+1)(q^4+q^3+q^2+q+1)\\ p_5(q) &=& -(q^2-1)^2(q^2+1)^3(q^4-q^2+1)\\ p_6(q) &=& -q^2(q^{12}-q^{10}+2q^6-q^2+1)\\ p_7(q) &=& q^{16}+1\\ p_8(q) &=& (q^4-q^2+1)(q^8-q^4+1)\\ p_9(q) &=& q^2-1\\ p_{10}(q) &=& (q^2-1)(q^4-q^2-1)\\ p_{11}(q) &=& q^2(q^6-q^2+1)\\ p_{12}(q) &=& -q^6+q^2-1\\ p_{13}(q) &=& q^6-q^4+1\\ p_{14}(q) &=& q^4-q^2+1 \end{array}$$

The matrix Λ is the diagonal matrix with entries

$$\begin{bmatrix} 1, f_1(q), q^4 f_2(q), q^{10} f_1(q), q^8 f_3(q), q^{10} f_3(q), q^{12} f_3(q), q^{12} f_3(q), q^{16} f_3(q), q^{18} f_3(q), q^{20} f_3(q) \end{bmatrix}$$
where

$$\begin{array}{rcl} f_1(q) &=& (q^2-1)(q^2+1)^2(q^4+1)(q^4-q^2+1)(q^8-q^4+1) \\ f_2(q) &=& (q^4-1)(q^4+1)^2(q^4-q^2+1)(q^8-q^4+1) \\ f_3(q) &=& (q^2-1)^2(q^2+1)^2(q^4+1)^2(q^4-q^2+1)(q^8-q^4+1) \end{array}$$

Moreover, for every $(u, \phi) \in \mathcal{N}_0^{F_0}$, the characteristic function $X_{u,\phi}$ obtained using this algorithm coincides with the uniform almost character $R_{\tilde{\rho}_{u,\phi}}$ on the unipotent elements of $\mathbf{G}_0^{F_0}$.

4 The disconnected cases

We keep the same notation as in the preceding sections.

4.1 Uniform almost characters

Digne-Michel [8] and Malle [17] have generalized the Deligne-Lusztig theory to disconnected groups. We briefly recall the construction: let $\widetilde{\mathbf{T}}'$ be a rational "torus" of $\widetilde{\mathbf{G}}$ contained in a "Borel" $\widetilde{\mathbf{B}}'$. We then have (see [8, Proposition 1.5]) $\widetilde{\mathbf{B}}' = \mathbf{U}' \rtimes \widetilde{\mathbf{T}}'$, where \mathbf{U}' is the unipotent radical of $\widetilde{\mathbf{B}}'^\circ$. We introduce the variety $\mathbf{Y}_{\mathbf{U}'} = \{x \in \mathbf{G} \mid x^{-1}F(x) \in \mathbf{U}'\}$. The groups \mathbf{G}^F and $\widetilde{\mathbf{T}}'^F$ act on $\mathbf{Y}_{\mathbf{U}'}$ and these actions commute. This induces linear actions on $H^i_c(\mathbf{Y}_{\mathbf{U}'}, \overline{\mathbb{Q}}_\ell)$, where ℓ is a prime number distinct from the characteristic. Let $\theta \in \operatorname{Irr}(\widetilde{\mathbf{T}}'^F)$, we define the associated generalized Deligne-Lusztig characters by (see [8, §2])

$$R_{\widetilde{\mathbf{T}}'}^{\widetilde{\mathbf{G}}}(\theta)(g) = \frac{1}{|\widetilde{\mathbf{T}}'^{F}|} \sum_{t \in \widetilde{\mathbf{T}}'^{F}} \theta(t^{-1}) \operatorname{Tr}\left((g,t) | H_{c}^{*}(\mathbf{Y}_{\mathbf{U}'}, \overline{\mathbb{Q}}_{\ell})\right) \quad \text{for } g \in \widetilde{\mathbf{G}}.$$

We now give a parametrization (up to conjugacy) of the rational "tori" of $\widetilde{\mathbf{G}}$ containing τ . We recall that the "tori" of $\widetilde{\mathbf{G}}$ are conjugate in \mathbf{G} ; see [8]. Let $\widetilde{\mathbf{T}}'$

be a "torus" of $\widetilde{\mathbf{G}}$ containing τ , then $\widetilde{\mathbf{T}}'^{\circ} = \widetilde{\mathbf{T}}' \cap \mathbf{G}^{\circ}$ is a maximal torus of \mathbf{G} and we have $\widetilde{\mathbf{T}}' = \widetilde{\mathbf{T}}'^{\circ} \rtimes \langle \tau \rangle$. We have

Proposition 4.1 We fix a maximal rational torus \mathbf{T}_0 of \mathbf{G}_0 and we set $\mathbf{T} = \mathbf{T}_0 \times \mathbf{T}_0$ and $\widetilde{\mathbf{T}} = \mathbf{T} \rtimes \langle \tau \rangle$. The $\widetilde{\mathbf{G}}^F$ -classes of rational "tori" of $\widetilde{\mathbf{G}}$ are in correspondence with the $\mathbf{G}_0^{F_0}$ -classes of maximal rational tori of \mathbf{G}_0 . We fix a system W_{F_0} of representatives of F_0 -classes of W_0 . Let $w \in W_{F_0}$ and n_w be a corresponding element in $\mathbf{N}_{\mathbf{G}_0}(\mathbf{T}_0)$. We set

$$\mathbf{T}_{0,w} = x \mathbf{T}_0 x^{-1} \quad and \quad \widetilde{\mathbf{T}}_w = (x,x) \widetilde{\mathbf{T}}(x,x)^{-1},$$

where x is such that $n_w = x^{-1}F_0(x)$. Then the sets $\{\mathbf{T}_{0,w} \mid w \in W_{F_0}\}$ and $\{\widetilde{\mathbf{T}}_w \mid w \in W_{F_0}\}$ are respectively representative systems of $\mathbf{G}_0^{F_0}$ -classes of maximal rational tori of \mathbf{G}_0 and $\widetilde{\mathbf{G}}^F$ -classes of rational "tori" of $\widetilde{\mathbf{G}}$.

Proof — Write X for the set of "tori" of $\widetilde{\mathbf{G}}$. The connected group \mathbf{G} acts transitively on X. We denote by $F': X \to X, \widetilde{\mathbf{T}}' \mapsto F(\widetilde{\mathbf{T}}')$. We have $F'(x\widetilde{\mathbf{T}}') = F(x)F'(\widetilde{\mathbf{T}}')$ and $N_{\mathbf{G}^{\diamond}}(\mathbf{T}')$ is a closed subset. We can apply Theorem 2.1. The "torus" $\widetilde{\mathbf{T}} = (\mathbf{T}_0 \times \mathbf{T}_0) \rtimes \langle \tau \rangle$ is a rational "torus" of X. We have

$$N_{\mathbf{G}}(\widetilde{\mathbf{T}}) = N_{\mathbf{G}}(\mathbf{T}_0 \times \mathbf{T}_0) \cap N_{\mathbf{G}}(\tau) = \mu \left(N_{\mathbf{G}_0}(\mathbf{T}_0) \right),$$

it follows that μ is an isomorphism of algebraic groups between $N_{\mathbf{G}_0}(\mathbf{T}_0)$ and $N_{\mathbf{G}}(\mathbf{T})$. Since $C_{\mathbf{G}}(\mathbf{\widetilde{T}}) = \mu(\mathbf{T}_0)$, we deduce that

$$N_{\mathbf{G}}(\widetilde{\mathbf{T}})^{\circ} = C_{\mathbf{G}}(\widetilde{\mathbf{T}}).$$

Moreover, we have

$$N_{\mathbf{G}}(\mathbf{T})/C_{\mathbf{G}}(\mathbf{T}) \simeq \mu \left(N_{\mathbf{G}_0}(\mathbf{T}_0)/\mathbf{T}_0 \right),$$

where μ is the map $w\mathbf{T}_0 \mapsto \mu(w\mathbf{T}_0)$. We then have

$$\forall w \in \mathrm{N}_{\mathbf{G}_0}(\mathbf{T}_0) / \mathbf{T}_0, \quad F(\mu(w)) = \mu(F_0(w)).$$

Let $w \in N_{\mathbf{G}_0}(\mathbf{T}_0)/\mathbf{T}_0$ and n_w be a representative of w in $N_{\mathbf{G}_0}(\mathbf{T}_0)$. Let $x \in \mathbf{G}_0$ be such that $n_w = x^{-1}F_0(x)$; We have

$$(x,x)^{-1}F(x,x) = \left(x^{-1}F_0(x), x^{-1}F_0(x)\right) = (n_w, n_w) = \mu(n_w),$$

The result is then a consequence of Theorem 2.1.

Remark 4.1 We have to find an analogue of Proposition 1.40 of [8] in our special case (with explicit representatives for the classes of rational "tori"). As a consequence of Proposition 4.1, we obtain Proposition 1.38 of [8], which says that every $\widetilde{\mathbf{G}}^{F}$ -class of "torus" of $\widetilde{\mathbf{G}}$ has a representative that contains τ .

The Weyl group of \mathbf{G} is $W = N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$, where $\mathbf{T} = \mathbf{T}_0 \times \mathbf{T}_0$. Following Malle in [17], we give a definition of the uniform almost characters of $\widetilde{\mathbf{G}}^F$: let ρ be an irreducible character of W^{τ} and $\widetilde{\rho}$ be an extension of ρ to $\widetilde{W} = W^{\tau} \rtimes \langle F \rangle$. We define the uniform almost character corresponding to $\widetilde{\rho}$ as the restriction of the class function onto the coset $\mathbf{G}^F \tau$: (we adapt here the definition [17, Definition 2] given in the case where F acts trivially on the Weyl group):

$$R_{\tilde{\rho}} = \frac{1}{|W^{\tau}|} \sum_{w \in W^{\tau}} \tilde{\rho}(wF) R_{\tilde{\mathbf{T}}_w}^{\tilde{\mathbf{G}}}(1_{\tilde{\mathbf{T}}_w^F}).$$

Note that W^{τ} is isomorphic to W_0 and the operation of F on W^{τ} is the same as the one of F_0 on W_0 .

In order to compute the decomposition of characters $R_{\tilde{\mathbf{T}}_w}^{\tilde{\mathbf{G}}}(1_{\tilde{\mathbf{T}}_w}^F)$ in irreducible constituents, we recall Proposition 4.8 of [8]:

Proposition 4.2 Let $\widetilde{\mathbf{T}}'$ and $\widetilde{\mathbf{T}}''$ be two rational "tori" containing τ ; suppose that $\theta' \in \operatorname{Irr}\left(\widetilde{\mathbf{T}}'^F\right)$ and $\theta'' \in \operatorname{Irr}\left(\widetilde{\mathbf{T}}''^F\right)$ satisfy $\theta'_0 = \operatorname{Res}_{\widetilde{\mathbf{T}}'^\circ F}^{\widetilde{\mathbf{T}}'F}(\theta') \in \operatorname{Irr}\left(\widetilde{\mathbf{T}}'^{\circ F}\right)^{\tau}$ and $\theta''_0 = \operatorname{Res}_{\widetilde{\mathbf{T}}''^\circ F}^{\widetilde{\mathbf{T}}''}(\theta'') \in \operatorname{Irr}\left(\widetilde{\mathbf{T}}'^{\circ F}\right)^{\tau}$. We have

$$\langle R_{\widetilde{\mathbf{T}}'}^{\widetilde{\mathbf{G}}}(\theta), R_{\widetilde{\mathbf{T}}''}^{\widetilde{\mathbf{G}}}(\theta'') \rangle_{\tau} = 0,$$

if $(\widetilde{\mathbf{T}}'^{\circ}, \theta'_{0})$ and $(\widetilde{\mathbf{T}}''^{\circ}, \theta''_{0})$ are not $\left((\widetilde{\mathbf{G}}^{\tau})^{\circ}\right)^{F}$ -conjugate and

$$\left\langle R_{\widetilde{\mathbf{T}}'}^{\widetilde{\mathbf{G}}}(\theta'), R_{\mathbf{T}'}^{\widetilde{\mathbf{G}}}(\theta') \right\rangle_{\tau} = \frac{1}{\left| \left(\widetilde{\mathbf{T}}'^{\tau} \right)^{\circ F} \right|} \left| \left\{ n \in \mathrm{N}_{(\widetilde{\mathbf{G}}^{\tau})^{\circ F}}(\widetilde{\mathbf{T}}') \mid {}^{n}\theta'_{0} = \theta'_{0} \right\} \right|.$$

Here, $\langle \psi_1, \psi_2 \rangle_{\tau}$ denotes $\frac{1}{|\mathbf{G}^F|} \sum_{g \in \mathbf{G}^F} \psi_1(g\tau) \overline{\psi_2(g\tau)}$.

4.2 Generalized Springer correspondence

The generalized Springer correspondence is extended to disconnected groups in [15, II]. It is extended for the connected component $D = \mathbf{G}\tau$ of $\widetilde{\mathbf{G}}$: since τ is unipotent, we can associate to every pair (u, ϕ) of \mathcal{N}_D (*u* is a unipotent element lying in D and ϕ is an irreducible character of A(u) defined in §2.2) a 4-tuple $(\mathbf{L}, v, \psi, \rho)$ as in the connected case, where \mathbf{L} is a τ -stable Levi subgroup of \mathbf{G} , (v, ψ) is a cuspidal pair of $\mathcal{N}_{\mathbf{L}\tau}$ and ρ is an irreducible character of $\mathbf{N}_{\mathbf{G}\tau}(\mathbf{L})/\mathbf{L}$; see [15, II.8.8,II.11.10]. For the pairs corresponding to \mathbf{T} (denoted $\mathcal{N}_{D,0}$ as previously), the generalization to the disconnected case was independently introduced by Sorlin [24].

Proposition 4.3 Let \mathbf{G}_0 , \mathbf{G} and \mathbf{G} as in §2.1. Let u be a unipotent element of \mathbf{G}_0 and $(u, \phi) \in \mathcal{N}_0$ and let $\rho_{u,\phi} \in W_0$ be the Springer correspondent of (u, ϕ) .

Then $(f(u), \phi \circ \mu^{-1}) \in \mathcal{N}_{D,0}$ (where f and μ are the maps defined in Proposition 2.2) and the Springer correspondent corresponding to it is $\rho_{u,\phi}$ (viewed as a character of $N_{\mathbf{G}^{\tau}}(\mathbf{T})/\mathbf{T} \simeq W_0$).

Proof — This is a consequence of Proposition 2.2 and [19, Theorem 3.3], applied with σ being trivial.

Corollary 4.1 The Springer correspondence for the pairs lying in $\mathcal{N}_{D,0}$ for the disconnected groups of type B_2 and F_4 are given in Tables 1 and 9 respectively. In Table 1 (resp. Table 9), the representatives u_i (resp. x_i) are now the images under f defined in Proposition 2.2 of the corresponding elements of \mathbf{G}_0 .

Remark 4.2 In fact, for the disconnected cases of type B_2 and F_4 , the generalized Springer correspondence of the pairs lying in \mathcal{N}_D is in correspondence with the generalized correspondence of the pairs lying in \mathcal{N} . The generalized Springer correspondence for disconnected groups indeed satisfies the same properties as the generalized Springer correspondence for connected groups. These properties are recalled in [18, §2]. Moreover in [26], Spaltenstein proves that the generalized Springer correspondence for a simple group of type F_4 in characteristic 2 corresponding to the pairs which are not in \mathcal{N}_0 can be computed using only these properties.

4.3 The disconnected case of type B_2

We use the notation of [2]. Note that the symbol q in [2] is the symbol q^2 in this paper. The group \mathbf{G}^F has 4 unipotent characters $\mathbf{1}_{\mathbf{G}^F}$, θ_1 , θ_4 and θ_5 which extend to $\widetilde{\mathbf{G}}^F$. If θ is such a character, we denote by $\widetilde{\theta}$ the extension such that $\widetilde{\theta}(\tau) > 0$. The values of these extensions are given in [2, Table 9]. Since every class of $\widetilde{\mathbf{G}}$ lying in the coset $\mathbf{G}\tau$ is real, it follows from Remark 2.1 that the class of f(x) (where $x \in \mathbf{G}_0$ and f defined in Proposition 2.2) is the Shintani correspondent of x in $\mathbf{G}_0^{F_0^2} \rtimes \langle F_0 \rangle$ when we use the identifications of Proposition 2.1. This correspondence is explicitly described in [2, Proposition 4.1]. Following Proposition 4.1, the $\widetilde{\mathbf{G}}^F$ -classes of rational "tori" of $\widetilde{\mathbf{G}}$ are parametrized by the F_0 -classes of $W = W(\mathbf{G}_0)$ and a system of representatives is given in Table 2. For such a representative w, we denote by R_w the restriction of $R_{\widetilde{\mathbf{T}}_w}^{\widetilde{\mathbf{G}}}(\mathbf{1}_{\widetilde{\mathbf{T}}_w}^F)$ to $\mathbf{G}^F \tau$.

Proposition 4.4 We have on the coset $\mathbf{G}^F \tau$:

$$\begin{array}{rcl} R_1 & = & \mathbf{1}_{\widetilde{\mathbf{G}}} + \widetilde{\theta}_4 \\ R_{w_a} & = & \mathbf{1}_{\widetilde{\mathbf{G}}} - \widetilde{\theta}_1 - \widetilde{\theta}_5 - \widetilde{\theta}_4 \\ R_{w_a w_b w_a} & = & \mathbf{1}_{\widetilde{\mathbf{G}}} + \widetilde{\theta}_1 + \widetilde{\theta}_5 - \widetilde{\theta}_4 \end{array}$$

Proof — Using the proof of [2, Proposition 3.1], we deduce that $R_1 = 1_{\tilde{\mathbf{G}}} + \tilde{\theta}_4$ on $\mathbf{G}^F \tau$. Let $w \in W$; note that $\langle R_w, 1_{\tilde{\mathbf{G}}} \rangle_{\tau} = 1$, where \langle , \rangle_{τ} is the restricted scalar product introduced in Proposition 4.2. We indeed have

$$\langle R_{\widetilde{\mathbf{T}}_{w}}^{\widetilde{\mathbf{G}}}(1_{\widetilde{\mathbf{T}}_{w}^{F}}), 1_{\widetilde{\mathbf{G}}} \rangle_{\tau} = \langle 1_{\widetilde{\mathbf{T}}_{w}^{F}}, R_{\widetilde{\mathbf{T}}_{w}}^{\widetilde{\mathbf{G}}}(1_{\widetilde{\mathbf{T}}_{w}^{F}}) \rangle_{\mathbf{T}_{w}^{\circ F}\tau} = \langle 1_{\widetilde{\mathbf{T}}_{w}^{F}}, 1_{\widetilde{\mathbf{T}}_{w}^{F}} \rangle_{\mathbf{T}_{w}^{\circ F}\tau} = 1,$$

where ${}^*\!R^{\widetilde{\mathbf{G}}}_{\widetilde{\mathbf{T}}_w}$ is the adjoint of $R^{\widetilde{\mathbf{G}}}_{\widetilde{\mathbf{T}}_w}$. Using Proposition 4.2, we deduce that

We then deduce that

$$\langle \widetilde{\theta}_4, R_{w_a} \rangle_{\tau} = \langle \widetilde{\theta}_4, R_{w_a w_b w_a} \rangle_{\tau} = -\langle 1_{\widetilde{\mathbf{G}}}, R_{w_a} \rangle_{\tau} = -1.$$

Moreover, since the constituents of R_{w_a} and $R_{w_a w_b w_a}$ are extensions of unipotent characters of \mathbf{G}^F , we deduce that the other constituents of these characters are $\tilde{\theta}_1$ and $\tilde{\theta}_5$ (possibly with multiplicity 0). However, the τ -norm of these characters is 4. It then follows that $\tilde{\theta}_1$ and $\tilde{\theta}_5$ occur with multiplicity ± 1 . Now, using [8, Theorem 4.13], we have

$$R_{w_a}(\tau) = R_{\mathbf{T}_{0,w_a}}^{\mathbf{G}_0}(1) = -(q^2 - 1)(q^2 + \sqrt{2}q + 1),$$

and similarly $R_{w_a w_b w_a}(\tau) = -(q^2 - 1)(q^2 - \sqrt{2}q + 1).$ Since $\tilde{\theta}_1(\tau) = \tilde{\theta}_5(\tau) = \frac{1}{\sqrt{2}}q(q^2 - 1)$, the result holds.

Corollary 4.2 We choose the extensions of the F-stable characters of W^{τ} to $W^{\tau} \rtimes \langle F \rangle$ as in Table 2. We then have

$$\begin{array}{rcl} R_{\widetilde{1}} & = & 1_{\widetilde{\mathbf{G}}^{F}}, \\ R_{\widetilde{\varepsilon}} & = & \widetilde{\theta}_{4}, \\ R_{\widetilde{\chi}} & = & \frac{\sqrt{2}}{2} (\widetilde{\theta}_{1} + \widetilde{\theta}_{5}). \end{array}$$

Moreover, the values of these class functions on the unipotent elements lying in the coset $\mathbf{G}^F \tau$ are given in Table 3, where we replace u_1 , u_2 , ρ and ρ^{-1} by the elements denoted by $(1, \sigma)$, (x_{a+b}, σ) , (x_a, σ) and $(x_a x_{a+b}, \sigma)$ in [2] respectively.

Let $\mathcal{N}_{D,0}^F$ be the set of pairs of $\mathcal{N}_{D,0}$ which are F-stable. When $(u, \phi) \in \mathcal{N}_{D,0}^F$, we can associate to (u, ϕ) a class function $Y_{u,\phi}$ on the unipotent elements of $\mathbf{G}^F \tau$; see [15, IV.19.6, IV.19.8]. When u is a unipotent element of $\mathbf{G}^F \tau$, we denote by \mathcal{B}_u the variety of Borel subgroups of \mathbf{G} which are u-stable (for the conjugation in $\widetilde{\mathbf{G}}$). We now have:

• A Springer correspondence for the pairs lying in $\mathcal{N}_{D,0}^F$;

- A basis of the space of class functions defined on the unipotent elements of $\mathbf{G}^F \tau$.
- The integers d_u giving the dimensions of the varieties \mathcal{B}_u .

We can then formally apply Lusztig's Algorithm with these data as input. Remark that if u is a unipotent element of \mathbf{G}_0 , then the varieties \mathcal{B}_u and $\mathcal{B}_{f(u)}$ (with f defined in Proposition 2.2) are isomorphic. The isomorphism is given by

$$\mathbf{B}' \in \mathcal{B}_u \mapsto (\mathbf{B}', \mathbf{B}') \in \mathcal{B}_{f(u)}.$$

We have seen in Proposition 2.2 that f(u) lies in a rational unipotent class of $\mathbf{G}\tau$. Let u' be a rational element in the class of f(u) obtained by the process of the proof of Proposition 2.2. We have shown that the action of F on A(u') is the same as the one of F on A(f(u)). We thus obtain a bijection between the pairs $(u', \phi') \in \mathcal{N}_{D,0}^{F_0}$ and the pairs $(f(u), \phi \circ \mu^{-1})$, with $\phi \circ \mu^{-1}$ being an F-stable character of A(f(u)). Since the action of F on A(f(u)) is the same as the one of F_0 on $A_{\mathbf{G}_0}(u)$, it follows that $\mathcal{N}_0^{F_0}$ and $\mathcal{N}_{D,0}^F$ are in bijection. Using the proof of [15, IV, Lemma 9.7], we define the class functions $\varphi_{f(u),\phi}$ for $(u,\phi) \in \mathcal{N}_0^{F_0}$ by the class function on the set of unipotent elements of $\mathbf{G}^F \tau$ as follows: the conjugacy classes of $\widetilde{\mathbf{G}}^F$ lying in $\mathrm{Cl}(f(u)^F)$ are in bijection with the F-classes of A(f(u)) (see the proof of Proposition 2.2). If a is a representative of such a class, we denote by \widetilde{a} a representative of the corresponding class of $\widetilde{\mathbf{G}}^F$. Let $g \in \widetilde{\mathbf{G}}^F$, we set

$$\varphi_{f(u),\phi}(g) = \begin{cases} \widetilde{\phi}(aF) & \text{if } g \text{ is conjugate in } \widetilde{\mathbf{G}}^F \text{ to } \widetilde{a}, \\ 0 & \text{otherwise.} \end{cases}$$

Here $\tilde{\phi}$ denotes an extension of $\phi \circ \mu^{-1}$ to $A(f(u)) \rtimes \langle F \rangle$. We choose for $\tilde{\phi}$ the same extension as in Table 4. For the notation, we replace in Table 4 the functions $\varphi_{u,\phi}$ by $\varphi_{f(u),\phi}$ and the classes as in Corollary 4.2.

Proposition 4.5 The outputs obtained using Lusztig's algorithm with these data are precisely the uniform almost characters of $\mathbf{G}^F \tau$.

4.4 Conjecture for the disconnected case of type F_4

Since Lusztig's algorithm is valid for the disconnected case of type B_2 , we can suppose that this will also be the case for the disconnected type F_4 . We state a conjecture of the same type as Proposition 4.5.

Conjecture 4.1 Suppose \mathbf{G}_0 is a simple group of type F_4 and F_0 the generalized Frobenius map that defines a Ree group of type F_4 . Then the values of the uniform almost characters of the group $\mathbf{G}_0^{F_0^2} \rtimes \langle F_0 \rangle$ on the coset $\mathbf{G}_0^{F_0^2} F_0$ are given in Table 11, when we replace u_i by its Shintani correspondent $N_{F_0/F_0^2}(u_i)$.

Acknowledgment

I wish to thank heartily Cédric Bonnafé to orient me in this direction, and I wish to thank deeply Frank Lübeck and Gunter Malle for the many valuable discussions that they granted to me.

References

- C. BONNAFÉ, Sur les caractères des groupes réductifs finis à centre non connexe: applications aux groupes spéciaux linéaires et unitaires, to appear in Astérisque.
- [2] O. BRUNAT, The Shintani descents of Suzuki groups and their consequences, Journal of Algebra 303, 2006, 869–890.
- [3] O. BRUNAT, On the extension of $G_2(3^{2n+1})$ by the exceptional graph automorphism, submitted for publication.
- [4] R.W. CARTER, Simple groups of Lie Type, Wiley, New York, 1972.
- [5] R.W. CARTER, Finite Groups of Lie Type: Conjugacy classes and Complex Characters, Wiley, New York, 1985.
- [6] F. DIGNE, Descente de Shintani et restriction des scalaires, Journal of London Math. Society, vol. 59, No 3 (1999)
- [7] F. DIGNE AND J. MICHEL, Fonctions \mathcal{L} des variétés de Deligne-Lusztig et descente de Shintani, Bull. S.M.F., mémoires 20, 1985.
- [8] F. DIGNE AND J. MICHEL, Groupes réductifs non connexes, Ann. École Normale Supérieure, 4ième série, t. 27, 1994, 345–406.
- [9] THE GAP GROUP, Groups, Algorithms, and Programming, Version 4.4.7; http://www.gap-system.org, 2006.
- [10] M. GECK AND G. MALLE, Fourier transforms and Frobenius eigenvalues for finite Coxeter groups, Journal of Algebra 260, 2003. 162–193.
- [11] T. KONDO, The characters of the Weyl group of type F₄, J. Fac. Sci. Univ. Tokyo, 11, 1965, 145–153.
- [12] G. LUSZTIG, Characters of reductive groups over a finite field, Annals Math. Studies 107, Princeton University Press, 1984.
- [13] G. LUSZTIG, Intersection cohomology complexes on reductive group, Invent. Math., 75, 1984, 205–272.
- [14] G. LUSZTIG, Character sheaves, Adv. Math. 56 (1985), 193-237; II, Adv. Math. 57 (1985) 226-265; III, Adv. Math. 57 (1985), 266-315; IV, Adv. Math. 59 (1986), 1-63; V, Adv. Math. 61 (1986), 103-155.

- [15] G. LUSZTIG, Character sheaves on disconnected groups II, Represent. Theory 8 (2004), 72–124; IV, Represent. Theory 8 (2004), 145–178.
- [16] G. MALLE, Die unipotenten Charaktere von ${}^{2}F_{4}(q^{2})$, Comm. Algebra, 18, 1990, 2361–2381.
- [17] G. MALLE, Generalized Deligne-Lusztig Characters, Journal of Algebra 159, 1993, 64–97.
- [18] G. MALLE, Springer correspondence for disconnected exceptional groups, Bull. London Math. Soc. 37, 2005, 391–398.
- [19] G. MALLE AND K. SORLIN, Springer correspondence for disconnected groups, Math. Z. 246, 2004, 291–319.
- [20] K. SHINODA, The conjugacy classes of Chevalley groups of type (F_4) over finite fields of characteristic 2, J. Fac. Sci. Univ. Tokyo, 21, 1974, 133-159.
- [21] K. SHINODA, The conjugacy classes of the finite Ree groups of type (F₄), J. Fac. Sci. Univ. Tokyo, 22, 1975, 1–15.
- [22] T. SHOJI, Geometry of orbits and Springer correspondence, Soc. Math. de France, Astérisque 168, 1988, 61–140.
- [23] T. SHOJI, Character sheaves and almost characters of reductive groups, Adv Math. 111 (1995), 244–313;
- [24] K. SORLIN, Springer correspondence in non connected reductive groups, J. Reine Angew. math. 508, 2004, 197–234.
- [25] N. SPALTENSTEIN, Classes unipotentes et sous-groupes de Borel, Lecture Notes in Math. 946, Springer, Berlin Heidelberg New York, 1982
- [26] N. SPALTENSTEIN, On the generalized Springer correspondence for exceptional groups, Algebraic groups and related topics, Adv. Stud. Pure Math. 6 (kinokuniya and North-Holland, Tokyo and Amsterdam, 1985), 317–338.
- [27] M. SUZUKI, On a class of doubly transitive groups. I, Annals of Math. 75, 1962, 105–145.
- [28] H.N. WARD, On Ree's series of simple groups, Trans. Amer. Math. Soc., 121, 1966, 62–89.