Computing the distribution of the maximum in balls-and-boxes problems, with application to clusters of disease cases

Warren J. Ewens Department of Biology, University of Pennsylvania Philadelphia, PA 19104-6018

<wewens@sas.upenn.edu>

Herbert S. Wilf Department of Mathematics, University of Pennsylvania Philadelphia, PA 19104-6395

<wilf@math.upenn.edu>

September 30, 2018

Abstract

We present a rapid method for the exact calculation of the cumulative distribution function of the maximum of multinomially distributed random variables. The method runs in time O(mn), where m is the desired maximum and n is the number of variables. We apply the method to the analysis of two situations where an apparent clustering of cases of a disease in some locality has raised the possibility that the disease might be communicable, and this possibility has been discussed in the recent literature. We conclude that one of these clusters may be explained on purely random grounds, whereas the other may not.

1 Introduction

It happens, from time to time, that cases of a disease will cluster both geographically and in time, in a manner which seems not to be random, and which invites further epidemiological study regarding communicability of the disease.

Of course mathematics alone cannot answer serious questions of public health, but it can provide guidelines about what sort of clustering should be regarded as unusual, and what sort is to be expected. In particular, the calculation of a P-value is required for an objective assessment of any observed event. In this paper we provide a rapid and exact P-value calculation for the standard "balls-in-boxes" model appropriate to the disease clustering situation.

2 The model

Suppose that during a certain time period, a number r of cases of some disease arise randomly in some large population, such as that of the the U.S. Let N be size of that population and N_0 be the population of the community in which the seemingly large number of cases has occurred.

We think of the entire country as consisting of $n = N/N_0$ identical communities, or cells, each containing N_0 people, and we ask:

If r cases occur randomly in the populations of n communities of the same size, what is the probability that no community gets more than m cases of the disease?

The standard calculation required to answer this question involves the "balls-in boxes" model, discussed below. If, for example, it turns out that it is extremely likely that *some* community of equivalent size to that where the seemingly large number of cases occurred, purely by chance, we could conclude that the observed cluster would not be a cause for further investigation or suspicion of communicability. Likewise, if it turns out that it is extremely unlikely that, by chance, *any* community of the size of that of interest would have the observed number of cases of the disease, then support would be given to the possibility of a public health hazard.

3 The mathematics

Mathematically speaking, we have r "balls" (the disease cases) being dropped randomly into n labeled "boxes" (the communities). The relevant calculation thus concerns the Pvalue associated with the box (or boxes) having the largest number of balls in it. It is well known that the distribution function of the maximum of a number of random variables changes sharply near the mean of the maximum, so that an exact rather than an approximate calculation is needed to find this P-value. We provide this exact calculation in this paper.

The P-value associated with an observed value m of cases of the disease in the community of interest is the probability that the maximum number of balls in any box in m or more.

We find this probability by first finding the probability that no box contains more than m balls. Denote this probability by P(r, n, m).

Now, the probability that there are r_1 balls in box 1, and r_2 in box 2, and ..., and r_n in box n, is given by the well known multinomial distribution,

$$\Pr(r_1, r_2, \dots, r_n) = \frac{1}{n^r} \frac{r!}{r_1! r_2! \dots r_n!}.$$
 (1)

The probability that no box contains more than m balls (i.e., the cumulative distribution function of the maximum of the r_i , evaluated at m) is

$$P(r, n, m) \stackrel{\text{def}}{=} \Pr(\text{all } r_i \text{ are } \le m) = \sum_{\substack{0 \le r_1, r_2, \dots, r_n \le m \\ r_1 + r_2 + \dots + r_n = r}} \frac{1}{n^r} \frac{r!}{r_1! r_2! \dots r_n!}.$$
 (2)

4 The computation

At first sight the expression (2) seems appallingly complicated for exact computation, if r and n are large. Various approximations, such as the Poisson approximation, have been used by researchers in order to avoid the apparently tedious computation in (2).

However the exact calculation can be completely tamed by two steps. First we introduce the function

$$e_m(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \ldots + \frac{x^m}{m!},$$

which is simply the *m*th section of the exponential series. Then P(r, n, m) is $r!/n^r$ times the coefficient of x^r in the series $e_m(x)^n$. The question of computing a particular coefficient of a high power of a given power series is a well studied problem in computer science, and the following solution, which makes the computation quite rapid and easy to program, is taken from [5] (chap. 21).

Let $f(x) = \sum_j a_j x^j$ be a given power series and let $h(x) = f(x)^n$. The question is, if $h(x) = \sum_j h_j x^j$, how can we economically compute the h_j 's from the given a_j 's? We begin by taking logarithms of the equation $h = f^n$, to get $\log h(x) = n \log f(x)$. Now differentiate both sides with respect to x to obtain h'/h = nf'/f, and cross multiply to eliminate fractions, yielding fh' = nhf'. Next insert the power series expansions of the various functions into this equation, and multiply both sides by x, for cosmetic reasons, to get

$$\left(\sum_{j} a_{j} x^{j}\right) \left(\sum_{\ell} \ell h_{\ell} x^{\ell}\right) = n \left(\sum_{j} h_{j} x^{j}\right) \left(\sum_{\ell} \ell a_{\ell} x^{\ell}\right).$$

Finally, equate the coefficients of a given power of x, say x^s , on both sides of the last equation, which gives,

$$\sum_{\ell=0}^{s} \ell h_{\ell} a_{s-\ell} = n \sum_{\ell=0}^{s} \ell a_{\ell} h_{s-\ell}.$$

This is a recurrence relation. We can use it to compute the unknown h_j 's successively, in the order h_0, h_1, h_2, \ldots To make this explicit, we can rewrite the above in the form

$$h_s = \frac{1}{sa_0} \sum_{\ell=1}^{s} ((n+1)\ell - s)a_\ell h_{s-\ell}. \qquad (s = 1, 2, 3, \ldots)$$
(3)

In this form it is clear that each h_s is determined from $h_0, h_1, \ldots, h_{s-1}$.

In the particular case at hand, of powers of the truncated exponential series $e_k(x)$, we have $a_j = 1/j!$, for $0 \le j \le m$, and $a_j = 0$ for all other values of j. The recurrence takes the form

$$h_s = \frac{1}{s} \sum_{\ell=1}^{\min(s,m)} ((n+1)\ell - s) \frac{h_{s-\ell}}{\ell!}. \qquad (s = 1, 2, 3, \ldots)$$
(4)

We summarize the calculation procedure as follows. To compute P(r, n, m) as defined by eq. (2) above,

- Take $h_0 = 1$ and successively compute h_1, h_2, \ldots, h_r from (4).
- Then $P(r, n, m) = r!h_r/n^r$.

A remarkable feature of this algorithm is that the computation of each h_s requires the knowledge of only m earlier values, so the entire computation can be done with just m units of array storage. For example, it can find the probability that the maximum is ≤ 8 , for 15000 balls in 10000 boxes using only 8 array storage locations. In summary, it runs in time which is O(mn) and uses only O(m) storage.

We remark that as we have presented it this method works only for the situation in which the cells have equal probabilities. It can be extended, at only a small extra cost, to the case of unequal probabilities, which may be useful for power calculations.

5 Some related work

The problem of finding the distribution of the maximum occupancy in a balls-and-cells problem is very old. Already in Barton and David [1] one finds the first observation above, namely that the desired probability is a certain coefficient in a power of a given power series. In [2] this observation of Barton and David is cited, and is said to be "not in a form convenient for computing," which is true absent our second step above, in eq. (3) of vastly accelerating the computation of the high power of the given series.

Freeman's algorithm in [2] sought to economize the computation by grouping together vectors of occupancy numbers which, as unordered multisets, were the same. Hence he listed partitions with given largest size part, and counted the occupancies of that subset of all partitions. This is a large amount more labor than our method above, which requires computing time roughly proportional to the square of the number of "balls," whereas earlier methods required exponential time.

Likewise the recurrence (3) for computing powers of power series has a long history. Although we have followed [5] in our presentation, the recurrence method was certainly not invented by them, and is described in several earlier works. Nonetheless, the concatenation of the two methods in connnection with finding the distribution of the maximum cell occupancy seems to be new.

6 Leukemia: two examples

Example 1. We consider first the much discussed case (see [3] and [4]) of childhood leukemia in Niles, IL in the five year period 1956–1960. Heath [3] gives a total of eight cases in this town during this period, as compared to an expected number of 1.6. In 1960 the population of Niles was about 20,000 people. The total population of the U.S. in 1960 was approximately 180,000,000 people. Therefore the U.S. population in 1960 can be thought of as consisting of 9,000 cells, the population of each being 20,000 people. An expected number of 1.6 in Niles would then correspond to a total of about 14,400 cases in the U.S. in the five year period studied.

Using the formula above, we therefore computed the exact probability that if 14,400 balls are distributed randomly into 9000 cells, then no cell will get more than m balls, for each m = 6, ..., 12, and in particular for m = 8. We also computed the *P*-value for each of these values of m using the fact that the *P*-values corresponding to an observed maximum of mis given by 1 - P(14400, 9000, m - 1).

For comparison, we ran a Monte Carlo computer experiment in which we repeated 1000 times the operation of distributing 14400 balls randomly into 9000 cells, and recorded the frequencies of the maximum occupancy numbers, thus giving an empirical distribution function for m. (1000 replications are needed to give an estimate of the P-value for m = 8 that is accurate to within ± 0.01 with probability 0.95.) The results of the this simulation, and the exact P(14400, 9000, m) computations are shown below, together with the exact P-values.

m	P(14400, 9000, m)	Monte Carlo	P-value
6	0.000005	0.000	1.000000
7	0.095395	0.096	0.999995
8	0.664954	0.678	0.904605
9	0.937864	0.944	0.335046
10	0.990843	0.993	0.062136
11	0.998788	0.998	0.009157
12	0.999852	0.999	0.001212

The computation of the above table of exact values, on a PC running the computer algebra system Maple,¹ required less than five seconds. The Monte Carlo computation required about thirty minutes. In both the Monte Carlo simulation and the exact calculations we observe the expected rapid change of P-values as m increases, emphasizing the need for exact P-value calculations as discussed above.

We conclude from the above that the probability that some cell of population 20,000 would have gotten 8 or more cases in the five year period studied is about 90 percent. Thus the Niles data do not appear, so far as formal *P*-value calculations are concerned, to show a significant cluster of cases of childhood leukemia.

Example 2. Twelve cases of acute lymphocytic leukemia were observed [6] in Churchill County, NV, among persons who had been residents of the county at the time of diagnosis, in the three year period 1999-2001. Concern was expressed that this was due to the exposure to some agent associated with a nearby naval air station. At that time the county had a population of approximately 24,000. The entire U.S. had a population of approximately 288,000,000, equivalent to 12,000 units, or cells, each of the size of Churchill County. The State Epidemiologist, Dr. Randall Todd, estimated that, based on its population, about one case would be expected in Churchill County every five years. If we use that estimate, the incidence in the U.S. as a whole would be 12000 cases per five years, or 8000 cases per three year period.

In this case we need the distribution function of the maximum number of balls in any cell if 8000 balls are thrown at random into 12,000 cells. The results are as shown below.

¹Program available on request

m	P(8000, 12000, m)	P-value
4	0.000472	1.000000
5	0.436361	0.999528
6	0.925122	0.563639
$\overline{7}$	0.993604	0.074878
8	0.999528	0.006396

Clearly the observed incidence of twelve cases in Churchill county cannot reasonably be ascribed to chance, and further epidemiological investigation is warranted.

7 Further comments on the *P*-value

The *P*-value corresponding to any value of m in the balls-in-boxes case can in principle be calculated exactly using standard inclusion/exclusion formulæ. In practice this seems extremely difficult, because the alternating signs can cause catastrophic loss of significant digits. A Poisson approximation is also possible but may be inaccurate, particularly around the tails of the distribution. Our exact method, described in eq. (4) above, is fast and does not suffer from any of those problems.

A further comment about *P*-values is more wide-ranging. Many diseases might come to our attention because of an apparent clustering in some location in some time period. Also, many different time periods might be potentially observed. An overall *P*-value calculation, taking these matters into consideration, would be desirable, but in practice would probably be impossibly difficult, since no precise value can be attached to "the number of diseases that might come to our attention" or, possibly, to the number of time periods that we might have considered.

8 A disclaimer

Mathematics cannot prove or disprove the communicability of a disease process. It can only help to define the word "unusual." The benchmark given above seems like an appropriate one to use when investigating an outbreak which is localized spatially, temporally, or both. By this benchmark, the clustering of leukemia cases in Niles, Illinois, between 1956 and 1960 was not unusual. In fact some collection of that number of cases in some community the size of Niles, in a five year period of keeping records, was to be expected with high probability. On the other hand, the Churchill County data seem extremely significant.

References

- [1] David, F.N., and Barton, D.E., Combinatorial Chance, Griffin, London, 1962.
- [2] Freeman, P.R., Algorithm AS 145: Exact distribution of the largest multinomial frequency, Applied Statistics, 28, No. 3 (1979), 333–336.
- [3] Heath, Clark W., Jr., Community Clusters of Childhood Leukemia and Lymphoma; Evidence of Infection?, American Journal of Epidemiology, 162, no. 9 (2005), 1–6.
- [4] Heath, Clark W., Jr., Heath responds to "Do leukemia/lymphoma cases have an infectious cause?," American Journal of Epidemiology, 162, no. 9 (2005), 825–826.
- [5] Nijenhuis, Albert, and Wilf, Herbert S., Combinatorial Algorithms, second ed., Academic Press, 1978.
- [6] Nevada State Statistics Page, online at <http://health2k.state.nv.us/healthofficer/leukemia/statistics.htm>.
- [7] Churchill County (Fallon) Childhood Leukemia Update, Nevada State Health Division, <http://health2k.state.nv.us/healthofficer/leukemia/fallon.htm>.