On Regular Locally Scalar Representations of Graph \widetilde{D}_4 in Hilbert Spaces

S. A. Kruglyak [†], L. A. Nazarova, A. V. Roiter.

 † Institute of Mathematics of National Academy of Sciences of Ukraine, Tereshchenkovska str., 3, Kiev, Ukraine, ind. 01601


E-mail: red@imath.kiev.ua

Representations of quivers corresponding to extended Dynkin graphs are described up to equivalence in [1]. Locally scalar representations of graphs in the category of Hilbert spaces were introduced in [2], and such representations are naturally classified up to unitary equivalense.

Representations of *-algebras generated by linearly related orthogonal projections are studied in [3–10] and others. The connection between locally scalar representations of several graphs (trees, which include also Dynkin graphs) and representations of such *-algebras is stated in [11], and we further use this connection.

The present paper is dedicated to the classification of indecomposable regular (see [12]) locally scalar representations of the graph \widetilde{D}_4 (for \widetilde{D}_4 those are indecomposable locally scalar representations in the dimension (2; 1, 1, 1, 1)). The answer obtained for the corresponding *-algebra in [5,6], in our opinion, cannot be satisfactory and definitive. We will obtain explicit formulas, expressing matrix elements of a representation by a character (see [2]) of a locally scalar representation and two "free" real parameters.

1. Let \mathcal{H} be the category of finite-dimensional Hilbert spaces, and \widetilde{D}_4 be an extended Dynkin graph



Remind (see [2]) that a representation Π of the graph \widetilde{D}_4 associates a space $H_i \in \mathcal{H}$ to each vertex i ($i = \overline{0,4}$), and a pair of interadjoint linear operators $\Pi(\gamma_i) = \{\Gamma_{i0}; \Gamma_{0i}\}$ to each edge γ_i , where $\Gamma_{0i}: H_i \to H_0$, $\Gamma_{i0} = \Gamma_{0i}^*$.

A morphism $C:\Pi\to\widetilde{\Pi}$ is a family $C=\{C_i\}_{i=\overline{0,4}}$ of operators $C_i:\Pi(i)\to\widetilde{\Pi}(i)$ such that

the diagrams

$$H_{i} \xrightarrow{\Gamma_{ji}} H_{j}$$

$$C_{i} \downarrow \qquad C_{j} \downarrow$$

$$\widetilde{H}_{i} \xrightarrow{\widetilde{\Gamma}_{ji}} \widetilde{H}_{j}$$

are commutative, i. e. $C_j\Gamma_{ji}=\widetilde{\Gamma}_{ji}C_i$.

Let M_i be the set of vertices connected with a vertex i by an edge, $A_i = \sum_{j \in M_i} \Gamma_{ij} \Gamma_{ji}$. Representation Π is called locally scalar [2] if all operators A_i are scalar; $A_i = \alpha_i I_{H_i}$, where I_{H_i} is identity operator in a space H_i . Since A_i is a positive operator, $\alpha_i \geq 0$. A vector $\{\dim \Pi(i)\}$ is a dimension of a finite-dimensional representation Π_i ; if $A_i = f(i)I_{H_i}$ then $\{f(i)\}$ is called a character of locally scalar representation Π .

Further we will denote as Rep (\widetilde{D}_4, f) the category of finite-dimensional locally scalar representations of the graph \widetilde{D}_4 in \mathcal{H} with given character f.

We will assume that $\alpha_i = f(i) > 0$, $i = \overline{0,4}$ and the character is normalized: $f(0) = \alpha_0 = 1$. On the other hand, consider the following *-algebra over the field \mathbb{C} :

$$\mathcal{P}_{4,f} = \mathbb{C}\langle p_1, p_2, p_3, p_4 \, | \, p_i = p_i^* = p_i^2, \, \sum_{i=1}^4 \alpha_i p_i = e \rangle,$$

where $\alpha_i = f(i)$, e is the identity of the algebra, and the category Rep $\mathcal{P}_{4,f}$ of finite-dimensional *-representation of the algebra $\mathcal{P}_{4,f}$.

Let $\Pi \in \text{Rep}(\widetilde{D}_4, f)$, $\Pi(i) = H_i$, $\pi(\gamma_i) = \{\Gamma_{i,0}; \Gamma_{0,i}\}$. Let us construct a representation π of the algebra $\mathcal{P}_{4,f}$ by the following way: $\pi(p_i) = \frac{1}{\alpha_i}\Gamma_{0,i} \cdot \Gamma_{i,0} = P_i$. If $C: \Pi \to \widetilde{\Pi}$ is a morphism in Rep (\widetilde{D}_4, f) then $C_0: \pi \to \widetilde{\pi}$ is a morphism in the category Rep $\mathcal{P}_{4,f}$ (C_0 is the operator interlacing representations π and $\widetilde{\pi}$). Define a functor

$$\Phi: \operatorname{Rep}(\widetilde{D}_4, f) \to \operatorname{Rep} \mathcal{P}_{4,f}$$

putting $\Phi(\Pi) = \pi$, $\Phi(C) = C_0$. Clearly, the functor Φ is the equivalence of categories.

Let $\pi \in \text{Rep } \mathcal{P}_{4,f}$ be a representation in the space H_0 . Set $H_i = \text{Im } P_i$, $i = \overline{1,4}$; $\Gamma_{0,i} : H_i \to H_0$ is the natural injection of the space H_i into H_0 , then, putting $\Pi(\gamma_i) = \{\Gamma_{0,i}; \Gamma_{0,i}^*\}$, we obtain a representation from $\text{Rep } (\widetilde{D}_4, f)$. If $C_0 : \pi \to \widetilde{\pi}$ set $C_i = C_0|_{H_i}$. If $\Phi^{(-1)}(\pi) = \Pi$, $\Phi^{(-1)}(C_0) = \{C_i\}_{i=\overline{0,4}}$ then $\Phi\Phi^{(-1)} \sim I_{\text{Rep } \mathcal{P}_{4,f}}$, $\Phi^{(-1)}\Phi \sim I_{\text{Rep } (\widetilde{D}_4,f)}$.

2. Consider representations of the *-algebra $\mathcal{P}_{4,f}$ for

$$0 < \alpha_1 \le \alpha_2 \le \alpha_3 \le \alpha_4 < 1, \sum_{i=1}^4 \alpha_i = 2.$$

(in the other cases representations of the *-algebras $\mathcal{P}_{4,f}$ are reduced to the simplest representations by the Coxeter functors [13]).

Let us make the substitution of generators in the algebra $\mathcal{P}_{4,f}$ [5]:

$$x = \alpha_2 p_2 + \alpha_3 p_3 - \frac{1}{2} \beta_1 e, \qquad \beta_1 = (2 - \alpha_1 + \alpha_2 + \alpha_3 - \alpha_4)/2,$$

$$y = \alpha_1 p_1 + \alpha_3 p_3 - \frac{1}{2} \beta_2 e, \text{ where } \beta_2 = (2 + \alpha_1 - \alpha_2 + \alpha_3 - \alpha_4)/2,$$

$$z = \alpha_1 p_1 + \alpha_2 p_2 - \frac{1}{2} \beta_3 e, \qquad \beta_3 = (2 + \alpha_1 + \alpha_2 - \alpha_3 - \alpha_4)/2.$$
(1)

Denote also

$$\gamma_1 = (\alpha_1^2 - \alpha_2^2 - \alpha_3^2 + \alpha_4^2)/4,$$

$$\gamma_2 = (-\alpha_1^2 + \alpha_2^2 - \alpha_3^2 + \alpha_4^2)/4,$$

$$\gamma_3 = (-\alpha_1^2 - \alpha_2^2 + \alpha_2^2 + \alpha_4^2)/4.$$

It is easy to check that $\gamma_1 \leq \gamma_2 \leq \gamma_3$ and $0 \leq \gamma_2$.

The new generators x, y, z satisfy the system of relations

$$\{y, z\} = \gamma_1 e,
\{z, x\} = \gamma_2 e, \quad (x + y + z)^2 = \alpha_4^2 e.
\{x, y\} = \gamma_3 e,$$
(2)

The equalities (1) imply

$$p_{1} = \frac{-x+y+z}{2\alpha_{1}} + \frac{1}{2}e,$$

$$p_{2} = \frac{x-y+z}{2\alpha_{2}} + \frac{1}{2}e,$$

$$p_{3} = \frac{x+y-z}{2\alpha_{3}} + \frac{1}{2}e,$$

$$p_{4} = \frac{-x-y-z}{2\alpha_{4}} + \frac{1}{2}e.$$
(3)

3. Let $\gamma_3 = 0$, then $0 \le \gamma_2 \le \gamma_3$ implies $\gamma_2 = 0$, hence $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \frac{1}{2}$ (this case is considered in [4]), and so $\gamma_1 = 0$.

In this case the system of relation (2) has a form

$$\{y, z\} = 0,$$

 $\{z, x\} = 0,$ $x^2 + y^2 + z^2 = \frac{1}{4}e.$ (4)
 $\{x, y\} = 0,$

Let π be an indecomposable two-dimensional representation of the algebra $\mathcal{P}_{4,f}$ and $\pi(x) = X$, $\pi(y) = Y$, $\pi(z) = Z$.

a) Let Z=0. The matrix X can be diagonalized as a matrix of self-adjoint operator. The relations of anticommutation imply that the triple X, Y, Z=0 is indecomposable only in the case when the diagonalized matrix X equals $\begin{bmatrix} -\lambda & 0 \\ 0 & \lambda \end{bmatrix}$. Then $Y=\begin{bmatrix} 0 & y_{12} \\ \overline{y}_{12} & 0 \end{bmatrix}, y_{12} \neq 0$,

and the element y_{12} can be made positive by the admissible tranformations. Therefore, we obtain the case

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \mu \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad Z = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}; \quad \lambda > 0, \, \mu > 0.$$

b) Let $Z \neq 0$ and X = 0. Then the matrix Y can be diagonalized: $Y = \begin{bmatrix} -\mu & 0 \\ 0 & \mu \end{bmatrix}$, $\mu > 0$ (in the other cases the triple of matrices turn out to be decomposable). Then from $\{y,z\} = 0$ we obtain $Z = \begin{bmatrix} 0 & z_{12} \\ \overline{z}_{12} & 0 \end{bmatrix}$ and one can reduce the matrix Z (does not changing the Y) by the admissible transformations to the form $Z = \begin{bmatrix} 0 & i\nu \\ i\nu & 0 \end{bmatrix}$, $\nu > 0$. The triple of matrices X, Y, Z is reduced by means of a unitary matrix $U = \begin{bmatrix} \sqrt{2}/2 & \sqrt{2}/2 \\ -\sqrt{2}/2 & \sqrt{2}/2 \end{bmatrix}$ by the transformation UXU^*, UYU^*, UZU^* to the form

$$X = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad Y = \mu \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad Z = \nu \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}; \quad \mu > 0, \nu > 0.$$

c) Let $X \neq 0, \ Z \neq 0, \ Y = 0$. In this case the matrices $X, \ Y, \ Z$ can be reduced to the form

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad Z = \nu \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}; \quad \lambda > 0, \ \nu > 0.$$

d) $X \neq 0, Y \neq 0, Z \neq 0$. In this case the matrices X, Y, Z can be reduced to the form

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \mu \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad Z = \nu \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}; \quad \lambda > 0, \, \mu > 0, \, \nu \in \mathbb{R}, \, \nu \neq 0.$$

Thus, unifying these cases, we may consider that

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \mu \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad Z = \nu \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix},$$
 where $\lambda^2 + \mu^2 + \nu^2 = \frac{1}{4}$ (follows from (4)), and either $\lambda > 0, \ \mu > 0, \ \nu \in \mathbb{R}$; either $\lambda = 0, \ \mu > 0, \ \nu > 0$; or $\lambda > 0, \ \mu = 0, \ \nu > 0$.

Formulas (3) imply

$$P_{1} = \begin{bmatrix} \frac{1}{2} - \lambda & \mu + \nu i \\ \mu - \nu i & \frac{1}{2} + \lambda \end{bmatrix}, \quad P_{2} = \begin{bmatrix} \frac{1}{2} + \lambda & -\mu + \nu i \\ -\mu - \nu i & \frac{1}{2} - \lambda \end{bmatrix},$$

$$P_{3} = \begin{bmatrix} \frac{1}{2} + \lambda & \mu - \nu i \\ \mu + \nu i & \frac{1}{2} - \lambda \end{bmatrix}, \quad P_{4} = \begin{bmatrix} \frac{1}{2} - \lambda & -\mu - \nu i \\ -\mu + \nu i & \frac{1}{2} + \lambda \end{bmatrix}.$$

If $\mu + \nu i = \sqrt{\mu^2 + \nu^2} e^{i\varphi} = \sqrt{\frac{1}{4} - \lambda^2} e^{i\varphi}$ then, passing on to the unitary equivalent representation by means of the matrix $U = \begin{bmatrix} e^{-i\nu p} & 0 \\ 0 & 1 \end{bmatrix}$, we may consider that

$$P_{1} = \begin{bmatrix} \frac{1}{2} - \lambda & \sqrt{\frac{1}{4} - \lambda^{2}} \\ \sqrt{\frac{1}{4} - \lambda^{2}} & \frac{1}{2} + \lambda \end{bmatrix}, \quad P_{2} = \begin{bmatrix} \frac{1}{2} + \lambda & e^{i\chi}\sqrt{\frac{1}{4} - \lambda^{2}} \\ e^{-i\chi}\sqrt{\frac{1}{4} - \lambda^{2}} & \frac{1}{2} - \lambda \end{bmatrix},$$

$$P_{3} = \begin{bmatrix} \frac{1}{2} + \lambda & -e^{-i\chi}\sqrt{\frac{1}{4} - \lambda^{2}} \\ -e^{-i\chi}\sqrt{\frac{1}{4} - \lambda^{2}} & \frac{1}{2} - \lambda \end{bmatrix}, \quad P_{4} = \begin{bmatrix} \frac{1}{2} - \lambda & -\sqrt{\frac{1}{4} - \lambda^{2}} \\ -\sqrt{\frac{1}{4} - \lambda^{2}} & \frac{1}{2} + \lambda \end{bmatrix}, \quad (5)$$

$$0 \le \lambda < 1/2, \text{ and}$$

$$0 < \chi < \pi/2 \text{ when } \lambda = 0,$$

$$-\pi/2 < \chi < \pi/2 \text{ when } 0 < \lambda < 1/2.$$

4. Let $\gamma_3 \neq 0$, then $\{x,y\} = \gamma_3 e$ implies $X \neq 0$ and $Y \neq 0$. Moreover, X and Y has no zero eigenvalues. Indeed, let martix X has the form after digonalization: $X = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix}$. In the matrices Y and Z either $y_{12} \neq 0$ or $z_{12} \neq 0$ (or else the triple of matrices is decomposable). $\{x,y\} = \gamma_3 e, \ \{x,z\} = \gamma_2 e \ \text{imply} \ (\lambda_1 + \lambda_2) z_{12} = 0 \ \text{and} \ (\lambda_1 + \lambda_2) y_{12} = 0$. Therefore we can conclude that $-\lambda_1 = \lambda_2 = \lambda \neq 0$, $\lambda > 0$. The same reasoning is useful also for Y.

Let $X = \begin{bmatrix} -\lambda & 0 \\ 0 & \lambda \end{bmatrix}$. $\{x, y\} = \gamma_3 e$ implies $y_{11} = -\frac{\gamma_3}{2\lambda}$, $y_{22} = \frac{\gamma_3}{2\lambda}$; $\{x, z\} = \gamma_2 e$ implies $z_{11} = -\frac{\gamma_2}{2\lambda}$, $z_{22} = \frac{\gamma_2}{2\lambda}$:

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \frac{1}{2\lambda} \begin{bmatrix} -\gamma_3 & y_{12} \\ \overline{y}_{12} & \gamma_3 \end{bmatrix}, \quad Z = \frac{1}{2\lambda} \begin{bmatrix} -\gamma_2 & z_{12} \\ \overline{z}_{12} & \gamma_2 \end{bmatrix}.$$

 $\{y,z\} = \gamma_1 e$ implies

$$\frac{1}{4\lambda^2} \begin{bmatrix} 2\gamma_2\gamma_3 + y_{12}\overline{z}_{12} + \overline{y}_{12}z_{12} & 0\\ 0 & 2\gamma_2\gamma_3 + y_{12}\overline{z}_{12} + \overline{y}_{12}z_{12} \end{bmatrix} = \gamma_1 I,$$

hence

$$y_{12}\overline{z}_{12} + \overline{y}_{12}z_{12} = 4\gamma_1\lambda^2 - 2\gamma_2\gamma_3. \tag{6}$$

Let us turn to the unitary equivalent representation by means of the unitary matrix $U = \begin{bmatrix} e^{i\varphi} & 0 \\ 0 & 0 \end{bmatrix}$ so that $y_{12} + z_{12} = r_1$ would be real positive; at that $-y_{12} + z_{12}$ remains to be complex in general, $-y_{12} + z_{12} = r_2 e^{i\chi}$.

Then (2) implies

$$(X+Y+Z)^2 = \frac{1}{4\lambda^2} \begin{bmatrix} (2\lambda^2 + \gamma_2 + \gamma_3)^2 + (y_{12} + z_{12})^2 & 0\\ 0 & (2\lambda^2 + \gamma_2 + \gamma_3)^2 + (y_{12} + z_{12})^2. \end{bmatrix} = \alpha_4^2 I$$

and

$$r_1^2 = (y_{12} + z_{12})^2 = 4\alpha_4^2 \lambda^2 - (2\lambda^2 + \gamma_2 + \gamma_3)^2,$$

from which it easy to obtain

$$r_{1} = \sqrt{-4\lambda^{4} + 2(\alpha_{1}^{2} + \alpha_{4}^{2})\lambda^{2} - \frac{1}{4}(\alpha_{4}^{2} - \alpha_{1}^{2})^{2}} =$$

$$= \sqrt{-4\left(\lambda^{2} - \frac{(\alpha_{4}^{2} - \alpha_{1}^{2})^{2}}{4}\right)\left(\lambda^{2} - \frac{(\alpha_{4}^{2} + \alpha_{1}^{2})^{2}}{4}\right)},$$

$$\frac{\alpha_{4} - \alpha_{1}}{2} \leq \lambda \leq \frac{\alpha_{4} + \alpha_{1}}{2}.$$

$$\begin{cases} y_{12} + z_{12} = r_{1}, \\ -y_{12} + z_{12} = r_{2}e^{i\chi}, \end{cases}$$
therefore
$$\begin{cases} y_{12} = (r_{1} - r_{2}e^{i\chi})/2, \\ z_{12} = (r_{1} + r_{2}e^{i\chi})/2, \end{cases}$$

 $y_{12}\overline{z}_{12} + \overline{y}_{12}z_{12} = (r_1^2 - r_2^2)/2 = 4\gamma_1\lambda^2 - 2\gamma_2\gamma_3$ (the last equality follows from (6)). Thus,

$$r_2^2 = r_1^2 + 4\gamma_2\gamma_3 - 8\gamma_1\lambda^2 = -4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2 \ge 0$$

and

$$-y_{12} + z_{12} = r_2 e^{i\chi} = e^{i\chi} \sqrt{-4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2} =$$

$$= e^{i\chi} \sqrt{-4\left(\lambda^2 - \frac{(\alpha_3^2 - \alpha_2^2)^2}{4}\right)\left(\lambda^2 - \frac{(\alpha_3^2 + \alpha_2^2)^2}{4}\right)},$$

$$\frac{\alpha_3 - \alpha_2}{2} \le \lambda \le \frac{\alpha_3 + \alpha_2}{2}.$$

Now we can directly pass on to the determining of the operators of the representation P_1 , P_2 , P_3 , P_4 :

$$\begin{split} P_1 &= \frac{-X + Y + Z}{2\alpha_1} + \frac{1}{2}I = \\ &= \frac{1}{4\alpha_1\lambda} \begin{bmatrix} 2\lambda^2 + 2\alpha_1\lambda - \frac{1}{2}(\alpha_4^2 - \alpha_1^2) & \sqrt{-4\lambda^4 + 2(\alpha_1^2 + \alpha_4^2)\lambda^2 - \frac{1}{4}(\alpha_4^2 - \alpha_1^2)^2} \\ -2\lambda^2 + 2\alpha_1\lambda + \frac{1}{2}(\alpha_4^2 - \alpha_1^2) \end{bmatrix} = \\ &= \frac{1}{4\alpha_1\lambda} \begin{bmatrix} 2\left(\lambda - \frac{1}{2}(\alpha_4 - \alpha_1)\right)\left(\lambda + \frac{1}{2}(\alpha_4 + \alpha_1)\right) & \sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_4 - \alpha_1)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_4 + \alpha_1)^2\right)} \\ \sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_4 - \alpha_1)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_4 + \alpha_1)^2\right)} & -2\left(\lambda + \frac{1}{2}(\alpha_4 - \alpha_1)\right)\left(\lambda - \frac{1}{2}(\alpha_4 + \alpha_1)\right) \end{bmatrix}, \end{split}$$

$$P_2 &= \frac{X - Y + Z}{2\alpha_2} + \frac{1}{2}I = \\ &= \frac{1}{4\alpha_2\lambda} \begin{bmatrix} -2\lambda^2 + 2\alpha_2\lambda + \frac{1}{2}(\alpha_3^2 - \alpha_2^2) \\ e^{-i\chi}\sqrt{-4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2} \end{bmatrix} = \\ &= \frac{1}{4\alpha_2\lambda} \begin{bmatrix} -2\left(\lambda - \frac{1}{2}(\alpha_3 + \alpha_2)\right)\left(\lambda + \frac{1}{2}(\alpha_3 - \alpha_2)\right) \\ e^{-i\chi}\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)} \end{bmatrix} = \\ &= \frac{1}{4\alpha_2\lambda} \begin{bmatrix} -2\left(\lambda - \frac{1}{2}(\alpha_3 + \alpha_2)\right)\left(\lambda + \frac{1}{2}(\alpha_3 - \alpha_2)\right) \\ e^{-i\chi}\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)} \end{bmatrix} = \\ &= \frac{1}{4\alpha_2\lambda} \begin{bmatrix} -2\left(\lambda - \frac{1}{2}(\alpha_3 + \alpha_2)\right)\left(\lambda - \frac{1}{2}(\alpha_3 - \alpha_2)\right) \\ e^{-i\chi}\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)} \end{bmatrix} \\ &= \frac{1}{4\alpha_2\lambda} \begin{bmatrix} -2\left(\lambda - \frac{1}{2}(\alpha_3 + \alpha_2)\right)\left(\lambda - \frac{1}{2}(\alpha_3 - \alpha_2)\right) \\ e^{-i\chi}\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)} \end{bmatrix} \end{bmatrix}$$

$$\begin{split} P_3 &= \frac{X + Y - Z}{2\alpha_3} + \frac{1}{2}I = \\ &= \frac{1}{4\alpha_3\lambda} \begin{bmatrix} -2\lambda^2 + 2\alpha_3\lambda - \frac{1}{2}(\alpha_3^2 - \alpha_2^2) & -e^{-i\chi}\sqrt{-4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2} \\ -e^{-i\chi}\sqrt{-4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2} \end{bmatrix} = \\ &= \frac{1}{4\alpha_3\lambda} \begin{bmatrix} -2\left(\lambda - \frac{1}{2}(\alpha_3 + \alpha_2)\right)\left(\lambda - \frac{1}{2}(\alpha_3 - \alpha_2)\right) & -e^{-i\chi}\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)} \\ -e^{-i\chi}\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)} & 2\left(\lambda + \frac{1}{2}(\alpha_3 + \alpha_2)\right)\left(\lambda + \frac{1}{2}(\alpha_3 - \alpha_2)^2\right) \end{bmatrix}, \\ P_4 &= \frac{-X - Y - Z}{2\alpha_4} + \frac{1}{2}I = \\ &= \frac{1}{4\alpha_4\lambda} \begin{bmatrix} 2\lambda^2 + 2\alpha_4\lambda + \frac{1}{2}(\alpha_4^2 - \alpha_1^2) & -\sqrt{-4\lambda^4 + 2(\alpha_4^2 + \alpha_1^2)\lambda^2 - \frac{1}{4}(\alpha_4^2 - \alpha_1^2)^2} \\ -\sqrt{-4\lambda^4 + 2(\alpha_4^2 + \alpha_1^2)\lambda^2 - \frac{1}{4}(\alpha_4^2 - \alpha_1^2)^2} & -2\lambda^2 + 2\alpha_4\lambda - \frac{1}{2}(\alpha_4^2 - \alpha_1^2) \end{bmatrix} = \\ &= \frac{1}{4\alpha_4\lambda} \begin{bmatrix} 2\left(\lambda + \frac{1}{2}(\alpha_4 + \alpha_1)\right)\left(\lambda + \frac{1}{2}(\alpha_4 - \alpha_1)\right) & -\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_4 + \alpha_1)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_4 - \alpha_1)^2\right)} \\ -\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_4 + \alpha_1)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_4 - \alpha_1)^2\right)} & -2\left(\lambda - \frac{1}{2}(\alpha_4 + \alpha_1)\right)\left(\lambda - \frac{1}{2}(\alpha_4 - \alpha_1)\right) \end{bmatrix}. \end{split}$$

5. Let us construct, according to p. 1, isometries $\Gamma_{0,i}: H_i \to H_0$, $i = \overline{1,4}$ by projectors $P_i: H_0 \to H_0$ and obtain regular indecomposable locally scalar representations of the graph \widetilde{D}_4 :

$$\Gamma_{0,1} = \begin{bmatrix} \sqrt{\frac{\lambda^2 + \alpha_1 \lambda - (\alpha_4^2 - \alpha_1^2)/4}{2\alpha_1 \lambda}} \\ \sqrt{\frac{-\lambda^2 + \alpha_1 \lambda + (\alpha_4^2 - \alpha_1^2)/4}{2\alpha_1 \lambda}} \end{bmatrix}, \quad \Gamma_{0,2} = \begin{bmatrix} \sqrt{\frac{-\lambda^2 + \alpha_2 \lambda + (\alpha_3^2 - \alpha_2^2)/4}{2\alpha_2 \lambda}} \\ e^{-i\chi} \sqrt{\frac{\lambda^2 + \alpha_2 \lambda - (\alpha_3^2 - \alpha_2^2)/4}{2\alpha_2 \lambda}} \end{bmatrix}, \quad \Gamma_{0,3} = \begin{bmatrix} \sqrt{\frac{-\lambda^2 + \alpha_3 \lambda - (\alpha_3^2 - \alpha_2^2)/4}{2\alpha_3 \lambda}} \\ -e^{-i\chi} \sqrt{\frac{\lambda^2 + \alpha_3 \lambda + (\alpha_3^2 - \alpha_2^2)/4}{2\alpha_3 \lambda}} \end{bmatrix}, \quad \Gamma_{0,4} = \begin{bmatrix} \sqrt{\frac{\lambda^2 + \alpha_4 \lambda + (\alpha_4^2 - \alpha_1^2)/4}{2\alpha_4 \lambda}} \\ -\sqrt{\frac{-\lambda^2 + \alpha_4 \lambda - (\alpha_4^2 - \alpha_1^2)/4}{2\alpha_4 \lambda}} \end{bmatrix}. \quad 0 < (\alpha_4 - \alpha_1)/2 \le \lambda \le \min((\alpha_2 + \alpha_3)/2, (\alpha_1 + \alpha_4)/2), \quad -\pi < \chi \le \pi. \end{bmatrix}$$

Thus, indecomposable regular (not necessarily normalized) locally scalar representations of the graph \widetilde{D}_4 depend on 6 real parameters (on the 4 of 5 parameters α_1' , α_2' , α_3' , α_4' , α_0' , connected by the relation $\alpha_1' + \alpha_2' + \alpha_3' + \alpha_4' = 2\alpha_0'$, $\alpha_i = \frac{\alpha_i'}{\alpha_0'}$, and parameters λ and χ).

References

1. L. A. Nazarova Representations of quivers of infinite type. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 37 (1973), 752–791.

- 2. S. A. Kruglyak, A. V. Roiter Locally scalar representations of graphs in the category of Hilbert spaces. Funct. Anal. Appl. 39 (2005), no. 2, 91–105
- 3. Yu. N. Bespalov. Sets of orthoprojectors connected by relations. Ukrainian Math. J. 44 (1992), no. 3, 269–277.
- 4. V. Ostrovskyi and Yu. Samoilenko Introduction to the theory of representations of finitely presented *-algebras. I. Representations by bounded operators. Reviews in Mathematics and Mathematical Physics, 11, pt.1. Harwood Academic Publishers, Amsterdam, 1999. iv+261 pp. ISBN: 90-5823-042-2
- 5. D. V. Galinsky, S. A. Kruglyak Representations of *-algebras generated by linearly related orthogonal projections. (Ukrainian.) Visnyk Kyiv. derzh. Univ. No. 2, 24–31, 1999.
- 6. D. V. Galinsky. Representations of *-algebras generated by orthogonal projections satisfying a linear relation. Methods Funct. Anal. Topology 4 (1998), no. 3, 27–32.
- 7. D. V. Galinsky, M. A. Muratov On representations of algebras generated by sets of three and four orthoprojections. Spectral and evolutionary problems, Vol. 8 (Sevastopol, 1997), 15–22, Tavria Publ., Simferopol, 1998.
- 8. S. A. Kruglyak, V. I. Rabanovich, Yu. S. Samoilenko On sums of projections. Funct. Anal. Appl. 36 (2002), no. 3, 182–195
- 9. S. Kruglyak, A. Kyrychenko. On four orthogonal projections that satisfy the linear relation $\alpha_1 P_1 + \alpha_2 P_2 + \alpha_3 P_3 + \alpha_4 P_4 = I$, $\alpha_i > 0$. Symmetry in nonlinear mathematical physics, Part 1, 2 (Kyiv, 2001), 461–465, Pr. Inst. Mat. Nats. Akad. Nauk Ukr. Mat. Zastos., 43, Part 1, 2, Natsional. Akad. Nauk Ukrai"ni, Īnst. Mat., Kiev, 2002.
- 10. A. Strelets. Description of the representations of the algebras generated by four linearly related idempotents. J. Algebra Appl. 4 (2005), no. 6, 671–681.
- 11. S. Kruglyak, S. Popovich, Yu. Samoilenko. The spectral problem and *-representations of algebras associated with Dynkin graphs. J. Algebra Appl. 4 (2005), no. 6, 761–776.
- 12. I. K. Redchuk, A. V. Roiter. Singular locally scalar representations of quivers in Hilbert spaces, and separating functions. Ukrainian Math. J. 56 (2004), no. 6, 947–963.
- 13. S. A. Kruglyak. Coxeter functors for a certain class of *-quivers and *-algebras. Methods Funct. Anal. Topology 8 (2002), no. 4, 49–57.