### THE $\,\varepsilon$ - REVISED SYSTEM OF THE RIGID BODY WITH THREE LINEAR CONTROLS

#### Dan COMĂNESCU, Mihai IVAN and Gheorghe IVAN

**Abstract.** In this paper we introduce the  $\varepsilon$ -revised system associated to a Hamilton-Poisson system. The  $\varepsilon$ -revised system of the rigid body with three linear controls is defined and some of its geometrical and dynamical properties are investigated. <sup>1</sup>

#### 1 Introduction

It is well known that many dynamical systems can be formulated using a Poisson structure (see for instance, R. Abraham and J. E. Marsden [1] and M. Puta [11]).

The metriplectic systems was introduced by P. J. Morrison in the paper [8]. These systems combine both the conservative and dissipative systems.

A metriplectic system is a differential system of the form  $\dot{x} = PdH + gdC$ , where P is a Poisson tensor on a manifold M,  $\mathbf{g}$  is a symmetric tensor of type (2,0) on M, and H and C are two smooth functions on M with the additional requirements:

(a) 
$$PdC = 0$$
; (b)  $gdH = 0$  and (c)  $dC \cdot gdC \le 0$ .

The differential systems of the form  $\dot{x} = PdH + gdC$  which satisfies only the conditions (a) and (b) are called almost metriplectic systems (see Fish, [2]; Marsden, [7]; Ortega and Planas - Bielsa, [9]). An interesting class of almost metriplectic systems are so-called the revised dynamical systems associated to Hamilton-Poisson systems (see Gh. Ivan and D. Opriş, [5]).

The control of the rotation rigid body is one of the problems with a large practical applicability. For this reason, in this paper we study the  $\varepsilon$  - revised dynamical system associated to the rigid body with three linear controls.

### 2 Almost metriplectic systems

We start this section with the presentation of the concept of almost metriplectic manifold (see Ortega and Planas- Bielsa, [9]).

Let M be a smooth manifold of dimension n and let  $C^{\infty}(M)$  be the ring of smooth real-valued functions on M.

A Leibniz manifold is a pair  $(M, [\cdot, \cdot])$ , where  $[\cdot, \cdot]$  is a Leibniz bracket on M, that is  $[\cdot, \cdot]: C^{\infty}(M) \times C^{\infty}(M) \to C^{\infty}(M)$  is a  $\mathbf R$  - bilinear operation satisfying the following two conditions:

(i) the left Leibniz rule:

$$[f_1 \cdot f_2, f_3] = [f_1, f_3] \cdot f_2 + f_1 \cdot [f_2, f_3]$$
 for all  $f_1, f_2, f_3 \in C^{\infty}(M)$ ;

Key words and phrases: almost metriplectic system,  $\varepsilon$ - revised system, rigid body

<sup>&</sup>lt;sup>1</sup>2000 Mathematical Subject Classification: 58F05.

(ii) the right Leibniz rule:

$$[f_1, f_2 \cdot f_3] = [f_1, f_2] \cdot f_3 + f_2 \cdot [f_1, f_3]$$
 for all  $f_3, f_3, f_3 \in C^{\infty}(M)$ 

where "." denotes the ordinary multiplication of functions.

Let P and  $\mathbf{g}$  be two tensor fields of type (2,0) on M and  $\varepsilon \in \mathbf{R}$  be a parameter. We define the map  $[\cdot,(\cdot,\cdot)]_{\varepsilon}:C^{\infty}(M)\times (C^{\infty}(M)\times C^{\infty}(M))\to C^{\infty}(M)$  by:

$$[f,(h_1,h_2)]_{\varepsilon} = P(df,dh_1) + \varepsilon \mathbf{g}(df,dh_2), \quad \text{for all} \quad f,h_1,h_2 \in C^{\infty}(M).$$
 (1)

**Proposition 2.1.** The map  $[\cdot,(\cdot,\cdot)]_{\varepsilon}$  given by (1) satisfy the following relations:

- (i)  $[af_1 + bf_2, (h_1, h_2)]_{\varepsilon} = a[f_1, (h_1, h_2)]_{\varepsilon} + b[f_2, (h_1, h_2)]_{\varepsilon};$
- (ii)  $[f, a(h_1, h_2) + b(h'_1, h'_2)]_{\varepsilon} = a[f, (h_1, h_2)]_{\varepsilon} + b[f, (h'_1, h'_2)]_{\varepsilon};$
- (iii)  $[ff_1, (h_1, h_2)]_{\varepsilon} = f[f_1, (h_1, h_2)]_{\varepsilon} + f_1[f, (h_1, h_2)]_{\varepsilon};$
- $[f, h(h_1, h_2)]_{\varepsilon} = h[f, (h_1, h_2)]_{\varepsilon} + h_1 P(df, dh) + \varepsilon h_2 \mathbf{g}(df, dh),$

for all  $f, f_1, f_2, h_1, h_2, h'_1, h'_2 \in C^{\infty}(M)$  and  $a, b \in \mathbf{R}$ .

**Proof.** Applying the properties of the differential of functions and using that P and  $\mathbf{g}$  are  $\mathbf{R}$ - bilinear maps, it is easy to establish the relations (i) - (iv).

We consider the map  $[[\cdot,\cdot]]_{\varepsilon}: C^{\infty}(M)\times C^{\infty}(M)\to C^{\infty}(M)$  defined by:

$$[[f,h]]_{\varepsilon} = [f,(h,h)]_{\varepsilon}, \text{ for all } f,h \in C^{\infty}(M).$$
 (2)

Therefore, the map  $[[\cdot,\cdot]]_{\varepsilon}: C^{\infty}(M)\times C^{\infty}(M)\to C^{\infty}(M)$  is given by:

$$[[f,h]]_{\varepsilon} = P(df,dh) + \varepsilon \mathbf{g}(df,dh), \text{ for all } f,h \in C^{\infty}(M).$$
 (3)

**Proposition 2.2.** The bracket  $[[\cdot,\cdot]]_{\varepsilon}$  on M given by (3) verify the right Leibniz rule:

$$[[f, hh']]_{\varepsilon} = h[[f, h']]_{\varepsilon} + h'[[f, h]]_{\varepsilon}, \quad \text{for all} \quad f, h, h' \in C^{\infty}(M).$$

**Proof.** Indeed,  $[[f, hh']]_{\varepsilon} = [f, (hh', hh')]_{\varepsilon} = [f, h(h', h')]_{\varepsilon}$ . Putting  $h_1 = h_2 = h'$  in the relation (iv) from Proposition 2.1, we have succesive:  $[f, h(h', h')]_{\varepsilon} = h[f, (h', h')] + h'P(df, dh) + \varepsilon h'\mathbf{g}(df, dh) = h[f, (h', h')]_{\varepsilon} + h'(P(df, dh) + \varepsilon \mathbf{g}(df, dh)) = h[f, (h', h')]_{\varepsilon} + h'[f, (h, h)]_{\varepsilon} = h[[f, h']]_{\varepsilon}$ .

By Proposition 2.1. (i), (ii) and (iii) and Proposition 2.2, we have that the map  $[[\cdot,\cdot]]_{\varepsilon}$  given by (3) is a Leibniz bracket on M. Hence,  $[[\cdot,\cdot]]_{\varepsilon}$  defines a Leibniz structure on the manifold M and  $(M,P,\mathbf{g},[[\cdot,\cdot]]_{\varepsilon})$  is a Leibniz manifold for each  $\varepsilon \in \mathbf{R}$ .

A Leibniz manifold  $(M, P, \mathbf{g}, [[\cdot, \cdot]]_{\varepsilon})$  such that P is a skewsymmetric tensor field and  $\mathbf{g}$  is a symmetric tensor field is called almost metriplectic manifold. In other words, given a skewsymmetric tensor field P of type (2,0) and a symmetric tensor field  $\mathbf{g}$  of type (2,0) on a manifold M, we can define an almost metriplectic structure on M.

If the tensor field P is Poisson and the tensor field  $\mathbf{g}$  is nondegenerate, then  $(M, P, \mathbf{g}, [[\cdot, \cdot]]_{\varepsilon})$  is a metriplectic manifold, see Ortega & Planas - Bielsa [9].

**Proposition 2.3.** Let  $(M, P, \mathbf{g}, [[\cdot, \cdot]]_{\varepsilon})$  be an almost metriplectic manifold. If there exist  $h_1, h_2 \in C^{\infty}(M)$  such that  $P(df, dh_2) = 0$  and  $\mathbf{g}(df, dh_1) = 0$  for all  $f \in C^{\infty}(M)$ , then the bracket  $[[\cdot, \cdot]]$  given by (3) satisfies the relation:

$$[[f, h_1 + h_2]]_{\varepsilon} = [f, (h_1, h_2)]_{\varepsilon}, \quad \text{for all } f \in C^{\infty}(M).$$

**Proof.** Indeed,  $[[f, h_1 + h_2]]_{\varepsilon} = P(df, d(h_1 + h_2)) + \varepsilon \mathbf{g}(df, d(h_1 + h_2)) = P(df, dh_1 + dh_2) + \varepsilon \mathbf{g}(df, dh_1 + dh_2) = P(df, dh_1) + P(df, dh_2) + \varepsilon \mathbf{g}(df, dh_1) + \varepsilon \mathbf{g}(df, dh_2) = P(df, dh_1) + \varepsilon \mathbf{g}(df, dh_2) = [f, (h_1, h_2)]_{\varepsilon}.$ 

Let  $(M, P, \mathbf{g}, [[\cdot, \cdot]]_{\varepsilon})$  be an almost metriplectic manifold and let  $h_1, h_2 \in C^{\infty}(M)$  two functions such that  $P(df, dh_2) = 0$  and  $\mathbf{g}(df, dh_1) = 0$  for all  $f \in C^{\infty}(M)$ . The vector field  $X_{h_1h_2}$  given by:

$$X_{h_1h_2}(f) = [[f, h_1 + h_2]]_{\varepsilon}$$
 for any  $f \in C^{\infty}(M)$ 

is called the *Leibniz vector field* associated to the triple  $(h_1, h_2, \varepsilon)$  on M. Taking account into Proposition 2.3 and (1),  $X_{h_1h_2}$  is given by:

$$X_{h_1h_2}(f) = [f, (h_1, h_2)]_{\varepsilon} = P(df, dh_1) + \varepsilon \mathbf{g}(df, dh_2), \quad \text{for all } f \in C^{\infty}(M).$$
 (5)

In local coordinates on M, the differential system given by:

$$\dot{x}^{i} = [[x^{i}, h_{1} + h_{2}]]_{\varepsilon} = [x^{i}, (h_{1}, h_{2})]_{\varepsilon}$$
(6)

where

$$[x^{i},(h_{1},h_{2})]_{\varepsilon} = X_{h_{1}h_{2}}(x^{i}) = P^{ij}\frac{\partial h_{1}}{\partial x^{j}} + \varepsilon g^{ij}\frac{\partial h_{2}}{\partial x^{j}}, \ i,j = \overline{1,n}$$

$$(7)$$

with  $P^{ij} = P(dx^i, dx^j)$  and  $g^{ij} = \mathbf{g}(dx^i, dx^j)$ , is called the almost metriplectic system on M associated to the Leibniz vector field  $X_{h_1h_2}$  with the bracket  $[[\cdot, \cdot]]_{\varepsilon}$ .

We denote the matrix of the tensor fields P and  $\mathbf{g}$  respectively by  $P = (P^{ij})$  and  $g = (g^{ij})$ . We have that P is a skewsymmetric matrix and g is a symmetric matrix.

We give now a way for to produce almost metriplectic manifolds.

**Proposition 2.4.** For a skewsymmetric tensor P of type (2,0) on a manifold M and two functions  $h_1, h_2 \in C^{\infty}(M)$  such that  $P(df, dh_2) = 0$  for all  $f \in C^{\infty}(M)$ , there exists a symmetric tensor  $\mathbf{g}$  of type (2,0) on M such that  $\mathbf{g}(df, dh_1) = 0$  for all  $f \in C^{\infty}(M)$  and  $(M, P, \mathbf{g}, [[\cdot, \cdot]]_{\varepsilon})$  is an almost metriplectic manifold.

**Proof.** In a system of local coordinates on M, let  $g = (g^{ij})$  the matrix of the symmetric tensor  $\mathbf{g}$  which must to be determined. Then, the components  $g^{ij}$ ,  $i, j = \overline{1.n}$  verify the system of differential equations  $g^{ij} \frac{\partial h_1}{\partial x^j} = 0$ ,  $i, j = \overline{1, n}$ .

In a chart U such that  $\frac{\partial h_1}{\partial x^j}(x) \neq 0$ , the components  $g^{ij}$  are given by:

$$\begin{cases}
g^{ii}(x) = -\sum_{k=1, k\neq i}^{n} \left(\frac{\partial h_1}{\partial x^k}\right)^2 \\
g^{ij}(x) = \frac{\partial h_1}{\partial x^i} \frac{\partial h_1}{\partial x^j}, & \text{for } i \neq j
\end{cases}$$
(8)

Applying now Proposition 2.3 we obtain the result.

Proposition 2.4 is useful when we consider the  $\varepsilon$ - revised system of a Hamilton-Poisson system.

For this, let be a Hamilton-Poisson system on M described by the Poisson tensor P having the matrix  $P=(P^{ij})$  and by the Hamiltonian function  $h_1 \in C^{\infty}(M)$  with the Casimir function  $h_2 \in C^{\infty}(M)$  (i.e.  $P^{ij} \frac{\partial h_2}{\partial x^j} = 0$  for  $i, j = \overline{1, n}$ ). The differential equations of the Hamilton-Poisson system are the following:

$$\dot{x}^i = P^{ij} \frac{\partial h_1}{\partial x^j}, \quad i, j = \overline{1, n}. \tag{9}$$

Using (8), we determine the matrix  $g = (g^{ij})$  and we have:

$$g^{ij}\frac{\partial h_1}{\partial x^j} = 0, \quad i, j = \overline{1, n}. \tag{10}$$

Applying now Proposition 2.4, for each  $\varepsilon \in \mathbf{R}$ , we obtain an almost metriplectic structure on M associated to system (9). The differential system associated to this structure is called the  $\varepsilon$  - revised system of the Hamilton - Poisson system.

Hence, the  $\varepsilon$  - revised system of the Hamilton - Poisson system defined by (9) is:

$$\dot{x}^{i} = P^{ij} \frac{\partial h_{1}}{\partial x^{j}} + \varepsilon g^{ij} \frac{\partial h_{2}}{\partial x^{j}}, \quad i, j = \overline{1, n}.$$

$$(11)$$

The terms  $g^{ij}\frac{\partial h_2}{\partial x^j}$ ,  $i,j=\overline{1,n}$  from the  $\varepsilon$  - revised system (11) describe a *cube perturbation* of the Hamilton - Poisson system.

**Remark 2.1.** We observe that the 0- revised system (11) coincide with the Hamilton - Poisson system (9).  $\Box$ 

## 3 The $\varepsilon$ - revised system associated to the rigid body with three linear controls

The rigid body equations with three linear controls (see, M. Puta and D. Comănescu [12]) are given by:

$$\begin{cases} \dot{x}^1 = (a_3 - a_2)x^2x^3 + cx^2 - bx^3 \\ \dot{x}^2 = (a_1 - a_3)x^1x^3 - cx^1 + ax^3 \\ \dot{x}^3 = (a_2 - a_1)x^1x^2 + bx^1 - ax^2 \end{cases}$$
(12)

where  $x(t)=(x^1(t),x^2(t),x^3(t))\in \mathbf{R}^3$  and  $a_1=\frac{1}{I_1},\ a_2=\frac{1}{I_2},\ a_3=\frac{1}{I_3}$  with  $I_1>I_2>I_3>0$  (  $I_1,I_2,I_3$  being the principal moments of inertia of the body ) and  $a,b,c\in \mathbf{R}$  are feedback parameters. We have  $0< a_1< a_2< a_3$ .

The dynamics (12) is described by the Poisson tensor  $\Pi$  and by the Hamiltonian H on  $\mathbb{R}^3$  given by:

$$\Pi(x) = \begin{pmatrix} 0 & -x^3 & x^2 \\ x^3 & 0 & -x^1 \\ -x^2 & x^1 & 0 \end{pmatrix},$$
(13)

$$H(x) = \frac{1}{2} [a_1(x^1)^2 + a_2(x^2)^2 + a_3(x^3)^2] + ax^1 + bx^2 + cx^3.$$
 (14)

Using (13) and (14), the dynamics (12) can be written in the matrix form:

$$\dot{x}(t) = \Pi(x(t)) \cdot \nabla H(x(t)), \tag{15}$$

where  $\dot{x}(t) = (\dot{x}^1(t), \dot{x}^2(t), \dot{x}^3(t))^T$  and  $\nabla H(x(t))$  is the gradient of the Hamiltonian function H with respect to the canonical metric on  $\mathbf{R}^3$ .

Therefore, the dynamics (12) has the Hamilton-Poisson formulation  $(\mathbf{R}^3, \Pi, H)$ , where  $\Pi$  and H are given by (13) and (14).

The function  $C \in C^{\infty}(\mathbf{R}^3)$  given by:

$$C(x) = \frac{1}{2}[(x^1)^2 + (x^2)^2 + (x^3)^2]$$
(16)

is a Casimir of the configuration  $(\mathbf{R}^3,\Pi)$ , i.e.

$$C(x) \cdot \nabla H(x) = O. \tag{17}$$

Applying the relations (8) for  $P = \Pi$ ,  $h_1(x) = H(x)$  and  $h_2(x) = C(x)$ , the symmetric tensor **g** is given by the matrix:

$$g = \begin{pmatrix} -(a_2x^2 + b)^2 - (a_3x^3 + c)^2 & (a_1x^1 + a)(a_2x^2 + b) & (a_1x^1 + a)(a_3x^3 + c) \\ (a_1x^1 + a)(a_2x^2 + b) & -(a_1x^1 + a)^2 - (a_3x^3 + c)^2 & (a_2x^2 + b)(a_3x^3 + c) \\ (a_1x^1 + a)(a_3x^3 + c) & (a_2x^2 + b)(a_3x^3 + c) & -(a_1x^1 + a)^2 - (a_2x^2 + b)^2 \end{pmatrix}$$

since 
$$\frac{\partial h_1}{\partial x^1} = a_1 x^1 + a$$
,  $\frac{\partial h_1}{\partial x^2} = a_2 x^2 + b$ ,  $\frac{\partial h_1}{\partial x^3} = a_3 x^3 + c$ .

We have

$$g(x) \cdot \nabla h_2(x) = (v_1(x), v_2(x), v_3(x))^T$$
(18)

where

$$\begin{cases} v_1(x) = -[(a_2x^2 + b)^2 + (a_3x^3 + c)^2]x^1 + (a_1x^1 + a)[(a_2x^2 + b)x^2 + (a_3x^3 + c)x^3] \\ v_2(x) = -[(a_1x^1 + a)^2 + (a_3x^3 + c)^2]x^2 + (a_2x^2 + b)[(a_1x^1 + a)x^1 + (a_3x^3 + c)x^3] \\ v_3(x) = -[(a_1x^1 + a)^2 + (a_2x^2 + b)^2]x^3 + (a_3x^3 + c)[(a_1x^1 + a)x^1 + (a_2x^2 + b)x^2] \end{cases}$$

$$(19)$$

The  $\varepsilon$  - revised system associated to dynamics (12) is:

$$\begin{cases}
\dot{x}^{1} = [(a_{3} - a_{2})x^{2}x^{3} + cx^{2} - bx^{3}] + \varepsilon v_{1}(x) \\
\dot{x}^{2} = [(a_{1} - a_{3})x^{1}x^{3} - cx^{1} + ax^{3}] + \varepsilon v_{2}(x) \\
\dot{x}^{3} = [(a_{2} - a_{1})x^{1}x^{2} + bx^{1} - ax^{2}] + \varepsilon v_{3}(x)
\end{cases} (20)$$

The differential system (20) is called the  $\varepsilon$  - revised system of the rigid body with three linear controls. Taking a = b = c = 0 and  $\varepsilon = 1$  in (20), we obtain the revised system of the free rigid body, see [5].

Vector writing of the  $\varepsilon$ - revised system (20). We introduce the following notations:

$$\mathbf{x} = (x^1, x^2, x^3), \quad \mathbf{v} = (v_1, v_2, v_3), \quad \mathbf{a} = (a, b, c), \quad \mathbf{m}(\mathbf{x}) = (a_1 x^1 + a, a_2 x^2 + b, a_3 x^3 + c).$$

For all  $\mathbf{u} = (u_1, u_2, u_3), \mathbf{w} = (w_1, w_2, w_3) \in \mathbf{R}^3$ , the following relation holds:

$$\mathbf{u} \cdot [\mathbf{w} \times (\mathbf{u} \times \mathbf{w})] = (\mathbf{u} \times \mathbf{w})^2, \text{ with } (\mathbf{u} \times \mathbf{w})^2 = (\mathbf{u} \times \mathbf{w}) \cdot (\mathbf{u} \times \mathbf{w})$$
 (21)

where " $\times$ " and " $\cdot$ " denote the cross product resp. inner product in  $\mathbf{R}^3$ ; that is:  $\mathbf{u} \times \mathbf{w} = (u_2w_3 - u_3w_2, u_3w_1 - u_1w_3, u_1w_2 - u_2w_1), \quad \mathbf{u} \cdot \mathbf{w} = u_1w_1 + u_2w_2 + u_3w_3.$ 

With the above notations, the dynamics (12) has the vector form:

$$\dot{\mathbf{x}} = \mathbf{x} \times \mathbf{m}(\mathbf{x}). \tag{22}$$

It is not hard to verify the following equality:

$$\mathbf{v} = (\mathbf{x} \times \mathbf{m}(\mathbf{x})) \times \mathbf{m}(\mathbf{x}). \tag{23}$$

Using the relations (22), (23) and (20), we can written the  $\varepsilon$ - revised system (20) in the vector form:

$$\dot{\mathbf{x}} = \mathbf{x} \times \mathbf{m}(\mathbf{x}) + \varepsilon [(\mathbf{x} \times \mathbf{m}(\mathbf{x})) \times \mathbf{m}(\mathbf{x})]. \tag{24}$$

#### 4 The equilibrium points of the $\varepsilon$ - revised system

The equilibrium points of the Hamilton - Poisson system (12) (or (22)) are solutions of the vector equation:

$$\mathbf{x} \times \mathbf{m}(\mathbf{x}) = \mathbf{0}.\tag{25}$$

The equilibrium points of the  $\varepsilon$  - revised system (20) ( or (24) ) are solutions of the vector equation:

$$\mathbf{x} \times \mathbf{m}(\mathbf{x}) + \varepsilon[(\mathbf{x} \times \mathbf{m}(\mathbf{x})) \times \mathbf{m}(\mathbf{x})] = \mathbf{0}.$$
 (26)

**Theorem 4.1.** The Hamilton - Poisson system (12) and its revised system (20) have the same equilibrium points.

**Proof.** Let  $\mathbf{x}_0$  be an equilibrium point of the system (12). According with (25) follows  $\mathbf{x}_0 \times \mathbf{m}(\mathbf{x}_0) = \mathbf{0}$ . We have that  $\mathbf{x}_0$  is a solution of the vector equation (26), since  $x_0 \times \mathbf{m}(\mathbf{x}_0) + (\mathbf{x}_0 \times \mathbf{m}(\mathbf{x}_0)) \times \mathbf{m}(\mathbf{x}_0) = \mathbf{0} + \mathbf{0} \times \mathbf{m}(\mathbf{x}_0) = \mathbf{0}$ . Hence  $\mathbf{x}_0$  is an equilibrium point of the  $\varepsilon$ - revised system (20).

Conversely, let  $\mathbf{x}_0$  be an equilibrium point for (20). Using (25) it follows

(a)  $\mathbf{x}_0 \times \mathbf{m}(\mathbf{x}_0) + \varepsilon [\mathbf{x}_0 \times \mathbf{m}(\mathbf{x}_0)) \times \mathbf{m}(\mathbf{x}_0)] = \mathbf{0}$ 

The relation (a) can be written in the form:

(b)  $\mathbf{x}_0 \times \mathbf{m}(\mathbf{x}_0) - \varepsilon[\mathbf{m}(\mathbf{x}_0) \times (\mathbf{x}_0) \times \mathbf{m}(\mathbf{x}_0))] = \mathbf{0}$ 

Multiplying the relation (b) with the vector  $\mathbf{x}_0$ , we obtain:

 $(c) \quad \mathbf{x}_0 \cdot (\mathbf{x}_0 \times \mathbf{m}(\mathbf{x}_0)) - \varepsilon \mathbf{x}_0 \cdot [\mathbf{m}(\mathbf{x}_0) \times (\mathbf{x}_0) \times \mathbf{m}(\mathbf{x}_0))] = \mathbf{0}.$ 

Using the equality (21), the relation (c) is equivalent with:

 $(d) \qquad -\varepsilon(\mathbf{x}_0 \times \mathbf{m}(\mathbf{x}_0))^2 = \mathbf{0}.$ 

From (d) (if  $\varepsilon \neq 0$ ), follows  $\mathbf{x}_0 \times \mathbf{m}(\mathbf{x}_0) = \mathbf{0}$ , that is  $\mathbf{x}_0$  is an equilibrium point for (20).

The equilibrium points of the dynamics (12) are well-known (see M. Puta and D. Comănescu, [12]) and these are presented in the following proposition.

**Proposition 4.1.** ([12]) The equilibrium points of the Hamilton - Poisson system (12) are the following:

(i)  $e_1 = (0,0,0);$ 

(ii) 
$$e_2 = (\frac{a}{\lambda - a_1}, \frac{b}{\lambda - a_2}, \frac{c}{\lambda - a_3})$$
 for  $\lambda \in \mathbf{R} \setminus \{a_1, a_2, a_3\}$ ;

(iii) 
$$e_3 = (\alpha, -\frac{b}{a_2 - a_1}, \frac{c}{a_1 - a_3})$$
 for  $\alpha \in \mathbf{R}$ , if  $a = 0$ ;

(iv) 
$$e_4 = (\frac{a}{a_2 - a_1}, \alpha, -\frac{c}{a_3 - a_2})$$
 for  $\alpha \in \mathbf{R}$ , if  $b = 0$ ;

$$(v)$$
  $e_5 = \left(-\frac{a}{a_1 - a_2}, \frac{b}{a_3 - a_2}, \alpha\right) \quad for \quad \alpha \in \mathbf{R}, \quad if \quad c = 0.$ 

By Theorem 4.1, the equilibrium points of the  $\varepsilon$ - revised system (20) are  $e_1, ..., e_5$  indicated in the Proposition 4.1.

It is well-known that the dynamics (12) have the first integrals H and C given by (14) and (16). These first integrals may be written thus:

$$H(x^1, x^2, x^3) = \frac{1}{2} \mathbf{x} \cdot \mathbf{I}^{-1} \mathbf{x} + \mathbf{a} \cdot \mathbf{x} \quad \text{and} \quad C(x^1, x^2, x^3) = \frac{1}{2} \mathbf{x}^2$$
 (27)

where  $\mathbf{I}$  is inertia tensor and  $\mathbf{I}^{-1}$  is its inverse. We have:

$$\frac{dH}{dt}(\mathbf{x}) = \mathbf{m}(\mathbf{x}) \cdot \dot{\mathbf{x}} \quad \text{and} \quad \frac{dC}{dt}(\mathbf{x}) = \mathbf{x} \cdot \dot{\mathbf{x}}.$$
 (28)

Indeed, 
$$\frac{dH}{dt} = (a_1x^1 + a)\dot{x}^1 + (a_2x^2 + b)\dot{x}^2 + (a_3x^3 + c)\dot{x}^3 = \mathbf{m}(\mathbf{x}) \cdot \dot{\mathbf{x}}$$
 and  $\frac{dC}{dt} = x^1\dot{x}^1 + x^2\dot{x}^2 + x^3\dot{x}^3 = \mathbf{x} \cdot \dot{\mathbf{x}}.$ 

**Theorem 4.2.** (i) For each  $\varepsilon \in \mathbb{R}$ , the function H given by (14) is a first integral for the  $\varepsilon$ - revised system (20).

(ii) If  $\mathbf{x}: \mathbf{R} \to \mathbf{R}^3$  is a solution of the  $\varepsilon$ - revised system, then:

$$\frac{d}{dt}(\frac{1}{2}\mathbf{x}^2) = -\varepsilon(\mathbf{x} \times \mathbf{m}(\mathbf{x}))^2. \tag{29}$$

(iii) For  $\varepsilon \in \mathbf{R}^*$ , the function C is not a first integral for the  $\varepsilon$ - revised system.

**Proof.** (i) Multiplying the relation (24) with the vector  $\mathbf{m}(\mathbf{x})$ , we have:  $\mathbf{m}(\mathbf{x}) \cdot \dot{\mathbf{x}} = \mathbf{m}(\mathbf{x}) \cdot (\mathbf{x} \times \mathbf{m}(\mathbf{x})) + \varepsilon \mathbf{m}(\mathbf{x}) \cdot [(\mathbf{x} \times \mathbf{m}(\mathbf{x})) \times \mathbf{m}(\mathbf{x})] = \mathbf{0}$ . Applying now (28), we obtain  $\frac{dH}{dt} = \mathbf{m}(\mathbf{x}) \cdot \dot{\mathbf{x}} = \mathbf{0}$ . Hence H is a first integral for the system (20). (ii) Multiplying the relation (24) with the vector  $\mathbf{x}$ , we have

 $\mathbf{x} \cdot \dot{\mathbf{x}} = \mathbf{x} \cdot (\mathbf{x} \times \mathbf{m}(\mathbf{x})) + \varepsilon \mathbf{x} \cdot [(\mathbf{x} \times \mathbf{m}(\mathbf{x})) \times \mathbf{m}(\mathbf{x})] = -\varepsilon \mathbf{x} \cdot [\mathbf{m}(\mathbf{x}) \times (\mathbf{x} \times \mathbf{m}(\mathbf{x}))].$ Using now the equality (21), we obtain  $\mathbf{x} \cdot \dot{\mathbf{x}} = -\varepsilon (\mathbf{x} \times \mathbf{m}(\mathbf{x}))^2$ . Then, we have  $\frac{d}{dt} (\frac{1}{2} \mathbf{x}^2) = \mathbf{x} \cdot \dot{\mathbf{x}} = -(\mathbf{x} \times \mathbf{m}(\mathbf{x}))^2.$ 

(iii) This assertion follows from the second relation of (28) and (ii).

**Remark 4.1.** The function H given by (14) can be put in the equivalent form:

$$H(x^{1}, x^{2}, x^{3}) = \frac{1}{2} \left[ a_{1}(x^{1} + \frac{a}{a_{1}})^{2} + a_{2}(x^{2} + \frac{b}{a_{2}})^{2} + a_{3}(x^{3} + \frac{c}{a_{3}})^{2} \right] - \frac{1}{2} \left( \frac{a^{2}}{a_{1}} + \frac{b^{2}}{a_{2}} + \frac{c^{2}}{a_{3}} \right).$$
(30)

For a given constant  $k \in \mathbf{R}$ , the geometrical image of the surface:

$$H(x^1, x^2, x^3) = k$$

is an ellipsoid, since  $a_1 > 0, a_2 > 0, a_3 > 0$ .

**Proposition 4.2.** The set of equilibrium points which belong to the ellipsoid  $H(x^1, x^2, x^3) = k$ , is finite.

**Proof.** Following the description of the equilibrium points given in Proposition 4.1, we remark that:

- (i) the equilibrium points of the form  $e_3$  (similarly, for  $e_4$  and  $e_5$ ) make a straight line; the intersection between a straight line and an ellipsoid have at most two points; we deduce that on the chosen ellipsoid there exist at most two points of the form  $e_3$ .
- (ii) the equilibrium points of the form  $e_2$  can be obtained by solving with respect  $\lambda$  the following equation:

$$\frac{1}{2}\left[a_1\left(\frac{a}{\lambda - a_1}\right)^2 + a_2\left(\frac{b}{\lambda - a_2}\right)^2 + a_3\left(\frac{c}{\lambda - a_3}\right)^2\right] + \frac{a^2}{\lambda - a_1} + \frac{b^2}{\lambda - a_2} + \frac{c^2}{\lambda - a_3} = k.$$

The above equation is equivalent with the determination of roots of a polynomial of degree at most 6; therefore on the chosen ellipsoid there exist at most 6 equilibrium points of the form  $e_2$ .

#### 5 The behaviour of the solutions of the $\varepsilon$ - revised system

**Theorem 5.1** (i) The solutions of the  $\varepsilon$ -revised system are bounded.

- (ii) The maximal solutions of the  $\varepsilon$ -revised system are globally solutions (i.e. these are defined on  $\mathbf{R}$ ).
- **Proof** (i) Given a solution of (20), there exists a constant k such that its trajectory lie on the ellipsoid  $H(x^1, x^2, x^3) = k$ . From this we deduce that all solutions are bounded
- (ii) Let  $\mathbf{x}:(m,M)\subset\mathbf{R}\to\mathbf{R}^3$  be a maximal solution. We assume that  $\mathbf{x}$  is not globally. It follows  $m>-\infty$  or  $M<\infty$ . In these situations, we known that there exists  $k\in\mathbf{R}$  such that  $H(x^1,x^2,x^3)=k$  for all  $t\in\mathbf{R}$  and the graph of the solution is contained in a compact domain. According with [6] (theorem 3.2.5, p.141) we obtain a contradiction with the fact that  $\mathbf{x}$  admit a prolongation on the right or the left (also, can be applied the theorem of Chilingworth (1976), see theorem 1.0.3, p.7 in [3]).

In the sequel we study the asymptotic behaviour of the globally solutions of the  $\varepsilon$ -revised system.

Denote by **E** the set of equilibrium points of the  $\varepsilon$ -revised system (20) and by  $\Gamma$  the trajectory of a solution  $\mathbf{x} : \mathbf{R} \to \mathbf{R}^3$  of (20). By theory of differential equations (see [10] p. 174-176), the  $\omega$ -limit set and  $\alpha$ -limit set of  $\Gamma$  are:

$$\omega(\Gamma) = \{ \mathbf{y} \in \mathbf{R}^3 \ / \ \exists t_n \to \infty \text{ such that } \mathbf{x}(t_n) \to \mathbf{y} \},$$
 
$$\alpha(\Gamma) = \{ \mathbf{z} \in \mathbf{R}^3 \ / \ \exists t_n \to -\infty \text{ such that } \mathbf{x}(t_n) \to \mathbf{z} \}.$$

**Theorem 5.2** Let  $\mathbf{x}: \mathbf{R} \to \mathbf{R}^3$  be a solution of the  $\varepsilon$ -revised system with  $\varepsilon \neq 0$ . There exist the equilibrium points  $\mathbf{x}_m, \mathbf{x}_M \in \mathbf{R}^3$  of the system (20) such that  $\lim_{t \to -\infty} \mathbf{x}(t) = \mathbf{x}_M$  and  $\lim_{t \to \infty} \mathbf{x}(t) = \mathbf{x}_m$ .

**Proof** The theorem is proved in the following steps:

- (i)  $\alpha(\Gamma) \neq \emptyset$  and  $\omega(\Gamma) \neq \emptyset$ .
- (ii)  $\alpha(\Gamma) \cap \omega(\Gamma) \subset \mathbf{E}$ .
- (iii) The sets  $\alpha(\Gamma)$  and  $\omega(\Gamma)$  contains exactly one element.

Taking account into that each solution is bounded (hence it is contained in a compact domain) and applying theorem 1, p. 175 in [10], we obtain immediately the assertions (i).

(ii) For demonstration consider the case when  $\varepsilon > 0$ . Using the relation (29), we deduce that the function  $t \to \mathbf{x}^2(t)$  is a strictly decreasing function. Being bounded it follows that there exists  $\lim_{t\to\infty} \mathbf{x}^2(t) = L$  and L is finite.

For each  $\mathbf{y} \in \omega(\Gamma)$  there exists the sequence  $t_n \to \infty$  such that  $\mathbf{x}(t_n) \to \mathbf{y}$ . Then  $\mathbf{x}^2(t_n) \to \mathbf{y}^2$  and hence  $\mathbf{y}^2 = L$ .

By theorem 2, p.176 in [10], we have that the trajectory  $\Gamma_{\mathbf{y}}$  of the solution  $\mathbf{x}_{\mathbf{y}}$  which verifies the initial condition  $\mathbf{x}_{\mathbf{y}}(0) = \mathbf{y}$ , satisfies the relation  $\Gamma_{\mathbf{y}} \subset \omega(\Gamma)$ .

If we assume that  $\mathbf{y}$  is not an equilibrium point, then we deduce (using the relation (29)) that for t > 0 we have  $\mathbf{x}_{\mathbf{y}}^2(t) < L$  and this is in contradiction with the above result. Therefore, we have  $\omega(\Gamma) \subset \mathbf{E}$ .

Similarly, we prove that  $\alpha(\Gamma) \subset \mathbf{E}$ . Hence the assertion (ii) holds.

The case  $\varepsilon < 0$  is similar.

(iii) There exists a constant k such that the sets  $\alpha(\Gamma)$  and  $\omega(\Gamma)$  are included in the ellipsoid  $H(x^1, x^2, x^3) = k$ . By (ii), we deduce that  $\alpha(\Gamma)$  and  $\omega(\Gamma)$  are included in the set of equilibrium points which lies of the above ellipsoid. On the other hand, applying Proposition 4.2 and using the fact that  $\alpha(\Gamma)$  and  $\omega(\Gamma)$  are connected (see theorem 1, p.175 in [10]), we obtain that  $\alpha(\Gamma)$  and  $\omega(\Gamma)$  are formed by only one element.

**Remark 5.1** Using the relation (29) it is easy to observe that the following assertions hold:

(i) if 
$$\varepsilon > 0$$
 then  $\mathbf{x}_M^2 > \mathbf{x}_m^2$ ;  
(ii) if  $\varepsilon < 0$  then  $\mathbf{x}_M^2 < \mathbf{x}_m^2$ .

As an immediate consequence we obtain the following theorem.

**Theorem 5.3** If  $\varepsilon \neq 0$ , then for each solution  $\mathbf{x} : \mathbf{R} \to \mathbf{R}^3$  of the  $\varepsilon$ -revised system we have:

$$\begin{cases} \text{ if } t \to \infty \Rightarrow d(\mathbf{x}(t), \mathbf{E}) \to 0\\ \text{ if } t \to -\infty \Rightarrow d(\mathbf{x}(t), \mathbf{E}) \to 0. \end{cases}$$
(31)

**Remark 5.2** From Theorem 5.3 follows that the set  $\mathbf{E}$  of the equilibrium points is an attracting set (see definition 2, p.178 in [10]) and also is a reppeling set (see [3], p.34). Thus, the space  $\mathbf{R}^3$  is simultaneously a domain of attraction and a domain of repulsion of  $\mathbf{E}$ .

# 6 The Lyapunov stability of equilibrium points of the $\varepsilon$ -revised system in the case $\varepsilon > 0$

The stability of the point  $e_1 = (0,0,0)$ . We have the following results.

**Theorem 6.1** The equilibrium point  $e_1$  is Lyapunov stable.

**Proof** Let  $\gamma > 0$ ,  $t_0 \in \mathbf{R}$  and  $\mathbf{x}_0 \in \mathbf{R}^3$  such that  $|\mathbf{x}_0| < \gamma$ , where  $|\cdot|$  denotes the euclidian norm in  $\mathbf{R}^3$ . Denote by  $t \to \mathbf{x}(t, t_0, \mathbf{x}_0)$  the solution of the  $\varepsilon$ -revised system which verifies the initial condition  $\mathbf{x}(0, t_0, \mathbf{x}_0) = \mathbf{x}_0$ .

Using the relation  $|\mathbf{x}(t, t_0, \mathbf{x}_0)| = \sqrt{\mathbf{x}^2(t, t_0, \mathbf{x}_0)}$  and according with the relation (29), we observe that the function  $t \to \mathbf{x}(t, t_0, \mathbf{x}_0)$  is a decreasing function and hence we have:

$$|\mathbf{x}(t,t_0,\mathbf{x}_0)| \leq |\mathbf{x}_0| < \gamma \text{ for } t > t_0.$$

Then (see [4], p.22) we have that  $e_1$  is a Lyapunov stable equilibrium point.  $\Box$ 

**Remark 6.1** The equilibrium point  $e_1$  is not asymptotical stable.

Indeed, if a = b = c = 0 then the coordinates axis are formed from equilibrium points. If al least of one of the numbers a, b, c is non null, then:

$$\text{if}\, |\lambda| \to \infty \, \Rightarrow \, (\frac{a}{\lambda - a_1}, \frac{b}{\lambda - a_2}, \frac{c}{\lambda - a_3}) \to (0, 0, 0).$$

Hence, in all neighbourhood of  $e_1$  there exist an infinity of equilibrium points.

The stability of the point  $\overline{\mathbf{x}_0} = (-\frac{a}{a_1}, -\frac{b}{a_2}, -\frac{c}{a_3})$ . The equilibrium point  $\overline{\mathbf{x}_0}$  is an equilibrium point of the form  $e_2$  and it is obtained for  $\lambda = 0$ .

**Theorem 6.2** The equilibrium point  $\overline{\mathbf{x}_0}$  is Lyapunov stable.

**Proof** Using the relation (30) and the inequality  $0 < a_1 < a_2 < a_3$ , we deduce:

$$\frac{a_1}{2}|\mathbf{x} - \overline{\mathbf{x}_0}| \le H(\mathbf{x}) - H(\overline{\mathbf{x}_0}) \le \frac{a_3}{2}|\mathbf{x} - \overline{\mathbf{x}_0}|$$

For  $t_0 \in \mathbf{R}$  and  $\mathbf{x}_0 \in \mathbf{R}^3$  denote with  $\mathbf{x}(t, t_0, \mathbf{x}_0)$  a solution of  $\varepsilon$ -revised system which verifies the initial condition  $\mathbf{x}(0, t_0, \mathbf{x}_0) = \mathbf{x}_0$ .

Let  $\gamma > 0$  and  $\delta(\gamma) = \frac{2\gamma}{a_1}$ . Let  $\mathbf{x}_0 \in \mathbf{R}^3$  such that:

$$H(\mathbf{x}_0) - H(\overline{\mathbf{x}_0}) \le \delta(\gamma).$$

From the fact that H is a first integral we deduce that:

$$H(\mathbf{x}(t, t_0, \mathbf{x}_0) - H(\overline{\mathbf{x}_0}) = H(\mathbf{x}_0) - H(\overline{\mathbf{x}_0}).$$

Hence for all  $t \in \mathbf{R}$  the following inequality holds:

$$\frac{a_1}{2}|\mathbf{x} - \overline{\mathbf{x}_0}| \le \delta(\gamma)$$

and we obtain that  $\overline{\mathbf{x}_0}$  is Lyapunov stable.

**Remark 6.2** The stable equilibrium point  $\overline{\mathbf{x}_0}$  realizes the absolute minimum of the function H.

The unstability of equilibrium points of the form  $e_2$  with  $\lambda \in (0, a_1)$ . For the demonstration of this results we use the Theorem 6.3 and Lemma 6.1.

**Theorem 6.3** If  $\mathbf{x}_0 \in \mathbf{E}$  such that there exists  $\mathbf{y} \in \mathbf{E}$  with the properties:

(i) 
$$H(\mathbf{y}) = H(\mathbf{x}_0)$$
 and (ii)  $|\mathbf{y}| < |\mathbf{x}_0|$ 

then  $\mathbf{x}_0$  is an unstable equilibrium point.

**Proof** For  $k \in \mathbf{R}$  denote by  $\mathbf{E}_k = \{\mathbf{x} \in \mathbf{E} / H(x) = k\}$ . The set  $\mathbf{E}_{H(\mathbf{X}_0)}$  is finite (by Proposition 4.2). We denote:

$$\gamma_0 = \min\{|\mathbf{x} - \mathbf{x}_0|/\mathbf{x} \in E_{H(\mathbf{X}_0)} - \{\mathbf{x}_0\}\}\$$

Let  $\mathbf{z} \in \mathbf{R}^3$  such that  $H(\mathbf{z}) = H(\mathbf{x}_0)$  and  $|\mathbf{z}| < |\mathbf{x}_0|$ . Then:

$$\lim_{t \to \infty} \mathbf{x}(t, 0, \mathbf{z}) \in \mathbf{E}$$

and

if 
$$t > 0 \Rightarrow |\mathbf{x}(t, 0, \mathbf{z})| < |\mathbf{z}|$$

and we deduce that there exists  $t_z > 0$  such that:

$$|\mathbf{x}(t,0,\mathbf{z}) - \mathbf{x}_0| > rac{\gamma_0}{2}$$
 if  $t > t_z$ 

It follows that  $\mathbf{x}_0$  is unstable.

We assume that  $(a, b, c) \neq (0, 0, 0)$  and we introduce the notation:

$$e_{2\lambda}=(\frac{a}{\lambda-a_1},\frac{b}{\lambda-a_2},\frac{c}{\lambda-a_3}) \text{ for all } \lambda \in \mathbf{R}-\{a_1,a_2,a_3\}$$

- **Lemma 6.1** (i) If  $\sigma < \mu < a_1$ , then  $|e_{2\sigma}| < |e_{2\mu}|$ . (ii) If  $\sigma, \mu > 0$ ,  $(\frac{\mu}{\sigma})^2 > \frac{a_3}{a_1}$  and  $H(e_{2\sigma}) = H(e_{2\mu})$ , then  $|e_{2\sigma}| > |e_{2\mu}|$ . (iii) If  $0 < \sigma < a_1 < a_3 < \mu$  and  $H(e_{2\sigma}) = H(e_{2\mu})$ , then  $|e_{2\sigma}| > |e_{2\mu}|$ .

**Proof** (i) Consider the function  $g:(-\infty,a_1)\to \mathbf{R}$  given by:

$$g(\lambda) = (\frac{a}{\lambda - a_1})^2 + (\frac{b}{\lambda - a_2})^2 + (\frac{c}{\lambda - a_3})^2.$$

The derivative of the function g is:

$$g'(\lambda) = -\frac{2a^2}{(\lambda - a_1)^3} - \frac{2b^2}{(\lambda - a_2)^3} - \frac{2c^2}{(\lambda - a_3)^3}$$

We observe that  $g'(\lambda) > 0$  and we obtain that g is a strictly increasing function. We have:

$$g(\sigma) = |e_{2\sigma}|^2, \ g(\mu) = |e_{2\mu}|^2$$

and we obtain the desired result.

(ii) From hypothesis  $H(e_{2\sigma}) = H(e_{2\mu})$  follows that there exists a constant q > 0with the following properties:

$$\frac{1}{a_1} \frac{a^2}{(\sigma - a_1)^2} + \frac{1}{a_2} \frac{b^2}{(\sigma - a_2)^2} + \frac{1}{a_3} \frac{c^2}{(\sigma - a_3)^2} = \frac{q}{\sigma^2}$$

$$\frac{1}{a_1} \frac{a^2}{(\mu - a_1)^2} + \frac{1}{a_2} \frac{b^2}{(\mu - a_2)^2} + \frac{1}{a_3} \frac{c^2}{(\mu - a_3)^2} = \frac{q}{\mu^2}$$

Using  $a_1 < a_2 < a_3$ , we obtain the inequalities:

$$|e_{2\sigma}|^2 > \frac{a_1q}{\sigma}, |e_{2\mu}|^2 < \frac{a_3q}{\mu}$$

and we observe that the assertion (ii) holds.

(iii) This assertion follows immediately from (ii).

**Theorem 6.4** The equilibrium point  $e_{2\lambda}$  with  $0 < \lambda < a_1$  is unstable.

**Proof** Consider the function  $h: (-\infty, a_1) \bigcup (a_3, \infty) \to \mathbf{R}$  given by:

$$h(\sigma) = H(e_{2\sigma}).$$

Using the relation (30) for H, we find:

$$h(\sigma) = \frac{\sigma^2}{2} \left[ \frac{a^2}{a_1(\sigma - a_1)^2} + \frac{b^2}{a_2(\sigma - a_2)^2} + \frac{c^2}{a_3(\sigma - a_3)^2} \right] - \frac{1}{2} \left( \frac{a^2}{a_1} + \frac{b^2}{a_2} + \frac{c^2}{a_3} \right)$$

The function h have the following properties:

- 0 is an absolute minimum point.
- $\lim_{\sigma \to -\infty} h(\sigma) = \lim_{\sigma \to \infty} h(\sigma) = 0.$
- $\lim_{\sigma \to a_1} h(\sigma) = \lim_{\sigma \to a_3} h(\sigma) = \infty$ .
- h is strictly decreasing on  $(-\infty, 0)$ , strictly increasing on  $(0, a_1)$  and strictly decreasing on  $(a_3, \infty)$ .

The demonstrations divided on three cases.

(I) Assume that  $h(\lambda) < 0$ . In this situation there exists  $\sigma < 0 < \lambda < a_1$  such that  $h(\lambda) = h(\sigma)$  and imply  $H(e_{2\lambda}) = H(e_{2\sigma})$ . Hence the equilibrium points  $e_{2\lambda}$  and  $e_{2\sigma}$  belong to same ellipsoid.

On the other hand, by Lemma 6.1 (ii), follows  $|e_{2\sigma}| < |e_{2\lambda}|$ . Applying now Theorem 6.3, deduce that  $e_{2\lambda}$  is an unstable equilibrium point.

- (II) Assume that  $h(\lambda) = 0$  we have  $H(e_{2\lambda}) = H(0,0,0)$  and it is clearly that  $|(0,0,0)| < |e_{2\lambda}|$ . By Theorem 6.3 we find the desired result.
- (III) Assume that  $h(\lambda) > 0$ . Then there exists  $\sigma > a_3$  such that  $h(\lambda) = h(\sigma)$  and hence  $H(e_{2\lambda}) = H(e_{2\sigma})$ . Applying Lemma 6.1 (iii) follows  $|e_{2\lambda}| > |e_{2\sigma}|$  and by Theorem 6.3 we deduce that  $e_{2\lambda}$  is unstable.

The stability of equilibrium points of the form  $e_2$  with  $\lambda < 0$ .

**Theorem 6.5** The equilibrium points of the form  $e_2$  with  $\lambda < 0$  are Lyapunov stables.

**Proof** Let  $\lambda < 0$  and the equilibrium point  $\mathbf{x}_0 = (\frac{a}{\lambda - a_1}, \frac{b}{\lambda - a_2}, \frac{c}{\lambda - a_3})$  of the form  $e_2$ . It is well-known that the study of stability of  $\mathbf{x}_0$  in the Lyapunov sense is equivalent with the study of stability of the null solution (0,0,0) for the differential system obtained from the  $\varepsilon$ -revised system by transformation of variables:

$$z = x - x_0$$

The system obtained in this manner is called the **perturbed**  $\varepsilon$ -revised system.

Consider the function  $K: \mathbf{R}^3 \to \mathbf{R}$  given by

$$K(\mathbf{z}) = \frac{1}{2}\mathbf{z} \cdot \mathbf{I}^{-1}\mathbf{z} - \frac{\lambda}{2}\mathbf{z}^2$$

Since the tensor  $\mathbf{I}^{-1}$  is strictly positive definite and  $\lambda < 0$  we obtain that K is a quadratic form strictly positive definite.

Next we prove that if  $\mathbf{z}: \mathbf{R} \to \mathbf{R}^3$  is a solution for the perturbed  $\varepsilon$ -revised system, then:

$$\frac{d}{dt}K(\mathbf{z}(t)) < 0$$

By a direct computation and taking account into the relations (27) we have:

$$K(\mathbf{z}) = H(\mathbf{x}) - \lambda C(\mathbf{x}) - \frac{1}{2}\mathbf{x}_0 \cdot \mathbf{I}^{-1}\mathbf{x}_0 - \mathbf{a} \cdot \mathbf{x}_0 + \frac{\lambda}{2}\mathbf{x}_0^2$$

Applying now Theorem 4.2, we obtain:

$$\frac{d}{dt}K(\mathbf{z}(t)) = \varepsilon \lambda(\mathbf{x} \times \mathbf{m}(\mathbf{x}))^2$$

and follows that  $\frac{d}{dt}K(\mathbf{z}(t)) < 0$ , since  $\varepsilon > 0$ ,  $\lambda < 0$ .

It is easy to see that  $K^*(t) = K(\mathbf{z}(t))$  is a strictly decreasing function. By theorem 1.1, p.21 in the paper [4] we deduce that  $\mathbf{x}_0$  is Lyapunov stable.

## Conclusion - the stability problem for the Hamilton Poisson system (12) versus the $\varepsilon$ - revised system (20) with $\varepsilon > 0$

Concerning to the equilibrium points of the system (12) are established the following results (see, theorem 1.1, [12]):

- (1)  $e_1$  is Lyapunov stable;
- (2)  $e_2$  are Lyapunov stables for  $\lambda \in (-\infty, a_1) \cup (a_3, \infty)$ ;
- (3)  $e_3$  are Lyapunov stables;
- (4)  $e_4$  are unstables;
- (5)  $e_5$  are Lyapunov stables.

By Remark 4.1 (see, [12]), there exist cases for which the problem to decide the nonlinear stability or unstability are not discussed.

For the stability of equilibrium points of the  $\varepsilon$ - revised system (20) with  $\varepsilon > 0$  have proved the following assertions:

- (1)  $e_1$  is Lyapunov stable;
- (2) the equilibrium points of the form  $e_2$  with  $\lambda \leq 0$  are Lyapunov stables;
- (3) the equilibrium points of the form  $e_2$  with  $0 < \lambda < a_1$  are unstables.

#### References

- [1]. R. Abraham, J.E. Marsden, Foundations of Mechanics. Second Edition. Addison-Wesley, 1978.
- [2]. D. Fish, Dissipative perturbation of 3D Hamiltonian systems. Metriplectic systems. Preprint, arXiv:math-ph/0506047, v1,2005.
- [3]. J. Guckenheimer, P. Holmes, Nonlinear oscilations, dynamical systems and bifurcations of vector fields. Springer- Verlag, New York, 1990.
- [4]. A. Halanay, Teoria calitativă a ecuațiilor diferențiale. Ed. Academiei, București, 1963.
- [5] Gh. Ivan, D. Opriş, Dynamical systems on Leibniz algebroids. Differential Geometry Dynamical Systems, 8( 2006 ), 127 137.
- [6]. St. Mirică, Ecuații diferențiale și cu derivate parțiale I. Litografia Univ. București, 1989.
- [7] J.E. Marsden, Lectures on Mechanics. London Mathematical Society, Lectures Note Series, vol.174, 2 nd edition, Cambridge University Press, 1992.
- [8] P.J. Morrison, A paradigm for joined Hamiltonian and dissipative systems. Physica, 18D (1986), 410 419.
- [9]. J.- P. Ortega, V. Planas -Bielsa, Dynamics on Leibniz manifolds. Preprint, arXiv:math. DS/0309263,2003.
- [10]. L. Perko, Differential equations and dynamical systems. Springer- Verlag, New York, 1991.
- [11]. M. Puta, Hamiltonian mechanics and geometric quantization. Mathematics and its Applications, vol. 260, Kluwer, 1993.
- [12]. M. Puta, D. Comănescu, On the rigid body with three linear controls. Analele Univ. din Timișoara, Seria Matematică Informatică, vol. 35 (1), 1997, p.-.
- [13]. M. Puta, D. Comănescu, S. Chirici, Elemente de mecanică hamiltoniană. Ed. Mirton, Timișoara, 2004.

Seminarul de Geometrie și Topologie

West University of Timişoara

Bd-ul V. Pârvan no.4, 300223, Timişoara

Romania

E-mail: comanescu@math.uvt.ro, ivan@math.uvt.ro and mihai31ro@yahoo.com