Defensive k-alliances in graphs

Juan A. Rodríguez-Velázquez*and Ismael G. Yero[†] Department of Computer Engineering and Mathematics Rovira i Virgili University of Tarragona Av. Països Catalans 26, 43007 Tarragona, Spain José M. Sigarreta[‡] Departamento de Matemáticas Universidad Carlos III de Madrid

Avda. de la Universidad 30, 28911 Leganés (Madrid), Spain

Abstract

Let G = (V, E) be a simple graph of order n and degree sequence $\delta_1 \geq \delta_2 \geq \cdots \geq \delta_n$. For a nonempty set $X \subseteq V$, and a vertex $v \in V$, $\delta_X(v)$ denotes the number of neighbors v has in X. A nonempty set $S \subseteq V$ is a defensive k-alliance in G = (V, E) if $\delta_S(v) \geq \delta_{\bar{S}}(v) + k, \forall v \in S$. The defensive k-alliance number of G, denoted by $a_k(G)$, is defined as the minimum cardinality of a defensive k-alliance in G. We study the mathematical properties of $a_k(G)$. We show that $\left\lceil \frac{\delta_n + k + 2}{2} \right\rceil \leq a_k(G) \leq n - \left\lfloor \frac{\delta_n - k}{2} \right\rfloor$ and $a_k(G) \geq \left\lceil \frac{n(\mu + k + 1)}{n + \mu} \right\rceil$, where μ is the algebraic connectivity of G and $k \in \{-\delta_n, \ldots, \delta_1\}$. Moreover, we show that for every $k, r \in \mathbb{Z}$ such that $-\delta_n \leq k \leq \delta_1$ and $0 \leq r \leq \frac{k + \delta_n}{2}$, $a_{k-2r}(G) + r \leq a_k(G)$ and, as a consequence, we show that for every $k \in \{-\delta_n, \ldots, 0\}, a_k(G) \leq \left\lceil \frac{n + k + 1}{2} \right\rceil$. In the case of the line graph $\mathcal{L}(G)$ of a simple graph G, we obtain bounds on $a_k(\mathcal{L}(G))$ and, as a consequence of the study, we show that for any δ -regular graph, $\delta > 0$,

^{*}juanalberto.rodriguez@urv.cat

 $^{^{\}dagger} is mael.gonzalez @urv.cat$

[‡]josemaria.sigarreta@uc3m.es

and for every $k \in \{2(1-\delta), \ldots, 0\}, a_k(\mathcal{L}(G)) = \delta + \lceil \frac{k}{2} \rceil$. Moreover, for any (δ_1, δ_2) -semiregular bipartite graph $G, \delta_1 > \delta_2$, and for every $k \in \{2 - \delta_1 - \delta_2, \ldots, \delta_1 - \delta_2\}, a_k(\mathcal{L}(G)) = \lceil \frac{\delta_1 + \delta_2 + k}{2} \rceil$.

Keywords: Alliances in graphs; Algebraic connectivity; Graph eigenvalues; Line graph.

AMS Subject Classification numbers: 05C69; 05A20; 05C50

1 Introduction

The mathematical properties of alliances in graphs were first studied by P. Kristiansen, S. M. Hedetniemi and S. T. Hedetniemi [7]. They proposed different types of alliances: namely, defensive alliances [5, 6, 7, 11], offensive alliances [2, 9, 10] and dual alliances or powerful alliances [1]. A generalization of these alliances called k-alliances was presented by K. H. Shafique and R. D. Dutton [12, 13].

In this paper, we study the mathematical properties of defensive kalliances. We begin by stating the terminology used. Throughout this article, G = (V, E) denotes a simple graph of order |V| = n and size |E| = m. We denote two adjacent vertices u and v by $u \sim v$. For a nonempty set $X \subseteq V$, and a vertex $v \in V$, $N_X(v)$ denotes the set of neighbors v has in $X: N_X(v) := \{u \in X : u \sim v\}$ and the degree of v in X will be denoted by $\delta_X(v) = |N_X(v)|$. We denote the degree of a vertex $v_i \in V$ by $\delta(v_i)$ (or by δ_i for short) and the degree sequence of G by $\delta_1 \geq \delta_2 \geq \cdots \geq \delta_n$. The subgraph induced by $S \subset V$ will be denoted by $\langle S \rangle$ and the complement of the set Sin V will be denoted by \bar{S} .

A nonempty set $S \subseteq V$ is a *defensive* k-alliance in $G = (V, E), k \in \{-\delta_1, \ldots, \delta_1\}$, if for every $v \in S$,

$$\delta_S(v) \ge \delta_{\bar{S}}(v) + k. \tag{1}$$

A vertex $v \in S$ is said to be *k*-satisfied by the set S if (1) holds. Notice that (1) is equivalent to

$$\delta(v) \ge 2\delta_{\bar{S}}(v) + k. \tag{2}$$

A defensive (-1)-alliance is a *defensive alliance* and a defensive 0-alliance is a *strong defensive alliance* as defined in [7]. A defensive 0-alliance is also known as a *cohesive set* [14].

Defensive alliances are the mathematical model of web communities. Adopting the definition of Web community proposed recently by Flake, Lawrence, and Giles [3], "a Web community is a set of web pages having more hyperlinks (in either direction) to members of the set than to non-members".

2 Defensive *k*-alliance number

The defensive k-alliance number of G, denoted by $a_k(G)$, is defined as the minimum cardinality of a defensive k-alliance in G. Notice that

$$a_{k+1}(G) \ge a_k(G). \tag{3}$$

The defensive (-1)-alliance number of G is known as the alliance number of G and the defensive 0-alliance number is known as the strong alliance number, [7, 5, 6]. For instance, in the case of the 3-cube graph, $G = Q_3$, every set composed by two adjacent vertices is a defensive alliance of minimum cardinality and every set composed by four vertices whose induced subgraph is isomorphic to the cycle C_4 is a strong defensive alliance of minimum cardinality. Thus, $a_{-1}(Q_3) = 2$ and $a_0(Q_3) = 4$.

Notice that if every vertex of G has even degree and k is odd, k = 2l - 1, then every defensive (2l - 1)-alliance in G is a defensive (2l)-alliance. Hence, in such a case, $a_{2l-1}(G) = a_{2l}(G)$. Analogously, if every vertex of G has odd degree and k is even, k = 2l, then every defensive (2l)-alliance in G is a defensive (2l + 1)-alliance. Hence, in such a case, $a_{2l}(G) = a_{2l+1}(G)$.

For some graphs, there are some values of $k \in \{-\delta_1, \ldots, \delta_1\}$, such that defensive k-alliances do not exist. For instance, for $k \geq 2$ in the case of the star graph S_n , defensive k-alliances do not exist. By (2) we conclude that, in any graph, there are defensive k-alliances for all $k \in \{-\delta_1, \ldots, \delta_n\}$. For instance, a defensive (δ_n) -alliance in G = (V, E) is V. Moreover, if $v \in V$ is a vertex of minimum degree, $\delta(v) = \delta_n$, then $S = \{v\}$ is a defensive kalliance for every $k \leq -\delta_n$. As $a_k(G) = 1$ for $k \leq -\delta_n$, hereafter we only will consider the cases $-\delta_n \leq k \leq \delta_1$. Moreover, the bounds showed in this paper on $a_k(G)$, for $\delta_n \leq k \leq \delta_1$, are obtained by supposing that the graph G contains defensive k-alliances.

It was shown in [7] that for any graph G of order n and minimum degree δ_n ,

$$a_{-1}(G) \le n - \left\lceil \frac{\delta_n}{2} \right\rceil$$
 and $a_0(G) \le n - \left\lfloor \frac{\delta_n}{2} \right\rfloor$.

Here we generalize the previous result to defensive k-alliances and we obtain lower bounds.

Theorem 1. For every $k \in \{-\delta_n, \ldots, \delta_1\}$,

$$\left\lceil \frac{\delta_n + k + 2}{2} \right\rceil \le a_k(G) \le n - \left\lfloor \frac{\delta_n - k}{2} \right\rfloor.$$

Proof. Let $X \subseteq V$ be a defensive k-alliance in G. In this case, for every $v \in X$ we have

$$\delta(v) = \delta_X(v) + \delta_{\bar{X}}(v)$$
$$\delta(v) \le \delta_X(v) + \frac{\delta(v) - k}{2}$$
$$\frac{\delta(v) + k}{2} \le \delta_X(v) \le |X| - 1$$
$$\frac{\delta_n + k + 2}{2} \le |X|.$$

Hence, the lower bound follows.

On the other hand, if $X \subseteq V$ is a defensive k-alliance in G for $\delta_n \leq k \leq \delta_1$, then $a_k(G) \leq |X| \leq n \leq n - \lfloor \frac{\delta_n - k}{2} \rfloor$. Suppose $-\delta_n \leq k \leq \delta_n$. Let $S \subseteq V$ be a set of cardinality $n - \lfloor \frac{\delta_n - k}{2} \rfloor$. For every vertex $v \in S$ we have $\frac{\delta(v) - k}{2} \geq \lfloor \frac{\delta_n - k}{2} \rfloor \geq \delta_{\bar{S}}(v)$. Hence, S is a defensive k-alliance and $a_k(G) \leq |S| = n - \lfloor \frac{\delta_n - k}{2} \rfloor$.

We denote by K_n the complete graph of order n.

Corollary 2. For every $k \in \{1 - n, \dots, n - 1\}$, $a_k(K_n) = \left\lceil \frac{n + k + 1}{2} \right\rceil$. The sense 2. For every $k \in \{7, \dots, n - 1\}$, $a_k(K_n) = \left\lceil \frac{n + k + 1}{2} \right\rceil$.

Theorem 3. For every $k, r \in \mathbb{Z}$ such that $-\delta_n \leq k \leq \delta_1$ and $0 \leq r \leq \frac{k+\delta_n}{2}$,

 $a_{k-2r}(G) + r \le a_k(G).$

Proof. Let $S \subset V$ be a defensive k-alliance of minimum cardinality in G. By Theorem 1, $\frac{\delta_n+k+2}{2} \leq |S|$, then we can take $X \subset S$ such that |X| = r. Hence, for every $v \in Y = S - X$,

$$\delta_Y(v) = \delta_S(v) - \delta_X(v)$$

$$\geq \delta_{\bar{S}}(v) + k - \delta_X(v)$$

$$= \delta_{\bar{Y}}(v) + k - 2\delta_X(v)$$

$$\geq \delta_{\bar{Y}}(v) + k - 2r.$$

Therefore, Y is a defensive (k - 2r)-alliance in G and, as a consequence, $a_{k-2r}(G) \leq a_k(G) - r$.

Notice that, according to the result in Corollary 2, the bound for $a_k(G)$ in Theorem 3 is attained for the complete graph K_n for every n, k, r with its respective restrictions. From the above theorem we derive some interesting consequences.

Corollary 4. Let $t \in \mathbb{Z}$.

- If $\frac{1-\delta_n}{2} \le t \le \frac{\delta_1-1}{2}$, then $a_{2t-1}(G) + 1 \le a_{2t+1}(G)$.
- If $\frac{2-\delta_n}{2} \le t \le \frac{\delta_1}{2}$, then $a_{2(t-1)}(G) + 1 \le a_{2t}(G)$.

Corollary 5. For every $k \in \{0, \ldots, \delta_n\}$,

- if k is even, then $a_{-k}(G) + \frac{k}{2} \le a_0(G) \le a_k(G) \frac{k}{2}$,
- if k is odd, then $a_{-k}(G) + \frac{k-1}{2} \le a_{-1}(G) \le a_k(G) \frac{k+1}{2}$.

It was shown in [5, 7] that for any graph G of order n,

$$a_{-1}(G) \le \left\lceil \frac{n}{2} \right\rceil$$
 and $a_0(G) \le \left\lfloor \frac{n}{2} \right\rfloor + 1.$ (4)

By Corollary 5 and (4) we obtain the following result.

Theorem 6. For every $k \in \{-\delta_n, \ldots, 0\}$, $a_k(G) \leq \left\lceil \frac{n+k+1}{2} \right\rceil$.

Notice that the above bound is attained, for instance, for the complete graph $G = K_n$.

3 Algebraic connectivity and defensive k-alliance number

It is well-known that the second smallest Laplacian eigenvalue of a graph is probably the most important information contained in the Laplacian spectrum. This eigenvalue, frequently called *algebraic connectivity*, is related to several important graph invariants and imposes reasonably good bounds on the values of several parameters of graphs which are very hard to compute. The algebraic connectivity of G, μ , satisfies the following equality shown by Fiedler [4] on weighted graphs

$$\mu = 2n \min\left\{\frac{\sum_{v_i \sim v_j} (w_i - w_j)^2}{\sum_{v_i \in V} \sum_{v_j \in V} (w_i - w_j)^2} : w \neq \alpha \mathbf{j} \text{ for } \alpha \in \mathbb{R}\right\}, \quad (5)$$

where $V = \{v_1, v_2, ..., v_n\}$, $\mathbf{j} = (1, 1, ..., 1)$ and $w \in \mathbb{R}^n$.

The following theorem shows the relationship between the algebraic connectivity of a graph and its defensive k-alliance number.

Theorem 7. For any connected graph G and for every $k \in \{-\delta_n, \ldots, \delta_1\}$,

$$a_k(G) \ge \left\lceil \frac{n(\mu+k+1)}{n+\mu} \right\rceil.$$

Proof. If S denotes a defensive k-alliance in G, then

$$\delta_{\bar{S}}(v) + k \le |S| - 1, \quad \forall v \in S.$$
(6)

By (5), taking $w \in \mathbb{R}^n$ defined as

$$w_i = \begin{cases} 1 & \text{if } v_i \in S; \\ 0 & \text{otherwise,} \end{cases}$$

we have

$$\mu \le \frac{n \sum_{v \in S} \delta_{\bar{S}}(v)}{|S|(n-|S|)}.$$
(7)

Thus, (6) and (7) lead to

$$\mu \le \frac{n(|S| - k - 1)}{n - |S|}.$$
(8)

Therefore, solving (8) for |S|, and considering that it is an integer, we obtain the bound on $a_k(G)$.

The above bound is sharp as we can see in the following example. As the algebraic connectivity of the complete graph $G = K_n$ is $\mu = n$, the above theorem gives the exact value of $a_k(K_n) = \left\lceil \frac{n+k+1}{2} \right\rceil$.

Theorem 8. For any connected graph G and for every $k \in \{-\delta_n, \ldots, \delta_1\}$,

$$a_k(G) \ge \left\lceil \frac{n(\mu - \lfloor \frac{\delta_1 - k}{2} \rfloor)}{\mu} \right\rceil.$$

Proof. If S denotes a defensive k-alliance in G, then $\delta_1 \ge \delta(v) \ge 2\delta_{\bar{S}}(v) + k$, $\forall v \in S$. Therefore,

$$\left\lfloor \frac{\delta_1 - k}{2} \right\rfloor \ge \delta_{\bar{S}}(v), \quad \forall v \in S.$$
(9)

Hence, by (7) and (9) the result follows.

The bound is attained for every k in the case of the complete graph $G = K_n$.

The reader is referred to [8] for more details on the spectral study of offensive alliances and dual alliances.

4 Defensive *k*-alliance number and line graph

Hereafter, we denote by $\mathcal{L}(G) = (V_l, E_l)$ the line graph of a simple graph G. The degree of the vertex $e = \{u, v\} \in V_l$ is $\delta(e) = \delta(u) + \delta(v) - 2$. If the degree sequence of G is $\delta_1 \geq \delta_2 \geq \cdots \geq \delta_n$, then the maximum degree of $\mathcal{L}(G)$, denoted by Δ_l , is bounded by

$$\Delta_l \le \delta_1 + \delta_2 - 2 \tag{10}$$

and the minimum degree of $\mathcal{L}(G)$, denoted by δ_l , is bounded by

$$\delta_l \ge \delta_n + \delta_{n-1} - 2. \tag{11}$$

In this section we obtain some results on $a_k(\mathcal{L}(G))$ in terms of the degree sequence of G.

Theorem 9. For any simple graph G of maximum degree δ_1 , and for every $k \in \{2(1 - \delta_1), \ldots, 0\},\$

$$a_k(\mathcal{L}(G)) \le \delta_1 + \left\lceil \frac{k}{2} \right\rceil$$

7

Proof. Suppose k is even. Let $v \in V(G)$ be a vertex of maximum degree in G and let $S_v = \{e \in E : v \in e\}$. Let $Y_k \subset S_v$ such that $|Y_k| = -\frac{k}{2}$ and let $X_k = S_v - Y_k$. Thus, $\langle S_v \rangle \cong K_{\delta_1}$ and, as a consequence,

$$\delta_{X_k}(e) = \delta_1 - 1 + \frac{k}{2} \ge \delta_2 - 1 + \frac{k}{2} \ge \delta_{\bar{X_k}}(e) + k, \quad \forall e \in X_k.$$

Hence, $X_k \subset V_l$ is a defensive k-alliance in $\mathcal{L}(G)$. So, for k even we have $a_k(\mathcal{L}(G)) \leq \delta_1 + \lceil \frac{k}{2} \rceil$. Moreover, if k is odd, then $a_k(\mathcal{L}(G)) \leq a_{k+1}(\mathcal{L}(G)) \leq \delta_1 + \lceil \frac{k+1}{2} \rceil = \delta_1 + \lceil \frac{k}{2} \rceil$. \Box

One advantage of applying the above bound is that it requires only little information about the graph G; just the maximum degree. The above bound is tight, as we will see below, for any δ -regular graph, and $k \in \{2(1 - \delta), \ldots, 0\}$, $a_k(\mathcal{L}(G)) = \delta + \lceil \frac{k}{2} \rceil$. Even so, we can improve this bound for the case of nonregular graphs. The drawback of such improvement is that we need to know more about G.

Theorem 10. Let G be a simple graph, whose degree sequence is $\delta_1 \geq \delta_2 \geq \cdots \geq \delta_n$. Let $v \in V$ such that $\delta(v) = \delta_1$, let $\delta_v = \max_{u:u \sim v} \{\delta(u)\}$ and let $\delta_* = \min_{v:\delta(v)=\delta_1} \{\delta_v\}$. For every $k \in \{2 - \delta_* - \delta_1, \dots, \delta_1 - \delta_*\}$,

$$a_k(\mathcal{L}(G)) \le \left\lceil \frac{\delta_1 + \delta_* + k}{2} \right\rceil.$$

Moreover, for every $k \in \{2 - \delta_1 - \delta_2, \dots, \delta_1 + \delta_2 - 2\},\$

$$\left\lceil \frac{\delta_n + \delta_{n-1} + k}{2} \right\rceil \le a_k(\mathcal{L}(G))$$

Proof. Let $v \in V$ a vertex of maximum degree $\delta(v) = \delta_1$ such that v is adjacent to a vertex of degree δ_* . Let $S_v = \{e \in E : v \in e\}$. Suppose $\delta_1 + \delta_* + k$ is even. Therefore, taking $S \subset S_v$ such that $|S| = \frac{\delta_1 + \delta_* + k}{2}$, we obtain $\langle S \rangle \cong K_{\frac{\delta_1 + \delta_* + k}{2}}$. Thus, $\forall e \in S$,

$$\delta_S(e) - k = \frac{\delta_1 + \delta_* + k}{2} - 1 - k \ge \delta_1 + \delta_* - 1 - \frac{\delta_1 + \delta_* + k}{2}$$

On the other hand, as $\delta(e) = \delta_S(e) + \delta_{\bar{S}}(e)$, we have

$$\delta_1 + \delta_* - 2 \ge \frac{\delta_1 + \delta_* + k}{2} - 1 + \delta_{\bar{S}}(e) \Leftrightarrow \delta_1 + \delta_* - 1 - \frac{\delta_1 + \delta_* + k}{2} \ge \delta_{\bar{S}}(e).$$

So, $\delta_S(e) - k \geq \delta_{\bar{S}}(e)$ and S is a defensive k-alliance in $\mathcal{L}(G)$ and, as a consequence, $a_k(\mathcal{L}(G)) \leq \left\lceil \frac{\delta_1 + \delta_* + k}{2} \right\rceil$. If $\delta_1 + \delta_* + k$ is odd, then $\delta_1 - \delta_* > k$. Therefore, $a_k(\mathcal{L}(G)) \leq a_{k+1}(\mathcal{L}(G)) \leq \left\lceil \frac{\delta_1 + \delta_* + k + 1}{2} \right\rceil = \left\lceil \frac{\delta_1 + \delta_* + k}{2} \right\rceil$.

The lower bound follows from Theorem 1 and (11)

Corollary 11. For any δ -regular graph, $\delta > 0$, and for every $k \in \{2(1 - \delta)\}$ δ), ..., 0},

$$a_k(\mathcal{L}(G)) = \delta + \left\lceil \frac{k}{2} \right\rceil.$$

We recall that a graph G = (V, E) is a $(\delta_1, \delta_2$ -semiregular bipartite graph if the set V can be partitioned into two disjoint subsets V_1, V_2 such that if $u \sim v$ then $u \in V_1$ and $v \in V_2$ and also, $\delta(v) = \delta_1$ for every $v \in V_1$ and $\delta(v) = \delta_2$ for every $v \in V_2$.

Corollary 12. For any (δ_1, δ_2) -semiregular bipartite graph $G, \delta_1 > \delta_2$, and for every $k \in \{2 - \delta_1 - \delta_2, \dots, \delta_1 - \delta_2\},\$

$$a_k(\mathcal{L}(G)) = \left\lceil \frac{\delta_1 + \delta_2 + k}{2} \right\rceil.$$

We should point out that from the results obtained in the other sections of this article on $a_k(G)$, we can derive some new results on $a_k(\mathcal{L}(G))$. The reader is referred to [11] for more details on $a_{-1}(\mathcal{L}(G))$ and $a_0(\mathcal{L}(G))$.

References

- [1] R. Brigham; R. Dutton and S.Hedetniemi, A sharp lower bound on the powerful alliance number of $C_m \times C_n$. Congr. Numer. 167 (2004) 57-63.
- [2] O. Favaron, G. Fricke, W. Goddard, S. Hedetniemi, S. T. Hedetniemi, P. Kristiansen, R. C. Laskar, R. D. Skaggs, Offensive alliances in graphs. Discuss. Math. Graphs Theory 24 (2) (2004) 263-275.
- [3] G. W. Flake, S. Lawrence, and C. L. Giles, Efficient Identification of Web Communities. In Proceedings of the 6th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining (KDD-2000), (2000) 150-160.

- [4] M. Fiedler, A property of eigenvectors of nonnegative symmetric matrices and its application to graph theory. *Czechoslovak Math. J.* 25 (100) (1975) 619-633.
- [5] G. H. Fricke, L. M. Lawson, T. W. Haynes, S. M. Hedetniemi and S. T. Hedetniemi, A Note on Defensive Alliances in Graphs. *Bull. Inst. Combin. Appl.* 38 (2003) 37-41.
- [6] T. W. Haynes, S. T. Hedetniemi and M. A. Henning, Global defensive alliances in graphs. *Electron. J. Combin.* 10 (2003) 139-146.
- [7] P. Kristiansen, S. M. Hedetniemi and S. T. Hedetniemi, Alliances in graphs. J. Combin. Math. Combin. Comput. 48 (2004) 157-177.
- [8] J. A. Rodríguez and J. M. Sigarreta, Spectral study of alliances in graphs. Discuss. Math. Graphs Theory 27 (1) (2007) 143-157.
- [9] J. A. Rodríguez and J. M. Sigarreta, Offensive alliances in cubic graphs. Int. Math. Forum 1 (36) (2006) 1773-1782.
- [10] J. A. Rodríguez-Velázquez and J. M. Sigarreta, Global offensive alliances in graphs. *Electron. Notes Discrete Math.* 25 (2006) 157-164.
- [11] J. M. Sigarreta and J. A. Rodríguez, On defensive alliances and line graphs. Appl. Math. Lett. 19 (12) (2006) 1345-1350.
- [12] K. H. Shafique and R. D. Dutton, Maximum alliance-free and minimum alliance-cover sets. *Congr. Numer.* 162 (2003) 139-146.
- [13] K. H. Shafique and R. Dutton, A tight bound on the cardinalities of maximum alliance-free and minimum alliance-cover sets. J. Combin. Math. Combin. Comput. 56 (2006) 139-145.
- [14] K. H. Shafique and R. D. Dutton, On satisfactory partitioning of graphs. Congr. Numer. 154 (2002) 183-194.