

# PRINCIPAL GROUPOID $C^*$ -ALGEBRAS WITH BOUNDED TRACE

LISA ORLOFF CLARK AND ASTRID AN HUEF

ABSTRACT. Suppose  $G$  is a second countable, locally compact, Hausdorff, principal groupoid with a fixed left Haar system. We define a notion of integrability for groupoids, and show  $G$  is integrable if and only if the groupoid  $C^*$ -algebra  $C^*(G)$  has bounded trace.

## 1. INTRODUCTION

Let  $H$  be a locally compact, Hausdorff group acting continuously on a locally compact, Hausdorff space  $X$ , so that  $(H, X)$  is a transformation group. A lovely theorem of Green says that if  $H$  acts freely on  $X$ , then the associated transformation-group  $C^*$ -algebra  $C_0(X) \rtimes H$  has continuous trace if and only if the action of  $H$  on  $X$  is proper [5, Theorem 17]. Muhly and Williams defined a notion of proper groupoid, and proved that for principal groupoids  $G$ , the groupoid  $C^*$ -algebra  $C^*(G)$  has continuous trace if and only if the groupoid is proper [8, Theorem 2.3]. Of course, when  $G = H \times X$  is the transformation-group groupoid, then  $G$  is proper if and only if the action of  $H$  on  $X$  is proper.

In [13] Rieffel introduced a notion of an integrable action of a group  $H$  on a  $C^*$ -algebra  $A$ . This notion of integrability for  $A = C_0(X)$  turned out to characterize when  $C_0(X) \rtimes H$ , arising from a free action of  $H$  on  $X$ , has bounded trace [6, Theorem 4.8]. In this paper we define a notion of integrability for groupoids (see Definition 3.1) which, when  $G = H \times X$  is the transformation-group groupoid, reduces to an integrable action of  $H$  on  $X$  (see Example 3.3). We then prove that for principal groupoids  $G$ ,  $C^*(G)$  has bounded trace if and only if the groupoid is integrable (see Theorem 4.4). This theorem is thus very much in the spirit of [8, Theorem 2.4], [4, Theorem 7.9], [4, Theorem 4.1] (see also [3, Corollary 5.9]) and [4, Theorem 5.3], which characterize when principal-groupoid  $C^*$ -algebras are, respectively continuous-trace, Fell, CCR and GCR  $C^*$ -algebras. The key technical tools used to prove Theorem 4.4 are first, a homeomorphism of the spectrum of  $C^*(G)$  onto the orbit space [4, Proposition 5.1], and second, a generalisation to groupoids of the notion of  $k$ -times convergence in the orbit space of a transformation group from [1].

## 2. PRELIMINARIES

Let  $A$  be a  $C^*$ -algebra. An element  $a$  of the positive cone  $A^+$  of  $A$  is called a bounded-trace element if the map  $\pi \mapsto \text{tr}(\pi(a))$  is bounded on the spectrum  $\hat{A}$  of  $A$ ; the linear span of the bounded-trace elements is a two-sided  $*$ -ideal in  $A$ . We say  $A$  has *bounded trace* if the ideal of (the span of) the bounded-trace elements is dense in  $A$ .

Throughout,  $G$  is a locally compact, Hausdorff groupoid; in our main results  $G$  is assumed to be second-countable and principal. We denote the unit space of  $G$  by  $G^0$ , and the range and source maps  $r, s : G \rightarrow G^0$  are  $r(\gamma) = \gamma\gamma^{-1}$  and  $s(\gamma) = \gamma^{-1}\gamma$ , respectively. We let  $\pi : G \rightarrow G^0 \times G^0$  be the map  $\pi(\gamma) = (r(\gamma), s(\gamma))$ ; recall that  $G$  is principal if  $\pi$  is injective. In order to define the groupoid  $C^*$ -algebra, we also assume that  $G$  is equipped with a fixed left Haar system: a set  $\{\lambda^x : x \in G^0\}$  of non-negative Radon measures on  $G$  such that

- (1)  $\text{supp } \lambda^x = r^{-1}(\{x\})$ ;
- (2) for  $f \in C_c(G)$ , the function  $x \mapsto \int f d\lambda^x$  on  $G^0$  is in  $C_c(G^0)$ ; and

---

*Date:* August 22, 2006.

*2000 Mathematics Subject Classification.* 46L05, 46L55.

*Key words and phrases.* Locally compact groupoid,  $C^*$ -algebra, bounded trace.

This research was supported by the Australian Research Council and an AWM-NSF Mentoring Travel Grant.

(3) for  $f \in C_c(G)$  and  $\gamma \in G$ , the following equation holds:

$$\int f(\gamma\alpha) d\lambda^{s(\gamma)}(\alpha) = \int f(\alpha) d\lambda^{r(\gamma)}(\alpha).$$

Condition (3) implies that  $\lambda^{s(\gamma)}(\gamma^{-1}E) = \lambda^{r(\gamma)}(E)$  for measurable sets  $E$ . The collection  $\{\lambda_x : x \in G^0\}$ , where  $\lambda_x(E) := \lambda^x(E^{-1})$ , gives a fixed right Haar system such that the measures are supported on  $s^{-1}(\{x\})$  and

$$\int f(\gamma\alpha) d\lambda_{r(\alpha)} = \int f(\gamma) d\lambda_{s(\alpha)}$$

for  $f \in C_c(G)$  and  $\gamma \in G$ . We will move freely between these two Haar systems.

If  $N \subseteq G^0$  then the *saturation* of  $N$  is  $r(s^{-1}(N)) = s(r^{-1}(N))$ . In particular, we call the saturation of  $\{x\}$  the *orbit* of  $x \in G^0$  and denote it by  $[x]$ .

If  $G$  is principal and all the orbits are locally closed, then by [4, Proposition 5.1] the orbit space  $G^0/G = \{[x] : x \in G^0\}$  and the spectrum  $C^*(G)^\wedge$  of the groupoid  $C^*$ -algebra  $C^*(G)$  are homeomorphic. This homeomorphism is induced by the map  $x \mapsto L^x : G^0 \rightarrow C^*(G)^\wedge$ , where  $L^x : C^*(G) \rightarrow B(L^2(G, \lambda_x))$  is given by

$$L^x(f)\xi(\gamma) = \int f(\gamma\alpha)\xi(\alpha^{-1})d\lambda^x(\alpha)$$

for  $f \in C_c(G)$  and  $\xi \in L^2(G, \lambda_x)$ .

### 3. INTEGRABLE GROUPOIDS AND CONVERGENCE IN THE ORBIT SPACE OF A GROUPOID

The following definition is motivated by the notion of integrable action of a locally compact, Hausdorff group on a space from [6, Definition 3.2].

**Definition 3.1.** A locally compact, Hausdorff groupoid  $G$  is *integrable* if for every compact subset  $N$  of  $G^0$ ,

$$(3.1) \quad \sup_{x \in N} \{\lambda^x(s^{-1}(N))\} < \infty,$$

or, equivalently,  $\sup_{x \in N} \{\lambda_x(r^{-1}(N))\} < \infty$ .

**Remarks 3.2.** We could have taken the supremum in (3.1) over the whole unit space, that is

$$\sup_{x \in G^0} \{\lambda^x(s^{-1}(N))\} = \sup_{x \in N} \{\lambda^x(s^{-1}(N))\}.$$

To see this, first note that if  $y$  is not in the saturation  $r(s^{-1}(N)) = s(r^{-1}(N))$  of  $N$  then  $s^{-1}(N) \cap r^{-1}(\{y\}) = \emptyset$ , and hence  $\lambda^y(s^{-1}(N)) = 0$ . Second, if  $y$  is in the saturation of  $N$  then there exists a  $\gamma \in G$  such that  $s(\gamma) = y$  and  $r(\gamma) \in N$ . Then

$$r^{-1}(\{y\}) \cap s^{-1}(N) = \gamma^{-1}\gamma(r^{-1}(\{y\}) \cap s^{-1}(N)) = \gamma^{-1}(r^{-1}(\{r(\gamma)\}) \cap s^{-1}(N)),$$

and now

$$\begin{aligned} \lambda^y(s^{-1}(N)) &= \lambda^y(r^{-1}(\{y\}) \cap s^{-1}(N)) = \lambda^{r(\gamma^{-1})}(r^{-1}(\{r(\gamma)\}) \cap s^{-1}(N)) \\ &= \lambda^{s(\gamma^{-1})}(r^{-1}(\{r(\gamma)\}) \cap s^{-1}(N)) = \lambda^{r(\gamma)}(s^{-1}(N)) \end{aligned}$$

with  $r(\gamma) \in N$ .

**Example 3.3.** Let  $(H, X)$  be a locally compact, Hausdorff transformation group with  $H$  acting on the left of the space  $X$ . Then  $G = H \times X$  with

$$G^2 = \{((h, x), (k, y)) \in G \times G : y = h^{-1} \cdot x\}$$

and operations  $(h, x)(k, h^{-1} \cdot x) = (hk, x)$  and  $(h, x)^{-1} = (h^{-1}, h^{-1} \cdot x)$  is called the *transformation-group groupoid*. We identify the unit space  $\{e\} \times X$  with  $X$ , and then the range and source maps  $r, s : G \rightarrow X$  are  $s(h, x) = h^{-1} \cdot x$  and  $r(h, x) = x$ . If  $\delta_x$  is the point-mass measure on  $X$  and  $\mu$  is a left Haar measure on  $H$ , then  $\{\lambda^x := \mu \times \delta_x : x \in X\}$  is a left Haar system for  $G$ . Now

$$\lambda^x(s^{-1}(N)) = \mu(\{h \in H : h^{-1} \cdot x \in N\})$$

and hence

$$\sup_{x \in N} \{\lambda^x(s^{-1}(N))\} = \sup_{x \in N} \{\mu(\{h \in H : h^{-1} \cdot x \in N\})\},$$

that is, Definition 3.1 reduces to [6, Definition 3.2].

The following characterization of integrability will be important later. In the case of a transformation-group groupoid, Lemma 3.4 reduces to a special case of [1, Lemma 3.5].

**Lemma 3.4.** *Let  $G$  be a locally compact, Hausdorff groupoid. Then  $G$  is integrable if and only if, for each  $z \in G^0$ , there exists an open neighborhood  $U$  of  $z$  in  $G^0$  such that*

$$\sup_{x \in U} \{\lambda^x(s^{-1}(U))\} < \infty.$$

*Proof.* The proof is exactly the same as the proof of [1, Lemma 3.5]. □

If a groupoid fails to be integrable, there exists a  $z \in G^0$  such that

$$\sup_{x \in U} \{\lambda^x(s^{-1}(U))\} = \infty$$

for every open neighborhood  $U$  of  $z$ ; we then say that the *groupoid fails to be integrable at  $z$* .

It is evident from [1, 2] that integrability and  $k$ -times convergence in the orbit space of a transformation group are closely related. Moreover, Lemma 2.6 of [8] says that, if a principal groupoid fails to be proper and the orbit space  $G^0/G$  is Hausdorff, then there exists a sequence that converges 2-times in  $G^0/G$  in the sense of Definition 3.5.

**Definition 3.5.** A sequence  $\{x_n\}$  in the unit space of a groupoid  $G$  converges  $k$ -times in  $G^0/G$  to  $z \in G^0$  if there exist  $k$  sequences

$$\{\gamma_n^{(1)}\}, \{\gamma_n^{(2)}\}, \dots, \{\gamma_n^{(k)}\} \subseteq G$$

such that

- (1)  $r(\gamma_n^{(i)}) \rightarrow z$  as  $n \rightarrow \infty$  for  $1 \leq i \leq k$ ;
- (2)  $s(\gamma_n^{(i)}) = x_n$  for  $1 \leq i \leq k$ ;
- (3) if  $1 \leq i < j \leq k$  then  $\gamma_n^{(j)}(\gamma_n^{(i)})^{-1} \rightarrow \infty$  as  $n \rightarrow \infty$ , in the sense that  $\{\gamma_n^{(j)}(\gamma_n^{(i)})^{-1}\}$  admits no convergent subsequence.

**Remarks 3.6.** (a) Condition (2) in Definition 3.5 is needed so that the composition in (3) makes sense.

(b) Definition 3.5 does not require that  $x_n \rightarrow z$ , but as in the transformation-group case ([2, Definition 2.2]), this can be arranged by changing the sequence which converges  $k$ -times: replace  $x_n$  by  $r(\gamma_n^{(1)})$  and replace  $\gamma_n^{(j)}$  by  $\gamma_n^{(j)}(\gamma_n^{(1)})^{-1}$ .

(c) Part (3) of Definition 3.5 means  $\gamma_n^{(j)}(\gamma_n^{(i)})^{-1}$  is eventually outside every compact set. In particular, if  $LL^{-1}$  is compact,  $L\gamma_n^{(i)} \cap L\gamma_n^{(j)} = \emptyset$  eventually.

**Example 3.7.** Let  $G = H \times X$  be a transformation-group groupoid (see Example 3.3) and suppose that  $\{x_n\} \subseteq G^0$  is a sequence converging 2-times in  $G^0/G$  to  $z \in G^0$ . Then there exist two sequences

$$\{\gamma_n^{(1)}\} = \{(s_n, y_n)\} \quad \text{and} \quad \{\gamma_n^{(2)}\} = \{(t_n, z_n)\}$$

in  $G$  such that (1)  $y_n \rightarrow z$  and  $z_n \rightarrow z$ ; (2)  $s_n^{-1} \cdot y_n = x_n$  and  $t_n^{-1} \cdot z_n = x_n$ ; and (3)  $(t_n s_n^{-1}, z_n) \rightarrow \infty$  as  $n \rightarrow \infty$ . To see that the sequence  $\{x_n\}$  converges 2-times in  $X/H$  to  $z$  in the sense of [2, §4], consider the two sequences  $\{s_n\}$  and  $\{t_n\}$  in  $H$ . We have  $s_n \cdot x_n \rightarrow z$  and  $t_n \cdot x_n \rightarrow z$  using (1) and (2). Also, since  $z_n \rightarrow z$  by (1), (3) implies that  $t_n s_n^{-1} \rightarrow \infty$  in  $H$ .

In §4 we will prove that a principal groupoid  $G$  is integrable if and only if  $C^*(G)$  has bounded trace. For the “only if” direction we will need to know that the orbits are locally closed so that [4, Proposition 5.1] applies and  $x \mapsto L^x$  induces a homeomorphism of  $G^0/G$  onto  $C^*(G)^\wedge$ ; Corollary 3.11 below establishes that if  $G$  is integrable then the orbits are locally closed. We will prove the contrapositive of the “if” direction, and a key observation for the proof is Proposition 3.12: if a groupoid fails to be integrable at some  $z$  then there is a non-trivial sequence  $\{x_n\}$  which converges  $k$ -times in  $G^0/G$  to  $z$ , for every  $k \in \mathbb{N} \setminus \{0\}$ . The next three lemmas are needed to establish Corollary 3.11 and Proposition 3.12.

Recall that a neighborhood  $W$  of  $G^0$  is called *conditionally compact* if the sets  $WW$  and  $VW$  are relatively compact for every compact set  $V$  in  $G$ .

**Lemma 3.8.** *Let  $G$  be a second countable, locally compact, Hausdorff groupoid.*

- (1) *Let  $z \in G^0$  and let  $K$  be a relatively compact neighborhood of  $z$  in  $G$ . There exist  $a \in \mathbb{R}$  and a neighborhood  $U$  of  $z$  in  $G^0$  such that  $0 < a \leq \lambda_x(K)$  for all  $x \in U$ .*
- (2) *Let  $Q$  be a conditionally compact neighborhood in  $G$ . Given any relatively compact neighborhood  $V$  in  $G^0$  such that  $QV \neq \emptyset$ , there exists  $c \in \mathbb{R}$  such that  $c > 0$  and  $\lambda_x(Q) \leq c$  for all  $x \in V$ .*

*Proof.* (1) Suppose not. Let  $\{U_i\}$  be a decreasing sequence of open neighborhoods of  $z$  in  $G^0$ . There exists an increasing sequence  $i_1 < i_2 < \dots < i_n < \dots$  and  $x_n \in U_{i_n}$  such that  $\lambda_{x_n}(K) < 1/n$  for each  $n \geq 1$ . Note that  $x_n \rightarrow z$ .

Let  $f \in C_c(G)$  such that  $0 \leq f \leq 1$ ,  $f(z) = 1$  and  $\text{supp } f \subseteq K$ ; note that  $\int f(\gamma) d\lambda_z(\gamma) > 0$ . By the continuity of the Haar system,

$$\frac{1}{n} > \lambda_{x_n}(K) \geq \int f(\gamma) d\lambda_{x_n}(\gamma) \rightarrow \int f(\gamma) d\lambda_z(\gamma) \text{ as } n \rightarrow \infty,$$

which is impossible since the left-hand side converges to 0 and  $\int f(\gamma) d\lambda_z(\gamma) > 0$ .

(2) Let  $V$  be any relatively compact neighborhood in  $G^0$  such that  $QV \neq \emptyset$ . Let  $f \in C_c(G)$  such that  $0 \leq f \leq 1$  and  $f$  is identically one on the relatively compact subset  $QV$ . The function  $w \mapsto \int f(\gamma) d\lambda_w(\gamma)$  is in  $C_c(G^0)$ , so achieves a maximum  $c > 0$ . Then, for  $x \in V$ ,

$$\lambda_x(Q) = \lambda_x(Qx) \leq \int f(\gamma) d\lambda_x(\gamma) \leq c.$$

□

**Lemma 3.9.** *Let  $G$  be a locally compact, Hausdorff groupoid and let  $z \in G^0$ . Suppose the orbit  $[z]$  is not locally closed and let  $V$  be an open neighborhood of  $z$  in  $G^0$ . For every compact neighborhood  $K$  of  $z$  in  $G$  there exists  $\gamma_K \in G \setminus K$  such that  $s(\gamma_K) = z$  and  $r(\gamma_K) \in V$ .*

*Proof.* Suppose not, that is, suppose that there exist an open neighborhood  $V$  of  $z$  in  $G^0$  and a compact neighborhood  $K$  of  $z$  in  $G$  such that for all  $\gamma \notin K$ , either  $s(\gamma) \neq z$  or  $r(\gamma) \notin V$ . Since the orbit  $[z]$  is not locally closed,  $(\overline{[z]} \setminus [z]) \cap V \neq \emptyset$ . Let  $y \in (\overline{[z]} \setminus [z]) \cap V$ . Then there exists  $\{\gamma_i\} \subseteq G$  such that  $s(\gamma_i) = z$ ,  $r(\gamma_i) \rightarrow y$ . Moreover, since  $y \in V$  and  $V$  is open,  $r(\gamma_i) \in V$  eventually. So by assumption,  $\gamma_i \in K$  eventually. By passing to a subsequence and relabeling we may assume that  $\gamma_i \rightarrow \gamma \in K$ . But now  $s(\gamma) = z$  and  $r(\gamma) = y$ , which means that  $y \in [z]$ , a contradiction. □

Lemma 3.10 generalises [2, Lemma 2.1] from transformation groups to principal groupoids.

**Lemma 3.10.** *Let  $G$  be a second countable, locally compact, Hausdorff, principal groupoid and let  $z \in G^0$ . Then the following four conditions are equivalent.*

- (1) *the orbit  $[z]$  is not locally closed;*
- (2) *for every  $k \in \mathbb{N} \setminus \{0\}$ , the sequence  $\{z, z, \dots\}$  converges  $k$ -times in  $G^0/G$  to  $z$ ;*
- (3) *for every open neighborhood  $V$  of  $z$  in  $G^0$ ,  $\lambda_z(r^{-1}(V)) = \infty$ ;*
- (4) *for every open neighborhood  $V$  of  $z$  in  $G^0$ ,  $r^{-1}(V) \cap s^{-1}(\{z\})$  is not relatively compact.*

*Proof.* (1)  $\implies$  (2). Suppose that  $[z]$  is not open in  $\overline{[z]}$  and fix  $k \in \mathbb{N} \setminus \{0\}$ . Then, for every neighborhood  $W$  of  $z$ ,

$$(\overline{[z]} \setminus [z]) \cap W \neq \emptyset.$$

Let  $\{V_n\}$  be a decreasing sequence of basic open neighborhoods of  $z$  in  $G^0$ , and  $K_n$  an increasing sequence of compact neighborhoods of  $z$  in  $G$  such that  $G = \bigcup_{n \geq 1} \text{Int}(K_n)$ .

Now we (temporarily) fix  $n$  and choose  $\gamma_n^{(1)}, \dots, \gamma_n^{(k)}$  as follows. Set  $\gamma_n^{(1)} = z$ . By Lemma 3.9 there exists  $\gamma_n^{(2)} \in G \setminus (K_n \gamma_n^{(1)})$  such that  $s(\gamma_n^{(2)}) = z$  and  $r(\gamma_n^{(2)}) \in V_n$ . Next, note that  $K_n \gamma_n^{(1)} \cup K_n \gamma_n^{(2)}$  is compact, so by Lemma 3.9 there exists

$$\gamma_n^{(3)} \in G \setminus (K_n \gamma_n^{(1)} \cup K_n \gamma_n^{(2)})$$

with  $s(\gamma_n^{(3)}) = z$  and  $r(\gamma_n^{(3)}) \in V_n$ . Continue.

By construction,  $r(\gamma_n^{(i)}) \rightarrow z$  for  $1 \leq i \leq k$  as  $n \rightarrow \infty$ . Also, if  $1 \leq i < j \leq k$  then  $\gamma_n^{(j)} \notin \cup_{i=1}^{j-1} K_n \gamma_n^{(i)}$ ; in particular

$$\gamma_n^{(j)}(\gamma_n^{(i)})^{-1} \notin K_n.$$

Now let  $Q$  be any compact set in  $G$ . Choose  $n_0$  such that  $Q \subseteq K_n$  whenever  $n > n_0$ . Since  $\gamma_n^{(j)}(\gamma_n^{(j)})^{-1} \notin K_n$  we have  $\gamma_n^{(j)}(\gamma_n^{(j)})^{-1} \notin Q$  whenever  $n > n_0$ . Hence  $\gamma_n^{(j)}(\gamma_n^{(j)})^{-1} \rightarrow \infty$  as  $n \rightarrow \infty$ . Thus  $\{z, z, \dots\}$  converges  $k$ -times in  $G^{(0)}/G$  to  $z$ . Finally, note that  $k$  was arbitrary.

(2)  $\implies$  (3). Suppose that for every  $k \in \mathbb{N} \setminus \{0\}$ , the sequence  $\{z, z, \dots\}$  converges  $k$ -times in  $G^0/G$  to  $z$ . Let  $V$  be an open neighborhood of  $z$  in  $G^{(0)}$  and let  $M \in \mathbb{N} \setminus \{0\}$ . There exists an open neighborhood  $U$  of  $z$  in  $G^0$  and a compact symmetric neighborhood  $K$  of  $z$  in  $G$  such that  $r(KU) \subseteq V$ . By Lemma 3.8(1) and by shrinking  $U$  if necessary, there exists  $c > 0$  such that  $\lambda_u(K) \geq c$  for  $u \in U$ . Choose  $k \in \mathbb{N} \setminus \{0\}$  such that  $kc > M$ . Next, choose  $k$  sequences  $\{\gamma_n^{(i)}\} \subseteq G$  such that, for

- (a)  $s(\gamma_n^{(i)}) = z$  and  $r(\gamma_n^{(i)}) \rightarrow z$  as  $n \rightarrow \infty$  for  $1 \leq i \leq k$ , and
- (b) if  $1 \leq i < j \leq k$  then  $\gamma_n^{(j)}(\gamma_n^{(i)})^{-1} \rightarrow \infty$  as  $n \rightarrow \infty$ .

So there exists  $n_0$  such that  $r(\gamma_{n_0}^{(i)}) \in U$  for  $1 \leq i \leq k$  and  $\gamma_{n_0}^{(j)}(\gamma_{n_0}^{(i)})^{-1} \in G \setminus K^2$ . Now  $r(K\gamma_{n_0}^{(i)}) \subseteq r(KU) \subseteq V$  and  $K\gamma_{n_0}^{(i)} \cap K\gamma_{n_0}^{(j)} = \emptyset$  when  $i \neq j$ . By our choice of  $n_0$  we have

$$\lambda_z(r^{-1}(V)) \geq \lambda_z\left(\bigcup_{i=1}^k K\gamma_{n_0}^{(i)}\right) = \sum_{i=1}^k \lambda_z(K\gamma_{n_0}^{(i)}) = \sum_{i=1}^k \lambda_{r(\gamma_{n_0}^{(i)})}(K) \geq kc > M.$$

Since  $M$  was arbitrary,  $\lambda_z(r^{-1}(V)) = \infty$ .

(3)  $\implies$  (4) If  $\lambda_z(r^{-1}(V)) = \infty$  then

$$\lambda_z(r^{-1}(V) \cap s^{-1}(\{z\})) = \infty$$

because the support of  $\lambda_z$  is  $s^{-1}(\{z\})$ . Since  $\lambda_z$  is a Radon measure  $r^{-1}(V) \cap s^{-1}(\{z\})$  cannot be relatively compact.

(4)  $\implies$  (1) We prove the contrapositive. Assume  $[z]$  is locally closed in  $G$ . Then there exists an open set  $U$  in  $G$  such that  $U \cap \overline{[z]} = [z]$ . Let  $N$  be a compact neighborhood of  $z$  in  $G$  such that  $N \subset U$  (such  $N$  exists because  $G$  is locally compact Hausdorff). Since  $G^0$  is closed, we may assume  $N \subset G^0$  (otherwise replace  $N$  by  $N \cap G^0$ ). Since  $N \cap \overline{[z]}$  is compact in  $G$  and  $N \cap \overline{[z]} = N \cap [z]$ ,  $N \cap [z]$  is a compact neighborhood of  $z$  in  $[z]$ .

Consider  $G|_{[z]} = \{\gamma \in G : s(\gamma) \in [z] \text{ and } r(\gamma) \in [z]\}$ , the restriction of  $G$  to  $[z]$ . Since  $[z]$  is locally closed, it is locally compact, and hence  $G|_{[z]}$  is locally compact. Thus  $G|_{[z]}$  satisfies the hypothesis of [11, Theorem 2.1]: it is a second countable, locally compact, Hausdorff principal groupoid with (one) locally closed orbit  $(G|_{[z]})^0 = [z]$ . Thus  $r| : s^{-1}(\{z\}) \rightarrow [z]$  is a homeomorphism. (Notice that  $s^{-1}(\{z\}) \subseteq G|_{[z]}$ .) Now

$$A := r|^{-1}(N \cap [z]) = \{\gamma \in G : s(\gamma) = z \text{ and } r(\gamma) \in N \cap [z]\}$$

is a compact neighborhood of  $z$  in  $s^{-1}(\{z\})$ . So  $A$  is a compact subset of  $G$  as well. Let  $V$  be the interior of  $N$  in  $G^0$ . Then  $V$  is an open neighborhood of  $z$  in  $G^0$  such that

$$s^{-1}(\{z\}) \cap r^{-1}(V \cap [z]) = \pi^{-1}((V \cap [z]) \times \{z\})$$

is relatively compact in  $G$ . □

**Corollary 3.11.** *Let  $G$  be a second countable, locally compact, Hausdorff, principal groupoid. If  $G$  is integrable then all orbits are locally closed.*

*Proof.* We show the contrapositive. Suppose there exists  $z \in G^0$  such that  $[z]$  is not locally closed. Let  $V$  be any open relatively compact neighborhood of  $z$  in  $G$ . By Lemma 3.10  $\lambda_z(r^{-1}(V)) = \infty$  and hence

$$\sup\{\lambda^x(s^{-1}(\overline{V})) : x \in \overline{V}\} \geq \sup\{\lambda_x(r^{-1}(V)) : x \in V\} = \infty.$$

So  $G$  is not integrable. □

**Proposition 3.12.** *Let  $G$  be a locally compact, Hausdorff groupoid. Let  $z \in G^0$  and suppose that  $G$  fails to be integrable at  $z$ . Then there exists a sequence  $\{x_n\}$  in  $G^0$  such that  $x_n \rightarrow z$ , and  $\{x_n\}$  converges  $k$ -times in  $G^0/G$  to  $z$ , for every  $k \in \mathbb{N} \setminus \{0\}$ . In addition, if  $G$  is second countable, principal and the orbit  $[z]$  is locally closed, then  $x_n \neq z$  eventually.*

*Proof.* Suppose the groupoid fails to be integrable at  $z$ . Fix  $k \in \mathbb{N} \setminus \{0\}$ . Let  $\{U_n\}$  be a decreasing sequence of open relatively compact neighborhoods of  $z$  in  $G^0$ . By Lemma 3.4

$$\sup_{y \in U_n} \{\lambda^y(s^{-1}(U_n))\} = \infty$$

for each  $n$ . So we can choose a sequence  $\{x_n\}$  such that  $x_n \in U_n$  and  $\lambda^{x_n}(s^{-1}(U_n)) > n$ . Note that  $x_n \rightarrow z$  as  $n \rightarrow \infty$ .

Let  $Q$  be an open symmetric conditionally compact neighborhood of  $z$  in  $G$  and  $V$  an open relatively compact neighborhood of  $z$  in  $G^0$ . By Lemma 3.8(2) there exist  $c > 0$  such that  $\lambda_v(Q^2) \leq c$  whenever  $v \in V$ . Choose  $n_0$  such that  $n_0 > (k-1)c$  and  $U_{n_0} \subseteq V$ . Temporarily fix  $n > n_0$ . Set  $\gamma_n^{(1)} = x_n$ . For  $k \geq 2$  choose  $k-1$  elements  $\gamma_n^{(2)}, \dots, \gamma_n^{(k)}$  as follows. Note that since  $x_n = r(\gamma_n^{(1)}) \in V$  we have

$$\begin{aligned} \lambda_{x_n}(r^{-1}(U_n) \setminus Q^2\gamma_n^{(1)}) &\geq \lambda_{x_n}(r^{-1}(U_n)) - \lambda_{x_n}(Q^2\gamma_n^{(1)}) \\ &= \lambda_{x_n}(r^{-1}(U_n) \cap s^{-1}(\{x_n\})) - \lambda_{r(\gamma_n^{(1)})}(Q^2) \\ &> (k-1)c - c = (k-2)c \geq 0. \end{aligned}$$

So there exists

$$\gamma_n^{(2)} \in (r^{-1}(U_n) \cap s^{-1}(\{x_n\})) \setminus Q^2\gamma_n^{(1)};$$

note that  $r(\gamma_n^{(2)}) \in U_n \subset V$  and  $s(\gamma_n^{(2)}) = x_n$ . Next,

$$\begin{aligned} \lambda_{x_n}((r^{-1}(U_n) \setminus (Q^2\gamma_n^{(1)} \cup Q^2\gamma_n^{(2)}))) &\geq \lambda_{x_n}(r^{-1}(U_n)) - \lambda_{x_n}(Q^2\gamma_n^{(1)}) - \lambda_{x_n}(Q^2\gamma_n^{(2)}) \\ &\geq \lambda_{x_n}(r^{-1}(U_n) \cap s^{-1}(\{x_n\})) - \lambda_{r(\gamma_n^{(1)})}(Q^2) - \lambda_{r(\gamma_n^{(2)})}(Q^2) \\ &> (k-3)c \geq 0. \end{aligned}$$

Continue until  $\gamma_n^{(1)}, \dots, \gamma_n^{(k)}$  have been chosen in this way.

If  $n > n_0$ , then by construction  $s(\gamma_n^{(i)}) = x_n$  and  $r(\gamma_n^{(i)}) \in U_n$  for each  $n$ ; so  $r(\gamma_n^{(i)}) \rightarrow z$  as  $n \rightarrow \infty$  for  $1 \leq i \leq k$ . Moreover  $\gamma_n^{(j)}(\gamma_n^{(i)})^{-1} \notin Q^2$  for  $1 \leq i < j \leq k$  and  $n > n_0$ . To see that  $\{\gamma_n^{(j)}(\gamma_n^{(i)})^{-1}\}$  tends to infinity, suppose that it doesn't. Then,  $\gamma_n^{(j)}(\gamma_n^{(i)})^{-1} \rightarrow \gamma$  by passing to a subsequence and relabelling. But then  $s(\gamma_n^{(j)}(\gamma_n^{(i)})^{-1}) = r(\gamma_n^{(i)}) \rightarrow z$  and  $r(\gamma_n^{(j)}(\gamma_n^{(i)})^{-1}) = r(\gamma_n^{(j)}) \rightarrow z$  implies  $\gamma = z$ , which is impossible because  $\gamma_n^{(j)}(\gamma_n^{(i)})^{-1} \notin Q^2$  and  $Q$  contains  $G^0$ . Hence  $\{x_n\}$  converges  $k$ -times in  $G^0/G$  to  $z$ .

We claim that if  $G$  is second countable and principal, then  $x_n \neq z$  eventually. To see this, suppose  $x_n = z$  frequently. Then  $\lambda^z(s^{-1}(U_n)) > n$  frequently, and hence

$$(3.2) \quad \lambda^z(s^{-1}(U_1)) = \infty.$$

The orbits are locally closed and  $G$  is second countable and principal, so the source map restricts to a homeomorphism  $s| : r^{-1}(\{z\}) \rightarrow [z]$ . Since  $U_1$  is relatively compact,  $s^{-1}([z] \cap U_1)$  is relatively compact in  $r^{-1}(\{z\})$  because  $s| : r^{-1}(\{z\}) \rightarrow [z]$  is a homeomorphism. But now  $\lambda^z(s^{-1}([z] \cap U_1)) = \lambda^z(s^{-1}(U_1)) < \infty$ , contradicting (3.2).  $\square$

#### 4. INTEGRABILITY OF $G$ AND TRACE PROPERTIES OF $C^*(G)$

**Proposition 4.1.** *Let  $G$  be a second-countable, locally compact, Hausdorff, principal groupoid. If  $C^*(G)$  has bounded trace then  $G$  is integrable.*

The proof of Proposition 4.1 is based on that of [8, Theorem 2.3]. There, Muhly and Williams choose a sequence  $\{x_n\} \subseteq G^0$  with  $x_n \rightarrow z$  which witnesses the failure of the groupoid to be proper. They then carefully construct a function  $f \in C_c(G)$  to obtain an element  $d$  of the Pedersen ideal of  $C^*(G)$  such that  $\text{tr}(L^{x_n}(d))$  does not converge to  $\text{tr}(L^z(d))$ . Since the Pedersen ideal is the minimal dense ideal [9, Theorem 5.6.1], the ideal of continuous-trace elements cannot be dense, so  $C^*(G)$  does not have

continuous trace. We adopt the same strategy, use exactly the same function  $f$ , but adapt the proof of [8, Theorem 2.3] using ideas from [6, Proposition 3.5].

*Proof of Proposition 4.1.* Fix  $M \in \mathbb{N} \setminus \{0\}$ . We will show that there is an element  $d$  of the Pedersen ideal of  $C^*(G)$ , a sequence of representations  $\{L^{x_n}\}$  and  $n_0 > 0$  such that  $\text{tr}(L^{x_n}(d)) > M$  whenever  $n > n_0$ . Since  $M$  is arbitrary,  $C^*(G)$  cannot have bounded trace.

If  $G$  is not integrable, then the integrability fails at some  $z \in G^0$  by Lemma 3.4. If the orbits are not closed then  $C^*(G)$  cannot be CCR by [4, Theorem 4.1], hence cannot have bounded trace. So from now on we may assume that the orbits are closed. By Proposition 3.12, there exists a sequence  $\{x_n\}$  such that  $x_n \neq z$ ,  $x_n \rightarrow z$ , and  $\{x_n\}$  converges  $k$ -times in  $G^0/G$  to  $z$ , for every  $k \in \mathbb{N} \setminus \{0\}$ .

Since we will use exactly the same function  $f$  that was used in the proof of [8, Theorem 2.3], our first task is to briefly outline its construction. Fix a function  $g \in C_c(G^0)$  such that  $0 \leq g \leq 1$  and  $g$  is identically one on a neighborhood  $U$  of  $z$ . Let  $N = \text{supp } g$  and

$$\begin{aligned} F_z^N &:= s^{-1}(\{z\}) \cap r^{-1}([z] \cap N) = s^{-1}(\{z\}) \cap r^{-1}(N) \\ F_N^z &:= r^{-1}(\{z\}) \cap s^{-1}([z] \cap N) = r^{-1}(\{z\}) \cap s^{-1}(N). \end{aligned}$$

There exist symmetric, open, conditionally compact neighborhoods  $W_0$  and  $W_1$  in  $G$  such that

$$G^0 \subseteq W_0 \subseteq \overline{W_0} \subseteq W_1 \quad \text{and} \quad F_N^z \cup F_z^N \subseteq W_0.$$

Thus  $\overline{W_1^7}z \setminus W_0z \subseteq r^{-1}(G^0 \setminus N)$ . (The reason for using  $\overline{W_1^7}$  becomes clear at (4.4) below). By a compactness argument, there exist open, symmetric, relatively compact neighborhoods  $V_0 \subseteq G^0$  and  $V_1$  of  $z$  in  $G$  such that  $\overline{V_0} \subset V_1$  and

$$(4.1) \quad \overline{W_1^7} \overline{V_0} \setminus W_0 V_0 \subseteq r^{-1}(G^0 \setminus N).$$

Now note that if  $\gamma \in \overline{W_1^7} \overline{V_1} \overline{W_1^7} \setminus W_0 V_0 W_0$  then  $r(\gamma) \in r(\overline{W_1^7} \overline{V_0} \setminus W_0 V_0) \subseteq G^0 \setminus N$ . It follows that the function  $g^{(1)} : G \rightarrow [0, 1]$  defined by

$$g^{(1)}(\gamma) = \begin{cases} g(r(\gamma)) & \text{if } \gamma \in \overline{W_1^7} \overline{V_1} \overline{W_1^7}, \\ 0 & \text{if } \gamma \notin W_0 V_0 W_0 \end{cases}$$

is well-defined and continuous with compact support in  $G$ . By construction

$$(W_0 V_0 W_0)^2 = W_0 V_0 W_0^2 V_0 W_0 \subseteq W_0^4 V_0 W_0^4 \subseteq \overline{W_0^4} \overline{V_0} \overline{W_0^4} \subseteq W_1^4 V_1 W_1^4 \subseteq \overline{W_1^4} \overline{V_1} \overline{W_1^4}.$$

So there exists a function  $b \in C_c(G)$  such that  $0 \leq b \leq 1$ ,  $b$  is identically one on  $W_0 V_0 W_0^2 V_0 W_0$  and is identically zero on the complement of  $\overline{W_1^4} \overline{V_1} \overline{W_1^4}$ . Further, we can replace  $b$  with  $(b + b^*)/2$  to ensure that  $b$  is self-adjoint. Set

$$f(\gamma) = g(r(\gamma))g(s(\gamma))b(\gamma);$$

note that  $f \in C_c(G)$  is self-adjoint.

For  $\xi \in L^2(G, \lambda_u)$  and  $\gamma \in G$  we have

$$\begin{aligned} L^u(f)\xi(\gamma) &= \int f(\gamma\alpha)\xi(\alpha^{-1}) d\lambda^u(\alpha) \\ &= \int g(r(\gamma))g(s(\alpha))b(\gamma\alpha)\xi(\alpha^{-1}) d\lambda^u(\alpha) \\ &= g(r(\gamma)) \int g(s(\alpha))b(\gamma\alpha)\xi(\alpha^{-1}) d\lambda^u(\alpha) \\ (4.2) \quad &= g(r(\gamma)) \int g(r(\alpha))b(\gamma\alpha^{-1})\xi(\alpha) d\lambda_u(\alpha). \end{aligned}$$

By [8, Lemma 2.8],  $g^{(1)}$  is an eigenvector for  $L^{x_n}(f)$  with eigenvalue

$$\mu_{x_n}^{(1)} = \int g(r(\alpha))g^{(1)}(\alpha) d\lambda_{x_n}(\alpha) = \int_{W_0 V_0 W_0} g(r(\alpha))^2 d\lambda_{x_n}(\alpha).$$

By [8, Lemma 2.9], there exist an open  $V_2 \subseteq V_0$  and a conditionally compact neighborhood  $Y$  of  $G^0$  so that  $Y \subseteq W_0$  and if  $v \in V_2$ , then  $r(Yv) \subseteq U$ . Notice that  $YV_2Y$  is a relatively compact subset of  $W_0V_0W_0$ . By Lemma 3.8(1) there exist an open neighborhood  $V_3$  of  $z$  and  $a > 0$  such that

$$(4.3) \quad \lambda_v(YV_2Y) \geq a \text{ whenever } v \in V_3.$$

Now, if  $\alpha \in YV_2Y$  then  $r(\alpha) \in U$  and hence  $g(r(\alpha)) = 1$ ; it follows that

$$\mu_{x_n}^{(1)} \geq \int_{YV_2Y} g(r(\alpha))^2 d\lambda_{x_n}(\alpha) = \lambda_{x_n}(YV_2Y) \geq a > 0$$

whenever  $x_n \in V_3$ .

So far our set-up is the one from [8]. Now choose  $l \in \mathbb{N} \setminus \{0\}$  such that  $la^2 > M$ . (Note that  $a$  is independent of  $l$ !) The sequence  $\{x_n\}$  converges  $k$ -times in  $G/G^0$  to  $z$  for every  $k \in \mathbb{N} \setminus \{0\}$ , so it certainly converges  $l$  times. So there exist  $l$  sequences

$$\{\gamma_n^{(1)}\}, \{\gamma_n^{(2)}\}, \dots, \{\gamma_n^{(l)}\} \subseteq G$$

such that

- (1)  $r(\gamma_n^{(i)}) \rightarrow z$  as  $n \rightarrow \infty$  for  $1 \leq i \leq l$ ;
- (2)  $s(\gamma_n^{(i)}) = x_n$  for  $1 \leq i \leq k$ ;
- (3) if  $1 \leq i < j \leq l$  then  $\gamma_n^{(j)}(\gamma_n^{(i)})^{-1} \rightarrow \infty$ .

Moreover, by construction (see Proposition 3.12), we may take  $\gamma_n^{(1)} = x_n$ . Temporarily fix  $n$ . Set  $g_n^{(1)} := g^{(1)}$ , and for  $2 \leq j \leq l$  set

$$\begin{aligned} g_n^{(j)}(\gamma) &:= \begin{cases} g^{(1)}(\gamma(\gamma_n^{(j)})^{-1}), & \text{if } s(\gamma) = s(\gamma_n^{(j)}); \\ 0, & \text{otherwise} \end{cases} \\ &= \begin{cases} g(r(\gamma)), & \text{if } \gamma \in \overline{W_1^7 V_1 W_1^7} \gamma_n^{(j)}; \\ 0, & \text{otherwise} \end{cases} \\ &= \begin{cases} g(r(\gamma)), & \text{if } \gamma \in \overline{W_1^7 V_1 W_1^7} \gamma_n^{(j)}; \\ 0, & \text{if } \gamma \notin W_0 V_0 W_0 \gamma_n^{(j)}. \end{cases} \end{aligned}$$

Each  $g_n^{(i)}$  ( $1 \leq j \leq l$ ) is a well-defined function in  $C_c(G)$  with support contained in  $W_0 V_0 W_0 \gamma_n^{(j)}$ . For  $1 \leq i < j \leq l$ ,  $\gamma_n^{(j)}(\gamma_n^{(i)})^{-1} \notin (W_0 V_0 W_0)^2$  eventually, so there exists  $n_0 > 0$  such that, for every  $0 \leq i, j \leq l$ ,  $i \neq j$ ,

$$W_0 V_0 W_0 \gamma_n^{(j)} \cap W_0 V_0 W_0 \gamma_n^{(i)} = \emptyset$$

whenever  $n > n_0$ .

We now prove a generalization of [8, Lemma 2.8] which, together with (4.2), immediately implies that each  $g_n^{(j)}$  is an eigenvector of  $L^{x_n}(f)$  for  $1 \leq j \leq l$ .

**Lemma 4.2.** *With the choices made above, for all  $\alpha, \gamma \in G$  and  $1 \leq j \leq l$ ,*

$$g(r(\gamma))g(r(\alpha))b(\gamma\alpha^{-1})g_n^{(j)}(\alpha) = g_n^{(j)}(\gamma)g(r(\alpha))g_n^{(j)}(\alpha).$$

*Proof.* If  $\alpha \notin W_0 V_0 W_0 \gamma_n^{(j)}$ , then both sides are zero. So we may assume throughout that  $\alpha \in W_0 V_0 W_0 \gamma_n^{(j)}$ .

If  $\gamma \in W_0 V_0 W_0 \gamma_n^{(j)}$ , then  $g_n^{(j)}(\gamma) = g(r(\gamma))$  and  $\gamma\alpha^{-1} \in W_0 V_0 W_0^2 V_0 W_0$ , so  $b(\gamma\alpha^{-1}) = 1$  and both sides agree.

If  $\gamma \in \overline{W_1^7 V_1 W_1^7} \gamma_n^{(j)} \setminus W_0 V_0 W_0 \gamma_n^{(j)}$ , then  $g(r(\gamma)) = 0 = g_n^{(j)}(\gamma)$ , so both sides are zero.

Finally, if  $\gamma \notin \overline{W_1^7 V_1 W_1^7} \gamma_n^{(j)}$ , then  $g_n^{(j)}(\gamma) = 0$  so the right-hand side zero. On the other hand, if  $\gamma\alpha^{-1} \in \overline{W_1^4 V_1 W_1^4} (= \text{supp } b)$  then

$$(4.4) \quad \gamma \in \overline{W_1^4 V_1 W_1^4} \gamma_n^{(j)} \subseteq \overline{W_1^7 V_1 W_1^7} \gamma_n^{(j)}.$$

So  $\gamma \notin \overline{W_1^7 V_1 W_1^7} \gamma_n^{(j)}$  implies  $\gamma\alpha^{-1} \notin \text{supp } b$ , so the left-hand side is zero as well.  $\square$

Let  $\mu_n^{(j)}$  be the eigenvalue corresponding to the eigenvector  $g_n^{(j)}$ . Using (4.3)

$$\mu_n^{(j)} = \int_{W_0 V_0 W_0 \gamma_n^{(j)}} g(r(\alpha))^2 d\lambda_{x_n}(\alpha) \geq \lambda_{x_n}(YV_2Y\gamma_n^{(j)}) = \lambda_{r(\gamma_n^{(j)})}(YV_2Y) \geq a$$

whenever  $r(\gamma_n^{(j)}) \in V_3$ . Choose  $n_1 > n_0$  such that  $n > n_1$  implies  $x_n \in V_3$  and  $r(\gamma_n^{(j)}) \in V_3$  for  $1 \leq j \leq l$ . Then  $L^{x_n}(f * f)$  is a positive compact operator with  $l$  eigenvalues  $\mu_n^{(j)} \geq a^2$  for  $1 \leq j \leq l$ . To push  $f * f$  into the Pedersen ideal, let  $r \in C_c(0, \infty)$  be any function satisfying

$$r(t) = \begin{cases} 0, & \text{if } t < \frac{a^2}{3}; \\ 2t - \frac{2a^2}{3}, & \text{if } \frac{a^2}{3} \leq t < \frac{2a^2}{3}; \\ t, & \text{if } \frac{2a^2}{3} \leq t \leq \|f * f\|. \end{cases}$$

Set  $d := r(f * f)$ . Now  $d$  is a positive element of the Pedersen ideal of  $C^*(G)$  with  $\text{tr}(L^{x_n}(d)) \geq la^2 > M$  whenever  $n > n_1$ . Since  $M$  was arbitrary,  $L^x \mapsto \text{tr}(L^x(d))$  is unbounded on  $C^*(G)^\wedge$ . Thus  $C^*(G)$  does not have bounded trace.  $\square$

**Proposition 4.3.** *Suppose  $G$  is a second countable, locally compact, Hausdorff, principal groupoid. If  $G$  is integrable then  $C^*(G)$  has bounded trace.*

*Proof.* Since  $G$  is principal and integrable, the orbits are locally closed by Corollary 3.11, and  $x \mapsto L^x$  induces a homeomorphism of  $G^0/G$  onto  $C^*(G)^\wedge$  by [4, Proposition 5.1]. To show that  $C^*(G)$  has bounded trace, it suffices to see that for a fixed  $u \in G^0$  and all  $f \in C_c(G)$ ,  $\text{tr}(L^u(f * f))$  is bounded independent of  $u$ .

Fix  $u \in G^0$  and let  $\xi \in L^2(G, \lambda_u)$ . Since

$$L^u(f)\xi(\gamma) = \int f(\gamma\alpha^{-1})\xi(\alpha) d\lambda_u(\alpha),$$

$L^u(f)$  is a kernel operator on  $L^2(G, \lambda_u)$  with kernel  $k_f$  given by  $k_f(\gamma, \alpha) = f(\gamma\alpha^{-1})$ . We will show that  $k_f \in L^2(G \times G, \lambda_u \times \lambda_u)$  and find a bound on  $k_f$  independent of  $u$ . This will complete the proof since  $\text{tr}(L^u(f * f)) = \|k_f\|^2$  by, for example, [10, Theorem 3.4.16].

Notice that

$$\begin{aligned} \|k_f\|^2 &= \int_{G \times G} |k_f(\gamma, \alpha)|^2 d(\lambda_u \times \lambda_u)(\gamma, \alpha) = \int_G \int_G |f(\gamma\alpha^{-1})|^2 d\lambda_u(\gamma) d\lambda_u(\alpha) \\ (4.5) \qquad &= \int_G \int_G |f(\gamma)|^2 d\lambda_{r(\alpha)}(\gamma) d\lambda_u(\alpha) \end{aligned}$$

by Tonelli's Theorem and right invariance. For a fixed  $\alpha$ , the inner integral

$$\int_G |f(\gamma)|^2 d\lambda_{r(\alpha)}(\gamma) \leq \|f\|_\infty^2 \lambda_{r(\alpha)}(\text{supp } f)$$

and is zero unless  $r(\alpha) \in s(\text{supp } f)$ . The outer integral is zero unless  $s(\alpha) = u$ . Let

$$K = r^{-1}(s(\text{supp } f)) \cap s^{-1}(\{u\}).$$

So

$$\begin{aligned} (4.5) &\leq \int_K \|f\|_\infty^2 \lambda_{r(\alpha)}(\text{supp } f) d\lambda_u(\alpha) \\ &\leq \|f\|_\infty^2 \sup \{ \lambda_{r(\alpha)}(\text{supp } f) : r(\alpha) \in s(\text{supp } f) \} \lambda_u(K) \\ &\leq \|f\|_\infty^2 \sup \{ \lambda_x(\text{supp } f) : x \in s(\text{supp } f) \} \sup \{ \lambda_x(r^{-1}(s(\text{supp } f))) : x \in G^0 \}. \end{aligned}$$

Since  $s(\text{supp } f)$  is a compact subset of  $G^0$ , by integrability there exists  $N > 0$  such that

$$\sup \{ \lambda_x(r^{-1}(s(\text{supp } f))) : x \in G^0 \} < N$$

(see also Remark 3.2). Note that  $N$  does not depend on  $u$ . By Lemma 3.8(2), applied to the conditionally compact neighborhood  $\text{supp } f$  and the relatively compact neighborhood  $s(\text{supp } f)$ , there exists  $M > 0$  such that  $\lambda_x(\text{supp } f) < M$  for all  $x \in s(\text{supp } f)$ , that is,

$$\sup \{ \lambda_x(\text{supp } f) : x \in s(\text{supp } f) \} < M.$$

Note that  $M$  does not depend on  $u$ .

Thus  $\|k_f\|^2 < \|f\|_\infty^2 MN$ , so  $k_f \in L^2(G \times G, \lambda_u \times \lambda_u)$  as claimed, and

$$\mathrm{tr}(L^u(f^* * f)) = \|k_f\|^2 < \|f\|_\infty^2 MN,$$

which is a bound on  $\mathrm{tr}(L^u(f^* * f))$  independent of  $u$ .  $\square$

Combining Propositions 4.1 and 4.3 we have:

**Theorem 4.4.** *Suppose  $G$  is a second countable, locally compact, Hausdorff, principal groupoid. Then  $G$  is integrable if and only if  $C^*(G)$  has bounded trace.*

By Corollary 3.11, integrable groupoids have locally closed orbits; we can now improve this result using Proposition 4.3 (which relied on Corollary 3.11). The  $C^*$ -algebras of integrable groupoids have bounded trace by Proposition 4.3, hence are CCR. But if  $C^*(G)$  is CCR then the orbit space  $G^0/G$  is  $T_1$  by [4, Theorem 4.1], that is, the orbits are closed.

**Corollary 4.5.** *Suppose  $G$  is a second countable, locally compact, Hausdorff, principal groupoid. If  $G$  is integrable then the orbits of  $G$  are closed.*

We have so far been unable to prove directly that integrability implies that the orbits are closed.

#### REFERENCES

- [1] R.J. Archbold and K. Deicke, *Bounded trace  $C^*$ -algebras and integrable actions*, Math. Zeit. **250** (2005), 393–410.
- [2] R.J. Archbold and A. an Huef, *Strength of convergence in the orbit space of a transformation group*, to appear in J. Funct. Anal.
- [3] R.J. Archbold and D.W.B. Somerset, *Transition probabilities and trace functions for  $C^*$ -algebras*, Math. Scand. **73** (1993), 81–111.
- [4] L.O. Clark, *Classifying the type of principal groupoid  $C^*$ -algebras*, to appear in J. Operator Theory.
- [5] P. Green,  *$C^*$ -algebras of transformation groups with smooth orbit space*, Pacific J. Math. **72** (1977), 71–97.
- [6] A. an Huef, *Integrable actions and the transformation groups whose  $C^*$ -algebras have bounded trace*, Indiana Univ. Math. J. **51** (2002), 1197–1233.
- [7] P. S. Muhly, J. Renault, and D. P. Williams, *Equivalence and isomorphism for groupoid  $C^*$ -algebras*, J. Operator Theory **17** (1987), 3–22.
- [8] P.S. Muhly and D.P. Williams, *Continuous trace groupoid  $C^*$ -algebras*, Math. Scand. **66** (1990), 231–241.
- [9] G.K. Pedersen,  *$C^*$ -algebras and their automorphism groups*, Academic Press, London, 1979.
- [10] G. K. Pedersen, *Analysis Now*, Springer-Verlag, New York, 1989.
- [11] A. Ramsay, *The Mackey-Glimm dichotomy for foliations and other Polish groupoids*, J. Funct. Anal. **94** (1990), 358–374.
- [12] J. Renault, *A groupoid approach to  $C^*$ -algebras*, Lecture Notes in Mathematics, No. **793**, Springer-Verlag, New York, 1980.
- [13] M.A. Rieffel, *Integrable and proper actions on  $C^*$ -algebras, and square-integrable representations of groups*, Expo. Math. **22** (2004), 1–53.

DEPARTMENT OF MATHEMATICAL SCIENCES, SUSQUEHANNA UNIVERSITY, SELINGROVE, PA 17870, USA  
*E-mail address:* clarklisa@susqu.edu

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW SOUTH WALES, SYDNEY, NSW 2052, AUSTRALIA  
*E-mail address:* astrid@unsw.edu.au