Global defensive k-alliances in graphs

J. A. Rodríguez-Velázquez¹ and J. M. Sigarreta²

¹Department of Computer Engineering and Mathematics Rovira i Virgili University of Tarragona

Av. Països Catalans 26, 43007 Tarragona, Spain

e-mail: juanalberto.rodriguez@urv.cat

²Departamento de Matemáticas, Universidad Carlos III de Madrid, Avda. de la Universidad 30, 28911 Leganés (Madrid) Spain.

e-mail:josemaria.sigarreta@uc3m.es

Abstract

Let $\Gamma = (V, E)$ be a simple graph. For a nonempty set $X \subseteq V$, and a vertex $v \in V$, $\delta_X(v)$ denotes the number of neighbors v has in X. A nonempty set $S \subseteq V$ is a *defensive k-alliance* in $\Gamma = (V, E)$ if $\delta_S(v) \geq \delta_{\bar{S}}(v) + k, \forall v \in S$. A defensive k-alliance S is called global if it forms a dominating set. The global defensive k-alliance number of Γ , denoted by $\gamma_k^a(\Gamma)$, is the minimum cardinality of a defensive k-alliance in Γ . We study the mathematical properties of $\gamma_k^a(\Gamma)$.

Keywords: Defensive alliances, alliances in graphs, domination. *AMS Subject Classification numbers:* 05C69; 05A20

1 Introduction

Since (defensive, offensive and dual) alliances were first introduced by P. Kristiansen, S. M. Hedetniemi and S. T. Hedetniemi [12], several authors have studied their mathematical properties [2, 4, 3, 6, 9, 13, 14, 16, 18, 20, 22] as well as the complexity of computing minimum-cardinality of alliances [1, 7, 10, 11]. The minimum-cardinality of a defensive (respectively, offensive

or dual) alliance in a graph Γ is called the defensive (respectively, offensive or dual) alliance number of Γ . The mathematical properties of defensive alliances were first studied in [12] where several bounds on the defensive alliance number were given. The particular case of global (strong) defensive alliances was investigated in [9] where several bounds on the global (strong) defensive alliance number were obtained. The dual alliances were introduced as powerful alliances in [2, 3]. In [14] there were obtained several tight bounds on the defensive (offensive and dual) alliance number. In particular, there was investigated the relationship between the alliance numbers of a graph and its algebraic connectivity, its spectral radius, and its Laplacian spectral radius. Moreover, the study of global defensive (offensive and dual) alliances in planar graph was initiated in [16] and the study of defensive alliances in the line graph of a simple graph was initiated in cite [22]. The particular case of global alliances in trees has been investigated in [4]. For many properties of offensive alliances, readers may refer to [6, 13, 23].

A generalization of (defensive and offensive) alliances called k-alliances was presented by K. H. Shafique and R. D. Dutton [18, 19] where was initiated the study of k-alliance free sets and k-alliance cover sets. The aim of this work is to study mathematical properties of defensive k-alliances. We begin by stating the terminology used. Throughout this article, $\Gamma = (V, E)$ denotes a simple graph of order |V| = n and size |E| = m. We denote two adjacent vertices u and v by $u \sim v$. For a nonempty set $X \subseteq V$, and a vertex $v \in V$, $N_X(v)$ denotes the set of neighbors v has in $X: N_X(v) := \{u \in X : u \sim v\}$, and the degree of v in X will be denoted by $\delta_X(v) = |N_X(v)|$. We denote the degree of a vertex $v_i \in V$ by $\delta(v_i)$ (or by d_i for short) and the degree sequence of Γ by $d_1 \ge d_2 \ge \cdots \ge d_n$. The subgraph induced by $S \subset V$ will be denoted by $\langle S \rangle$ and the complement of the set S in V will be denoted by \overline{S} .

A nonempty set $S \subseteq V$ is a defensive k-alliance in $\Gamma = (V, E), k \in \{-d_1, \ldots, d_1\}$, if for every $v \in S$,

$$\delta_S(v) \ge \delta_{\bar{S}}(v) + k. \tag{1}$$

A vertex $v \in S$ is said to be *k*-satisfied by the set S if (1) holds. Notice that (1) is equivalent to

$$\delta(v) \ge 2\delta_{\bar{S}}(v) + k. \tag{2}$$

A defensive (-1)-alliance is a *defensive alliance* and a defensive 0-alliance is a *strong defensive alliance* as defined in [12]. A defensive 0-alliance is also known as a *cohesive set* [21].

The defensive k-alliance number of Γ , denoted by $a_k(\Gamma)$, is defined as the minimum cardinality of a defensive k-alliance in Γ . Notice that

$$a_{k+1}(\Gamma) \ge a_k(\Gamma). \tag{3}$$

The defensive (-1)-alliance number of Γ is known as the alliance number of Γ and the defensive 0-alliance number is known as the strong alliance number, [12, 8, 9]. For instance, in the case of the 3-cube graph, $\Gamma = Q_3$, every set composed by two adjacent vertices is a defensive alliance of minimum cardinality and every set composed by four vertices whose induced subgraph is isomorphic to the cycle C_4 is a strong defensive alliance of minimum cardinality. Thus, $a_{-1}(Q_3) = 2$ and $a_0(Q_3) = 4$.

For some graphs, there are some values of $k \in \{-d_1, \ldots, d_1\}$, such that defensive k-alliances do not exist. For instance, for $k \geq 2$ in the case of the star graph S_n , defensive k-alliances do not exist. By (2) we conclude that, in any graph, there are defensive k-alliances for $k \in \{-d_1, \ldots, d_n\}$. For instance, a defensive (d_n) -alliance in $\Gamma = (V, E)$ is V. Moreover, if $v \in V$ is a vertex of minimum degree, $\delta(v) = d_n$, then $S = \{v\}$ is a defensive k-alliance for every $k \leq -d_n$. Therefore, $a_k(\Gamma) = 1$, for $k \leq -d_n$. For the study of the mathematical properties of $a_k(\Gamma)$, $k \in \{d_n, \ldots, d_1\}$, we cite [17].

A set $S \subset V$ is a *dominating set* in $\Gamma = (V, E)$ if for every vertex $u \in \overline{S}$, $\delta_S(u) > 0$ (every vertex in \overline{S} is adjacent to at least one vertex in S). The *domination number* of Γ , denoted by $\gamma(\Gamma)$, is the minimum cardinality of a dominating set in Γ .

A defensive k-alliance S is called global if it forms a dominating set. The global defensive k-alliance number of Γ , denoted by $\gamma_k^a(\Gamma)$, is the minimum cardinality of a defensive k-alliance in Γ . Clearly,

$$\gamma_{k+1}^{a}(\Gamma) \ge \gamma_{k}^{a}(\Gamma) \ge \gamma(\Gamma) \quad \text{and} \quad \gamma_{k}^{a}(\Gamma) \ge a_{k}(\Gamma).$$
 (4)

The global defensive (-1)-alliance number of Γ is known as the global alliance number of Γ and the global defensive 0-alliance number is known as the global strong alliance number [9]. For instance, in the case of the 3-cube graph, $\Gamma = Q_3$, every set composed by four vertices whose induced subgraph is isomorphic to the cycle C_4 is a global (strong) defensive alliance of minimum cardinality. Thus, $\gamma_{-1}^a(Q_3) = \gamma_0^a(Q_3) = 4$.

For some graphs, there are some values of $k \in \{-d_1, \ldots, d_1\}$, such that global defensive k-alliances do not exist. For instance, for $k = d_1$ in the case of nonregular graphs, defensive k-alliances do not exist. Therefore, the bounds

showed in this paper on $\gamma_k^a(\Gamma)$, for $k \leq d_1$, are obtained by supposing that the graph Γ contains defensive k-alliances. Notice that for any graph Γ , every dominating set is a global defensive $(-d_1)$ -alliance. Hence, $\gamma_{-d_1}^a(\Gamma) = \gamma(\Gamma)$. Moreover, for any d_1 -regular graph of order n, $\gamma_{d_1-1}^a(\Gamma) = \gamma_{d_1}^a(\Gamma) = n$.

2 Global defensive k-alliance number

Theorem 1. Let S be a global defensive k-alliance of minimum cardinality in Γ . If $W \subset S$ is a dominating set in Γ , then for every $r \in \mathbb{Z}$ such that $0 \leq r \leq \gamma_k^a(\Gamma) - |W|,$

$$\gamma^a_{k-2r}(\Gamma) + r \le \gamma^a_k(\Gamma).$$

Proof. We can take $X \subset S$ such that |X| = r. Hence, for every $v \in Y = S - X$,

$$\delta_Y(v) = \delta_S(v) - \delta_X(v)$$

$$\geq \delta_{\bar{S}}(v) + k - \delta_X(v)$$

$$= \delta_{\bar{Y}}(v) + k - 2\delta_X(v)$$

$$\geq \delta_{\bar{Y}}(v) + k - 2r.$$

Therefore, Y is a defensive (k - 2r)-alliance in Γ . Moreover, as $W \subset Y$, Y is a dominating set and, as a consequence, $\gamma_{k-2r}^a(\Gamma) \leq \gamma_k^a(\Gamma) - r$.

Notice that if every vertex of Γ has even degree and k is odd, k = 2l - 1, then every defensive (2l - 1)-alliance in Γ is a defensive (2l)-alliance. Hence, in such a case, $a_{2l-1}(\Gamma) = a_{2l}(\Gamma)$ and $\gamma_{2l-1}^a(\Gamma) = \gamma_{2l}^a(\Gamma)$. Analogously, if every vertex of Γ has odd degree and k is even, k = 2l, then every defensive (2l)-alliance in Γ is a defensive (2l + 1)-alliance. Hence, in such a case, $a_{2l}(\Gamma) = a_{2l+1}(\Gamma)$ and $\gamma_{2l}^a(\Gamma) = \gamma_{2l+1}^a(\Gamma)$. For instance, for the complete graph of order n we have

$$n = \gamma_{n-1}^{a}(K_{n}) = \gamma_{n-2}^{a}(K_{n})$$

$$\geq \gamma_{n-3}^{a}(K_{n}) = \gamma_{n-4}^{a}(K_{n}) = n - 1$$

...

$$\geq \gamma_{2-n}^{a}(K_{n}) = \gamma_{3-n}^{a}(K_{n}) = 2$$

$$\geq \gamma_{1-n}^{a}(K_{n}) = 1.$$

Therefore, for every $k \in \{1 - n, \dots, n - 1\}$, and for every $r \in \{0, \dots, \frac{k+n-1}{2}\}$,

$$\gamma_{k-2r}^{a}(K_{n}) + r = \gamma_{k}^{a}(K_{n}).$$

$$\tag{5}$$

Moreover, notice that for every $k \in \{1-n, \dots, n-1\}, \gamma_k^a(K_n) = \left\lceil \frac{n+k+1}{2} \right\rceil$.

It was shown in [9] that

$$\frac{\sqrt{4n+1}-1}{2} \le \gamma_{-1}^a(\Gamma) \le n - \left\lceil \frac{d_n}{2} \right\rceil$$
(6)

and

$$\sqrt{n} \le \gamma_0^a(\Gamma) \le n - \left\lfloor \frac{d_n}{2} \right\rfloor.$$
 (7)

Here we generalize the previous results to defensive k-alliances.

Theorem 2. For any graph
$$\Gamma$$
, $\frac{\sqrt{4n+k^2}+k}{2} \leq \gamma_k^a(\Gamma) \leq n - \left\lfloor \frac{d_n-k}{2} \right\rfloor$.

Proof. If $d_n \leq k$, then $\gamma_k^a(\Gamma) \leq n \leq n - \lfloor \frac{d_n - k}{2} \rfloor$. Otherwise, consider $u \in V$ such that $\delta(u) \geq \lfloor \frac{d_n + d_1}{2} \rfloor$. Let $X \subset V$ be the set of neighbors u has in Γ , $X = \{w \in V : w \sim u\}$. Let $Y \subset X$ be a vertex set such that $|Y| = \lfloor \frac{d_n - k}{2} \rfloor$. In such a case, the set V - Y is a global defensive k-alliance in Γ . That is, V - Y is a dominating set and for every $v \in V - Y$ we have $\frac{\delta(v) - k}{2} \geq \lfloor \frac{d_n - k}{2} \rfloor \geq \delta_Y(v)$. Therefore, $\gamma_k^a(\Gamma) \leq n - \lfloor \frac{d_n - k}{2} \rfloor$.

On the other hand, let $S \subseteq V$ be a dominating set in Γ . Then,

$$n - |S| \le \sum_{v \in S} \delta_{\bar{S}}(v). \tag{8}$$

Moreover, if S is a defensive k-alliance in Γ ,

$$k|S| + \sum_{v \in S} \delta_{\bar{S}}(v) \le \sum_{v \in S} \delta_{S}(v) \le |S|(|S| - 1).$$
(9)

Hence, solving

$$0 \le |S|^2 - k|S| - n \tag{10}$$

we deduce the lower bound.

5

The upper bound is attained, for instance, for the complete graph $\Gamma = K_n$ for every $k \in \{1 - n, \dots, n - 1\}$. The lower bound is attained, for instance, for the 3-cube graph $\Gamma = Q_3$, in the following cases: $2 \leq \gamma_{-3}^a(Q_3)$ and $4 \leq \gamma_1(Q_3) = \gamma_0(Q_3)$.

It was shown in [9] that for any bipartite graph Γ of order n and maximum degree d_1 ,

$$\gamma_{-1}^{a}(\Gamma) \ge \left\lceil \frac{2n}{d_1+3} \right\rceil \quad \text{and} \quad \gamma_{0}^{a}(\Gamma) \ge \left\lceil \frac{2n}{d_1+2} \right\rceil$$

Here we generalize the previous bounds to defensive k-alliances. Moreover, we show that the result is not restrictive to the case of bipartite graphs.

Theorem 3. For any graph
$$\Gamma$$
, $\gamma_k^a(\Gamma) \ge \left\lceil \frac{n}{\lfloor \frac{d_1-k}{2} \rfloor + 1} \right\rceil$.

Proof. If S denotes a defensive k-alliance in Γ , then

$$d_1 \ge \delta(v) \ge 2\delta_{\bar{S}}(v) + k, \quad \forall v \in S.$$

Therefore,

$$\left\lfloor \frac{d_1 - k}{2} \right\rfloor \ge \delta_{\bar{S}}(v), \quad \forall v \in S.$$
(11)

Hence,

$$|S| \left\lfloor \frac{d_1 - k}{2} \right\rfloor \ge \sum_{v \in S} \delta_{\bar{S}}(v).$$
(12)

Moreover, if S is a dominating set, S satisfies inequality (8). The result follows by (8) and (12). \Box

The above bound is tight. For instance, for the Petersen graph the bound is attained for every $k: 3 \leq \gamma_{-3}^{a}(\Gamma), 4 \leq \gamma_{-2}^{a}(\Gamma) = \gamma_{-1}^{a}(\Gamma), 5 \leq \gamma_{0}(\Gamma) = \gamma_{1}(\Gamma)$ and $10 \leq \gamma_{2}(\Gamma) = \gamma_{3}(\Gamma)$. For the 3-cube graph $\Gamma = Q_{3}$, the above theorem leads to the following exact values of $\gamma_{k}^{a}(Q_{3}): 2 \leq \gamma_{-3}^{a}(Q_{3}), 4 \leq \gamma_{0}(Q_{3}) = \gamma_{1}(Q_{3})$ and $8 \leq \gamma_{2}(Q_{3}) = \gamma_{3}(Q_{3})$.

Hereafter, we denote by $\mathcal{L}(\Gamma) = (V_l, E_l)$ the line graph of a simple graph Γ . The degree of the vertex $e = \{u, v\} \in V_l$ is $\delta(e) = \delta(u) + \delta(v) - 2$. If the degree sequence of Γ is $d_1 \geq d_2 \geq \cdots \geq d_n$, then the maximum degree of $\mathcal{L}(\Gamma)$, denoted by Δ_l , is bounded by $\Delta_l \leq d_1 + d_2 - 2$.

Corollary 4. For any graph Γ of size m and maximum degrees $d_1 \geq d_2$,

$$\gamma_k^a(\mathcal{L}(\Gamma)) \ge \left\lceil \frac{m}{\left\lfloor \frac{d_1+d_2-2-k}{2} \right\rfloor + 1} \right\rceil.$$

The above bound is attained for $k \in \{-3, -2, -1, 2, 3\}$ in the case of the complete bipartite graph $\Gamma = K_{1,4}$. Notice that $\mathcal{L}(K_{1,4}) = K_4$ and $\gamma_{-3}^a(K_4) = 1, \ \gamma_{-2}^a(K_4) = \gamma_{-1}^a(K_4) = 2, \ \gamma_2^a(K_4) = \gamma_3^a(K_4) = 4.$

In the case of cubic graphs¹ $\gamma(\Gamma) = \gamma_{-3}^{a}(\Gamma) \leq \gamma_{-2}^{a}(\Gamma) = \gamma_{-1}^{a}(\Gamma) \leq \gamma_{0}^{a}(\Gamma) = \gamma_{1}^{a}(\Gamma) \leq \gamma_{2}^{a}(\Gamma) = \gamma_{3}^{a}(\Gamma) = n$. So, in this case we only study, $\gamma_{-1}^{a}(\Gamma)$ and $\gamma_{0}^{a}(\Gamma)$.

Theorem 5. For any cubic graph Γ , $\gamma_{-1}^{a}(\Gamma) \leq 2\gamma(\Gamma)$.

Proof. Let S be a dominating set of minimum cardinality in Γ . Let $X \subseteq S$ be the set composed by all $v_i \in S$ such that $\delta_S(v_i) = 0$. For each $v_i \in X$ we take a vertex $u_i \in \overline{S}$ such that $u_i \sim v_i$. Let $Y \subseteq \overline{S}$ defined as $Y = \bigcup_{v_i \in X} \{u_i\}$. Then we have $|Y| \leq \gamma(\Gamma)$ and the set $S \cup Y$ is a global defensive (-1)-alliance in Γ .

The above bound is tight. For instance, in the case of the 3-cube graph we have $\gamma_{-1}^{a}(Q_3) = 2\gamma(Q_3) = 4$.

A set $S \subset V$ is a total dominating set if every vertex in V has a neighbor in S. The total domination number $\gamma_t(\Gamma)$ is the minimum cardinality of a total dominating set in Γ . Notice that if Γ is a cubic graph, then

$$\gamma_{-1}^{a}(\Gamma) = \gamma_{t}(\Gamma). \tag{13}$$

It was shown in [5] that if Γ is a connected graph of order $n \geq 3$, then

$$\gamma_t(\Gamma) \le \frac{2n}{3}.\tag{14}$$

Moreover, by Theorem 3 we have

$$\frac{n}{3} \le \gamma_{-1}^{a}(\Gamma) \quad \text{and} \quad \frac{n}{2} \le \gamma_{0}^{a}(\Gamma).$$
(15)

 $^{^1\}mathrm{A}$ cubic graph is a 3-regular graph.

3 Defensive *k*-alliances in planar graphs

It is well-known that the size of a planar graph Γ of order $n \geq 3$ is bounded by $m \leq 3(n-2)$. Moreover, in the case of triangle-free graphs $m \leq 2(n-2)$. This inequalities allow us to obtain tight bounds for the studied parameters.

Theorem 6. Let $\Gamma = (V, E)$ be a graph of order n. If Γ has a global defensive k-alliance S such that the subgraph $\langle S \rangle$ is planar.

- (i) If n > 2(2-k), then $|S| \ge \left\lceil \frac{n+12}{7-k} \right\rceil$.
- (ii) If n > 2(2-k) and $\langle S \rangle$ is a triangle-free graph, then $|S| \ge \left\lceil \frac{n+8}{5-k} \right\rceil$.

Proof.

(i) If $|S| \leq 2$, for every $v \in S$ we have $\delta_{\bar{S}}(v) \leq 1 - k$. Thus, $n \leq 2(2 - k)$. Therefore, $n > 2(2 - k) \Rightarrow |S| > 2$.

If $\langle S \rangle$ is planar and |S| > 2, the size of $\langle S \rangle$ is bounded by

$$\frac{1}{2}\sum_{v\in S}\delta_S(v) \le 3(|S|-2).$$
(16)

If S is a global defensive k-alliance in Γ ,

$$k|S| + (n - |S|) \le k|S| + \sum_{v \in S} \delta_{\bar{S}}(v) \le \sum_{v \in S} \delta_{S}(v).$$
(17)

By (16) and (17) the result follows.

(ii) If $\langle S \rangle$ is a triangle-free graph, then

$$\frac{1}{2} \sum_{v \in S} \delta_S(v) \le 2(|S| - 2).$$
(18)

The result follows by (17) and (18).

Corollary 7. For any planar graph Γ of order n.

(a) If n > 2(2-k), then $\gamma_k^a(\Gamma) \ge \left\lceil \frac{n+12}{7-k} \right\rceil$.

(b) If n > 2(2-k) and Γ is a triangle-free graph, then $\gamma_k^a(\Gamma) \ge \left\lceil \frac{n+8}{5-k} \right\rceil$.

The above bounds are tight. In the case of the graph of Figure 1, the set $S = \{1, 2, 3\}$ is a global defensive k-alliance for k = -2, k = -1 and k = 0, and Corollary 7-(a) leads to $\gamma_k^a(\Gamma) \geq 3$. Moreover, if $\Gamma = Q_3$, the 3-cube graph, Corollary 7-(b) leads to the following exact values of $\gamma_k^a(Q_3)$: $2 \leq \gamma_{-3}^{a}(Q_3)$, $4 \leq \gamma_{0}^{a}(Q_3) = \gamma_{1}^{a}(Q_3)$ and $8 \leq \gamma_{3}^{a}(Q_3)$.

Theorem 8. Let Γ be a graph of order n. If Γ has a global defensive kalliance S such that the subgraph $\langle S \rangle$ is planar connected with f faces. Then,

$$|S| \ge \left\lceil \frac{n-2f+4}{3-k} \right\rceil.$$

Proof. By Euler's formula, $\sum_{v \in S} \delta_S(v) = 2(|S| + f - 2)$, and (17) we deduce the result.

In the case of the graph of Figure 1, the set $S = \{1, 2, 3\}$ is a global defensive k-alliance for k = -1, k = 0 and k = 2. Moreover, $\langle S \rangle$ has two faces. In such a case, Theorem 8 leads to $|S| \ge 3$.



3.1Defensive k-alliances in trees

In this section we study global defensive k-alliances in trees but we impose a condition on the number of connected components of the subgraphs induced by the alliances.

Theorem 9. Let T be a tree of order n. Let S be a global defensive k-alliance in T such that the subgraph $\langle S \rangle$ has c connected components. Then,

$$|S| \ge \left\lceil \frac{n+2c}{3-k} \right\rceil$$

Proof. As the subgraph $\langle S \rangle$ is a forest with c connected components,

$$\sum_{v \in S} \delta_S(v) = 2(|S| - c).$$
(19)

The bound of |S| follows from (17) and (19).



Figure 2:

The above bound is attained, for instance, for the left hand side graph of Figure 2, where $S = \{1, 2, 3, 4\}$ is a global defensive (-1)-alliance and $\langle S \rangle$ has two connected components. Moreover, the bound is attained in the case of the right hand side graph of Figure 2, where $S = \{1, 2, 3, 4, 5\}$ is a global defensive 0-alliance and $\langle S \rangle$ has two connected components.

Corollary 10. For any tree T of order n, $\gamma_k^a(T) \ge \left\lceil \frac{n+2}{3-k} \right\rceil$.

The above bound is attained for $k \in \{-4, -3, -2, 0, 1\}$ in the case of $\Gamma = K_{1,4}$. As a particular case of above theorem we obtain the bounds obtained in [9]:

$$\gamma_{-1}^{a}(T) \ge \left\lceil \frac{n+2}{4} \right\rceil$$
 and $\gamma_{0}^{a}(T) \ge \left\lceil \frac{n+2}{3} \right\rceil$

4 Global connected defensive *k*-alliances

It is clear that a defensive k-alliance of minimum cardinality must induce a connected subgraph. But we can have a global defensive k-alliance of minimum cardinality with nonconnected induced subgraph. We say that a defensive k-alliance S is connected if $\langle S \rangle$ is connected. We denote by $\gamma_k^{ca}(\Gamma)$ the minimum cardinality of a global connected defensive k-alliance in Γ . Obviously, $\gamma_k^{ca}(\Gamma) \geq \gamma_k^a(\Gamma)$. For instance, for the left hand side graph of Figure 2 we have $\gamma_{-1}^{ca}(\Gamma) = 5 > 4 = \gamma_{-1}^a(\Gamma)$ and for the right hand side graph of Figure 2 we have $\gamma_{0}^{ca}(\Gamma) = 6 > 5 = \gamma_0^a(\Gamma)$.

Theorem 11. For any connected graph Γ of diameter $D(\Gamma)$,

(i)
$$\gamma_k^{ca}(\Gamma) \ge \left\lceil \frac{\sqrt{4(D(\Gamma)+n-1)+(1-k)^2}+(k-1)}{2} \right\rceil$$

(ii) $\gamma_k^{ca}(\Gamma) \ge \left\lceil \frac{n+D(\Gamma)-1}{\lfloor \frac{\Delta-k}{2} \rfloor+2} \right\rceil$.

Proof. If S is a dominating set in Γ such that $\langle S \rangle$ is connected, then $D(\Gamma) \leq D(\langle S \rangle) + 2$. Hence,

$$D(\Gamma) \le |S| + 1. \tag{20}$$

Moreover, if S is a global defensive k-alliance in Γ , then |S| satisfies (10). The first result follows by (10) and (20).

As a consequence of (8), (11) and (20) we obtain the second result. \Box

Both bounds in Theorem 11 are tight. For instance, both bounds are attained for $k \in \{-2, -1, 0\}$ for the graph of Figure 1. In such a case, both bounds lead to $\gamma_k^{ca}(\Gamma) \geq 3$. Moreover, both bounds lead to the exact values of $\gamma_k^{ca}(K_{3,3})$ in the following cases: $2 \leq \gamma_{-3}^{ca}(K_{3,3}) = \gamma_{-2}^{ca}(K_{3,3}) = \gamma_{-1}^{ca}(K_{3,3})$. Furthermore, notice that bound (ii) leads to the exact values of $\gamma_k^{ca}(Q_3)$ in the cases $4 \leq \gamma_0^{ca}(Q_3) = \gamma_1^{ca}(Q_3)$, while bound (i) only gives $3 \leq \gamma_0^a(Q_3)$ and $3 \leq \gamma_1^{ca}(Q_3)$.

By Theorem 11, and taking into a count that $D(\Gamma) - 1 \leq D(\mathcal{L}(\Gamma))$, we obtain the following result on the global connected k-alliance number of the line graph of Γ in terms of some parameters of Γ .

Corollary 12. For any connected graph Γ of size m, diameter $D(\Gamma)$, and maximum degrees $d_1 \geq d_2$,

(i)
$$\gamma_k^{ca}(\mathcal{L}(\Gamma)) \ge \left\lceil \frac{\sqrt{4(D(\Gamma)+m-2)+(1-k)^2}-(1-k)}{2} \right\rceil$$

(ii) $\gamma_k^{ca}(\mathcal{L}(\Gamma)) \ge \left\lceil \frac{2(m+D(\Gamma)-2)}{d_1+d_2-k+1} \right\rceil$.

References

- A. Cami, H. Balakrishnan, N. Deo, and R. D. Dutton, On the complexity of finding optimal global alliances. J. Combin. Math. Combin. Comput. 58 (2006) 23-31.
- [2] R. C. Brigham, R. D. Dutton and S. T. Hedetniemi, A sharp lower bound on the powerful alliance number of $C_m \times C_n$. Congr. Numer. 167 (2004), 57-63.
- [3] R. C. Brigham, R. D. Dutton, T. W. Haynes, and S. T. Hedetniemi, Powerful alliances in graphs. *Discrete Appl. Math.*, to appear.
- [4] M. Chellali and T. Haynes, Global alliances and independence in trees. Discussiones Mathematicae Graph Theory 27 (1) (2007) 19-27.
- [5] E. J. Cockayne, R. Dawes and S. T. Hedetniemi, Total domination in graphs, *Networks* 10 (1980), 211-215.
- [6] O. Favaron, G. Fricke, W. Goddard, S. M. Hedetniemi, S. T. Hedetniemi, P. Kristiansen, R. C. Laskar, R. D. Skaggs, Offensive alliances in graphs. *Discussiones Mathematicae Graph Theory* 24 (2) (2004) 263-275.
- [7] H. Fernau and D. Raible, Alliances in Graphs: a Complexity-Theoretic Study. SOFSEM 2 (2007) 61-70.
- [8] G. H. Fricke, L. M. Lawson, T. W. Haynes, S. M. Hedetniemi and S. T. Hedetniemi, A Note on Defensive Alliances in Graphs. Bull. Inst. Combin. Appl. 38 (2003), 37-41.
- [9] T. W. Haynes, S. T. Hedetniemi and M. A. Henning, Global defensive alliances in graphs. *Electron. J. Combin.* 10 (2003) 139-146.
- [10] L. H. Jamieson, S. T. Hedetniemi, and A. A. McRae, The algorithmic complexity of alliances in graphs. J. Combin. Math. Combin. Comput., to appear, 2007.

- [11] L. H. Jamieson. Algorithms and Complexity for Alliances and Weighted Alliances of Different Types. PhD thesis, Clemson University, 2007.
- [12] P. Kristiansen, S. M. Hedetniemi and S. T. Hedetniemi, Alliances in graphs. J. Combin. Math. Combin. Comput. 48 (2004) 157-177.
- [13] J. A. Rodríguez and J. M. Sigarreta, Offensive alliances in cubic graphs. International mathematical forum 1 (36) (2006) 1773-1782.
- [14] J. A. Rodríguez and J. M. Sigarreta, Spectral study of alliances in graphs. Discussiones Mathematicae Graph Theory 27 (1) (2007) 143-157.
- [15] J. A. Rodríguez-Velázquez and J. M. Sigarreta, Global offensive alliances in graphs. *Electronic Notes in Discrete Mathematics* 25 (2006) 157-164.
- [16] J. A. Rodríguez-Velázquez, J. M. Sigarreta, Global alliances in planar graphs. AKCE Int. J. Graphs Comb. 4 (1) (2007) 83-98.
- [17] J. A. Rodríguez-Velázquez, I. G. Yero and J. M. Sigarreta, Defensive k-alliances in graphs. Submitted.
- [18] K. H. Shafique and R. D. Dutton, Maximum alliance-free and minimum alliance-cover sets. *Congr. Numer.* 162 (2003) 139-146.
- [19] K. H. Shafique and R. Dutton, A tight bound on the cardinalities of maximun alliance-free and minimun alliance-cover sets. J. Combin. Math. Combin. Comput. 56 (2006), 139-145.
- [20] K. H. Shafique, Partitioning a Graph in Alliances and its Application to Data Clustering. Ph. D. Thesis, 2004.
- [21] K. H. Shafique and R. D. Dutton, On satisfactory partitioning of graphs. Congr. Numer. 154 (2002) 183-194.
- [22] J. M. Sigarreta and J. A. Rodríguez, On defensive alliance and line graphs. Appied Mathematics Letters 19 (12) (2006) 1345-1350.
- [23] J. M. Sigarreta and J. A. Rodríguez, On the global offensive alliance number of a graph. Submitted.



This figure "f1a.jpg" is available in "jpg" format from:

This figure "f1b.jpg" is available in "jpg" format from:

This figure "f2a.jpg" is available in "jpg" format from:

This figure "f2b.jpg" is available in "jpg" format from: