

# THE MINIMAL LOG DISCREPANCY

FLORIN AMBRO

ABSTRACT. We survey the known and expected properties of the minimal log discrepancy, the local invariant of a log variety.

## CONTENTS

Introduction	1
1. Background	2
2. Log varieties, minimal log discrepancies	5
3. Problems on minimal log discrepancies	8
4. The log canonical threshold	11
References	13

## INTRODUCTION

The minimal log discrepancy is a fundamental invariant of the singularities that appear in the birational classification of algebraic varieties. Introduced by Shokurov [52] in connection to the termination of a sequence of flips, it appears in the local context of the classification of singularities, or the global context of Fujita's conjecture on adjoint linear systems. The minimal log discrepancy measures stable vanishing orders of sections of canonical graded rings, or the rate of growth of certain subspaces in the spaces of jets of a singularity. It has an arithmetic flavour, being related to the first minimum of Minkowski in the geometry of numbers.

In this note we introduce the minimal log discrepancy, present some basic open problems, and illustrate them with toric examples. We hope to reinforce the original connection of Reid [50] between discrepancies of singularities on the one hand, and stable vanishing orders of sections of canonical graded rings on the other hand.

The minimal log discrepancy is the local invariant of a *log variety*. We recall in §1 the construction of canonical models and discrepancies, and their logarithmic version. This is the natural motivation for log varieties with *log canonical singularities*, which locally are just open subsets of log canonical models. We give the rigorous definition of log varieties and minimal log discrepancies in §2, and present explicit combinatorial formulas in the toric case. We present problems on minimal log discrepancies in §3, and discuss their toric case and some methods, old and new. In §4, we discuss the log canonical threshold and its applications to the problem of constructing *flat log structures*.

---

<sup>1</sup>The author is supported by a 21st Century COE Kyoto Mathematics Fellowship, and by the JSPS Grant-in-Aid No 17740011.

## 1. BACKGROUND

1-A. **Zariski decomposition.** The origin of Zariski decomposition is the problem of computing the Hilbert function  $\varphi(n) = \dim_{\mathbb{C}} R(X, D)_n$  of the graded ring

$$R(X, D) = \bigoplus_{n=0}^{\infty} H^0(X, nD),$$

for a given divisor  $D$  on a complex manifold  $X$  [61]. We recall the solution to this problem in the case when  $R(X, D)$  is finitely generated and  $D$  is big. We use Zariski's notation

$$H^0(X, D) = \{a \in \mathbb{C}(X)^{\times}; (a) + D \geq 0\} \cup \{0\},$$

which makes sense even if  $D$  does not have integer coefficients.

First, if  $D$  is an ample divisor, there exists a polynomial  $P \in \mathbb{Q}[T]$  of degree  $d = \dim(X)$ , and a positive integer  $n_0$ , such that  $\varphi(n) = P(n)$  for  $n \geq n_0$ . In general, finite generation means that there exists a positive integer  $r$  such that the natural map  $S^l H^0(X, rD) \rightarrow H^0(X, lrD)$  is surjective for every  $l \geq 1$ . By Hironaka's resolution of the base locus of a linear system, there exists a birational modification  $f_r: X_r \rightarrow X$  such that if  $F_r$  is the fixed divisor of  $|f_r^*(rD)|$ , the mobile part  $M_r = f_r^*(rD) - F_r$  defines a linear system without base points. After taking the Stein factorization,  $|M_r|$  defines a morphism with connected fibers  $g_r: X_r \rightarrow Y_r$ . The variety  $Y_r$  has normal singularities and  $M_r = g_r^*(A_r)$ , for a normally generated ample Cartier divisor  $A_r$  on  $Y$ .

$$\begin{array}{ccc} & X_r & \\ f_r \swarrow & & \searrow g_r \\ X & & Y_r \end{array}$$

We have  $R(X, D) = R(X_r, f_r^*(D))$ , the natural inclusion  $R(X_r, \frac{1}{r}M_r) \subseteq R(X_r, f_r^*(D))$  becomes an identity, and  $R(X_r, \frac{1}{r}M_r) = R(Y_r, \frac{1}{r}A_r)$ . In particular,  $Y_r = \text{Proj } R(X, D)$  and

$$\varphi(n) = \dim_{\mathbb{C}} H^0(Y_r, \frac{n}{r}A_r).$$

There are polynomials  $P_0, \dots, P_{r-1} \in \mathbb{Q}[T]$ , of degree  $d$ , and a positive integer  $n_0$ , such that  $\varphi(n) = P_{n-r \lfloor \frac{n}{r} \rfloor}(n)$  for  $n \geq n_0$ . The growth rate of  $\varphi$  is

$$\lim_{n \rightarrow \infty} \frac{\varphi(n)}{n^d/d!} = \frac{(A_r^d)}{r^d} \in \frac{1}{r^d} \mathbb{N}.$$

The decomposition  $f_r^*(D) = g_r^*(\frac{1}{r}A) + \frac{1}{r}F_r$  is the prototype of a Zariski decomposition. In general, we say that  $D = P + F$  is a *Zariski decomposition* on a normal variety  $X$ , if  $F$  is an effective  $\mathbb{R}$ -divisor and  $P$  is a nef  $\mathbb{R}$ -divisor such that the induced inclusion  $R(X, P) \subseteq R(X, D)$  is surjective. The Zariski decomposition is unique if  $D$  is big, and in this case  $F = \lim_{n \rightarrow \infty} \frac{1}{n}F_n$ , where  $F_n$  is the fixed part of  $|nD|$ . There are several notions of Zariski decomposition at present, but they all coincide for big divisors (see [61, 29, 22, 15, 46]).

Two useful examples of Zariski decomposition are as follows. For a nef  $\mathbb{R}$ -divisor  $P$ , the decomposition  $P = P + 0$  is a Zariski decomposition. If  $D = P + F$  is a Zariski decomposition on  $X$  and  $F'$  is an effective  $\mathbb{R}$ -divisor supported by the exceptional locus of a birational map  $\mu: X' \rightarrow X$ , then  $\mu^*D + E = \mu^*P + (\mu^*F + F')$  is a Zariski decomposition.

**1-B. Canonical models, discrepancies.** Let  $X$  be a complex projective manifold of general type, with *canonical divisor*  $K_X$ . The *canonical ring*  $R(X, K_X)$  is expected to be finitely generated, and if it is, we would obtain a natural birational map

$$\Phi: X \dashrightarrow Y := \text{Proj } R(X, K_X).$$

The birational model  $Y$  is called the *canonical model* of  $X$ . It depends only on the birational class of  $X$  and it has a canonical polarization, but it has singularities in general. For example,  $Y$  may have some Du Val singularities in dimension two. The singularities that may appear on  $Y$  were coined *canonical singularities* by Reid [50].

To get to the formal definition of canonical singularities, let us take a closer look at what  $\Phi$  does for  $K_X$ . By Hironaka's resolution of singularities, there is a *Hironaka hut*

$$\begin{array}{ccc} & X' & \\ f \swarrow & & \searrow g \\ X & \xrightarrow{\Phi} & Y \end{array}$$

that is  $X'$  is a projective manifold,  $f, g$  are birational morphisms and  $\Phi = g \circ f^{-1}$ . By definition,  $K_X$  is the divisor ( $\omega$ ) of zeros and poles of a non-zero top rational differential form  $\omega \in \wedge^{\dim(X)} \Omega_X^1 \otimes_{\mathbb{C}} \mathbb{C}(X)$ . Denote  $K_{X'} = (f^*\omega)$  and  $K_Y = (g_*f^*\omega)$ . The latter is a well defined Weil divisor, since  $Y$  is normal. Since  $X$  has no singularities, the divisor  $A_f = K_{X'} - f^*(K_X)$  is effective and supported by the exceptional locus of  $f$ . Equivalently, the natural map  $f_*: R(X', K_{X'}) \rightarrow R(X, K_X)$  is an isomorphism. In particular,  $g: X' \rightarrow Y$  is the canonical model of  $X'$ . Since  $g$  is a morphism and  $K_{X'}$  is a big divisor, it follows that there exists  $r \geq 1$  such that  $|rK_Y|$  defines a projectively normal embedding, and  $A_g = \frac{1}{r}(rK_{X'} - g^*(rK_Y))$  is effective and supported by the exceptional locus of  $g$ . In particular,  $g_*: R(X', K_{X'}) \rightarrow R(Y, K_Y)$  is also an isomorphism:

$$\begin{array}{ccc} & R(X', K_{X'}) & \\ \simeq \swarrow & & \searrow \simeq \\ R(X, K_X) & & R(Y, K_Y) \end{array}$$

Reid [50] called a normal germ  $P \in Y$  a canonical singularity if  $A_g$  is well defined and effective, for a resolution of singularities  $g: X' \rightarrow Y$ . The coefficients of the  $\mathbb{Q}$ -divisor  $A_g$  are called *discrepancies*. To understand discrepancies in terms of the manifolds that we started with, we go back to our global setting and note that

$$K_{X'} = g^*(K_Y) + A_g$$

is a Zariski decomposition of  $K_{X'}$ , with positive part  $g^*(K_Y)$  and fixed part  $A_g$ . Since  $|rK_Y|$  defines a linear system free of base points,  $rA_g$  is the fixed part of the linear system  $|rK_{X'}|$ . Finally, it turns out that  $f^*(K_X) = g^*(K_Y) + (A_g - A_f)$  is a Zariski decomposition.

**1-C. Log canonical models of open manifolds.** Let  $U$  be a complex quasi-projective manifold of general type, in the sense of Iitaka [25]. By Hironaka's resolution of singularities, there exists an open embedding  $U \subset X$  such that  $X$  is a proper manifold, and the complement  $X \setminus U = \sum_i E_i$  is a divisor with simple normal crossings. The general type assumption means that the *log canonical divisor*  $K_X + \sum_i E_i$  is big. The *log canonical ring*  $R(X, K_X + \sum_i E_i)$  is independent of the choice of compactification, and in fact depends

only on the (proper) birational class of  $U$ . It is expected to be finitely generated, and if it is, we would obtain a natural birational map

$$\Phi: X \dashrightarrow Y := \text{Proj } R(X, K + \sum_i E_i).$$

As before, we can find a Hironaka hut with the extra property that  $\text{Exc}(f) \cup f_*^{-1}(\sum_i E_i)$  is a simple normal crossings divisor  $\sum_{i'} E_{i'}$ . Denote  $B_Y = g_*(\sum_{i'} E_{i'})$ . We imitate the arguments in the compact case, and obtain isomorphisms

$$\begin{array}{ccc} & R(X', K_{X'} + \sum_{i'} E_{i'}) & \\ \swarrow \cong & & \searrow \cong \\ R(X, K_X + \sum_i E_i) & & R(Y, K_Y + B_Y) \end{array}$$

Again,  $|r(K_Y + B_Y)|$  defines a normally generated embedding for some  $r \geq 1$ , and we have Zariski decompositions  $K_{X'} + \sum_{i'} E_{i'} = g^*(K_Y + B_Y) + A_g$  and  $f^*(K_X + \sum_i E_i) = g^*(K_Y + B_Y) + (A_g - A_f)$ . One can see that  $\Phi^{-1}$  contracts no divisors of  $Y$ , and  $\Phi_*(\sum_i E_i) = B_Y$ . The pair  $(Y, B_Y)$  is log canonically polarized, and its singularities are log canonical, as we will see shortly. The pair  $(Y, B_Y)$  is called the *log canonical model* of  $U$ .

For example, let  $U \rightarrow M_g$  be a resolution of singularities of the moduli space of smooth curves of genus  $g \geq 2$ . Then the log canonical model of  $U$  is  $(\overline{M}_g, \delta)$ , where  $\overline{M}_g$  is the moduli space of stable curves of genus  $g$ , and  $\delta = \overline{M}_g \setminus M_g$  [14].

**1-D. Log canonical models of log manifolds, log discrepancies.** *Log manifolds* are the bridge between open and compact manifolds. They are pairs  $(X, \sum_i b_i E_i)$ , where  $X$  is nonsingular, the  $E_i$ 's are nonsingular divisors intersecting transversely, and  $b_i \in [0, 1] \cap \mathbb{Q}$  for all  $i$ . We call  $\sum_i b_i E_i$  the boundary of the log manifold, and denote it by  $B$ . Suppose moreover that  $(X, B)$  is of general type, that is the log canonical divisor  $K_X + B$  is big. The log canonical ring  $R(X, K_X + B)$  is expected to be finitely generated, and if it is, we would obtain a birational map

$$\Phi: X \dashrightarrow Y := \text{Proj } R(X, K_X + B).$$

Again, we construct a Hironaka hut with the extra property that  $\text{Exc}(f) \cup f_*^{-1}(\sum_i E_i)$  is a simple normal crossings divisor. Let  $\cup_j F_j$  be the exceptional locus of  $f$  and denote  $B_Y = g_*(f_*^{-1}B + \sum_j F_j)$ . We imitate the previous argument, and obtain isomorphisms

$$\begin{array}{ccc} & R(X', K_{X'} + f_*^{-1}B + \sum_j F_j) & \\ \swarrow \cong & & \searrow \cong \\ R(X, K_X + B) & & R(Y, K_Y + B_Y) \end{array}$$

There exists  $r \in \mathbb{Z}_{\geq 1}$  such that  $|r(K_Y + B_Y)|$  defines a normally generated embedding, and we have Zariski decompositions

$$K_{X'} + f_*^{-1}B + \sum_j F_j = g^*(K_Y + B_Y) + A_g$$

$$f^*(K_X + B) = g^*(K_Y + B_Y) + (A_g - A_f).$$

One can see that  $\Phi^{-1}$  contracts no divisors of  $Y$ , and  $\Phi_*(B) = B_Y$ . The birational model  $\Phi: (X, B) \dashrightarrow (Y, B_Y)$  is called the *log canonical model* of  $(X, B)$ . It is polarized by the

*log canonical divisor*  $K_Y + B_Y$  (a  $\mathbb{Q}$ -divisor), and its singularities are called *log canonical singularities*. The coefficients of the  $\mathbb{Q}$ -divisor  $A_g$  are called *log discrepancies*.

## 2. LOG VARIETIES, MINIMAL LOG DISCREPANCIES

Log varieties with log canonical singularities are locally open subsets of log canonical models. For technical purposes, it is better to work in a slightly more general context, such as non-rational boundaries (to take convex hulls and limits of divisors), or even non-log canonical singularities (to construct a flat log structure at a prescribed point of a polarized manifold, cf §4-A).

**Definition 2.1.** A *log variety*  $(X, B)$  is a complex normal variety  $X$  endowed with an effective  $\mathbb{R}$ -Weil divisor  $B = \sum_i b_i E_i$  such that  $K_X + B$  is  $\mathbb{R}$ -Cartier.

Recall that the canonical divisor  $K_X = (\omega)$  is the Weil divisor of zeros and poles of a non-zero top rational differential form  $\omega$  (it depends on the choice of  $\omega$ , but only up to linear equivalence). The  $E_i$ 's are prime divisors and the  $b_i$ 's are non-negative real numbers. The  $\mathbb{R}$ -Cartier assumption means that locally on  $X$ ,  $K_X + B$  equals a finite sum  $\sum_i r_i(\varphi_i)$ , where  $r_i \in \mathbb{R}$  and  $\varphi_i \in \mathbb{C}(X)^\times$ .

Let now  $\mu: X' \rightarrow X$  be birational morphism, and  $E \subset X'$  a prime divisor. We use the same form to define the canonical class of  $X'$ , that is  $K_{X'} = (f^*\omega)$ . The *log discrepancy* of  $(X, B)$  at  $E$  is defined as

$$a(E; X, B) = \text{mult}_E(K_{X'} + E - \mu^*(K_X + B)) \in \mathbb{R}.$$

The log discrepancy depends only on the valuation that  $E$  induces on  $\mathbb{C}(X)$ . We call such valuations *geometric*, and denote  $c_X(E) = \mu(E)$ . For example, if  $E$  is a prime divisor in  $X$ , then  $a(E; X, B) = 1 - \text{mult}_E(B)$ .

**Definition 2.2.** The *minimal log discrepancy* of a log variety  $(X, B)$  at a Grothendieck point  $\eta \in X$  is defined as

$$a(\eta; X, B) = \inf\{a(E; X, B); c_X(E) = \bar{\eta}\} \in \{-\infty\} \cup \mathbb{R}_{\geq 0}.$$

The *global minimal log discrepancy* is  $a(X, B) = \inf_{\eta \in X} a(\eta; X, B)$ .

**Example 2.3.**  $a(0; \mathbb{C}^d) = d$ .

The reader may check that  $a(\eta; X, B) < 0$  implies  $a(\eta; X, B) = -\infty$ . Otherwise,  $a(\eta; X, B)$  is a non-negative real number, and the infimum is a minimum. If  $\eta$  has codimension one, then  $a(\eta; X, B) = 1 - \text{mult}_\eta(B)$ . Otherwise, construct a log resolution  $\mu: X' \rightarrow X$  such that  $\mu^{-1}(\bar{\eta})$  is a divisor, and  $\mu^{-1}(\bar{\eta}), \mu_*^{-1}B$  and  $\text{Exc}(\mu) = \cup_j F_j$  are all supported by a simple normal crossings divisor. Then  $a(\eta; X, B) = \min_{\mu(F_j)=\eta} a(F_j; X, B)$ . The log pullback formula

$$\mu^*(K_X + B) = K_{X'} + \mu_*^{-1}B + \sum_j (1 - a(F_j; X, B))F_j$$

shows that  $a(\eta; X, B) \in \frac{1}{r}\mathbb{Z}$  if  $r(K_X + B)$  is a Cartier divisor near  $\eta$ .

**Definition 2.4.** A log variety  $(X, B)$  has *log canonical singularities* at  $\eta$  if  $a(\eta; X, B) \geq 0$ . We say that  $(X, B)$  has *log canonical singularities* if  $a(X, B) \geq 0$ .

**Example 2.5.** Consider the log variety  $(X, \sum_i b_i E_i)$ , where  $X$  is a manifold,  $\sum_i E_i$  a simple normal crossings divisor and  $b_i \in [0, 1]$  for all  $i$ . Then  $a(X, \sum_i b_i E_i) = \min_i(1 - b_i)$ .

**Example 2.6.** Let  $X$  be a toric variety and  $X \setminus T = \cup_i E_i$  the complement of the torus. Then  $(X, \sum_i E_i)$  is a log variety with  $a(X, \sum_i E_i) = 0$  and  $K_X + \sum_i E_i = 0$ .

If  $a(X, B) \geq 0$ , the infimum in its definition is also a minimum. With the notation above,  $a(X, B) = \min(\min_i a(E_i; X, B), \min_j a(F_j; X, B))$ . Moreover, the formula

$$K_{X'} + \mu_*^{-1} B + \sum_j F_j = \mu^*(K_X + B) + \sum_j a(F_j; X, B) F_j$$

becomes a Zariski decomposition of the relative log manifold of general type  $(X', (\mu^{-1})_* B + \sum_j F_j) \rightarrow X$ . Its log canonical model in the sense of §1 is  $(X/X, B)$ , where  $X/X$  is the identity morphism.

**Definition 2.7.** Let  $(X/S, B)$  be a log variety  $(X, B)$  with  $a(X, B) \geq 0$ , endowed with a projective morphism  $\pi: X \rightarrow S$ . We say that  $(X/S, B)$  is a

- *log Fano* if  $-(K_X + B)$  is  $\pi$ -ample.
- *log Calabi-Yau* if  $K_X + B$  is  $\pi$ -numerically trivial.
- *log canonical model* if  $K_X + B$  is  $\pi$ -ample.

It is useful to observe that a log variety  $(X, B)$  has log canonical singularities at  $\eta$  and only if  $(X/X, B)$  is a log canonical model in a neighborhood of  $\eta$  (cf. Example 4.6). Note however that the three geometric types above coincide if  $\pi$  is the identity map.

**Definition 2.8.** A germ of log variety  $P \in (X, B)$  is called *flat* if  $a(P; X, B) = 0$ .

The typical example of flat germ is  $0 \in (\mathbb{C}^d, \sum_{i=1}^d H_i)$ , where  $H_i$  are the coordinate hyperplanes. Our terminology is inspired by an analogy between germs and projective manifolds, where if the minimal log discrepancy corresponds to the Kodaira dimension, flat germs correspond to manifolds with Kodaira dimension zero (an elliptic curve, for example). Also, note that  $0 \in (\mathbb{C}, 1 \cdot 0)$  is the only flat log germ in dimension one.

**2-A. Examples of minimal log discrepancies.** Minimal log discrepancies can be easily computed for log varieties  $(X, B)$  such that  $X$  is a toric variety and  $B$  is supported by the complement of the torus (see [47] for standard terminology on toric varieties). We only consider here  $\mathbb{Q}$ -factorial, log canonical toric germs of log varieties

$$P \in (X, B) = (T_N \text{emb}(\sigma), \sum_{i=1}^d b_i H_i).$$

They are in one-to-one correspondence with the following data:

- $\sigma = \{x \in \mathbb{R}^d; x_1, \dots, x_d \geq 0\}$ .
- $N \subset \mathbb{R}^d$  is a lattice, containing  $(1, 0, \dots, 0), \dots, (0, \dots, 0, 1)$  as primitive vectors.
- $(b_1, \dots, b_d) \in [0, 1]^d$ .

The following basic facts provide lots of examples of minimal log discrepancies:

- (a)  $a(\eta_{H_i}; X, B) = 1 - b_i$ .
- (b) Let  $x \in N^{\text{prim}} \cap \sigma$  be a primitive vector. Then  $x$  defines a barycentric subdivision  $\Delta_x$  of  $\sigma$ , and the exceptional locus of the induced birational map  $T_N \text{emb}(\Delta_x) \rightarrow T_N \text{emb}(\sigma)$  is a prime divisor  $E_x$ . Then  $a(E_x; X, B) = \sum_{i=1}^d (1 - b_i) x_i$ .
- (c) Log resolutions exists in the toric category. Therefore minimal log discrepancies can be computed using only valuations  $E_x$  as in (b).

- (d) The point  $P$  is the unique fixed point of the torus action. Its minimal log discrepancy is computed as follows

$$a(P; X, B) = \min\left\{\sum_{i=1}^d (1 - b_i)x_i; x \in N \cap \text{int}(\sigma)\right\}.$$

- (e) Let  $P \in C \subset X$  be the toric cycle corresponding to a face  $\tau \prec \sigma$ . The minimal log discrepancy at its generic point is

$$a(\eta_C; X, B) = \min\left\{\sum_{i=1}^d (1 - b_i)x_i; x \in N \cap \text{relint}(\tau)\right\}.$$

- (f) The global minimal log discrepancy is computed as follows

$$a(X, B) = \min\left\{\sum_{i=1}^d (1 - b_i)x_i; x \in N \cap \sigma \setminus 0\right\}.$$

- (g) In all minimums above, it suffices to consider only the finitely many lattice points  $x \in N \cap [0, 1]^d$ .

**Example 2.9.** Suppose  $N = \mathbb{Z}^d$ , that is  $X = \mathbb{C}^d$  and the  $H_i$ 's are the coordinate hyperplanes. For the cycle  $C : (x_1 = \cdots = x_s = 0)$ , we have  $a(\eta_C; X, B) = s - b_1 - \cdots - b_s$ .

**Example 2.10.** Suppose  $B = 0$ . Since  $\sigma$  is fixed, only the lattice  $N$  varies.

- (i) Take  $N = \mathbb{Z}^2 + \mathbb{Z}(\frac{1}{q}, \frac{q-1}{q})$ , for some integer  $q \geq 2$ . The surface germ  $P \in X$  is a  $A_{q-1}$ -singularity. We compute  $N \cap (0, 1]^2 = \{(\frac{k}{q}, \frac{q-k}{q}); 1 \leq k \leq q-1\} \cup \{(1, 1)\}$ , so  $a(P; X) = 1$ .
- (ii) Take  $N = \mathbb{Z}^2 + \mathbb{Z}(\frac{1}{k}, \frac{1}{k})$ , for some positive integer  $k$ . As above, we compute  $a(P; X) = \frac{2}{k}$ .
- (iii) Take  $N = \mathbb{Z}^3 + \mathbb{Z}\frac{1}{q}(1, p, q-p)$ , where  $p, q$  are integers with  $1 \leq p \leq q-1$ ,  $\gcd(p, q) = 1$ . Then  $P \in X$  is a terminal 3-fold singularity, with  $a(P; X) = 1 + \frac{1}{q}$ .
- (iv) Take  $N = \mathbb{Z}^3 + \mathbb{Z}\frac{1}{2q}(1, q, 1+q)$ , with  $q \geq 1$ . Then  $P \in X$  is a 3-fold singularity with  $a(P; X) = 1 + \frac{1}{q}$ . This germ has the minimal log discrepancy of a terminal singularity, but it's not terminal, since it is not an isolated singularity. The singular locus of  $X$  is  $C_2 : (x_1 = x_3 = 0)$ , and  $a(\eta_{C_2}; X) = \frac{2}{q}$ .

**Example 2.11.** For each point  $x \in \mathbb{Q}^d \cap (0, 1]^d$ , we construct a  $d$ -dimensional germ of toric log variety  $P_x \in (X_x, B_x)$ , with minimal log discrepancy

$$a(P_x; X_x, B_x) = \min_{n \geq 0} \sum_{i=1}^d (1 + nx_i - \lceil nx_i \rceil).$$

Choose a positive integer  $q$  such that  $qx \in \mathbb{Z}^d$ , and define integers

$$n_j = \frac{\gcd(q, qx_1, \dots, q\widehat{x_j}, \dots, qx_d)}{\gcd(q, qx_1, \dots, qx_d)} \quad (1 \leq j \leq d).$$

These integers are independent of the choice of  $q$ . Let  $\sigma \subset \mathbb{R}^d$  be the standard positive cone, and  $P_x \in X_x := T_{\mathbb{Z}^d + \mathbb{Z} \cdot x} \text{emb}(\sigma)$  the unique point fixed by the torus. The primitive

vectors of the lattice  $\mathbb{Z}^d + \mathbb{Z} \cdot x$  along the rays of  $\sigma$  are

$$\frac{1}{n_1}(1, 0, \dots, 0), \dots, \frac{1}{n_d}(0, \dots, 0, 1).$$

They define invariant prime divisors  $H_1, \dots, H_d$  in  $X$ . Set  $B_x = \sum_{i=1}^d (1 - \frac{1}{n_i})H_i$ .

It is natural to “compactify”  $\mathbb{Q}^d \cap (0, 1]^d$  to  $\mathbb{Q}^d \cap [0, 1]^d \setminus 0$ , in order to study the limiting behaviour of  $a(P_x; X_x, B_x)$  [9, 5].

The minimal log discrepancy of a toric germ is the “local” version of Minkowski’s first minimum of a convex body about the origin (see [13] for the definition and basic properties of the latter invariant). In our setting, if we denote by  $\Delta$  the convex set  $\{x \in \mathbb{R}^d; x_1, \dots, x_d \geq 0, \sum_{i=1}^d (1 - b_i)x_i \leq 1\}$ , then

$$a(P; X, B) = \sup\{t \geq 0; N \cap \text{int}(t\Delta) = \emptyset\}.$$

See [11] for more on minimal log discrepancies versus lattice-point-free convex bodies.

### 3. PROBLEMS ON MINIMAL LOG DISCREPANCIES

Minimal log discrepancies originate in the problem of termination of log flips: starting with a given log variety, can we perform log flips infinitely many times? Log flips are surgery operations which preserve codimension one cycles, and improve the singularities of higher codimensional cycles. As a measure of this improvement, log discrepancies may only increase after a log flip, and some of them increase strictly [51]. This is the heuristic behind the termination of a sequence of log flips, and it lead Shokurov [51, 52, 55] to question the existence of an infinite increasing sequence of minimal log discrepancies.

First, we fix a log variety  $(X, B)$ , and investigate the set of minimal log discrepancies of all prime cycles of  $X$  [4]. The formula  $a(\eta_C; X, B) = a(P; X, B) - \dim(C)$ , for a general closed point  $P$  of a given prime cycle  $C \subset X$ , shows that closed points contain the essential information. Consider now the minimal log discrepancy  $a(P; X, B)$  as a function on the set of the closed points  $P \in X$ . This function has a finite image, and in particular the set of minimal log discrepancies of all prime cycles of  $X$  is *finite*. Moreover, the level sets  $\{P \in X; a(P; X, B) \leq t\}$  ( $t \geq 0$ ) are constructible. Simple examples, such as a Du Val singularity  $P \in X$ , with  $a(x; X) = 2$  for  $x \neq P$ , and  $a(P; X) = 1$ , suggest that these level sets are in fact closed.

**Conjecture 3.1** ([3]). *The minimal log discrepancy  $a(P; X, B)$  is lower semi-continuous as a function on the closed points  $P$  of  $X$ .*

This behaviour is confirmed in several cases: a)  $\dim(X) \leq 3$  [3, 4]; b)  $(X, B)$  is a toric log variety [4]; c)  $X$  is a local complete intersection [19, 18]. Also, it is equivalent to the inequality  $a(P; X, B) \leq a(\eta_C; X, B) + 1$ , for every closed point on a curve in  $X$  [4].

Now consider the general case, when log flips change the log variety  $(X, B)$  in codimension at least two. The coefficients of the boundary are preserved, so we may assume that they belong to a given finite set. More generally, let  $\mathcal{B} \subset [0, 1]$  be a set satisfying the *descending chain condition* ( $\mathcal{B} = \{1 - \frac{1}{n}; n \geq 1\} \cup \{1\}$  is the typical example), and define

$$\text{Mld}(d, \mathcal{B}) = \{a(P; X, B); \dim(X) = d, \text{ coefficients of } B \text{ belong to } \mathcal{B}\}.$$

The set  $\text{Mld}(1, \mathcal{B}) = \{1 - b; b \in \mathcal{B}\}$  clearly satisfies the ascending chain condition.

**Conjecture 3.2** (Shokurov [52, 55]). *The following properties hold:*

- (1)  $\text{Mld}(d, \mathcal{B})$  satisfies the ascending chain condition.
- (2)  $a(P; X, B) \leq \dim(X)$ . Moreover, if  $a(P; X, B) > \dim(X) - 1$ , then  $P \in X$  is a nonsingular point and  $a(P; X, B) = \dim(X) - \text{mult}_P(B)$ .
- (3) Assume that  $\mathcal{B} \cap [0, 1 - \frac{1}{k}]$  is a finite set for every  $k \geq 1$ . Then the accumulation points of  $\text{Mld}(d, \mathcal{B})$  are included in  $\text{Mld}(d-1, \mathcal{B}')$ , for a suitable set  $\mathcal{B}'$ .

This conjecture was confirmed for surfaces [53, 1], and toric log varieties [9, 5]. By the classification of terminal 3-fold singularities,  $\text{Mld}(3, \{0\}) \cap (1, +\infty) = \{1 + \frac{1}{q}; q \geq 1\} \cup \{3\}$  [32, 41]. Also, (2) holds if  $X$  is a local complete intersection [19, 18]. Shokurov [57] reduced the global problem of the termination of flips to the two local problems 3.1 and 3.2.(1). See [7] for more on this.

An interesting problem posed by Shokurov is to relate the minimal log discrepancy and the Cartier index of a singularity. Suppose  $P \in X$  is the germ of a  $d$ -fold with  $a(P; X) \geq 0$ . If  $nK_X \sim 0$  and Conjecture 3.2.(2) holds, then the minimal log discrepancy can take at most finitely many values:  $a(P; X) \in \{0, \frac{1}{n}, \dots, \frac{nd}{n}\}$ . Conversely, is there an integer  $n$ , depending only on  $d$  and  $a(P; X)$ , such that  $nK_X \sim 0$ ? The answer is positive if  $d = 2$  (Shokurov, unpublished). An important special case of this problem is  $a(P; X) = 0$ . If  $d = 2$ , then  $n \in \{1, 2, 3, 4, 6\}$  [54]. If  $d = 3$ , then  $\varphi(n) \leq 20$  and  $n \neq 60$ , where  $\varphi$  is the Euler number [26]. See also [21] for a higher dimensional reduction to a global problem on log Calabi-Yau varieties in one dimension less. The boldest conjecture here is the following.

**Conjecture 3.3** (Shokurov [56]). *Let  $P \in (X, B)$  be a log germ such that  $a(P; X, B) = 0$  and  $B$  has coefficients in a set  $\mathcal{B} \subset [0, 1] \cap \mathbb{Q}$  satisfying the descending chain condition. Then  $n(K + B) \sim 0$  for some positive integer  $n$ , depending only on  $\dim(X)$  and  $\mathcal{B}$ .*

A useful formula in inductive arguments in the log category is a comparison of minimal log discrepancies under adjunction, called precise inversion of adjunction.

**Conjecture 3.4** (Shokurov [54], Kollár [33]). *Let  $P \in S \subset (X, B)$  be the germ of a log variety and a normal prime divisor  $S$  with  $\text{mult}_S(B) = 1$ . By adjunction, we have  $(K_X + B)|_S = K_S + B_S$ . Then  $a(P; X, B) = a(P; S, B_S)$ .*

It follows from the Log Minimal Model Program if  $a(P; X, B) \leq 1$  [33], and it holds if  $X$  is a local complete intersection [19, 18].

Minimal log discrepancies appear naturally in global contexts, such as Fujita's Conjecture on adjoint linear systems or the boundedness of log Fano varieties.

**Conjecture 3.5** (A. and L. Borisov [8]-Alexeev [2]). *Let  $\epsilon \in (0, 1]$  and  $d \in \mathbb{Z}_{\geq 1}$ . Then log varieties  $X$  with  $-K_X$  ample,  $a(X) \geq \epsilon$  and  $\dim(X) = d$ , form a bounded family.*

This is known in several cases: a)  $X$  is toric [8]; b)  $X$  nonsingular [34]; c)  $d = 2$  [2]; d)  $d = 3$ ,  $\epsilon = 1$  [31, 38]; e)  $d = 3$ , and the index of  $K_X$  is fixed [12].

**3-A. Toric case.** In the assumptions and notations of § 2-A, we illustrate some of the local problems on minimal log discrepancies. For *lower semi-continuity*, it is enough to see that  $a(P; X, B) \leq a(\eta_C; X, B) + 1$  for a torus-invariant curve  $P \in C$ . Suppose  $C$  corresponds to the face  $\tau = \sigma \cap (x_d = 0)$ . There exists  $(x', 0) \in N^{\text{prim}} \cap \text{relint}(\sigma)$  such that  $a(\eta_C; X, B) = \sum_{i=1}^{d-1} (1 - b_i)x_i$ . Then  $(x', 1) \in N \cap \text{int}(\sigma)$  and there exists  $x \in N^{\text{prim}}$

and a positive integer  $l$  with  $lx = (x', 1)$ . We have

$$a(E_x; X, B) \leq la(E_x; X, B) = \sum_{i=1}^{d-1} (1 - b_i)x'_i + 1 - b_d \leq a(\eta_C; X, B) + 1.$$

Therefore  $a(P; X, B) \leq a(\eta_C; X, B) + 1$ .

For *precise inversion of adjunction*, suppose  $B = \sum_{i=1}^{d-1} b_i H_i + H_d$ . Then  $S = H_d$  is the toric variety  $T_{N_d} \text{emb}(\sigma_d)$ , where  $\sigma_d = \{x \in \mathbb{R}^{d-1}; x_1, \dots, x_{d-1} \geq 0\}$  and  $N_d = \{x \in \mathbb{R}^{d-1}; \exists t \in \mathbb{R}, (x, t) \in N\}$ . To bring this to the normal form in § 2-A, note that there are positive integers  $n_1, \dots, n_{d-1}$  such that  $\frac{1}{n_i}(0, \dots, \overset{i}{1}, \dots, 0)$  are primitive vectors of  $N_d^{\text{prim}}$ . Then  $S = T_{N'} \text{emb}(\sigma')$ , where  $N' = \{x' \in \mathbb{R}^{d-1}; (n_1 x'_1, \dots, n_{d-1} x'_{d-1}) \in N_d\}$  and  $\sigma'$  is the usual positive cone. Let  $H'_1, \dots, H'_{d-1}$  the torus invariant prime divisors of  $S$ . The key observation is that the log canonical divisor  $K_X + B = \sum_{i=1}^{d-1} -(1 - b_i)H_i$  is independent of  $H_d$ . The boundary induced on  $S$  by adjunction is  $B_S = \sum_{i=1}^{d-1} (1 - \frac{1-b_i}{n_i})H'_i$ , and the equality  $a(P; X, B) = a(P; S, B_S)$  is clear.

Finally, for the *ascending chain condition*, assume by contradiction that we have a strictly increasing sequence  $a^1 < a^2 < a^3 < \dots$ , where  $a^n = a(P^n; T_{N^n} \text{emb}(\sigma))$  for  $n \geq 1$ . For simplicity, assume that the boundary is zero, so only the lattice changes. We may find  $x^n \in (0, 1]^d \cap N^n$  such that  $a^n = \sum_{i=1}^d x_i^n$ . In particular,  $a^n \leq d$  for all  $n$ . Consider now the strictly increasing sequence of open sets  $U^n = \{x \in (0, +\infty)^d; \sum_{i=1}^d x_i < a^n\}$ . By [40],  $G^n = \{x \in \mathbb{R}^d; U^n \cap (\mathbb{Z}^d + \mathbb{Z}x) = \emptyset\}$  is the union of finitely many closed subgroups containing  $\mathbb{Z}^d$  (the Flatness Theorem of Khinchin [28] gives an alternative proof). We have  $G^n \supseteq G^{n+1}$  since  $a^n < a^{n+1}$  and  $x^n \in G^n \setminus G^{n+1}$ , so we obtain a strictly decreasing sequence of finite unions of closed subgroups containing  $\mathbb{Z}^d$ . This is impossible, since the latter set satisfies the descending chain condition.

**3-B. Methods.** The toric case (see [43, 10]) suggests that behind the ascending chain condition of minimal log discrepancies lies a deeper fact, the boundedness of singularities with minimal log discrepancy bounded away from zero. Some log canonical singularities are classified in low dimension, but in general we could only expect general structure theorems and boundedness results in terms of minimal log discrepancies. For example, Du Val singularities are classified as follows:  $A_n, D_n, E_6, E_7, E_8$ . From the above point of view, Du Val singularities are nothing but surface singularities having minimal log discrepancy at least 1, and they come in two types: a 1-dimensional series with two components ( $A$  and  $D$ ), and a 0-dimensional series ( $E$ ).

The known method for bounding germs  $P \in X$  is to study the singularities at  $P$  of the linear systems  $| -mK_X |$  ( $m \geq 1$ ), and reduce this local problem to the global problem of bounding log Fano or log Calabi-Yau varieties in one dimension less [56, 48]. Given that §1 suggests that minimal log discrepancies are invariants of objects of general type, it also natural to investigate the singularities at  $P$  of the linear systems  $|mK_X|$  ( $m \geq 1$ ) and  $|\mathfrak{m}_{P,X}^m|$  ( $m \geq 1$ ), and relate germs with log canonical models in one dimension less.

Also, it is possible that minimal log discrepancies can be understood from several points of view: analytic, birational, motivic or p-adic. The motivic interpretation of minimal log discrepancies is known in the case when the canonical divisor is  $\mathbb{Q}$ -Cartier [44, 60]. As for the analytic side, the description of log discrepancies as the coefficients of a Zariski decomposition suggests an interpretation of minimal log discrepancies in terms of Lelong

numbers. For example, the upper bound of Conjecture 3.2.(2) is essentially equivalent to the following problem: let  $(X, B)$  be a log manifold of general type having a Zariski decomposition  $K_X + B = P + F$ , such that  $F$  is supported by the components of  $B$  with coefficient one. Then some coefficient of  $F$  is at most  $\dim(X)$ .

#### 4. THE LOG CANONICAL THRESHOLD

**Definition 4.1.** Let  $(X, B)$  be a log variety,  $\eta \in X$  a Grothendieck point such that  $a(\eta; X, B) \geq 0$ , and  $D$  an effective  $\mathbb{R}$ -Cartier  $\mathbb{R}$ -divisor passing through  $\eta$ . The *log canonical threshold* of  $D$  with respect to  $\eta \in (X, B)$  is defined as

$$\text{lct}(\eta \in (X, B); D) = \sup\{t \geq 0; a(\eta; X, B + tD) \geq 0\}.$$

The *global log canonical threshold* is  $\text{lct}(X, B; D) = \min_{\eta \in X} \text{lct}(\eta \in (X, B); D)$ .

Note that  $a(\eta; X, B) \geq 0$  if and only if  $(X, B)$  has log canonical singularities at  $\eta$ . Since the support of  $B + tD$  is a fixed divisor  $S$ , we may compute the log canonical threshold on a log resolution of  $X$  and  $S$ . In particular,  $\text{lct}(\eta \in (X, B); D) \in \mathbb{Q}$  if  $B$  and  $D$  are rational. The inequality  $\text{lct}(P \in (X, B); D) \leq \text{lct}(\eta_C \in (X, B); D)$  holds for a closed point of a prime cycle  $P \in C \subset X$ , with equality if  $P$  is general in  $C$ .

**Example 4.2.** Let  $\dim(X) = 1$  and  $P \in X$ . Then  $\text{lct}(P \in (X, B); D) = \frac{a(P; X, B)}{\text{mult}_P(D)}$ .

**Example 4.3** ([16, 23]). Let  $0 \in H: (f = 0) \subset \mathbb{C}^d$  be a hypersurface. Then

$$\text{lct}(0 \in \mathbb{C}^d; H) = \sup\{t \geq 0; |f|^{-t} \text{ is } L^2 \text{ near } 0\}.$$

The reciprocal number  $\mu(0; H) = 1/\text{lct}(0 \in \mathbb{C}^d; H)$ , called Arnold multiplicity, satisfies the inequalities  $\mu(0; H) \leq \text{mult}(0; H) \leq d \cdot \mu(0; H)$ .

**Example 4.4.** Let  $H_1, \dots, H_d$  be the coordinate hyperplanes in  $\mathbb{C}^d$ ,  $b_1, \dots, b_d \in [0, 1]$  and  $n_1, \dots, n_d \in \mathbb{Z}_{\geq 1}$ . Then

- (i)  $\text{lct}(0 \in (\mathbb{C}^d, \sum_{i=1}^d b_i H_i); (x_1^{n_1} \cdots x_d^{n_d})) = \min_{i=1}^d \frac{1-b_i}{n_i}$ .
- (ii)  $\text{lct}(0 \in (\mathbb{C}^d, \sum_{i=1}^d b_i H_i); (x_1^{n_1} + \cdots + x_d^{n_d})) = \min(1, \sum_{i=1}^d \frac{1-b_i}{n_i})$ .

**Example 4.5.** Log canonical thresholds of non-degenerate hypersurfaces in toric varieties have a combinatorial description. Let  $P \in (X, B) = (T_N \text{ emb}(\sigma), \sum_{i=1}^d b_i H_i)$  be a toric germ as in §2-A, and

$$f = \sum_{\alpha} \lambda_{\alpha} \chi^{m_{\alpha}} \in \mathfrak{m}_{P, X} \setminus 0.$$

Here  $m_{\alpha}$  are finitely many points in  $M \cap \sigma^{\vee}$ , where  $M$  is the lattice dual to  $N$  and  $\sigma \subset M_{\mathbb{R}}$  is the cone dual to  $\sigma$ . Note that  $(1 - b_1, \dots, 1 - b_d) \in \sigma^{\vee}$ . Let  $\square$  be the convex hull of the  $\cup_{\alpha} (m_{\alpha} + \sigma^{\vee})$ . The ray  $\mathbb{R}_{\geq 0}(1 - b_1, \dots, 1 - b_d)$  intersects  $\square$  for the first time in a point  $\mu(1 - b_1, \dots, 1 - b_d)$ . We have the following inequality

$$\text{lct}(P \in (X, B); (f = 0)) \leq \min(1, \frac{1}{\mu}),$$

and equality holds if the coefficients  $\lambda_{\alpha}$  are sufficiently general. The generality condition can be made explicit, as in [6].

**Example 4.6.** The log canonical threshold  $\text{lct}(P \in (X, B); D)$  is the largest  $t \geq 0$  such that  $(X/X, B + tD)$  is a relative log canonical model near  $P$ . For example, consider the germ of the cuspidal curve  $0 \in C: (x^2 - y^3 = 0) \subset \mathbb{C}^2$ , with log canonical threshold  $\frac{5}{6}$ . The cusp can be resolved by a composition of three blow-ups  $\mu: X \rightarrow \mathbb{C}^2$ . Let  $E_1, E_2, E_3$  be the proper transforms on  $X$  of the exceptional divisors, in the order of their appearance. We obtain a singular model  $X \rightarrow Y \rightarrow \mathbb{C}^2$  by contracting  $E_1$  and  $E_2$ , and let  $C'$  and  $E'_3$  be the proper transforms of  $C$  and  $E_3$  on  $Y$ . The relative log variety  $(X, E_1 + E_2 + E_3 + t\mu_*^{-1}C) \rightarrow \mathbb{C}^2$  is of general type for  $t \in [0, 1] \cap \mathbb{Q}$ , and its log canonical model is  $(\mathbb{C}^2, tC) \rightarrow \mathbb{C}^2$  for  $t \leq \frac{5}{6}$ , and  $(Y, E'_3 + tC') \rightarrow \mathbb{C}^2$  for  $t > \frac{5}{6}$ . On the other hand,  $(\mathbb{C}^2, tC) \rightarrow \mathbb{C}^2$  is a log canonical model on  $\mathbb{C}^2 \setminus 0$ , for every  $t \in [0, 1] \cap \mathbb{Q}$ .

**4-A. Problems on log canonical thresholds.** One can classify a singularity by constructing a *flat log structure* on it, in an effective way. Log canonical thresholds appear naturally in this construction, as the coefficients of the boundary of the flat log structure.

For example, consider a log germ  $P \in (X, B)$  with  $a(P; X, B) \geq 0$ . If  $a(P; X, B) = 0$ , the log germ is flat. Otherwise, for a general hypersurface  $D_1 \in |\mathfrak{m}_{P, X}|$ , the log canonical threshold  $\gamma_1 = \text{lct}(P \in (X, B); D_1)$  is defined in such a way so that  $a(P; X, B + \gamma D_1) \geq 0$ , and there exists a minimal cycle  $C \ni P$  such that  $a(\eta_C; X, B + \gamma D) = 0$ . If  $C = \{P\}$ , then  $P \in (X, B + \gamma D)$  is flat. Otherwise, we repeat this process (at most  $\dim(C)$  times) until we obtain a flat log structure  $0 \in (X, B + \sum_i \gamma_i D_i)$ . Flat germs with certain restrictions on their boundaries are expected to be bounded in a certain sense (cf. §3), so for effective results we should also control the  $\gamma_i$ 's. Shokurov observed that these coefficients satisfy the ascending chain condition in dimension two, and used this to construct 3-fold log flips.

**Conjecture 4.7** (Shokurov [54], Kollár [37]). *For a positive integer  $d$  and a set  $0 \in \mathcal{B} \subset [0, 1]$  satisfying the descending chain condition, define*

$$\text{Lct}(d, \mathcal{B}) = \{\text{lct}(P \in (X, B); D); \dim(X) = d, \text{coefficients of } B \text{ belong to } \mathcal{B}, D \in |\mathfrak{m}_{P, X}|\}.$$

*The following properties hold:*

- (1) *The set  $\text{Lct}(d, \mathcal{B})$  satisfies the ascending chain condition.*
- (2) *Assume that  $\mathcal{B} \cap [1, 1 - \frac{1}{k}]$  is a finite set for every  $k \geq 1$ . Then the set of accumulation points of  $\text{Lct}(d, \mathcal{B})$  is  $\text{Lct}(d - 1, \mathcal{B}') \setminus \{1\}$ , for a suitable set  $\mathcal{B}' \subset [0, 1]$ .*

This limiting behaviour is known in several cases: a)  $d = 2$  [54]; b)  $d = 3$  [2, 42]; c)  $P \in X$  is toric,  $B$  is invariant and  $D \in |\mathfrak{m}_{P, X}|$  is a general hypersurface [27]. The set  $\text{Lct}_d := \text{Lct}(d, \{0\})$  has an explicit description in some cases: a)  $\text{Lct}_1 = \{\frac{1}{n}; n \in \mathbb{Z}_{\geq 1}\}$ ; b)  $\text{Lct}_2 \cap [\frac{1}{2}, 1] = \{\frac{1}{2} + \frac{1}{n}; n \geq 3\} \cup \{1\}$  [39]; c)  $\text{Lct}_3 \cap [\frac{41}{42}, 1] = \{\frac{41}{42}, 1\}$  [35]. Conjecture 4.7 would imply that the number  $1 - \epsilon_d = \max \text{Lct}_d \cap (0, 1)$  is well defined, and Kollár [35] suggests that  $\epsilon_d$  is the minimal degree of a log canonical model  $(Y, B_Y)$  of dimension  $d - 1$ , with  $B_Y$  having coefficients in  $\{1 - \frac{1}{n}; n \geq 1\} \cup \{1\}$ . For example,  $\epsilon_1 = \frac{1}{2}, \epsilon_2 = \frac{1}{6}, \epsilon_3 = \frac{1}{42}$ . For the relationship between Conjectures 3.2, 3.5 and 4.7, see [7].

Flat log structures have global applications as well. Given a closed point  $P$  on a polarized manifold  $(X, H)$ , does there exist  $n \geq 1$  and  $D \in |nH|$  with  $a(P; X, \frac{1}{n}D) = 0$ ? If such a flat structure exists, then  $P$  is not in the base locus of any divisor  $L$  such that  $L - K_X - H$  ample. This is a powerful technique to study adjoint line bundles, parallel to the  $L^2$ -methods for singular hermitian metrics in complex geometry (see [30, 17, 37, 3] and [58, 16] for the algebraic and analytic side of the story, respectively). As above, log

canonical thresholds appear naturally. For a log variety  $(X, B)$ , an ample  $\mathbb{Q}$ -divisor  $H$  and a Grothendieck point  $\eta \in X$ , define

$$\gamma(\eta \in (X, B); H) = \inf\{n \operatorname{lct}(\eta \in (X, B); D); D \in |nH|, n \geq 1\}$$

and  $\gamma(X, B; H) = \inf_{\eta \in X} \gamma(\eta \in (X, B); H)$  (cf. [59]).

**Example 4.8.** Suppose  $\dim(X) = 1$ . For every closed point  $P \in X$  we have

$$\gamma(\eta \in (X, B); H) = \frac{a(P; X, B)}{\deg(H)} \leq \frac{1}{\deg(H)},$$

and  $\gamma(X, B; H) = \frac{a(X, B)}{\deg(H)}$ .

**Lemma 4.9** ([51, 37]).  $\gamma(P \in (X, B); H) \sqrt[d]{H^d} \leq d$  for every closed point  $P \in X^d$ .

For Fujita's Conjecture on adjoint linear systems,  $\gamma(P \in (X, B); H)$  has to be small. This is achieved by Lemma 4.9, if  $\sqrt[d]{H^d}$  is large. On the other hand,  $\gamma(P \in (X, B); H)$  cannot be too small in a bounded context, and this can be used to bound  $\sqrt[d]{H^d}$  from above. Some known cases are: a)  $\gamma(\eta_X \in X; -K_X) \geq \frac{1}{d+1}$  if  $X$  is a Fano manifold of dimension  $d$  [23]; b)  $\gamma(G; H) \geq 1$  if  $\mathcal{O}_G(H)$  is the Plücker line bundle on the Grassmanian [24]; c)  $\gamma(A; \Theta) \geq 1$  for a principally polarized abelian variety [36]. Pukhlikov [49] suggests that a Fano variety  $X$  with large  $\gamma(X; -K_X)$  is birationally rigid. Hwang [23] suggests that for a log canonical model  $X$ ,  $\gamma(\eta_X \in X; K_X) \sqrt[d]{K_X^d}$  is also bounded away from zero, only in terms of dimension. The level sets and critical points of the functional  $\gamma(\eta \in (X, B); H)$ , and an extension of Faltings' product theorem [20, 45, 23], should play a key role in constructing flat log structures.

## REFERENCES

- [1] Alexeev, V., *Two two-dimensional terminations*. Duke Math. J. 69 (1993), no. 3, 527–545.
- [2] Alexeev, V., *Boundedness and  $K^2$  for log surfaces*. Internat. J. Math. 5 (1994), no. 6, 779–810.
- [3] Ambro, F., *The Adjunction Conjecture and its applications*, preprint math.AG/9903060.
- [4] Ambro, F., *On minimal log discrepancies*, Math. Res. Letters **6**, (1999) 573–580.
- [5] Ambro, F., *The set of toric minimal log discrepancies*. Cent. Eur. J. Math. 4 (2006), no. 3, 358–370.
- [6] Arnold, V. I.; Gusen-Zade, S. M.; Varchenko, A. N., *Singularities of differentiable maps. Vol. II. Monodromy and asymptotics of integrals. Monographs in Mathematics, 83. Birkhäuser Boston, Inc., Boston, MA, 1988*.
- [7] Birkar, C.; Shokurov, V., V., *Mld's vs thresholds and flips*. preprint math.AG/0609539.
- [8] Borisov, A. A.; Borisov, L. A., *Singular toric Fano three-folds*. (Russian) Mat. Sb. 183 (1992), no. 2, 134–141; translation in Russian Acad. Sci. Sb. Math. 75 (1993), no. 1, 277–283.
- [9] Borisov, A., *Minimal discrepancies of toric singularities*. Manuscripta Math. 92 (1997), no. 1, 33–45.
- [10] Borisov, A., *On classification of toric singularities*. Algebraic geometry, 9 J. Math. Sci. (New York) 94 (1999), no 1, 1111–1113.
- [11] Borisov, A., *Convex lattice polytopes and cones with few lattice points inside, from a birational geometry viewpoint*. preprint math.AG/0001109.
- [12] Borisov, A., *Boundedness of Fano threefolds with log-terminal singularities of given index*. J. Math. Sci. Univ. Tokyo 8 (2001), no. 2, 329–342.
- [13] Cassels, J. W. S., *An introduction to Diophantine approximation*. Cambridge Tracts in Mathematics and Mathematical Physics, No. 45. Cambridge University Press, New York, 1957.
- [14] Cornalba, M.; Harris, J., *Divisor classes associated to families of stable varieties, with applications to the moduli space of curves*. Ann. Sci. École Norm. Sup. (4) 21 (1988), no. 3, 455–475.

- [15] Cutkosky, S. D., *Zariski decomposition of divisors on algebraic varieties*. Duke Math. J. 53 (1986), no. 1, 149–156.
- [16] Demailly, J.-P., *Multiplier ideal sheaves and analytic methods in algebraic geometry*. School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000), 1–148, ICTP Lect. Notes, 6, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2001.
- [17] Ein, L., *Multiplier ideals, vanishing theorems and applications*. Algebraic geometry—Santa Cruz 1995, 203–219, Proc. Sympos. Pure Math., 62, Part 1, Amer. Math. Soc., Providence, RI, 1997.
- [18] Ein, L.; Mustață, M.; Yasuda, T., *Jet schemes, log discrepancies and inversion of adjunction*. Invent. Math. 153 (2003), no. 3, 519–535.
- [19] Ein, L.; Mustață, M., *Inversion of adjunction for local complete intersection varieties*. Amer. J. Math. 126 (2004), no. 6, 1355–1365.
- [20] Faltings, G., *Diophantine approximation on abelian varieties*. Ann. of Math. (2) 133 (1991), no. 3, 549–576.
- [21] Fujino, O., *The indices of log canonical singularities*. Amer. J. Math. 123 (2001), no. 2, 229–253.
- [22] Fujita, T., *Zariski decomposition and canonical rings of elliptic threefolds*. J. Math. Soc. Japan 38 (1986), no. 1, 19–37.
- [23] Hwang, J.-M., *Arnold multiplicity of divisors on rational homogeneous spaces*. Proceedings of Algebraic Geometry Symposium (Kinosaki 2005), A. Ishii (Ed.), pp. 78 – 81.
- [24] Hwang, J.-M., *Log canonical thresholds of divisors on Grassmannians*. Math. Ann. 334 (2006), no. 2, 413–418.
- [25] Itaka, S., *Birational geometry of algebraic varieties*. Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983), 727–732, PWN, Warsaw, 1984.
- [26] Ishii, S., *The quotients of log-canonical singularities by finite groups*. Singularities—Sapporo 1998, 135–161, Adv. Stud. Pure Math., 29, Kinokuniya, Tokyo, 2000.
- [27] Ishii, S., *Log canonical thresholds for pairs  $\mathbb{C}^n$  and non-degenerate hypersurfaces*. preprint 2002.
- [28] Kannan, R.; Lovász, L., *Covering minima and lattice-point-free convex bodies*. Ann. of Math. (2) 128 (1988), no. 3, 577–602.
- [29] Kawamata, Y., *The Zariski decomposition of log-canonical divisors*. Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), 425–433, Proc. Sympos. Pure Math., 46, Part 1, Amer. Math. Soc., Providence, RI, 1987.
- [30] Kawamata Y., Matsuda K., Matsuki K., *Introduction to the minimal model program*, Algebraic Geometry, Sendai, Advanced Studies in Pure Math. 10 (1987), 283–360.
- [31] Kawamata, Y., *Boundedness of  $\mathbb{Q}$ -Fano threefolds*. Proceedings of the International Conference on Algebra, Part 3 (Novosibirsk, 1989), 439–445, Contemp. Math., 131, Part 3, Amer. Math. Soc., Providence, RI, 1992.
- [32] Kawamata, Y., *The minimal discrepancy of a 3-fold terminal singularity*, An appendix to [54].
- [33] Kollár, J., *Adjunction and discrepancies*. In *Flips and abundance for algebraic threefolds* (Kollár, J. Ed.), Astérisque 211, 1992.
- [34] Kollár, J.; Miyaoka, Y.; Mori, S., *Rational connectedness and boundedness of Fano manifolds*. J. Differential Geom. 36 (1992), no. 3, 765–779.
- [35] Kollár, J., *Log surfaces of general type; some conjectures*. Classification of algebraic varieties (L’Aquila, 1992), 261–275, Contemp. Math., 162, Amer. Math. Soc., Providence, RI, 1994.
- [36] Kollár, J., *Shafarevich Maps and Automorphic Forms*, Princeton Univ. Press, 1995.
- [37] Kollár, János, *Singularities of pairs*. Algebraic geometry—Santa Cruz 1995, 221–287, Proc. Sympos. Pure Math., 62, Part 1, Amer. Math. Soc., Providence, RI, 1997.
- [38] Kollár, J.; Miyaoka, Y.; Mori, S.; Takagi, H., *Boundedness of canonical  $\mathbb{Q}$ -Fano 3-folds*. Proc. Japan Acad. Ser. A Math. Sci. 76 (2000), no. 5, 73–77.
- [39] Kuwata, T., *On log canonical thresholds of reducible plane curves*. Amer. J. Math. 121 (1999), no. 4, 701–721.
- [40] Lawrence, J., *Finite unions of closed subgroups of the  $n$ -dimensional torus*. Applied geometry and discrete mathematics, 433–441, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 4, Amer. Math. Soc., Providence, RI, 1991.
- [41] Markushevich, D., *Minimal discrepancy for a terminal  $cDV$  singularity is 1*, J. Math. Sci. Univ. Tokyo 3 (1996), no. 2, 445–456.

- [42] McKernan, J.; Prokhorov, Y., *Threefold thresholds*. Manuscripta Math. 114 (2004), no. 3, 281–304.
- [43] Mori, S.; Morrison, D. R.; Morrison, I., *On four-dimensional terminal quotient singularities*. Math. Comp. 51 (1998), no. 184, 769–786.
- [44] Mustață, M., *Singularities of pairs via jet schemes*. J. Amer. Math. Soc. 15 (2002), no. 3, 599–615.
- [45] Nadel, A. M., *The boundedness of degree of Fano varieties with Picard number one*. J. Amer. Math. Soc. 4 (1991), no. 4, 681–692.
- [46] Nakayama, N., *Zariski-decomposition and abundance*. MSJ Memoirs, 14. Mathematical Society of Japan, Tokyo, 2004.
- [47] Oda, T., *Convex bodies and algebraic geometry. An introduction to the theory of toric varieties*. Ergebnisse der Mathematik und ihrer Grenzgebiete (3) 15. Springer-Verlag, Berlin, 1988.
- [48] Prokhorov, Y. G., *Lectures on complements on log surfaces*. MSJ Memoirs, 10. Mathematical Society of Japan, Tokyo, 2001.
- [49] Pukhlikov, A. V., *Birational geometry of Fano direct products*. preprint math.AG/0405011.
- [50] Reid, M., *Canonical 3-folds*. Journées de Géométrie Algébrique d’Angers, Juillet 1979/Algebraic Geometry, Angers, 1979, pp. 273–310, Sijthoff & Noordhoff, Alphen aan den Rijn—Germantown, Md., 1980.
- [51] Shokurov, V. V., *A nonvanishing theorem*. Izv. Akad. Nauk SSSR Ser. Mat. 49 (1985), no. 3, 635–651.
- [52] Shokurov, V.V., *Problems about Fano varieties*, Birational Geometry of Algebraic Varieties, Open Problems - Katata 1988, 30–32.
- [53] Shokurov, V.V., *A.c.c. in codimension 2*, preprint 1991.
- [54] Shokurov, V. V., *Three-dimensional log perestroikas. (Russian) With an appendix in English by Yujiro Kawamata*. Izv. Ross. Akad. Nauk Ser. Mat. 56 (1992), no. 1, 105–203; translation in Russian Acad. Sci. Izv. Math. 40 (1993), no. 1, 95–202.
- [55] Shokurov, V. V., *3-fold log models*. Algebraic geometry, 4., J. Math. Sci. 81 (1996), no. 3, 2667–2699.
- [56] Shokurov, V. V., *Complements on surfaces. Algebraic geometry, 10*. J. Math. Sci. (New York) 102 (2000), no. 2, 3876–3932.
- [57] Shokurov, V. V., *Letters of a bi-rationalist. V. Minimal log discrepancies and termination of log flips. (Russian) Tr. Mat. Inst. Steklova 246 (2004), Algebr. Geom. Metody, Svyazi i Prilozh., 328–351; translation in Proc. Steklov Inst. Math. 2004, no. 3 (246), 315–336.*
- [58] Siu, Y.-T., *The Fujita conjecture and the extension theorem of Ohsawa-Takegoshi*. Geometric complex analysis (Hayama, 1995), 577–592, World Sci. Publ., River Edge, NJ, 1996.
- [59] Viehweg, E., *Quasi-projective moduli for polarized manifolds*. Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 30. Springer-Verlag, Berlin, 1995.
- [60] Yasuda, T., *Dimensions of jet schemes of log singularities*. Amer. J. Math. 125 (2003), no. 5, 1137–1145.
- [61] Zariski, O., *The theorem of Riemann-Roch for high multiples of an effective divisor on an algebraic surface*. Ann. of Math. (2) 76 (1962), 560–615.

RIMS, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN.

*E-mail address:* ambro@kurims.kyoto-u.ac.jp