A Congruence Theorem for Minimal Surfaces in S^5 with Constant Contact Angle

Rodrigo Ristow Montes and Jose A. Verderesi *

Departamento de Matemática, Universidade Federal da Paraíba,

BR- 58.051-900 João Pessoa, P.B., Brazil and
Departamento de Matemática Pura, Instituto de Matemática e Estatística, Universidade de São Paulo, Caixa Postal 66281,
BR-05315-970 São Paulo, S.P., Brazil

Abstract

We provide a congruence theorem for minimal surfaces in S^5 with constant contact angle using Gauss-Codazzi-Ricci equations. More precisely, we prove that Gauss-Codazzi-Ricci equations for minimal surfaces in S^5 with constant contact angle satisfy an equation for the Laplacian of the holomorphic angle. Also, we will give a characterization of flat minimal surfaces in S^5 with constant contact angle.

Keywords: contact angle, holomorphic angle, Clifford torus, congruence theorem.

2000 Math Subject Classification: 53C42 - 53D10 - 53D35.

^{*}ristow@mat.ufpb.br and javerd@ime.usp.br

1 Introduction

We will establish a condition in order to investigate minimal surfaces in S^5 with constant contact angle using Gauss-Codazzi-Ricci equations. We define α to be the angle given by $\cos \alpha = \langle ie_1, v \rangle$, where e_1 and v are defined in section 2. The holomorphic angle α is the analogue of the Kähler angle introduced by Chern and Wolfson in [3] and [7].

In [5] we introduced the notion of contact angle, that can be considered as a new geometric invariant useful to investigate the geometry of immersed surfaces in S^3 . Geometrically, the contact angle (β) is the complementary angle between the contact distribution and the tangent space of the surface. Also in [5], we deduced formulas for the Gaussian curvature and the Laplacian of an immersed minimal surface in S^3 , and we gave a characterization of the Clifford Torus as the only minimal surface in S^3 with constant contact angle.

Recently, in [6], we construct a family of minimal tori in S^5 with constant contact angle and constant holomorphic angle. These tori are parametrized by the following circle equation

$$a^{2} + \left(b - \frac{\cos\beta}{1 + \sin^{2}\beta}\right)^{2} = 2\frac{\sin^{4}\beta}{(1 + \sin^{2}\beta)^{2}},$$
(1)

where a and b are given in section 3 (equation (10)). In particular, when a = 0 in (1), we recover the examples found by Kenmotsu, in [4]. These examples are defined for $0 < \beta < \frac{\pi}{2}$. Also, when b = 0 in (1), we find a new family of minimal tori in S^5 , and these tori are defined for $\frac{\pi}{4} < \beta < \frac{\pi}{2}$. For $\beta = \frac{\pi}{2}$, we give an alternative proof of this classification of a Theorem from Blair in [2], and Yamaguchi, Kon and Miyahara in [8] for Legendrian minimal surfaces in S^5 with constant Gaussian curvature.

In this paper, we will establish a congruence theorem for minimal surfaces in S^5 with constant contact angle (β). Using Gauss-Codazzi-Ricci equations, we prove the following theorem:

Theorem 1. Consider S a riemannian surface, $\alpha : S \rightarrow]0, \frac{\pi}{2}[$ a function over S that verifies the following equation:

$$\Delta(\alpha) = \cot \alpha \csc^3(\beta) |\nabla \alpha|^2 + a^2 \cot \alpha \cot^4(\beta) - 2a \cot \alpha \csc \beta \cot^2 \beta \alpha_2 -2 \cos \alpha (\cot \beta - \tan \beta) \tan^2 \beta \alpha_1 + \sin \alpha \cos \alpha (5 - \cot^4 \beta - 3 \csc^2 \beta)$$
(2)

then there exist one and only one minimal immersion of S into S^5 with constant contact angle such that α is the holomorphic angle of this immersion, where a is given in section 2, and its determined as a function of α and β in section 5.

As a immediately consequence of this method, we have a classification of flat minimal surfaces in S^5

Corollary 1. Suppose that the contact angle (β) is constant and suppose that S is a flat minimal surface in S⁵ with constant principal curvature in the direction e_3 , i.e. a is constant, then the holomorphic angle (α) must be constant

In general we have

Remark 1. The Theorem 1 implies a more general classification in [6], in the sense that, in [6] we need the condition that both angles are constant.

2 Contact Angle for Immersed Surfaces in S^{2n+1}

Consider in \mathbb{C}^{n+1} the following objects:

- the Hermitian product: $(z, w) = \sum_{j=0}^{n} z^{j} \bar{w}^{j};$
- the inner product: $\langle z, w \rangle = Re(z, w);$
- the unit sphere: $S^{2n+1} = \{ z \in \mathbb{C}^{n+1} | (z, z) = 1 \};$
- the *Reeb* vector field in S^{2n+1} , given by: $\xi(z) = iz$;
- the contact distribution in S^{2n+1} , which is orthogonal to ξ :

$$\Delta_z = \left\{ v \in T_z S^{2n+1} | \langle \xi, v \rangle = 0 \right\}.$$

We observe that Δ is invariant by the complex structure of \mathbb{C}^{n+1} .

Let now S be an immersed orientable surface in S^{2n+1} .

Definition 1. The contact angle β is the complementary angle between the contact distribution Δ and the tangent space TS of the surface.

Let (e_1, e_2) be a local frame of TS, where $e_1 \in TS \cap \Delta$, where e_1 is the characteristic field and introduced by Bennequin in [1].

Then $\cos \beta = \langle \xi, e_2 \rangle$. Finally, let v be the unit vector in the direction of the orthogonal projection of e_2 on Δ , defined by the following relation

$$e_2 = \sin\beta v + \cos\beta\xi. \tag{3}$$

3 Equations for Gaussian curvature and Laplacian of a minimal surface in S^5

In this section, we deduce the equations for the Gaussian curvature and for the Laplacian of a minimal surface in S^5 in terms of the contact angle and the holomorphic angle. Consider the normal vector fields

$$e_{3} = i \csc \alpha e_{1} - \cot \alpha v$$

$$e_{4} = \cot \alpha e_{1} + i \csc \alpha v$$

$$e_{5} = \csc \beta \xi - \cot \beta e_{2}$$
(4)

where $\beta \neq 0, \pi$ and $\alpha \neq 0, \pi$. We will call $(e_j)_{1 \leq j \leq 5}$ an *adapted frame*.

Using (3) and (4), we get

$$v = \sin \beta e_2 - \cos \beta e_5, \quad iv = \sin \alpha e_4 - \cos \alpha e_1 \tag{5}$$
$$\xi = \cos \beta e_2 + \sin \beta e_5$$

It follows from (4) and (5) that

$$ie_1 = \cos\alpha \sin\beta e_2 + \sin\alpha e_3 - \cos\alpha \cos\beta e_5$$
(6)

$$ie_2 = -\cos\beta z - \cos\alpha \sin\beta e_1 + \sin\alpha \sin\beta e_4$$

Consider now the dual basis (θ^j) of (e_j) . The connection forms (θ^j_k) are given by

$$De_j = \theta_j^k e_k,$$

and the second fundamental form with respect to this frame are given by

$$II^{j} = \theta_{1}^{j}\theta^{1} + \theta_{2}^{j}\theta^{2}; \quad j = 3, ..., 5.$$

Using (6) and differentiating v and ξ on the surface S, we get

$$D\xi = -\cos\alpha\sin\beta\theta^2 e_1 + \cos\alpha\sin\beta\theta^1 e_2 + \sin\alpha\theta^1 e_3 + \sin\alpha\sin\beta\theta^2 e_4 -\cos\alpha\cos\beta\theta^1 e_5,$$

$$Dv = (\sin\beta\theta_2^1 - \cos\beta\theta_5^1)e_1 + \cos\beta(d\beta - \theta_5^2)e_2 + (\sin\beta\theta_2^3 - \cos\beta\theta_5^3)e_3 + (\sin\beta\theta_4^2 - \cos\beta\theta_5^4)e_4 + \sin\beta(d\beta + \theta_5^5)e_5.$$

$$(7)$$

Differentiating e_3 , e_4 and e_5 , we have

$$\begin{aligned}
\theta_3^1 &= -\theta_1^3 \\
\theta_3^2 &= \sin \beta (d\alpha + \theta_4^1) - \cos \beta \sin \alpha \theta^1 \\
\theta_3^4 &= \csc \beta \theta_1^2 - \cot \alpha (\theta_1^3 + \csc \beta \theta_2^4) \\
\theta_3^5 &= \cot \beta \theta_2^3 - \csc \beta \sin \alpha \theta^1 \\
\theta_4^1 &= -d\alpha - \csc \beta \theta_2^3 + \sin \alpha \cot \beta \theta^1 \\
\theta_4^2 &= -\theta_2^4 \\
\theta_4^3 &= \csc \beta \theta_2^1 + \cot \alpha (\theta_1^3 + \csc \beta \theta_2^4) \\
\theta_5^4 &= \cot \beta \theta_2^4 - \sin \alpha \theta^2 \\
\theta_5^1 &= -\cos \alpha \theta^2 - \cot \beta \theta_2^1 \\
\theta_5^2 &= d\beta + \cos \alpha \theta^1 \\
\theta_5^3 &= -\cot \beta \theta_2^3 + \csc \beta \sin \alpha \theta^1 \\
\theta_5^4 &= -\cot \beta \theta_2^4 + \sin \alpha \theta^2
\end{aligned}$$
(8)

The conditions of minimality and of symmetry are equivalent to the following equations:

$$\theta_1^{\lambda} \wedge \theta^1 + \theta_2^{\lambda} \wedge \theta^2 = 0 = \theta_1^{\lambda} \wedge \theta^2 - \theta_2^{\lambda} \wedge \theta^1.$$
(9)

On the surface S, we consider

$$\theta_1^3 = a\theta^1 + b\theta^2$$

It follows from (9) that

$$\begin{aligned}
\theta_1^3 &= a\theta^1 + b\theta^2 \\
\theta_2^3 &= b\theta^1 - a\theta^2 \\
\theta_1^4 &= d\alpha + (b\csc\beta - \sin\alpha\cot\beta)\theta^1 - a\csc\beta\theta^2 \\
\theta_2^4 &= d\alpha \circ J - a\csc\beta\theta^1 - (b\csc\beta - \sin\alpha\cot\beta)\theta^2 \\
\theta_1^5 &= d\beta \circ J - \cos\alpha\theta^2 \\
\theta_2^5 &= -d\beta - \cos\alpha\theta^1
\end{aligned}$$
(10)

where J is the complex structure of S is given by $Je_1 = e_2$ and $Je_2 = -e_1$. Moreover, the normal connection forms are given by:

$$\begin{aligned}
\theta_3^4 &= -\sec\beta d\beta \circ J - \cot\alpha \csc\beta d\alpha \circ J + a\cot\alpha \cot^2\beta \theta^1 \\
&+ (b\cot\alpha \cot^2\beta - \cos\alpha \cot\beta \csc\beta + 2\sec\beta \cos\alpha)\theta^2 \\
\theta_3^5 &= (b\cot\beta - \csc\beta \sin\alpha)\theta^1 - a\cot\beta\theta^2 \\
\theta_4^5 &= \cot\beta (d\alpha \circ J) - a\cot\beta \csc\beta \theta^1 + (-b\csc\beta \cot\beta + \sin\alpha (\cot^2\beta - 1))\theta^2,
\end{aligned}$$
(11)

while the Gauss equation is equivalent to the equation:

$$d\theta_2^1 + \theta_k^1 \wedge \theta_2^k = \theta^1 \wedge \theta^2.$$
(12)

Therefore, using equations (10) and (12), we have

$$K = 1 - |\nabla\beta|^2 - 2\cos\alpha\beta_1 - \cos^2\alpha - (1 + \csc^2\beta)(a^2 + b^2) + 2b\sin\alpha\csc\beta\cot\beta + 2\sin\alpha\cot\beta\alpha_1 - |\nabla\alpha|^2 + 2a\csc\beta\alpha_2 - 2b\csc\beta\alpha_1 - \sin^2\alpha\cot^2\beta = 1 - (1 + \csc^2\beta)(a^2 + b^2) - 2b\csc\beta(\alpha_1 - \sin\alpha\cot\beta) + 2a\csc\beta\alpha_2$$
(13)
$$-|\nabla\beta + \cos\alpha e_1|^2 - |\nabla\alpha - \sin\alpha\cot\beta e_1|^2$$

Using (8) and the complex structure of S, we get

$$\theta_2^1 = \tan\beta (d\beta \circ J - 2\cos\alpha\theta^2) \tag{14}$$

Differentiating (14), we conclude that

$$d\theta_2^1 = (-(1 + \tan^2 \beta) |\nabla \beta|^2 - \tan \beta \Delta \beta - 2 \cos \alpha (1 + 2 \tan^2 \beta) \beta_1 + 2 \tan \beta \sin \alpha \alpha_1 - 4 \tan^2 \beta \cos^2 \alpha) \theta^1 \wedge \theta^2$$

where $\Delta = tr \nabla^2$ is the Laplacian of S. The Gaussian curvature is therefore given by:

$$K = -(1 + \tan^2 \beta) |\nabla \beta|^2 - \tan \beta \Delta \beta - 2 \cos \alpha (1 + 2 \tan^2 \beta) \beta_1 + 2 \tan \beta \sin \alpha \alpha_1 - 4 \tan^2 \beta \cos^2 \alpha.$$
(15)

From (13) and (15), we obtain the following formula for the Laplacian of S:

$$\tan \beta \Delta \beta = (1 + \csc^2 \beta)(a^2 + b^2) + 2b \csc \beta(\alpha_1 - \sin \alpha \cot \beta) - 2a \csc \beta \alpha_2 - \tan^2 \beta(|\nabla \beta + 2 \cos \alpha e_1|^2 - |\cot \beta \nabla \alpha + \sin \alpha (1 - \cot^2 \beta) e_1|^2) + \sin^2 \alpha (1 - \tan^2 \beta)$$
(16)

4 Gauss-Codazzi-Ricci equations for a minimal surface in S^5 with constant Contact angle β

In this section, we will compute Gauss-Codazzi-Ricci equations for a minimal surface in S^5 with constant Contact angle β .

Assume that the fundamental second form at the direction of the normal vector field e_3 diagonal. Then we have the following connection forms:

$$\begin{aligned}
\theta_1^3 &= a\theta^1 \\
\theta_2^3 &= -a\theta^2 \\
\theta_1^4 &= d\alpha + (-\sin\alpha\cot\beta)\theta^1 - a\csc\beta\theta^2 \\
\theta_2^4 &= d\alpha \circ J - a\csc\beta\theta^1 - (-\sin\alpha\cot\beta)\theta^2 \\
\theta_1^5 &= -\cos\alpha\theta^2 \\
\theta_2^5 &= -\cos\alpha\theta^1
\end{aligned}$$
(17)

Normal connection forms are:

$$\begin{aligned}
\theta_3^4 &= -\cot\alpha\csc\beta d\alpha \circ J + a\cot\alpha\cot^2\beta\theta^1 \\
&+ (-\cos\alpha\cot\beta\csc\beta + 2\sec\beta\cos\alpha)\theta^2 \\
\theta_3^5 &= (-\csc\beta\sin\alpha)\theta^1 - a\cot\beta\theta^2 \\
\theta_4^5 &= \cot\beta(d\alpha\circ J) - a\cot\beta\csc\beta\theta^1 + (\sin\alpha(\cot^2\beta - 1))\theta^2,
\end{aligned}$$
(18)

Using the connection form (17) and (18) in the Codazzi-Ricci equations, we have

$$d\theta_1^3 + \theta_2^3 \wedge \theta_1^2 + \theta_4^3 \wedge \theta_1^4 + \theta_5^3 \wedge \theta_1^5 = 0$$

This implies that

$$-a_{2} + a^{2}(\cot\alpha\csc\beta\cot^{2}\beta) - a\cot\alpha(\csc^{2}\beta + \cot^{2}\beta)\alpha_{2}$$

$$-\cos\alpha\csc\beta(2(\cot\beta - \tan\beta)\alpha_{1} - \sin\alpha(\cot^{2}\beta - 3)) + \cot\alpha\csc\beta|\nabla\alpha|^{2} = 0$$
(19)

Replacing the following (17) and (18) in the Codazzi-Ricci equations

$$\begin{aligned} d\theta_2^3 &+ \theta_1^3 \wedge \theta_2^1 + \theta_4^3 \wedge \theta_2^4 + \theta_5^3 \wedge \theta_2^5 &= 0 \\ d\theta_1^4 &+ \theta_2^4 \wedge \theta_1^2 + \theta_3^4 \wedge \theta_1^3 + \theta_5^4 \wedge \theta_1^5 &= 0 \\ d\theta_3^5 &+ \theta_1^5 \wedge \theta_3^1 + \theta_2^5 \wedge \theta_3^2 + \theta_4^5 \wedge \theta_3^4 &= 0 \end{aligned}$$

We get

$$a_1 + a(\cot \alpha \alpha_1 + 6 \tan \beta \cos \alpha) -2 \sec \beta \cos \alpha \alpha_2 = 0$$
(20)

Using the connection form (17) and (18) in the Codazzi-Ricci equations

$$\begin{aligned} d\theta_2^4 &+ \theta_1^4 \wedge \theta_2^1 + \theta_3^4 \wedge \theta_2^3 + \theta_5^4 \wedge \theta_2^5 &= 0 \\ d\theta_4^5 &+ \theta_1^5 \wedge \theta_4^1 + \theta_2^5 \wedge \theta_4^2 + \theta_3^5 \wedge \theta_4^3 &= 0 \\ d\theta_3^4 &+ \theta_1^4 \wedge \theta_3^1 + \theta_2^4 \wedge \theta_3^2 + \theta_5^4 \wedge \theta_3^5 &= 0 \end{aligned}$$

We have

$$a_{2} - a^{2}(\cot\alpha\sin\beta\cot^{2}\beta) + a\cot\alpha\alpha_{2}$$

$$+2\cos\alpha(\cot\beta - 3\tan\beta)) + 2\cos\alpha\sin\beta(\cot\beta - \tan\beta)\alpha_{1}$$

$$+\sin\alpha\cos\alpha\sin\beta(5 - \cot^{2}\beta) + \sin\beta\Delta\alpha = 0$$

$$(21)$$

Codazzi-Ricci equations

$$\begin{aligned} d\theta_1^2 &+ \theta_3^2 \wedge \theta_1^3 + \theta_4^2 \wedge \theta_1^4 + \theta_5^2 \wedge \theta_1^5 &= \theta^2 \wedge \theta^1 \\ d\theta_1^5 &+ \theta_2^5 \wedge \theta_1^2 + \theta_3^5 \wedge \theta_1^3 + \theta_4^5 \wedge \theta_1^4 &= 0 \end{aligned}$$

give the following equation

$$a^{2}(1 + \csc^{2}\beta) - 2a \csc \beta \alpha_{2}$$

+ $|\nabla \alpha|^{2} + 2 \sin \alpha (\tan \beta - \cot \beta) \alpha_{1} - 4 \tan^{2} \beta \cos^{2} \alpha$
- $\sin^{2} \alpha (1 - \cot^{2} \beta) = 0$ (22)

The following Codazzi equation is automatically verified

$$d\theta_2^5 + \theta_1^5 \wedge \theta_2^1 + \theta_3^5 \wedge \theta_2^3 + \theta_4^5 \wedge \theta_2^4 = 0$$

5 Proof of the Results

In this section, we will prove the results using the equations of the sections 3 and 4

5.1 Proof of Theorem 1

Using equations (19) and (21) we have:

$$\sin^2 \beta \Delta \alpha - \cot \alpha \csc \beta |\nabla \alpha|^2 - a^2 \cot \alpha \cos^2 \beta \cot^2 \beta + 2a \cot \alpha \sin \beta \cot^2 \beta \alpha_2 + 2\cos \alpha (\cot \beta - \tan \beta) \cos^2 \beta \alpha_1 - \sin \alpha \cos \alpha (5\sin^2 \beta - \cos^2 \beta \cot^2 \beta - 3) = 0$$

The equation above can be rewritten as the following equation

$$\Delta(\alpha) = \cot \alpha \csc^{3}(\beta) |\nabla \alpha|^{2} + a^{2} \cot \alpha \cot^{4}(\beta) - 2a \cot \alpha \csc \beta \cot^{2} \beta \alpha_{2} -2 \cos \alpha (\cot \beta - \tan \beta) \tan^{2} \beta \alpha_{1} + \sin \alpha \cos \alpha (5 - \cot^{4} \beta - 3 \csc^{2} \beta)$$
(23)

Using the equation (22) we get a as a function of α and β

$$a = \frac{2\csc\beta\alpha_2 + \sqrt{f}(\alpha,\beta)}{2(1 + \csc^2\beta)}$$

where $f(\alpha, \beta)$ is given by

$$f(\alpha,\beta) = 4\alpha_2^2 \cot^2 \beta - 4(1 + \csc^2 \beta)\alpha_1^2 -4(1 + \csc^2 \beta)(2\sin\alpha(\tan\beta - \cot\beta)\alpha_1 - 4\tan^2\beta\cos^2\alpha - \sin^2\alpha(1 - \cot^2\beta))$$

which prove Theorem 1.

5.2 Proof of Corollary 1

Suppose that K = 0 at the equation (15), we determine α_1 as a function of α

$$\alpha_1 = 2tg\beta \cot\alpha \cos\alpha \tag{24}$$

Now suppose that a = k, where k is constant, at the equation (20), we determine α_2 as a function of α_1

$$\alpha_2 = k(\frac{\cos\beta\csc\alpha\alpha_1}{2} + 3\sin\beta) \tag{25}$$

Using (24) at the equation above, we have α_2 as a function of α

$$\alpha_2 = k \sin\beta (\cot^2 \alpha + 3) \tag{26}$$

Finally, using α_1 and α_2 at the equation (19), we get α as a function of β , and therefore constant, which prove Corollary 1.

References

- B. Aebischer: Sympletic Geometry, Progress in Mathematics, Vol. 124 Springer-Verlag, Berlin-New York, 1992.
- [2] D. Blair: Contact Manifolds in Riemannian Geometry, Lecture Notes in Mathematics, Vol. 509, Berlin-Heidelberg-New York, Springer 1976.

- [3] S.S. Chern and J.G. Wolfson: Minimal surfaces by moving frames, American J. Math. 105 (1983), 59-83.
- [4] K. Kenmotsu: On a parametrization of minimal immersions R² into S⁵, Tohoku Math. J. 27 (1975), 83-90.
- [5] R.R. Montes and J.A. Verderesi: A new characterization of the Clifford Torus, Pre-Print, 2002.
- [6] R.R. Montes and J.A. Verderesi: Contact Angle for Immersed Surfaces in S^{2n+1} , to appear Differential Geometry and its Applications, 2007.
- [7] J.G. Wolfson: *Minimal surfaces in complex manifolds*, Ph.D. Thesis, University of California Berkeley, (1982).
- [8] S. Yamaguchi; M. Kon; Y. Miyahara: A theorem on C-totally real minimal surface, Proc. American Math. Soc. 54 (1976), 276-280.