

Attraction time for strongly reinforced walks

Codina Cotar*

Institut für Mathematik
Technische Universität Berlin
Strasse des 17. Juni 136
D-10623 Berlin, Germany

Vlada Limic[‡]

CNRS and Université de Provence
LATP UMR 6632, Centre de Mathématiques et Informatique
39, rue F. Joliot Curie
13453 Marseille Cedex 13, France

December 2, 2024

Abstract

We consider a class of strongly edge reinforced random walks, where the corresponding reinforcement weight function is non-decreasing. It is known by Limic and Tarrès (2006) that the attracting edge emerges with probability 1, whenever the underlying graph is locally bounded. We study the asymptotic behavior of the tail distribution of the (random) time of attraction. In particular, we obtain exact (up to multiplicative constant) asymptotics if the underlying graph has two edges. Next we show some extensions in the setting of finite and bounded degree infinite graphs. As a corollary we obtain that if the reinforcement weight has the form $w(k) = k^\rho$, $\rho > 1$, then (universally over finite graphs) the expected time to attraction is infinite if and only if $\rho \leq 1 + \frac{1+\sqrt{5}}{2}$.

AMS 2000 Subject Classification. 60G50, 60J10, 60K35

Key words and phrases. reinforced walk, time of attraction, attracting edge, strong reinforcement

Running Head: Attraction time under strong reinforcement

*Research conducted while on a postdoctoral fellow position at University of British Columbia, Vancouver

[†]Research supported in part by NSERC research grant, and by Alfred P. Sloan Research Fellowship

[‡]on leave from University of British Columbia, Canada

1 Introduction

Let \mathcal{G} be a locally finite graph with the edge set $E(\mathcal{G})$ and the vertex set $V(\mathcal{G})$. We will assume without further mention that \mathcal{G} is connected. We call any two vertices u, v connected by an edge *adjacent* (or *neighboring*), in this case we write $u \sim v$ and denote by $\{u, v\} = \{v, u\}$ the edge connecting them. We will denote by

$$|\mathcal{G}| = |E(\mathcal{G})|$$

the number of edges of \mathcal{G} , and by

$$\#\mathcal{G} = |V(\mathcal{G})|$$

the number of vertices of \mathcal{G} . Finally, denote by $D(\mathcal{G}) = \sup_{v \in V(\mathcal{G})} \text{degree}(v)$ the degree of \mathcal{G} , where for any $v \in V(\mathcal{G})$ $\text{degree}(v)$ equals the number of edges incident to v .

Let $(\ell_0^e, e \in E(\mathcal{G}))$ be given integers, and assume $\ell_0^e \geq 0, e \in E(\mathcal{G})$. Given a *reinforcement weight function* $w : \{0, 1, 2, \dots\} \mapsto (0, \infty)$, the edge-reinforced random walk (ERRW) on \mathcal{G} records nearest neighbor step transitions of a particle in $V(\mathcal{G})$. That is:

- (i) if currently at vertex $v \in V(\mathcal{G})$, in the next step the particle jumps to a vertex $u \in V(\mathcal{G})$ adjacent to v .
- (ii) the probability if a jump to u is *w-proportional* to the number of previous traversals of the edge $\{v, u\}$.

The more formal definition is as follows. If \mathcal{G} is a finite graph it seems natural from the point of notation to construct and study the edge reinforced random walk started at the *initial time*

$$t_0 := \sum_{e \in E(\mathcal{G})} \ell_0^e \geq 0,$$

a process starting at time 0 is obtained by a time shift. If \mathcal{G} is an infinite graph, just set $t_0 := 0$. Denote by I_n the (random) position of the edge reinforced random walk at time n . Then $I_{t_0} \in V(\mathcal{G})$ is the initial position, and $\{I_n, I_{n+1}\} \in E(\mathcal{G})$ for all $n \geq t_0$, almost surely. Let \mathcal{F}_n be the filtration

$$\mathcal{F}_n = \sigma\{I_k, k = 0, \dots, n, (\ell_0^e, e \in E(\mathcal{G}))\}. \quad (1)$$

Moreover, the dynamics of the edge reinforced random walk is prescribed according to the rule:

$$P(I_{n+1} = v | \mathcal{F}_n) 1_{\{I_n = u\}} = \frac{w(X_n^{\{u, v\}})}{\sum_{y \sim u} w(X_n^{\{u, y\}})} 1_{\{I_n = u, u \sim v\}},$$

where for any $e \in E(\mathcal{G})$,

$$X_n^e = \ell_0^e + \sum_{i=t_0}^{n-1} 1_{\{e \text{ was traversed at } i\text{th step}\}} = \ell_0^e + \sum_{i=t_0}^{n-1} 1_{\{\{I_i, I_{i+1}\} = e\}} \quad (2)$$

equals the *initial weight* ℓ_0^e incremented by the total number of (undirected) traversals of edge e prior to time n . Note that t_0 is chosen so that, whenever $V(\mathcal{G}) < \infty$, $\sum_{e \in E(\mathcal{G})} X_k^e = k$ for all $k \geq t_0$, almost surely. The starting weights $X_{t_0} := \ell_0^e$ are specified as deterministic above but one could use random variables instead in applications, and the definition (1) accounts for this possibility. Our results would then hold conditionally on the starting weights.

We denote by \mathcal{G}_1 the *range* of the edge reinforced random walk on \mathcal{G} . More precisely, we let

$$\mathcal{G}_1 = (V(\mathcal{G}_1), E(\mathcal{G}_1))$$

be the random subgraph of \mathcal{G} where for any $v \in V(\mathcal{G})$ we have

$$v \in V(\mathcal{G}_1) \Leftrightarrow \exists n \geq t_0 \text{ s.t. } I_n = v,$$

and, for any $e \in E(\mathcal{G})$,

$$e \in E(\mathcal{G}_1) \Leftrightarrow \exists n \geq t_0 \text{ s.t. } \{I_n, I_{n+1}\} = e.$$

Apart from the behavior analogous to recurrence or transience of Markov chains (see, for example, [10] or [7] or [9] theorems 5.2 and 5.6), ERRW may exhibit a very different asymptotic behavior as time increases. For example, it is easy to see, [11], [5] that the following assumption

$$\sum_k \frac{1}{w(k)} < \infty \tag{A0}$$

is sufficient for the event

$$\{\mathcal{G}_1 \text{ is a finite graph}\}$$

to have probability 1, whenever $D(\mathcal{G}) < \infty$. It is easy to find examples of locally bounded trees with $D(\mathcal{G}) = \infty$ such that (A0) holds but that \mathcal{G}_1 is infinite with positive probability. Sellke [11] provides (slightly peculiar) examples of edge reinforced random walks on \mathbb{Z} where $\sum_k 1/w(k)$ is finite over even k and infinite over odd k , but where \mathcal{G}_1 is still a finite graph, almost surely.

We next briefly discuss the link of our work to the recent literature. For a detailed review of a number of interesting results on edge reinforced random walk, we refer the reader to a recent survey of Pemantle [9] on stochastic reinforcement processes.

A result of Sellke [11] (the argument is also described in detail in [5], Section 2) implies that (A0) is sufficient and necessary for

$$P(\text{the walk ultimately traverses a single edge}) = 1, \tag{3}$$

whenever the underlying graph is bipartite and of bounded degree. Limic [5] proves that (A0) implies (3) on any graph of bounded degree, where the

reinforcement weight is a reciprocally summable power function. In a recent work, Limic and Tarrès [6] show that for a fairly general class of reinforcement weights (in particular, whenever w is a non-decreasing function satisfying (A0)) (3) holds on any graph of bounded degree. We will refer to any weight w satisfying condition (A0) as *strong*, and to the corresponding ERRW as *strongly reinforced walk*.

The current paper assumes the setting of [6], and is devoted to the study of the tail behavior of the *time of attraction*

$$\begin{aligned} T &= \inf\{k \geq 0 : \exists e \in E(\mathcal{G}) \text{ s.t. } \forall f \neq e \ X_k^f = \max_{m \geq k} X_m^f\} \\ &= \inf\{k \geq 0 : \{I_n, I_{n+1}\} = \{I_{n+1}, I_{n+2}\}, \forall n \geq k\}, \end{aligned} \quad (4)$$

that is, the first time after which only the *attracting* edge is traversed. This random variable is an important statistic, useful for applications (e.g. [4] or [3]). In this paper we make a few connections to the literature on behavioral science of social insects, and refer the reader to [9] for a diverse list of potential applications.

It is clear that the sequence of tail probabilities $(P(T > k), k \geq 1)$ depends on the structure of the underlying graph \mathcal{G} , the weight function w , the initial weights ℓ_0 , and the initial position I_{t_0} . However, the results of sections 3.1-3.2 verify an interesting universality-type behavior. Namely, fix w satisfying (A0), let \mathcal{G} be an arbitrary finite graph with some prescribed initial edge weights and initial position, and let \mathcal{G}' be the simple two-edge graph from Section 2 with initial weights equal to 1 on both edges. Then if $P^{\mathcal{G}}$ (resp. $P^{1,1}$) denotes the law of the corresponding ERRW on \mathcal{G} (resp. \mathcal{G}'), the asymptotic order of magnitude of $P^{\mathcal{G}}(T > k)$ is induced by that of $P^{1,1}(T > k)$. To some extent this holds also on infinite trees of bounded degree, cf. Corollary 18.

Definition 1 For sequences a_k, b_k of real numbers, we write $a_k \asymp b_k$ iff

$$\frac{a_k}{b_k} \in [c, C], \quad k \geq 1 \text{ for some } c, C \in (0, \infty).$$

The rest of the paper is organized as follows. Section 2 is devoted to the careful study of the two-edge setting. In particular, in Section 2.1 we prove Corollary 5, stated less precisely as follows: Assume that $E(\mathcal{G})$ consists of two elements, and let

$$Z_\infty := \#\{\text{times the non-attracting edge is traversed}\},$$

then under assumption (A0),

$$P(Z_\infty = \ell) \asymp \frac{1}{w(\ell)}, \quad \ell \rightarrow \infty.$$

A stronger statement, due to Robin Pemantle (personal communication)

$$\lim_{\ell \rightarrow \infty} w(\ell)P(Z_\infty = \ell) \in (0, \infty), \quad (5)$$

holds in this simple setting. The limit will become apparent in the course of the proof sketched at the end Section 2.2.

Lemma 6 of Section 2.3 provides the initial order of magnitude estimates on the tail probability $P(T > k)$, as $k \rightarrow \infty$. More precisely, we prove that

$$P(T > n) \asymp \sum_{k=n+1}^{\infty} \sum_{\ell \leq k/2} \left[\frac{1}{w(k-\ell)} \prod_{j=0}^{\infty} \frac{w(k+j-\ell)}{w(k+j-\ell) + w(\ell+1)} + \frac{1}{w(\ell)} \prod_{j=0}^{\infty} \frac{w(j+\ell-1)}{w(j+\ell-1) + w(k+1-\ell)} \right].$$

Such an expression seems awkward for application, and we work further to find simplifications. In particular, Theorem 9 shows a simpler looking asymptotics of the tail distribution of T , under the additional assumption (A1) that w is a non-decreasing function. The main idea is simple: the event which with overwhelming probability contributes to the event $\{T = k+1\}$ of interest is the one where at time k the weaker edge (i.e. the edge with lower current number of traversals) is traversed, and at all future times the stronger edge is traversed. Therefore (Z_k denotes the number of traversals of the less traversed edge at time k)

$$P(T = k+1) \asymp \sum_{\ell \leq k/2} P(Z_k = \ell) \frac{w(\ell)}{w(k-\ell) + w(\ell)} P(Z_{\infty} = \ell+1 | Z_{k+1} = \ell+1).$$

For ℓ close to $k/2$ it is plausible that $P(Z_k = \ell)$ is sufficiently small so that the contribution in the above sum vanishes asymptotically. For ℓ small the middle term $w(\ell)/(w(k-\ell) + w(\ell))$ is again small. In order to estimate well the above sum, one then needs to find the interval of indices ℓ which make up the most of the contribution. As shown in Proposition 2, $P(Z_k = \ell) \asymp \frac{w(k-\ell)+w(\ell)}{w(k-\ell)w(\ell)}$, so it is plausible that the overwhelming contribution to the sum comes approximately from the range of indices where $P(Z_{\infty} = \ell+1 | Z_{k+1} = \ell+1) \asymp 1$. For formal estimates see Section 2.3.

In Section 2.4 we include specific calculations for cases of w that already have been used (or might be used) in applications, cf. [2], [4], that satisfy the assumptions (A0)-(A1). In particular we paraphrase

Theorem 10 (a) *If $w(k) = k^{\rho}$ for some fixed $\rho > 1$, and if $\rho' = (\rho - 1)/\rho$*

$$P(T > k) \asymp \frac{1}{k^{\rho-\rho'-1}}.$$

In particular $E(T)$ is infinite if $\rho \leq 1 + \frac{1+\sqrt{5}}{2}$ and finite if $\rho > 1 + \frac{1+\sqrt{5}}{2}$.

This type of result should be particularly interesting for applications. In fact, in [4], for a similar model, the reinforcement weight is set to $w(k) = k^{\rho}$ and real life data is compared to different values of ρ and initial configurations. More precisely, the authors study a colony of ants, which explores a chemically unmarked territory randomly, starting from the nest. The exploration

is done on two branches A and B . Initially both branches are equally likely to be chosen. However, each ant that passes along one of the two branches leaves an additional pheromone mark and in this way influences the following ant's decision in choosing A or B . In the real-life experiment it is observed that after initial fluctuations, one of the two branches becomes more or less completely preferred to the other. In their (reinforcement) model, k represents the number of ants that have chosen a particular branch and ρ determines the degree of non-linearity. The model is used for further study of the explorer movement pattern in two-dimensional space.

Section 3 is devoted to analysis on general graphs of bounded degree. In particular, we are interested in a universality-type behavior of the tail distribution $P^{\mathcal{G}}(T > \cdot)$ over graphs, once the reinforcement weight function w is fixed. In the course of our analysis we also obtain exponential bounds (cf. Lemma 25) on the tail distribution of $|\mathcal{G}_1|$, and in particular some information on the distance of the attracting edge from the starting point. Providing a universal lower bound on $P^{\mathcal{G}}(T > \cdot)$ in terms of the corresponding quantity in the two-edge graph setting turns out to be simple (cf. Lemma 12), however finding an analogous upper bound is not as simple. Section 3.1 is devoted to analysis on trees, here initial universality-type behavior is demonstrated using slick comparison (coupling) arguments. Section 3.2 is devoted to the finite graph setting. By generalizing the technique of Section 2, a fairly general universality-type behavior is shown, under an additional assumption (A2). Finally, Section 3.3 discusses extensions to infinite graph setting.

In the rest of the paper we assume that all edges have “**trivial**” **initial weight** $\ell_0 \equiv 1$, unless otherwise specified. Also, we will denote by $a \wedge b$ (resp. $a \vee b$) the minimum (resp. the maximum) of two numbers a and b , and by $\lfloor a \rfloor$ the integer part of a number a .

2 Two-edge case

The ERRW on graph \mathcal{G} that contains only two edges is the prototype model of interest. Several interesting qualitative features, specific to edge-reinforcement with particular reinforcement weight function w , are already observed and usually relatively easy to verify. This process also corresponds to a generalized urn model, see for example [1] or [11]. A recent study of Oliveira and Spencer [8] concerns finer properties of this urn model in the case where $w(k) = k^\rho$ for some $\rho > 1$.

We will initially assume that \mathcal{G} contains two vertices 0 and 1, and two edges, *green* and *red*, connecting them. Abbreviate

$$G_n := X_n^{\text{green}}, \quad R_n := X_n^{\text{red}}.$$

In the rest of this section, we also assume that the initial configuration on the two edges is $G_2 = R_2 = 1$, unless otherwise specified. We use the notation

$P^{a,b}$ for the law of the system with the initial configuration $G_{a+b} = a, R_{a+b} = b$. When not in risk of confusion, we use simply P for the law $P^{1,1}$.

The other natural choice of a graph with two edges is the one spanned by three vertices $-1, 0, 1$ and a green edge that connects 0 and -1 and a red one that connects 0 and 1 . In the study of this model we mainly concentrate on the case where the initial weights a and b are of opposite parity. We denote by $\bar{P}^{a,b}$ the law of the ERRW on the two-edge graph spanned by $-1, 0, 1$ started (without loss of generality) at the initial position 0 . Note that the study of $\bar{P}^{a,b}$ is necessary as it will be needed later for the subsequent analyses of the time of attraction of the ERRW on trees and on finite graphs. The main results in Section 3 are expressed in terms of both $P^{a,b}$ and $\bar{P}^{a,b}$.

Observe that under $\bar{P}^{a,b}$ we have $G_{a+b} = a, R_{a+b} = b$ and $G_{a+b+2j} - G_{a+b+2j-2} \in \{0, 2\}$, $R_{a+b+2j} - R_{a+b+2j-2} + G_{a+b+2j} - G_{a+b+2j-2} = 2$, for all $j \geq 1$.

2.1 Some preliminary estimates

Due to monotonicity, $R_\infty := \lim_{k \rightarrow \infty} R_k$ and $G_\infty := \lim_{k \rightarrow \infty} G_k$ exist almost surely as $(0, \infty]$ -valued random variables. Define

$$Z_k = \min\{G_k, R_k\}.$$

Note that $Z_k \leq Z_{k+1}$ and that the limit

$$Z_\infty := \lim_{k \rightarrow \infty} Z_k = R_\infty \wedge G_\infty \quad (6)$$

is an almost surely finite random variable, since the reinforcement is strong.

Proposition 2 Define $c := \max_{k \geq 2} \left(\frac{w(k-1)+w(1)}{w(k-1)} \right)$.

(a) For any $1 \leq \ell < \frac{k}{2}$, we have

$$\frac{w(1)}{c} P(Z_\infty = 1) \frac{w(k-\ell) + w(\ell)}{w(k-\ell)w(\ell)} \leq P(Z_k = \ell) \leq w(1) \frac{w(k-\ell) + w(\ell)}{w(k-\ell)w(\ell)}.$$

(b) For any $\ell \geq 1$,

$$\frac{w(1)}{c} \cdot \frac{P(Z_\infty = 1)}{w(\ell)} \leq P(Z_{2\ell} = \ell) \leq \frac{w(1)}{w(\ell)}.$$

Note that the lower bounds are interesting only for strongly reinforced walks, where $P(Z_\infty < \infty) = 1$ and $P(Z_\infty = 1) > 0$. A careful reader of the proof will note that all the above inequalities are strict, however we do not anticipate any use of this fact.

Proof. We will prove the upper bounds by induction, and the lower bounds will follow in a similar way as indicated at the end of the proof. Note first that for $\ell \leq \frac{k}{2} - 1$

$$P(Z_k = \ell) = P(Z_{k-1} = \ell) \frac{w(k-\ell-1)}{w(k-\ell-1)+w(\ell)} + P(Z_{k-1} = \ell-1) \frac{w(\ell-1)}{w(\ell-1)+w(k-\ell)}. \quad (7)$$

Similarly, we also have in the special cases $k = 2\ell$

$$P(Z_{2\ell} = \ell) = P(Z_{2\ell-1} = \ell - 1) \frac{w(\ell - 1)}{w(\ell - 1) + w(\ell)}, \quad (8)$$

and $k = 2\ell + 1$

$$P(Z_{2\ell+1} = \ell) = P(Z_{2\ell} = \ell) + P(Z_{2\ell} = \ell - 1) \frac{w(\ell - 1)}{w(\ell - 1) + w(\ell + 1)}. \quad (9)$$

Since any probability is bounded by 1, we have trivially

$$P(Z_k = 1) < w(1) \frac{w(k - 1) + w(1)}{w(k - 1)w(1)},$$

an observation that will be used in the base and in each step of the induction.

The base of induction is the case $\ell = 1$, $k = 2\ell + 1 = 3$, and the statement here is trivial as noted above.

Let us assume now that the upper bound inequalities in the statements (a) and (b) of the theorem hold for all $i \leq k - 1$ and $\ell \leq \frac{i}{2}$. For the induction step we need to show the bounds hold for $i = k$ and $\ell \leq \frac{k}{2}$. Suppose first that $\ell \leq \lfloor \frac{k}{2} \rfloor - 1$. Then by (7) and the induction hypothesis, we have

$$P(Z_k = \ell) \leq \frac{w(1)}{w(\ell)} + \frac{w(1)}{w(k - \ell)} = w(1) \frac{w(k - \ell) + w(\ell)}{w(k - \ell)w(\ell)}$$

For the two atypical cases $k = 2\ell$ and $k = 2\ell + 1$ we have similarly by (8)

$$P(Z_{2\ell} = \ell) = P(Z_{2\ell-1} = \ell - 1) \frac{w(\ell - 1)}{w(\ell - 1) + w(\ell)} < \frac{w(1)}{w(\ell)},$$

and by (9)

$$P(Z_{2\ell+1} = \ell) < \frac{w(1)}{w(\ell)} + \frac{w(1)}{w(\ell + 1)} = w(1) \frac{w(\ell) + w(\ell + 1)}{w(\ell)w(\ell + 1)}.$$

The proof of the lower bounds is symmetric. Note that $P(Z_k = 1) \geq P(Z_\infty = 1)$, and the choice of c was precisely made so that the lower bound holds both in (a) for any $k \geq 3$ and $\ell = 1$, and in (b) for $\ell = 1$. Given these initial bounds, the above argument by induction on k will carry over to yield the lower bound of (a) and (b). \square

The result above under the law $P^{1,1}$ generalizes to the setting of law $\bar{P}^{1,2}$ on two-edge graph with three vertices in the following way.

Proposition 3 *Define*

$$\bar{c} := \max_{k \geq 2 \text{ even}} \left(\frac{w(k) + w(1)}{w(k)} \right) \vee \max_{k \geq 1 \text{ odd}} \left(\frac{w(k) + w(2)}{w(k)} \right).$$

For any $k \geq 1$, $1 \leq o, e \leq 2k$, such that o is odd and e is even and $o + e = 2k + 1$ we have

$$\bar{P}^{1,2}(G_{2k+1} = o, R_{2k+1} = e) \leq (w(1) \vee w(2)) \frac{w(o) + w(e)}{w(o)w(e)},$$

and

$$\frac{w(1) \wedge w(2)}{\bar{c}} \cdot \frac{w(o) + w(e)}{w(o)w(e)} \prod_{j=1}^2 \bar{P}^{1,2}(Z_\infty = j) \leq \bar{P}^{1,2}(G_{2k+1} = o, R_{2k+1} = e).$$

Proof. Abbreviate $\bar{P}^{1,2}$ by P . We first concentrate on the upper bound. If either $o = 1$ or $e = 2$ (or both) the upper bound is trivial, so the base of induction is verified. Now, as in the previous proposition, if both $o > 1$ and $e > 2$ we apply induction step using

$$P(G_{2k} = o, R_{2k} = e) = P(G_{2k-2} = o-2, R_{2k-2} = e) \frac{w(o-2)}{w(o-2) + w(e)} + \\ P(G_{2k-2} = o, R_{2k-2} = e-2) \frac{w(e-2)}{w(o-2) + w(e-2)}.$$

Similarly, note that $\bar{P}^{1,2}(G_{2k+1} = 1, R_{2k+1} = 2k) \geq \bar{P}^{1,2}(G_\infty = j) = \bar{P}^{1,2}(Z_\infty = j)$ and $\bar{P}^{1,2}(G_{2k+1} = 2k-1, R_{2k+1} = 2) \geq \bar{P}^{1,2}(R_\infty = 2) = \bar{P}^{1,2}(Z_\infty = 2)$, and so the lower bound holds for any $k \geq 1$ whenever $o = 1$ or $e = 2$, with the above choice of \bar{c} . Given these initial bounds, one applies the induction once again to prove the general lower bound. \square

Moreover, using the same technique as above one arrives to the following general result.

Theorem 4 *Let $a, b \geq 1$ and define*

$$c(a, b) \equiv c(a, b, w) := \max_{k \geq a+b} \left(\frac{w(k-a) + w(a)}{w(k-a)} \right) \text{ and} \\ \bar{c}(a, b) \equiv c(a, b, w) := \max_{k \geq b, k \text{ even}} \left(\frac{w(k) + w(a)}{w(k)} \right) \vee \max_{k \geq a, k \text{ odd}} \left(\frac{w(k) + w(b)}{w(k)} \right).$$

Then for any $k \geq a + b$ and $\ell \geq a \wedge b$ we have

$$\frac{w(a)}{c(a, b)} P^{a,b}(Z_\infty = 1) \frac{w(k-\ell) + w(\ell)}{w(k-\ell)w(\ell)} \leq P^{a,b}(Z_k = \ell) \leq w(a) \frac{w(k-\ell) + w(\ell)}{w(k-\ell)w(\ell)},$$

and, assuming that a is odd while b is even,

$$\frac{w(a) \wedge w(b)}{\bar{c}(a, b)} \prod_{j \in \{a, b\}} \bar{P}^{a,b}(Z_\infty = j) \frac{w(k-\ell) + w(\ell)}{w(k-\ell)w(\ell)} \\ \leq \bar{P}^{a,b}(Z_k = \ell) \leq (w(a) \vee w(b)) \frac{w(k-\ell) + w(\ell)}{w(k-\ell)w(\ell)}.$$

These pre-asymptotic estimates will be useful in further analysis.

Before continuing note that a direct consequence is the following result, already announced in the introduction.

Corollary 5 *Assuming (A0), under any of the laws from Theorem 4 there exist $c, C \in (0, \infty)$ depending on the choice of the law, such that*

$$P(Z_\infty = \ell) \in \left(\frac{c}{w(\ell)}, \frac{C}{w(\ell)} \right).$$

2.2 The “time-line” construction

Here we briefly recall the construction of the edge reinforced walk using independent families of exponentials, see [1],[11], or [5]. In the current work we will use it mainly in the context of trees. To simplify the notation we focus on two cases, where \mathcal{G} is either a two-edge graph or a “star” with m fingers, the reader can easily do the general case.

First assume that \mathcal{G} contains two vertices 0 and 1, and two edges e_G and e_R connecting them. Fix initial weights $\ell_0^{e_G} = a$ and $\ell_0^{e_R} = b$ and let $I_{t_0} = 0$, where $t_0 = a + b$. Note that the corresponding edge reinforced random walk has the law $P^{a,b}$.

For each $k \geq 1$ let E_k^G, E_k^R be exponential (rate $w(k)$) random variables, and let $\{E_k^G, k \geq 1\}, \{E_k^R, k \geq 1\}$ be two independent families of independent random variables. Let

$$T_\infty^G := \sum_{k \geq 0} E_{a+k}^G, \quad T_\infty^R := \sum_{k \geq 0} E_{b+k}^R. \quad (10)$$

Note that T_∞^G and T_∞^R are independent and finite (due to (A0)) almost surely. In Figure 1 intervals between subsequent dots have length E_{a+k}^G or E_{b+k}^R corresponding to the edge and to the index of the chronological order k , and the limits T_∞^G, T_∞^R are indicated as well.




Figure 1

Here is how one can construct a realization of the edge reinforced random walk on \mathcal{G} from the above data, or (informally) from the figure. Find the minimum of E_a^G and E_b^R by “simultaneously erasing at rate 1 in the chronological direction” the time-lines corresponding to both edges until encountering the first dot. In the figure this happens to be the first dot on the time-line corresponding to edge e_G , i.e. $E_a^G < E_b^R$. Thus the particle moves from 0 to 1 traversing the edge e_G in the first step. Note that due to the properties of exponentials, the probability of this move is exactly $w(a)/(w(a)+w(b))$. Continue by simultaneous erasing (the previously non-erased parts of) time-lines corresponding both edges until the next dot. In the figure, it appears on the time-line of e_R . Hence the particle traverses the edge e_R in the second step

to go back from 1 to 0. Due to the memoryless properties of exponentials, the (residual) length of the interval until the first dot on the time-line of e_R is again distributed as exponential (rate $w(b)$) random variable, independent of all other data. Therefore, again the probability of this transition (namely, $w(b)/(w(b) + w(a+1))$) matches the one of the edge reinforced random walk. Continue the above procedure of simultaneous erasure of the time-lines. In this way, the steps of the corresponding edge reinforced random walk are generated inductively.

As its by-product, a “continuized” version of the edge reinforced random walk arises: here the particle makes the jumps exactly at times when the dots are encountered. If we denote the position of the particle (in the new process) at time s by $\tilde{I}(s)$ and if $\tau_0 = 0$ and $0 < \tau_1 < \tau_2 < \dots$ are the subsequent jump times of the particle, then the discrete-time edge reinforced random walk constructed above, and its continuized version are coupled as follows

$$I_k \equiv \tilde{I}(\tau_k), \quad k \geq 0.$$

One typically says that I is the skeleton process of \tilde{I} .

Now let \mathcal{G} be a labeled tree with the central vertex 0 which is connected via edge e_i to each leaf vertex i , $i = 1, \dots, m$. We call such \mathcal{G} a star with m fingers. Fix initial weights $\ell_0^{e_i} = \ell_0^i$ and let $I_{t_0} = 0$, where $t_0 = \sum_{i=1}^m \ell_0^i$. In particular, note that if $m = 2$ and $\ell_0^1 = a$, $\ell_0^2 = b$ then the corresponding edge reinforced random walk has the law $\bar{P}^{a,b}$.

Similarly to the previous construction, for each $i = 0, \dots, m$ and $k \geq 1$ let E_k^i be exponential (rate $w(k)$) random variable and let $\{E_k^i, i = 0, \dots, m, k \geq 1\}$ be a family of independent random variables. Denote by

$$T_\infty^i := \sum_{k \geq 0} E_{\ell_0^i + 2k}^i, \quad i = 0, \dots, m.$$

The multiple “2” in the subscript comes from the fact that the particle traverses each edge twice before coming back to the central vertex. Note again that the above m random variables are continuous, independent and finite almost surely. In Figure 2 intervals between subsequent dots have length of $E_{\ell_0^i + 2k}^i$ for the corresponding i and k .




Figure 2

One constructs a realization of the corresponding edge reinforced random walk from the above data analogously to the two-edge setting, the only difference being now that in every second step, when not at the central vertex 0, the particle jumps almost surely back to 0. From the above figure one can read off the first four steps of the walk as $I_{t_0+1} = 1$, $I_{t_0+1} = 0$, $I_{t_0+1} = 2$, and $I_{t_0+1} = 0$. The reader will quickly verify that, due to the properties of exponentials, the probability of transitions match those of the edge reinforced random walk.

Again a continuized version of the edge reinforced random walk emerges, where there are various possibilities to account for the “singular” behavior of the walk at the leaves of \mathcal{G} . For example, one could use the random variables $E_{\ell_0+2k+1}^i$ (that did not play any role in the construction of the walk) as subsequent waiting times at the leaf i , for each $i = 1, \dots, m$.

Proof of (5). We concentrate on the case $P = P^{1,1}$ and show that the limit in (5) equals $2 \int_0^\infty f(x)^2 dx$, where f is the (continuous) density of $T_\infty^G = \sum_{k=1}^\infty E_k^G$. Let $S_\ell^R := \sum_{k=1}^{\ell-1} E_k^R$ and f_ℓ be the density of S_ℓ^R . Then since $P(Z_\infty = \ell) = P(G_\infty = \ell) + P(R_\infty = \ell) = 2P(R_\infty = \ell)$ by symmetry, we have

$$\begin{aligned} w(\ell)P(Z_\infty = \ell) &= 2w(\ell)P(S_\ell^R < T_\infty^G < S_{\ell+1}^R) \\ &= 2w(\ell)E(1_{\{S_\ell^R < T_\infty^G\}}P(S_{\ell+1}^R > T_\infty^G | S_\ell^R, T_\infty^G)) \\ &= 2w(\ell)E(1_{\{S_\ell^R < T_\infty^G\}}e^{-w(\ell)(T_\infty^G - S_\ell^R)}) \\ &= 2 \int_0^\infty dt f(t) \int_0^t ds f_\ell(s) w(\ell) e^{-w(\ell)(t-s)}, \end{aligned} \quad (11)$$

where the first identity is clear from the graphical construction above, the second is a simple conditioning relation, the third uses the fact that $S_{\ell+1}^R - S_\ell^R = E_\ell^R$ is exponential (rate $w(\ell)$), independent from the σ -field generated by S_ℓ^R, T_∞^G , and the last one is the same expression written in terms of densities of S_ℓ^R and T_∞^G .

In order to prove that the integral in (11) converges to $\int_0^\infty f(x)^2 dx$, it suffices to show that, as $\ell \rightarrow \infty$,

$$\int_0^t ds f_\ell(s) w(\ell) e^{-w(\ell)(t-s)} \rightarrow f(t), \quad (12)$$

and that the left-hand side above is uniformly bounded in ℓ and in t . In fact, $f_\ell(s)$ is bounded by a fixed constant in both ℓ and s , as a convolution of an exponential and another density. Moreover,

$$|f_\ell(s_1) - f_\ell(s_2)| \leq \int |f_1(s_1 - u) - f_1(s_2 - u)| g_\ell(u) du,$$

where f_1 is the (exponential) density of S_1^R , and g_ℓ is the density of $S_\ell^R - S_1^R$. We conclude that $(f_\ell, \ell \geq 1)$ is a uniformly continuous family of functions.

The convergence in (12) is now not difficult to show by writing $f_\ell(s) = (f_\ell(s) - f_\ell(t)) + (f_\ell(t) - f(t)) + f(t)$ and using the fact that $S_\ell^R \nearrow T_\infty^R \stackrel{d}{=} T_\infty^G$, as well as $w(\ell) \rightarrow \infty$, so that the integral concentrates around t , as $\ell \rightarrow \infty$.

Note that one can modify the above proof to show analogous statement under any law $P^{a,b}$, $a, b \geq 1$, by using instead $S_\ell^G := \sum_{k=a}^{\ell-1} E_k^G$, $S_\ell^R := \sum_{k=b}^{\ell-1} E_k^R$, $T_\infty^G := \sum_{k \geq 0} E_{a+k}^G$, and $T_\infty^R := \sum_{k \geq 0} E_{b+k}^R$. Namely, one can easily verify using the above method that

$$\lim_{\ell \rightarrow \infty} w(\ell) P(Z_\infty = \ell) = 2 \int f_{T_\infty^G}(t) f_{T_\infty^R}(t) dt,$$

where $f_{T_\infty^G}$ and $f_{T_\infty^R}$ are densities of T_∞^G and T_∞^R , respectively.

2.3 Time of attraction

Next consider the time of attraction

$$T := \min\{k \geq 1 : Z_l = Z_{l+1}, \text{ for all } l \geq k\}.$$

Note that $\{T = k + 1\}$ is a disjoint union of A_k^s and A_k^w , where

$A_k^s = \{\text{the less (or equally) traversed edge is chosen at time } k \text{ and the remaining edge is chosen at all later times}\},$

$A_k^w = \{\text{the more traversed edge is chosen at time } k \text{ and the remaining edge is chosen at all later times}\}.$

Given $\{Z_k = \ell\}$ for some $\ell < k/2$, the event A_k^s happens with probability

$$\frac{w(\ell)}{w(\ell) + w(k - \ell)} \prod_{j=k}^{\infty} \frac{w(j - \ell)}{w(j - \ell) + w(\ell + 1)}, \quad (13)$$

while A_k^w happens with probability

$$\frac{w(k - \ell)}{w(k - \ell) + w(\ell)} \prod_{j=\ell}^{\infty} \frac{w(j)}{w(j) + w(k - \ell + 1)}. \quad (14)$$

Finally, if $2\ell = k$, then $P(A_k^s | Z_k = \ell)$ is an expression analogous to (13)

$$\prod_{j=k}^{\infty} \frac{w(j - k/2)}{w(j - k/2) + w(k/2 + 1)}.$$

It will be useful to abbreviate

$$W_k(\ell) := \prod_{j=0}^{\infty} \frac{w(k + j - \ell)}{w(k + j - \ell) + w(\ell)}, \quad \overline{W}_k(\ell) := \prod_{j=0}^{\infty} \frac{w(k + 2j - \ell)}{w(k + 2j - \ell) + w(\ell)}. \quad (15)$$

Note that if $\ell < k/2$ then

$$W_k(\ell) = P(Z_\infty = \ell | Z_k = \ell). \quad (16)$$

Then the identities (13)-(15) yield

$$P(A_k^s | Z_k = \ell) = \frac{w(\ell)}{w(\ell) + w(k - \ell)} W_{k+1}(\ell + 1), \quad \ell < k/2, \quad (17)$$

$$P(A_k^s | Z_k = \ell) = W_{k+1}(k/2 + 1), \quad k = 2\ell, \quad (18)$$

$$P(A_k^w | Z_k = \ell) = \frac{w(k - \ell)}{w(k - \ell) + w(\ell)} W_{k+1}(k + 1 - \ell), \quad (19)$$

and similar identities hold under the laws $P^{a,b}$ and $\bar{P}^{a,b}$ (with \bar{W} used in place of W), for $a, b \geq 1$.

We have, as discussed above,

$$\begin{aligned} P(T = k + 1) &= \sum_{j=1}^{\lceil k/2 \rceil - 1} P(Z_k = j) (P(A_k^s | Z_k = j) + P(A_k^w | Z_k = j)) \\ &\quad + P(Z_k = k/2) P(A_k^s | Z_k = k/2), \end{aligned}$$

Now (17)-(19) together with Theorem 4 imply the following asymptotic formula:

Lemma 6 *Under the law $P^{a,b}$ we have*

$$P(T = k + 1) \asymp \sum_{\ell=a \wedge b}^{k/2} \left[\frac{1}{w(k - \ell)} W_{k+1}(\ell + 1) + \frac{1}{w(\ell)} W_{k+1}(k + 1 - \ell) \right].$$

Similarly under $\bar{P}^{a,b}$, where a is odd and b is even, and k is odd (since the initial time is $a + b$ and the initial position is 0),

$$P(T = k + 2) \asymp \sum_{\ell=a, \text{ odd}}^{k-b} \frac{1}{w(k - \ell)} \bar{W}_{k+2}(\ell + 2) + \sum_{\ell=b, \text{ even}}^{k-a} \frac{1}{w(k - \ell)} \bar{W}_{k+2}(\ell + 2).$$

From now on we will also assume that

$$w(k) \text{ is non-decreasing in } k. \quad (A1)$$

This will be useful for future estimates since then

Lemma 7 (a) $W_k(\cdot)$ is a non-increasing function for each $k \geq 1$, on the interval $[1, k/2]$,

(b) $W_k(\ell)$ is a non-decreasing function for each ℓ on the interval $[2\ell, \infty)$.

(c) For any $k, \ell \geq 1$ such that $\ell \leq k/2$ we have $W_k(\ell) \geq W_{k+1}(\ell + 1)$,

(d) For each fixed ℓ we have $\lim_{k \rightarrow \infty} W_k(\ell) \rightarrow 1$.

Proof. Note that

$$W_k(\ell) = \prod_{j=0}^{\infty} \left(1 + \frac{w(\ell)}{w(k+j-\ell)} \right)^{-1}, \quad W_{k+1}(\ell+1) = \prod_{j=0}^{\infty} \left(1 + \frac{w(\ell+1)}{w(k+j-\ell)} \right)^{-1}. \quad (20)$$

(a) We need to show that $W_k(\ell) \geq W_k(\ell+1)$, for all $\ell \in [1, k/2]$. By assumption (A1), we have

$$w(k+j-\ell)w(\ell+1) \geq w(k+j-\ell-1)w(\ell), \quad \text{for all } j \geq 0,$$

from which we get that for all $j \geq 0$

$$\left(1 + \frac{w(\ell)}{w(k+j-\ell)} \right)^{-1} \geq \left(1 + \frac{w(\ell+1)}{w(k+j-\ell-1)} \right)^{-1}.$$

(b) Here we need to show that $W_k(\ell) \leq W_{k+1}(\ell)$, for all $k \geq 1$. By assumption (A1), we have $w(k+j-\ell) \leq w(k+1+j-\ell)$, $\forall j \geq 0$, which together with (20) directly implies the above inequality.

(c) Again by assumption (A1) we have $w(\ell) \leq w(\ell+1)$, $\forall \ell \geq 1$, so that representation (20) implies the claim.

(d) This is an easy consequence of (6) using probabilistic interpretation (16) (equivalently, one can use the algebraic definition and (A0)). \square

Corollary 8 *Assume (A1). Then for $\ell < k/2$, we have*

$$P(A_k^w | Z_k = \ell) \leq 2P(A_k^s | Z_k = \ell).$$

Proof. Use (17) and (19). Note that

$$w(k-\ell) + w(\ell+1) \leq 2(w(k+1-\ell) + w(\ell)),$$

for all k , due to assumption (A1), and that for each $i \geq 1$, the i th term

$$\frac{w(\ell+i)}{w(\ell+i) + w(k-\ell+1)}$$

in the infinite product of (14) is bounded above by the i th term

$$\frac{w(k+i-\ell)}{w(\ell+1) + w(k+i-\ell)},$$

in the infinite product of (13), again since (A1) holds. \square

Therefore, to obtain asymptotic (in the sense of relation \asymp) upper and lower bounds on $P(T = k+1)$ it suffices to study only

$$\begin{aligned} P(T = k+1, A_k^s) &= \sum_{\ell=1}^{k/2} P(Z_k = \ell) P(A_k^s | Z_k = \ell) \\ &\asymp \sum_{\ell=1}^{k/2} \frac{1}{w(k-\ell)} W_{k+1}(\ell+1), \end{aligned}$$

implying

Theorem 9 *If (A0), (A1) hold and if P is $P^{a,b}$ for some fixed $a, b \geq 1$ then*

$$P(T = k + 1) \asymp \sum_{\ell=1}^{k/2} \frac{1}{w(k - \ell)} W_{k+1}(\ell + 1), \quad (21)$$

and

$$P(T > n) \asymp \sum_{k=n+1}^{\infty} \sum_{\ell=1}^{k/2} \frac{1}{w(k - \ell)} W_{k+1}(\ell + 1). \quad (22)$$

Analogous statements are valid if P is $\bar{P}^{a,b}$, where W needs to be replaced by \bar{W} . Namely, assume that a and b are of opposite parity. Then one shows in a similar fashion to Corollary 8 that for $k \geq a + b$ odd we have $\bar{P}^{a,b}(\bar{A}_k^w | Z_k = \ell) \leq 2\bar{P}^{a,b}(\bar{A}_k^s | Z_k = \ell)$, where \bar{A}_k^s is the event on which the walk traverses the less (or equally) traversed edge at time k , and the remaining (stronger) edge from time $k + 2$ onwards, and where $\bar{A}_k^w = \{T = k + 2\} \setminus \bar{A}_k^s$. Putting this together with Lemma 6 one obtains

$$\bar{P}(T = k + 2) \asymp \sum_{\ell=1}^{k/2} \frac{1}{w(k - \ell)} \bar{W}_{k+2}(\ell + 2),$$

and

$$\bar{P}(T > n) \asymp \sum_{\substack{k=n+1, \\ k \text{ odd}}}^{\infty} \sum_{\ell=1}^{k/2} \frac{1}{w(k - \ell)} \bar{W}_{k+2}(\ell + 2).$$

2.4 Examples

Let $a, b \geq 1$ be fixed integers, and denote by P either of the laws $P^{a,b}$ or $\bar{P}^{a,b}$.

Theorem 10 *Suppose $\rho > 1$, let $\rho' := (\rho - 1)/\rho$, $\alpha \in (0, \infty)$ and let $\epsilon > 0$ be arbitrarily fixed.*

- (a) *If $w(k) = k^\rho$, $k \geq 1$ then there exist finite positive $c_1(\rho), c_2(\rho)$ such that for all $k \geq 1$*

$$\frac{c_1(\rho)}{k^{\rho-\rho'}} \leq P(T = k + 1) \leq \frac{c_2(\rho)}{k^{\rho-\rho'}}.$$

- (b) *If $w(k) = k^\rho \log^\alpha k$, $k \geq 1$, $\alpha > 1$ then there exist finite positive $c_1(\alpha, \rho), c_2(\alpha, \rho)$ such that for all $k \geq 1$*

$$\frac{c_1(\alpha, \rho)}{k^{\rho-\rho'} \log^\alpha k} \leq P(T = k + 1) \leq \frac{c_2(\alpha, \rho)}{k^{\rho-\rho'} \log^\alpha k}.$$

- (c) *If $w(k) = k e^{\log^\alpha k}$, $k \geq 1$, $0 < \alpha < 1$ then there exist finite positive $c_1(\alpha), c_2(\alpha)$ such that for all $k \geq 1$ and all $\epsilon > 0$*

$$\frac{c_1(\alpha)}{k(\log k)^{1-\alpha} e^{\log^{\alpha^2} k - \beta \log^{2\alpha^2 - \alpha} k}} \leq P(T = k + 1) \leq \frac{c_2(\alpha)}{k(\log k)^{1-\alpha} e^{\log^{\alpha^2} k - (\alpha + \epsilon) \log^{2\alpha^2 - \alpha} k}},$$

where $\beta = \alpha$ if $0 < \alpha \leq 2/3$ and $\beta = \alpha - \epsilon$ otherwise.

In particular, if $0 < \alpha \leq 1/2$ then $\exp\{-\log^{2\alpha^2-\alpha} k\} \asymp 1$, so

$$P(T = k + 1) \asymp \frac{c_2(\alpha)}{k(\log k)^{1-\alpha} e^{\log^{\alpha^2} k}},$$

(d) If $w(k) = e^{k^\epsilon}$, for some $0 < \epsilon \leq 1$, then there exist finite positive $c_1(\epsilon), c_2(\epsilon)$ such that for all $k \geq 1$

$$\frac{c_1(\epsilon)k^{1-\epsilon}}{e^{(k-k^*)^\epsilon}} \leq P(T = k + 1) \leq \frac{c_2(\epsilon)k^{1-\epsilon}}{e^{(k-k^*)^\epsilon}},$$

where $k^* = \frac{k}{2} - \frac{(1-\epsilon)}{\epsilon^{2^{2-\epsilon}}} k^{1-\epsilon} \log k$.

(e) If $w(k) = e^{\alpha k}$, then there exist finite positive $c_1(\alpha), c_2(\alpha)$ such that

$$c_1(\alpha)e^{-\alpha k/2} \leq P(T = k + 1) \leq c_2(\alpha)e^{-\alpha k/2}.$$

Remark All the constants of the form $c_1(\cdot), c_2(\cdot)$ featuring in the statements above depend in addition on the initial weights a and b due to Theorems 4 and 9, for an example see (26) below.

Proof. We concentrate on the case where $P = P^{a,b}$, the other case $P = \bar{P}^{a,b}$ can be done similarly. Without loss of generality, we assume $\ell \leq k/2$. We are going to use the following inequality

$$e^{-w(\ell+1) \sum_{j=k}^{\infty} \frac{1}{w(j-\ell)}} \leq W_{k+1}(\ell + 1) \leq e^{-w(\ell+1) \sum_{j=k}^{\infty} \frac{1}{2w(j-\ell)}}, \quad (23)$$

which is a direct consequence of the fact

$$e^{-x} \leq (1+x)^{-1} \leq e^{-x/2}, \quad 0 \leq x \leq 1.$$

(a) For $w(k) = k^\rho$, (23) becomes

$$e^{-(\ell+1)^\rho \sum_{j=k}^{\infty} \frac{1}{(j-\ell)^\rho}} \leq W_{k+1}(\ell + 1) \leq e^{-(\ell+1)^\rho \sum_{j=k}^{\infty} \frac{1}{2(j-\ell)^\rho}}.$$

Using

$$\int_{k+1}^{\infty} \frac{dx}{(x-\ell)^\rho} \leq \sum_{j=k}^{\infty} \frac{1}{(j-\ell)^\rho} \leq \int_k^{\infty} \frac{dx}{(x-\ell)^\rho},$$

we get a lower and an upper bound

$$\frac{1}{\rho-1} \frac{1}{(k+1-\ell)^{\rho-1}} \leq \sum_{j=k}^{\infty} \frac{1}{(j-\ell)^\rho} \leq \frac{1}{\rho-1} \frac{1}{(k-\ell)^{\rho-1}}. \quad (24)$$

Therefore we have for $1 \leq \ell \leq k/2$

$$e^{-\frac{2^{\rho-1}}{\rho-1} \frac{(\ell+1)^\rho}{k^{\rho-1}}} \leq e^{-\frac{1}{\rho-1} \frac{(\ell+1)^\rho}{(k-\ell)^{\rho-1}}} \leq W_{k+1}(\ell+1) \leq e^{-\frac{1}{2^{\rho-2}} \frac{(\ell+1)^\rho}{(k+1-\ell)^{\rho-1}}} \leq e^{-\frac{1}{2^{\rho-2}} \frac{\ell^\rho}{k^{\rho-1}}}. \quad (25)$$

Now Theorem 9 implies

$$\begin{aligned}
P(T = k + 1) &\geq c(a, b) \sum_{\ell=1}^{k^{\rho'}} \frac{1}{(k - \ell)^{\rho}} e^{-\frac{2^{\rho-1}}{\rho-1} \frac{(\ell+1)^{\rho}}{k^{\rho-1}}} \\
&\geq c(a, b) e^{-\frac{2^{\rho-1}}{\rho-1}} \sum_{\ell=1}^{k^{\rho'}} \frac{1}{(k - \ell)^{\rho}} \\
&\geq c(a, b) e^{-\frac{2^{\rho-1}}{\rho-1}} k^{\rho' - \rho}
\end{aligned} \tag{26}$$

$$\tag{27}$$

where $c(a, b)$ is a finite positive constant. For the upper bound again use Theorem 9 and (25). Now note that, since $k \geq 2\ell$,

$$\sum_{\ell=k^{\rho'}}^{k/2} \frac{1}{(k - \ell)^{\rho}} e^{-\frac{1}{2^{\rho-2}} \frac{\ell^{\rho}}{k^{\rho-1}}} \leq \left(\frac{2}{k}\right)^{\rho} \sum_{\ell=k^{\rho'}}^{k/2} e^{-\frac{1}{2^{\rho-2}} \frac{\ell^{\rho}}{k^{\rho-1}}}, \tag{28}$$

To bound the last term above, we split the interval $[k^{\rho'}, k/2]$ into subintervals of equal width $k^{\rho'}$, with the last subinterval possibly having smaller width $k/2 - \lfloor k^{1-\rho'}/2 \rfloor k^{\rho'}$. We abbreviate $a_k := \lfloor k^{1-\rho'}/2 \rfloor$. Then we have

$$\begin{aligned}
\sum_{\ell=k^{\rho'}}^{k/2} e^{-\frac{1}{2^{\rho-2}} \frac{\ell^{\rho}}{k^{\rho-1}}} &\leq k^{\rho'} e^{-\frac{1}{2^{\rho-2}}} + k^{\rho'} e^{-\frac{2^{\rho}}{2^{\rho-2}}} + \dots + k^{\rho'} e^{-\frac{a_k^{\rho}}{2^{\rho-2}}} \\
&\leq k^{\rho'} \left[e^{-\frac{1}{2^{\rho-2}}} + e^{-\frac{2}{2^{\rho-2}}} + \dots + e^{-\frac{a_k}{2^{\rho-2}}} \right] \leq \frac{k^{\rho'}}{1 - e^{-\frac{1}{2^{\rho-2}}}},
\end{aligned}$$

which together with (28) concludes the proof of part (a).

(b) Using (23), and the fact that

$$\sum_{j=k}^{\infty} \frac{1}{(j - \ell)^{\rho} \log^{\alpha}(j - \ell)}$$

is up to a constant multiple of order $1/((k - \ell)^{\rho-1} \log^{\alpha}(k - \ell))$ one obtains

$$e^{-c_1(\alpha, \rho) \frac{(\ell+1)^{\rho}}{(k-\ell)^{\rho-1}} \frac{\log^{\alpha}(\ell+1)}{\log^{\alpha}(k-\ell)}} \leq W_{k+1}(\ell + 1) \leq e^{-c_2(\alpha, \rho) \frac{(\ell+1)^{\rho}}{k^{\rho-1}} \frac{\log^{\alpha}(\ell+1)}{\log^{\alpha} k}}, \tag{29}$$

where $c_1(\alpha, \rho), c_2(\alpha, \rho)$ are finite positive constants. As in the case (a), one gets the lower bound by evaluating the order or

$$\sum_{\ell=\xi}^{\zeta} \frac{1}{w(k - \ell)} W_{k+1}(\ell + 1), \tag{30}$$

for $\xi = 1, \zeta = k^{\rho'}$, and the upper bound by evaluating the order of the sum for $\xi = 1, \zeta = k^{\rho'}$ and for $\xi = k^{\rho'}, \zeta = k/2$, separately.

(c) We have

$$\int_{k+1}^{\infty} \frac{dx}{(x-\ell)e^{\log^{\alpha}(x-\ell)}} \leq \sum_{j=k}^{\infty} \frac{1}{(j-\ell)e^{\log^{\alpha}(j-\ell)}} \leq \int_k^{\infty} \frac{dx}{(x-\ell)e^{\log^{\alpha}(x-\ell)}}.$$

Since

$$\begin{aligned} \int_k^{\infty} \frac{dx}{(x-\ell)e^{\log^{\alpha}(x-\ell)}} &= \frac{\log^{1-\alpha}(k-\ell)}{\alpha e^{\log^{\alpha}(k-\ell)}} + \frac{1-\alpha}{\alpha} \int_k^{\infty} \frac{dx}{(x-\ell) \log^{\alpha}(x-\ell) e^{\log^{\alpha}(x-\ell)}} \\ &= \frac{\log^{1-\alpha}(k-\ell)}{\alpha e^{\log^{\alpha}(k-\ell)}} (1 + o_{k-\ell}(1)), \end{aligned}$$

where $o_k(1) \rightarrow 0$ as $k \rightarrow \infty$, using the fact that $1 \leq \ell \leq \frac{k}{2}$ above, we get the inequality

$$c_1(\alpha) \frac{(\log k)^{1-\alpha}}{e^{\log^{\alpha} k}} \leq \sum_{j=k}^{\infty} \frac{1}{(j-\ell)e^{\log^{\alpha}(j-\ell)}} \leq c_2(\alpha) \frac{(\log k)^{1-\alpha}}{e^{\log^{\alpha} k}},$$

for some $c_1(\alpha), c_2(\alpha) \in (0, \infty)$. By applying now (23), we obtain

$$e^{-c_1(\alpha)(\ell+1)e^{\log^{\alpha}(\ell+1)} \frac{\log^{1-\alpha} k}{e^{\log^{\alpha} k}}} \leq W_{k+1}(\ell+1) \leq e^{-c_2(\alpha)(\ell+1)e^{\log^{\alpha}(\ell+1)} \frac{\log^{1-\alpha} k}{e^{\log^{\alpha} k}}}.$$

As in parts (a) and (b), we find a convenient breaking point and approximate the sums (30) separately. We take

$$\zeta = e^{\log^{\alpha} k - \log^{\alpha^2} k + \beta \log^{2\alpha^2 - \alpha} k} / \log^{1-\alpha} k,$$

where $\beta = \alpha$ for $0 < \alpha \leq 2/3$ and $\beta = \alpha - \epsilon$ otherwise.

To verify the lower bound, we need to show that we can bound

$$\exp \left\{ -c_1(\alpha)(\zeta+1)e^{\log^{\alpha}(\zeta+1)} \frac{\log^{1-\alpha} k}{e^{\log^{\alpha} k}} \right\}$$

from below by a positive constant. Hence we estimate (the constant c below is finite and positive, and possibly changes from line to line)

$$\begin{aligned} &\frac{(\zeta+1) \log^{1-\alpha} k}{\exp(\log^{\alpha} k)} \exp(\log^{\alpha}(\zeta+1)) \\ &\leq c \exp[\beta \log^{2\alpha^2 - \alpha} k - \log^{\alpha^2} k] \end{aligned} \tag{31}$$

$$\begin{aligned} &\times \exp[(\log^{\alpha} k - \log^{\alpha^2} k + \beta \log^{2\alpha^2 - \alpha} k - (1-\alpha) \log \log k)^{\alpha}] \\ &\leq c \exp[\beta \log^{2\alpha^2 - \alpha} k - \log^{\alpha^2} k + \log^{\alpha^2} k (1 - \log^{\alpha^2 - \alpha} k + \beta \log^{2\alpha^2 - 2\alpha} k)^{\alpha}] \\ &\leq c \exp[\beta \log^{2\alpha^2 - \alpha} k - \log^{\alpha^2} k + \log^{\alpha^2} k (1 - \alpha \log^{\alpha^2 - \alpha} k + \beta \alpha \log^{2\alpha^2 - 2\alpha} k)] \\ &= c \exp(\beta \log^{2\alpha^2 - \alpha} k - \alpha \log^{2\alpha^2 - \alpha} k + \beta \alpha \log^{3\alpha^2 - 2\alpha} k), \end{aligned} \tag{32}$$

where β is chosen as above, and where for the third inequality we use that $(1-x)^{\epsilon} \leq 1 - \epsilon x$ for $0 \leq x < 1$ and $0 \leq \epsilon \leq 1$. Note now that

if $\alpha \leq 2/3$, then $\log^{3\alpha^2-2\alpha} k$ is bounded from above by a constant, so $\beta = \alpha$ is sufficient to bound from above by a constant the expression (32). If $\alpha > 2/3$, then $\log^{3\alpha^2-2\alpha} k \rightarrow \infty$, and $3\alpha^2 - 2\alpha < 2\alpha^2 - \alpha$, so taking $\beta = \alpha - \epsilon$ for any $\epsilon > 0$ will suffice to make (32) bounded by a constant. By Lemma 7 (a), $W_{k+1}(\ell + 1) \geq W_{k+1}(\zeta + 1) \geq e^{-c'}$, for $\ell \leq \zeta$, so that

$$\sum_{\ell=1}^{\zeta} \frac{1}{w(k-\ell)} W_{k+1}(\ell + 1) \geq e^{-c'} \frac{\zeta}{w(k)} \geq \frac{c_1(\alpha)}{k \log^{1-\alpha} k e^{\log^{\alpha^2} k - \beta \log^{2\alpha^2 - \alpha} k}} \quad (33)$$

which proves the lower bound.

For the upper bound, break the summation in turn at the point

$$\zeta := e^{\log^{\alpha} k - \log^{\alpha^2} k + (\alpha + \epsilon) \log^{2\alpha^2 - \alpha} k} / \log^{1-\alpha} k,$$

We have, since $W_{k+1}(\ell + 1) \leq 1$,

$$\sum_{\ell=1}^{\zeta} \frac{1}{w(k-\ell)} W_{k+1}(\ell + 1) \leq \sum_{\ell=1}^{\zeta} \frac{1}{w(k-\ell)} \leq \frac{\zeta}{w(k-\zeta)},$$

and since $w(k-\zeta) \asymp w(k)$ we can bound from above this term by the term of the order stated in the formulation of part (c). Hence, it suffices to bound $\sum_{\ell=\zeta}^{k/2} W_{k+1}(\ell + 1)/w(k-\ell)$ as follows.

$$\begin{aligned} \sum_{\ell=\zeta}^{k/2} \frac{1}{w(k-\ell)} W_{k+1}(\ell + 1) &\leq \frac{1}{w(k/2)} \sum_{\ell=\zeta}^{k/2} W_{k+1}(\ell + 1) \\ &\leq \frac{c_2}{k e^{\log^{\alpha} k}} \int_{\zeta}^{\infty} \frac{dx}{e^{c_2(\alpha) x e^{\log^{\alpha} x \frac{\log^{1-\alpha} k}{e^{\log^{\alpha} k}}}}} \\ &= \frac{c_2}{k e^{\log^{\alpha} k}} \cdot \frac{1}{\log^{1-\alpha} k (1 + \alpha \log^{\alpha-1}(\zeta)) e^{\log^{\alpha}(\zeta)}} \\ &\quad \cdot \frac{1}{e^{c_2(\alpha) \cdot \zeta \cdot e^{\log^{\alpha}(\zeta) \frac{\log^{1-\alpha} k}{e^{\log^{\alpha} k}}}}} (1 + o_{\zeta}(1)), \end{aligned} \quad (34)$$

where the last term is obtained by integration by parts.

First estimate

$$\begin{aligned} &\frac{\zeta \cdot \log^{1-\alpha} k}{\exp(\log^{\alpha} k)} \exp(\log^{\alpha}(\zeta)) \\ &= \exp[(\alpha + \epsilon) \log^{2\alpha^2 - \alpha} k - \log^{\alpha^2} k] \\ &\quad \times \exp[(\log^{\alpha} k - \log^{\alpha^2} k + (\alpha + \epsilon) \log^{2\alpha^2 - \alpha} k - (1 - \alpha) \log \log k)^{\alpha}] \\ &\geq c \exp[(\alpha + \epsilon) \log^{2\alpha^2 - \alpha} k - \log^{\alpha^2} k + \log^{\alpha^2} k (1 - \log^{\alpha^2 - \alpha} k)^{\alpha}] \\ &\geq c \exp[(\alpha + \epsilon) \log^{2\alpha^2 - \alpha} k - \log^{\alpha^2} k + \log^{\alpha^2} k (1 - (\alpha + \epsilon) \log^{\alpha^2 - \alpha} k)] \geq c' \end{aligned}$$

where in the last inequality above we used

$$(1-x)^\alpha \geq 1 - (\alpha + \epsilon)x, \quad \epsilon > 0, \quad x \rightarrow 0. \quad (35)$$

Therefore we can bound the multiple (34) from above by a constant c . Furthermore,

$$\begin{aligned} \frac{c_2}{k e^{\log^\alpha k}} &\cdot \frac{e^{\log^\alpha k}}{\log^{1-\alpha} k (1 + \alpha \log^{\alpha-1}(\zeta)) e^{\log^\alpha(\zeta)}} \\ &\leq \frac{c_2 \exp\{-(\log^\alpha k - \log^{\alpha^2} k + (\alpha + \epsilon) \log^{2\alpha^2-\alpha} k - (1-\alpha) \log \log k)^\alpha\}}{k (\log k)^{1-\alpha}} \\ &\leq \frac{c_2 \exp\{-(\log^\alpha k - \log^{\alpha^2} k)^\alpha\}}{k (\log k)^{1-\alpha}} \\ &\leq \frac{c_2 \exp\{-\log^{\alpha^2} k (1 - (\alpha + \epsilon) \log^{\alpha^2-\alpha} k)\}}{k (\log k)^{1-\alpha}}, \end{aligned}$$

where in the last inequality above we again used (35).

(d) We have

$$\int_{k-\ell+1}^{\infty} \frac{dx}{e^{x^\epsilon}} \leq \sum_{j=k-\ell}^{\infty} \frac{1}{e^{j^\epsilon}} \leq \int_{k-\ell}^{\infty} \frac{dx}{e^{x^\epsilon}}.$$

Note that

$$\int_{k-\ell}^{\infty} \frac{dx}{e^{x^\epsilon}} = \frac{(k-\ell)^{1-\epsilon}}{\epsilon e^{(k-\ell)^\epsilon}} + \frac{1-\epsilon}{\epsilon} \int_{k-\ell}^{\infty} \frac{dx}{x^\epsilon e^{x^\epsilon}} = \frac{(k-\ell)^{1-\epsilon}}{\epsilon e^{(k-\ell)^\epsilon}} (1 + o_{k-\ell}(1)),$$

where again $o_k(1) \rightarrow 0$ as $k \rightarrow \infty$. From the above formulas we get the inequality

$$c_2(\epsilon) \frac{(k-\ell)^{1-\epsilon}}{e^{(k-\ell)^\epsilon}} \leq \sum_{j=k-\ell}^{\infty} \frac{1}{e^{j^\epsilon}} \leq c_1(\epsilon) \frac{(k-\ell)^{1-\epsilon}}{e^{(k-\ell)^\epsilon}}, \quad (36)$$

for some $c_1(\epsilon), c_2(\epsilon) \in (0, \infty)$. By applying now (23), we obtain

$$e^{-c_1(\epsilon) e^{\ell^\epsilon} \frac{(k-\ell)^{1-\epsilon}}{e^{(k-\ell)^\epsilon}}} \leq W_{k+1}(\ell+1) \leq e^{-c_2(\epsilon) e^{\ell^\epsilon} \frac{(k-\ell)^{1-\epsilon}}{e^{(k-\ell)^\epsilon}}}. \quad (37)$$

Now take $k^* = \frac{k}{2} - \frac{(1-\epsilon)}{2\epsilon 2^{1-\epsilon}} k^{1-\epsilon} \log k$, then

$$W_{k+1}(k^*+1) \geq e^{-c_1(\epsilon) e^{k^{*\epsilon}} \frac{(k-k^*)^{1-\epsilon}}{e^{(k-k^*)^\epsilon}}}.$$

Note that

$$\begin{aligned} k^{*\epsilon} - (k-k^*)^\epsilon &= (k-k^*)^\epsilon \left[\left(1 - \frac{(1-\epsilon) k^{1-\epsilon} \log k}{\epsilon 2^{1-\epsilon} (k-k^*)} \right)^\epsilon - 1 \right] \\ &\leq -(1-\epsilon) \log k + O(k^{-\epsilon/2}), \end{aligned}$$

where the last inequality is obtained using $(1-x)^\epsilon \leq 1-\epsilon x$ for $0 \leq x < 1$ and $0 \leq \epsilon \leq 1$, together with $k - k^* \geq k/2(1 + O(\log k/k^\epsilon))$. Therefore $\liminf_k W_{k+1}(k^* + 1) > 0$, and by Lemma 7 (a)

$$W_{k+1}(\ell + 1) \geq W_{k+1}(k^* + 1) \geq c(\epsilon), \quad \ell \leq k^* \text{ for some } c(\epsilon) > 0.$$

Therefore, recalling $c_1(\epsilon), c_2(\epsilon)$ from (36),

$$\begin{aligned} \sum_{\ell=1}^{k^*} \frac{1}{w(k-\ell)} W_{k+1}(\ell + 1) &\geq c(\epsilon) \sum_{\ell=1}^{k^*} \frac{1}{e^{(k-\ell)\epsilon}} \geq c(\epsilon) \int_{k-k^*+1}^{k+1} \frac{dx}{e^{x\epsilon}} \\ &\geq c(\epsilon) \left(\frac{c_1(\epsilon)(k-k^*)^{1-\epsilon}}{e^{(k-k^*)\epsilon}} - \frac{c_2(\epsilon)k^{1-\epsilon}}{e^{k\epsilon}} \right) \\ &\geq \frac{c(\epsilon)c_1(\epsilon)(k-k^*)^{1-\epsilon}}{e^{(k-k^*)\epsilon}}, \end{aligned}$$

where constant $c(\epsilon)$ may change from line to line by a positive finite multiple. This proves the lower bound.

To get the corresponding upper bound, first observe that $W_{k+1}(\ell + 1) \leq 1$, so using (36) we can simply bound

$$\sum_{\ell=1}^{k^*} \frac{1}{w(k-\ell)} W_{k+1}(\ell + 1) \leq \sum_{j=k-k^*}^{\infty} \frac{1}{w(j)} \leq c_2(\epsilon) \frac{(k-k^*)^{1-\epsilon}}{e^{(k-k^*)^{1-\epsilon}}},$$

and we proceed to bound $\sum_{\ell=k^*}^{k/2} \frac{1}{w(k-\ell)} W_{k+1}(\ell + 1)$.

Due to (36) and (37) we can write, for $\ell \in [k^*, k/2]$,

$$\exp \left\{ -c'_1(\epsilon) w(\ell) \int_{k-\ell}^{\infty} \frac{dx}{e^{x\epsilon}} \right\} \leq W_{k+1}(\ell + 1) \leq \exp \left\{ -c'_2(\epsilon) w(\ell) \int_{k-\ell}^{\infty} \frac{dx}{e^{x\epsilon}} \right\}, \quad (38)$$

for some $c'_1(\epsilon), c'_2(\epsilon) \in (0, \infty)$. Since $W_{k+1}(k^* + 1) \asymp 1$, as we showed in the proof of the lower bound, we have

$$\frac{1}{w(k^*)} \asymp \int_{k-k^*}^{\infty} \frac{dx}{e^{x\epsilon}},$$

and since

$$\int_{k-\ell}^{\infty} \frac{dx}{e^{x\epsilon}} \geq \int_{k-k^*}^{\infty} \frac{dx}{e^{x\epsilon}}, \quad \ell \geq k^*,$$

we conclude from (38) that

$$W_{k+1}(\ell + 1) \leq \exp\{-c w(\ell)/w(k^*)\}, \quad \ell \in [k^*, k/2].$$

Therefore,

$$\sum_{\ell=k^*}^{k/2} \frac{1}{w(k-\ell)} W_{k+1}(\ell + 1) \leq \sum_{\ell=k^*}^{k/2} \frac{e^{-c \frac{w(\ell)}{w(k^*)}}}{w(k-\ell)} \leq \sum_{\ell=k^*}^{k/2} \frac{\bar{c} w^2(k^*)}{w^2(\ell) w(k-\ell)},$$

where $\bar{c} \in (0, \infty)$ is such that $e^{-cx} \leq \bar{c}x^{-2}$ for all $x \geq 1$, and where we use the fact $w(\ell) \geq w(k^*)$, for $\ell \geq k^*$. Finally, it is easy to check that $\ell \mapsto w(k - \ell)w(\ell)$, $\ell \geq k^*$, is a non-decreasing function so

$$\frac{w(k^*)}{w(\ell)w(k - \ell)} \leq \frac{1}{w(k - k^*)},$$

and therefore

$$\begin{aligned} \sum_{\ell=k^*}^{k/2} \frac{w^2(k^*)}{w^2(\ell)w(k - \ell)} &\leq \frac{w(k^*)}{w(k - k^*)} \sum_{\ell=k^*}^{k/2} \frac{1}{w(\ell)} \leq \frac{w(k^*)}{w(k - k^*)} \int_{k^*-1}^{k/2} \frac{dx}{w(x)} \\ &\leq c_2(\epsilon) \frac{(k^*)^{1-\epsilon}}{w(k - k^*)} \end{aligned}$$

which gives the upper bound due to $k^* < k/2 < k - k^*$.

- (e) A direct consequence of part (d), but its direct proof (left to an interested reader) is much easier, this fact is related to the following property: among all the weights in (d), it is only the case of $w(k) = e^k$ where the edge reinforced random walk gets attracted at any particular time with probability uniformly bounded away from zero.

□

As a consequence of Theorem 10, we now have

Corollary 11 *Suppose that $w(k)$ is as in (a) or (b) in Theorem 10. Then $E(T)$ is infinite if $\rho \leq 1 + \frac{1+\sqrt{5}}{2}$ and finite if $\rho > 1 + \frac{1+\sqrt{5}}{2}$.*

3 Analysis on general graphs

Assume that \mathcal{G} is a connected graph with $D(\mathcal{G}) < \infty$. Recall that $P^{\mathcal{G}}$ is the law of the reinforced random walk on \mathcal{G} .

We start with an easy lower bound in terms of the tail distribution of T under two-edge law \bar{P} . In fact, in the following comparison arguments, it will be convenient to consider instead the law of

$$(T)^+ \equiv T^+ := T - t_0.$$

Unless otherwise stated, in this section we will assume that $\ell_0^e, e \in E(\mathcal{G})$ forms the (general) initial configuration of weights on edges, such that $\ell_0^e < \infty$, $e \in E(\mathcal{G})$.

Lemma 12 *There exists $c = c(w, \mathcal{D}(\mathcal{G})) \in (0, \infty)$, and $a, b \in \mathbb{N}$ such that*

$$P^{\mathcal{G}}(T^+ > k) \geq c\bar{P}^{a,b}(T^+ > k).$$

Proof. Let $I_{t_0} = v \in \mathcal{G}$ be the initial position. Without loss of generality, assume that at least two edges e and f meet at v . Otherwise, at least two edges must meet at the unique neighbor of v , and the argument is similar. Recall that \mathcal{G}_1 denotes the range of the walk. Define event

$$A_{e,f} := \{\mathcal{G}_1 \subset \text{graph spanned by } e, f\},$$

and note that, due to (A0), event $A_{e,f}$ has positive probability for any given bounded degree graph and any fixed configuration $\ell_0^e, e \in E(\mathcal{G})$. At the same time

$$\{T^+ > k\} \cap A_{e,f} \subset \{T^+ > k\}.$$

Denote by v_e and v_f the two vertices such that $e = \{v, v_e\}$ and $f = \{v, v_f\}$. We will verify below the existence of a positive constant β that depends on \mathcal{G}, w and the initial weights, such that for each (possibly infinite) path $v = i_0 \sim i_1 \sim \dots$ of vertices where $i_n \in \{v, v_e, v_f\}$, $n \geq 0$, we have

$$P^{\mathcal{G}}(I_{t_0} = i_0, I_{t_0+1} = i_1, \dots) \geq \beta \bar{P}^{\ell_0^e, \ell_0^f}(I_{t_0} = i_0, I_{t_0+1} = i_1, \dots). \quad (39)$$

Note that t_0 equals ℓ_0^e, ℓ_0^f under the law $\bar{P}^{\ell_0^e, \ell_0^f}$, but as commented earlier the edge reinforced random walk can be redefined by time-shift to start from any fixed initial time, and this does not change the probability of it taking any particular path. Clearly (39) implies that $P^{\mathcal{G}}(B \cap A_{e,f}) \geq \beta \bar{P}^{\ell_0^e, \ell_0^f}(B)$, for any event B in the σ -field generated by the walk. In particular,

$$P^{\mathcal{G}}(T^+ > k) \geq P^{\mathcal{G}}(\{T^+ > k\} \cap A_{e,f}) \geq \beta \bar{P}^{a,b}(T^+ > k),$$

as claimed.

It suffices to verify (39) for each infinite path $i_0 \sim i_1 \sim \dots$ specified above. For $n \geq t_0$, define $x_n^e := \ell_0^e + \#\{j \leq n : \{i_{j-1}, i_j\} = e\}$ and $x_n^f := \ell_0^f + \#\{j \leq n : \{i_{j-1}, i_j\} = f\}$. The probability on the right-hand side of (39) equals

$$\prod_{n=0}^{\infty} \frac{w(x_{t_0+2n}^{\{i_{2n}, i_{2n+1}\}})}{w(x_{t_0+2n}^e) + w(x_{t_0+2n}^f)}. \quad (40)$$

Define

$$c(v) := \sum_{u: u \sim v, u \neq v_e, v_f} w(\ell_0^{\{u, v\}}),$$

$$c(v_e) := \sum_{u: u \sim v_e, u \neq v} w(\ell_0^{\{u, v_e\}}), \quad c(v_f) := \sum_{u: u \sim v_f, u \neq v} w(\ell_0^{\{u, v_f\}}).$$

The probability on the left-hand side of (39) equals

$$\prod_{n=0}^{\infty} \frac{w(x_{t_0+2n}^{\{i_{2n}, i_{2n+1}\}})}{w(x_{t_0+2n}^e) + w(x_{t_0+2n}^f) + c(v)} \prod_{n=0}^{\infty} \frac{w(x_{t_0+2n+1}^{\{i_{2n+1}, i_{2n+2}\}})}{w(x_{t_0+2n+1}^{\{i_{2n+1}, i_{2n+2}\}}) + c(v_{\{i_{2n+1}, i_{2n+2}\}})}, \quad (41)$$

where the first infinite product accounts for all the steps originating from the middle vertex v , while the second infinite product accounts for all the steps originating from the “boundary vertices” v_e and v_f . Since

$$\sum_{n=0}^{\infty} \frac{c(v_{\{i_{2n+1}, i_{2n+2}\}})}{w(x_{t_0+2n+1}^{\{i_{2n+1}, i_{2n+2}\}}) + c(v_{\{i_{2n+1}, i_{2n+2}\}})} \leq 2 \sum_{n=0}^{\infty} \frac{c(v_e) \vee c(v_f)}{w(n) + (c(v_e) \wedge c(v_f))} < \infty$$

by (A0), a well-known calculus fact implies that the second product is uniformly (over infinite paths) bounded away from 0. The ratio of the first product in (40) and the probability in (42) is again uniformly bounded away from 0 since

$$\sum_{n=0}^{\infty} \frac{c(v)}{w(x_{t_0+2n}^e) + w(x_{t_0+2n}^f) + c(v)} \leq 2 \sum_{n=0}^{\infty} \frac{c(v)}{w(n) + c(v)} < \infty. \quad (42)$$

□

Getting a corresponding upper bound on the tails of distribution of T seems more difficult. As a warm-up we study the tree setting next, and the general finite graph and the infinite graph settings respectively in following subsections.

The following fact, complementary in spirit to conditioning on event $A_{e,f}$ in the proof of Lemma 12, will prove useful soon.

Lemma 13 *Suppose \mathcal{G}^* is a finite connected subgraph of \mathcal{G} . Then for each (possibly infinite) path $i_0 \sim i_1 \sim \dots$ of vertices all contained in \mathcal{G}^* we have, assuming $P^{\mathcal{G}}(I_{t_0} = i_0) = P^{\mathcal{G}^*}(I_{t_0} = i_0) = 1$,*

$$P^{\mathcal{G}}(I_{t_0} = i_0, I_{t_0+1} = i_1, \dots) \leq P^{\mathcal{G}^*}(I_{t_0} = i_0, I_{t_0+1} = i_1, \dots).$$

Proof. At each step k where all the neighbors of the current position i_k are contained in \mathcal{G}^* the probability of the transition from i_k to i_{k+1} is the same under both laws $P^{\mathcal{G}}$ and $P^{\mathcal{G}^*}$.

At each step k where at least one neighbor of the current position i_k is an element of $V(\mathcal{G}) \setminus V(\mathcal{G}^*)$, note that the probability of the transition from i_k to i_{k+1} under $P^{\mathcal{G}}$ is strictly smaller than that under $P^{\mathcal{G}^*}$. □

3.1 Analysis on trees

In this subsection we assume that \mathcal{G} is a tree such that $D(\mathcal{G}) < \infty$, and we derive some upper bound estimates on the tail distribution of T under $P^{\mathcal{G}}$.

First let \mathcal{G} be the star with m fingers as defined in Section 2.2. For concreteness, we assume that all the initial weights $\ell_0^{e_i}$ are equal to 1, and that $I_{t_0} = I_m = 0$. A similar statement applies for more general initial configurations.

Lemma 14 $P^{\mathcal{G}}(T^+ > k) \leq \binom{m}{2} \bar{P}^{1,1}(T^+ > k/\binom{m}{2})$.

Proof. We will show that

$$T^+ \leq \sum_{1 \leq i < j \leq m} T^{e_i, e_j, restr, +}, \text{ almost surely,} \quad (43)$$

where $T^{e_i, e_j, restr} = T^{e_j, e_i, restr}$ is a random variable to be defined, corresponding to the pair of edges e_i, e_j such that its law under $P^{\mathcal{G}}$ is the law of T under $\bar{P}^{1,1}$, and where $T^{e_i, e_j, restr, +} = (T^{e_i, e_j, restr})^+$. The reason for (43) is as follows. Suppose that $f \in \{e_1, \dots, e_m\}$ is the attracting edge for the walk. The steps away from the central vertex 0 up to time T are naturally split into K_e many steps traversing edge $e \neq f$ (so in total there are $2K_e$ many steps along any $e \neq f$). Up to time T , there are therefore $T - 2\sum_{e \neq f} K_e$ steps across f . For $e \neq f$, define

$$Y^{e,f} := \text{time of the last traversal of } e,$$

and

$$\begin{aligned} T_n^{e,f} &:= \ell_0^e + \ell_0^f + \#\text{traversals of } e \text{ or } f \text{ up to time } n, \\ T_n^{e,f,+} &:= T_n^{e,f} - (\ell_0^e + \ell_0^f). \end{aligned}$$

Recall the time-line construction of Section 2.2, using m independent “time-lines” (one corresponding to each edge).

Now fix arbitrary edges e and g . By ignoring all the time-lines except the ones corresponding to edges e and g one obtains the construction of the reinforced random walk under the law $\bar{P}^{1,1}$. Call this process the *restriction to edges e and g* . Define

$$\begin{aligned} T^{e,g, restr} &:= \text{time of attraction for the restriction to } e \text{ and } g, \text{ and} \\ T^{e,g, restr, +} &:= T^{e,g, restr} - (\ell_0^e + \ell_0^g). \end{aligned}$$

In particular, $T^{e,g, restr, +}$ under $P^{\mathcal{G}}$ has the law of T^+ under $\bar{P}^{1,1}$. Next observe that in the case where $g = f$ is the attracting edge, we have

$$T^{e,f, restr} \equiv T_{Y^{e,f}}^{e,f} + 1 \text{ and } T^{e,f, restr, +} \equiv T_{Y^{e,f}}^{e,f,+} + 1, \quad (44)$$

where the extra 1 on the right-hand side accounts for the traversal of edge f at the attraction time $T^{e,f, restr}$. In addition, note that for each $e \neq f$ the number $2K_e$ of steps traversing e before time T equals the number of steps traversing e before time $T_{Y^{e,f}}^{e,f}$. By similar reasoning, the number of steps traversing f strictly before time T equals the number of steps traversing f before time $T_{Y^{g,f}}^{g,f}$, for at least one $g \neq f$ (precisely, g is the edge traversed at time $T - 1$). The last two claims imply

$$T^+ \leq \sum_{e: e \neq f} T_{Y^{e,f}}^{e,f,+} + 1, \quad (45)$$

where again the extra 1 accounts for the traversal of f at time T . By (44),

$$\sum_{e:e \neq f} T_{Y_{e,f}}^{e,f,+} + 1 = \sum_{e:e \neq f} (T^{e,f, \text{restr},+} - 1) + 1 \leq \frac{1}{2} \sum_{e,g:e \neq g} T^{e,g, \text{restr},+},$$

so (45) implies

$$T^+ \leq \frac{1}{2} \sum_{e,g:e \neq g} T^{e,g, \text{restr},+}, \quad (46)$$

in particular yielding (43). As noted already, the $\binom{m}{2}$ different random variables $T^{e,g, \text{restr},+}$ are (identically) distributed under the law $P^{\mathcal{G}}$ as T^+ is under the law $\bar{P}^{1,1}$. The statement of the lemma is now a standard consequence of (43). \square

Now consider \mathcal{G} a finite tree. Let $m(\mathcal{G})$ be the total number of pairs of edges in \mathcal{G} that meet at a vertex. For example, the star with m fingers has $m(\mathcal{G}) = \binom{m}{2}$. Using the same reasoning (for each v , consider separately the star created by restricting the tree to v and all u , $u \sim v$) as in Lemma 14 one quickly obtains

Lemma 15 *Assume $\ell_0^e = 1$, $e \in E(\mathcal{G})$. Then*

$$P^{\mathcal{G}}(T^+ > k) \leq m(\mathcal{G})(\bar{P}^{1,1}(T > k/m(\mathcal{G})) + \bar{P}^{1,2}(T > k/m(\mathcal{G}))).$$

Here $\bar{P}^{1,2}(T > k/m(\mathcal{G}))$ appears due to parity considerations. Namely, for any two edges e, f that meet at a vertex v say, at the first time the walk visits v the configuration of weights is either 1, 1 or 1, 2 or 2, 1.

We will soon show analogous results for the walk on a general finite graph. Before this we quickly turn to the case where \mathcal{G} is an infinite tree of bounded degree. Recall that $\#\mathcal{G}_1$ denotes the total number of vertices ever visited by the edge reinforced random walk on \mathcal{G} . Here again we assume that $\ell_0^e = 1$, $e \in E(\mathcal{G})$. The next lemma can be proved in an analogous (but simpler) way to Lemma 25, we leave its verification to an interested reader.

Lemma 16 *There exists $p > 0$, depending on the weight w and the degree $D(\mathcal{G})$ of \mathcal{G} only, such that $\#\mathcal{G}_1/2$ is stochastically bounded by Z where Z is a geometric random variable with success probability p .*

Corollary 17 *Let \mathcal{G} be an infinite tree, such that $D(\mathcal{G}) < \infty$. Then for any $c > 1$ we have*

$$P^{\mathcal{G}}(T^+ > k) \leq O\left(\frac{1}{k^{c \log(1/(1-p))/2}}\right) + \max_{\mathcal{G}_{k,c}} P^{\mathcal{G}_{k,c}}(T^+ > k)$$

where the above maximum is taken over all trees $\mathcal{G}_{k,c}$ having fewer than $cD(\mathcal{G}) \log k$ vertices and degree bounded by $D(\mathcal{G})$.

Proof. Due to the last lemma, with probability $(1-p)^{(c \log k)/2}$ the range \mathcal{G}_1 of the walk is a subtree of \mathcal{G} containing initial position I_{t_0} and $c \log k$ or more vertices. On the opposite event, denoted by $B_{c \log k}$, we have $\mathcal{G}_1 \subset \mathcal{G}_{c \log k}^*$, where $\mathcal{G}_{c \log k}^*$ is a (non-random) subtree of \mathcal{G} generated by all vertices v of \mathcal{G} such that the graph distance of v and I_{t_0} is smaller or equal to $c \log k$. Therefore, by Lemma 13, we can bound

$$P^{\mathcal{G}}(\{T^+ > k\} \cap B_{c \log k}) \leq P^{\mathcal{G}}(\{T^+ > k\} \cap \{\mathcal{G}_1 \subset \mathcal{G}_{c \log k}^*\}) \leq P^{\mathcal{G}_{c \log k}^*}(T^+ > k).$$

□

Corollary 18 *Let \mathcal{G} be an infinite tree of bounded degree, and let $w(\cdot)$ be as in the examples of Theorem 10 (a)-(b). Then*

$$P^{\mathcal{G}}(T > k) = P^{1,1}(T > k) O(\log^q k),$$

for any $q > \rho - \rho'$.

Proof. For any tree $\mathcal{G}_{k,c}$ of bounded degree with fewer than $O(\log k)$ vertices one also has $m(\mathcal{G}_{k,c}) = O(\log k)$. Use previous corollary with $c \log(1/(1-p))/2 > \rho - \rho' - 1$. Finally note that under the assumptions of Theorem 10 (a) (resp. (b)) $m(\mathcal{G}_{k,c}) P^{1,1}(T^+ > k/m(\mathcal{G}_{k,c}))$ is of order $\log k \cdot P^{1,1}(T^+ > k)(\log k)^{\rho-\rho'-1}$ (resp. $\log k \cdot P^{1,1}(T^+ > k)(\log k)^{\rho-\rho'-1}(\log \log k)^\alpha$). □

3.2 Analysis on finite graphs

Assume that (A0) holds. Let \mathcal{G} be a finite graph, and abbreviate $\bar{n} = |E(\mathcal{G})|$. Moreover, denote the edges of \mathcal{G} by $E(\mathcal{G}) = \{e_1, e_2, \dots, e_{\bar{n}}\}$. If v is an arbitrary vertex of the graph, let $n_v = \text{degree}(v)$, and let $\mathcal{N}_v := \{e_1^v, e_2^v, \dots, e_{n_v}^v\}$ be the set of edges incident to v . Recall that X_k^e equals the initial weight ℓ_0^e incremented by the number of times edge e has been visited by time k .

As before, we will start the walk at time $\sum_{e \in E(\mathcal{G})} \ell_0^e$. Fix the initial position I_{t_0} at some arbitrary vertex v_0 . Then the following proposition holds:

Proposition 19 *Let $k \geq \sum_{e \in E(\mathcal{G})} \ell_0^e$ and $v \in V(\mathcal{G})$ and denote by $A_{v,k}$ be the event $\{I_k = v\}$. Then for any ℓ^e , $e \in E(\mathcal{G})$ such that $\ell^e \geq \ell_0^e$, $e \in E(\mathcal{G})$ and $\sum_{e \in E(\mathcal{G})} \ell^e = k$, we have*

$$P^{\mathcal{G}}(X_k^e = \ell^e, e \in E(\mathcal{G}), A_{v,k}) \leq \frac{\prod_{e \in E(\mathcal{G})} w(\ell_0^e)}{\min_{e \in \mathcal{N}_{v_0}} w(\ell_0^e)} \cdot \frac{\sum_{e \in \mathcal{N}_v} w(\ell^e)}{\prod_{e \in E(\mathcal{G})} w(\ell^e)}. \quad (47)$$

Remark Inequality (47) holds trivially when the conditions of the propositions do not hold, since the left-hand side then equals 0.

Proof. As in the two-edge case, we will use induction on $\sum_e \ell^e = k$ to prove the above inequality. The base of induction at the initial time $\sum_e \ell_0^e$ clearly

holds, since when the left-hand-side is 0 the right-hand-side is positive, and when the left-hand-side is 1 the right-hand-side is greater than 1.

Now take $k > \sum_e \ell_0^e$ and consider the event on the left-hand-side. For each $i = 1, 2, \dots, n_v$, let $v_i \in V(\mathcal{G})$ be the neighbor of v such that $e_i^v = \{v, v_i\}$. In order for the event $\{X_k^e = \ell^e, e \in E(\mathcal{G}), A_{v,k}\}$ to happen, it must be $I_{k-1} = v_i$ for some $v_i \sim v$ such that $\ell^{\{v, v_i\}} = \ell_i^{e_i^v} > \ell_0^{e_i^v}$, and furthermore it must be $\{I_{k-1}, I_k\} = e_i^v$. Therefore,

$$\begin{aligned} P^{\mathcal{G}}(X_k^e = \ell^e, e \in E(\mathcal{G}), A_{v,k}) \\ = \sum_{i=1, \ell_i^{e_i^v} > \ell_0^{e_i^v}}^{n_v} P^{\mathcal{G}}(X_{k-1}^e = \ell^e, \forall e \neq e_i^v, X_{k-1}^{e_i^v} = \ell_i^{e_i^v} - 1, A_{v_i, k-1}) \\ \times \frac{w(\ell_i^{e_i^v} - 1)}{w(\ell_i^{e_i^v} - 1) + \sum_{e \neq e_i^v \in \mathcal{N}_{v_i}} w(\ell^e)}. \end{aligned}$$

Similarly to Propositions 2 and 3, the proof follows immediately by induction.

□

From now on, denote by

$$\bar{w}_0(\bar{n}) := \frac{\prod_{e \in E(\mathcal{G})} w(\ell_0^e)}{\min_{e \in \mathcal{N}_{v_0}} w(\ell_0^e)},$$

the constant (insuring appropriate scale invariant behavior with respect to w) from the above proposition.

Set $S_1(k) := 1/w(k)$, $k \geq 1$, and for each $n \geq 2$ and $k \geq n$, define

$$S_n(k) := \sum_{\ell_1 + \ell_2 + \dots + \ell_n = k} \frac{1}{w(\ell_1)w(\ell_2) \dots w(\ell_n)}, \quad (48)$$

where the indices ℓ_i , $i = 1, \dots, n$ in the above summation are all greater than or equal to 1. If $k < n$ set simply $S_n(k) := 0$. Then note that, for $k \geq n \geq 2$,

$$\begin{aligned} S_n(k) &= \frac{1}{w(1)} S_{n-1}(k-1) + \frac{1}{w(2)} S_{n-1}(k-2) + \dots \\ &+ \dots + \frac{1}{w(k-n+1)} S_{n-1}(n-1). \end{aligned} \quad (49)$$

Subsequently, we will make use of the following assumption on $w(k)$:

$$\sum_{i=1}^{k-1} \frac{1}{w(i)w(k-i)} \leq \frac{C_w}{w(1)} \cdot \frac{1}{w(k)}, \quad k \geq 1, \quad (A2)$$

where $C_w < \infty$ depends on $w(\cdot)$ up to scaling.

Remark The examples of Theorem 10 (a)-(c) all satisfy (A2). The next lemma will be useful in deriving Corollary 22 below.

Lemma 20 *If (A2) holds, then for all $k \geq n \geq 2$,*

$$S_n(k) \leq \frac{(C_w)^n}{(w(1))^n w(k)}.$$

Proof. We prove the statement inductively. The case $n = 2$ is direct consequence of assumption (A2). Suppose that for some $n > 2$ and for all $k \geq n$ we have $S_n(k) \leq \frac{(C_w)^n}{(w(1))^n w(k)}$. Then, assumption (A2) and identity (49) imply together with inductive hypothesis that for each $k \geq n + 1$

$$S_{n+1}(k) \leq \frac{(C_w)^n}{(w(1))^n} \sum_{j=1}^{k-n} \frac{1}{w(j)w(k-j)} \leq \frac{(C_w)^{n+1}}{(w(1))^{n+1} w(k)}.$$

□

The next result is in the spirit of Lemma 6. It applies in the following setting: fix three different vertices ω, v and u such that $\omega \sim v$ and $v \sim u$. Recall the notation from the beginning of this section. Furthermore, we assume without loss of generality that

$$e_1^v = e_1^u = \{u, v\}, \quad e_1^\omega = e_2^v = \{\omega, v\}. \quad (50)$$

Assume that $n_\omega = q$, $n_v = p$ and $n_u = m$ (recall these are the degrees of the corresponding vertices). We specify the following notation, to be used in the next theorem,

$$P^{\mathcal{G}}(\omega, u, v; k) := P^{\mathcal{G}}(I_k = \omega, I_{k+2i+1} = v, I_{k+2i+2} = u, i \geq 0).$$

Theorem 21 *In the setting of Proposition 19 we have*

$$\begin{aligned} P^{\mathcal{G}}(\omega, u, v; k) &\leq \bar{w}_0(\bar{n}) \cdot \sum_{\ell^e: \sum_e \ell^e = k} \frac{w(\ell^{e_v^v})}{\prod_e w(\ell^e)} \\ &\quad \times \prod_{i=0}^{\infty} \frac{w(\ell^{e_1^v} + 2i)}{w(\ell^{e_1^v} + 2i) + w(\ell^{e_2^v} + 1) + w(\ell^{e_3^v}) + \dots + w(\ell^{e_p^v})} \\ &\quad \times \prod_{i=0}^{\infty} \frac{w(\ell^{e_1^v} + 2i + 1)}{w(\ell^{e_1^v} + 2i + 1) + w(\ell^{e_2^u}) + \dots + w(\ell^{e_m^u})}. \end{aligned} \quad (51)$$

Proof. This is a direct consequence of Proposition 19 and repeated conditioning. Namely, given a particular configuration of weights $\ell^e, e \in E(\mathcal{G})$, (47) estimates the probability for the walk to realize this configuration at time k and to end up at vertex ω at time k , the probability of the next step is

$$w(\ell^{e_1^\omega}) / (w(\ell^{e_1^\omega}) + \dots + w(\ell^{e_q^\omega})) = w(\ell^{e_2^v}) / (w(\ell^{e_1^v}) + \dots + w(\ell^{e_q^v})),$$

and that of the infinitely many steps, each traversing $\{v, u\}$, is given by the two infinite products in the statement. Note that we made use of notation

(50). □

There are various ways to simplify (and lose precision in doing so) the above bound. We chose a particularly simple one for illustration, since we could not find a good enough simplification that would “eliminate” the exponential term in the size \bar{n} of the graph in Corollary 23 below. From now on assume that both (A1) and (A2) hold.

Note that we can bound the sum (51) by

$$\sum_{\ell^e: \sum_e \ell^e = k} \frac{w(\ell^{e_v^v})}{\prod_e w(\ell^e)} \cdot \prod_{i=0}^{\infty} \frac{w(\ell^{e_i^v} + 2i)}{w(\ell^{e_i^v} + 2i) + w(\ell^{e_i^v} + 1)}. \quad (52)$$

Next, rearranging (52) according to the value $s = \ell^{e_1^v} + \ell^{e_2^v}$ yields (recall definitions (48) and (15))

$$\begin{aligned} P\mathcal{G}(\omega, v, u; k) &\leq \bar{w}_0(\bar{n}) \sum_{s=\ell_0^{e_1^v} + \ell_0^{e_2^v}}^{k - \sum_{e \neq e_1^v, e_2^v} \ell_0^e} S_{\bar{n}-2}(k-s) \sum_{j=1}^{s-1} \frac{1}{w(s-j)} \overline{W}_{s+1}(j+1) \\ &\leq \bar{w}_0(\bar{n}) \sum_{s=2}^{k-1} S_{\bar{n}-2}(k-s) \sum_{j=1}^{s-1} \frac{1}{w(s-j)} \overline{W}_{s+1}(j+1) \\ &\leq \bar{w}_0(\bar{n}) \sum_{s=2}^{k-1} S_{\bar{n}-2}(k-s) \sum_{j=1}^{s-1} \frac{1}{w(s-j)} \overline{W}_{k+1}(j+1), \end{aligned} \quad (53)$$

where for the very last inequality we used (A1) that implies $\overline{W}_{s+1}(j+1) \leq \overline{W}_{k+1}(j+1)$, as in Lemma 7 (b). Interchanging the order of summation, applying Lemma 20 and (A2) gives now

$$P\mathcal{G}(\omega, v, u; k) \leq \frac{\bar{w}_0(\bar{n})(C_w)^{\bar{n}-2}}{(w(1))^{\bar{n}-2}} \sum_{j=1}^{k-1} \sum_{s=j+1}^{k-1} \frac{1}{w(k-s)} \frac{1}{w(s-j)} \overline{W}_{k+1}(j+1).$$

and, in turn,

$$P\mathcal{G}(\omega, v, u; k) \leq \frac{\bar{w}_0(\bar{n})(C_w)^{\bar{n}-1}}{(w(1))^{\bar{n}-1}} \sum_{j=1}^{k-1} \frac{1}{w(k-1-(j-1))} \overline{W}_{k+1}(j+1),$$

which comparing with the expression for $\bar{P}^{a,b}$ in Lemma 6, accounting for various possibilities of parity, finally implies

$$P\mathcal{G}(\omega, v, u; k) \leq \bar{w}_1 \frac{\bar{w}_0(\bar{n})(C_w)^{\bar{n}-1}}{(w(1))^{\bar{n}-1}} \sum_{a,b \in \{1,2\}} \bar{P}^{a,b}(T = k+1),$$

where $\bar{w}_1 \in (0, \infty)$ accounts for the “ \asymp ” equivalence of Lemma 6.

Remark It will be convenient for the comparison arguments in the next

corollary to refer to $\bar{P}^{a,b}$ even when a and b are of the same parity. In this case the reader has an option of either noting that the parity does not influence the arguments for Theorem 4, Lemma 6 and Theorem 10, or noting that if $a - b$ is an even number, then the law $\bar{P}^{a,b}$ with reinforcement weight w corresponds to the law $P^{a,b}$ with reinforcement weight \bar{w} , where $\bar{w}(a + j) = w(a + 2j)$, $j \geq 0$.

Corollary 22 *Assuming (A0)-(A2) we have*

$$P^{\mathcal{G}}(T = k + 1) \leq \bar{w}_1 \frac{2\bar{n}D(\mathcal{G})\bar{w}_0(\bar{n})(C_w)^{\bar{n}-1}}{(w(1))^{\bar{n}-1}} \sum_{a,b \in \{1,2\}} \bar{P}^{a,b}(T = k + 1).$$

Proof. Sum over all possible choices of vertex v , edge $\{u, v\}$, and neighbor ω of v , and note that there are at most $2\bar{n}D(\mathcal{G})$ terms of type $P^{\mathcal{G}}(\omega, v, u; k)$ contributing. \square

Corollary 23 *Assuming (A0)-(A2), and $\ell^e = 1$, for all $e \in E(\mathcal{G})$,*

$$P^{\mathcal{G}}(T = k + 1) \leq 2\bar{w}_1\bar{n}D(\mathcal{G})(C_w)^{\bar{n}-1} \sum_{a,b \in \{1,2\}} \bar{P}^{a,b}(T = k + 1).$$

As noted earlier, the examples (a)-(c) from Theorem 10 satisfy (A0)-(A2). In particular, Corollaries 11 and 22 now imply

Corollary 24 *Let \mathcal{G} be a finite graph. Suppose that $w(k)$ is as in (a) or (b) in Theorem 10. Then $E^{\mathcal{G}}(T)$ is infinite if $\rho \leq 1 + \frac{1+\sqrt{5}}{2}$ and finite if $\rho > 1 + \frac{1+\sqrt{5}}{2}$.*

The examples (d)-(e) of Theorem 10 do not satisfy (A2). Here, one could use (53) with separately derived bounds on $S_n(k)$ to obtain bounds on $P^{\mathcal{G}}(T = k + 1)$ as in Corollary 22. In particular, if $w(k) = e^{k^\varepsilon}$, $\varepsilon \in (0, 1]$, then $S_n(k) \leq \binom{k-1}{n-1}/w(k)$, and the above reasoning, together with Theorem 10 (d), gives

$$P^{\mathcal{G}}(T = k + 1) = O(k^{\bar{n}}) \frac{k^{1-\varepsilon}}{e^{(k/2)^\varepsilon}}, \quad (54)$$

which is $\bar{P}^{1,1}(T = k + 1)$ up to a polynomial correction.

3.3 Extensions to bounded degree graphs

Let \mathcal{G} be an infinite graph of bounded degree and as usual let assumption (A0) hold. We wish to estimate

$$P(\#\mathcal{G}_1 > k),$$

where we recall that $\#\mathcal{G}_1$ denotes the number of vertices in the range of the walk. Since $D(\mathcal{G}) < \infty$, note that the above estimate will imply an estimate on $P(|\mathcal{G}_1| > k)$.

Lemma 25 *The random variable $\#\mathcal{G}_1$ is stochastically bounded by $2 \cdot Z$ where Z has geometric distribution with success probability $p \in (0, 1)$, where p depends only on $w(\cdot)$, $D(\mathcal{G})$, and the initial configuration of weights $\ell_0^e, e \in E(\mathcal{G})$.*

As a consequence we obtain that whenever \mathcal{G} has bounded degree, both $\#\mathcal{G}_1$ and $|\mathcal{G}_1|$ have exponential tails.

Proof. We will construct a coupling of \mathcal{G}_1 and \mathcal{G}_1^* such that $\mathcal{G}_1 \subset \mathcal{G}_1^*$, almost surely, and such that the claim of the lemma holds for \mathcal{G}_1^* . Denote by T_v the time of the first visit to the vertex v of \mathcal{G} , where T_v is infinite if the walk never visits v . If $\{T_v = n\}$, then either

- (i) at least one neighbor v' of v has not been visited by the walk before time n , or
- (ii) all the neighbors of v were visited by the walk up to time n .

First suppose that case (i) happens. Then “add” to \mathcal{G}_1^* both vertices v and v' as well as just traversed edge leading to v and the edge $\{v, v'\}$. Due to the assumptions, with probability p , uniformly bounded away from 0 that depends only on $D(\mathcal{G})$, $w(\cdot)$, and $\ell_0^e, e \in E(\mathcal{G})$ (in fact only ℓ_0^f on edges f incident to v'), the walk keeps traversing solely the edge $\{v, v'\}$ after time n . In symbols, on the event of case (i),

$$P(\{I_k, I_{k+1}\} = \{v, v'\}, k \geq n | \mathcal{F}_n) > p.$$

If the above event $\{I_k, I_{k+1}\} = \{v, v'\}, k \geq n$ does not occur, the walk will keep exploring the graph elsewhere. Either it will get attracted to an edge before encountering another new vertex or it will encounter another new vertex prior to getting attracted.

If the case (ii) happens note that \mathcal{G}_1^* already contains vertex v . Namely, let u be the neighbor of v such that

$$T_u = \max_{v' \sim v} T_{v'} < T_v.$$

Then v must have been added to \mathcal{G}_1^* as part of case (i) procedure, before or at time T_u .

Therefore, $\mathcal{G}_1 \subset \mathcal{G}_1^*$, almost surely, by induction. Moreover, from the construction it is clear that $\#\mathcal{G}_1$ is stochastically bounded by $2Z$. \square

As a conclusion, we offer the following weak universality-type result.

Corollary 26 *Assume $\ell^e = 1$, for all $e \in E(\mathcal{G})$.*

If $w(k) = k^\rho$, $\rho > 1$, there exists $p > 0$ such that $P^{\mathcal{G}}(T > k) \leq \frac{1}{k^p}$,

Proof. Use the idea of Corollary 17 together with the previous lemma and the bound of Corollary 23. Namely, split the event $\{T > k\}$ according to whether or not the walk reaches distance d_k from I_0 . Choose $d_k = c \log k$ where c is such that $(C_w)^{c \log k} \ll k^{\rho - \rho' - 1}$. \square

Remark If $w(k) = e^k$, the walk gets attracted at any particular step with probability bounded away from 0, so there exists $c > 0$ such that $P^{\mathcal{G}}(T >$

$k) = O\left(\frac{1}{e^{ck}}\right)$. Somewhat disappointingly, the bound of type (54) is too weak to provide an alternative derivation (analogous to the proof of the last corollary) of the above bound. Indeed, the question of finding exact (up to multiplicative constant) behavior of the tail distribution of T on general bounded degree graphs, even in the case of examples in Theorem 10, remains open.

Acknowledgment

C.C. is very grateful to David Brydges for his advice and support during her time as a postdoctoral fellow at University of British Columbia. We wish to thank the anonymous referee for many helpful comments and suggestions.

References

- [1] B. Davis. Reinforced random walk. *Probab. Theo. Rel. Fields*, 84:203–229, 1990.
- [2] R. Beckers, J.-L. Deneubourg, and S. Goss. Trails and U-turns in the selection of a path by the ant *Lasius Niger*. *J. Theor. Biol.*, 159:397-415, 1992.
- [3] E. Bonabeau, M. Dorigo, and G. Theraulaz. *Swarm Intelligence, From Natural to Artificial Systems*. Oxford University Press, 1999.
- [4] J.-L. Deneubourg, S. Aron, S. Goss, and J.M. Pasteels. The self-organising exploratory pattern of the argentine ant. *Journal of insect behavior*, 3:159-168, 1990.
- [5] V. Limic. Attracting edge property for a class of reinforced random walks. *Ann. Probab.*, 31:1615–1654, 2003.
- [6] V. Limic and P. Tarrès. Attracting edge and strongly reinforced random walks. *Ann. Probab.*, to appear, 2007.
- [7] F. Merkl and S. Rolles. Edge-reinforced random walk on a ladder. *Ann. Probab.*, 33:2051-2093, 2005.
- [8] R. Oliveira and J. Spencer. Avoiding defeat in a balls-in-bins process with feedback. preprint, 2005.
- [9] R. Pemantle. A survey of random proceses with reinforcement. *Probab. Surv.*, 4:1-79, 2007.
- [10] R. Pemantle. Phase transition of reinforced random walk and rwre on trees. *Ann. Probab.*, 16:1229-1241, 1998.

- [11] T. Sellke. Reinforced random walks on the d –dimensional integer lattice. *EJP*, to appear, 2007.
- [12] P. Tarrès. VRRW on \mathbb{Z} eventually gets stuck at a set of five points. *Ann. Probab.*, 32:2650–2701, 2004.