# Option Pricing without Price Dynamics: A Probabilistic Approach

Dimitris Bertsimas \* Natasha Bushueva <sup>†</sup>

#### Abstract

Employing probabilistic techniques we compute best possible upper and lower bounds on the price of an option on one or two assets with continuous piecewise linear payoff function based on prices of simple call options of possibly distinct maturities and the no-arbitrage condition, but without any assumption on the price dynamics of underlying assets. We show that the problem reduces to solving linear optimization problems that we explicitly characterize. We report numerical results that illustrate the effectiveness of the algorithms we develop.

<sup>\*</sup>Boeing Professor of Operations Research, Sloan School of Management, Rm. E53-363, Massachusetts Institute of Technology, Cambridge, Mass. 02139. dbertsim@mit.edu. Research partially supported by the MIT-Singapore Alliance.

<sup>&</sup>lt;sup>†</sup>Work was done while being a graduate student at the Mathematics department of Massachusetts Institute of Technology and is part of her PhD thesis (Apr 2003; http://hdl.handle.net/1721.1/29349). natacha@alum.mit.edu.

## 1 Introduction

One of the central questions in financial economics is to find the price of a derivative security given information on the underlying asset. Pricing of a derivative is usually realized by solving the Black-Scholes equation [4], which is based on the assumptions of a Geometric Brownian motion for the price of the underlying asset and no-arbitrage in the market. One of the parameters in the equation, the volatility of returns on the underlying asset, is assumed to be defined and constant, while in reality it is varying in time and not known in advance. Practitioners typically use the implied volatility to value a derivative security by the Black-Scholes formula, that is they extract the volatility from prices of other options in the market, based on the assumption that the Black-Scholes equation is valid and other options are correctly priced.

A natural question that arises is to determine the range of values for the price of an option based only on prices of other options and the no-arbitrage assumption, but with no assumption on the price dynamics of the underlying asset.

In their seminal work, Cox and Ross [7] and Harrison and Kreps [12] show that the condition of no-arbitrage is equivalent to the existence a probability measure Q, equivalent to the original measure P, with respect to which all discounted securities processes are martingales. Rubinstein [18] and Longstaff [16] introduce the idea of deducing the martingale measure from observed European call prices by solving a quadratic optimization problem.

Lo [15] derives best possible closed form bounds on the price of a European call option given the mean and variance of the underlying stock price under risk neutral measure. Grundy [11] extends Lo's work for the case when the first and the *k*th moments of the stock price are known. Bertsimas and Popescu [3] derive best possible bounds of the price of a European call option, as well as on moments of the prices of the asset, given prices of other similar options, using a convex optimization approach. D'Aspremont and El Ghaoui [1] address the problem of computing upper and lower bounds on the price of a European basket call option, given prices on other similar baskets. They introduce a linear programming relaxation of the problem and show that this relaxation is best possible in some special cases. All the above problems are solved for the case of a **single maturity** and single underlying stock.

In this paper, we consider the problem of determining upper and lower bounds on the price of a European option given prices of other options of **different maturities**. In particular, the contributions of the article are the following:

(a) We establish necessary and sufficient conditions that European call options of different maturities should jointly satisfy, so that there exists no arbitrage. From this structural property and using geometric arguments, one can derive best bounds on the price of a European call option with a given strike price and maturity. As we

present more general methods in later sections we do not present this method here (see Bertsimas and Bushueva [2]).

- (b) Given a set of European call options of different strike prices and different maturities we determine best possible upper and lower bounds on the price of an option with a continuous piecewise linear payoff function. Based on direct probabilistic methods, we solve the problem as a linear optimization problem.
- (c) Given European call options on two individual assets we determine best possible upper and lower bounds on the price of an option with a payoff being a continuous piecewise linear function of prices of the two assets at the time of option's maturity. Again, options are allowed to have different maturities. We present two algorithms based on linear optimization. The first is an exact formulation and involves a linear optimization problem of potentially exponential size; the second is asymptotically exact and involves a linear optimization problem of polynomial size.
- (d) We determine upper and lower bounds on the price of a European basket call option, given prices on other similar baskets solving exactly the problem addressed in D'Aspremont and El Ghaoui [1] approximately.

Throughout the paper, we refer to these problems as Problems (a), (b), (c), (d). The paper is structured as follows: in Sections 2, 3, 4, 5 we present our solutions for Problems (a), (b), (c) and (d) respectively.

## 2 Characterization theorem for the case of multiple maturities

In this section, we determine best possible bounds on the price of a European call option given prices of options on the same underlying security, but with potentially different maturities.

The price of a European call option with a strike price k and maturity t is given by

$$C = e^{-\int_{0}^{t} r(s)ds} E_{Q}\left[ (S_{t} - k)^{+} \right] = E_{Q}\left[ \left( X_{t} - ke^{-\int_{0}^{t} r(s)ds} \right)^{+} \right], \quad (1)$$

where r(s) is the instantaneous riskless rate of return,  $\{S_t\}_{t\geq 0}$  is the price process of the underlying security defined on some probability space  $(\Omega, \mathcal{B}, P)$ , and Q is a measure equivalent to P, such that  $\{X_t\}_{t\geq 0} := \{S_t e^{-\int_0^t r(s)ds}\}_{t\geq 0}$  is a martingale process under Q. Two measures are equivalent if they have the same null sets.

Thus, the problem we address in this section can be reformulated as follows: given the set

$$\mathcal{S} := \left\{ (k_{ij}, C_{ij}, t_i) \in \mathbb{R}^3_+ \mid t_1 < t_2 < \ldots < t_n, i \in \{1, 2, \ldots, n\}, \ j \in \{1, 2, \ldots, U(i)\} \right\}$$

find the sufficient and necessary condition that values of the set should jointly satisfy so that there exists a nonnegative martingale  $\{X_{t_1}, X_{t_2}, \ldots, X_{t^*}, \ldots, X_{t_n}\}$ , such that

$$E\left[\left(X_{t_i} - k_{ij}e^{-\int_0^{t_i} r(s)ds}\right)^+\right] = C_{ij} \tag{2}$$

for all points of the set.

#### Remarks:

- (a) We can set without loss of generality  $r(s) \equiv 0$ , since we can define  $\tilde{k}_{ij} = k_{ij}e^{-\int_0^{t_i} r(s)ds}$ , and rewrite Eq. (2) as  $E\left[\left(X_{t_i} \tilde{k}_{ij}\right)^+\right] = C_{ij}$ .
- (b) Since we can assume  $r(s) \equiv 0$ , only the order of  $t_1, t_2, \ldots, t_n$  is important, but not the specific values that the sequence takes.
- (c) The price of an option with a strike price 0 is given by  $E[X_t] = X_0$ , which is the current stock price and, thus, is always known.

In light of the above remarks, a simpler formulation of the problem is as follows.

#### Problem (a)

Given an ordered set

$$\mathcal{S} := \left\{ (k_{tj}, C_{tj}) \in \mathbb{R}^2_+ \mid t \in \{1, 2, \dots, n\}, \ j \in \{1, 2, \dots, U(t)\} \right\}$$
(3)

and  $X_0 \in \mathbb{R}_+$  find the sufficient and necessary condition that values of the set should jointly satisfy so that there exists a nonnegative martingale  $\{X_1, \ldots, X_{t^*}, \ldots, X_n\}$  such that

$$E\left[\left(X_t - k_{tj}\right)^+\right] = C_{tj} \text{ for all } (k_{tj}, C_{tj}) \in \mathcal{S},$$
(4)

and

$$E[X_t] = X_0$$
 for all  $t = 1, \ldots, n$ .

Let  $\mathcal{C}$  designate the class of probability laws with support in  $\mathbb{R}_+$  and finite expectations. We associate with each element  $\Pi$  of  $\mathcal{C}$  the transform  $\Psi_{\Pi}$ :  $\mathbb{R}_+ \to \mathbb{R}_+$ , defined as

$$\Psi_{\Pi}(t) = E_{\Pi}\left[(x-t)^{+}\right] = \int_{0}^{\infty} (x-t)^{+} d\Pi(x).$$
(5)

We next show that  $\Psi_{\Pi}(t)$  uniquely determines  $\Pi$ .

**Theorem 1.** (a) The right derivative  $\Psi_{\Pi}'(t+)$  of  $\Psi_{\Pi}(t)$  exists for all  $t \ge 0$  and

$$\Psi_{\Pi}'(t+) = -\Pi((t, +\infty)).$$

(b) Let  $\Pi_1$ ,  $\Pi_2 \in \mathcal{C}$  and for each  $t \in \mathbb{R}_+$ ,  $\Psi_{\Pi_1}(t) = \Psi_{\Pi_2}(t)$ . Then,  $\Pi_1 = \Pi_2$ .

*Proof.* (a) Let  $t \ge 0$  and h > 0. We have

$$\frac{\Psi_{\Pi}(t+h) - \Psi_{\Pi}(t)}{h} = \int_0^\infty \frac{(x - (t+h))^+ - (x-t)^+}{h} \, d\Pi(x) = \int_0^\infty f_{t,h}(x) \, d\Pi(x),$$

where the function  $f_{t,h}(x)$  is defined as follows (see also Figure 1):

$$f_{t,h}(x) := \frac{(x - (t + h))^{+} - (x - t)^{+}}{h} = \begin{cases} 0, & x \le t, \\ -\frac{x - t}{h}, & t \le x \le t + h, \\ -1, & x \ge t + h. \end{cases}$$



Figure 1: The function  $f_{t,h}(x)$ .

Since the sequence of functions  $f_{t,h}(x)$  is monotone (in h), and  $f_{t,h}(x)$  converges as  $h \to 0^+$ , then by by the monotone convergence theorem [9, p.100],  $\int_0^{+\infty} f_{t,h}(x) d\Pi(x) \to -\Pi((t, +\infty))$  as  $h \to 0^+$ . Thus  $\Psi'_{\Pi}(t+)$  exists and equals  $-\Pi((t, +\infty))$ . (b) Immediately follows from (a).

The following proposition summarizes the properties of the  $\Psi$ -transform.

**Proposition 1.** Let  $\Pi \in \mathcal{C}$ . Then  $\Psi_{\Pi}(t)$  satisfies:

- (a)  $\Psi_{\Pi}(t)$  is a nonincreasing function of  $t \in \mathbb{R}_+$ ;
- (b)  $\Psi_{\Pi}(t)$  is convex;
- (c)  $\lim_{t\to+\infty} \Psi_{\Pi}(t) = 0$ . In particular, if  $\Pi$  has a bounded support, then  $\Psi_{\Pi}(t) = 0$  for t large;
- (d)  $\Psi_{\Pi}(t) \ge \Psi_{\Pi}(0) t.$

*Proof.* (a) Notice, that  $\Psi_{\Pi}(t)$  is a continuous function:

$$|\Psi_{\Pi}(t+h) - \Psi_{\Pi}(t)| = \left| \int_{t+h}^{t} (x-t)d\Pi(x) \right| \le |h|.$$

Since  $\Psi_{\Pi}(t)$  is continuous and  $\Psi_{\Pi}'(t+) = -\Pi((t, +\infty)) \leq 0$ , then  $\Psi_{\Pi}(t)$  is nonincreasing.

(b)  $f_x(t) := (x - t)^+$  is convex in t for each x. Thus,  $\Psi_{\Pi}(t)$  is convex as a convex combination of convex functions.

$$\Psi_{\Pi}(t) = \int_{0}^{+\infty} (x-t)^{+} d\Pi(x) = \int_{t}^{+\infty} (x-t)^{+} d\Pi(x) \le \int_{t}^{+\infty} x d\Pi(x).$$

Since  $\int_0^{+\infty} x d\Pi(x) < \infty$ , then  $\int_t^{+\infty} x d\Pi(x) \to 0$  as  $t \to \infty$ . Thus,  $\Psi_{\Pi}(t) \to 0$  as  $t \to +\infty$ . (d)  $\Psi_{\Pi}(t) = \int_0^{\infty} (x-t)^+ d\Pi(x) \ge \int_0^{\infty} (x-t) d\Pi(x) = \Psi_{\Pi}(0) - t$ .

We next show that the properties in Proposition 1 are also sufficient for a function  $\Psi$  to be a  $\Psi$ -transform of some distribution  $\Pi$ .

**Theorem 2.** Let  $g : \mathbb{R}_+ \to \mathbb{R}_+$  be a function such that (a) g(x) is nonincreasing, (b) g(x) is convex, (c)  $\lim_{x\to+\infty} g(x) = 0$ , and (d)  $g(x) \ge g(0) - x$ . Then, there exists  $\Pi \in \mathcal{C}$  such that  $g(x) = \Psi_{\Pi}(x)$ .

*Proof.* From the convexity of g it follows that the right derivative of g, g'(x+), is a right continuous nondecreasing function. Since  $g(x) \ge g(0) - x$  and g(x) is nonincreasing, then  $-1 \le g'(x+) \le 0$ . Also we have  $g'(x+) \to 0$  as  $x \to +\infty$ . To justify the last statement, notice that g'(x+) is a monotone bounded function, and consequently, it must converge to some finite value a as  $x \to \infty$ . However,  $a \ne 0$  would contradict the requirement in (c).

Consider the function F(x) := 1 + g'(x+) defined on  $\mathbb{R}_+$ . From the above statements, it follows that (i)  $0 \le F(x) \le 1$ , (ii) F(x) is a right-continuous and nondecreasing, and (iii)  $F(x) \to 1$  as  $x \to \infty$ . Thus, F(x) satisfies all the properties of a distribution function. Denote by  $\Pi$  the law with the distribution function F(x). Then,

$$\Psi_{\Pi}(0) = E_{\Pi}[X] = \int_{0}^{+\infty} (1 - F(x))dx = \int_{0}^{+\infty} (-g'(x+)) \, dx \stackrel{*}{=} g(0) - \lim_{x \to +\infty} g(x) = g(0).$$

The second to last equality (identified with "\*") holds since g(x) is a convex function and thus its right and left derivatives might differ only on a countable set of points.

Since  $\Psi_{\Pi}'(x+) = -(1 - F(x)) = g'(x+)$ ,  $\Psi_{\Pi}(x)$  and g(x) are both continuous and  $\Psi_{\Pi}(0) = g(0)$ , it follows that  $\Psi_{\Pi}(x) = g(x)$  for any  $x \in \mathbb{R}_+$ .

Since there is a one to one correspondence between laws of random variables and their  $\Psi$ -transforms, it follows that conditions that a sequence of laws of random variables should satisfy can always be reformulated in terms of conditions on the sequence of

 $\Psi$ -transforms of these random variables. Theorem 3 establishes necessary and sufficient conditions that the sequence of  $\Psi$ -transforms should satisfy so that the corresponding random variables form a martingale. However prior to proving that theorem, we will need the following auxiliary results.

**Definition 1.** Let  $\Pi_1$ ,  $\Pi_2 \in C$ . Define the relation  $\Pi_1 \mathcal{R} \Pi_2$  if and only if there exist random variables  $X_1$  and  $X_2$ , defined on the same probability space, such that  $\{X_1, X_2\}$ is a martingale and  $X_1$  has a marginal law  $\Pi_1$ , while  $X_2$  has a marginal law  $\Pi_2$ .

Proposition 2. The following properties hold:

- (a) If  $\Pi_1 \mathcal{R} \Pi_2$  and  $\Pi_2 \mathcal{R} \Pi_1$  then  $\Pi_1 = \Pi_2$ ;
- (b)  $\Pi_1 \mathcal{R} \Pi_1;$
- (c) If  $\Pi_1 \mathcal{R} \Pi_2$  and  $\Pi_2 \mathcal{R} \Pi_3$ , then  $\Pi_1 \mathcal{R} \Pi_3$ .

Proof. (a) If  $\{X_1, \mathcal{B}_1\}$ ,  $\{X_2, \mathcal{B}_2\}$  is a martingale with marginal laws  $\Pi_1$  and  $\Pi_2$  correspondingly, and f is a convex function, defined on  $\mathbb{R}$ , then  $f(X_1) = f(E[X_2|\mathcal{B}_1]) \leq E[f(X_2)|\mathcal{B}_1]$  by the conditional Jensen's inequality, and thus  $E[f(X_1)] \leq E[f(X_2)]$ .  $(X-t)^+$  is a convex function of X, so  $\Psi_{\Pi_1}(t) \leq \Psi_{\Pi_2}(t)$  for each  $t \in \mathbb{R}_+$ . From  $\Pi_2 \mathcal{R} \Pi_1$  by the same argument we obtain  $\Psi_{\Pi_2}(t) \leq \Psi_{\Pi_1}(t)$  for each  $t \in \mathbb{R}_+$ . Thus  $\Psi_{\Pi_1}(t) = \Psi_{\Pi_2}(t)$  and by Theorem 1,  $\Pi_1 = \Pi_2$ .

(b)  $\{X_1, X_1\}$  is always a martingale.

(c) Let  $\{X_1, X_2\}$  be a martingale with the law  $\mathcal{L}(X_1, X_2)$  defined on  $\mathbb{R} \times \mathbb{R}$  with support in  $\mathbb{R}_+ \times \mathbb{R}_+$  and marginal laws  $\Pi_1$  and  $\Pi_2$  respectively. Let  $\{\tilde{X}_2, X_3\}$  be a martingale with the law  $\mathcal{L}(\tilde{X}_2, X_3)$  defined on  $\mathbb{R} \times \mathbb{R}$  with support in  $\mathbb{R}_+ \times \mathbb{R}_+$  and marginal laws  $\Pi_2$  and  $\Pi_3$ . Then on  $\mathbb{R}$  there exist conditional distributions  $\mathcal{L}(X_1|X_2)$  for  $\mathcal{L}(X_1, X_2)$ and  $\mathcal{L}(X_3|\tilde{X}_2)$  for  $\mathcal{L}(\tilde{X}_2, X_3)$ . Since the marginal laws of  $X_2$  and  $\tilde{X}_2$  are equal, then by Vorob'ev-Berkes-Philipp theorem [10, p.7], we can define a law  $\mathcal{L}(X_1, X_2, X_3)$  on  $\mathbb{R} \times \mathbb{R} \times \mathbb{R}$ , such that  $X_1$  and  $X_3$  are conditionally independent given  $X_2$ , that is

$$\mathcal{L}((X_1, X_3)|X_2) = \mathcal{L}(X_1|X_2) \times \mathcal{L}(X_3|X_2).$$

Denote by  $\mathcal{B}_i$ , i = 1, 2, 3, the smallest  $\sigma$ -algebra for which all  $X_k$ ,  $k \leq i$ , are measurable. Then  $\{X_i, \mathcal{B}_i\}_{i=1}^3$  is a martingale with marginal laws  $\Pi_1, \Pi_2$  and  $\Pi_3$  correspondingly. Note that the martingale thus constructed is also a Markov process.

**Theorem 3.** Let  $\Pi_1, \Pi_2, \ldots, \Pi_n \in C$  and  $E_{\Pi_i}[X] = E_{\Pi_1}[X]$  for each  $i = 1, \ldots, n$ . Then, there exists a martingale  $\{X_1, X_2, \ldots, X_n\}$ , such that  $X_i$  has marginal law  $\Pi_i$  for each i if and only if the sequence of functions  $\Psi_{\Pi_1}, \Psi_{\Pi_2}, \ldots, \Psi_{\Pi_n}$  is nondecreasing.

**Remark:** The theorem was obtained in Kertz and Rosler [14]. The key step in their proof is a result of Strassen [19]. Our approach is completely different.

*Proof.* The only if part is a direct consequence of the fact that  $(X - t)^+$  is a convex function of X and Jensen's inequality for convex functions of martingales [9, p.277]. The proof of this statement is also in the proof of part (a) of Proposition 2.

In the other direction, we have to show that it is possible to define a joint distribution of  $X_1, X_2, \ldots, X_n$  with the given marginal laws (since they are uniquely determined by  $\Psi_{\Pi_1}, \Psi_{\Pi_2}, \ldots, \Psi_{\Pi_n}$ ), so that this sequence is a martingale. It is enough to show how to define a joint distribution of  $X_1$  and  $X_2$ , since afterwards we can proceed recursively to form a martingale which is a Markov process as in the proof of Proposition 2.

We first define a sequence  $X_1, Y_1, Y_2, \ldots, Y_n, \ldots$ , so that this sequence is a martingale and  $\Psi_{Y_n} \to \Psi_{\Pi_2}$ . We will then show that there exists  $Y_\infty$ , such that  $\{Y_n\}_{1 \le n \le \infty}$  is a martingale and  $Y_\infty$  has a distribution  $\Pi_2$ . Then  $X_2$  can be defined as  $X_2 = Y_\infty$ .

Let  $Y_0 = X_1$  and let us form a martingale  $Y_0, Y_1, Y_2, \ldots, Y_n$  such that it is also a Markov process, and thus, we only need to define the joint distribution of  $Y_i$  and  $Y_{i+1}$ for each *i*. Now, the joint distribution of  $Y_i$  and  $Y_{i+1}$  is uniquely determined as soon as we know for each  $y \in \mathbb{R}_+$  the conditional distribution of  $Y_{i+1}$  given  $Y_i = y$ . Thus, we are going to define the conditional distribution for  $Y_1$ , then for  $Y_2$  and so on. We



Figure 2: Construction of  $\Psi_{Y_1}$ .

know that  $\Psi_{X_1} = \Psi_{\Pi_1} \leq \Psi_{\Pi_2}$ . Now  $Y_1$  will have its  $\Psi$ -transform,  $\Psi_{Y_1}$ , equal to  $\Psi_{X_1}$ everywhere except on (a, b) where it will be a straight line (Figure 2.) We choose (a, b) so that the line which goes through the points  $(a, \Psi_{X_1}(a)), (b, \Psi_{X_1}(b))$  is below the function  $\Psi_{\Pi_2}(t)$  (the specific choice of (a, b) is discussed later.) What distribution will have its  $\Psi$ -transform equal to the transform of  $\Pi_1$  everywhere except (a, b) and being a straight line on (a, b)? From Theorem 1 (a), we see that  $\Psi_{Y_1}$  should have no weight on (a, b) but the weight of [a, b] should be the same as for the distribution  $\Pi_1$ . Thus, we take all the weight of (a, b) and redistribute it between  $\{a\}$  and  $\{b\}$ . We have to do it in such a way that  $\{X_1, Y_1\}$  is a martingale.

The law  $\mathcal{L}(X_1)$  is uniquely determined by  $\Psi_{X_1}$  and the law  $\mathcal{L}(Y_1)$  is uniquely determined by  $\Psi_{Y_1}$ . We define the law  $P := \mathcal{L}(X_1, Y_1)$  with support in  $\mathbb{R}_+ \times \mathbb{R}_+$  in the following way. As the marginal law of  $X_1$  take  $\Pi_{X_1}$ . Then for each  $X_1 \in \mathbb{R}_+$  define the conditional distribution of  $Y_1$  given  $X_1$ . First, if  $X_1 \in \{\mathbb{R}_+ \setminus (a, b)\}$ , take  $Y_1 = X_1$ . Now, if  $X_1 \in (a, b)$  we should have  $P(Y_1 \in \{a\} \cup \{b\} | X_1 \in (a, b)) = 1$  for  $\Psi_{Y_1}(t)$  to be a straight line on (a, b) and equal to  $\Psi_{X_1}(t)$  everywhere else. Let us denote  $\pi_{X_1}(X_1, Y_1) := X_1$ .

Thus, for any  $c \in (a, b)$  we want the joint distribution of  $X_1, Y_1$  to be such that the following holds

$$aP(Y_1 = a|X_1 = c) + bP(Y_1 = b|X_1 = c) = c,$$
  
 $P(Y_1 = a|X_1 = c) + P(Y_1 = b|X_1 = c) = 1.$ 

Consequently,

$$P(Y_1 = a | X_1 = c) = \frac{b - c}{b - a}, \quad P(Y_1 = b | X_1 = c) = \frac{c - a}{b - a}$$

Thus, the conditional distribution of  $Y_1$  given  $X_1$  is defined and  $\{X_1, Y_1\}$  is a martingale. In the same manner we can define the conditional distribution of  $Y_{i+1}$  given  $Y_i$ , for i = 1, 2, ...

The next step is to show that  $\Psi_{X_2}(t)$  can be approached by  $\Psi_{Y_n}(t)$  with any given precision, that is that we can choose  $Y_1, Y_2, \ldots, Y_n$  so that  $\Psi_{Y_n}(t) \to \Psi_{X_2}(t)$ . For each  $i, \Psi_{Y_{i+1}}(t)$  is different from  $\Psi_{Y_i}(t)$  only on a certain interval  $(a_i, b_i)$ , where  $\Psi_{Y_{i+1}}(t)$  is a straight line. Thus, the choice of the sequence  $\{Y_n\}$  is equivalent to the choice of the sequence  $\{(a_n, b_n)\}$ .



Figure 3: The choice of  $(a_1, b_1)$ .

Let

$$U := \{ t \in \mathbb{R}_+ : \Psi_{X_2}(t) > \Psi_{X_1}(t) \} \,.$$

Let  $\{t_n\}$  be a countable dense set in U. At each  $t_i$ , i = 1, 2, ..., draw a tangent line  $y_i(t)$  to  $\Psi_{X_2}(t)$ . (See Figure 3.) Let us first assume that  $\Psi_{X_2}(t_i) \neq 0$ . Then  $\Psi_{X_2}'(t_i+) < 0$  and the tangent line intersects  $\Psi_{Y_{i-1}}(t)$  at two different points,  $a_i \in \mathbb{R}_+$  and  $b_i \in \mathbb{R}_+$ , since  $\Psi_{Y_i}(t) \to 0$  as  $t \to 0$  and  $\Psi_{Y_i}(0) = 1$ , while  $y_i(t)$  is below  $\Psi_{X_2}(t)$  and thus  $y_i(0) \leq 1$ . Thus, the choice of  $\{t_n\}$  is equivalent to the choice of  $\{(a_n, b_n)\}$ . We define  $\Psi_{Y_{i+1}} = \max\{y_i(t), \Psi_{Y_i}(t)\}$ . In case  $\Psi_{X_2}(t_i) = 0$ ,  $\Psi_{X_2'}(t_i) = 0$  and we take  $\Psi_{Y_{i+1}} := \Psi_{Y_i}$ .

Thus, the sequence  $\{\Psi_{Y_n}\}$  is an increasing sequence of continuous functions converging to a continuous function  $\Psi_{X_2}$  on a countable dense set. Consequently,  $\Psi_{Y_n} \to \Psi_{X_2}$ pointwise on  $\mathbb{R}_+$ . Since  $\Psi_{X_1}(t) \to 0$  and  $\Psi_{X_2} \to 0$  as  $t \to +\infty$ , then for any  $\epsilon$  there exist  $T \in \mathbb{R}_+$ such that for any t > T,  $|\Psi_{X_2}(t) - \Psi_{X_1}(t)| < \epsilon$ . Then for any n and any t > T,  $|\Psi_{X_2}(t) - \Psi_{Y_n}(t)| < \epsilon$ . Taking into account that  $\{\Psi_{Y_n}\}$  converges uniformly to  $\Psi_{X_2}$  on [0,T], we conclude that  $\Psi_{Y_n}$  converges uniformly to  $\Psi_{X_2}$  on  $\mathbb{R}_+$ .

Now, since  $\Psi_{Y_n} \to \Psi_{X_2}$  uniformly on  $\mathbb{R}_+$  and the right derivatives of  $\Psi_{Y_n}(t)$ ,  $n = 1, 2, \ldots$ , and  $\Psi_{X_2}$  are monotone functions, then also  $\Psi_{Y_n}'(t+) \to \Psi_{X_2}'(t+)$ . Thus, the distribution functions of  $Y_n$ ,  $n = 1, 2, \ldots$ , converge pointwise to the distribution function of  $X_2$ .

We next show that  $\{Y_n\}$  is uniformly integrable, i.e., for any  $\epsilon > 0$  there exists  $t_0 > 0$ , such that for all  $t > t_0$ ,

$$\sup_{n} E\left[Y_n \mathbb{1}_{\{Y_n > t\}}\right] < \epsilon.$$

For each t and n we have  $\Psi_{Y_n}(t) \leq \Psi_{X_2}(t)$ . Therefore,

$$\int_{0}^{+\infty} (Y_n - t)^+ d\Pi_{Y_n} \le \int_{0}^{+\infty} (X_2 - t)^+ d\Pi_{X_2},$$
$$\int_{t}^{+\infty} Y_n \ d\Pi_{Y_n} \le \int_{t}^{+\infty} X_2 \ d\Pi_{X_2} + t \left| \Pi_{Y_n} \left( [t, +\infty] \right) - \Pi_{X_2} \left( [t, +\infty] \right) \right|.$$

Since  $\int_t^{+\infty} X_2 \, d\Pi_{X_2} \to 0$  as  $t \to +\infty$ , then we can take  $t_0$  such that  $\int_t^{+\infty} X_2 \, d\Pi_{X_2} < \epsilon/3$  for all  $t > t_0$ . Now since  $\Pi_{Y_n} ([t_0, +\infty]) \to \Pi_{X_2} ([t_0, +\infty])$ , there exists  $n_0$  such that for all  $n > n_0$ ,  $|\Pi_{Y_n} ([t_0, +\infty]) - \Pi_{X_2} ([t_0, +\infty])| < \frac{\epsilon}{3t_0}$ . Thus, for any  $n > n_0$  and  $t > t_0$  we have

$$E\left[Y_n \mathbb{1}_{\{Y_n > t\}}\right] \le E\left[Y_n \mathbb{1}_{\{Y_n > t_0\}}\right] < \frac{\epsilon}{3} + t_0 \frac{\epsilon}{3t_0} = \frac{2\epsilon}{3}.$$

Therefore,

$$\sup_{n>n_0} E\left[Y_n \mathbb{1}_{\{Y_n>t\}}\right] < \frac{2\epsilon}{3}.$$

For  $Y_1, Y_2, \ldots, Y_{n_0}$  we can always choose  $t_1$  big enough so that for all  $t > t_1$ ,  $E\left[Y_i \mathbb{1}_{\{Y_i > t\}}\right]$  $< \frac{\epsilon}{3}, i = 1, 2, \ldots, n_0$ , since all  $Y_i$  have finite expectations. Consequently, if we denote  $t^* := \max\{t_0, t_1\}$ , then

$$\sup_{n} E\left[Y_n \mathbb{1}_{\{Y_n > t\}}\right] < \epsilon,$$

for all  $t > t^*$ .

Thus,  $Y_1, Y_2, \ldots, Y_n$  is uniformly integrable, and, consequently [9, p.283], right closable, which implies that there exists a random variable  $Y_{\infty}$  such that  $E[Y_{\infty}|\mathcal{B}_n] = Y_n$  for all n, where  $\mathcal{B}_n$  designates the smallest  $\sigma$ -algebra for which  $Y_1, \ldots, Y_n$  are all measurable. Then by Doob's theorem [9, p.285]  $Y_n$  converges to  $Y_{\infty}$  a.s., and consequently the distribution functions of  $Y_1, Y_2, \ldots, Y_n$  also converge to the distribution function of  $Y_{\infty}$ . But then  $Y_{\infty}$  and  $X_2$  have the same distribution function, and thus the same  $\Psi$ -transform. So we take  $X_2 = Y_{\infty}$  and  $X_1, X_2$  is a martingale.

Now we are ready to formulate the conditions that the set S defined in (3) should satisfy, so that there exists a martingale  $\{X_1, X_2, \ldots, X_n\}$ , such that Eq. (4) holds.

**Proposition 3.** The set S, as defined in (3), satisfies the condition of martingale existence if and only if there exists a nondecreasing sequence of convex functions,  $\{g_t(x)\}_{t \in \{1,...,n\}}$ :  $\mathbb{R}_+ \to \mathbb{R}_+$ , such that for each  $t \in \{1, 2, ..., n\}$ 

(a) 
$$g_t(0) = g_1(0);$$

- **(b)**  $g_t(x) \ge g_t(0) x;$
- (c)  $g_t(x) \to 0$  as  $x \to \infty$ ;
- (d)  $g_t(k_{tj}) = C_{tj}$  for each  $j \in \{1, 2, \dots, U(t)\}$ .

*Proof.* " $\implies$ " By Theorem 2 for each  $g_t$  there exists  $\Pi_t \in \mathcal{C}$  such that  $\Psi_{\Pi_t} = g_t$ . Then, since the sequence of functions  $\Psi_{\Pi_1}, \Psi_{\Pi_1}, \dots, \Psi_{\Pi_n}$  is nondecreasing and  $\Psi_{\Pi_t}(0) = \Psi_{\Pi_1}(0)$  for each t, from Theorem 3 it follows that the martingale satisfying all the points of the set  $\mathcal{S}$  exists.

"  $\Leftarrow$ " Suppose that there exists a martingale  $\{X_1, X_2, \ldots, X_n\}$ , such that (4) holds. Then we can take  $g_t(x) := \Psi_{X_t}(x)$  for  $t = 1, 2, \ldots, n$ . By Theorem 3 the sequence of functions  $\{g_t\}_{t=1,\ldots,n}$  is nondecreasing and by Proposition 1 conditions (a)-(c) hold. By definition,  $g_t(k_{tj}) = \Psi_{X_t}(k_{tj}) = C_{tj}$  for all  $(k_{tj}, C_{tj}) \in \mathcal{S}$ .

Designate by

$$S_t := \{ (k_{lj}, C_{lj}) \mid (k_{lj}, C_{lj}) \in S, \ l = t \}$$
(6)

the set of options of maturity t and by

$$\mathcal{S}_{\geq t} := \cup_{l \geq t} \mathcal{S}_l \tag{7}$$

the set of options of maturities t and higher. Let also

$$S_{\geq t}^{\infty} := S_{\geq t} \cup \{(0, +\infty)\} \cup \{(+\infty, 0)\}.$$
(8)

Finally, let  $\partial S_{\geq t} : \mathbb{R}_+ \longrightarrow \mathbb{R}_+$  represent the lower boundary of the convex hull of  $S_{>t}^{\infty} \cup \{(0, X_0)\}$ , that is

$$\partial \mathcal{S}_{\geq t}(x) = \min\left\{ y | (x, y) \in \text{convex hull of } \{ \mathcal{S}_{\geq t}^{\infty} \cup (0, X_0) \} \right\}.$$
(9)

**Theorem 4.** The set S satisfies the no-arbitrage condition if and only if

(a)  $C_{tj} \ge X_0 - k_{tj}$  for each  $(k_{tj}, C_{tj}) \in S$ ;

(b) for each t, none of the points of  $\mathcal{S}_t$  is in the interior of the convex hull of  $\mathcal{S}_{\geq t}^{\infty}$ .

*Proof.* " $\implies$ " Define functions  $\tilde{g}_t(k) : \mathbb{R}_+ \longrightarrow \mathbb{R}_+$  for each  $t = 1, \ldots, n$ , such that  $\tilde{g}_t(k) = \partial S_{\geq t}(k) \ \forall k \geq 0$ . Then the sequence of functions  $\{\tilde{g}_t\}$  satisfies all the properties

of Proposition 3 except of property (c). Each function  $\tilde{g}_t$  is piecewise linear with slopes of all, but the last, line segments being negative. The last line segment has a slope zero. Let  $x_t := \min \{x \mid x \in \mathbb{R}_+, \ \tilde{g}'_t(x) = 0\}$ . Denote by **s** the set of all slopes of functions  $\tilde{g}_t$ ,  $t = 1, 2, \ldots, n$ , and let  $s^* := \max \{s \mid s \in \mathbf{s}, s < 0\}$ . Define the function

$$g_t(x) := \begin{cases} \tilde{g}_t(x), & x \le x_t, \\ \tilde{g}_t(x_t) + s^*(x - x_t), & \{x \ge x_t \mid g_t(x_t) + s^*(x - x_t) \ge 0\}, \\ 0, & \text{otherwise.} \end{cases}$$

Then the sequence of function  $g_t(x)$  satisfies all the properties of Proposition 3 and, thus, the martingale exists.

"  $\Leftarrow$ " If  $X_1, X_2, \ldots, X_n$  is a martingale, satisfying all the points of S, then for any (k, C) which belongs to the convex hull of  $S_{\geq 1}$ , it must be that  $C \geq \Psi_{X_1}(k) \geq X_0 - k$  by Proposition 1 and Theorem 3. Assume that there exists a point  $(k_{t^*j^*}, C_{t^*j^*})$ ,  $t^* \in \{1, \ldots, n\}, j^* \in 1, \ldots, U(t^*)$ , which is in the interior of the convex hull of  $S_{\geq t^*}^\infty$  (Figure 4.) Take points  $(k^l, C^l) \in S_{\geq t}^\infty$ ,  $(k^r, C^r) \in S_{\geq t}^\infty$  such that  $\partial S_{\geq t^*}(k^l) = C^l$  and  $\partial S_{\geq t^*}(k^r) = C^r$  and  $k^l$  is the closest to  $k_{t^*j^*}$  from the left, while  $k^r$  is the closest to  $k_{t^*j^*}$  from the right. Then  $(k_{t^*j^*}, C_{t^*j^*})$  lies strictly above the interval connecting  $(k^l, C^l)$  and  $(k^r, C^r)$ . Since  $\Psi_{X_{t^*}} \leq \Psi_{X_{t^*+1}} \leq \ldots \leq \Psi_{X_n}$  by Theorem 3, then  $\Psi_{X_{t^*}}(k^l) \leq C^l$  and  $\Psi_{X_{t^*}}(k^l)$  and  $(k^r, \Psi_{X_{t^*}}(k^r))$ , which contradicts to the condition that  $\Psi_{X_t^*}$  is a convex function.



Figure 4: The point  $(k_{t^*j^*}, C_{t^*j^*}), t^* \in \{1, \ldots, n\}, j^* \in \{1, \ldots, U(t^*) \text{ is in the interior of the convex hull of } \mathcal{S}_{>t^*}^{\infty}$ .

From the last theorem, we can find bounds on the price of a call option, given the set of priced options based on a geometric approach, see Bertsimas and Bushueva [2]. However, we will suggest another approach, which leads to the linear optimization problem, and that is extendable to multiple dimensions.

#### **Restriction to Markov martingales**

The idea of our approach is to reduce the class of martingale distributions over which we maximize (or minimize) as much as possible, while making sure that the optimal solution does not change. In this section, we show that it is enough to consider martingales with the Markov property only.

**Proposition 4.** Let  $X_1, X_2, \ldots, X_n$  be a martingale with values in  $\mathbb{R}^d$  and let  $\mu_i$  be the marginal distribution of  $X_i$ . Then, there exists a martingale with the Markov property  $\tilde{X}_1, \tilde{X}_2, \ldots, \tilde{X}_n$ , such that the marginal distribution of  $\tilde{X}_i$  is  $\mu_i$  for all  $i = 1, 2, \ldots, n$ .

*Proof.* For specifying the distribution of a discrete time Markov process, it is enough to specify the transition probabilities as well as the initial distribution.

Let  $\nu = \mathcal{L}(X_1)$ ,  $\mu_{1,2} = \mathcal{L}(X_2|X_1)$ ,...,  $\mu_{n-1,n} = \mathcal{L}(X_n|X_{n-1})$ . Then by the existence theorem for Markov processes due to Kolmogorov [13, p. 120] there exists a Markov process  $\tilde{X}_1, \ldots, \tilde{X}_n$  such that  $\mathcal{L}(X_1) = \nu$ ,  $\mathcal{L}(\tilde{X}_2|\tilde{X}_1) = \mu_{1,2}, \ldots, \mathcal{L}(\tilde{X}_{n-1}|\tilde{X}_n) = \mu_{n-1,n}$ .

It remains to show that the so-defined process  $\tilde{X}_1, \tilde{X}_2, \ldots, \tilde{X}_n$  is also a martingale. Let  $\sigma_k$  be the smallest  $\sigma$ -algebra for which  $\tilde{X}_1, \tilde{X}_2, \ldots, \tilde{X}_k$  are all measurable. We need to prove that  $E[\tilde{X}_{k+1}|\sigma_k] = \tilde{X}_k$ . From the Markov property of  $\tilde{X}_1, \ldots, \tilde{X}_k$ , it follows that  $E[\tilde{X}_{k+1}|\sigma_k] = E[\tilde{X}_{k+1}|\tilde{X}_k]$ . By definition  $E[\tilde{X}_{k+1}|\tilde{X}_k] = E[X_{k+1}|X_k]$ .

On the other hand, since  $X_1, X_2, \ldots, X_n$  is a martingale, we have

 $E[E[X_{k+1}|X_k, X_{k-1}, \dots, X_1]|X_k] = E[X_k|X_k] = X_k. \text{ Also}$  $E[E[X_{k+1}|X_k, X_{k-1}, \dots, X_1]|X_k] = E[X_{k+1}|X_k]. \text{ Thus } E[X_{k+1}|X_k] = X_k. \text{ Consequently, } E[\tilde{X}_{k+1}|\tilde{X}_k] = \tilde{X}_k \text{ and } E[\tilde{X}_{k+1}|\sigma_k] = \tilde{X}_k.$ 

**Definition 2.** A stochastic process which is a martingale and a Markov process will be called a Markov martingale.

**Corollary 1.** If we restrict our consideration to Markov martingales only, the optimal solution to our problems does not change.

## 3 The One Dimensional Case

In this section, we address Problem (b), i.e., we find best possible upper and lower bounds on the price of a European style option with a continuous piecewise linear payoff function, given the set of European call options of different maturities.

The price of a European call option with a strike price k and maturity t is given by Eq. (1). The problem we address in this section can be formulated as follows: given a continuous piecewise linear function  $g : \mathbb{R}_+ \to \mathbb{R}_+$  and a set

$$\mathcal{S} := \{ (k_{ij}, C_{ij}, t_i) \mid \{ k_{ij}, C_{ij}, t_i \} \in \mathbb{R}_+, \ t_1 < t_2 \dots < t_n, \\ i \in \{ 1, 2, \dots, n \}, \ j \in \{ 1, 2, \dots, U(i) \} \}$$

find the domain of values of

$$e^{-\int_{0}^{t^{*}} r(s)ds} E\left[g\left(e^{\int_{0}^{t^{*}} r(s)ds} X_{t^{*}}\right)\right]$$

under the condition, that  $\{X_{t_1}, X_{t_2}, \dots, X_{t^*}, \dots, X_{t_n}\}$  is a nonnegative martingale such that

$$E\left[\left(X_{t_i} - k_{ij}e^{-\int_0^{t_i} r(s)ds}\right)^+\right] = C_{ij}$$

It is assumed that such martingale exists, that is the set of given options satisfies the no-arbitrage condition.

#### **Remarks:**

(a) As before, we can assume  $r(s) \equiv 0$ , since we can take  $\tilde{k}_{ij} = k_{ij}e^{-\int_0^{t_i} r(s)ds}$  instead of  $k_{ij}$ , and rewrite all the equalities as  $E\left[\left(X_{t_i} - \tilde{k}_{ij}\right)^+\right] = C_{ij}$ . Also, we can take the function

$$\tilde{g}(X_{t^*}) := e^{-\int_0^{t^*} r(s)ds} g\left(e^{\int_0^{t^*} r(s)ds} X_{t^*}\right)$$

instead of  $g(X_{t^*})$ . Notice that if  $g(X_{t^*})$  is a continuous piecewise linear function, so is  $\tilde{g}(X_{t^*})$ .

- (b) Thus, again, only the order of  $t_1, t_2, \ldots, t_n$  is important, but not the specific values that the sequence takes. Thus we can assume  $\{t_1, t_2, \ldots, t_n\} \equiv \{1, 2, \ldots, n\}$  for convenience.
- (c) Recall that we must have  $E[X_t] = X_0$ , where  $X_0$  is the current price of the stock.

In light of the above remarks, Problem (b) can be reformulated as follows:

#### Problem (b)

Given a continuous piecewise linear function  $g: \mathbb{R}_+ \to \mathbb{R}_+$  and an ordered set

$$\mathcal{S} := \left\{ (k_{tj}, C_{tj}) \mid (k_{tj}, C_{tj}) \in \mathbb{R}^2_+, \ t \in \{1, 2, \dots, n\}, \ j \in \{1, 2, \dots, U(t)\} \right\}$$
(10)

find the domain of values of

$$E\left[g(X_{t^*})\right] \tag{11}$$

for some  $t^* \in \{1, 2, ..., n\}$ , under the condition, that  $\{X_1, ..., X_{t^*}, ..., X_n\}$  is a nonnegative martingale such that

$$E\left[\left(X_t - k_{tj}\right)^+\right] = C_{tj} \text{ for all } (k_{tj}, C_{tj}) \in \mathcal{S},$$
(12)

and

$$E[X_t] = X_0$$
 for all  $t = 1, \ldots, n$ .

#### **3.1** Treatment of future conditions

Let

$$\mathcal{F} = \{ (k_{tj}, C_{tj}) | t^* < t \le n, \ (k_{tj}, C_{tj}) \text{ is a vertex of the convex hull of } \mathcal{S}_{>t^*}^{\infty} \}$$

represent future points which are on the border of the convex hull of  $S_{\geq t^*}$ . And let  $(k_{fut,j}, C_{fut,j}), j = 1, 2, \ldots, U(f_{ut})$ , be the elements of  $\mathcal{F}$ . Then we have the following lemma:

Lemma 1. If we reduce conditions (12) in Problem (b) to the set of conditions

$$E\left[\left(X_{t}-k_{tj}\right)^{+}\right] = C_{tj}, \quad t \le t^{*}, \ j = 1, 2..., U(t),$$
(13)

$$E\left[\left(X_{t^*} - k_{fut,j}\right)^+\right] \le C_{fut,j}, \quad j = 1, 2, \dots, U(f_{ut}),$$
 (14)

the solution to Problem (b) will not change.

*Proof.* Suppose that we want to maximize  $E[g(X_{t^*})]$  and  $g_1^*$  is the optimal solution under conditions (12) and  $g_2^*$  is the optimal solution under conditions (13) and (14). Then  $g_1^* \leq g_2^*$ . Let us show that  $g_1^* \geq g_2^*$ .

Let  $\Pi^*$  be the distribution that maximizes  $\int_0^{+\infty} g(x) d\Pi(x)$  over distributions  $\Pi$  of  $X_{t^*}$ under conditions (13)& (14). Then there exists a martingale  $X_1, \ldots, X_{t^*}$  which satisfies the part of conditions (12) for  $t \leq t^*$  and with the distribution  $\Pi^*$  of  $X_{t^*}$ , such that  $\Psi_{\Pi^*}(k_{fut,j}) \leq C(k_{fut,j})$  for all  $j = 1, \ldots, U(f_{ut})$ . Since  $\Psi_{\Pi^*}$  has the properties stated in Proposition 1, none of the points  $(k, \Psi_{\Pi^*}(k)), k \geq 0$ , is in the interior of the convex hull of  $S_{\geq t^*}^{\infty}$  and  $\Psi(k) \leq X_0 - k$ . Then, since also the set  $S_{\geq t^*}$  by itself satisfies the no-arbitrage condition, by Theorem 4 there exists a martingale  $\tilde{X}_t^*, \ldots, \tilde{X}_{t_n}$  which satisfies the set  $\{S_{\geq t^*}\} \cup \{(k, \Psi_{\Pi^*}(k))\}_{k\geq 0}$ . Notice that the distribution of  $\tilde{X}_{t^*}$  is!  $\Psi_{\Pi^*}$ . As in the proof of Proposition 4, it can be shown that the martingale  $X_1, X_2, \ldots, X_{t^*}$  can be extended to the martingale  $X_1, \ldots, X_{t^*}, \ldots, X_n$  by specifying transition probabilities  $\mathcal{L}(X_{t^*+1}|X_{t^*}) = \mathcal{L}(\tilde{X}_{t^*+1}|\tilde{X}_{t^*}), \ldots, \mathcal{L}(X_n|X_{n-1}) = \mathcal{L}(\tilde{X}_n|\tilde{X}_{n-1})$ . This extended martingale satisfies all the conditions (12). Thus  $g_1^* \geq g_2^*$ .

#### **3.2** Restriction to discrete distributions

From now on we make an additional assumption that

$$P(X_i \le L, i = 1, \dots, t^*) = 1.$$

for some L > 0. Let

$$\mathcal{K}_{S} = \{0\} \cup \{L\} \cup \{k_{tj} \mid (k_{tj}, C_{tj}) \in \mathcal{S}, t \le t^{*}\} \cup \{k_{tj} \mid (k_{tj}, C_{tj}) \in \mathcal{F}\}$$

and

$$\mathcal{K}_g = \{k \in [0, L] \mid g'(k-) \neq g'(k+)\}$$

be the set of points, where g changes its derivative. Define:

$$\mathcal{K} = \mathcal{K}_S \cup \mathcal{K}_g.$$

**Theorem 5.** Let  $X_1, \ldots, X_{t^*}$  be a nonnegative Markov martingale with support in [0, L]. Then there exists a martingale  $\tilde{X}_1, \ldots, \tilde{X}_{t^*}$  with support in  $\mathcal{K}$ , such that

$$E\left[\left(X_{t}-k_{tj}\right)^{+}\right] = E\left[\left(\tilde{X}_{t}-k_{tj}\right)^{+}\right], \quad t = 1, 2, \dots, t^{*}; \quad j = 1, 2, \dots, U(t)$$
$$E\left[\left(X_{t^{*}}-k_{fut,j}\right)^{+}\right] = E\left[\left(\tilde{X}_{t^{*}}-k_{fut,j}\right)^{+}\right], \quad j = 1, 2, \dots, U(f_{ut})$$

and

$$E\left[g(X_{t^*})\right] = E\left[g(\tilde{X}_{t^*})\right].$$

We will use the following two lemmas to prove Theorem 5.

**Lemma 2.** Let  $X_1, X_2$  be a martingale with values in  $\mathbb{R}^d$ . Then there exists a continuous time martingale  $\{\tilde{X}_t\}_{t\in[1,2]}$  with continuous paths, such that  $\tilde{X}_1 = X_1$  and  $\tilde{X}_2 = X_2$ .

*Proof.* By the martingale version of Skorohod embedding theorem [13, p.229], there exists a Brownian motion B and optional time  $\tau \geq 0$ , such that  $X_2 - X_1 = B_{\tau}$  a.s. Let  $M_t = X_1 + B_{t \wedge \tau}$ . Then by optional stopping theorem  $M_t$  is a martingale. This martingale is bounded and, thus, uniformly integrable. Consequently, there exists  $M_{\infty}$ 

such that  $\{M_t\}_{t \in [0, +\infty]}$  is a martingale and  $M_t$  converges to  $M_\infty$  almost surely by Doob's theorem [9, p.285].

Let  $h: [0, +\infty] \to [1, 2]$  be a nondecreasing continuous function and let  $\overline{M}_t \equiv M_{h(t)}$ ,  $t \in [1, 2]$ . Then  $\{\overline{M}_1, \overline{M}_2\}$  has the same joint distribution as  $\{X_1, X_2\}$  and  $\{\overline{M}_t\}_{t \in [1, 2]}$  is a continuous time martingale with continuous paths.

**Lemma 3.** Let  $\{X_t\}_{t\in[1,2]}$  be a continuous time martingale with values in  $\mathbb{R}^d$ . Let  $h: \mathbb{R}^d \to \mathbb{R}$ , be a continuous piecewise linear function. Let  $\mathbb{R}^d = \bigcup_i D_i$ , and h is affine on each  $D_i$ , i = 1, 2, ..., n. Then for each i

$$E\left[h(X_1)|\{X_t\}_{t\in[1,2]}\in D_i\right] = E\left[h(X_2)|\{X_t\}_{t\in[1,2]}\in D_i\right].$$

*Proof.* The result immediately follows from conditional Jensen's inequality applied to h and -h restricted to  $D_i$  [9, p.277].

Now let us return to the proof of Theorem 5:

Proof. We start at time  $t^*$  and proceed recursively in the following way. Let  $B_t, t \in [0, +\infty)$ , be a Brownian motion starting at 0. Let  $\dot{X}_t = X_{t^*} + B_{t-t^*}$  for all  $t \geq t^*$ . Then  $X_1, X_2, \ldots, X_{t^*}, \dot{X}_t$  is a martingale. If  $\tau_1$  is the first hitting time of  $\dot{X}_t$  on  $\mathcal{K}$ , then by the optional stopping theorem  $X_1, X_2, \ldots, X_{t^*}, \dot{X}_{(t^*+1)\wedge\tau_1}, \dot{X}_{(t^*+2)\wedge\tau_1}, \ldots$  is also a martingale. Since this martingale is bounded, it is uniformly integrable, and thus [9, p.283, p.285] there exists  $\dot{X}_{\infty}$ , such that  $E\left[\dot{X}_{\infty}|\mathcal{B}_n\right] = \dot{X}_n$  for all  $n = t^*, t^* + 1, \ldots$ , and  $\dot{X}_n$  converges to  $\dot{X}_{\infty}$  a.s. Let  $\tilde{X}_{t^*} = \dot{X}_{\infty}$ . Then  $X_1, X_2, \ldots, X_{t^*-1}, \tilde{X}_{t^*}$  is a martingale and  $P\left(\tilde{X}_{t^*} \in \mathcal{K}\right) = 1$ .

Now let us make the discrete time martingale  $X_1, X_2, \ldots, X_{t^*-1}, \tilde{X}_{t^*}$  be continuous on  $[t^*-1, t^*]$  and have continuous paths. That is, by Lemma 2 we can define the martingale  $\left\{\dot{X}_t\right\}_{t\in[t^*-1,t^*]}$ , such that  $\dot{X}_{t^*-1} = X_{t^*-1}, \dot{X}_{t^*} = \tilde{X}_{t^*}$  and it is continuous. Let  $\tau_2$  be the stopping time which stops the martingale when it first hits  $\mathcal{K}$  after  $t^* - 1$ . Then  $\tau_2$  is bounded and  $X_{t^*-1}, X_{\tau_2}$  is a martingale by the optional stopping theorem. By Lemma 3,  $X_{t^*-1}\left(=\dot{X}_{t^*-1}\right)$  and  $\dot{X}_{\tau_2}$  takes the same values on functionals but  $P\left(\dot{X}_{\tau_{t^*-1}} \in \mathcal{K}\right) = 1$ . Since  $X_1, X_2, \ldots, X_{t^*-1}, \dot{X}_{\tau_2}, \tilde{X}_{t^*}$  is a martingale, so is  $X_1, X_2, \ldots, X_{t^*-2}, \dot{X}_{\tau_2}, \tilde{X}_{t^*}$ . We let  $\tilde{X}_{t^*-1} = \dot{X}_{\tau_2}$  and proceed in this way until time 1.

We next introduce Problem (b') that is equivalent to Problem (b).

#### Problem (b')

Find the domain of values of

 $E\left[g(X_{t^*})\right]$ 

under the condition, that  $\{X_1, \ldots, X_{t^*}\}$  is a nonnegative martingale with support in  $\mathcal{K} := \mathcal{K}_{\mathcal{S}} \cup \mathcal{K}_g$  such that

$$E\left[\left(X_{t} - k_{tj}\right)^{+}\right] = C_{tj}, \quad t \le t^{*}, \ j = 1, 2..., U(t),$$

$$E\left[\left(X_{t^{*}} - k_{fut,j}\right)^{+}\right] \le C_{fut,j}, \quad j = 1, 2, ..., U(f_{ut}).$$
(15)

From Lemma 1 and Theorem 5 we obtain

**Corollary 2.** If the price of the stock is bounded by L, as in (15), then Problem (b) and Problem (b') have the same solutions.

**Proposition 5.** If there is no bound on the price of the stock, then as  $L \to \infty$ , the optimal solution of Problem (b') converges monotonically to the optimal solution of Problem (b).

Proof. Notice, that since g is a continuous piecewise-linear function,  $E[g(X_{t^*})]$  is always finite once  $E[X_{t^*}]$  is finite. Thus, the best possible upper bound on  $E[g(X_{t^*})]$ ,  $C^*$ , for Problem (b) is finite. If  $C_L^*$  is the best possible upper bound on  $E[g(X_{t^*})]$  for Problem (b'), then  $C_L^* \leq C^*$  and  $C_L^*$  approaches monotonically  $C^*$  as  $L \to \infty$  according to Corollary 2. Thus  $C_L^* \to C^*$  as  $L \to \infty$ . The same is true for the lower bound.

#### 3.3 Algorithm for the one dimensional case

We have shown that in order to solve problem (b), it is enough to consider Markov martingales with support in  $\mathcal{K}$ . The distribution of a discrete time Markov process is fully determined by joint distributions of each two consecutive states.

Let  $\mathcal{K} = \{k_1, k_2, \dots, k_n, \dots, k_N\}$  and  $\mathcal{N}$  denote the set of all the indexes  $1, 2, \dots, N$ . Then the distribution of the Markov process with support in  $\mathcal{K}$  is determined by the collection of

$$P_{t;\mathfrak{n}_1,\mathfrak{n}_2} = P\left(X_t = k_{\mathfrak{n}_1}, X_{t+1} = k_{\mathfrak{n}_2}\right), \ t = 1, \dots, t^*, \ \mathfrak{n}_1, \mathfrak{n}_2 \in \mathcal{N}.$$

Since most of the constraints that we consider are constraints on marginal distributions of the process, for convenience we also introduce

$$P_{t;\mathfrak{n}} = P\left(X_t = k_{\mathfrak{n}}\right) \quad t = 1, \dots, t^*, \quad \mathfrak{n} \in \mathcal{N}.$$

For P to be a measure, it must satisfy the countable additivity condition:

$$\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n}_1,\mathbf{n}} = P_{t;\mathbf{n}_1} \quad \text{for all } t \le t^*, \ \mathbf{n}_1 \in \mathcal{N},$$

$$\sum_{\mathfrak{n} \in \mathcal{N}} P_{t;\mathfrak{n},\mathfrak{n}_1} = P_{t+1;\mathfrak{n}_1} \quad \text{for all } t < t^*, \ \mathfrak{n}_1 \in \mathcal{N},$$

and nonnegativity condition:

$$P_{t;\mathfrak{n}} \ge 0, \quad P_{t;\mathfrak{n}_1,\mathfrak{n}_2} \ge 0, \quad \text{for all } t < t^*, \ \mathfrak{n}, \mathfrak{n}_1, \mathfrak{n}_2 \in \mathcal{N}.$$

In addition, for the probability measure we must have

$$\sum_{\mathfrak{n}\in\mathcal{N}}P_{1;\mathfrak{n}}=1.$$

The martingale condition can be written as

$$\sum_{\mathfrak{n}\in\mathcal{N}} P_{t;\mathfrak{n}_1,\mathfrak{n}}k_{\mathfrak{n}} = P_{t;\mathfrak{n}_1}k_{\mathfrak{n}_1} \quad \text{for all } t \leq t^*, \ \mathfrak{n}_1 \in \mathcal{N}$$

For the martingale to satisfy prices of options with maturities less than or equal to  $t^*$ , we must have:

$$\sum_{\mathfrak{n}\in\mathcal{N}} P_{t;\mathfrak{n}}(k_{\mathfrak{n}} - k_{tj})^+ = C_{tj} \quad \text{for all } t \le t^*, \ j = 1, 2..., U(t)$$

As was proved in Theorem 5, for prices of options of maturities greater than  $t^*$  to be satisfied, it is enough to have

$$\sum_{\mathfrak{n}\in\mathbb{N}} P_{t^*;\mathfrak{n}}(k_\mathfrak{n} - k_{fut,j})^+ \le C_{fut,j} \quad \text{for all } j = 1, 2..., U(f_{ut}).$$

Thus the problem of determining best possible upper (lower) bounds on the price of an option with maturity  $t^*$  and a payoff function g, consists of finding the collection of  $P_{t;n}$  and  $P_{t;n_1,n_2}$ , such that

$$\begin{aligned} \max(\min) \quad &\sum_{\mathbf{n}\in\mathcal{N}} P_{t^*;\mathbf{n}} \ g(k_{\mathbf{n}}) \\ \text{s.t.} \quad &\sum_{\mathbf{n}\in\mathcal{N}} P_{1;\mathbf{n}} = 1, \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n}_1,\mathbf{n}} = P_{t;\mathbf{n}_1} \quad \text{for all } t \leq t^*, \ \mathbf{n}_1 \in \mathcal{N}, \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n},\mathbf{n}_1} = P_{t+1;\mathbf{n}_1} \quad \text{for all } t < t^*, \ \mathbf{n}_1 \in \mathcal{N}, \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n}_1,\mathbf{n}}k_{\mathbf{n}} = P_{t;\mathbf{n}_1}k_{\mathbf{n}_1} \quad \text{for all } t < t^*, \ \mathbf{n}_1 \in \mathcal{N}, \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n}}(k_{\mathbf{n}} - k_{tj})^+ = C_{tj} \quad \text{for all } t \leq t^*, \ j = 1, 2 \dots, U(t), \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t^*;\mathbf{n}}(k_{\mathbf{n}} - k_{fut,j})^+ \leq C_{fut,j} \quad \text{for all } j = 1, 2 \dots, U(f_{ut}), \\ &P_{t;\mathbf{n}_1,\mathbf{n}_2} \geq 0 \quad \text{for all } t \leq t^*, \ \mathbf{n}_1, \mathbf{n}_2 \in \mathcal{N}. \end{aligned}$$

This is a linear optimization problem with  $O(N^2)$ , where N is the number of elements of  $\mathcal{K}$ .

## 4 The Two Dimensional Case

In this section, we address Problem (c), i.e., the problem of determining bounds on options on two different assets if prices of options on individual assets are known.

Let  $\{S^{I}(t)\}_{t\geq 0}$  be a stochastic process describing the price process of the first asset, and  $\{S^{II}(t)\}_{t\geq 0}$  that of the second asset. If  $(k_{ij}^{h}, C_{ij}^{h}, t_{i})$ ,  $i = 1, \ldots, n, j = 1, 2, \ldots, U(t, h)$ , represent the set of options on asset  $S^{h}$ , with  $t_{i}$  being the maturity of an option,  $k_{ij}^{h}$  its strike price, and  $C_{ij}^{h}$  its price, then we must have

$$C_{ij}^{h} = e^{-\int_{0}^{t_{i}} r(s)ds} E_{Q} \left[ \left( S^{h}(t_{i}) - k_{ij}^{h} \right)^{+} \right] = E_{Q} \left[ \left( X_{t_{i}}^{h} - k_{ij}^{h} e^{-\int_{0}^{t_{i}} r(s)ds} \right)^{+} \right], \quad (16)$$

where  $\{X_t^h\}_{t\geq 0}$ , is a martingale under Q. As in the single dimensional case, we can assume without loss of generality that the risk-free rate r(s) can be assumed 0. Thus, we can change times  $t_i$  to their indexes.

Let  $\mathcal{T} := \{(h, t, j) | h \in \{I, II\}, t \in \{1, 2, \dots, n\}, j = 1, 2, \dots, U(t, h)\}$ . We define a family of measurable functions  $f_{tj}^h : \mathbb{R}^2_+ \to \mathbb{R}_+$  as

$$f_{tj}^{h}(\mathbf{X}_{t}) = \left(X_{t}^{h} - k_{tj}^{h}\right)^{+}, \ (h, t, j) \in \mathcal{T}.$$
(17)

Then the problem can be formulated as follows:

#### Problem (c)

Given the set  $(k_{tj}^h, C_{tj}^h)$ ,  $(h, t, j) \in \mathcal{T}$ , and a continuous piecewise linear function  $g : \mathbb{R}^2 \to \mathbb{R}$ , find the maximal and minimal possible values of  $E[g(\mathbf{X}_{t^*})]$ , for a given  $t^* \in \{1, 2, ..., n\}$ , under the condition that  $\mathbf{X}_1, \mathbf{X}_2, ..., \mathbf{X}_n$  is a two dimensional nonnegative martingale, such that

$$E\left[f_{tj}^{h}(\mathbf{X}_{t})\right] = C_{tj}^{h}, \ (h, t, j) \in \mathcal{T}$$

$$\tag{18}$$

**Definition 3.** We call function  $g : \mathbb{R}^d \to \mathbb{R}$  piecewise-linear, if there exists a partition of  $\mathbb{R}^d$  by a finite number of hyperplanes (lines if d = 2) into nonoverlapping subsets (regions if d = 2), such that in the interior of each subset g is an affine function.

Notice, that by Proposition 4 we might restrict our consideration to Markov martingales only.

#### 4.1 Treatment of future conditions

Notice that the function  $f_{tj}^h$  depends only on the *h*-coordinate of  $\mathbf{X}_t$ . With this property of f-constraints, we can show that future conditions have a "simple effect", in the sense

that conditions (18) for  $t > t^*$  can be reduced to separate conditions on the first and second coordinates of  $\mathbf{X}_{t^*}$ . Let

$$\begin{split} \mathcal{S}^{h}_{\geq t^{*}} &= \left\{ \left( k^{h}_{tj}, C^{h}_{tj} \right) \middle| t^{*} \leq t \leq n, t \in \{1, 2, \dots, U(t, h)\} \right\}, \ h =_{I,II} \\ \mathcal{S}^{h\infty}_{\geq t^{*}} &= \mathcal{S}^{h}_{\geq t^{*}} \cup \{(0, +\infty), (+\infty, 0)\}, \ h =_{I,II}, \end{split}$$

and

$$\mathcal{F}^{h} = \left\{ \left( k_{tj}^{h}, C_{tj}^{h} \right) | t^{*} < t \le n, \ \left( k_{tj}^{h}, C_{tj}^{h} \right) \text{ is a vertex of the convex hull of } \mathcal{S}_{\ge t^{*}}^{h\infty} \right\}$$

represent future points which are on the border of the convex hull of  $\mathcal{S}_{>t^*}^{h\infty}$ . Let

$$\left(k_{fut,j}^h, C_{fut,j}^h\right), \quad j = 1, 2, \dots, U(_{fut}, h)$$

be enumerated elements of  $\mathcal{F}^h$ .

In formulating all further results we will assume that  $\begin{pmatrix} k_{tj}^{I}, C_{tj}^{I} \end{pmatrix}$ ,  $({}_{I}, t, j) \in \mathcal{T}$  and  $\begin{pmatrix} k_{tj}^{II}, C_{tj}^{II} \end{pmatrix}$ ,  $({}_{II}, t, j) \in \mathcal{T}$  separately do satisfy the no-arbitrage condition, that is that there exist one-dimensional martingales  $X_{t}^{I}$  and  $X_{t}^{II}$  such that

$$E\left[\left(X_t^h - k_{tj}^h\right)^+\right] = C_{tj}^h, \quad h = I, II$$

Notice that this condition ensures that the two dimensional martingale which satisfies conditions (18) also exists.

**Theorem 6.** Let  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{t^*}$  be a two-dimensional martingale satisfying  $E\left[f_{tj}^h(\mathbf{X}_t)\right] = C_{tj}^h$ , for all  $\{(h, t, j) \in \mathcal{T} \mid t \leq t^*\}$ . If also for each  $h \in \{I, II\}$ 

$$E\left[\left(X_{t^*}^h - k_{fut,j}^h\right)^+\right] \le C_{fut,j}^h$$

is satisfied for all  $(k_{fut,j}^h, C_{fut,j}^h) \in \mathcal{F}^h$ , then there exist two-dimensional random variables  $\mathbf{X}_{t^*+1}, \ldots, \mathbf{X}_n$  such that

- (a)  $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}, \mathbf{X}_{t^*+1}, \ldots, \mathbf{X}_n$  is a martingale;
- (b) all the conditions  $E\left[f_{ti}^{h}(\mathbf{X}_{t})\right] = C_{ti}^{h}$  are satisfied for all  $(h, t, j) \in \mathcal{T}$ .

*Proof.* As in the proof of Lemma 1, we can show that for each  $h \in \{I, II\}$  there exists a one-dimensional martingale  $\tilde{X}_{t^*}^h, \tilde{X}_{t^*+1}^h, \ldots, \tilde{X}_n^h$ , such that it satisfies the set  $(k_{tj}^h, C_{tj}^h), t \geq t^*, j = 1, \ldots, U(t, h)$  and  $\Psi_{\tilde{X}_{t^*}}^h = \Psi_{X_{t^*}}^h$ .

We will extend the martingale  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{t^*}$  to the martingale  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{t^*}, \mathbf{X}_{t^*+1}, \dots, \mathbf{X}_n$ , such that  $\mathbf{X}_{t^*}, \mathbf{X}_{t^*+1}, \dots, \mathbf{X}_n$  is a Markov process. To define a joint distribution of  $\mathbf{X}_t$  and  $\mathbf{X}_{t+1}$  for  $t \geq t^*$ , allow each coordinate to evolve independently of each other. That is, for each  $h =_{I,II}$  define  $\mathcal{L}(X_t^h|X_{t+1}^h)$ , as  $\mathcal{L}(\tilde{X}_t^h|\tilde{X}_{t+1}^h)$  for  $t = t^*, t^* + 1, \dots, n$ .

Theorem 6 allows us to reduce "future conditions" to conditions at time  $t^*$ . This is a significant simplification of the problem, since difficulties arise from the requirement that the martingale satisfies  $E\left[f_{tj}^h(\mathbf{X}_t)\right] = C_{tj}^h, (h, t, j) \in \mathcal{T}$ .

**Corollary 3.** Our main problem is equivalent to the following: find the maximum (minimum) of  $E[g(\mathbf{X}_{t^*})]$  under the condition that  $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}$  is a Markov Martingale such that

(a) 
$$E\left[\left(X_t^h - k_{tj}^h\right)^+\right] = C_{tj}^h$$
, for all  $t \le t^*$ ,  $j \le U(t,h)$  and  
(b)  $E\left[\left(X_{t^*}^h - k_{fut,j}^h\right)^+\right] \le C_{fut,j}^h$  for all  $\left(k_{fut,j}^h, C_{fut,j}^h\right) \in \mathcal{F}^h$ 

**Definition 4.** Let  $f_{fut,j}^h : \mathbb{R}^2_+ \to \mathbb{R}_+$ ,  $f_{fut,j}^h(\mathbf{X}) = \left(X^h - k_{fut,j}^h\right)^+$ , be a family of functions, where each function corresponds to some  $k_{fut,j}^h$ , such that  $\left(k_{fut,j}^h, C_{fut,j}^h\right) \in \mathcal{F}^h$ , h = I, II.

Constraints from the past introduce much more difficulty to the problem than future constraints. It is not possible to reduce past martingale conditions to separate conditions on  $X_{t^*}^{I}$  and  $X_{t^*}^{II}$ . In other words, for a given distribution of  $\mathbf{X}_{t^*}$ , and two sequences of marginal laws  $\{\Pi_t\}_{t=1,...,t^*}$  and  $\{Q_t\}_{t=1,...,t^*}$  of one dimensional processes, such that  $\Pi_{t^*} = \mathcal{L}\left(X_{t^*}^{I}\right)$  and  $Q_{t^*} = \mathcal{L}\left(X_{t^*}^{II}\right)$ , the existence of one-dimensional martingales with the above sequences of marginal laws, is not sufficient for the existence of a two-dimensional martingale  $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}$  such that  $\mathcal{L}\left(X_t^{I}\right) = \Pi_t$  and  $\mathcal{L}\left(X_t^{II}\right) = Q_t$ for  $t = 1, \ldots, t^*$ .

**Counterexample**. Suppose that  $\mathbf{X}_2$  has an atomic distributions with two atoms a and b, as shown on Figure 5, where each atom has weight of 1/2. Suppose that the marginal law of  $X_1^I$  is  $P\left(X_1^I = x_c\right) = 1$ , where  $x_c = \frac{1}{2}(x_a + x_b)$  and the marginal law of  $X_1^{II}$  is  $P\left(X_1^{II} = y_a\right) = P\left(X_1^{II} = y_b\right) = \frac{1}{2}$ . Obviously, we can define a joint distribution of  $X_1^I, X_2^I$ , so that it is a one-dimensional martingale. The same is true for  $X_1^{II}, X_2^{II}$ . However, it is not possible to define a joint distribution of  $\mathbf{X}_1, \mathbf{X}_2$ , so that it is a two-dimensional martingale.

Thus, the condition that  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_{t^*}$  is a martingale, is truly a "joint" condition on  $\{X_t^I\}$  and  $\{X_t^{II}\}$ .

#### 4.2 **Restriction to discrete distributions**

In what follows, function g is defined as  $g : \mathbb{R}^2_+ \to \mathbb{R}_+$ ,  $(x, y) \to |\alpha x + \beta y - k_g|^+$ . However, an analogous solution would apply if  $g : \mathbb{R}^2_+ \to \mathbb{R}$  is any continuous piecewise-linear function.

As in the one dimensional case we add an additional constraint that stock prices are bounded, that is, there exists L > 0, such that for h = I, II

$$P(X_1^h, X_2^h, \dots, X_n^h \le L) = 1.$$



Figure 5: Counterexample.

**Definition 5.** For each  $t = 1, 2, ..., t^* - 1$  we define Graph  $G_t$  (Figure 6), which is contained in  $[0, L] \times [0, L]$  and has vertices on intersections of the lines: (a)  $x = k_{tj}^I$ ,  $j \leq U(t, I)$ , (b)  $y = k_{tj}^{II}$ ,  $j \leq U(t, II)$ , (c) segments of Q: from (0, 0) to (0, L), (0, L) to (L, L), (L, L) to (L, 0), (0, 0) to (L, 0).

For  $t^*$  we define Graph  $G_{t^*}$  (Figure 7) which has vertices on intersections of the lines (a)  $x = k_{t^*j}^I$ ,  $j \le U(t, I)$ , (a')  $x = k_{fut,j}^I$ ,  $j \le U(_{fut,I})$ , (b)  $y = k_{t^*j}^{II}$ ,  $j \le U(t, II)$ , (b')  $y = k_{fut,j}^{II}$ ,  $j \le U(_{fut,II})$ , (c) segments of Q: from (0,0) to (0, L), (0, L) to (L, L), (L, L) to (L, 0), (0, 0) to (L, 0), (d)  $\alpha x + \beta y = k_g$ .

Each edge of  $G_t$  belongs to one of the corresponding lines (or segments). The intersection between two edges is either empty or consists of one vertex point.



Figure 6: The form of Graph  $G_t$  for  $t = 1, 2, \ldots, t^* - 1$ .

Figure 7: The form of Graph  $G_{t^*}$ .

Let

•  $G_t^V$  designate all the vertices of Graph  $G_t$ ,

- $G_t^E$  denote the set of points which lie on edges of  $G_t$  and
- $G_t^R$  denote regions of  $\mathbb{R}^2$  over which edges of Graph  $G_t$  splits  $[0, L] \times [0, L]$ .

Let us enumerate regions of  $G_t^R$ . We will thus write r for the region of  $G_t^R$  numbered r. Let  $\mathcal{R}_t$  be the set of all integers such that each of them represents some region of  $G_t^R$ and  $\mathcal{V}$  be the set of all integers such that each of them represents some vertex of  $G_{t^*}^N$ . Let  $\mathfrak{Reg}_t : [0, L] \times [0, L] \to \mathcal{R}_t$ , be the function which attributes to each point of  $[0, L] \times [0, L]$ the index of the region that contains it, and let  $\mathcal{V}er : [0, L] \times [0, L] \to \mathcal{V}$ , be the function which attributes to each vertex of  $G_{t^*}^V$  its index v. Since edges are contained by several regions, we have to specify which region corresponds to each edge.

**Definition 13.** We will say that a random process  $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}$  has property  $P_G$  if and only if all of the following hold

(a)  $P(\mathbf{X}_{t^*} \in G_{t^*}^V) = 1;$ 

(b)  $\mathbf{X}_1$  has an atomic distribution with at most one atom per region of  $G_1^R$ ;

(c) If  $r_1 \in \mathcal{R}_1, r_2 \in \mathcal{R}_2 \dots, r_t \in \mathcal{R}_t$  with  $t < t^*$ , then

$$\mathcal{L}\left(\mathbf{X}_{t+1} \left| \left[ \mathfrak{Reg}_t(\mathbf{X}_1), \mathfrak{Reg}_t(\mathbf{X}_2), \dots, \mathfrak{Reg}_t(\mathbf{X}_t) \right] = [r_1, r_2, \dots, r_t] \right)$$

has an atomic measure with at most one atom per region of  $G_t^R$ .

**Theorem 7.** Let  $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}$  be a nonnegative Markov Martingale bounded by L. Then there exists a nonnegative Markov Martingale  $\{\bar{\mathbf{X}}_1, \bar{\mathbf{X}}_2, \ldots, \bar{\mathbf{X}}_{t^*}\}$ , bounded by L, such that it has property  $P_G$  and satisfies

$$E\left[f_{tj}^{h}(\mathbf{X}_{t})\right] = E\left[f_{tj}^{h}\left(\bar{\mathbf{X}}_{t}\right)\right], \ t \leq t^{*}, \ (h, t, j) \in \mathcal{T},$$
$$E\left[f_{fut, j}^{h}(\mathbf{X}_{t^{*}})\right] = E\left[f_{fut, j}^{h}\left(\bar{\mathbf{X}}_{t^{*}}\right)\right], \ j = 1, 2, \dots, U(f_{ut}, h), \ h \in \{I, II\},$$
$$E\left[g\left(\mathbf{X}_{t^{*}}\right)\right] = E\left[g\left(\bar{\mathbf{X}}_{t^{*}}\right)\right] = E\left[g\left(\bar{\mathbf{X}}_{t^{*}}\right)\right].$$

*Proof.* For  $i = 1, ..., t^*$ , let  $\sigma_i$  designate the smallest  $\sigma$ -algebra for which  $\mathbf{X}_1, \mathbf{X}_2, ..., \mathbf{X}_i$  are all measurable.

The proof goes by induction. Let  $\bar{\sigma}_1$  be the smallest  $\sigma$ -algebra for which the random variable  $\mathfrak{Reg}_1(\mathbf{X}_1)$  is measurable. Then  $\bar{\sigma}_1 \subset \sigma_1$ . Define  $\mathbf{\bar{X}}_1 = E[\mathbf{X}_1|\bar{\sigma}_1]$ . Then  $\{\mathbf{\bar{X}}_1, \mathbf{X}_1; \bar{\sigma}_1, \sigma_1\}$  is a martingale.  $\mathbf{\bar{X}}_1$  has an atomic distribution with at most one atom  $\mathbf{Z}_{1;r} = E[\mathbf{X}_1 | \mathfrak{Reg}_1(\mathbf{X}_1) = r]$  per each region r of  $G_1^R$  and takes the same values on functionals  $f_{1i}^h, j = 1, 2, \ldots, U(1, h), h \in \{\tau, \tau_I\}$ , as  $\mathbf{X}_1$  by Lemma 3.

Now,  $\mathbf{\bar{X}}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}$  is also a martingale. Let  $\bar{\sigma}_2$  be the smallest  $\sigma$ -algebra which includes  $\bar{\sigma}_1$  and for which  $\mathfrak{Reg}_2(\mathbf{X}_2)$  is measurable. Define  $\mathbf{\bar{X}}_2 = E[\mathbf{X}_2|\bar{\sigma}_2]$ . Then  $\{\mathbf{\bar{X}}_1, \mathbf{\bar{X}}_2, \mathbf{X}_2; \bar{\sigma}_1, \bar{\sigma}_2, \sigma_2\}$  is a martingale. Indeed,  $E[\mathbf{\bar{X}}_2|\bar{\sigma}_1] = E[E[\mathbf{X}_2|\bar{\sigma}_2]|\bar{\sigma}_1] =$  $E[\mathbf{X}_2|\bar{\sigma}_1] = \mathbf{\bar{X}}_1$ , since  $\bar{\sigma}_1 \subset \bar{\sigma}_2$ , and  $E[\mathbf{X}_2|\bar{\sigma}_2] = \mathbf{\bar{X}}_2$  by definition.  $\mathcal{L}(\mathbf{\bar{X}}_2 |\mathfrak{Reg}_1(\mathbf{X}_1) = r)$ has an atomic measure with at most one atom  $\mathbf{Z}_{2;r,\tilde{r}} = E[\mathbf{X}_2|\mathfrak{Reg}_1(\mathbf{X}_1) = r, \mathfrak{Reg}_2(\mathbf{X}_2) = \tilde{r}]$  per each region  $\tilde{r}$  of  $G_2^R$ . Also by Lemma 3  $\bar{\mathbf{X}}_2$  takes the same values on the functionals  $f_{2j}^h, j = 1, 2, \ldots, U(1, h), h \in \{I, II\}$ , as  $\mathbf{X}_2$ .

Continuing in the same way, we can form a Markov Martingale  $\bar{\mathbf{X}}_1, \ldots, \bar{\mathbf{X}}_{t^*-1}, \bar{\mathbf{X}}_{t^*}$ such that  $\mathcal{L}(\bar{\mathbf{X}}_t | [\mathfrak{Reg}_1(\mathbf{X}_1), \mathfrak{Reg}_2(\mathbf{X}_2), \ldots, \mathfrak{Reg}_{t-1}(\mathbf{X}_{t-1})] = [r_1, r_2, \ldots, r_{t-1}])$  has an atomic measure with at most one atom per region of  $\mathcal{R}_t$  for all  $t = 1, 2, \ldots, t^*$ .

Now, let  $\mathbf{B}_{t-t^*}$  be a two-dimensional Brownian motion starting at 0. We define for  $t \geq t^* \, \hat{\mathbf{X}}_t = \bar{\mathbf{X}}_{t^*} + \mathbf{B}_{t-t^*}$  and let  $\tau$  be the first hitting time of  $\hat{\mathbf{X}}_t$  on  $G_{t^*}^E$ . Then  $\bar{\mathbf{X}}_{t^*}, \hat{\mathbf{X}}_{t^* \wedge \tau}, \hat{\mathbf{X}}_{(t^*+1)\wedge \tau}, \dots$ , is a martingale by the optional stopping theorem. This martingale is bounded, thus there exists  $\hat{\mathbf{X}}_{\infty}$ , such that  $\hat{\mathbf{X}}_{(t^*+1)\wedge \tau}, \hat{\mathbf{X}}_{(t^*+2)\wedge \tau}, \dots, \hat{\mathbf{X}}_{\infty}$  is a martingale and  $\hat{\mathbf{X}}_{(t^*+1)\wedge \tau}, \hat{\mathbf{X}}_{(t^*+2)\wedge \tau}, \dots$ , converges to  $\hat{\mathbf{X}}_{\infty}$  a.s. by Doob's theorem. Let  $\dot{\mathbf{X}}_{t^*} = \hat{\mathbf{X}}_{\infty}$ .

By Lemma 3,  $P\left(\dot{\mathbf{X}}_{t^*} \in G_{t^*}^E\right) = 1$  and  $\dot{\mathbf{X}}_{t^*}$  takes the same values on functionals  $E[f_{t^*j}^h(.)]$ ,  $E[f_{fut,j}^h(.)]$  and E[g(.)] as  $\bar{\mathbf{X}}_{t^*}$ . In the same way, but by adding an one dimensional Brownian motion to  $\dot{\mathbf{X}}_{t^*}$  over the edge to which  $\dot{\mathbf{X}}_{t^*}$  belongs, we can reduce the support of the martingale at time  $t^*$  to  $G_{t^*}^V$ .

Thus, to solve optimization problem (c), it is enough to consider only Markov martingales  $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}$  satisfying the property  $P_G$ . Consequently, it is a finite dimensional optimization problem (distributions with the  $P_G$  property can be described with finitely many parameters.) General measures belong to an infinite-dimensional space which is even uncountable. So the above result is a significant simplification of the problem. However we will see that the number of parameters grows exponentially with  $t^*$ . So solution methods described below will work only if  $t^*$  is not very large.

#### 4.3 Algorithm for the two dimensional case

By Theorem 7 in order to solve Problem (c) it is enough to consider martingales with property  $P_G$ . That means that the distribution of the initial state (t = 1) is atomic with at most one atom per each region of  $G_1^R$ , the distribution of the final state  $(t = t^*)$ is atomic with atoms at the vertices of  $G_{t^*}$ , and the conditional distribution of each intermediate state  $(1 < t < t^*)$  is atomic with at most one atom per each region of  $G_t^R$ , conditioned that the sequence of all prior visited regions is known.

The distribution of the described process is fully determined by the probabilities

$$P_{t^*;r_1,\dots,r_{t^*-1},r_{t^*}} = P\left(\left[\mathfrak{Reg}_1(\mathbf{X}_1),\dots,\mathfrak{Reg}_{t^*-1}(\mathbf{X}_{t^*-1}),\mathcal{V}er(\mathbf{X}_{t^*})\right] = [r_1,\dots,r_{t^*-1},v]\right) (19)$$

of visiting all possible sequences of regions (vertices at the final state), and by the unique places of visits of each particular region, conditioned on the prior sequence of regions. Let  $\mathbf{Z}_{t;r_1,r_2,...,r_t} \in \mathbb{R}^2_+$  denote the place of the unique atom of the conditional distribution

$$\mathcal{L}\left(\mathbf{X}_{t} \mid \left[\mathfrak{Reg}_{1}(\mathbf{X}_{1}), \ldots, \mathfrak{Reg}_{t}(\mathbf{X}_{t})\right] = \left[r_{1}, r_{2}, \ldots, r_{t}\right]\right),$$

where  $r_1 \in \mathcal{R}_1, r_2 \in \mathcal{R}_2, \ldots, r_t \in \mathcal{R}_t, t < t^*$ . Since all regions of  $\mathcal{R}_t$  for  $t = 1, 2, \ldots, t^* - 1$ are rectangular,  $\mathbf{Z}_{t;r_1,r_2,\ldots,r_t}$  can be represented as

$$\mathbf{Z}_{t;r_{1},r_{2},...,r_{t}} = \mathbf{u}_{r_{t}} + \mathbf{w}_{r_{t}} z_{t;r_{1},r_{2},...,r_{t}}^{I} + \mathbf{v}_{r_{t}} z_{t;r_{1},r_{2},...,r_{t}}^{II},$$
(20)

where  $\mathbf{u}_{r_t}, \mathbf{w}_{r_t}, \mathbf{v}_{r_t} \in \mathbb{R}^2_+$  are vectors uniquely defined by the rectangular  $r_t$  (see Figure 8) and  $z^h_{t;r_1,r_2,...,r_t} \in [0,1], h = r_{r_t}$ .



Figure 8: Parametrization of a rectangular.

Notice that  $f_{tj}^{I}$ ,  $f_{tj}^{II}$  are affine when restricted to one of the regions of  $G_{t}^{R}$ , and

$$f_{tj}^{h}\left(\mathbf{Z}_{t;r_{1},\ldots,r_{t-1},r}\right) = a_{tj}^{h}(r)z_{t;r_{1},r_{2},\ldots,r_{t-1},r}^{h} + b_{tj}^{h}(r), \quad r \in \mathcal{R}_{t}$$
(21)

for some coefficients  $a_{tj}^h(r)$ ,  $b_{tj}^h(r)$ , which are easy to determine.

For convenience we also introduce

$$P_{t;r_1,\ldots,r_t} = P\left([\mathfrak{Reg}_1(\mathbf{X}_1),\ldots,\mathfrak{Reg}_t(\mathbf{X}_t)] = [r_1,\ldots,r_t]\right)$$

for each  $t < t^*$ , which are probabilities of visiting a particular sequence of regions prior to time t.

In order to satisfy the conditions of a probability measure, in addition to nonnegativity, we must have

$$\sum_{r \in \mathcal{R}_1} P_{1;r} = 1, \tag{22}$$

$$P_{t-1;r_1,r_2,\ldots,r_{t-1}} = \sum_{r \in \mathcal{R}_t} P_{t;r_1,r_2,\ldots,r_{t-1},r}, \text{ for all } r_1 \in \mathcal{R}_1,\ldots,r_{t-1} \in \mathcal{R}_{t-1}, t = 2,\ldots,t^* - 1, (23)$$

$$P_{t^*-1;r_1,r_2,\dots,r_{t^*-1}} = \sum_{r \in \mathcal{R}_t} P_{t;r_1,r_2,\dots,r_{t^*-1},v}, \text{ for all } r_1 \in \mathcal{R}_1,\dots,r_{t^*-1} \in \mathcal{R}_{t^*-1}, v \in \mathcal{V}.$$
(24)

To satisfy the martingale condition we must have

$$\mathbf{Z}_{t-1;r_1r_2,\dots,r_{t-1}} P_{t-1;r_1,r_2,\dots,r_{t-1}} = \sum_{r \in \mathcal{R}} \mathbf{Z}_{t;r_1r_2,\dots,r_{t-1},r} P_{t;r_1,r_2,\dots,r_{t-1},r}$$
(25)  
for all  $r_1 \in \mathcal{R}_1,\dots,r_{t-1} \in \mathcal{R}_{t-1}, \quad t = 2,3,\dots,t^* - 1,$ 



Figure 9: Linear inequalities on  $\hat{z}_{t;r_1,r_2,\ldots,r_{t-1},r_t}$  and  $P_{t;r_1,r_2,\ldots,r_t}$ .

$$\mathbf{Z}_{t^*-1;r_1r_2,\dots,r_{t^*-1}}P_{t-1;r_1,r_2,\dots,r_{t^*-1}} = \sum_{v \in \mathcal{V}} \mathbf{Z}_{t^*;r_1r_2,\dots,r_{t^*-1},v}P_{t^*;r_1,r_2,\dots,r_{t-1},v} \quad (26)$$
  
for all  $r_1 \in \mathcal{R}_1,\dots,r_{t^*-1} \in \mathcal{R}_{t^*-1}, v \in \mathcal{V}.$ 

Notice, that  $\mathbf{Z}_{t;r_1r_2,...,r_{t-1},r}$  is defined in terms of  $z_{t;r_1r_2,...,r_{t-1},r}^I$  and  $z_{t;r_1r_2,...,r_{t-1},r}^{II}$  in (20).

Finally, the measure must satisfy the prices of options. We use the representation (21) of payoff functions to write down the constraints from options prices. For satisfying past constraints we must have

$$\sum_{r_1 \in \mathcal{R}_1, \dots, r_t \in \mathcal{R}_t} \left( a_{tj}^h(r_t) z_{t;r_1, r_2, \dots, r_{t-1}, r_t}^h + b_{tj}{}^h(r_t) \right) P_{t;r_1, r_2, \dots, r_t} = C_{tj}^h, \qquad (27)$$
  
for  $\{(h, t, j) \in \mathcal{T} | t < t^*.\}$ 

The present constraint are

$$\sum_{r_1 \in \mathcal{R}_1, \dots, r_{t^*-1} \in \mathcal{R}_{t^*-1}, v \in \mathcal{V}} f^h_{t^*j}(v) P_{t^*; r_1, \dots, r_{t^*-1}, v} = C^h_{t^*j}, \qquad (28)$$
  
for  $\{(h, t, j) \in \mathcal{T} | t = t^*.\}$ 

Finally, constraints from options of maturities larger than  $t^*$  are

$$\sum_{r_1 \in \mathcal{R}_1, \dots, r_{t^*-1} \in \mathcal{R}_{t^*-1}, v \in \mathcal{V}} f^h_{fut,j}(v) P_{t^*; r_1, r_2, \dots, r_{t^*-1}, v} \le C^h_{fut,j}$$
(29)  
for  $\{(h, t, j) \in \mathcal{T} | t > t^*.\}$ 

Thus problem (c) can be written as :

r

Find the collection of  $P_{t;r_1,r_2,...,r_t} \ge 0$  and  $z_{t;r_1,...,r_t}^h \in [0,1]$ ,  $r_1 \in \mathcal{R}_1, \ldots, r_t \in \mathcal{R}_t$ ,  $t = 1, 2, \ldots, t^* - 1$ , h = I, II, such that it satisfies conditions (22)-(29) and

$$\sum_{1 \in \mathcal{R}_1, \dots, r_{t^*-1} \in \mathcal{R}_{t^*-1}, v \in \mathcal{V}} g(v) P_{t^*; r_1, \dots, r_{t^*-1}, v}$$

is maximized (minimized.)

The above system is not linear. By making the change of variables

$$\hat{z}_{t;r_1,r_2,\dots,r_{t-1},r_t} = z_{t;r_1,r_2,\dots,r_{t-1},r_t} P_{t;r_1,r_2,\dots,r_t}$$
(30)

for  $t = 1, 2, ..., t^* - 1$ , we make it linear. The domain for  $(\hat{z}_{t;r_1,r_2,...,r_t}, P_{t;r_1,r_2,...,r_t})$  is represented on Figure 9. Thus all the constraints as well as the objective function are linear.

#### 4.4 Computational Example

The data on VerizonCM and Cisco call options (Table 1) is taken from The Wall Street Journal of October 22 2002 to derive best possible bounds on the price of an option with a payoff function  $(V + C - k)^+$ , where V is the price of the VerizonCM stock at the maturity, C is the price of Cisco stock and k is the strike price. The prices of VerizonCM and Cisco stocks on that day were \$37.75 and \$11.22 correspondingly.

Table 2 represents best possible bounds on the price of the option with maturity 52, if all given options are taken into account. Table 3 gives bounds on the price of the same option, if only options of maturity 52 are taken into account. As expected, these bounds are looser.

As can be noticed, bounds for strike prices smaller than 20 are extremely tight. That makes sense, since the probability that the sum of prices of the two given stocks will go below 20, is almost zero. Thus, the option will be exercised almost surely and buying the option is almost equivalent to buying the two stocks.

#### 4.5 Approximation approach

In this section, we develop an approximation approach which allows to overcome the exponential growth (in  $t^*$ ) of the number of variables.

Recall that we consider only martingales  $\mathbf{X}_t$  such that  $P(\mathbf{X}_t \leq L) = 1$  for some L > 0. For now we will allow  $g : \mathbb{R}^2_+ \to \mathbb{R}$ , to be any function continuous on  $[0, L] \times [0, L]$  and develop a general approximation algorithm. Then we will show how the efficiency of the algorithm can be improved, if g is a continuous piecewise linear function. Choose  $\epsilon > 0$ and designate by  $L_{\epsilon}$  the set of all vertices of the  $\epsilon$ -square lattice in  $[0, L] \times [0, L] \subset \mathbb{R}^2$ .

**Theorem 8.** Let  $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}$  be a two-dimensional Markov Martingale with values in  $[0, L] \times [0, L]$ . Then there exists a two-dimensional Markov Martingale  $\tilde{\mathbf{X}}_1, \tilde{\mathbf{X}}_2, \ldots, \tilde{\mathbf{X}}_{t^*}$ such that for all  $t = 1, \ldots, t^*, P\left(\tilde{\mathbf{X}}_t \in L_{\epsilon}^t\right) = 1$  and

$$P\left(\left|\tilde{X}_{t}^{h}-X_{t}^{h}\right|\leq t\epsilon\right)=1,\quad h=T,T$$

Here  $L_{\epsilon}^{t}$  is an  $\epsilon$ -square lattice in  $[0, L + (t-1)\epsilon] \times [0, L + (t-1)\epsilon]$ .

	VERIZON		CISCO				
maturity	strike price	price	maturity	strike price	price		
17	30.0	7.80	17	10.0	1.65		
17	35.0	3.70	17	12.5	0.35		
17	37.5	2.05	52	12.5	0.65		
17	40.0	0.95	80	7.50	3.75		
17	42.5	0.35	80	10.0	2.15		
52	40.0	1.80	80	12.5	0.80		
80	30.0	8.60	80	15.0	0.30		
80	35.0	5.10	171	15.0	0.65		
80	40.0	2.20					
171	42.5	2.00					

Table 1: Data.

strike price	0	10	20	30	40	45	50	55	60	70
lower bound	48.97	38.98	28.99	19.09	9.02	4.03	0.89	0.57	0.04	0.01
upper bound	48.97	39.02	29.31	19.47	10.56	7.17	3.91	2.08	1.93	1.62

Table 2: Best possible lower and upper bounds on the price of an option on the linear combination of two stocks. Maturity is 52. Options of all maturities are taken into account.

strike price	0	10	20	30	40	45	50	55	60	70
lower bound	48.97	38.98	28.97	18.97	8.97	3.97	0.25	0.00	0.00	0.00
upper bound	48.97	39.97	30.99	21.99	13.01	8.79	4.56	2.37	2.23	1.92

Table 3: Best possible lower and upper bounds on the price of an option on the linear combination of two stocks. Maturity is 52. Only options of this maturity are taken into account. *Proof.* Let  $\mathcal{B}_1 \subset \mathcal{B}_2 \subset \cdots \subset \mathcal{B}_{t^*}$  be the family of  $\sigma$ -algebras corresponding to the martingale  $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}$ . Let  $\mathbf{\Delta}_t := \mathbf{X}_t - \mathbf{X}_{t-1}$  for  $t = 2, 3, \ldots, t^*$ . Then

$$\mathbf{X}_t = \mathbf{X}_1 + \mathbf{\Delta}_2 + \mathbf{\Delta}_3 + \dots + \mathbf{\Delta}_t$$

and  $E\left[\mathbf{\Delta}_t | \mathcal{B}_{t-1}\right] = 0.$ 



Figure 10: Lattice  $\tilde{L}_{\epsilon}$ .

Let  $\tilde{L}_{\epsilon}$  designate the set of all vertices of the  $\epsilon$ -square lattice in  $[-L, L] \times [-L, L] \in \mathbb{R}^2$ (Figure 10.) And let  $S^j$ :  $[-L, L] \times [-L, L] \rightarrow \tilde{L}_{\epsilon}$ , j = 1, 2, 3, 4, be the family of functions, such that each function puts in correspondence to  $x \in [-L, L] \times [-L, L]$  one of the closest to it four vertices of  $\tilde{L}_{\epsilon}$ .

Notice that  $\Delta_t \in [-L, L] \times [-L, L]$  for all  $t = 2, \ldots, t^*$ . Let us define a random variable  $\tilde{\Delta}_t$  such that conditioned on  $\Delta_t$ , it can take only four values:  $\{S^j(\Delta_t) - \Delta_t, j = 1, 2, 3, 4\}$  and such that  $E\left[\tilde{\Delta}_t \middle| \mathcal{B}_t\right] = 0$ . Moreover, we add a requirement, that conditioned on  $\mathcal{B}_t$ ,  $\tilde{\Delta}_t$  is independent of any other random variables in consideration, including  $\tilde{\Delta}_1, \tilde{\Delta}_2, \ldots, \tilde{\Delta}_{t-1}$ . While  $\tilde{\Delta}_2, \tilde{\Delta}_3, \ldots, \tilde{\Delta}_{t^*}$  can be defined based on  $\Delta_2, \Delta_3, \ldots, \Delta_{t^*}$  as described above, define  $\tilde{\Delta}_1$  in a similar way, but take  $\Delta_1 = \mathbf{X}_1$ .

Then we have  $\mathbf{\Delta}_t + \tilde{\mathbf{\Delta}}_t \in \tilde{L}_{\epsilon}$  and  $P\left(\left|\tilde{\Delta}_t^h\right| \leq \epsilon\right) = 1, h = I, II, t = 1, 2..., t^*$ . Let  $\sigma\left(\tilde{\mathbf{\Delta}}_1, \tilde{\mathbf{\Delta}}_2, \ldots, \tilde{\mathbf{\Delta}}_t\right)$  denote the smallest  $\sigma$ -algebra for which  $\tilde{\mathbf{\Delta}}_1, \tilde{\mathbf{\Delta}}_2, \ldots, \tilde{\mathbf{\Delta}}_t$  are all measurable and

$$\begin{split} \tilde{\mathcal{B}}_t &:= \sigma \left( \mathcal{B}_t \cup \sigma \left( \tilde{\Delta}_1, \tilde{\Delta}_2, \dots, \tilde{\Delta}_t \right) \right), t = 1, 2 \dots, t^* \\ & \tilde{\mathbf{X}}_1 = \mathbf{X}_1 + \tilde{\Delta}_1, \\ & \tilde{\mathbf{X}}_2 = \mathbf{X}_2 + \tilde{\Delta}_1 + \tilde{\Delta}_2, \\ & \dots \\ & \tilde{\mathbf{X}}_{t^*} = \mathbf{X}_{t^*} + \tilde{\Delta}_1 + \tilde{\Delta}_2 + \dots + \tilde{\Delta}_{t^*}. \end{split}$$

 $\tilde{\mathbf{X}}_{\mathbf{1}} \in L_{\epsilon}$  is obvious. Now

$$\tilde{\mathbf{X}}_{\mathbf{2}} = \mathbf{X}_{2} + \tilde{\mathbf{\Delta}}_{\mathbf{1}} + \tilde{\mathbf{\Delta}}_{\mathbf{2}} = \mathbf{X}_{1} + (\mathbf{X}_{2} - \mathbf{X}_{1}) + \tilde{\mathbf{\Delta}}_{\mathbf{1}} + \tilde{\mathbf{\Delta}}_{\mathbf{2}} = \left(\mathbf{X}_{1} + \tilde{\mathbf{\Delta}}_{\mathbf{1}}\right) + \left(\mathbf{\Delta}_{2} + \tilde{\mathbf{\Delta}}_{\mathbf{2}}\right) \in \tilde{L}_{\epsilon}.$$

But since also  $P\left(\left|\tilde{X}_{2}^{h}-X_{2}^{h}\right|\leq 2\epsilon\right)=1, h=I,II$ , and  $\mathbf{X}_{2}\in L_{\epsilon}\subset \tilde{L}_{\epsilon}$ , it must be that  $\tilde{\mathbf{X}}_{2}\in L_{\epsilon}^{2}$ . Continuing recursively the theorem follows.

If  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n$  is a two-dimensional martingale, that satisfies  $E\left[f_{tj}^h(\mathbf{X}_t)\right] = C_{tj}^h$  for  $(t, j, h) \in \mathcal{T}$ , and  $\tilde{\mathbf{X}}_1, \tilde{\mathbf{X}}_2, \dots, \tilde{\mathbf{X}}_n$  is a two-dimensional martingale, such that  $P\left(\left|\tilde{X}_t^h - X_t^h\right| \le t\epsilon\right) = 1, h =_{I,II}, t = 1, \dots, n$ , then we must have

$$\left| E \left[ f_{tj}^{h} \left( \tilde{\mathbf{X}}_{\mathbf{t}} \right) \right] - C_{tj}^{h} \right| \le \epsilon t, \quad h = \iota, \iota \iota, \quad t = 1, \dots, n$$
(31)

and

$$\left| E\left[g\left(\tilde{\mathbf{X}}_{\mathbf{t}^*}\right)\right] - E\left[g\left(\mathbf{X}_{t^*}\right)\right] \right| \le M\sqrt{2}\epsilon t^*$$
(32)

where M is the smallest number, such that

$$|g(\mathbf{x}_2) - g(\mathbf{x}_1)| \le M \|\mathbf{x}_1 - \mathbf{x}_2\|_2$$

for all  $\mathbf{x}_1, \mathbf{x}_2 \ge 0$ , Notice that if  $g(x, y) = |\alpha x + \beta y - k|^+$ , then  $\sqrt{2}M = \alpha + \beta$ .

**Theorem 9.** The following problem has its optimal solution converging to the optimal solution of problem (18) as  $\epsilon \to 0$ :

Minimize (maximize)  $E\left[g\left(\tilde{\mathbf{X}}_{\mathbf{t}^*}\right)\right]$  over all Markov martingales  $\tilde{\mathbf{X}}_1, \tilde{\mathbf{X}}_2, \dots, \tilde{\mathbf{X}}_{\mathbf{t}^*}$ , such that  $P\left(\tilde{\mathbf{X}}_{\mathbf{t}} \in L_{\epsilon}^t\right) = 1$  and

$$\left| E\left[ f_{tj}^{h}\left( \tilde{\mathbf{X}}_{\mathbf{t}} \right) \right] - C_{tj}^{h} \right| \le \epsilon t, \quad \left\{ (h, t, j) \in \mathcal{T} \mid t \le t^{*} \right\},$$
(33)

$$E\left[f_{fut,j}^{h}\left(\tilde{\mathbf{X}}_{\mathbf{t}^{*}}\right)\right] - C_{fut,j}^{h} \leq \epsilon t^{*}, \quad \left\{\left(h,_{fut},j\right) \in \mathcal{T} \mid \left(k_{fut,j}^{h}, C_{fut,j}^{h}\right) \in \mathcal{F}^{h}\right\}.$$
(34)

*Proof.* Let us prove the theorem for the case when we want to maximize  $E[g(\mathbf{X}_{t^*})]$ . For the minimization problem the proof is similar.

Let  $g^*$  be the optimal solution to the original problem (c) and  $g^s$  the supremum as  $\epsilon \to 0$  of optimal solutions to problems with martingales' state spaces restricted to  $L^t_{\epsilon}$  and subject to the constraints (33) and (34). From Theorem 8 and (32) it immediately follows that  $g^s \ge g^*$ . Suppose that there exists  $\delta > 0$ , such that  $g^s \ge g^* + \delta$ . Then there exists a sequence of  $\epsilon$ 's,  $\{\{\epsilon_n\} \mid \epsilon_1 > \epsilon_2 > \cdots < \epsilon_n > \cdots; \lim_{n \to \infty} \epsilon_n \to 0\}$ , such that the limit of optimal solutions to problems corresponding to  $\epsilon_1, \epsilon_2, \ldots, \epsilon_n, \ldots$ , is equal to  $g^s$ .

Let  $\{\mathbf{X}_t^{\epsilon_n}\}_{n\geq 1}$  be the sequence of martingales corresponding to the sequence  $\{\epsilon_n\}$ , such that  $\{\mathbf{X}_t^{\epsilon_n}\}_{n\geq 1}$  defines an optimal solution to the  $\epsilon_n$ -problem.

By definition of  $\{\mathbf{X}_t^{\epsilon_n}\}$ , it takes values in  $L_{\epsilon_n}^{t^*}$  and

$$|E[f_{tj}(\mathbf{X}_t^{\epsilon_n})] - C_{tj}| \le \epsilon_n t, \quad t = 1, 2, \dots, t^*.$$

Let  $\mu^{\epsilon_n}$  designate the law of  $\{\mathbf{X}_t^{\epsilon_n}\}_{t=1,\dots,t^*}$ . Then  $\mu_{\epsilon_1}, \mu_{\epsilon_2}, \dots, \mu_{\epsilon_n}, \dots$  have support in  $[0, L + t^* \epsilon_1]^2$ , which is a compact. It follows that the sequence of laws  $\{\mu_{\epsilon_n}\}_{n\geq 1}$  is uniformly tight. Consequently, there exists a subsequence  $\mu_{\epsilon_n(\epsilon)} \to \mu$  for some law  $\mu$  [9, p.230]. But then we have an admissible solution to our original problem with an optimal value greater than  $g^* + \delta$ , which is a contradiction.

#### 4.6 Algorithm for the approximation approach

Let us enumerate all the nodes of  $L_{\epsilon}^{t^*}$  and denote by  $\mathbb{N}$  the set of all the indexes. Designate by  $\mathbf{Z}_{\mathfrak{n}} \in L_{\epsilon}^{t^*}$ , the  $\mathfrak{n}$ -th node of  $L_{\epsilon}^{t^*}$ . Let

$$P_{t;\mathfrak{n}} = P\left(\tilde{\mathbf{X}}_{\mathbf{t}} = \mathbf{Z}_{\mathfrak{n}}\right) \text{ and}$$
$$P_{t;\mathfrak{n}_{1},\mathfrak{n}_{2}} = P\left(\tilde{\mathbf{X}}_{\mathbf{t}} = \mathbf{Z}_{\mathfrak{n}_{1}}; \ \tilde{\mathbf{X}}_{\mathbf{t}+1} = \mathbf{Z}_{\mathfrak{n}_{2}}\right).$$

Notice that the distribution of  $\tilde{X}_1, \tilde{X}_2, \ldots, \tilde{X}_{t^*}$  is uniquely determined by  $P_{t;\mathfrak{n}_1,\mathfrak{n}_2}, t \in \{1, 2, \ldots, t^*\}, \mathfrak{n}_1, \mathfrak{n}_2 \in \mathcal{N}.$ 

Following the same set of arguments as in the description of the algorithm for the exact solution to the one dimensional case, we can state that the problem consists in finding the collection of  $P_{t;n}$  and  $P_{t;n_1,n_2}$ ,  $t \in \{1, 2, ..., t^*\}$ ,  $\mathfrak{n}, \mathfrak{n}_1, \mathfrak{n}_2 \in \mathcal{N}$ , which solves the linear optimization problem:

$$\begin{aligned} \max \ (\min) \quad &\sum_{\mathbf{n}\in\mathcal{N}} P_{t^*;\mathbf{n}} \ g(\mathbf{Z}_{\mathbf{n}}) \\ \text{s.t.} \quad &\sum_{\mathbf{n}\in\mathcal{N}} P_{1;\mathbf{n}} = 1, \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n}_1,\mathbf{n}} = P_{t;\mathbf{n}_1} \quad \text{for all } t < t^*, \ \mathbf{n}_1 \in \mathcal{N}, \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n},\mathbf{n}_1} = P_{t+1;\mathbf{n}_1} \quad \text{for all } t < t^*, \ \mathbf{n}_1 \in \mathcal{N}, \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n}_1,\mathbf{n}} \mathbf{Z}_{\mathbf{n}} = P_{t;\mathbf{n}_1} \mathbf{Z}_{\mathbf{n}_1} \quad \text{for all } t < t^*, \ \mathbf{n}_1 \in \mathcal{N}, \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n}} f_{tj}^h(\mathbf{Z}_{\mathbf{n}}) - C_{tj}^h \leq t\epsilon, \quad &\sum_{\mathbf{n}\in\mathcal{N}} P_{t;\mathbf{n}} f_{tj}^h(\mathbf{Z}_{\mathbf{n}}) - C_{tj}^h \geq -t\epsilon \\ &\text{for } \{(h,t,j)\in\mathcal{T} \mid t \leq t^*\}, \\ &\sum_{\mathbf{n}\in\mathcal{N}} P_{t^*;\mathbf{n}} f_{fut,j}^h(\mathbf{Z}_{\mathbf{n}}) - C_{fut,j}^h \leq t^*\epsilon \\ &\text{for } \{(h,t,j)\in\mathcal{T} \mid t > t^*\}, \\ &P_{t;\mathbf{n}_1,\mathbf{n}_2} \geq 0 \quad \text{for all } t \leq t^*, \ \mathbf{n}_1, \mathbf{n}_2 \in \mathcal{N}. \end{aligned}$$

As before this is a linear optimization problem.

## 4.7 Approximation approach for the special case of a payoff function

In this section, we show how the efficiency of the approximation algorithm can be improved, if g is a continuous piecewise linear function.

Let  $G^E = G_1^E \cup G_2^E \cup \cdots \cup G_{t^*}^E$  and let graph G be the graph with points of edges represented by  $G^E$ . Then, if we restrict the state space of martingales to  $G^E$ , the solution to problem (c) will not change. More precisely:

**Proposition 6.** Let  $\mathbf{X}_1, \mathbf{X}_2, \ldots, \mathbf{X}_{t^*}$  be a nonnegative two dimensional Markov Martingale bounded by L. Then there exists a nonnegative Markov Martingale  $\{\tilde{\mathbf{X}}_1, \tilde{\mathbf{X}}_2, \ldots, \tilde{\mathbf{X}}_{t^*}\}$ , with the state space in  $G^E$ , such that it satisfies

$$E\left[f_{tj}^{h}(\mathbf{X}_{t})\right] = E\left[f_{tj}^{h}\left(\tilde{\mathbf{X}}_{\mathbf{t}}\right)\right], \ t \leq t^{*}, \ (h, t, j) \in \mathcal{T},$$
$$E\left[f_{fut,j}^{h}(\mathbf{X}_{t})\right] = E\left[f_{fut,j}^{h}\left(\tilde{\mathbf{X}}_{\mathbf{t}}\right)\right], \ j = 1, 2, \dots, U(f_{ut}, t^{*}),$$
$$E\left[g\left(\mathbf{X}_{t^{*}}\right)\right] = E\left[g\left(\tilde{\mathbf{X}}_{t^{*}}\right)\right].$$

*Proof.* The proof is similar to the proof of Theorem 5 except that we deal with two dimensional random variables and, thus, a two dimensional Brownian motion should be introduced instead of one dimensional.  $\Box$ 

Let

$$\Lambda_{\epsilon}^{t} := \left\{ (x, y) \in L_{\epsilon}^{t} \mid \exists (x_{0}, y_{0}) \in G_{E} : |x - x_{0}| \le t\epsilon, |y - y_{0}| \le t\epsilon \right\}.$$

Then Theorem 9 will hold if we take  $\Lambda_{\epsilon}^{t}$  instead of  $L_{\epsilon}^{t}$ . The number of variables in this case grows only linear versus quadratically in the case of a general function g.

#### 4.8 Generalization to a multi-dimensional case

The exact solution, the approximation approach for an arbitrary payoff function g and the approximation approach for a continuous piecewise linear payoff function g, can all be extended from a two dimensional case to a multiple dimensional case. All theorems and definitions of this chapter can be reformulated for an n-dimensional case, and proofs will be identical up to a dimensionality. However the number of variables in the linear optimization problem will grow exponentially in the dimension.

## 4.9 Generalization to the case of options with continuous piecewise linear payoff functions

Across the paper we considered the case when prices of simple call options are given and bounds on prices of options with continuous piecewise linear functions must be found. From financial point of view this formulation of the problem can be motivated by the fact that simple call and put options are more liquid and prices on them are readily available in the market, compared to more complex exotic options. Notice, that conditions on European put options can be easily expressed as conditions on call options.

However, the problem could be extended to a more general case when options with continuous piecewise linear functions are given and bounds on similar type of options are to be determined. If there were no options of maturities larger than  $t^*$ , then all that has to be changed in the solution is definitions of the set  $\mathcal{K}$  in the one dimensional case and graphs  $G_t$  in the two dimensional case. In particular, in the one dimensional case, in section 3.2,  $\mathcal{K}$  must include all points of  $\mathbb{R}^+$  where payoff functions of given and target options change their derivatives. Similarly, for the two dimensional case, in section 4.2, lines defined in (a) and (b) of Definition 5 by strike prices of call options would have to be defined by points of  $\mathbb{R}^+$  where payoff functions of set  $\mathcal{K}$  and graphs  $G_t$  would remain valid.

However, the simplification that was made for treatment of future conditions (sections 3.1 and 4.1) is not possible if payoff functions of options of maturities larger than  $t^*$ are piecewise linear, since these functions are not necessarily convex. As a result, the martingale has to be constructed not only up to time  $t^*$ , but up to the time of the maximum options maturity, and future and present constraints have to be handled in the same way as past constraints. In the one dimensional case, since the state space of the martingale at each time is the same, there is no principal difference. All that one has to do in addition to changes in the definition of set  $\mathcal{K}$  is to extend the set of unknowns  $P_{t:n}$ and  $P_{t;n_1,n_2}$  up to the maximum time t and change inequalities to equalities in the future constraints. The problem remains a linear optimization problem. In the two dimensional case, however, there would not necessarily exist a martingale that takes values at the vertices of graph  $G_{t^*}$  at time  $t^*$  and satisfy all future constraints. As was shown in the proof of Theorem 7, there does exist a martingale that satisfies all given constraints and at time  $t^*$  has an atomic conditional distribution with at most one atom per each region of  $G_{t^*}^R$ , conditioned on the sequence of all prior visited regions. However, since regions of  $G_{t^*}^R$  are not necessarily rectangular, the simple form of parametrization (20) and the change of variables (30) can not be applied. Therefore, we don't think it is possible to reduce present constraints and the objective function to the linear form in this case.

It is easy to see that the approximation approach for the two dimensional case can be applied to the solution of the problem, when prices of options with continuous piecewise linear payoff functions are given. In this case, one only needs to extend the set of unknowns  $P_{t;n}$  and  $P_{t;n_1,n_2}$  from time  $t^*$  to the maximum maturity and use equalities instead of inequalities for future constraints.

### 5 Static arbitrage bounds on basket option prices

In this section, we address Problem (c) defined as follows:

#### Problem (d)

Given  $\mathbf{C} \in \mathbb{R}^m_+$ ,  $\mathbf{k} \in \mathbb{R}^m_+$ ,  $\omega_i \in \mathbb{R}^n$ , i = 1, ..., m and  $k_0 > 0$ ,  $w_0 \in \mathbb{R}^m_+$ , find the upper and lower bounds on

$$E\left(\omega_0^T \mathbf{X} - k_0\right)^+,\tag{35}$$

with respect to distributions of an n-dimensional random variable **X** with finite expectation and support in  $\mathbb{R}^n_+$  under the condition

$$E\left(\omega_i^T \mathbf{X} - k_i\right)^+ = C_i, \quad i = 1, \dots, m.$$
(36)

To solve the problem we make an additional assumption that the support of **X** is bounded, that is, there exists L > 0, such that  $P(X^h \le L) = 1$  for any h = 1, 2, ..., n.

Let  $f_i : \mathbb{R}^n_+ \to \mathbb{R}_+$  be the family of measurable functions defined as

$$f_i(\mathbf{X}) = \left(\mathbf{w}_i^T \mathbf{X} - k_i\right)^+, \quad i = 0, 1, \dots, m$$

Let  $\bigcup_j D_j = [0, L]^n$  be the partition of  $[0, L]^n$ , into subsets  $D_j$  of  $\mathbb{R}^n_+$  such that in the interior of each subset  $D_j$  all functions  $f_i$ ,  $i = 0, 1, \ldots, m$  are affine.

Let  $G \subset [0, L]^n$  be the graph in  $\mathbb{R}^n_+$  formed by intersecting hyperplanes

$$\omega_i^T \mathbf{X} = k_i, \quad i = 1, 2, ..., m,$$
  

$$X^h = 0, \quad h = 1, 2, ..., n,$$
  

$$X^h = L, \quad h = 1, 2, ..., n.$$
(37)

Let us enumerate these hyperplanes and call them  $h_1, h_2, \ldots, h_{2n+m}$  correspondingly. We call a vector  $\mathbf{r} \in [0, L]^n$  a vertex of G if there are n independent hyperplanes of (37) which intersect at  $\mathbf{r}$ . Let  $G_V$  designate the set of all vertices. Notice, that this set contains no more than  $C_{2n+m}^n = \frac{(2n+m)!}{(n+m)!n!}$  elements.

**Corollary 4.** For each random variable  $\mathbf{X}$  with support in  $[0, L]^n$  there exists a random variable  $\mathbf{\bar{X}}$  such that  $P(\mathbf{\bar{X}} \in G_V) = 1$  and  $E[f_i(\mathbf{X})] = E[f_i(\mathbf{\bar{X}})], i = 0, 1, ..., m$ .

Thus, to find optimal bounds we solve a linear optimization problem with unknown variables being weights of the atomic distribution. The atoms are located in the vertices of the defined graph.

## References

- A. d'Aspremont and L. El Ghaoui, Static arbitrage bounds on basket option prices, submitted to Operations Research, 2003.
- [2] D. Bertsimas and N. Bushueva, *Option Pricing without Price Dynamics: A geometric approach*, Operations Research Center, MIT, working paper, 2004.
- [3] D. Bertsimas and I. Popescu, On the relation between option and stock prices: a convex optimization approach, Operations Research 50 (2), p. 358-374, 2002.
- [4] F. Black and M.Scholes, *The pricing of options and corporate liabilities*, Journal of Political Economy, p. 637-54, 1973.
- [5] P. Boyle, X. Lin, Bounds on contingent claims based on several assets, Journal of Financial Economics, 46, p. 383-400, 1997.
- [6] J. Cox and C. Huang, Optimal consumption and portfolio policies when asset prices follow a diffusion process, Journal of Economic Theory 49, p. 33-83, 1989.
- J. Cox and S. Ross, The valuation of options for alternative stochastic processes, Journal of Financial Economics 3, p. 145-166, 1976.
- [8] J. Cox, S. Ross and M. Rubinstein, Option pricing: a simplified approach, Journal of Financial Economics, 7, No.3, p. 229-263, 1979.
- [9] R. Dudley, Real Analysis and Probability, New York: Chapman & Hall, 1989.
- [10] R. Dudley, Uniform Central Limit Theorems, Cambridge, MA: Cambridge University Press, 1999.
- B. Grundy, Option prices and the underlying asset's return distribution, Journal of Finance, 46(3), p. 1045-1070, 1991.
- [12] M. Harrison and D. Kreps, Martingales and arbitrage in multiperiod security markets, Journal of Economic Theory, 20, p. 381-408, 1979.
- [13] O. Kallenberg, Foundations of Modern Probability, Applied Probability Trust, 1997.
- [14] R. Kertz and U. Rosler, Stochastic and convex orders and lattices of probability measures, with a martingale interpretation, Israel Journal of Mathematics, 77, p. 129-164, 1992.
- [15] A. Lo, Semiparametric upper bounds for option prices and expected payoffs, Journal of Financial Economics, 19, p. 373-388, 1987.
- [16] F. Longstaff, Martingale restriction tests of option pricing models, version 1, Working Paper, University of California, Los Angeles, 1990.

- [17] S. Ross, Options and efficiency, Quarterly Journal of Economics, 90, p. 75-89, 1976.
- [18] M. Rubinstein, Implied binomial trees, Journal of Finance, 49(3): 771:819, 1994.
- [19] V. Strassen, The existence of probability measures with given marginals, Ann. Math. Statist., 36, p. 423-439, 1965.