

A SCHNEIDER TYPE THEOREM FOR HOPF ALGEBROIDS

A. ARDIZZONI, G. BÖHM, AND C. MENINI

ABSTRACT. Comodule algebras of a Hopf algebroid $\mathcal{H}$ with a bijective antipode, i.e. algebra extensions $B \subseteq A$ by $\mathcal{H}$, are studied. Assuming that a lifted canonical map is a split epimorphism of modules of the (non-commutative) base algebra of $\mathcal{H}$, *relative injectivity* of the $\mathcal{H}$ -comodule algebra A is related to the *Galois property* of the extension $B \subseteq A$ and also to the *equivalence* of the category of relative Hopf modules to the category of B -modules. This extends a classical theorem by H.-J. Schneider on Galois extensions by a Hopf algebra. Our main tool is an observation that relative injectivity of a comodule algebra is equivalent to *relative separability* of a forgetful functor, a notion introduced and analysed hereby.

CONTENTS

1. Introduction	1
2. Preliminaries. Coring extensions, entwining structures and Hopf algebroids	3
3. Relative separable functors	12
4. Application to entwining structures	18
5. Comodule algebras of Hopf algebroids	22
6. A Schneider type theorem	25
7. Equivariant injectivity and projectivity	29
Acknowledgements	33
References	33

1. INTRODUCTION

Galois extensions of non-commutative algebras by a Hopf algebra generalise Galois extensions of commutative rings by groups and are known as the algebraic (dual) versions of (non-commutative) principal bundles. By a Hopf Galois extension the following structure is meant. Comodules over a Hopf algebra H form a monoidal category $\mathfrak{M}^H$, whose monoids are called *comodule algebras*. This means an algebra and H -comodule A , such that the coaction $\rho^A : A \rightarrow A \otimes H$ is an algebra map (with respect to the tensor product algebra structure of the codomain). It can be looked at as a notion dual to the action of a group on a manifold. Dualising the notion of invariant points, *coinvariants* of A are defined as those elements on which coaction is trivial, i.e. the elements of the subalgebra

$$B := \{ b \in A \mid \rho^A(b) = b \otimes 1_H \}.$$

In this situation the algebra A is called an *extension of B by H* . The algebra extension $B \subseteq A$ is said to be *H -Galois* if in addition the so called *canonical map*

$$(1.1) \quad \text{can} : A \otimes_B A \rightarrow A \otimes H, \quad a \otimes_B a' \mapsto a\rho^A(a')$$

is bijective (hence an isomorphism of left A -modules and right H -comodules). This is a dual formulation of the condition that a group action on a manifold is free.

Date: February 26, 2019.

1991 *Mathematics Subject Classification.* Primary 16W30; Secondary 18A22.

Key words and phrases. Relative separable functors, relative injective comodule algebras, Hopf algebroids, Galois extensions.

(Right-right) *relative Hopf modules* are (right) modules for an H -comodule algebra A and (right) comodules for the Hopf algebra H , satisfying a compatibility condition with the H -coaction in A . In the case of an H -Galois extension $B \subseteq A$, relative Hopf modules are canonically identified with descent data for the extension $B \subseteq A$. Hence if A is faithfully flat as a left B -module, it follows by the Faithfully Flat Descent Theorem that the category $\mathfrak{M}_A^H$ of right-right relative Hopf modules is equivalent to the category $\mathfrak{M}_B$ of right B -modules.

In the study of Hopf Galois extensions, important tools are provided by theorems, stating that in appropriate situations surjectivity of the canonical map (1.1) implies its bijectivity. One group of such results (e.g. [KT, Theorem 1.7], [Scha1, Corollary 2.4.8 1] [SS, Theorem 3.1], [Bö2, Theorem 4.2 and Corollary 4.3]) can be called ‘*Kreimer-Takeuchi type*’ theorems (as their first representative was proven in [KT, Theorem 1.7]). In this group of theorems projectivity of the regular comodule of the coacting Hopf algebra is assumed. The other group involves ‘*Schneider type*’ theorems (after [Schn, Theorem I], see e.g. [SS, Theorem 4.9], [Brz2, Theorem 4.6], [MM1, Theorem 3.15], [MM2, Theorem 3.9]). Here relative injectivity of the Hopf comodule algebra in question is assumed.

The starting point of our work is an observation that the proofs of all above theorems share a common philosophy. Related to a comodule algebra A of a Hopf algebra H over a commutative ring k , there are forgetful functors

$$(1.2) \quad \mathfrak{M}_A^H \xrightarrow{\mathbb{R}} \mathfrak{M}^H \xrightarrow{\mathbb{U}} \mathfrak{M}_k.$$

If H is a projective k -module then the codomain of the canonical map (1.1) is a projective A -module. Then it follows from the surjectivity of the canonical map that its lifted version

$$(1.3) \quad \widetilde{\text{can}} : A \otimes_k A \rightarrow A \otimes_k H, \quad a \otimes_k a' \mapsto a\rho^A(a')$$

has a k -linear right inverse, i.e. it is a retraction of k -modules. The various Schneider type theorems give sufficient conditions for the forgetful functor $\mathbb{U}$ to reflect (certain) retractions. Then bijectivity of the canonical map (1.1) follows by a result of Schauenburg [Scha1, Corollary 2.4.8] stating that – under the additional assumption that all H -coinvariants of the obvious right H -comodule $A \otimes_k A$ are elements of $A \otimes_k B$ – the canonical map is bijective, provided that its lifted version (1.3) is a retraction of H -comodules.

In the present paper we introduce the notion of *separability* of a functor $\mathbb{U} : \mathfrak{A} \rightarrow \mathfrak{B}$, *relative* to a functor $\mathbb{R} : \mathfrak{A} \rightarrow \mathfrak{A}$ (not to be mixed with separability of the second kind in [CM]). An $\mathbb{R}$ -relative separable functor $\mathbb{U}$ reflects retractions in the sense that, for a morphism f in $\mathfrak{A}$ such that $\mathbb{U}\mathbb{R}(f)$ is a retraction, $\mathbb{R}(f)$ is a retraction. As it turns out, the conditions of all known Schneider type theorems imply the separability of the forgetful functor $\mathbb{U}$ in (1.2), relative to $\mathbb{R}$.

Our strategy, of tracing back Schneider type theorems to properties of a forgetful functor, can be compared to that of Caenepeel, Ion, Militaru and Zhu, when in [CIMZ] they explained all known Maschke type theorems by the separability of a forgetful functor.

The motivation of our work comes from a wish to prove a Schneider type theorem for more general algebra extensions by a Hopf algebroid, replacing the Hopf algebra H above. A Kreimer-Takeuchi type theorem was proven in [Bö2]. In that paper similar methods have been used as in [SS]: the entwining structure (over a non-commutative base), determined by a comodule algebra of a Hopf algebroid, has been studied. It turns out that this framework is not sufficient to obtain a Schneider type theorem for extensions by Hopf algebroids. Recall that a Hopf algebroid $\mathcal{H}$ consists of two related coring (and bialgebroid) structures, over two different base algebras L and R . A right $\mathcal{H}$ -comodule algebra A is in particular an R -ring. Together with the R -coring underlying $\mathcal{H}$ they form an entwining structure over R . Hence, by general results, $A \otimes_R \mathcal{H}$ possesses an A -coring structure. Its comodules are actually the relative Hopf modules. What is more, the L -coring underlying $\mathcal{H}$ is a right extension of the A -coring $A \otimes_R \mathcal{H}$, in the sense of [Brz3, Definition 2.1]. The proper approach to a Schneider type theorem for Hopf algebroids turns out to be a study of the forgetful functors

$$\mathfrak{M}_A^{\mathcal{H}} \xrightarrow{\mathbb{R}} \mathfrak{M}^{\mathcal{H}} \xrightarrow{\mathbb{U}} \mathfrak{M}_L,$$

related to this coring extension, incorporating both coring structures present in a Hopf algebroid.

The paper is organised as follows. Section 2 collects preliminary results about entwining structures (over arbitrary non-commutative algebras), coring extensions (in the sense of [Brz3]) and Hopf algebroids. In Section 3 the notion of a *separable* functor $\mathbb{U}$, *relative* to functors $\mathbb{L} : \mathfrak{L} \rightarrow \mathfrak{A}$ and $\mathbb{R} : \mathfrak{R} \rightarrow \mathfrak{A}$, is introduced and investigated. Section 4 concerns relative separability of a forgetful functor $\mathfrak{M}^{\mathcal{D}} \rightarrow \mathfrak{M}_L$, associated to an entwining structure $(A, \mathcal{D}, \psi)$ over an algebra L . If $\mathcal{D}$ possesses a grouplike element, relative separability of the forgetful functor is shown to imply relative injectivity of A as a $\mathcal{D}$ -comodule and, in the case when in addition the entwining map is bijective, also relative injectivity of A as an entwined module (see Theorem 4.3 and Proposition 4.4). A comodule algebra A for a Hopf algebroid $\mathcal{H}$ (over base algebras L and R) determines a coring extension which does not correspond to any entwining structure. In Section 5 separability of the forgetful functor $\mathfrak{M}^{\mathcal{H}} \rightarrow \mathfrak{M}_L$, relative to the forgetful functor from the category of relative Hopf modules to the category of $\mathcal{H}$ -comodules, is studied. In the case when the antipode of $\mathcal{H}$ is bijective, it is shown to be *equivalent* to relative injectivity of the $\mathcal{H}$ -comodule A (see Theorem 5.1). This result enables us to answer a question posed in [Bö2]. That is, in Proposition 5.2 we prove that, in a Galois extension $B \subseteq A$ by a finitely generated and projective Hopf algebroid $\mathcal{H}$ with a bijective antipode, A is faithfully flat as a left B -module if and only if it is faithfully flat as a right B -module. The main result is a Schneider type theorem in Section 6. Recall that Schneider's classical Theorem I in [Schn] deals with an algebra extension $B \subseteq A$ by a k -Hopf algebra H with a bijective antipode. It is assumed that H is a projective k -module and the canonical map (1.1) is surjective. Clearly, in this case the lifted canonical map (1.3) is a split epimorphism of k -modules. As a proper generalisation to an algebra extension $B \subseteq A$ by a Hopf algebroid $\mathcal{H}$, in Theorem 6.7 we assume that some lifted canonical map is a split epimorphism of modules for the (non-commutative) base algebra L of $\mathcal{H}$. This assumption is related to surjectivity of the canonical map and some projectivity conditions in Remark 6.4. Under the assumption that, for an algebra extension $B \subseteq A$ by a Hopf algebroid $\mathcal{H}$ with a bijective antipode, the lifted canonical map is a split epimorphism of L -modules, the Galois property of the extension is related to relative injectivity of the $\mathcal{H}$ -comodule A and to the equivalence of the category $\mathfrak{M}_A^{\mathcal{H}}$ of relative Hopf modules to the category of B -modules. Section 7 is devoted to a study of (relative) equivariant injectivity and projectivity properties.

2. PRELIMINARIES. CORING EXTENSIONS, ENTWINING STRUCTURES AND HOPF ALGEBROIDS

2.1. Throughout this paper the term **algebra** is used for an associative and unital but not necessarily commutative algebra over a fixed commutative ring k . Multiplication is denoted by juxtaposition and the unit element is denoted by 1.

The category of right (respectively, left) modules for an algebra A is denoted by $\mathfrak{M}_A$ (respectively, ${}_A\mathfrak{M}$). The set of morphisms between two A -modules M and M' is denoted by $\text{Hom}_A(M, M')$ (respectively, ${}_A\text{Hom}(M, M')$). The category of A - A bimodules is denoted by ${}_A\mathfrak{M}_A$ and its Hom sets by ${}_A\text{Hom}_A(M, M')$.

For an algebra A , the opposite algebra A^{op} means the same k -module A with opposite multiplication $a \otimes a' \mapsto a'a$. The tensor product $A \otimes_k A^{op}$ is an algebra by factorwise multiplication.

2.2. For a k -algebra A , an **A -ring** T means an algebra (or monoid) in the monoidal category of A - A bimodules. More explicitly, it consists of an A - A bimodule T , equipped with a bilinear associative product $\mu : T \otimes_A T \rightarrow T$ and a bilinear unit map $\eta : A \rightarrow T$. An A -ring T is equivalent to a k -algebra T and a k -algebra map $\eta : A \rightarrow T$.

For an A -ring (T, μ, η) , the opposite means the A^{op} -ring T^{op} , with A^{op} - A^{op} bimodule structure

$$A^{op} \otimes T \otimes A^{op} \rightarrow T, \quad a \otimes t \otimes a' \mapsto a'ta,$$

product $t \otimes_{A^{op}} t' \mapsto t't$ and unit $\eta : A^{op} \rightarrow T^{op}$.

An A -ring T determines a monad $\bullet \otimes_A T$ on $\mathfrak{M}_A$. By right T -modules we mean algebras for this monad. This notion coincides with the one of right modules for the k -algebra T . Left modules are defined symmetrically.

2.3. A **coring** over an algebra A means a coalgebra (or comonoid) in the monoidal category of A - A bimodules. More explicitly, it consists of an A - A bimodule $\mathcal{C}$, equipped with a bilinear coassociative coproduct $\Delta : \mathcal{C} \rightarrow \mathcal{C} \otimes_A \mathcal{C}$ and a bilinear counit map $\epsilon : \mathcal{C} \rightarrow A$. For the coproduct we use a Sweedler type index notation $\Delta(c) = c^{(1)} \otimes_A c^{(2)}$, for $c \in \mathcal{C}$, where implicit summation is understood. This extends the notion of a coalgebra.

For an A -coring $(\mathcal{C}, \Delta, \epsilon)$, the co-opposite means the A^{op} -coring $\mathcal{C}_{cop}$, with A^{op} - A^{op} bimodule $\mathcal{C}$,

$$A^{op} \otimes \mathcal{C} \otimes A^{op} \rightarrow \mathcal{C}, \quad a \otimes c \otimes a' \mapsto a'ca,$$

coproduct $\Delta^{op} : c \mapsto c^{(2)} \otimes_{A^{op}} c^{(1)}$ and counit ϵ .

An A -coring $\mathcal{C}$ determines a comonad $\bullet \otimes_A \mathcal{C}$ on $\mathfrak{M}_A$. By right $\mathcal{C}$ -comodules we mean coalgebras for this comonad. That is, a right $\mathcal{C}$ -comodule is a right A -module M , equipped with a right A -linear coassociative and counital coaction ϱ^M . For a right $\mathcal{C}$ -coaction we use an index notation of the form $\varrho^M(m) = m^{[0]} \otimes_A m^{[1]}$, for $m \in M$, implicit summation understood. Right $\mathcal{C}$ -comodule maps are right A -linear maps which are compatible with the coactions. The category of right $\mathcal{C}$ -comodules is denoted by $\mathfrak{M}^{\mathcal{C}}$ and its Hom sets by $\text{Hom}^{\mathcal{C}}(M, M')$. Left $\mathcal{C}$ -comodules are defined symmetrically. The coaction is denoted by ${}^M\varrho(m) = m^{[-1]} \otimes_A m^{[0]}$, for a left $\mathcal{C}$ -comodule M and $m \in M$, implicit summation understood. The category of left $\mathcal{C}$ -comodules is denoted by ${}^{\mathcal{C}}\mathfrak{M}$ and its Hom sets by ${}^{\mathcal{C}}\text{Hom}(M, M')$.

The forgetful functor $\mathfrak{M}^{\mathcal{C}} \rightarrow \mathfrak{M}_A$ possesses a right adjoint, the functor $\bullet \otimes_A \mathcal{C} : \mathfrak{M}_A \rightarrow \mathfrak{M}^{\mathcal{C}}$. The unit of the adjunction is given by the right coaction $\varrho^M : M \rightarrow M \otimes_A \mathcal{C}$, for $M \in \mathfrak{M}^{\mathcal{C}}$, and the counit of the adjunction is given in terms of the counit ϵ of $\mathcal{C}$ as $N \otimes_A \epsilon : N \otimes_A \mathcal{C} \rightarrow N$, for $N \in \mathfrak{M}_A$, cf. [BW, 18.13 (2)].

Recall that a morphism $f : C_1 \rightarrow C_2$ in a category $\mathfrak{C}$ is called a *split monomorphism* or *section* if there exists a morphism $\tilde{f} : C_2 \rightarrow C_1$ in $\mathfrak{C}$ such that $\tilde{f} \circ f = C_1$. A right comodule M for an A -coring $\mathcal{C}$ is called **relative injective** if any $\mathcal{C}$ -comodule map of domain M , which is a section of A -modules, is a section of $\mathcal{C}$ -comodules, too. By [BW, 18.18], M is a relative injective $\mathcal{C}$ -comodule if and only if the coaction ϱ^M is a section of $\mathcal{C}$ -comodules.

By [Brz1, Lemma 5.1], the right regular A -module extends to a comodule for an A -coring $\mathcal{C}$ if and only if there exists a **grouplike element** g in $\mathcal{C}$ (meaning that $\Delta(g) = g \otimes_A g$ and $\epsilon(g) = 1_A$). A bijective correspondence between grouplike elements g in $\mathcal{C}$ and right $\mathcal{C}$ -coactions ϱ^A in A is given by $g \mapsto (\varrho^A : a \mapsto ga)$. A similar equivalence holds between grouplike elements and left $\mathcal{C}$ -comodule structures in A .

For an A -coring $\mathcal{C}$ with a grouplike element g , the **coinvariants** with respect to g of a right $\mathcal{C}$ -comodule M are defined as the elements of the set

$$M^{co\mathcal{C}} = \{ m \in M \mid \varrho^M(m) = m \otimes_A g \}.$$

Coinvariants of left $\mathcal{C}$ -comodules are defined symmetrically. In particular, the coinvariants of A , both as a right $\mathcal{C}$ -comodule and a left $\mathcal{C}$ -comodule, are the elements of the subalgebra

$$B = \{ b \in A \mid gb = bg \}.$$

A grouplike element g in $\mathcal{C}$ determines an adjunction $(\bullet \otimes_B A, (\bullet)^{co\mathcal{C}})$, between the categories $\mathfrak{M}_B$ and $\mathfrak{M}^{\mathcal{C}}$. The unit and counit are given by the maps

$$(2.1) \quad u_N : N \rightarrow (N \otimes_B A)^{co\mathcal{C}}, \quad n \mapsto n \otimes_B 1_A, \quad \text{and}$$

$$(2.2) \quad n_M : M^{co\mathcal{C}} \otimes_B A \rightarrow M, \quad m \otimes_B a \mapsto ma,$$

respectively, for any $N \in \mathfrak{M}_B$ and $M \in \mathfrak{M}^{\mathcal{C}}$, cf. [BW, 28.8]. There is a symmetrical adjunction between the categories ${}_B\mathfrak{M}$ and ${}^{\mathcal{C}}\mathfrak{M}$.

An A -coring $\mathcal{C}$ with a grouplike element g is called a **Galois coring** if the canonical map

$$(2.3) \quad \text{can} : A \otimes_B A \rightarrow \mathcal{C}, \quad a \otimes a' \mapsto aga'$$

is bijective. For more information about corings we refer to the monograph [BW].

2.4. Let $\mathcal{D}$ be a coring over a base k -algebra L and $\mathcal{C}$ a coring over a k -algebra A . Assume that $\mathcal{C}$ is a $\mathcal{C}$ - $\mathcal{D}$ bicomodule with the left regular $\mathcal{C}$ -coaction $\Delta_{\mathcal{C}}$ and some right $\mathcal{D}$ -coaction $\tau_{\mathcal{C}}$. By definition [BW, 22.1], this means that $\tau_{\mathcal{C}}$ is left A -linear (hence $\mathcal{C} \otimes_A \mathcal{C}$ is also a right $\mathcal{D}$ -comodule with coaction $\mathcal{C} \otimes_A \tau_{\mathcal{C}}$) and the coproduct $\Delta_{\mathcal{C}}$ is right $\mathcal{D}$ -colinear. Equivalently, the coproduct $\Delta_{\mathcal{C}}$ is right L -linear (hence $\mathcal{C} \otimes_L \mathcal{D}$ is a left $\mathcal{C}$ -comodule with coaction $\Delta_{\mathcal{C}} \otimes_L \mathcal{D}$) and the $\mathcal{D}$ -coaction $\tau_{\mathcal{C}}$ is left $\mathcal{C}$ -colinear. In this case, following [Brz3, Definition 2.1], we say that $\mathcal{D}$ is a **right extension** of $\mathcal{C}$, for the following reason. In [Brz3, Theorem 2.6] this definition has been shown to be equivalent to the existence of a k -linear functor $\mathbb{R} : \mathfrak{M}^{\mathcal{C}} \rightarrow \mathfrak{M}^{\mathcal{D}}$, making the following diagram, involving four forgetful functors, commutative.

$$(2.4) \quad \begin{array}{ccc} \mathfrak{M}^{\mathcal{C}} & \xrightarrow{\quad \mathbb{R} \quad} & \mathfrak{M}^{\mathcal{D}} \\ \searrow \mathbb{U}^{\mathcal{C}} & & \swarrow \mathbb{U}^{\mathcal{D}} \\ \mathfrak{M}_A & & \mathfrak{M}_L \\ \searrow \mathbb{F}^A & & \swarrow \mathbb{F}^L \\ \mathfrak{M}_k & & \end{array}$$

Indeed, using the right $\mathcal{D}$ -coaction $\tau_{\mathcal{C}} : c \mapsto c_{[0]} \otimes_L c_{[1]}$, for $c \in \mathcal{C}$ (note our convention to use character τ for $\mathcal{D}$ -coactions and lower indices of the Sweedler type to denote components of the coproduct and coactions of $\mathcal{D}$), any right $\mathcal{C}$ -comodule M can be equipped with a right $\mathcal{D}$ -comodule structure with right L -action

$$(2.5) \quad ml := m^{[0]} \epsilon_{\mathcal{C}}(m^{[1]}l), \quad \text{for } m \in M \text{ and } l \in L,$$

and $\mathcal{D}$ -coaction

$$(2.6) \quad \tau_M : M \rightarrow M \otimes_L \mathcal{D}, \quad m \mapsto m_{[0]} \otimes_L m_{[1]} := m^{[0]} \epsilon_{\mathcal{C}}(m^{[1]}_{[0]}) \otimes_L m^{[1]}_{[1]}, \quad \text{for } m \in M,$$

where $\varrho^M : m \mapsto m^{[0]} \otimes_A m^{[1]}$ denotes the $\mathcal{C}$ -coaction on M (note our convention to use character ϱ for $\mathcal{C}$ -coactions and upper indices of the Sweedler type to denote components of the coproduct and coactions of $\mathcal{C}$). It is straightforward to check that with this definition any right $\mathcal{C}$ -comodule map is $\mathcal{D}$ -colinear. In particular, a right $\mathcal{C}$ -coaction, being $\mathcal{C}$ -colinear by coassociativity, is $\mathcal{D}$ -colinear.

2.5. An **entwining structure** over a (not necessarily commutative) algebra L consists of an L -ring A , with multiplication μ and unit η , an L -coring $\mathcal{D}$, with comultiplication Δ and counit ϵ , and an L - L bilinear map $\psi : \mathcal{D} \otimes_L A \rightarrow A \otimes_L \mathcal{D}$, satisfying the following compatibility conditions.

$$\begin{aligned} \psi \circ (\mathcal{D} \otimes_L \mu) &= (\mu \otimes_L \mathcal{D}) \circ (A \otimes_L \psi) \circ (\psi \otimes_L A) \\ \psi \circ (\mathcal{D} \otimes_L \eta) &= \eta \otimes_L \mathcal{D} \\ (A \otimes_L \Delta) \circ \psi &= (\psi \otimes_L \mathcal{D}) \circ (\mathcal{D} \otimes_L \psi) \circ (\Delta \otimes_L A) \\ (A \otimes_L \epsilon) \circ \psi &= \epsilon \otimes_L A. \end{aligned}$$

In complete analogy with [Brz1, Proposition 2.2], $A \otimes_L \mathcal{D}$ is an A -coring. Its bimodule structure is given by

$$a_1(a \otimes_L d)a_2 = a_1a\psi(d \otimes_L a_2), \quad \text{for } a_1, a_2 \in A, a \otimes_L d \in A \otimes_L \mathcal{D}.$$

The coproduct is equal to $A \otimes_L \Delta : A \otimes_L \mathcal{D} \rightarrow A \otimes_L \mathcal{D} \otimes_L \mathcal{D} \cong (A \otimes_L \mathcal{D}) \otimes_A (A \otimes_L \mathcal{D})$ and the counit is $A \otimes_L \epsilon : A \otimes_L \mathcal{D} \rightarrow A$. Via the canonical isomorphism $M \otimes_A A \otimes_L \mathcal{D} \cong M \otimes_L \mathcal{D}$, for any right A -module M , right comodules for the A -coring $A \otimes_L \mathcal{D}$ are identified with **entwined modules**. A right-right entwined module means a right A -module and right $\mathcal{D}$ -comodule M , with coaction $\tau_M : m \mapsto m_{[0]} \otimes_L m_{[1]}$, such that

$$\tau_M(ma) = m_{[0]}\psi(m_{[1]} \otimes_L a), \quad \text{for } m \in M, a \in A.$$

Morphisms of entwined modules are A -linear and $\mathcal{D}$ -colinear maps. The category of right-right entwined modules is denoted by $\mathfrak{M}_A^{\mathcal{D}}(\psi)$.

Entwining structures $(A, \mathcal{D}, \psi)$ over an algebra L provide examples of coring extensions. Namely, the associated A -coring $\mathcal{C} := (A \otimes_L \mathcal{D}, A \otimes_L \Delta, A \otimes_L \epsilon)$ is a right $\mathcal{D}$ -comodule with coaction $\tau_{\mathcal{C}} := A \otimes_L \Delta : A \otimes_L \mathcal{D} \rightarrow A \otimes_L \mathcal{D} \otimes_L \mathcal{D}$. By the coassociativity of the coproduct Δ in $\mathcal{D}$, $\tau_{\mathcal{C}}$ is left $\mathcal{C}$ -colinear. This means that the L -coring $\mathcal{D}$ is right extension of the A -coring $\mathcal{C}$. In this situation the functor $\mathbb{R}$ in Figure (2.4) can be identified with the forgetful functor $\mathfrak{M}^{\mathcal{C}} \cong \mathfrak{M}_A^{\mathcal{D}}(\psi) \rightarrow \mathfrak{M}^{\mathcal{D}}$.

Let $(A, \mathcal{D}, \psi)$ be an entwining structure over an algebra L and $\mathcal{C} := A \otimes_L \mathcal{D}$ the associated A -coring. If e is a grouplike element in $\mathcal{D}$ then $1_A \otimes_L e$ is a grouplike element in $\mathcal{C}$. In this case A is a right $\mathcal{C}$ -comodule hence a right-right entwined module. The $\mathcal{D}$ -coaction in A comes out as

$$(2.7) \quad A \rightarrow A \otimes_L \mathcal{D}, \quad a \mapsto \psi(e \otimes a).$$

The coinvariants of a right $\mathcal{C}$ -comodule (i.e. entwined module) M with respect to $1_A \otimes_L e$ can be identified with $\text{Hom}^{\mathcal{C}}(A, M)$, and the coinvariants of M as a right $\mathcal{D}$ -comodule with respect to e can be identified with $\text{Hom}^{\mathcal{D}}(L, \mathbb{R}(M))$ (cf. [BW, 28.4]). Since in this case the forgetful functor $\mathbb{R} : \mathfrak{M}^{\mathcal{C}} \cong \mathfrak{M}_A^{\mathcal{D}}(\psi) \rightarrow \mathfrak{M}^{\mathcal{D}}$ possesses a left adjoint, $\bullet \otimes_L A$, it follows that $\text{Hom}^{\mathcal{D}}(L, \mathbb{R}(M)) \cong \text{Hom}^{\mathcal{C}}(A, M)$. That is to say, the coinvariants of a right $\mathcal{C}$ -comodule (i.e. entwined module) with respect to $1_A \otimes_L e$ are the same as its coinvariants as a right $\mathcal{D}$ -comodule with respect to e .

If the entwining map ψ is bijective then it induces an A -coring structure in $\mathcal{D} \otimes_L A$. Its left comodules are identified with left A -modules and left $\mathcal{D}$ -comodules, satisfying a compatibility condition with ψ . If there exists a grouplike element e in $\mathcal{D}$ then the corresponding left $\mathcal{D}$ -coaction in A is given by

$$(2.8) \quad A \rightarrow \mathcal{D} \otimes_L A, \quad a \mapsto \psi^{-1}(a \otimes_L e).$$

2.6. The notion of a **bialgebroid** over an algebra L was introduced by Takeuchi in [Ta] under the original name $\times_L$ -**bialgebra**. Takeuchi's definition was shown by Brzeziński and Militaru in [BM] to be equivalent to the structure introduced in [Lu]. As a k -bialgebra consists of compatible algebra and coalgebra structures on the same k -module, an L -bialgebroid comprises compatible $L \otimes_k L^{op}$ -ring and L -coring structures. More explicitly, a **left bialgebroid** is given by the data $(H, L, s_L, t_L, \gamma_L, \pi_L)$. Here H and L are k -algebras and $s_L : L \rightarrow H$ and $t_L : L^{op} \rightarrow H$ are algebra maps, called the source and target maps, respectively. The map

$$L \otimes_k L^{op} \rightarrow H, \quad l \otimes_k l' \mapsto s_L(l)t_L(l')$$

is required to be an algebra map, equipping H with the structure of an $L \otimes_k L^{op}$ -ring. The L - L bimodule H , with actions

$$(2.9) \quad lhl' = s_L(l)t_L(l')h, \quad \text{for } l, l' \in L, h \in H,$$

is required to be an L -coring with coproduct γ_L and counit π_L . For the coproduct we use a Sweedler type index notation with *lower* indexes, $\gamma_L(h) = h_{(1)} \otimes_L h_{(2)}$, for $h \in H$, where implicit summation is understood. The compatibility axioms between the $L \otimes_k L^{op}$ -ring and L -coring structures are the following. Consider the subset of the L -module tensor square of the bimodule (2.9), the so called Takeuchi product

$$H \times_L H = \{ \sum_i h_i \otimes_L h'_i \in H \otimes_L H \mid \forall l \in L \sum_i h_i t_L(l) \otimes_L h'_i = \sum_i h_i \otimes_L h'_i s_L(l) \}.$$

Note that $H \times_L H$ is an $L \otimes_k L^{op}$ -ring, with factorwise multiplication and unit map

$$L \otimes_k L^{op} \rightarrow H \times_L H, \quad l \otimes_k l' \mapsto s_L(l) \otimes_L t_L(l').$$

The first bialgebroid axiom asserts that the coproduct corestricts to a map of $L \otimes_k L^{op}$ -rings $H \rightarrow H \times_L H$. The requirement, that the range of the coproduct lies within $H \times_L H$, is referred to as the **Takeuchi axiom**. Further axioms require the counit to preserve the unit and satisfy

$$\pi_L(h s_L \circ \pi_L(h')) = \pi_L(h h') = \pi_L(h t_L \circ \pi_L(h')),$$

for all $h, h' \in H$.

The L - L bimodule (2.9) is defined in terms of left multiplication by the source and target maps. Symmetrically, one defines **right bialgebroids** by interchanging the roles of left and right

multiplications. Explicitly, a right bialgebroid is given by the data $(H, R, s_R, t_R, \gamma_R, \pi_R)$, where H and R are k -algebras and $s_R : R \rightarrow H$ and $t_R : R^{op} \rightarrow H$ are algebra maps, called the source and target maps, respectively. H is required to be an $R \otimes_k R^{op}$ -ring with unit

$$R \otimes_k R^{op} \rightarrow H, \quad r \otimes_k r' \mapsto s_R(r)t_R(r'),$$

and an R -coring, with bimodule structure

$$(2.10) \quad rhr' = hs_R(r')t_R(r), \quad \text{for } r, r' \in R, h \in H,$$

coproduct γ_R and counit π_R . For the coproduct we use a Sweedler type index notation with *upper* indices, $\gamma_R(h) = h^{(1)} \otimes_R h^{(2)}$, for $h \in H$, where implicit summation is understood. The coproduct is required to be a map of $R \otimes_k R^{op}$ -rings from H to the Takeuchi product

$$H \times_R H = \left\{ \sum_i h_i \otimes_R h'_i \in H \otimes_R H \mid \forall r \in R \sum_i s_R(r)h_i \otimes_R h'_i = \sum_i h_i \otimes_R t_R(r)h'_i \right\},$$

where the R -module tensor product is taken with respect to the bimodule structure (2.10). The counit is defined to preserve the unit and satisfy

$$\pi_R(s_R \circ \pi_R(h)h') = \pi_R(hh') = \pi_R(t_R \circ \pi_R(h)h'),$$

for all $h, h' \in H$. For more details we refer to [KSz].

The co-opposite $(H, L^{op}, t_L, s_L, \gamma_L^{op}, \pi_L)$ of a left bialgebroid $(H, L, s_L, t_L, \gamma_L, \pi_L)$ is a left bialgebroid too. The opposite $(H^{op}, L, t_L, s_L, \gamma_L, \pi_L)$ is a right bialgebroid.

2.7. A right comodule of a right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$ means a right comodule of the R -coring (H, γ_R, π_R) (with bimodule structure (2.10)). The category of right $\mathcal{H}_R$ -comodules is denoted by $\mathfrak{M}^{\mathcal{H}_R}$. By a definition in Section 2.3, a right $\mathcal{H}_R$ -comodule is in particular a right R -module. Using the bialgebroid structure of $\mathcal{H}_R$ (not its coring structure alone), one can introduce also a left R -module structure in a right $\mathcal{H}_R$ -comodule M ,

$$(2.11) \quad rm := m^{[0]} \pi_R(s_R(r)m^{[1]}), \quad \text{for } m \in M, r \in R.$$

This makes M an R - R bimodule such that the (so called Takeuchi) identity

$$rm^{[0]} \otimes_R m^{[1]} = m^{[0]} \otimes_R t_R(r)m^{[1]}$$

holds, for all $m \in M$ and $r \in R$. Any $\mathcal{H}_R$ -comodule map is R - R bilinear. This amounts to saying that there is a forgetful functor $\mathfrak{M}^{\mathcal{H}_R} \rightarrow {}_R\mathfrak{M}_R$. It was observed in [Scha2, Proposition 5.6] that the forgetful functor $\mathfrak{M}^{\mathcal{H}_R} \rightarrow {}_R\mathfrak{M}_R$ is strict monoidal. That is, $\mathfrak{M}^{\mathcal{H}_R}$ is a monoidal category via the R -module tensor product. The coaction in the product $M \otimes_R N$ of two right $\mathcal{H}_R$ -comodules M and N is

$$(m \otimes_R n)^{[0]} \otimes_R (m \otimes_R n)^{[1]} = (m^{[0]} \otimes_R n^{[0]}) \otimes_R m^{[1]}n^{[1]}, \quad \text{for } m \otimes_R n \in M \otimes_R N.$$

The monoidal unit is R with coaction given by the source map s_R . A **right $\mathcal{H}_R$ -comodule algebra** is an algebra in the monoidal category $\mathfrak{M}^{\mathcal{H}_R}$ (hence it is in particular an R -ring). Explicitly, it means an R -ring and right $\mathcal{H}_R$ -comodule A whose coaction ϱ^A satisfies

$$\varrho^A(1_A) = 1_A \otimes_R 1_H, \quad \varrho^A(aa') = a^{[0]}a'^{[0]} \otimes_R a^{[1]}a'^{[1]}, \quad \text{for } a, a' \in A.$$

The R -coring (H, γ_R, π_R) underlying $\mathcal{H}_R$ possesses a grouplike element 1_H . Coinvariants of a right $\mathcal{H}_R$ -comodule are meant always with respect to the distinguished grouplike element 1_H . By the R - R bilinearity of the coaction ϱ^A in a right $\mathcal{H}_R$ -comodule algebra A , for any element r in R and any coinvariant b in A , the unit map $\eta : R \rightarrow A$ satisfies

$$\varrho^A(b\eta(r)) = b \otimes_R s_R(r) = \varrho^A(\eta(r)b).$$

Hence the elements $b \in A^{co\mathcal{H}_R}$ and $\eta(r)$, for $r \in R$, commute in A .

Left comodules of a right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$ (i.e. of the R -coring (H, γ_R, π_R)) are treated symmetrically. Their category is denoted by ${}^{\mathcal{H}_R}\mathfrak{M}$. A left $\mathcal{H}_R$ -comodule

M (which is a priori a left R -module) can be equipped with an R - R bimodule structure with right R -action

$$mr = \pi_R(s_R(r)m^{[-1]})m^{[0]} \quad \text{for } m \in M, r \in R.$$

The forgetful functor $\mathcal{H}_R\mathfrak{M} \rightarrow {}_{R^{op}}\mathfrak{M}_{R^{op}}$ is strict monoidal. For two left $\mathcal{H}_R$ -comodules M and N , the left and right R -actions and the left $\mathcal{H}_R$ -coaction in the product $M \otimes_{R^{op}} N$ take the form

$$r(m \otimes_{R^{op}} n)r' = mr' \otimes_{R^{op}} rn, \quad (m \otimes_{R^{op}} n)^{[-1]} \otimes_R (m \otimes_{R^{op}} n)^{[0]} = m^{[-1]}n^{[-1]} \otimes_R (m^{[0]} \otimes_{R^{op}} n^{[0]}),$$

for $r, r' \in R$ and $m \otimes_{R^{op}} n \in M \otimes_{R^{op}} N$. The monoidal unit is R^{op} , with coaction given by the target map t_R . A **left $\mathcal{H}_R$ -comodule algebra** is defined as an algebra in the monoidal category $\mathcal{H}_R\mathfrak{M}$. It is in particular an R^{op} -ring. Explicitly, a left $\mathcal{H}_R$ -comodule algebra is an R^{op} -ring and left $\mathcal{H}_R$ -comodule A , whose coaction ${}^A\varrho$ satisfies

$${}^A\varrho(1_A) = 1_H \otimes_R 1_A, \quad {}^A\varrho(aa') = a^{[-1]}a'^{[-1]} \otimes_R a^{[0]}a'^{[0]}, \quad \text{for } a, a' \in A.$$

Coinvariants of left $\mathcal{H}_R$ -comodules are meant always with respect to the distinguished grouplike element 1_H .

Comodules of left bialgebroids can be described symmetrically. For a right bialgebroid $\mathcal{H}_R$, the categories $(\mathcal{H}_R)^{cop}\mathfrak{M}$ and $\mathfrak{M}^{\mathcal{H}_R}$ are monoidally isomorphic. The categories $\mathfrak{M}^{(\mathcal{H}_R)^{op}}$ and $\mathfrak{M}^{\mathcal{H}_R}$ are anti-monoidally isomorphic.

2.8. Let $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$ be a right bialgebroid and A a right $\mathcal{H}_R$ -comodule algebra with coaction $a \mapsto a^{[0]} \otimes_R a^{[1]}$. Recall from Section 2.7 that A possesses an R -ring structure. The R -ring A , the R -coring (H, γ_R, π_R) and the R - R bimodule map

$$(2.12) \quad \psi_R : H \otimes_R A \rightarrow A \otimes_R H, \quad h \otimes_R a \mapsto a^{[0]} \otimes_R ha^{[1]}$$

form an entwining structure over R . This implies that $A \otimes_R H$ is an A -coring, with bimodule structure

$$a_1(a \otimes_R h)a_2 = a_1aa_2^{[0]} \otimes_R ha_2^{[1]}, \quad \text{for } a_1, a_2 \in A \text{ and } a \otimes_R h \in A \otimes_R H,$$

coproduct $A \otimes_R \gamma_R : A \otimes_R H \rightarrow A \otimes_R H \otimes_R H \cong (A \otimes_R H) \otimes_A (A \otimes_R H)$ and counit $A \otimes_R \pi_R : A \otimes_R H \rightarrow A$. Via the canonical isomorphism $M \otimes_A (A \otimes_R H) \cong M \otimes_R H$, for $M \in \mathfrak{M}_A$, right comodules for the A -coring $(A \otimes_R H, A \otimes_R \gamma_R, A \otimes_R \pi_R)$ can be identified with right-right entwined modules for the entwining structure (A, H, ψ_R) . Such entwined modules are also called **right-right $(A, \mathcal{H}_R)$ -relative Hopf modules**. They can be described equivalently as right modules for the algebra A in the category of right $\mathcal{H}_R$ -comodules. That is, right A -modules and right $\mathcal{H}_R$ -comodules M , such that the A -action is $\mathcal{H}_R$ -colinear, in the sense that the compatibility condition

$$(ma)^{[0]} \otimes_R (ma)^{[1]} = m^{[0]}a^{[0]} \otimes_R m^{[1]}a^{[1]}$$

holds, for $m \in M$, $a \in A$. The category of right-right $(A, \mathcal{H}_R)$ -relative Hopf modules will be denoted by $\mathfrak{M}_A^{\mathcal{H}_R}$. As it is explained in Section 2.5, in the R -entwining structure (A, H, ψ_R) the R -coring (H, γ_R, π_R) is a right extension of the A -coring $(A \otimes_R H, A \otimes_R \gamma_R, A \otimes_R \pi_R)$. For this coring extension, the functor $\mathbb{R}$ on Figure (2.4) can be identified with the forgetful functor $\mathfrak{M}_A^{\mathcal{H}_R} \rightarrow \mathfrak{M}^{\mathcal{H}_R}$.

A right $\mathcal{H}_R$ -comodule algebra A is called an **$\mathcal{H}_R$ -Galois extension** of its coinvariant subalgebra B if the associated A -coring $(A \otimes_R H, A \otimes_R \gamma_R, A \otimes_R \pi_R)$, with grouplike element $1_A \otimes_R 1_H$, is a Galois coring, i.e. the canonical map

$$(2.13) \quad \text{can} : A \otimes_B A \rightarrow A \otimes_R H, \quad a \otimes_B a' \mapsto aa'^{[0]} \otimes_R a'^{[1]}$$

is bijective.

For a left comodule algebra A' for a left bialgebroid $\mathcal{H}_L$ one defines left-left $(A', \mathcal{H}_L)$ -relative Hopf modules in a symmetrical way.

2.9. A **Hopf algebroid** [BSz], [Bö1] is a triple $\mathcal{H} = (\mathcal{H}_L, \mathcal{H}_R, S)$. It consists of a left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$ and a right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$ on the *same* total algebra H . They are subject to the following compatibility axioms

$$(2.14) \quad \begin{aligned} s_R \circ \pi_R \circ t_L &= t_L & s_L \circ \pi_L \circ t_R &= t_R \\ t_R \circ \pi_R \circ s_L &= s_L & t_L \circ \pi_L \circ s_R &= s_R \end{aligned}$$

and

$$(2.15) \quad (\gamma_R \otimes_L H) \circ \gamma_L = (H \otimes_R \gamma_L) \circ \gamma_R, \quad (\gamma_L \otimes_R H) \circ \gamma_R = (H \otimes_L \gamma_R) \circ \gamma_L.$$

The k -linear map $S : H \rightarrow H$ is called the **antipode**. It is required to be R - L bilinear in the sense that

$$S(t_L(l)ht_R(r)) = s_R(r)S(h)s_L(l), \quad \text{for } l \in L, r \in R, h \in H.$$

The antipode axioms read as

$$\mu \circ (H \otimes_R S) \circ \gamma_R = s_L \circ \pi_L, \quad \mu \circ (S \otimes_L H) \circ \gamma_L = s_R \circ \pi_R,$$

where μ denotes the multiplication both in the L -ring $s_L : L \rightarrow H$ and the R -ring $s_R : R \rightarrow H$.

In a Hopf algebroid there are two bialgebroid (hence two coring) structures present. Throughout this paper we insist on using upper indices of the Sweedler type to denote components of the coproduct and coactions of the right bialgebroid $\mathcal{H}_R$, and lower indices in the case of the left bialgebroid $\mathcal{H}_L$.

Similarly to the case of Hopf algebras, the antipode of a Hopf algebroid $\mathcal{H} = (\mathcal{H}_L, \mathcal{H}_R, S)$ is an anti-algebra map on the total algebra H . That is,

$$S(1_H) = 1_H \quad \text{and} \quad S(hh') = S(h')S(h), \quad \text{for } h, h' \in H.$$

It is also an anti-coring map $\mathcal{H}_L \rightarrow \mathcal{H}_R$ and $\mathcal{H}_R \rightarrow \mathcal{H}_L$. That is,

$$\begin{aligned} \pi_R \circ S &= \pi_R \circ s_L \circ \pi_L \quad \text{and} \quad S(h)^{(1)} \otimes_R S(h)^{(2)} = S(h_{(2)}) \otimes_R S(h_{(1)}), \\ \pi_L \circ S &= \pi_L \circ s_R \circ \pi_R \quad \text{and} \quad S(h)_{(1)} \otimes_L S(h)_{(2)} = S(h^{(2)}) \otimes_L S(h^{(1)}), \end{aligned} \quad \text{for } h \in H.$$

For a Hopf algebroid $\mathcal{H} = (\mathcal{H}_L, \mathcal{H}_R, S)$, also the opposite-co-opposite $\mathcal{H}_{cop}^{op} = ((\mathcal{H}_R)_{cop}^{op}, (\mathcal{H}_L)_{cop}^{op}, S)$ is a Hopf algebroid. If the antipode S is bijective then so are the opposite $\mathcal{H}^{op} = ((\mathcal{H}_R)^{op}, (\mathcal{H}_L)^{op}, S^{-1})$ and the co-opposite $\mathcal{H}_{cop} = ((\mathcal{H}_L)_{cop}, (\mathcal{H}_R)_{cop}, S^{-1})$, too.

2.10. Note that axiom (2.14) implies that the base algebras R and L in a Hopf algebroid $(\mathcal{H}_L, \mathcal{H}_R, S)$ are anti-isomorphic via the map $\pi_R \circ t_L : L^{op} \rightarrow R$. This gives rise to a monoidal isomorphism, given by the ‘restriction of scalars’ functors

$$L^{op} \mathfrak{M}_{L^{op}} \xrightleftharpoons[\mathbb{V}]{\tilde{\mathbb{V}}} R \mathfrak{M}_R.$$

Furthermore, axiom (2.15) implies that the L -coring (H, γ_L, π_L) is a right extension of the R -coring (H, γ_R, π_R) and vice versa. By considerations in Section 2.4, this gives rise to a couple of functors

$$\mathfrak{M}^{\mathcal{H}_L} \xrightleftharpoons[\mathbb{R}]{\tilde{\mathbb{R}}} \mathfrak{M}^{\mathcal{H}_R},$$

which were proven to be inverse monoidal isomorphisms in [BB2, Theorem 2.2]. Explicitly, the functor $\mathbb{R}$ maps a right $\mathcal{H}_R$ -comodule M , with coaction $m \mapsto m^{[0]} \otimes_R m^{[1]}$, to the right $\mathcal{H}_L$ -comodule M , with right L -action and $\mathcal{H}_L$ -coaction

$$(2.16) \quad ml = \pi_R \circ t_L(l)m, \quad \text{and} \quad m \mapsto m_{[0]} \otimes_L m_{[1]} \equiv m^{[0]} \pi_R(m^{[1]}_{(1)}) \otimes_L m^{[1]}_{(2)}.$$

Analogously to (2.11), the right $\mathcal{H}_L$ -comodule (2.16) can be made an L - L bimodule with left L -action

$$lm := m_{[0]} \pi_L(m_{[1]} s_L(l)) = m \pi_R \circ t_L(l).$$

The functor $\tilde{\mathbb{R}}$ maps a right $\mathcal{H}_L$ -comodule N , with coaction $n \mapsto n_{[0]} \otimes_L n_{[1]}$, to the right $\mathcal{H}_R$ -comodule N , with right R -action and $\mathcal{H}_R$ -coaction

$$(2.17) \quad nr = \pi_L \circ s_R(r)n, \quad \text{and} \quad n \mapsto n^{[0]} \otimes_R n^{[1]} \equiv n_{[0]} \pi_L(n_{[1]}^{(1)}) \otimes_R n_{[1]}^{(2)}.$$

As we have seen in Section 2.7, N is an R - R bimodule with left R -action (2.11),

$$rn := n^{[0]} \pi_R(s_R(r)n^{[1]}) = n \pi_L \circ s_R(r).$$

Summarising, we conclude on the commutativity of the following diagram

$$(2.18) \quad \begin{array}{ccc} \mathfrak{M}^{\mathcal{H}_L} & \begin{array}{c} \xrightarrow{\tilde{\mathbb{R}}} \\ \xleftarrow{\mathbb{R}} \end{array} & \mathfrak{M}^{\mathcal{H}_R} \\ \downarrow & & \downarrow \\ {}_{L^{op}}\mathfrak{M}_{{}_L^{op}} & \begin{array}{c} \xrightarrow{\tilde{\mathbb{V}}} \\ \xleftarrow{\mathbb{V}} \end{array} & {}_R\mathfrak{M}_R \end{array}$$

where the vertical arrows denote the forgetful functors, described in Section 2.7. Since in this way the categories $\mathfrak{M}^{\mathcal{H}_L}$ and $\mathfrak{M}^{\mathcal{H}_R}$ are canonically isomorphic (i.e. the isomorphism functors $\mathbb{R}$ and $\tilde{\mathbb{R}}$ are compatible with the forgetful functors to $\mathfrak{M}_k$, cf. Section 2.4), in what follows we do not distinguish them. We speak simply about the category $\mathfrak{M}^{\mathcal{H}}$ of right comodules of the Hopf algebroid $\mathcal{H}$. It is straightforward to see that coinvariants of a right $\mathcal{H}$ -comodule M with respect to $\mathcal{H}_L$ and $\mathcal{H}_R$ are the same. We refer to the equivalent notions of an $\mathcal{H}_L$ -comodule algebra and an $\mathcal{H}_R$ -comodule algebra as an $\mathcal{H}$ -comodule algebra.

2.11. Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$, and antipode S , and let A be a right $\mathcal{H}$ -comodule algebra. In addition to its A -coring structure, described in Section 2.8, $A \otimes_R H$ is also a right comodule for the L -coring (H, γ_L, π_L) , with coaction $A \otimes_R \gamma_L : A \otimes_R H \rightarrow A \otimes_R H \otimes_L H$. The coproduct $A \otimes_R \gamma_R$ in $A \otimes_R H$ is right (H, γ_L, π_L) -colinear. That is to say, the L -coring (H, γ_L, π_L) is a *right extension* of the A -coring $(A \otimes_R H, A \otimes_R \gamma_R, A \otimes_R \pi_R)$. As we identify the categories of comodules for the R -coring (H, γ_R, π_R) and for the L -coring (H, γ_L, π_L) , the functor $\mathbb{R}$ on Figure (2.4) for this coring extension can be identified with the forgetful functor $\mathfrak{M}_A^{\mathcal{H}} \rightarrow \mathfrak{M}^{\mathcal{H}}$ as well.

2.12. Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$, and antipode S , and let A be a right $\mathcal{H}$ -comodule algebra. This means in particular that A is a right comodule algebra for the right R -bialgebroid $\mathcal{H}_R$, with coaction $a \mapsto a^{[0]} \otimes_R a^{[1]}$. Hence one can consider right-right $(A, \mathcal{H}_R)$ -relative Hopf modules as in Section 2.8. What is more, since A is a right comodule algebra for the left bialgebroid $\mathcal{H}_L$ as well, with coaction $a \mapsto a_{[0]} \otimes_L a_{[1]}$, related to the $\mathcal{H}_R$ -coaction as in Section 2.10, the opposite algebra A^{op} is a right comodule algebra for the right L -bialgebroid $(\mathcal{H}_L)^{op}$. Hence in addition to the R -entwining structure (2.12), A determines also an L -entwining structure. It consists of the L -ring A^{op} (with unit, expressed in terms of the unit η of the R -ring A as $\eta \circ \pi_R \circ t_L$), the L -coring (H, γ_L, π_L) , and the entwining map

$$(2.19) \quad \psi_L : H \otimes_L A \rightarrow A \otimes_L H, \quad h \otimes_L a \mapsto a_{[0]} \otimes_L a_{[1]} h.$$

Therefore there is an associated A^{op} -coring structure on $A \otimes_L H$, whose right comodules can be identified with entwined modules for the L -entwining structure (A^{op}, H, ψ_L) . These entwined modules are the same as the right-right $(A^{op}, (\mathcal{H}_L)^{op})$ -relative Hopf modules and can be characterised equivalently as left modules for the algebra A in the category of right $\mathcal{H}$ -comodules. Their category is denoted by ${}_A\mathfrak{M}^{\mathcal{H}}$.

For a left $\mathcal{H}$ -comodule algebra A' (which is in particular an L -ring) one can consider left-left $(A', \mathcal{H}_L)$ -, and $(A'^{op}, (\mathcal{H}_R)^{op})$ -relative Hopf modules in a symmetrical way.

Note that the entwining map (2.19) is bijective with inverse $\psi_L^{-1}(a \otimes_L h) = S(a^{[1]})h \otimes_L a^{[0]}$. Hence $H \otimes_L A$ has a unique A^{op} -coring structure such that (2.19) is an isomorphism of corings. Clearly, by the existence of grouplike elements, A^{op} is a left comodule for the A^{op} -corings $A \otimes_L H \cong H \otimes_L A$. Hence it is a left $\mathcal{H}$ -comodule algebra. The left L -action in A is determined by the unit map of the corresponding L -ring A^{op} , $\eta \circ \pi_R \circ t_L : L \rightarrow A^{op}$ (see above), and the left $\mathcal{H}_L$ -coaction in A is given as

$$(2.20) \quad a \mapsto S(a^{[1]}) \otimes_L a^{[0]}.$$

Left comodules for the isomorphic A^{op} -corings $A \otimes_L H \cong H \otimes_L A$ can be identified with left-left $(A^{op}, \mathcal{H}_L)$ -relative Hopf modules, i.e. left modules for the algebra A^{op} in the monoidal category

of left $\mathcal{H}$ -comodules. This means right A -modules left $\mathcal{H}$ -comodules, such that the A -action is left $\mathcal{H}$ -colinear. Their category will be denoted by ${}^{\mathcal{H}}\mathfrak{M}_A$.

2.13. Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$, and a *bijective* antipode S . Let M be a right $\mathcal{H}$ -comodule with $\mathcal{H}_R$ -coaction $m \mapsto m^{[0]} \otimes_R m^{[1]}$ and $\mathcal{H}_L$ -coaction $m \mapsto m_{[0]} \otimes_L m_{[1]}$, related via (2.16)-(2.17). Then M has a left $\mathcal{H}_R$ -comodule structure with left R -action (2.11) and coaction

$$(2.21) \quad m \mapsto S^{-1}(m_{[1]}) \otimes_R m_{[0]}.$$

The isomorphism functor between the categories of left $\mathcal{H}_R$ -comodules and left $\mathcal{H}_L$ -comodules (i.e. a left comodules version of the functor $\mathbb{R}$ in Section 2.10) maps the left $\mathcal{H}_R$ -comodule M in (2.21) to a left $\mathcal{H}_L$ -comodule M , with left L -action $lm := m\pi_R \circ s_L(l)$, for $l \in L$ and $m \in M$, and coaction

$$(2.22) \quad m \mapsto S^{-1}(m^{[1]}) \otimes_L m^{[0]}.$$

Above considerations lead to the following. If the antipode of $\mathcal{H}$ is bijective, the commutative diagram (2.18) extends to the following commutative diagram.

$$(2.23) \quad \begin{array}{ccc} \mathfrak{M}^{\mathcal{H}_L} & \xrightleftharpoons[\mathbb{R}]{\tilde{\mathbb{R}}} & \mathfrak{M}^{\mathcal{H}_R} \\ \downarrow (2.21) & \swarrow L^{op}\mathfrak{M}_{L^{op}} \quad \tilde{\mathbb{V}} \quad \searrow \mathbb{V} & \downarrow (2.22) \\ & R\mathfrak{M}_R & \\ \uparrow (\mathcal{H}_R)^{op}\mathfrak{M} & \nwarrow \mathbb{W} \quad \tilde{\mathbb{W}} \quad \nearrow L^{op}\mathfrak{M}_{L^{op}} & \uparrow (\mathcal{H}_L)^{op}\mathfrak{M} \\ & \xleftarrow[\mathbb{L}]{\tilde{\mathbb{L}}} & \end{array}$$

The unlabeled functors are forgetful functors. The vertical arrows denote monoidal isomorphisms, induced by relations (2.21) and (2.22) between coactions. The functors $\mathbb{L}$ and $\tilde{\mathbb{L}}$ are the analogues of $\mathbb{R}$ and $\tilde{\mathbb{R}}$ for left comodules. The functors $\mathbb{W}$ and $\tilde{\mathbb{W}}$ are restriction of scalars functors, induced by the inverse algebra isomorphisms

$$\pi_R \circ s_L : L^{op} \rightarrow R, \quad \text{and} \quad \pi_L \circ t_R : R \rightarrow L^{op},$$

respectively.

2.14. Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$, and a *bijective* antipode S . Let A be a right $\mathcal{H}$ -comodule algebra with $\mathcal{H}_R$ -coaction $a \mapsto a^{[0]} \otimes_R a^{[1]}$ and $\mathcal{H}_L$ -coaction $a \mapsto a_{[0]} \otimes_L a_{[1]}$, related via (2.16)-(2.17). Denote the unit map in the corresponding R -ring by $\eta : R \rightarrow A$. By considerations in Section 2.13, A defines an algebra in all of the isomorphic monoidal comodule categories in Figure (2.23). This implies that the two isomorphic A^{op} -corings $A \otimes_L H \cong H \otimes_L A$ in Section 2.12 are anti-isomorphic to the A -corings $A \otimes_R H \cong H \otimes_R A$. An anti-isomorphism is given by a bijection in [Bö2, Lemma 3.3],

$$(2.24) \quad A \otimes_R H \rightarrow A \otimes_L H, \quad a \otimes_R h \mapsto a_{[0]} \otimes_L a_{[1]} S(h),$$

and an isomorphism $H \otimes_R A \rightarrow A \otimes_R H$ is given by the entwining map (2.12), with inverse

$$(2.25) \quad a \otimes_R h \mapsto h S^{-1}(a_{[1]}) \otimes_R a_{[0]},$$

cf. [Bö2, Lemma 4.1]. By the existence of grouplike elements, A is a left comodule for all these corings. This amounts to saying that A is a left comodule algebra for the opposite Hopf algebroid $\mathcal{H}^{op}$. The R -ring structure $\eta : R \rightarrow A$ is the same underlying the right $\mathcal{H}_R$ -comodule algebra structure of A and the left $\mathcal{H}_R$ -coaction in A is given as

$$(2.26) \quad a \mapsto S^{-1}(a_{[1]}) \otimes_R a_{[0]}.$$

The isomorphism functor $\tilde{\mathbb{L}}$ in Figure (2.23) maps the left $\mathcal{H}_R$ -comodule A in (2.26) to a left $\mathcal{H}_L$ -comodule A , with L^{op} -ring structure $\eta \circ \pi_R \circ s_L : L^{op} \rightarrow A$ and coaction

$$(2.27) \quad a \mapsto S^{-1}(a^{[1]}) \otimes_L a^{[0]}.$$

The left $\mathcal{H}_L$ -coaction (2.27) differs from (2.20) by the isomorphism $S^2 \otimes_L A$.

Left comodules for the isomorphic A -corings $A \otimes_R H \cong H \otimes_R A$ can be identified with left-left $(A, (\mathcal{H}_R)^{op})$ -relative Hopf modules. That is, left A -modules left $\mathcal{H}_R$ -comodules M , such that

$$(2.28) \quad (am)^{[-1]} \otimes_R (am)^{[0]} = m^{[-1]} S^{-1}(a_{[1]}) \otimes_R a_{[0]} m^{[0]}, \quad \text{for } a \in A, m \in M.$$

Since this means actually left A -modules and left $\mathcal{H}$ -comodules M with left $\mathcal{H}$ -colinear A -action, we denote their category by ${}^{\mathcal{H}}_A \mathfrak{M}$.

3. RELATIVE SEPARABLE FUNCTORS

We start by recalling some material about separable functors. For more information we refer to [HS, Chap. IX, page 307-312], [We, Chap. 8, page 279-281] and [CMZ]. Throughout the paper we use the following terminology, used already in Section 2.3. A morphism $f : C_1 \rightarrow C_2$ in a category $\mathfrak{C}$ is said to be a *split monomorphism* or *section* if it is *cosplit* by some morphism $h : C_2 \rightarrow C_1$ in $\mathfrak{C}$, i.e. $h \circ f = C_1$. Dually, f is called a *split epimorphism* or *retraction* provided that it is *split* by some morphism $g : C_2 \rightarrow C_1$ in $\mathfrak{C}$, i.e. $f \circ g = C_2$.

DEFINITION 3.1. Let $\mathfrak{C}$ be a category and let $\mathcal{S}$ be a class of morphisms in $\mathfrak{C}$. For a morphism $f : C_1 \rightarrow C_2$ in $\mathfrak{C}$, an object $P \in \mathfrak{C}$ is called *f-projective* if the map $\text{Hom}_{\mathfrak{C}}(P, f) : \text{Hom}_{\mathfrak{C}}(P, C_1) \rightarrow \text{Hom}_{\mathfrak{C}}(P, C_2)$ is surjective. P is *$\mathcal{S}$ -projective* if it is *f-projective* for every $f \in \mathcal{S}$.

Dually, an object $I \in \mathfrak{C}$ is called *f-injective* if it is *f^{op}-projective* in the opposite category $\mathfrak{C}^{op}$, where $f^{op} : C_2 \rightarrow C_1$ is considered to be a morphism $\mathfrak{C}^{op}$. I is called *$\mathcal{S}$ -injective* if it is *f-injective* for every $f \in \mathcal{S}$.

All results below about projective objects can be dualised to get their analogues for injective objects.

THEOREM 3.2. [Ar] Let $\mathbb{H} : \mathfrak{B} \rightarrow \mathfrak{A}$ be a covariant functor and consider a class of morphisms

$$(3.1) \quad \mathcal{E}_{\mathbb{H}} := \{g \in \mathfrak{B} \mid \mathbb{H}(g) \text{ is a split epimorphism in } \mathfrak{A}\}.$$

Assume that $\mathbb{T} : \mathfrak{A} \rightarrow \mathfrak{B}$ is a left adjoint of $\mathbb{H}$ and denote by $\varepsilon : \mathbb{T}\mathbb{H} \rightarrow \mathfrak{B}$ the counit of the adjunction. Then, for an object $P \in \mathfrak{B}$, the following assertions are equivalent.

- (a) P is $\mathcal{E}_{\mathbb{H}}$ -projective.
- (b) $\varepsilon_P : \mathbb{T}\mathbb{H}(P) \rightarrow P$ is a split epimorphism.
- (c) There is a split epimorphism $\pi : \mathbb{T}(X) \rightarrow P$, for a suitable object $X \in \mathfrak{A}$.

In particular, all objects of the form $\mathbb{T}(X)$, for $X \in \mathfrak{A}$, are $\mathcal{E}_{\mathbb{H}}$ -projective.

For completeness we include the dual version of Theorem 3.2.

THEOREM 3.3. [Ar] Let $\mathbb{T} : \mathfrak{A} \rightarrow \mathfrak{B}$ be a covariant functor and consider a class of morphisms

$$(3.2) \quad \mathcal{I}_{\mathbb{T}} := \{g \in \mathfrak{A} \mid \mathbb{T}(g) \text{ is a split monomorphism in } \mathfrak{B}\}.$$

Assume that $\mathbb{H} : \mathfrak{B} \rightarrow \mathfrak{A}$ is a right adjoint of $\mathbb{T}$ and denote by $\eta : \mathfrak{A} \rightarrow \mathbb{H}\mathbb{T}$ the unit of the adjunction. Then, for an object $I \in \mathfrak{A}$, the following assertions are equivalent.

- (a) I is $\mathcal{I}_{\mathbb{T}}$ -injective.
- (b) $\eta_I : I \rightarrow \mathbb{H}\mathbb{T}(I)$ is a split monomorphism.
- (c) There is a split monomorphism $i : I \rightarrow \mathbb{H}(Y)$, for a suitable object $Y \in \mathfrak{B}$.

In particular, all objects of the form $\mathbb{H}(Y)$, for $Y \in \mathfrak{B}$, are $\mathcal{I}_{\mathbb{T}}$ -injective.

Using the current terminology, relative injective right comodules of an A -coring $\mathcal{C}$, discussed in Section 2.3, can be characterised as $\mathcal{I}_{\mathbb{U}}$ -injective objects, where $\mathbb{U} : \mathfrak{M}^{\mathcal{C}} \rightarrow \mathfrak{M}_A$ denotes the forgetful functor. As recalled in Section 2.3, the forgetful functor $\mathbb{U}$ possesses a right adjoint, the functor $\bullet \otimes_A \mathcal{C}$. The unit of the adjunction is given by the $\mathcal{C}$ -coaction. Therefore, Theorem 3.3

(a) $\Leftrightarrow$ (b) includes the claim, recalled in Section 2.3, that a right $\mathcal{C}$ -comodule M is relative injective if and only if the coaction ϱ^M in it is a split monomorphism in $\mathfrak{M}^{\mathcal{C}}$.

Since any covariant functor preserves split epimorphisms and split monomorphisms, we immediately have that, for any two functors $\mathbb{F} : \mathfrak{A} \rightarrow \mathfrak{B}$ and $\mathbb{G} : \mathfrak{B} \rightarrow \mathfrak{C}$,

$$(3.3) \quad \mathcal{E}_{\mathbb{F}} \subseteq \mathcal{E}_{\mathbb{G}\mathbb{F}} \quad \text{and} \quad \mathcal{I}_{\mathbb{F}} \subseteq \mathcal{I}_{\mathbb{G}\mathbb{F}}.$$

As explained in the Introduction, in the field of Schneider type theorems one often faces the following problem. Consider an entwining structure $(A, \mathcal{D}, \psi)$ over an algebra L . Assume that some map in $\mathfrak{M}_A^{\mathcal{D}}(\psi)$ (practically the canonical map) is a retraction in $\mathfrak{M}_L$. Under what assumptions is it a retraction also in $\mathfrak{M}^{\mathcal{D}}$? Putting the question in a more functorial way, we can ask in which cases is $\mathcal{E}_{\mathbb{F}} = \mathcal{E}_{\mathbb{G}\mathbb{F}}$, for the forgetful functors $\mathbb{F} : \mathfrak{M}_A^{\mathcal{D}}(\psi) \rightarrow \mathfrak{M}^{\mathcal{D}}$ and $\mathbb{G} : \mathfrak{M}^{\mathcal{D}} \rightarrow \mathfrak{M}_L$. For these particular functors $\mathbb{F}$ and $\mathbb{G}$, property 1) in Proposition 3.4 below reduces to a similar (but somewhat weaker) assumption as in a Schneider type theorem [SS, Theorem 5.9] (see also [Brz2, Theorem 4.6]). Properties like in part 2) of Proposition 3.4 are assumed e.g. in [SS, Corollary 4.8].

PROPOSITION 3.4. *For two functors $\mathbb{F} : \mathfrak{A} \rightarrow \mathfrak{B}$ and $\mathbb{G} : \mathfrak{B} \rightarrow \mathfrak{C}$, $\mathcal{E}_{\mathbb{F}} = \mathcal{E}_{\mathbb{G}\mathbb{F}}$ whenever any of the following properties hold.*

- 1) $\mathbb{F}(A)$ is $\mathcal{E}_{\mathbb{G}}$ -projective, for every object $A \in \mathfrak{A}$.
- 2) $\mathfrak{A}, \mathfrak{B}$ and $\mathfrak{C}$ are abelian categories, $\mathbb{G}$ is left exact and reflects epimorphisms, $\mathbb{F}$ is left exact and $\mathbb{F}(A)$ is $\mathcal{I}_{\mathbb{G}}$ -injective, for every object $A \in \mathfrak{A}$.

Dually, $\mathcal{I}_{\mathbb{F}} = \mathcal{I}_{\mathbb{G}\mathbb{F}}$ whenever any of the following properties hold.

- 1^{op}) $\mathbb{F}(A)$ is $\mathcal{I}_{\mathbb{G}}$ -injective, for every object $A \in \mathfrak{A}$.
- 2^{op}) $\mathfrak{A}, \mathfrak{B}$ and $\mathfrak{C}$ are abelian categories, $\mathbb{G}$ is right exact and reflects monomorphisms, $\mathbb{F}$ is right exact and $\mathbb{F}(A)$ is $\mathcal{E}_{\mathbb{G}}$ -projective, for every object $A \in \mathfrak{A}$.

Proof. 1) Let $f : A_1 \rightarrow A_2$ be a morphism in $\mathcal{E}_{\mathbb{G}\mathbb{F}}$. Then $\mathbb{F}(f) : \mathbb{F}(A_1) \rightarrow \mathbb{F}(A_2)$ belongs to $\mathcal{E}_{\mathbb{G}}$ and hence, by hypothesis, it is a split epimorphism. Thus $f \in \mathcal{E}_{\mathbb{F}}$.

2) For $f \in \mathcal{E}_{\mathbb{G}\mathbb{F}}$, consider the exact sequence (kernel diagram)

$$0 \rightarrow K \xrightarrow{i} A_1 \xrightarrow{f} A_2$$

in $\mathfrak{A}$. The left exact functor $\mathbb{F}$ takes it to the exact sequence

$$0 \longrightarrow \mathbb{F}(K) \xrightarrow{\mathbb{F}(i)} \mathbb{F}(A_1) \xrightarrow{\mathbb{F}(f)} \mathbb{F}(A_2)$$

in $\mathfrak{B}$. Since f is an element of $\mathcal{E}_{\mathbb{G}\mathbb{F}}$, the morphism $\mathbb{G}\mathbb{F}(f)$ is a split epimorphism. Since $\mathbb{G}$ is left exact and $\mathfrak{C}$ is an abelian category, the sequence

$$0 \longrightarrow \mathbb{G}\mathbb{F}(K) \xrightarrow{\mathbb{G}\mathbb{F}(i)} \mathbb{G}\mathbb{F}(A_1) \xrightarrow{\mathbb{G}\mathbb{F}(f)} \mathbb{G}\mathbb{F}(A_2) \longrightarrow 0$$

in $\mathfrak{C}$ is split exact. Thus we deduce that $i \in \mathcal{I}_{\mathbb{G}\mathbb{F}}$. Moreover, since $\mathbb{G}$ reflects epimorphisms, $\mathbb{F}(f)$ is an epimorphism. So the sequence

$$0 \longrightarrow \mathbb{F}(K) \xrightarrow{\mathbb{F}(i)} \mathbb{F}(A_1) \xrightarrow{\mathbb{F}(f)} \mathbb{F}(A_2) \longrightarrow 0$$

in $\mathfrak{B}$ is exact too. Since i is an element of $\mathcal{I}_{\mathbb{G}\mathbb{F}}$, its image $\mathbb{F}(i)$ is in $\mathcal{I}_{\mathbb{G}}$. By assumption $\mathbb{F}(K)$ is $\mathcal{I}_{\mathbb{G}}$ -injective hence the monomorphism $\mathbb{F}(i)$ is split. Since $\mathfrak{B}$ is an abelian category, we conclude that $\mathbb{F}(f)$ is a split epimorphism, i.e. that $f \in \mathcal{E}_{\mathbb{F}}$.

Claims 1^{op}) and 2^{op}) follow by duality. $\square$

The most important notions of this section are introduced in the following definition.

DEFINITION 3.5. Consider the following diagram of functors.

$$\begin{array}{ccccc} & & \mathfrak{B} & & \\ & & \uparrow \text{U} & & \\ \mathfrak{C} & \xrightarrow{\text{L}} & \mathfrak{A} & \xleftarrow{\text{R}} & \mathfrak{A} \end{array}$$

They give rise to two functors

$$\mathrm{Hom}_{\mathfrak{A}}(\mathbb{L}(\bullet), \mathbb{R}(\bullet)) \quad \text{and} \quad \mathrm{Hom}_{\mathfrak{B}}(\mathbb{U}\mathbb{L}(\bullet), \mathbb{U}\mathbb{R}(\bullet)) : \mathfrak{L}^{op} \times \mathfrak{R} \rightarrow \underline{\underline{\mathbf{Sets}}}$$

and a natural transformation between them

$$(3.4) \quad \Phi(\mathbb{U}, \mathbb{L}, \mathbb{R}) : \mathrm{Hom}_{\mathfrak{A}}(\mathbb{L}(\bullet), \mathbb{R}(\bullet)) \rightarrow \mathrm{Hom}_{\mathfrak{B}}(\mathbb{U}\mathbb{L}(\bullet), \mathbb{U}\mathbb{R}(\bullet)), \quad \Phi(\mathbb{U}, \mathbb{L}, \mathbb{R})_{L,R}(f) := \mathbb{U}(f),$$

for all objects $L \in \mathfrak{L}, R \in \mathfrak{R}$ and for every morphism $f : \mathbb{L}(L) \rightarrow \mathbb{R}(R)$. We say that

- 1) $\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -*faithful* if $\Phi(\mathbb{U}, \mathbb{L}, \mathbb{R})_{L,R}$ is injective, for every objects $L \in \mathfrak{L}$ and $R \in \mathfrak{R}$.
- 2) $\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -*full* if $\Phi(\mathbb{U}, \mathbb{L}, \mathbb{R})_{L,R}$ is surjective, for every objects $L \in \mathfrak{L}$ and $R \in \mathfrak{R}$.
- 3) $\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -*separable* if $\Phi(\mathbb{U}, \mathbb{L}, \mathbb{R})$ is a split natural monomorphism.
- 4) $\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -*coseparable* if $\Phi(\mathbb{U}, \mathbb{L}, \mathbb{R})$ is a split natural epimorphism.

When both $\mathbb{L}$ and $\mathbb{R}$ are identity functors, we recover the classical definitions of a faithful, full, separable and naturally full (here called coseparable) functor. We are particularly interested in the case when either $\mathbb{L}$ or $\mathbb{R}$ is the identity functor. Anyway, some of our results can be stated for the general case.

REMARK 3.6. Following [Raf, page 1446], one can prove that Definition 3.5 3) can be reformulated as follows. It has the spirit of a characterization of separable functors in [NVV]. A functor $\mathbb{U} : \mathfrak{A} \rightarrow \mathfrak{B}$ is $(\mathbb{L}, \mathbb{R})$ -separable, for some functors $\mathbb{L} : \mathfrak{L} \rightarrow \mathfrak{A}$ and $\mathbb{R} : \mathfrak{R} \rightarrow \mathfrak{A}$, if and only if there is a map

$$\tilde{\Phi}(\mathbb{U}, \mathbb{L}, \mathbb{R})_{L,R} : \mathrm{Hom}_{\mathfrak{B}}(\mathbb{U}\mathbb{L}(L), \mathbb{U}\mathbb{R}(R)) \rightarrow \mathrm{Hom}_{\mathfrak{A}}(\mathbb{L}(L), \mathbb{R}(R)),$$

for all objects $L \in \mathfrak{L}$ and $R \in \mathfrak{R}$, satisfying the following identities.

- S1) $\tilde{\Phi}(\mathbb{U}, \mathbb{L}, \mathbb{R})_{L,R}(\mathbb{U}(f)) = f$, for any $f \in \mathrm{Hom}_{\mathfrak{A}}(\mathbb{L}(L), \mathbb{R}(R))$.
- S2) $\tilde{\Phi}(\mathbb{U}, \mathbb{L}, \mathbb{R})_{L',R'}(h') \circ \mathbb{L}(l) = \mathbb{R}(r) \circ \tilde{\Phi}(\mathbb{U}, \mathbb{L}, \mathbb{R})_{L,R}(h)$, for every commutative diagram in $\mathfrak{B}$ of the following form.

$$\begin{array}{ccc} \mathbb{U}\mathbb{L}(L) & \xrightarrow{h} & \mathbb{U}\mathbb{R}(R) \\ \mathbb{U}\mathbb{L}(l) \downarrow & & \downarrow \mathbb{U}\mathbb{R}(r) \\ \mathbb{U}\mathbb{L}(L') & \xrightarrow{h'} & \mathbb{U}\mathbb{R}(R') \end{array}$$

REMARK 3.7. Recall that faithful functors reflect mono, and epimorphisms. Analogously, for an $(\mathbb{L}, \mathbb{R})$ -faithful functor $\mathbb{U}$ the following hold true.

- 1) Assume that $\mathbb{R}$ is surjective on the objects and let $f : A \rightarrow \mathbb{L}(L)$ be a morphism in $\mathfrak{A}$. Then f is an epimorphism whenever $\mathbb{U}(f)$ is.
- 2) Assume that $\mathbb{L}$ is surjective on the objects and let $f : \mathbb{R}(R) \rightarrow A$ be a morphism in $\mathfrak{A}$. Then f is a monomorphism whenever $\mathbb{U}(f)$ is.

In the rest of the section we extend some standard results about separable functors to relative separable functors in Definition 3.5 3). Analogous results can be obtained for coseparable functors by a careful dualisation.

THEOREM 3.8. *Consider the following diagram of functors.*

$$\begin{array}{ccccccc} & & & \mathfrak{C} & & & \\ & & & \uparrow \mathbb{V} & & & \\ & & & \mathfrak{B} & & & \\ & & & \uparrow \mathbb{U} & & & \\ \mathfrak{L}' & \xrightarrow{\mathbb{L}'} & \mathfrak{L} & \xrightarrow{\mathbb{L}} & \mathfrak{A} & \xleftarrow{\mathbb{R}} & \mathfrak{R} \xleftarrow{\mathbb{R}'} \mathfrak{R}' \end{array}$$

The following assertions hold true.

- 1) If $\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -separable then $\mathbb{U}$ is $(\mathbb{L}\mathbb{L}', \mathbb{R}\mathbb{R}')$ -separable.
- 2) If $\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -separable and $\mathbb{V}$ is $(\mathbb{U}\mathbb{L}, \mathbb{U}\mathbb{R})$ -separable then $\mathbb{V}\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -separable.
- 3) If $\mathbb{V}\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -separable then $\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -separable.

Proof. The proof is similar to [CMZ, I.3 Proposition 46 and Corollary 9].

1) Since $\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -separable, there exists a natural retraction $\tilde{\Phi}(\mathbb{U}, \mathbb{L}, \mathbb{R})$ of the natural transformation (3.4). For any objects $L' \in \mathcal{L}'$ and $R' \in \mathcal{R}'$, the maps

$$\tilde{\Phi}(\mathbb{U}, \mathbb{L}, \mathbb{R})_{\mathbb{L}'(L'), \mathbb{R}'(R')} : \text{Hom}_{\mathfrak{B}}(\mathbb{U}\mathbb{L}\mathbb{L}'(L'), \mathbb{U}\mathbb{R}\mathbb{R}'(R')) \rightarrow \text{Hom}_{\mathfrak{A}}(\mathbb{L}\mathbb{L}'(L'), \mathbb{R}\mathbb{R}'(R'))$$

define a natural transformation which is a retraction of $\Phi(\mathbb{U}, \mathbb{L}\mathbb{L}', \mathbb{R}\mathbb{R}')$, defined analogously to (3.4).

2) The natural transformation

$$(3.5) \quad \Phi(\mathbb{V}\mathbb{U}, \mathbb{L}, \mathbb{R}) : \text{Hom}_{\mathfrak{A}}(\mathbb{L}(\bullet), \mathbb{R}(\bullet)) \rightarrow \text{Hom}_{\mathfrak{C}}(\mathbb{V}\mathbb{U}\mathbb{L}(\bullet), \mathbb{V}\mathbb{U}\mathbb{R}(\bullet)), \quad f \mapsto \mathbb{V}\mathbb{U}(f)$$

is a composite of the natural monomorphisms $\Phi(\mathbb{U}, \mathbb{L}, \mathbb{R})$, corresponding via (3.4) to the $(\mathbb{L}, \mathbb{R})$ -separable functor $\mathbb{U}$, and $\Phi(\mathbb{V}, \mathbb{U}\mathbb{L}, \mathbb{U}\mathbb{R})$, corresponding to the $(\mathbb{U}\mathbb{L}, \mathbb{U}\mathbb{R})$ -separable functor $\mathbb{V}$. Hence (3.5) is a natural monomorphism too, proving $(\mathbb{L}, \mathbb{R})$ -separability of $\mathbb{V}\mathbb{U}$.

3) Since the functor $\mathbb{V}\mathbb{U}$ is $(\mathbb{L}, \mathbb{R})$ -separable, the corresponding natural transformation (3.4) possesses a retraction $\tilde{\Phi}(\mathbb{V}\mathbb{U}, \mathbb{L}, \mathbb{R})$. The composite $\tilde{\Phi}(\mathbb{V}\mathbb{U}, \mathbb{L}, \mathbb{R}) \circ \Phi(\mathbb{V}, \mathbb{U}\mathbb{L}, \mathbb{U}\mathbb{R})$ is a natural retraction of $\Phi(\mathbb{U}, \mathbb{L}, \mathbb{R})$ in (3.4). $\square$

THEOREM 3.9 (Maschke type Theorem). *Let $\mathbb{U} : \mathfrak{A} \rightarrow \mathfrak{B}$, $\mathbb{L} : \mathcal{L} \rightarrow \mathfrak{A}$ and $\mathbb{R} : \mathfrak{R} \rightarrow \mathfrak{A}$ be functors.*

- 1) *If $\mathbb{U}$ is $(\mathfrak{A}, \mathbb{R})$ -separable then, for any objects $R \in \mathfrak{R}$ and $A \in \mathfrak{A}$, a morphism $f : \mathbb{R}(R) \rightarrow A$ is a split monomorphism whenever $\mathbb{U}(f)$ is a split monomorphism. Moreover, in this case $\mathcal{I}_{\mathbb{R}} = \mathcal{I}_{\mathbb{U}\mathbb{R}}$ and $\mathcal{E}_{\mathbb{R}} = \mathcal{E}_{\mathbb{U}\mathbb{R}}$.*
- 2) *If $\mathbb{U}$ is $(\mathbb{L}, \mathfrak{A})$ -separable then, for any objects $L \in \mathcal{L}$ and $A \in \mathfrak{A}$, a morphism $f : A \rightarrow \mathbb{L}(L)$ is a split epimorphism whenever $\mathbb{U}(f)$ is a split epimorphism. Moreover, in this case $\mathcal{I}_{\mathbb{L}} = \mathcal{I}_{\mathbb{U}\mathbb{L}}$ and $\mathcal{E}_{\mathbb{L}} = \mathcal{E}_{\mathbb{U}\mathbb{L}}$.*

Proof. Let A , R and f be as in part 1). Let $\tilde{\Phi}(\mathbb{U}, \mathfrak{A}, \mathbb{R})$ be a natural retraction of $\Phi(\mathbb{U}, \mathfrak{A}, \mathbb{R})$ in (3.4). In view of S2) in Remark 3.6, any retraction π of $\mathbb{U}(f)$ satisfies

$$\tilde{\Phi}(\mathbb{U}, \mathfrak{A}, \mathbb{R})_{A, R}(\pi) \circ f = \mathbb{R}(R).$$

That is, f is a split monomorphism. In particular, $f := \mathbb{R}(g)$ is a split monomorphism, for any $g \in \mathcal{I}_{\mathbb{U}\mathbb{R}}$. Together with (3.3) this proves $\mathcal{I}_{\mathbb{U}\mathbb{R}} = \mathcal{I}_{\mathbb{R}}$. Next take a morphism $g : R \rightarrow R'$ in $\mathcal{E}_{\mathbb{U}\mathbb{R}}$, and a section σ of $\mathbb{U}\mathbb{R}(g)$. Then, by naturality of $\tilde{\Phi}(\mathbb{U}, \mathfrak{A}, \mathbb{R})$,

$$\mathbb{R}(R') = \tilde{\Phi}(\mathbb{U}, \mathfrak{A}, \mathbb{R})_{\mathbb{R}(R'), R'}(\mathbb{U}\mathbb{R}(R')) = \tilde{\Phi}(\mathbb{U}, \mathfrak{A}, \mathbb{R})_{\mathbb{R}(R'), R'}(\mathbb{U}\mathbb{R}(g) \circ \sigma) = \mathbb{R}(g) \circ \tilde{\Phi}(\mathbb{U}, \mathfrak{A}, \mathbb{R})_{\mathbb{R}(R'), R}(\sigma).$$

This implies that $\mathbb{R}(g)$ is a split epimorphism, i.e. $g \in \mathcal{E}_{\mathbb{R}}$. In view of (3.3), we have $\mathcal{E}_{\mathbb{R}} = \mathcal{E}_{\mathbb{U}\mathbb{R}}$ proven.

Part 2) is proven by dual reasoning. $\square$

COROLLARY 3.10. *Let $(\mathbb{T}, \mathbb{H})$ be an adjunction of functors $\mathbb{T} : \mathfrak{A} \rightarrow \mathfrak{B}$ and $\mathbb{H} : \mathfrak{B} \rightarrow \mathfrak{A}$. For any functors $\mathbb{L} : \mathcal{L} \rightarrow \mathfrak{B}$ and $\mathbb{R} : \mathfrak{R} \rightarrow \mathfrak{A}$, the following hold.*

- 1) *If the functor $\mathbb{H}$ is $(\mathbb{L}, \mathfrak{B})$ -separable then $\mathbb{L}(L)$ is $\mathcal{E}_{\mathbb{H}}$ -projective for every $L \in \mathcal{L}$.*
- 2) *If the functor $\mathbb{T}$ is $(\mathfrak{A}, \mathbb{R})$ -separable then $\mathbb{R}(R)$ is $\mathcal{I}_{\mathbb{T}}$ -injective for every $R \in \mathfrak{R}$.*

Proof. Let $\eta : \mathfrak{A} \rightarrow \mathbb{H}\mathbb{T}$ be the unit and $\varepsilon : \mathbb{T}\mathbb{H} \rightarrow \mathfrak{B}$ be the counit of the adjunction $(\mathbb{T}, \mathbb{H})$.

1) For any object $L \in \mathcal{L}$, the epimorphism $\mathbb{H}(\varepsilon_{\mathbb{L}(L)})$ is split by $\eta_{\mathbb{H}\mathbb{L}(L)}$. Hence, by Theorem 3.9 2), $\varepsilon_{\mathbb{L}(L)}$ is a split epimorphism in $\mathfrak{B}$. By Theorem 3.2 (b) $\Rightarrow$ (a), $\mathbb{L}(L)$ is $\mathcal{E}_{\mathbb{H}}$ -projective.

2) For any object $R \in \mathfrak{R}$, the monomorphism $\mathbb{T}(\eta_{\mathbb{R}(R)})$ is split by $\varepsilon_{\mathbb{T}\mathbb{R}(R)}$. Hence the claim follows analogously to part 1), by Theorem 3.9 1) and Theorem 3.3. $\square$

In the following theorem functors preserving and reflecting relative projective (resp. injective) objects are studied.

THEOREM 3.11. *Let $(\mathbb{T}, \mathbb{H})$ and $(\mathbb{T}', \mathbb{H}')$ be adjunctions and consider the following (not necessarily commutative) diagrams of functors.*

$$\begin{array}{ccc} \mathfrak{A} & \xrightarrow{\mathbb{F}'} & \mathfrak{A}' \\ \mathbb{T} \downarrow & & \downarrow \mathbb{T}' \\ \mathfrak{B} & \xrightarrow{\mathbb{F}} & \mathfrak{B}' \end{array} \quad \begin{array}{ccc} \mathfrak{A} & \xrightarrow{\mathbb{F}'} & \mathfrak{A}' \\ \mathbb{H} \uparrow & & \uparrow \mathbb{H}' \\ \mathfrak{B} & \xrightarrow{\mathbb{F}} & \mathfrak{B}' \end{array}$$

If $\mathbb{T}'\mathbb{F}'$ and $\mathbb{F}\mathbb{T}$ are naturally equivalent, then the following hold.

- 1) If an object P in $\mathfrak{B}$ is $\mathcal{E}_{\mathbb{H}}$ -projective then $\mathbb{F}(P)$ is $\mathcal{E}_{\mathbb{H}'}$ -projective.
- 2^{op}) Assume that $\mathbb{F}'$ is $(\mathfrak{A}, \mathbb{R})$ -separable for some functor $\mathbb{R} : \mathfrak{R} \rightarrow \mathfrak{A}$. If, for an object $R \in \mathfrak{R}$, the object $\mathbb{F}'\mathbb{R}(R)$ is $\mathcal{I}_{\mathbb{T}'}$ -injective, then $\mathbb{R}(R)$ is $\mathcal{I}_{\mathbb{T}}$ -injective.

If $\mathbb{F}'\mathbb{H}$ and $\mathbb{H}'\mathbb{F}$ are naturally equivalent, then the following hold.

- 1^{op}) If an object I in $\mathfrak{A}$ is $\mathcal{I}_{\mathbb{T}}$ -injective, then $\mathbb{F}'(I)$ is $\mathcal{I}_{\mathbb{T}'}$ -injective.
- 2) Assume that $\mathbb{F}$ is $(\mathbb{L}, \mathfrak{B})$ -separable for some functor $\mathbb{L} : \mathfrak{L} \rightarrow \mathfrak{B}$. If, for an object $L \in \mathfrak{L}$, the object $\mathbb{F}\mathbb{L}(L)$ is $\mathcal{E}_{\mathbb{H}'}$ -projective then $\mathbb{L}(L)$ is $\mathcal{E}_{\mathbb{H}}$ -projective.

Proof. Denote by $\eta : \mathfrak{A} \rightarrow \mathbb{H}\mathbb{T}$ the unit and by $\varepsilon : \mathbb{T}\mathbb{H} \rightarrow \mathfrak{B}$ the counit of the adjunction $(\mathbb{T}, \mathbb{H})$.

1) By Theorem 3.2 (a) $\Rightarrow$ (b), $\mathcal{E}_{\mathbb{H}}$ -projectivity of P implies that $\varepsilon_P : \mathbb{T}\mathbb{H}(P) \rightarrow P$ is a split epimorphism. Hence also $\mathbb{F}(\varepsilon_P) : \mathbb{T}'\mathbb{F}'\mathbb{H}(P) \sim \mathbb{F}\mathbb{T}\mathbb{H}(P) \rightarrow \mathbb{F}(P)$ is a split epimorphism. Application of Theorem 3.2 (c) $\Rightarrow$ (a) to the adjunction $(\mathbb{T}', \mathbb{H}')$ completes the proof of $\mathcal{E}_{\mathbb{H}'}$ -projectivity of $\mathbb{F}(P)$.

2) For any object P in $\mathfrak{B}$, the monomorphism $\eta_{\mathbb{H}(P)}$ is split by $\mathbb{H}(\varepsilon_P)$. Hence $\mathbb{F}'\mathbb{H}(\varepsilon_P)$ is a split epimorphism. By the natural equivalence $\mathbb{F}'\mathbb{H} \sim \mathbb{H}'\mathbb{F}$, also $\mathbb{H}'\mathbb{F}(\varepsilon_P)$ is a split epimorphism, yielding that $\mathbb{F}(\varepsilon_P) : \mathbb{F}\mathbb{T}\mathbb{H}(P) \rightarrow \mathbb{F}(P)$ belongs to $\mathcal{E}_{\mathbb{H}'}$. In the case when $\mathbb{F}(P)$ is $\mathcal{E}_{\mathbb{H}'}$ -projective, we conclude that $\mathbb{F}(\varepsilon_P)$ is a split epimorphism in $\mathfrak{B}'$. Now put $P = \mathbb{L}(L)$, such that $\mathbb{F}\mathbb{L}(L)$ is $\mathcal{E}_{\mathbb{H}'}$ -projective as in the claim. Then, by Theorem 3.9 2), $\varepsilon_{\mathbb{L}(L)}$ is a split epimorphism in $\mathfrak{B}$ and hence $\mathbb{L}(L)$ is $\mathcal{E}_{\mathbb{H}}$ -projective by Theorem 3.2 (b) $\Rightarrow$ (a).

The remaining claims 1^{op}) and 2^{op}) follow by dual reasoning. $\square$

Let $(\mathbb{T}, \mathbb{H})$ be an adjunction of functors $\mathbb{T} : \mathfrak{A} \rightarrow \mathfrak{B}$ and $\mathbb{H} : \mathfrak{B} \rightarrow \mathfrak{A}$. Denote by $\varepsilon : \mathbb{T}\mathbb{H} \rightarrow \mathfrak{B}$ and $\eta : \mathfrak{A} \rightarrow \mathbb{H}\mathbb{T}$ the counit and the unit of the adjunction, respectively. Consider the canonical isomorphism

$$(3.6) \quad \phi_{A,B} : \text{Hom}_{\mathfrak{B}}(\mathbb{T}(A), B) \rightarrow \text{Hom}_{\mathfrak{A}}(A, \mathbb{H}(B)), \quad \phi_{A,B}(f) = \mathbb{H}(f) \circ \eta_A$$

with inverse

$$\phi_{A,B}^{-1} : \text{Hom}_{\mathfrak{A}}(A, \mathbb{H}(B)) \rightarrow \text{Hom}_{\mathfrak{B}}(\mathbb{T}(A), B), \quad \phi_{A,B}^{-1}(g) = \varepsilon_B \circ \mathbb{T}(g).$$

In terms of the natural transformations (3.4) and (3.6), for any functors $\mathbb{L} : \mathfrak{L} \rightarrow \mathfrak{A}$ and $\mathbb{R} : \mathfrak{R} \rightarrow \mathfrak{A}$, define a natural transformation

$$\Omega := \phi \circ \Phi(\mathbb{T}, \mathbb{L}, \mathbb{R}) : \text{Hom}_{\mathfrak{A}}(\mathbb{L}(\bullet), \mathbb{R}(\bullet)) \rightarrow \text{Hom}_{\mathfrak{A}}(\mathbb{L}(\bullet), \mathbb{H}\mathbb{T}\mathbb{R}(\bullet)).$$

Then, for every morphism $f : \mathbb{L}(L) \rightarrow \mathbb{R}(R)$, one has

$$(3.7) \quad \Omega_{L,R}(f) = \mathbb{H}\mathbb{T}(f) \circ \eta_{\mathbb{L}(L)} = \eta_{\mathbb{R}(R)} \circ f.$$

Dually, for functors $\mathbb{L} : \mathfrak{L} \rightarrow \mathfrak{B}$ and $\mathbb{R} : \mathfrak{R} \rightarrow \mathfrak{B}$, there is a natural transformation

$$\mathcal{U} := \phi^{-1} \circ \Phi(\mathbb{H}, \mathbb{L}, \mathbb{R}) : \text{Hom}_{\mathfrak{A}}(\mathbb{L}(\bullet), \mathbb{R}(\bullet)) \rightarrow \text{Hom}_{\mathfrak{A}}(\mathbb{T}\mathbb{H}\mathbb{L}(\bullet), \mathbb{R}(\bullet)),$$

mapping a morphism $f : \mathbb{L}(L) \rightarrow \mathbb{R}(R)$ to

$$\mathcal{U}_{L,R}(f) = \varepsilon_{\mathbb{R}(R)} \circ \mathbb{T}\mathbb{H}(f) = f \circ \varepsilon_{\mathbb{L}(L)}.$$

LEMMA 3.12. *On the category of functors and natural transformations consider the following end-ofunctor α . It maps a functor $\mathbb{F} : \mathfrak{A} \rightarrow \mathfrak{B}$ to the functor $\text{Hom}_{\mathfrak{B}}(\bullet, \mathbb{F}(\bullet)) : \mathfrak{B}^{op} \times \mathfrak{A} \rightarrow \underline{\mathbf{Sets}}$, and it maps a natural transformation $\sigma \in \underline{\mathbf{Nat}}(\mathbb{F}, \mathbb{G})$ to $\text{Hom}_{\mathfrak{B}}(\bullet, \sigma_{\bullet})$, i.e.*

$$\alpha(\sigma)_{B,A} : \text{Hom}_{\mathfrak{B}}(B, \mathbb{F}(A)) \rightarrow \text{Hom}_{\mathfrak{B}}(B, \mathbb{G}(A)), \quad g \mapsto \sigma_A \circ g.$$

The functor α is fully faithful.

Proof. The bijectivity of the maps

$$\alpha_{\mathbb{F}, \mathbb{G}} : \underline{\text{Nat}}(\mathbb{F}, \mathbb{G}) \rightarrow \underline{\text{Nat}}(\text{Hom}_{\mathfrak{B}}(\bullet, \mathbb{F}(\bullet)), \text{Hom}_{\mathfrak{B}}(\bullet, \mathbb{G}(\bullet))) \quad \sigma \mapsto \alpha(\sigma),$$

for any functors $\mathbb{F}, \mathbb{G} : \mathfrak{A} \rightarrow \mathfrak{B}$, is proven by constructing the inverse $(\alpha_{\mathbb{F}, \mathbb{G}})^{-1}(\mathcal{P})_A := \mathcal{P}_{\mathbb{F}(A), A}(\mathbb{F}(A))$, for $\mathcal{P} \in \underline{\text{Nat}}(\text{Hom}_{\mathfrak{B}}(\bullet, \mathbb{F}(\bullet)), \text{Hom}_{\mathfrak{B}}(\bullet, \mathbb{G}(\bullet)))$ and $A \in \mathfrak{A}$. It is straightforward to check that the naturality of $\mathcal{P}$ (i.e. the identity $\mathbb{G}(a) \circ \mathcal{P}_{B, A}(g) \circ b = \mathcal{P}_{B', A'}(\mathbb{F}(a) \circ g \circ b)$, for $a \in \text{Hom}_{\mathfrak{A}}(A, A')$, $b \in \text{Hom}_{\mathfrak{B}}(B', B)$ and $g \in \text{Hom}_{\mathfrak{B}}(B, \mathbb{F}(A))$) implies the naturality of $(\alpha_{\mathbb{F}, \mathbb{G}})^{-1}(\mathcal{P})$. Furthermore, (keeping the notation),

$$\alpha_{\mathbb{F}, \mathbb{G}}((\alpha_{\mathbb{F}, \mathbb{G}})^{-1}(\mathcal{P}))_{B, A}(g) = (\alpha_{\mathbb{F}, \mathbb{G}})^{-1}(\mathcal{P})_A \circ g = \mathcal{P}_{\mathbb{F}(A), A}(\mathbb{F}(A)) \circ g = \mathcal{P}_{B, A}(g),$$

where the last equality follows by the naturality of $\mathcal{P}$. Also,

$$(\alpha_{\mathbb{F}, \mathbb{G}})^{-1}(\alpha_{\mathbb{F}, \mathbb{G}}(\sigma))_A = \alpha_{\mathbb{F}, \mathbb{G}}(\sigma)_{\mathbb{F}(A), A}(\mathbb{F}(A)) = \sigma_A,$$

what completes the proof. $\square$

THEOREM 3.13. *Let $(\mathbb{T}, \mathbb{H})$ be an adjunction of functors $\mathbb{T} : \mathfrak{A} \rightarrow \mathfrak{B}$ and $\mathbb{H} : \mathfrak{B} \rightarrow \mathfrak{A}$, with unit η and counit ε . Consider any functor $\mathbb{R} : \mathfrak{A} \rightarrow \mathfrak{A}$ and a functor $\mathbb{L} : \mathfrak{L} \rightarrow \mathfrak{A}$ which is surjective on the objects (e.g. the identity functor $\mathbb{L} = \mathfrak{A}$). Then the following assertions hold.*

- 1) $\mathbb{T}$ is $(\mathbb{L}, \mathbb{R})$ -faithful if and only if it is $(\mathfrak{A}, \mathbb{R})$ -faithful and if and only if $\eta_{\mathbb{R}(R)}$ is a monomorphism, for every object $R \in \mathfrak{A}$.
- 2) $\mathbb{T}$ is $(\mathbb{L}, \mathbb{R})$ -full if and only if it is $(\mathfrak{A}, \mathbb{R})$ -full and if and only if $\eta_{\mathbb{R}(R)}$ is a split epimorphism, for every object $R \in \mathfrak{A}$.
- 3) (Rafael type Theorem) $\mathbb{T}$ is $(\mathfrak{A}, \mathbb{R})$ -separable if and only if $\eta_{\mathbb{R}(\bullet)}$ is a split natural monomorphism.
- 4) (Dual Rafael type Theorem) $\mathbb{T}$ is $(\mathfrak{A}, \mathbb{R})$ -coseparable if and only if $\eta_{\mathbb{R}(\bullet)}$ is a split natural epimorphism.

Proof. Recall that the natural transformation ϕ in (3.6) is an isomorphism.

1) $(\mathbb{L}, \mathbb{R})$ -faithfulness of $\mathbb{T}$, i.e. injectivity of the natural transformation $\Phi(\mathbb{T}, \mathbb{L}, \mathbb{R})_{L, R}$ in (3.4), for every object $L \in \mathfrak{L}$ and $R \in \mathfrak{A}$, is equivalent to injectivity of $\Omega_{L, R}$, for every $L \in \mathfrak{L}$ and $R \in \mathfrak{A}$. Since $\mathbb{L}$ is surjective on the objects, in light of (3.7) this is equivalent to saying that $\eta_{\mathbb{R}(R)}$ is a monomorphism for every $R \in \mathfrak{A}$.

2) $(\mathbb{L}, \mathbb{R})$ -fullness of $\mathbb{T}$, i.e. surjectivity of the natural transformation $\Phi(\mathbb{T}, \mathbb{L}, \mathbb{R})_{L, R}$ in (3.4), for every object $L \in \mathfrak{L}$ and $R \in \mathfrak{A}$, is equivalent to surjectivity of $\Omega_{L, R}$, for every $L \in \mathfrak{L}$ and $R \in \mathfrak{A}$. Let us prove that this is equivalent to saying that $\eta_{\mathbb{R}(R)}$ is a split epimorphism for every $R \in \mathfrak{A}$. In fact, since $\mathbb{L}$ is surjective on the objects, for every $R \in \mathfrak{A}$ there exists an object $L \in \mathfrak{L}$ such that $\mathbb{H}\mathbb{T}\mathbb{R}(R) = \mathbb{L}(L)$. Thus if $\Omega_{L, R}$ is surjective then, by $\mathbb{H}\mathbb{T}\mathbb{R}(R) \in \text{Hom}_{\mathfrak{A}}(\mathbb{H}\mathbb{T}\mathbb{R}(R), \mathbb{H}\mathbb{T}\mathbb{R}(R)) = \text{Hom}_{\mathfrak{A}}(\mathbb{L}(L), \mathbb{H}\mathbb{T}\mathbb{R}(R))$, there exists $\sigma \in \text{Hom}_{\mathfrak{A}}(\mathbb{L}(L), \mathbb{R}(R))$ such that $\eta_{\mathbb{R}(R)} \circ \sigma = \mathbb{H}\mathbb{T}\mathbb{R}(R)$. Conversely, let g be any morphism in $\text{Hom}_{\mathfrak{A}}(\mathbb{L}(L), \mathbb{H}\mathbb{T}\mathbb{R}(R))$, for some $L \in \mathfrak{L}$ and $R \in \mathfrak{A}$. Let σ be a section of $\eta_{\mathbb{R}(R)}$. Define $f \in \text{Hom}_{\mathfrak{A}}(\mathbb{L}(L), \mathbb{R}(R))$ by $f := \sigma \circ g$. Then $\Omega_{L, R}(f) = \eta_{\mathbb{R}(R)} \circ f = g$.

3) $(\mathfrak{A}, \mathbb{R})$ -separability of $\mathbb{T}$, i.e. natural cosplitting of $\Phi(\mathbb{T}, \mathfrak{A}, \mathbb{R})$, is equivalent to natural cosplitting of Ω . Note that Ω is the image of the natural transformation $\eta_{\mathbb{R}(\bullet)}$ under the functor α in Lemma 3.12. Hence the claim follows by Lemma 3.12, as a fully faithful functor preserves and reflects split monomorphisms.

4) $(\mathfrak{A}, \mathbb{R})$ -coseparability of $\mathbb{T}$, i.e. natural splitting of $\Phi(\mathbb{T}, \mathfrak{A}, \mathbb{R})$, is equivalent to natural splitting of Ω . Hence this claim follows by the same argument as 3) does, as a fully faithful functor preserves and reflects split epimorphisms as well. $\square$

Dually, one proves the following result.

THEOREM 3.14. *Let $(\mathbb{T}, \mathbb{H})$ be an adjunction of functors $\mathbb{H} : \mathfrak{A} \rightarrow \mathfrak{B}$ and $\mathbb{T} : \mathfrak{B} \rightarrow \mathfrak{A}$, with unit η and counit ε . Consider any functor $\mathbb{L} : \mathfrak{L} \rightarrow \mathfrak{A}$ and a functor $\mathbb{R} : \mathfrak{A} \rightarrow \mathfrak{A}$ which is surjective on the objects (e.g. the identity functor $\mathbb{R} = \mathfrak{A}$). Then the following assertions hold.*

- 1) $\mathbb{H}$ is $(\mathbb{L}, \mathbb{R})$ -faithful if and only if it is $(\mathbb{L}, \mathfrak{A})$ -faithful and if and only if $\varepsilon_{\mathbb{L}(L)}$ is an epimorphism for every object $L \in \mathfrak{L}$.

- 2) $\mathbb{H}$ is $(\mathbb{L}, \mathbb{R})$ -full if and only if it is $(\mathbb{L}, \mathfrak{A})$ -full and if and only if $\varepsilon_{\mathbb{L}(L)}$ is a split monomorphism for every object $L \in \mathfrak{L}$.
- 3) (Rafael type Theorem) $\mathbb{H}$ is $(\mathbb{L}, \mathfrak{A})$ -separable if and only if $\varepsilon_{\mathbb{L}(\bullet)}$ is a split natural epimorphism.
- 4) (Dual Rafael type Theorem) $\mathbb{H}$ is $(\mathbb{L}, \mathfrak{A})$ -coseparable if and only if $\varepsilon_{\mathbb{L}(\bullet)}$ is a split natural monomorphism.

A notion somewhat reminiscent to our *relative separability* of a functor was introduced in [CM] under the name of *separability of the second kind*. Our next task is to find a relation between the two notions.

DEFINITION 3.15. Let $\mathbb{R} : \mathfrak{A} \rightarrow \mathfrak{A}'$ and $\mathbb{T} : \mathfrak{A} \rightarrow \mathfrak{B}$ be covariant functors. Following [CM, Definition 2.1] and using the notation introduced in (3.4), $\mathbb{T}$ is called *$\mathbb{R}$ -separable of the second kind* if the natural transformation $\Phi(\mathbb{R}, \mathfrak{A}, \mathfrak{A})$ factors through $\Phi(\mathbb{T}, \mathfrak{A}, \mathfrak{A})$.

PROPOSITION 3.16. Let $(\mathbb{T}, \mathbb{H})$ and $(\mathbb{T}', \mathbb{H}')$ be adjunctions with respective units η and η' . Consider the following diagrams of functors.

$$\begin{array}{ccc} \mathfrak{A} & \xrightarrow{\mathbb{R}} & \mathfrak{A}' \\ \mathbb{T} \downarrow & & \downarrow \mathbb{T}' \\ \mathfrak{B} & & \mathfrak{B}' \end{array} \quad \begin{array}{ccc} \mathfrak{A} & \xrightarrow{\mathbb{R}} & \mathfrak{A}' \\ \mathbb{H} \uparrow & & \uparrow \mathbb{H}' \\ \mathfrak{B} & & \mathfrak{B}' \end{array}$$

The following assertions are equivalent.

- (a) $\mathbb{T}$ is $\mathbb{R}$ -separable of the second kind.
- (b) There exists a natural transformation $\nu : \mathbb{R}\mathbb{H}\mathbb{T} \rightarrow \mathbb{R}$, satisfying $\nu_A \circ \mathbb{R}(\eta_A) = \mathbb{R}(A)$, for any $A \in \mathfrak{A}$.

Assume that there exists a natural equivalence $\xi : \mathbb{H}'\mathbb{T}'\mathbb{R} \rightarrow \mathbb{R}\mathbb{H}\mathbb{T}$ such that

$$(3.8) \quad \xi \circ \eta'_{\mathbb{R}(\bullet)} = \mathbb{R}(\eta_{\bullet}).$$

Then the following assertion is also equivalent to the foregoing ones.

- (c) $\mathbb{T}'$ is $(\mathfrak{A}', \mathbb{R})$ -separable.

Proof. (a) $\Leftrightarrow$ (b) This equivalence was proven in [CM, Theorem 2.7].

(b) $\Leftrightarrow$ (c) This equivalence follows by Theorem 3.13 3), in view of (3.8). $\square$

4. APPLICATION TO ENTWINING STRUCTURES

In this section we study coring extensions, especially those ones which arise from entwining structures. We focus on the problem of $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separability of the functor $\mathbb{U}^{\mathcal{D}}$ in Figure (2.4).

The following first result is an easy generalisation of [Brz1, Corollary 3.6] to coring extensions.

PROPOSITION 4.1. Consider an L -coring $\mathcal{D}$ which is right extension of an A -coring $\mathcal{C}$, and the corresponding functors in Figure (2.4). The forgetful functor $\mathbb{U}^{\mathcal{D}}$ is $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separable if and only if the right $\mathcal{D}$ -coaction $\tau_{\mathcal{C}}$ in $\mathcal{C}$ is a split monomorphism of left $\mathcal{C}$ -comodules and right $\mathcal{D}$ -comodules.

Proof. The functor $\mathbb{U}^{\mathcal{D}}$ possesses a right adjoint, the functor $\bullet \otimes_L \mathcal{D} : \mathfrak{M}_L \rightarrow \mathfrak{M}^{\mathcal{D}}$ (cf. [BW, 18.13]). The unit of the adjunction is given by the $\mathcal{D}$ -coaction τ . Hence, by Theorem 3.13 3), $\mathbb{U}^{\mathcal{D}}$ is $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separable if and only if there exists a natural retraction ν of $\tau_{\mathbb{R}(\bullet)}$. Therefore if $\mathbb{U}^{\mathcal{D}}$ is $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separable then in particular $\tau_{\mathcal{C}}$ possesses a right $\mathcal{D}$ -colinear retraction $\nu_{\mathcal{C}}$. We claim that $\nu_{\mathcal{C}}$ is also left $\mathcal{C}$ -colinear. Indeed, for any right A -module N and $n \in N$, the map $\mathcal{C} \rightarrow N \otimes_A \mathcal{C}$, $c \mapsto n \otimes_A c$ is right $\mathcal{C}$ -colinear. Hence by the naturality of ν ,

$$\nu_{N \otimes_A \mathcal{C}}(n \otimes_A c \otimes_L d) = n \otimes_A \nu_{\mathcal{C}}(c \otimes_L d),$$

for $n \in N$, $c \in \mathcal{C}$ and $d \in \mathcal{D}$. In particular, taking $N = A$, we conclude on the left A -linearity of $\nu_{\mathcal{C}}$. Furthermore, a right $\mathcal{C}$ -coaction $\rho^M : m \mapsto m^{[0]} \otimes_A m^{[1]}$ (being coassociative) is $\mathcal{C}$ -colinear, hence the naturality of ν implies

$$\rho^M(\nu_M(m \otimes_L d)) = \nu_{M \otimes_A \mathcal{C}}(\rho^M(m) \otimes_L d),$$

for any $\mathcal{C}$ -comodule M , $m \in M$ and $d \in \mathcal{D}$. Therefore

$$\nu_M(m \otimes_L d)^{[0]} \otimes_A \nu_M(m \otimes_L d)^{[1]} = m^{[0]} \otimes_A \nu_{\mathcal{C}}(m^{[1]} \otimes_L d).$$

Taking $M = \mathcal{C}$ we have the left $\mathcal{C}$ -colinearity of $\nu_{\mathcal{C}}$ proven.

Conversely, let $\tilde{\nu}$ be a left $\mathcal{C}$ -colinear right $\mathcal{D}$ -colinear retraction of $\tau_{\mathcal{C}}$. The natural transformation ν is constructed as follows. For any right $\mathcal{C}$ -comodule M , put

$$(4.1) \quad \nu_M : M \otimes_L \mathcal{D} \rightarrow M, \quad m \otimes_L d \mapsto m^{[0]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m^{[1]} \otimes_L d).$$

Its naturality is obvious. It follows by the $\mathcal{D}$ -colinearity of a $\mathcal{C}$ -coaction ρ^M that ν_M is a retraction of $\tau_M : m \mapsto m_{[0]} \otimes_L m_{[1]}$. Indeed,

$$\nu_M \circ \tau_M(m) = m_{[0]}^{[0]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m_{[0]}^{[1]} \otimes_L m_{[1]}) = m^{[0]} \epsilon_{\mathcal{C}} \circ \tilde{\nu} \circ \tau_{\mathcal{C}}(m^{[1]}) = m.$$

It remains to check the $\mathcal{D}$ -colinearity of ν_M . For $m \otimes_L d \in M \otimes_L \mathcal{D}$,

$$\begin{aligned} \tau_M \circ \nu_M(m \otimes_L d) &= (m^{[0]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m^{[1]} \otimes_L d))_{[0]} \otimes_L (m^{[0]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m^{[1]} \otimes_L d))_{[1]} \\ &= (m^{[0]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m^{[1]} \otimes_L d))^{[0]} \epsilon_{\mathcal{C}} ((m^{[0]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m^{[1]} \otimes_L d))^{[1]})_{[0]} \otimes_L (m^{[0]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m^{[1]} \otimes_L d))^{[1]}_{[1]} \\ &= m^{[0]} \epsilon_{\mathcal{C}} ((m^{[1]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m^{[2]} \otimes_L d))_{[0]}) \otimes_L (m^{[1]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m^{[2]} \otimes_L d))_{[1]} \\ &= m^{[0]} \epsilon_{\mathcal{C}} (\tilde{\nu}(m^{[1]} \otimes_L d)_{[0]}) \otimes_L \tilde{\nu}(m^{[1]} \otimes_L d)_{[1]} \\ &= m^{[0]} \epsilon_{\mathcal{C}} \circ \tilde{\nu}(m^{[1]} \otimes_L d_{(1)}) \otimes_L d_{(2)} = (\nu_M \otimes \mathcal{D}) \circ (M \otimes \Delta_{\mathcal{D}})(m \otimes d), \end{aligned}$$

where the second equality follows by the explicit form (2.6) of the functor $\mathbb{R}$, relating τ_M to ϱ^M , the third one follows by the right A -linearity of a $\mathcal{C}$ -coaction, and the fourth and fifth equalities follow by the left $\mathcal{C}$ -colinearity and the right $\mathcal{D}$ -colinearity of $\tilde{\nu}$, respectively. $\square$

If the two corings $\mathcal{C}$ and $\mathcal{D}$ are equal and $\mathbb{R}$ is the identity functor, then Proposition 4.1 reduces to [Brz1, Corollary 3.6]. On the other hand, for an arbitrary coring extension $\mathcal{D}$ of $\mathcal{C}$, [Brz1, Corollary 3.6] together with Theorem 3.8 1) implies that if $\mathcal{D}$ is a coseparable coring then the functor $\mathbb{U}^{\mathcal{D}}$ in Figure (2.4) is $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separable. This fact follows alternatively by Proposition 4.1: if ζ is a $\mathcal{D}$ - $\mathcal{D}$ bilinear retraction of $\Delta_{\mathcal{D}}$, then $(\mathcal{C} \otimes_L \epsilon_{\mathcal{D}} \circ \zeta) \circ (\tau_{\mathcal{C}} \otimes_L \mathcal{D})$ is a $\mathcal{C}$ - $\mathcal{D}$ bilinear retraction of $\tau_{\mathcal{C}}$.

Note that, by Corollary 3.10 2), for any coring extension $\mathcal{D}$ of $\mathcal{C}$, $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separability of $\mathbb{U}^{\mathcal{D}}$ implies in particular that every right $\mathcal{C}$ -comodule is relative injective as a right $\mathcal{D}$ -comodule. In what follows we turn to analysing more consequences of $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separability of $\mathbb{U}^{\mathcal{D}}$, for coring extensions arising from entwining structures $(A, \mathcal{D}, \psi)$ over an algebra L . As a main result of the section, Theorem 4.3 and Proposition 4.4 show that if ψ is bijective and there exists a grouplike element in $\mathcal{D}$, then $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separability of $\mathbb{U}^{\mathcal{D}}$ implies that A is relative injective *also as an entwined module*. A key notion of our study is a following generalization of Doi's total integral in [Doi].

DEFINITION 4.2. Let $(A, \mathcal{D}, \psi)$ be an entwining structure over an algebra L . Assume that $\mathcal{D}$ possesses a grouplike element e so that A is a right $\mathcal{D}$ -comodule with coaction (2.7). A right $\mathcal{D}$ -comodule map $j : \mathcal{D} \rightarrow A$, satisfying the normalisation condition $j(e) = 1_A$, is called a *right total integral*.

In a bijective entwining structure $(A, \mathcal{D}, \psi)$ over an algebra L , such that $\mathcal{D}$ possesses a grouplike element e , a *left total integral* is defined as a right total integral in the L^{op} -entwining structure $(A^{op}, \mathcal{D}_{cop}, \psi^{-1})$. This is the same as a left $\mathcal{D}$ -comodule map $j : \mathcal{D} \rightarrow A$, with respect to the coaction (2.8), satisfying the normalisation condition $j(e) = 1_A$.

Consider an entwining structure $(A, \mathcal{D}, \psi)$ over an algebra L , and denote by $\mathcal{C}$ the associated A -coring $A \otimes_L \mathcal{D}$. Consider the following diagrams of functors

$$(4.2) \quad \begin{array}{ccc} \mathfrak{M}^{\mathcal{C}} \cong \mathfrak{M}_A^{\mathcal{D}}(\psi) & \xrightarrow{\mathbb{R}} & \mathfrak{M}^{\mathcal{D}} \\ \mathbb{T} = \mathbb{U}^{\mathcal{C}} \downarrow & & \downarrow \mathbb{T}' = \mathbb{U}^{\mathcal{D}} \\ \mathfrak{M}_A & & \mathfrak{M}_L \end{array} \quad \begin{array}{ccc} \mathfrak{M}^{\mathcal{C}} \cong \mathfrak{M}_A^{\mathcal{D}}(\psi) & \xrightarrow{\mathbb{R}} & \mathfrak{M}^{\mathcal{D}} \\ \uparrow \mathbb{H} = \bullet \otimes_L \mathcal{D} & & \mathbb{H}' = \bullet \otimes_L \mathcal{D} \uparrow \\ \mathfrak{M}_A & & \mathfrak{M}_L \end{array}$$

where $\mathbb{T} = \mathbb{U}^{\mathcal{C}}, \mathbb{T}' = \mathbb{U}^{\mathcal{D}}$ and $\mathbb{R}$ are forgetful functors (cf. Figure (2.4)). Note that $(\mathbb{T}, \mathbb{H})$ and $(\mathbb{T}', \mathbb{H}')$ are adjunctions and the respective units η and η' are given by the right $\mathcal{D}$ -coaction, in both cases (cf. Section 2.3). Hence they satisfy $\mathbb{R}(\eta_{\bullet}) = \eta'_{\mathbb{R}(\bullet)}$.

THEOREM 4.3. *Let $(A, \mathcal{D}, \psi)$ be an entwining structure over an algebra L . Consider the functors in Figure (4.2). The following assertions are equivalent.*

- (a) $\mathbb{T} = \mathbb{U}^{\mathcal{C}}$ is $\mathbb{R}$ -separable of the second kind.
- (b) There exists a natural transformation $\nu : \mathbb{R}\mathbb{H}\mathbb{T} \rightarrow \mathbb{R}$, satisfying $\nu_M \circ \mathbb{R}(\eta_M) = \mathbb{R}(M)$, for any $M \in \mathfrak{M}_A^{\mathcal{D}}(\psi)$.
- (c) $\mathbb{T}' = \mathbb{U}^{\mathcal{D}}$ is $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separable.
- (d) There exists a morphism $\theta \in {}_L\text{Hom}_L(\mathcal{D} \otimes_L \mathcal{D}, A)$ satisfying, for all $d, d' \in \mathcal{D}$,

$$(4.3) \quad \theta(d \otimes_L d'_{(1)}) \otimes_L d'_{(2)} = \psi(d_{(1)} \otimes_L \theta(d_{(2)} \otimes_L d')) \quad \text{and} \quad \theta(d_{(1)} \otimes_L d_{(2)}) = \eta_A \circ \epsilon_{\mathcal{D}}(d).$$

If these equivalent conditions hold, and in addition there exists a grouplike element in $\mathcal{D}$, then there exists a right total integral in the L -entwining structure $(A, \mathcal{D}, \psi)$.

Proof. The equivalence of assertions (a), (b) and (c) is a consequence of Proposition 3.16.

The equivalence of assertions (a) and (d) is proven by an easy extension of arguments in [CM, Proposition 4.12], about entwining structures over commutative rings, to non-commutative base.

Assume that there exists a grouplike element e in $\mathcal{D}$, hence A is a right $\mathcal{D}$ -comodule with coaction (2.7). In this situation the map

$$j : \mathcal{D} \rightarrow A \quad d \mapsto \theta(e \otimes_L d)$$

is right $\mathcal{D}$ -colinear and satisfies the normalisation condition $j(e) = 1_A$. That is, j is a right total integral in the sense of Definition 4.2. $\square$

Note that, following the proof of [CM, Proposition 4.12], a bijective correspondence can be obtained between maps θ as in (4.3) and left $\mathcal{C} = (A \otimes_L \mathcal{D})$ -colinear right $\mathcal{D}$ -colinear retractions of the $\mathcal{D}$ -coaction $A \otimes_L \Delta_{\mathcal{D}}$. The explicit relation is given by the same formulae as in [CM], in the paragraph preceding Proposition 4.12. Since in view of Proposition 4.1 the existence of a left $\mathcal{C} = (A \otimes_L \mathcal{D})$ -colinear right $\mathcal{D}$ -colinear retraction of the $\mathcal{D}$ -coaction $A \otimes_L \Delta_{\mathcal{D}}$ is equivalent to assertion (c) in Theorem 4.3, in [CM, Proposition 4.12] implicitly also the equivalence of assertions (a) and (c) in Theorem 4.3 is proven.

In contrast to [CM], in the current paper the term *total integral* is used only in the more restricted sense of Definition 4.2.

The following proposition extends [BB1, Proposition 4.2]. It clarifies the role of total integrals in bijective entwining structures with a grouplike element.

PROPOSITION 4.4. *Consider a bijective entwining structure $(A, \mathcal{D}, \psi)$ over an algebra L , such that there exists a grouplike element e in $\mathcal{D}$. Let $\mathcal{C} := A \otimes_L \mathcal{D}$ be the associated A -coring. The following assertions are equivalent.*

- (a) A is a relative injective right (resp. left) $\mathcal{C}$ -comodule.
- (b) A is a relative injective right (resp. left) $\mathcal{D}$ -comodule.
- (c) There exists a right (resp. left) total integral in the entwining structure $(A, \mathcal{D}, \psi)$.

If these equivalent conditions hold then $B := A^{\text{co}\mathcal{C}} = A^{\text{co}\mathcal{D}}$ is a direct summand of A as a right (resp. left) B -module.

Proof. (a) $\Rightarrow$ (b) For a relative injective right $\mathcal{C}$ -comodule M , the right $\mathcal{D}$ -coaction has a right A -linear right $\mathcal{D}$ -colinear retraction. Hence it is a relative injective right $\mathcal{D}$ -comodule.

(b) $\Rightarrow$ (a) Assume that A is a relative injective right $\mathcal{D}$ -comodule. Similarly to the proof of [SS, Lemma 4.1], in terms of a right $\mathcal{D}$ -colinear retraction ν_A of the $\mathcal{D}$ -coaction (2.7) in A , a right $\mathcal{C}$ -colinear retraction is given by

$$\mu_A \circ [\nu_A(1_A \otimes_L \bullet) \otimes_L A] \circ \psi^{-1} : A \otimes_L \mathcal{D} \rightarrow A.$$

The equivalence (b) $\Leftrightarrow$ (c) was proven in [BB1, Proposition 4.2], as follows. To a right $\mathcal{D}$ -colinear retraction ν_A of the $\mathcal{D}$ -coaction (2.7) in A , one associates a right total integral $j : d \mapsto \nu_A(1_A \otimes_L d)$. Conversely, in terms of a right total integral j , a right $\mathcal{D}$ -colinear retraction of the $\mathcal{D}$ -coaction (2.7) in A is constructed as $\nu_A := \mu_A \circ (j \otimes_L A) \circ \psi^{-1}$.

It remains to prove the last statement. By property (a), the right $\mathcal{C}$ -coaction in A is a split monomorphism in $\mathfrak{M}^{\mathcal{C}}$. Taking the $\mathcal{C}$ -coinvariants part (with respect to the grouplike element $1_A \otimes_L e$) of its retraction, we obtain right B -linear retraction of the inclusion $B \rightarrow A$.

In order to prove the claim about the left comodule structures, the same arguments can be applied to the entwining structure $(A^{op}, \mathcal{C}_{cop}, \psi^{-1})$ over the algebra L^{op} . $\square$

The following Lemma is a simple generalization of [SS, Lemma 4.1].

LEMMA 4.5. *Let $\mathcal{C}$ be an A -coring possessing a grouplike element g . Assume that A is a relative injective left $\mathcal{C}$ -comodule via the coaction $a \mapsto ag$, determined by g . Denote by $B := A^{co\mathcal{C}}$ the coinvariants of A with respect to g . Then the unit of the adjunction $(\bullet \otimes_B A, (\bullet)^{co\mathcal{C}})$, i.e. the natural transformation (2.1), is an isomorphism.*

Proof. Let M be a relative injective left $\mathcal{C}$ -comodule and ν_M a left $\mathcal{C}$ -colinear retraction of the coaction ${}^M\rho$. Introduce a further map $\xi^M : M \rightarrow \mathcal{C} \otimes_A M$, $m \mapsto g \otimes_A m$. We claim that

$$M \begin{array}{c} \xrightarrow{{}^M\rho} \\ \xleftarrow{\nu_M} \\ \xrightarrow{\xi^M} \end{array} \mathcal{C} \otimes_A M$$

is a contractible pair in ${}_B\mathfrak{M}$. Clearly, all morphisms ${}^M\rho$, ξ^M and ν_M are left B -linear. By definition $\nu_M \circ {}^M\rho = M$. Hence we conclude after observing that, for $m \in M$,

$${}^M\rho \circ \nu_M \circ \xi^M(m) = \nu_M(g \otimes_A m)^{[-1]} \otimes_A \nu_M(g \otimes_A m)^{[0]} = g \otimes_A \nu_M(g \otimes_A m) = \xi^M \circ \nu_M \circ \xi^M(m),$$

where in the second equality the left $\mathcal{C}$ -colinearity of ν_M has been used. In particular we deduce that the equaliser of ${}^A\rho : A \rightarrow \mathcal{C}$, $a \mapsto ag$ and $\xi^A : A \rightarrow \mathcal{C}$, $a \mapsto ga$ cosplits in ${}_B\mathfrak{M}$. Hence it is preserved by the functor $N \otimes_B \bullet : {}_B\mathfrak{M} \rightarrow \mathfrak{M}_k$, for any right B -module N . Recall that A is a right $\mathcal{C}$ -comodule with coaction ξ^A and $N \otimes_B A$ is a right $\mathcal{C}$ -comodule with coaction $N \otimes_B \xi^A$. Therefore

$$(N \otimes_B A)^{co\mathcal{C}} = \text{Ker}(N \otimes_B {}^A\rho - N \otimes_B \xi^A) = N \otimes_B \text{Ker}({}^A\rho - \xi^A) = N \otimes_B B \cong N.$$

This proves that (2.1) is a natural isomorphism, as stated. $\square$

The following proposition formulates a functorial criterion for a coring with a grouplike element to be a Galois coring.

PROPOSITION 4.6. *Let $\mathcal{C}$ be an A -coring possessing a grouplike element g . Denote by $B := A^{co\mathcal{C}}$ the coinvariants of A with respect to g . Consider the adjunction $(\bullet \otimes_B A, (\bullet)^{co\mathcal{C}})$ in Section 2.3 and the canonical map can in (2.3). The following statements hold.*

- 1) *can is an epimorphism if and only if the functor $(\bullet)^{co\mathcal{C}}$ is $(\bullet \otimes_A \mathcal{C}, \mathfrak{M}^{\mathcal{C}})$ -faithful.*
- 2) *can is a split monomorphism if and only if the functor $(\bullet)^{co\mathcal{C}}$ is $(\bullet \otimes_A \mathcal{C}, \mathfrak{M}^{\mathcal{C}})$ -full.*

In particular, $\mathcal{C}$ is a Galois coring if and only if the functor $(\bullet)^{co\mathcal{C}}$ is $(\bullet \otimes_A \mathcal{C}, \mathfrak{M}^{\mathcal{C}})$ -fully faithful.

Proof. Denote the counit of the coring $\mathcal{C}$ by ϵ . For any right A -module M , the counit n of the adjunction $(\bullet \otimes_B A, (\bullet)^{co\mathcal{C}})$ (cf. (2.2)) is subject to the equality of maps $(M \otimes_A \mathcal{C})^{co\mathcal{C}} \otimes_B A \rightarrow M \otimes_A \mathcal{C}$,

$$(M \otimes_A \text{can}) \circ (M \otimes_A \epsilon \otimes_B A) = n_{M \otimes_A \mathcal{C}}.$$

Since the restriction of $M \otimes_A \epsilon$ is an isomorphism $(M \otimes_A \mathcal{C})^{co\mathcal{C}} \rightarrow M \otimes_A A$, the claims follow by Theorem 3.14 1) and 2), respectively. $\square$

5. COMODULE ALGEBRAS OF HOPF ALGEBROIDS

Consider a Hopf algebroid $\mathcal{H}$, with constituent left bialgebroid $\mathcal{H}_L$ over the base algebra L and right bialgebroid $\mathcal{H}_R$ over R , and a right $\mathcal{H}$ -comodule algebra A . Recall (from Section 2.10) that latter means a right $\mathcal{H}_R$ -comodule algebra, or equivalently a right $\mathcal{H}_L$ -comodule algebra, A with coactions $a \mapsto a^{[0]} \otimes_R a^{[1]}$ and $a \mapsto a_{[0]} \otimes_L a_{[1]}$, respectively, related as in (2.16) and (2.17). It has been recalled in Section 2.8 that a right $\mathcal{H}$ -comodule algebra A determines an entwining structure over the algebra R , with R -ring A and R -coring (with grouplike element 1_H) underlying $\mathcal{H}_R$. Hence there is an associated A -coring $\mathcal{C} := A \otimes_R \mathcal{H}_R$. What is more, the L -coring $\mathcal{D}$ underlying $\mathcal{H}_L$ is a right extension of $\mathcal{C}$ (cf. Section 2.11). In this section we analyse the functors in Figure (2.4), associated to this coring extension $\mathcal{D}$ of $\mathcal{C}$. As before, we focus on $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separability of the forgetful functor $\mathbb{U}^{\mathcal{D}}$. Note that the above coring extension $\mathcal{D}$ of $\mathcal{C}$ does not arise from any $(L-)$ entwining structure. Since in this way the coring extension $\mathcal{D}$ of $\mathcal{C}$ is not of the type considered in Theorem 4.3, $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separability of $\mathbb{U}^{\mathcal{D}}$ can not be replaced by separability of the second kind of any other functor.

Application of Proposition 4.1 to the coring extension $\mathcal{D} = (H, \gamma_L, \pi_L)$ of $\mathcal{C} = (A \otimes_R H, A \otimes_R \gamma_R, A \otimes_R \pi_R)$ yields the following generalisation of Doi's theorem [Doi, Theorem 1.6].

THEOREM 5.1. *Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $(H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $(H, R, s_R, t_R, \gamma_R, \pi_R)$, and antipode S . Let A be a right $\mathcal{H}$ -comodule algebra. The following assertions are equivalent.*

- (a) *There exists a right total integral in the R -entwining structure (2.12) with grouplike element 1_H , i.e. a morphism $j \in \text{Hom}^{\mathcal{H}}(H, A)$, normalised as $j(1_H) = 1_A$.*
- (b) *There exists a right total integral in the bijective L -entwining structure (2.19) with grouplike element 1_H , i.e. a morphism $j^{op} \in \text{Hom}^{\mathcal{H}}(H, A)$, normalised as $j^{op}(1_H) = 1_A$.*
- (c) *A is a relative injective right comodule for the L -coring (H, γ_L, π_L) .*
- (d) *Any object in $\mathfrak{M}_A^{\mathcal{H}}$ is relative injective as a right comodule for the L -coring (H, γ_L, π_L) .*
- (e) *The forgetful functor $\mathfrak{M}^{\mathcal{H}} \rightarrow \mathfrak{M}_L$ is $(\mathfrak{M}^{\mathcal{H}}, \mathbb{R})$ -separable, where $\mathbb{R}$ denotes the forgetful functor $\mathfrak{M}_A^{\mathcal{H}} \rightarrow \mathfrak{M}^{\mathcal{H}}$.*

If the antipode of the Hopf algebroid $\mathcal{H}$ is bijective then the above assertions are equivalent also to the following ones.

- (f) *A is a relative injective right comodule for the R -coring (H, γ_R, π_R) .*
- (g) *Any object in ${}_A\mathfrak{M}^{\mathcal{H}}$ is relative injective as a right comodule for the R -coring (H, γ_R, π_R) .*
- (h) *The forgetful functor $\mathfrak{M}^{\mathcal{H}} \rightarrow \mathfrak{M}_R$ is $(\mathfrak{M}^{\mathcal{H}}, \mathbb{R}^{op})$ -separable, where $\mathbb{R}^{op}$ denotes the forgetful functor ${}_A\mathfrak{M}^{\mathcal{H}} \rightarrow \mathfrak{M}^{\mathcal{H}}$.*
- (i) *There exists a left total integral in the bijective R -entwining structure (2.12) with grouplike element 1_H , i.e. a left $\mathcal{H}$ -colinear map with respect to the coaction (2.27) in A , $j_{cop} : H \rightarrow A$, normalised as $j_{cop}(1_H) = 1_A$.*
- (j) *There exists a left total integral in the bijective L -entwining structure (2.19) with grouplike element 1_H , i.e. a left $\mathcal{H}$ -colinear map with respect to the coaction (2.20) in A , $j_{cop}^{op} : H \rightarrow A$, normalised as $j_{cop}^{op}(1_H) = 1_A$.*
- (k) *A (with coaction (2.26)) is a relative injective left comodule for the R -coring (H, γ_R, π_R) .*
- (l) *Any object in ${}^{\mathcal{H}}\mathfrak{M}_A$ is relative injective as a left comodule for the R -coring (H, γ_R, π_R) .*
- (m) *The forgetful functor ${}^{\mathcal{H}}\mathfrak{M} \rightarrow {}_R\mathfrak{M}$ is $({}^{\mathcal{H}}\mathfrak{M}, \mathbb{R}_{cop}^{op})$ -separable, where $\mathbb{R}_{cop}^{op}$ denotes the forgetful functor ${}^{\mathcal{H}}\mathfrak{M}_A \rightarrow {}^{\mathcal{H}}\mathfrak{M}$.*
- (n) *A (with coaction (2.20)) is a relative injective left comodule for the L -coring (H, γ_L, π_L) .*
- (o) *Any object in ${}^{\mathcal{H}}\mathfrak{M}_A$ is relative injective as a left comodule for the L -coring (H, γ_L, π_L) .*
- (p) *The forgetful functor ${}^{\mathcal{H}}\mathfrak{M} \rightarrow {}_L\mathfrak{M}$ is $({}^{\mathcal{H}}\mathfrak{M}, \mathbb{R}_{cop})$ -separable, where $\mathbb{R}_{cop}$ denotes the forgetful functor ${}^{\mathcal{H}}\mathfrak{M}_A \rightarrow {}^{\mathcal{H}}\mathfrak{M}$.*

Proof. (a) $\Leftrightarrow$ (b) This equivalence, with correspondence $j^{op} = j$, is obvious by [BB2, Theorem 2.2], cf. Section 2.10.

(b) $\Leftrightarrow$ (c) This equivalence follows by Proposition 4.4 (b) $\Leftrightarrow$ (c), since the entwining map (2.19) is bijective.

(a) $\Rightarrow$ (e) The claim is proven by applying Proposition 4.1 to the coring extension $\mathcal{D} = (H, \gamma_L, \pi_L)$ of $\mathcal{C} = (A \otimes_R H, A \otimes_R \gamma_R, A \otimes_R \pi_R)$. That is, we need to construct a left $(A \otimes_R H, A \otimes_R \gamma_R, A \otimes_R \pi_R)$ -colinear right $\mathcal{H}$ -colinear retraction of $A \otimes_R \gamma_L$. Take a right total integral j and put

$$(5.1) \quad \nu_{A \otimes_R H} : A \otimes_R H \otimes_L H \rightarrow A \otimes_R H, \quad a \otimes_R h \otimes_L k \mapsto aj(S(h)k^{(1)}) \otimes_R k^{(2)}.$$

Since j is right $\mathcal{H}$ -colinear, it is R - R bilinear in the sense that $j(s_R(r)hs_R(r')) = rj(h)r'$, for $h \in H$ and $r, r' \in R$. The antipode satisfies $S(ht_R(r)) = s_R(r)S(h)$ and $S(t_L(l)h) = S(h)s_L(l)$, for $h \in H$, $r \in R$ and $l \in L$. The coproduct γ_R is left L -linear. These considerations imply that $\nu_{A \otimes_R H}$ is a well defined map. It is a retraction of $A \otimes_R \gamma_L$ by the right $\mathcal{H}_R$ -colinearity of γ_L . That is,

$$\begin{aligned} \nu_{A \otimes_R H} \circ (A \otimes_R \gamma_L)(a \otimes_R h) &= aj(S(h_{(1)})h_{(2)}^{(1)}) \otimes_R h_{(2)}^{(2)} = aj(S(h^{(1)}_{(1)})h^{(1)}_{(2)}) \otimes_R h^{(2)} \\ &= aj \circ s_R \circ \pi_R(h^{(1)}) \otimes_R h^{(2)} = a \otimes_R h^{(2)} t_R \circ \pi_R(h^{(1)}) = a \otimes_R h, \end{aligned}$$

where the third equality follows by the antipode axiom and the penultimate equality follows by the left (or right) R -linearity and normalisation of j . Right $\mathcal{H}$ -colinearity of $\nu_{A \otimes_R H}$ follows by the coassociativity of γ_R . Its left A -linearity is obvious. Left $(A \otimes_R H, A \otimes_R \gamma_R, A \otimes_R \pi_R)$ -colinearity is checked as follows. (We freely omit canonical isomorphisms of the type $M \otimes_A (A \otimes_R H) \cong M \otimes_R H$.)

$$\begin{aligned} [(A \otimes_R H) \otimes_A \nu_{A \otimes_R H}] \circ (A \otimes_R \gamma_R \otimes_L H)(a \otimes_R h \otimes_L k) &= aj(S(h^{(2)})k^{(1)})^{[0]} \otimes_R h^{(1)} j(S(h^{(2)})k^{(1)})^{[1]} \otimes_R k^{(2)} \\ &= aj(S(h^{(2)}_{(2)})k^{(1)}) \otimes_R h^{(1)} S(h^{(2)}_{(1)})k^{(2)} \otimes_R k^{(3)} \\ &= aj(S(h_{(2)})k^{(1)}) \otimes_R h_{(1)}^{(1)} S(h_{(1)}^{(2)})k^{(2)} \otimes_R k^{(3)} \\ &= aj(S(h_{(2)})k^{(1)}) \otimes_R s_L \circ \pi_L(h_{(1)})k^{(2)} \otimes_R k^{(3)} \\ &= aj(S(h)k^{(1)}) \otimes_R k^{(2)} \otimes_R k^{(3)} \\ &= (A \otimes_R \gamma_R) \circ \nu_{A \otimes_R H}(a \otimes_R h \otimes_L k), \end{aligned}$$

where the second equality follows by the right $\mathcal{H}$ -colinearity of j and the anti-comultiplicativity of S , the third one does by the right $\mathcal{H}_L$ -colinearity of γ_R , and the fourth one does by the antipode axiom. The fifth equality follows by the Takeuchi axiom and anti-multiplicativity of S . Indeed, in a Hopf algebroid $s_L = t_R \circ \pi_R \circ s_L$ and $s_R = S \circ t_R$. Hence the Takeuchi axiom implies that $k^{(1)} \otimes_R s_L(l)k^{(2)} = S \circ s_L(l)k^{(1)} \otimes_R k^{(2)}$ for $k \in H$ and $l \in L$.

(e) $\Rightarrow$ (d) The forgetful functor $\mathfrak{M}^{\mathcal{H}} \rightarrow \mathfrak{M}_L$ has a right adjoint, the functor $\bullet \otimes_L H$. Hence the claim follows by Corollary 3.10 2).

(d) $\Rightarrow$ (c) This implication is trivial as A itself is an object in $\mathfrak{M}_A^{\mathcal{H}}$.

If the antipode is bijective then implications (b) $\Rightarrow$ (h) $\Rightarrow$ (g) $\Rightarrow$ (f) $\Leftrightarrow$ (a) follow by applying (a) $\Rightarrow$ (e) $\Rightarrow$ (d) $\Rightarrow$ (c) $\Leftrightarrow$ (b) to the opposite Hopf algebroid $\mathcal{H}^{op}$ and its right comodule algebra A^{op} (with $(\mathcal{H}_R)^{op}$ -coaction $a \mapsto a^{[0]} \otimes_R a^{[1]}$). Hence we have the equivalence of assertions (a)-(h) proven.

Recall that the coactions (2.20) and (2.27) differ by the isomorphism $S^2 \otimes_L A$. Hence A is a relative injective left $\mathcal{H}_L$ -comodule with respect to the coaction (2.20) if and only if it is relative injective with respect to the coaction (2.27). Similarly, relative injectivity of A as a left $\mathcal{H}_R$ -comodule with respect to the coactions $a \mapsto S(a_{[1]}) \otimes_R a_{[0]}$ and $a \mapsto S^{-1}(a_{[1]}) \otimes_R a_{[0]}$ are equivalent properties. Hence application of the implications (a) $\Rightarrow$ (e) $\Rightarrow$ (d) $\Rightarrow$ (c) $\Leftrightarrow$ (b) to the opposite Hopf algebroid $\mathcal{H}_{cop}$ and its right comodule algebra A^{op} , with $\mathcal{H}_L$ -coaction (2.27), implies (i) $\Rightarrow$ (p) $\Rightarrow$ (o) $\Rightarrow$ (n) $\Leftrightarrow$ (j). Similarly, applying (a) $\Rightarrow$ (e) $\Rightarrow$ (d) $\Rightarrow$ (c) $\Leftrightarrow$ (b) to the Hopf algebroid $\mathcal{H}_{cop}^{op}$ and its right comodule algebra A , with coaction (2.20), we conclude that (j) $\Rightarrow$ (m) $\Rightarrow$ (l) $\Rightarrow$ (k) $\Leftrightarrow$ (i). This proves the equivalence of assertions (i)-(p).

It remains to prove the equivalences $(a) \Leftrightarrow (i)$ and $(a) \Leftrightarrow (j)$. For a right total integral j as in part (a), consider the map $j_{cop} := j \circ S$. By the unitality of the antipode it is normalised, i.e. $j \circ S(1_H) = 1_A$. Its left $\mathcal{H}$ -colinearity with respect to the coaction (2.27) follows by the colinearity of j and the anti-comultiplicativity of S . That is, for $h \in H$,

$$S^{-1}((j \circ S)(h)^{[1]}) \otimes_L (j \circ S)(h)^{[0]} = S^{-1}(S(h)^{(2)}) \otimes_L j(S(h)^{(1)}) = h_{(1)} \otimes_L (j \circ S)(h_{(2)}).$$

In a similar way, for a left total integral j_{cop} as in part (i), $j_{cop} \circ S^{-1}$ is a right total integral as in part (a). This proves the equivalence $(a) \Leftrightarrow (i)$. A left total integral j_{cop}^{op} in part (j) is related to a right total integral j as in part (a) via the bijection $j \mapsto j_{cop}^{op} := j \circ S^{-1}$. $\square$

Theorem 5.1 makes us able to answer a question which was left open in [Bö2].

PROPOSITION 5.2. *Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $(H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $(H, R, s_R, t_R, \gamma_R, \pi_R)$, and a bijective antipode S . Assume that H is a projective left R -module via t_R and a projective right comodule for the R -coring (H, γ_R, π_R) via γ_R . (These assumptions hold e.g. if H is a finitely generated and projective left or right L or R -module, cf. [Bö2, Section 4].) Then the following assertions on a right $\mathcal{H}$ -Galois extension $B \subseteq A$ are equivalent.*

- (a) A is a faithfully flat right B -module.
- (b) B is a direct summand of the right B -module A .
- (c) The functors $A \otimes_B \bullet : {}_B\mathfrak{M} \rightarrow {}_A^{\mathcal{H}}\mathfrak{M}$ and ${}^{co\mathcal{H}}(\bullet) : {}_A^{\mathcal{H}}\mathfrak{M} \rightarrow {}_B\mathfrak{M}$ are inverse equivalences and $H \otimes_R A$ is a flat right A -module.
- (d) A is a projective generator in ${}_A^{\mathcal{H}}\mathfrak{M}$ and $H \otimes_R A$ is a flat right A -module.
- (e) A is a generator of right B -modules.
- (f) A is a faithfully flat left B -module.
- (g) B is a direct summand of the left B -module A .
- (h) The functors $\bullet \otimes_B A : \mathfrak{M}_B \rightarrow \mathfrak{M}_A^{\mathcal{H}}$ and $(\bullet)^{co\mathcal{H}} : \mathfrak{M}_A^{\mathcal{H}} \rightarrow \mathfrak{M}_B$ are inverse equivalences.
- (i) A is a projective generator in $\mathfrak{M}_A^{\mathcal{H}}$.
- (j) A is a generator of left B -modules.
- (k) A is a relative injective $\mathcal{H}$ -comodule. That is, any – and hence all – of the equivalent properties, that A is a relative injective left or right comodule for the L -coring (H, γ_L, π_L) or the R -coring (H, γ_R, π_R) , hold.

Proof. $(a) \Leftrightarrow (b)$ and $(f) \Leftrightarrow (g)$ These equivalences follow by [Row, 2.11.29], as A is a projective left and right B -module by [Bö2, Proposition 4.2].

$(b) \Rightarrow (k)$ and $(g) \Rightarrow (k)$ Since a comodule algebra for a Hopf algebroid with a bijective antipode determines bijective entwining structures (2.12) and (2.19), these implications follow by [BB1, Proposition 4.1] (which is a simple generalisation of [SS, Remark 4.2] to the case of non-commutative base algebras), and Theorem 5.1 $(c) \Leftrightarrow (f) \Leftrightarrow (k) \Leftrightarrow (n)$.

$(k) \Rightarrow (b)$ and $(k) \Rightarrow (g)$ These assertions follow by Proposition 4.4.

$(a) \Leftrightarrow (d) \Leftrightarrow (c)$ These equivalences follow by the *Galois Coring Structure Theorem* [BW, 28.19 (2)].

$(f) \Leftrightarrow (i) \Leftrightarrow (h)$ Since H is a projective left R -module by assumption, $A \otimes_R H$ is a projective (hence flat) left A -module. Therefore also these equivalences follow by the *Galois Coring Structure Theorem* [BW, 28.19 (2)].

$(b) \Rightarrow (e)$ and $(g) \Rightarrow (j)$ These implications are trivial.

$(e) \Rightarrow (b)$ and $(j) \Rightarrow (g)$ A is a generator of right (respectively, left) B -modules if and only if there exist finite sets $\{a_i\}$ in A and $\{\alpha_i\}$ in $\text{Hom}_B(A, B)$ (respectively, in ${}_B\text{Hom}(A, B)$), satisfying $\sum_i \alpha_i(a_i) = 1_B$. In terms of these elements, a right B -linear retraction of the inclusion $B \rightarrow A$ is given by the map $a \mapsto \sum_i \alpha_i(a_i a)$ (respectively, a left B -linear retraction of the inclusion $B \rightarrow A$ is given by the map $a \mapsto \sum_i \alpha_i(aa_i)$). $\square$

Applying Proposition 5.2 to the co-opposite Hopf algebroid $\mathcal{H}_{cop}$, we see that the claims in Proposition 5.2 – with the only modification that claims (h) and (i) need to be supplemented by the assertion that $A \otimes_R H$ is a flat left A -module – can be proven alternatively by replacing the

assumptions about the projectivity of H a left R -module (via t_R) and a right $\mathcal{H}$ -comodule (via the coproducts) with the assumptions that it is a projective right R -module (via s_R) and a projective left $\mathcal{H}$ -comodule (via the coproducts).

6. A SCHNEIDER TYPE THEOREM

This section contains the main result of the paper, Theorem 6.7. The starting point of our study is the following result [BTW, Theorem 2.1]. Recall that a right comodule P for an A -coring $\mathcal{C}$, which is a finitely generated and projective right A -module, is a *Galois comodule* if the canonical map

$$(6.1) \quad \text{can} : \text{Hom}_A(P, A) \otimes_S P \rightarrow \mathcal{C}, \quad \phi \otimes_S p \mapsto \phi(p^{[0]})p^{[1]}$$

is bijective, where $S := \text{End}^{\mathcal{C}}(P)$. Assume that S is a T -ring (e.g. T is a subalgebra of S). Denote $P^* = \text{Hom}_A(P, A)$. A symmetrical (and slightly extended) version of [BTW, Theorem 2.1], formulated for right comodules, is the following.

THEOREM 6.1. *The canonical map (6.1) is bijective and P^* is a T -relative projective right S -module provided that the following conditions hold true.*

- i) *The map $P^* \otimes_T S \rightarrow \text{Hom}^{\mathcal{C}}(P, P^* \otimes_T P)$, $\xi \otimes_T s \mapsto (p \mapsto \xi \otimes_T s(p))$ is an isomorphism (of right S -modules);*
- ii) *The lifted canonical map,*

$$(6.2) \quad \widetilde{\text{can}}^T : P^* \otimes_T P \rightarrow \mathcal{C}, \quad \phi \otimes_T p \mapsto \phi(p^{[0]})p^{[1]}$$

is a split epimorphism of right $\mathcal{C}$ -comodules.

Motivated by this result, in the present section we investigate how one can use the $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separability of the functor $\mathbb{U}^{\mathcal{D}} : \mathfrak{M}^{\mathcal{D}} \rightarrow \mathfrak{M}_L$ on Figure (2.4), to derive properties i) and ii) in Theorem 6.1 for a coring extension $\mathcal{D}$ of $\mathcal{C}$. As before, we are motivated by the case when the coring extension comes from a Hopf algebroid extension. We start with a lemma formulating sufficient conditions for property ii) in Theorem 6.1.

LEMMA 6.2. *Consider an L -coring $\mathcal{D}$ which is a right extension of an A -coring $\mathcal{C}$ and the functors in Figure (2.4). Assume the following properties.*

- a) *The forgetful functor $\mathbb{U}^{\mathcal{D}} : \mathfrak{M}^{\mathcal{D}} \rightarrow \mathfrak{M}_L$ is $(\mathfrak{M}^{\mathcal{D}}, \mathbb{R})$ -separable.*
- b) *The right regular $\mathcal{C}$ -comodule is $\mathcal{E}_{\mathbb{R}}$ -projective.*

If these properties hold, then the lifted canonical map (6.2) is a split epimorphism of right $\mathcal{C}$ -comodules if and only if it is a split epimorphism of right L -modules.

Proof. Note that the map (6.2) is right $\mathcal{C}$ -colinear. Hence, by assumption a), Theorem 3.9 1) implies that it is a split epimorphism in $\mathfrak{M}^{\mathcal{D}}$ whenever it is a split epimorphism in $\mathfrak{M}_L$. Then (6.2) is a split epimorphism of right $\mathcal{C}$ -comodules by assumption b). The converse implication is trivial since any right $\mathcal{C}$ -comodule map is right L -linear. $\square$

REMARK 6.3. If the coring extension $\mathcal{D}$ of $\mathcal{C} := A \otimes_L \mathcal{D}$ comes from a *bijective* L -entwining structure $(A, \mathcal{D}, \psi)$, then assumption b) in Lemma 6.2 holds true by Theorem 3.2. Indeed, the forgetful functor $\mathfrak{M}_A^{\mathcal{D}}(\psi) \rightarrow \mathfrak{M}^{\mathcal{D}}$ possesses a left adjoint, the functor $\mathbb{T} = \bullet \otimes_L A$, and $\mathcal{C} := A \otimes_L \mathcal{D}$ is isomorphic as a right-right entwined module to $\mathcal{D} \otimes_L A = \mathbb{T}(\mathcal{D})$ via ψ .

In particular, if $\mathcal{H}$ is a Hopf algebroid over the base algebras L and R with a bijective antipode, and A is a right $\mathcal{H}$ -comodule algebra, then the right regular comodule for the A -coring $A \otimes_R H$ is $\mathcal{E}_{\mathbb{R}}$ -projective, where $\mathbb{R}$ is the forgetful functor $\mathfrak{M}_A^{\mathcal{H}} \rightarrow \mathfrak{M}^{\mathcal{H}}$.

REMARK 6.4. Consider an L -coring $\mathcal{D}$ which is a right extension of an A -coring $\mathcal{C}$ and the corresponding functors in Figure (2.4). The property in Lemma 6.2, saying that the (right $\mathcal{C}$ -colinear) map (6.2) belongs to $\mathcal{E}_{\mathbb{U}^{\mathcal{D}} \mathbb{R}}$, i.e. that it is a split epimorphism of right L -modules, holds in various situations.

1) *If the (right L -linear) canonical map (6.1) is surjective and $\mathcal{C}$ is a projective right L -module.* In the particular case of a comodule algebra A of a Hopf algebroid $\mathcal{H}$ over base algebras L and

R , the corresponding A -coring $\mathcal{C} = A \otimes_R H$ is a projective right L -module provided that H is a projective right L -module (via the target map of the constituent left bialgebroid) and A is a projective right R -module.

2) If the (right A -linear) canonical map (6.1) is surjective, $\mathcal{C}$ is a projective right A -module and $\mathcal{E}_{\mathbb{U}^{\mathcal{C}}} \subseteq \mathcal{E}_{\mathbb{U}^{\mathcal{D}}}$.

Let $\mathcal{H}$ be a Hopf algebroid with a bijective antipode, over base algebras L and R . In the particular example of an $\mathcal{H}$ -comodule algebra A , the corresponding A -coring $\mathcal{C} = A \otimes_R H \cong H \otimes_R A$ is a projective right A -module provided that H is a projective right R -module (via the source map of the constituent right bialgebroid).

The condition $\mathcal{E}_{\mathbb{U}^{\mathcal{C}}} \subseteq \mathcal{E}_{\mathbb{U}^{\mathcal{D}}}$ holds whenever dealing with a comodule algebra A of a Hopf algebra $\mathcal{H}$ over a commutative ring k . Indeed, in this case the role of the coring $\mathcal{D}$ is played by the k -coalgebra underlying the Hopf algebra $\mathcal{H}$. The A -coring $\mathcal{C}$ is equal to $A \otimes_k H$. All functors $\mathbb{U}^{\mathcal{C}}$, $\mathbb{U}^{\mathcal{D}}$ and $\mathbb{R}$ are forgetful functors. A fourth forgetful functor $\mathfrak{M}_A \rightarrow \mathfrak{M}_k$ makes the following diagram commutative.

$$\begin{array}{ccc} \mathfrak{M}_A^{\mathcal{H}} & \xrightarrow{\mathbb{R}} & \mathfrak{M}^{\mathcal{H}} \\ \mathbb{U}^{\mathcal{C}} \downarrow & & \downarrow \mathbb{U}^{\mathcal{D}} \\ \mathfrak{M}_A & \longrightarrow & \mathfrak{M}_k \end{array}$$

This proves that in this case $\mathcal{E}_{\mathbb{U}^{\mathcal{C}}} \subseteq \mathcal{E}_{\mathbb{U}^{\mathcal{D}}}$, thus assumptions 2) hold e.g. in Schneider's theorem [Schn, Theorem I].

REMARK 6.5. In the particular case when the right $\mathcal{C}$ -comodule P in Theorem 6.1 is equal to the base algebra A , property i) reduces to $(A \otimes_T A)^{\text{co}\mathcal{C}} = A \otimes_T A^{\text{co}\mathcal{C}}$. Let us investigate this condition. Note that, for an A -coring $\mathcal{C}$ possessing a grouplike element g , and any right T -module V , $V \otimes_T A$ is a right $\mathcal{C}$ -comodule via the comodule structure of A . There is an obvious inclusion $V \otimes_T A^{\text{co}\mathcal{C}} \rightarrow (V \otimes_T A)^{\text{co}\mathcal{C}}$, which is an isomorphism in appropriate situations: e.g. if V is a flat T -module, or in the situation described in Lemma 4.5. Indeed, in the last case, by applying Lemma 4.5 to a right $B := A^{\text{co}\mathcal{C}}$ -module $V \otimes_T B$, for a right T -module V , we conclude that $(V \otimes_T A)^{\text{co}\mathcal{C}} = V \otimes_T B$ whenever A is a relative injective left $\mathcal{C}$ -comodule.

As it is explained in Section 2.14, to a right comodule algebra A of a Hopf algebroid $\mathcal{H}$ (with constituent left and right bialgebroids $(H, L, s_L, t_L, \gamma_L, \pi_L)$ and $(H, R, s_R, t_R, \gamma_R, \pi_R)$, and a bijective antipode S) one associates two isomorphic A -corings, on the k -modules $A \otimes_R H$ and $H \otimes_R A$, and two isomorphic A^{op} -corings, on the k -modules $A \otimes_L H$ and $H \otimes_L A$. These A - and A^{op} -corings are anti-isomorphic, cf. (2.24). The grouplike element 1_H , in the L - and R -corings underlying $\mathcal{H}$, determines grouplike elements in all associated A - and A^{op} -corings (preserved by the coring (anti-) isomorphisms (2.12), (2.19) and (2.24) between them). That is, A (or A^{op}) is a right comodule in each case. Corresponding to the four corings, there are four canonical maps of the type (6.1), which differ by the respective coring (anti-) isomorphisms in Section 2.14. Following Theorem 6.7 is formulated in terms of the A -coring $A \otimes_R H$ and the corresponding canonical map (2.13). Certainly, all claims can be reformulated in terms of any of the other three (anti-) isomorphic corings.

Let $\mathcal{H}$ be a Hopf algebroid over base algebras L and R . Recall that a right $\mathcal{H}$ -comodule algebra A is an R -ring. Assume that the coinvariant subalgebra $B := A^{\text{co}\mathcal{H}}$ is a T -ring (e.g. T is a k -subalgebra of B). Consider the lifted version of the canonical map (2.13)

$$(6.3) \quad \widetilde{\text{can}}^T : A \otimes_T A \rightarrow A \otimes_R H, \quad a \otimes_T a' \mapsto aa'^{[0]} \otimes_R a'^{[1]}.$$

Note that it is right L -linear with respect to the module structures

$$(6.4) \quad (a \otimes_T a')l = a \otimes_T \pi_R \circ t_L(l)a' \quad \text{and} \quad (a \otimes_R h)l = a \otimes_R t_L(l)h,$$

for $a \otimes_T a' \in A \otimes_T A$, $a \otimes_R h \in A \otimes_R H$ and $l \in L$. Moreover, the lifted canonical map (6.3) is also left L -linear with respect to the module structures

$$(6.5) \quad l(a \otimes_T a') = a\pi_R \circ s_L(l) \otimes_T a' \quad \text{and} \quad l(a \otimes_R h) = a \otimes_R s_L(l)h,$$

for $a \otimes_T a' \in A \otimes_T A$, $a \otimes_R h \in A \otimes_R H$ and $l \in L$.

LEMMA 6.6. *Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $(H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $(H, R, s_R, t_R, \gamma_R, \pi_R)$, and a bijective antipode S . Let A be a right $\mathcal{H}$ -comodule algebra which is a relative injective right $\mathcal{H}_R$ -comodule. Assume that $A^{\text{co}\mathcal{H}}$ is a T -ring (e.g. T is a subalgebra of $A^{\text{co}\mathcal{H}}$). Assume furthermore that the lifted canonical map (6.3) possesses a right L -module section ζ_0^T (with respect to the module structures (6.4)). Then (6.3) possesses a section in $\mathfrak{M}_A^{\mathcal{H}}$, given as*

$$\begin{aligned} \zeta^T : A \otimes_R H &\rightarrow A \otimes_T A, \\ a \otimes_R h &\mapsto \zeta_0^T(1_A \otimes_R h_{(1)} S^{-1}(a_{[1]})_{(1)})^{[0]} j \left(S(\zeta_0^T(1_A \otimes_R h_{(1)} S^{-1}(a_{[1]})_{(1)})^{[1]}) h_{(2)} S^{-1}(a_{[1]})_{(2)} \right) a_{[0]}, \end{aligned} \quad (6.6)$$

where $a \mapsto a_{[0]} \otimes_L a_{[1]}$ and $a \mapsto a^{[0]} \otimes_R a^{[1]}$ are the $\mathcal{H}_L$, and $\mathcal{H}_R$ -coactions in A , respectively, related via (2.16)-(2.17), and $\gamma_L(h) = h_{(1)} \otimes_L h_{(2)}$, for $h \in H$.

Proof. Theorem 5.1 (f) $\Rightarrow$ (e) implies that the forgetful functor $\mathfrak{M}^{\mathcal{H}} \rightarrow \mathfrak{M}_L$ is $(\mathfrak{M}^{\mathcal{H}}, \mathbb{R})$ -separable, where $\mathbb{R}$ is the forgetful functor $\mathfrak{M}_A^{\mathcal{H}} \rightarrow \mathfrak{M}^{\mathcal{H}}$. Furthermore, by Remark 6.3, the right regular comodule for the A -coring $A \otimes_R H$ is $\mathcal{E}_{\mathbb{R}}$ -projective. Thus the lifted canonical map (6.3) is a split epimorphism in $\mathfrak{M}_A^{\mathcal{H}}$, by Lemma 6.2. A section can be explicitly constructed as follows. By Theorem 3.9 1) the lifted canonical map (6.3) is a split epimorphism of right $\mathcal{H}$ -comodules. A right $\mathcal{H}$ -comodule section ζ_1^T can be constructed using arguments in the proof of the Rafael type theorem 3.13 3). Indeed, in terms of a right L -module section ζ_0^T and a natural retraction ν of $\tau_{\mathbb{R}}(\bullet)$ (where τ denotes the $\mathcal{H}_L$ -coaction, i.e. the unit of the adjunction of the forgetful functor $\mathfrak{M}^{\mathcal{H}} \rightarrow \mathfrak{M}_L$ and the induction functor $\bullet \otimes_L H : \mathfrak{M}_L \rightarrow \mathfrak{M}^{\mathcal{H}}$), one can put

$$\zeta_1^T := \nu_{A \otimes_T A} \circ (\zeta_0^T \otimes_L H) \circ (A \otimes_R \gamma_L).$$

The natural retraction ν was constructed in the proofs of Proposition 4.1 and Theorem 5.1 (a) $\Rightarrow$ (e). It follows by (4.1) that $\nu_{A \otimes_T A} = A \otimes_T \nu_A$ and $\nu_A = (A \otimes_R \pi_R) \circ \nu_{A \otimes_R H} \circ (\varrho^A \otimes_L H)$. The map $\nu_{A \otimes_R H}$ was related to a total integral $j : H \rightarrow A$ in (5.1). Its application yields the explicit form $\nu_A(a \otimes_L h) = a^{[0]} j(S(a^{[1]})h)$.

Furthermore, by Remark 6.3, $A \otimes_R H \in \mathfrak{M}_A^{\mathcal{H}}$ is isomorphic to $H \otimes_R A$ via the entwining map ψ_R in (2.12), with inverse (2.25). Hence $A \otimes_R H$ is $\mathcal{E}_{\mathbb{R}}$ -projective. Thus a right A -module right $\mathcal{H}$ -comodule section of the lifted canonical map (6.3) is given by

$$\zeta^T = (A \otimes_T \mu) \circ (\zeta_1^T \otimes_R A) \circ (\psi_R \otimes_R A) \circ (H \otimes_R \eta \otimes_R A) \circ \psi_R^{-1},$$

where $\eta : R \rightarrow A$ and $\mu : A \otimes_R A \rightarrow A$ are unit and multiplication maps in the R -ring A , respectively. The map ζ^T comes out explicitly as in (6.6). $\square$

THEOREM 6.7. *Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $(H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $(H, R, s_R, t_R, \gamma_R, \pi_R)$, and a bijective antipode S . Let A be a right $\mathcal{H}$ -comodule algebra and put $B := A^{\text{co}\mathcal{H}}$. Assume that B is a T -ring (e.g. T is a k -subalgebra of B).*

- 1) *If the lifted canonical map (6.3) is a split epimorphism of right L -modules (with respect to the module structures (6.4)) then the following assertions are equivalent.*
 - (a) *The canonical map (2.13) is bijective and B is a direct summand of A as a right B -module (hence A is a generator of right B -modules).*
 - (b) *A is a relative injective right comodule for the R -coring (H, γ_R, π_R) .*
 - (c) *The functor ${}^{\text{co}\mathcal{H}}(\bullet) : {}^{\mathcal{H}}\mathfrak{M} \rightarrow {}_B\mathfrak{M}$ is an equivalence, with inverse $A \otimes_B \bullet$, and B is a direct summand of A as a right B -module.*

Furthermore, if these equivalent statements hold then A is a T -relative projective right B -module.

- 2) *If the lifted canonical map (6.3) is a split epimorphism of left L -modules (with respect to the module structures (6.5)) then the following assertions are equivalent.*

- (a) The canonical map (2.13) is bijective and B is a direct summand of A as a left B -module.
- (b) A is a relative injective right comodule for the R -coring (H, γ_R, π_R) .
- (c) The functor $(\bullet)^{\text{co}\mathcal{H}} : \mathfrak{M}_A^{\mathcal{H}} \rightarrow \mathfrak{M}_B$ is an equivalence, with inverse $\bullet \otimes_B A$, and B is a direct summand of A as a left B -module.

Furthermore, if these equivalent statements hold then A is a T -relative projective left B -module.

Proof. Recall from Section 2.14 that bijectivity of the antipode S in the Hopf algebroid $\mathcal{H}$ implies that also the entwining map (2.12) is bijective.

(a) $\Rightarrow$ (b) By [BB1, Proposition 4.1] (see also [SS, Remark 4.2]), A is a relative injective right comodule for the R -coring (H, γ_R, π_R) .

(b) $\Rightarrow$ (a) The lifted canonical map (6.3) is a split epimorphism in $\mathfrak{M}_A^{\mathcal{H}}$ by Lemma 6.6. By considerations in Section 2.5, coinvariants of the right comodules A and $A \otimes_T A$ (latter one defined via the second tensorand) for the A -coring $A \otimes_R H$ coincide with the $\mathcal{H}$ -coinvariants in A and $A \otimes_T A$, respectively. By Theorem 5.1 (f) $\Rightarrow$ (k), A (with coaction (2.26)) is a relative injective left comodule for the R -coring (H, γ_R, π_R) . So that, by Proposition 4.4 (b) $\Rightarrow$ (a), A is a relative injective left comodule for the A -coring $A \otimes_R H$. Taking Remark 6.5 into account, it follows that $(A \otimes_T A)^{\text{co}\mathcal{H}} = A \otimes_T B$, hence all assumptions in Theorem 6.1 hold. Therefore the canonical map (2.13) is bijective and A is a T -relative projective right B -module by Theorem 6.1. It follows by Proposition 4.4 that the right regular B -module is a direct summand in A .

(b) $\Rightarrow$ (c) By Proposition 4.4 (b) $\Rightarrow$ (a), assertion (b) in the claim implies that A is a relative injective right comodule for the A -coring $A \otimes_R H$. The A -coring $A \otimes_R H$ possesses a grouplike element $1_A \otimes_R 1_H$, hence the unit of the adjunction $(A \otimes_B \bullet, {}^{\text{co}\mathcal{H}}(\bullet))$ (cf. Section 2.3) is an isomorphism, by a left-right symmetric version of Lemma 4.5 (recall that coinvariants with respect to $\mathcal{H}$ and $A \otimes_R H$ are the same, by arguments in Section 2.5). Let us construct the inverse of the counit,

$$n_M : A \otimes_B^{\text{co}\mathcal{H}} M \rightarrow M, \quad a \otimes m \mapsto am,$$

for $M \in {}^{\mathcal{H}}\mathfrak{M}$. Denote the related left $\mathcal{H}_R$ -, and $\mathcal{H}_L$ -coactions in M by $m \mapsto m^{[-1]} \otimes_R m^{[0]}$ and $m \mapsto m_{[-1]} \otimes_L m_{[0]}$, respectively. The canonical map (2.13) is bijective by part (a). Consider the map

$$(6.7) \quad M \rightarrow A \otimes_B M, \quad m \mapsto \text{can}^{-1}(1_A \otimes_R m^{[-1]})m^{[0]}.$$

By Lemma 6.6 the lifted canonical map (6.3) has a right $\mathcal{H}$ -colinear right A -linear section ζ^T in (6.6). The map (6.7) is equal to the composite of

$$(6.8) \quad M \rightarrow A \otimes_T M, \quad m \mapsto \zeta^T(1_A \otimes_R m^{[-1]})m^{[0]}$$

and the canonical epimorphism $A \otimes_T M \rightarrow A \otimes_B M$. We claim that the range of (6.8) is in $A \otimes_T^{\text{co}\mathcal{H}} M$. The right $\mathcal{H}_R$ -comodule A is relative injective by assumption. So by Theorem 5.1 (f) $\Rightarrow$ (o), the left $\mathcal{H}_L$ -coaction in M has a left $\mathcal{H}$ -colinear retraction. The ‘coinvariants part’ of this retraction yields a k -linear retraction of the inclusion ${}^{\text{co}\mathcal{H}}M \rightarrow {}^{\text{co}\mathcal{H}}(H \otimes_L M) \cong M$. Explicitly, in terms of a right total integral j (cf. Theorem 5.1 (a)), we obtain an idempotent map

$$E_M : M \rightarrow {}^{\text{co}\mathcal{H}}M, \quad m \mapsto j(m^{[-1]})m^{[0]}.$$

Consider the right L -module A with action $al := \pi_R \circ t_L(l)a$, for $l \in L$ and $a \in A$. Take a right total integral j as in part (a) of Theorem 5.1 and introduce a left B -module map

$$P_M : A \otimes_L M \rightarrow M, \quad a \otimes m \mapsto a^{[0]}j(S(a^{[1]})m^{[-1]})m^{[0]}.$$

It is well defined by the right L -linearity of the right $\mathcal{H}_R$ -coaction in A and the left L -linearity of the left $\mathcal{H}_R$ -coaction in M , and module map properties of S and j . Making use of the relative Hopf module structure of M , that is (2.28), one checks that $E_M \circ P_M = P_M$. This means that the

range of P_M is within ${}^{co\mathcal{H}}M$. Since the section (6.6) of the lifted canonical map (6.3) satisfies, for $m \in M$,

$$\zeta^T(1_A \otimes_R m^{[-1]})m^{[0]} = ((A \otimes_T P_M) \circ (\zeta_0^T \otimes_L M))(1_A \otimes_R m_{[-1]} \otimes_L m_{[0]}),$$

the range of (6.8) is in $A \otimes_T {}^{co\mathcal{H}}M$. This implies that the range of (6.7) is in $A \otimes_B {}^{co\mathcal{H}}M$. The proof is completed by showing that the corestriction of (6.7) to a map $\tilde{n}_M : M \rightarrow A \otimes_B {}^{co\mathcal{H}}M$ yields the inverse of n_M . Indeed, since $(A \otimes_R \pi_R) \circ \text{can}(a \otimes_B a') = aa'$, for $a, a' \in A$, and can is bijective, $n_M \circ \tilde{n}_M(m) = \pi_R(m^{[-1]})m^{[0]} = m$, for any $m \in M$. On the other hand, since M is an object in ${}^{\mathcal{H}}\mathfrak{M}$, it follows by (2.28) that $\tilde{n}_M \circ n_M(a \otimes_B m) = (\text{can}^{-1}(1_A \otimes_R S^{-1}(a_{[1]}))a_{[0]})m$, for $a \otimes_B m \in A \otimes_B {}^{co\mathcal{H}}M$. The right A -linearity of can implies that $\text{can}^{-1}(1_A \otimes_R S^{-1}(a_{[1]}))a_{[0]} = a \otimes_B 1_A$, for $a \in A$, which proves $\tilde{n}_M \circ n_M(a \otimes_B m) = a \otimes_B m$.

(c) $\Rightarrow$ (a) The left regular comodule of the A -coring $H \otimes_R A$ (cf. Section 2.14) determines an object in ${}^{\mathcal{H}}\mathfrak{M}$. The counit of the adjunction $(A \otimes_B \bullet, {}^{co\mathcal{H}}(\bullet))$, evaluated at the object $H \otimes_R A$, is the isomorphism

$$n_{H \otimes_R A} : A \otimes_B A \rightarrow H \otimes_R A, \quad a \otimes_B a' \mapsto S^{-1}(a_{[1]}) \otimes_R a_{[0]} a'.$$

The canonical map (2.13) is a composite of isomorphisms, $\text{can} = \psi_R \circ n_{H \otimes_R A}$, where ψ_R is the bijective entwining map (2.12). This proves bijectivity of the canonical map (2.13).

In view of Theorem 5.1 (f) $\Leftrightarrow$ (n), part 2) follows by applying part 1) to the co-opposite Hopf algebroid $\mathcal{H}_{cop}$ and its right comodule algebra A^{op} , with $(\mathcal{H}_R)_{cop}$ -coaction $a \mapsto a_{[0]} \otimes_{R^{op}} S^{-1}(a_{[1]})$. $\square$

Observe that T -relative projectivity and generator properties of the B -module A in Theorem 6.7 are as close to its faithful flatness as it is possible for an arbitrary base algebra T of B . If A is a projective T -module (e.g. $T = k$ is a field) then the equivalent assertions in part 1) or 2) imply that A is a projective right or left B -module. Hence the properties, that B is a direct summand in the right or left B -module A , in 1) (a) and (c) or 2) (a) and (c) imply that A is a faithfully flat right or left B -module, cf. [Row, 2.11.29].

If $\mathcal{H}$ is a coseparable Hopf algebroid (i.e. the underlying L -coring or, equivalently, the underlying R -coring is coseparable, cf. [Bö1, Theorem 3.2]) then the forgetful functor $\mathbb{U}^R : \mathfrak{M}^{\mathcal{H}} \rightarrow \mathfrak{M}_R$ is separable (cf. [Brz1, Corollary 3.6]). This implies that if the coseparability of the Hopf algebroid $\mathcal{H}$ is assumed (not only relative injectivity of the comodule algebra A) then the assumption in Theorem 6.7 1) about splitting of the lifted canonical map (6.3) as a right L -module map can be replaced by its splitting as a right R -module map. Latter assumption holds in particular if the canonical map (2.13) is surjective and H is projective as a right R -module (via the source map of the constituent right bialgebroid). Indeed, under this assumption, $A \otimes_R H \cong H \otimes_R A$ is a projective right A -module. So the surjectivity of the right A -module map (2.13) implies that its lifted version (6.3) is a retraction of right A -modules, and hence of right R -modules. By the separability of $\mathbb{U}^R$ it follows then that (6.3) is a retraction of right $\mathcal{H}$ -comodules and the proof can be completed as in Theorem 6.7. Thus we obtain the following.

THEOREM 6.8. *Let $\mathcal{H}$ be a coseparable Hopf algebroid with constituent left bialgebroid $(H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $(H, R, s_R, t_R, \gamma_R, \pi_R)$, and a bijective antipode S . Let A be a right comodule algebra and put $B := A^{co\mathcal{H}}$. Assume that the canonical map (2.13) is surjective.*

- 1) *If H is a projective right R -module (via s_R) then assertions (a), (b) and (c) in Theorem 6.7 1) are equivalent. Furthermore, if they hold then A is a k -relative projective right B -module.*
- 2) *If H is a projective left R -module (via t_R) then assertions (a), (b) and (c) in Theorem 6.7 2) are equivalent. Furthermore, if they hold then A is a k -relative projective left B -module.*

7. EQUIVARIANT INJECTIVITY AND PROJECTIVITY

The notion of equivariant projectivity of a Hopf Galois extension was introduced in the papers [DGH] and [HM]. Equivariant projectivity of a Hopf Galois extension is a crucial property from the non-commutative geometric point of view, as it turns out to be equivalent to the existence of

a strong connection – a non-commutative formulation of local triviality of a principal bundle (see [H]). In the context of Galois extensions $B \subseteq A$ by corings (or bialgebroids or Hopf algebroids), equivariant projectivity *relative* to some subalgebra of B was shown to be equivalent to the existence of more general strong connections in the paper [BB1].

In this section we look for conditions on a Galois extension by a Hopf algebroid under which it obeys (relative) equivariant injectivity and projectivity properties. Recall that having a Hopf algebra H over a commutative ring k and a right H -comodule algebra A , which is a relative injective right H -comodule, A was shown to be a $B(= A^{coH})$ -equivariantly injective H -comodule in [SS, Theorem 5.6]. (This result is extended to algebra extensions by Hopf algebroids in Theorem 7.2 below.) What is more, using the proven B -equivariant injectivity of a relative injective H -comodule algebra A , it was also shown in [SS, Theorem 5.6] that the B -module A is H -equivariantly projective if and only if it is k -relative projective. If A is a relative injective right comodule algebra of a Hopf algebra H with a bijective antipode, with coinvariants B , and the lifted canonical map (1.3) is a split epimorphism of k -modules, then A is an H -Galois extension of B and the B -module A is relative projective (cf. Theorem 6.7). Hence the B -module A is also H -equivariantly projective by the quoted result in [SS, Theorem 5.6]. A most naive generalisation of this result to Hopf algebroid Galois extensions seems not to hold. The reason is that – if working with a Hopf algebroid $\mathcal{H}$ over different non-commutative base algebras L and R – relative projectivity of the B -module A is not enough to prove its (relative) $\mathcal{H}$ -equivariant projectivity. One needs more: (relative) L -equivariant projectivity (see Theorem 7.3 below). As a matter of fact, for a relative injective right comodule algebra A of a Hopf algebroid with a bijective antipode, with coinvariants B , we were not able to deduce relative L -equivariant projectivity of the B -module A from the splitting of the lifted canonical map (6.3) as a left, or right L -module map, as assumed in Theorem 6.7. We needed a stronger assumption: splitting of the lifted canonical map (6.3) as an L - L bimodule map (see Proposition 7.4 below).

DEFINITION 7.1. Let $\mathcal{D}$ be an L -coring and B a T -ring. A left B -module and right $\mathcal{D}$ -comodule V , with left B -linear right $\mathcal{D}$ -coaction, is called a *T -relative $\mathcal{D}$ -equivariantly projective* left B -module if the left action $B \otimes_T V \rightarrow V$ is an epimorphism split by a left B -module right $\mathcal{D}$ -comodule map. We call V a *B -equivariantly injective* right $\mathcal{D}$ -comodule if the right coaction $V \rightarrow V \otimes_L \mathcal{D}$ is a monomorphism split by a left B -module right $\mathcal{D}$ -comodule map.

Analogous notions for right B -modules and left $\mathcal{D}$ -comodules, with a right B -linear left $\mathcal{D}$ -coaction, are defined symmetrically.

Considering an algebra L as a trivial L -coring $\mathcal{L}$, a T -relative $\mathcal{L}$ -equivariantly projective left B -module V is called simply *T -relative L -equivariantly projective*. Clearly, for an L -coring $\mathcal{D}$ and a T -ring B , a T -relative $\mathcal{D}$ -equivariantly projective left B -module V is necessarily T -relative L -equivariantly projective.

For a (left or right) bialgebroid $\mathcal{S}$ over an algebra L , and a T -ring B , a T -relative $\mathcal{S}$ -equivariantly projective B -module means a B -module which is T -relative equivariantly projective for the L -coring underlying $\mathcal{S}$. Consider a Hopf algebroid $\mathcal{H}$ with constituent left bialgebroid $\mathcal{H}_L$ and right bialgebroid $\mathcal{H}_R$, and a left B -module and right $\mathcal{H}$ -comodule V . In view of considerations in Section 2.10, the right $\mathcal{H}_L$ -coaction in V is left B -linear if and only if the right $\mathcal{H}_R$ -coaction is left B -linear. By the same token, V is a T -relative $\mathcal{H}_L$ -equivariantly projective left B -module if and only if it is a T -relative $\mathcal{H}_R$ -equivariantly projective left B -module. In this situation we call the B -module V simply *T -relative $\mathcal{H}$ -equivariantly projective*.

The following two theorems extend [SS, Theorem 5.6] to non-commutative base algebras.

THEOREM 7.2. *Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$, and antipode S . Let A be a right $\mathcal{H}$ -comodule algebra which is a relative injective right $\mathcal{H}_L$ -comodule. Denoting $B := A^{co\mathcal{H}}$, A is a B -equivariantly injective right $\mathcal{H}_L$ -comodule.*

Proof. Using a method in [SS, Lemma 4.1], one constructs a left B -linear right $\mathcal{H}$ -colinear retraction ϕ of the $\mathcal{H}_L$ -coaction $a \mapsto a_{[0]} \otimes_L a_{[1]}$ in A , in terms of an $\mathcal{H}$ -colinear retraction ν . Explicitly,

$$\phi : A \otimes_L H \rightarrow A, \quad a \otimes_L h \mapsto a^{[0]} \nu \left(1_A \otimes_L S(a^{[1]})h \right),$$

where $a \mapsto a^{[0]} \otimes_R a^{[1]}$ denotes the right $\mathcal{H}_R$ -coaction in A , related to the $\mathcal{H}_L$ -coaction via (2.17). Note that since ν is $\mathcal{H}$ -colinear, it is left R -linear and hence the map ϕ is well defined. $\square$

THEOREM 7.3. *Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$, and antipode S . Let A be a right $\mathcal{H}$ -comodule algebra and $B := A^{co\mathcal{H}}$. Assume that A is a B -equivariantly injective right $\mathcal{H}_L$ -comodule. If B is a T -ring (e.g. T is some k -subalgebra of B) then A is a T -relative $\mathcal{H}$ -equivariantly projective left B -module if and only if it is a T -relative L -equivariantly projective left B -module.*

Proof. If A is a T -relative $\mathcal{H}$ -equivariantly projective left B -module then it is obviously T -relative L -equivariantly projective. In order to see the converse implication, take a B - L bimodule section χ_0^T of the multiplication map $B \otimes_T A \rightarrow A$ and a left B -linear and right $\mathcal{H}$ -colinear retraction ϕ of the right $\mathcal{H}_L$ -coaction $\tau_A : A \rightarrow A \otimes_L H$ in A . Consider the left B -linear and right $\mathcal{H}$ -colinear map

$$\chi^T := (B \otimes_T \phi) \circ (\chi_0^T \otimes_L H) \circ \tau_A : A \rightarrow B \otimes_T A.$$

It follows by the left B -linearity of ϕ that χ^T is a section of the multiplication map $B \otimes_T A \rightarrow A$. $\square$

The message of Theorem 7.2 and Theorem 7.3 is to look for situations, in which the T -relative L -equivariant projectivity condition in Theorem 7.3 holds, for a relative injective right comodule algebra of a constituent left bialgebroid in a Hopf algebroid.

We have seen in Section 2.14 that if the antipode of a Hopf algebroid $\mathcal{H}$ (over base algebras L and R) is bijective then a right comodule algebra A has a canonical left $\mathcal{H}^{op}$ -comodule algebra structure, with coactions (2.26) and (2.27). Recall that the coaction (2.27) corresponds to the left L -module structure of A , which is related to its right R -module structure via

$$(7.1) \quad la = a\pi_R \circ s_L(l), \quad \text{for } l \in L, a \in A.$$

Since the right actions in A by R and $B := A^{co\mathcal{H}}$ commute (cf. Section 2.7), A is an L - B bimodule via the left L -action (7.1) and the obvious right B -action. The following proposition concerns equivariant projectivity of this L - B bimodule A .

PROPOSITION 7.4. *Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$, and a bijective antipode S . Let A be a right $\mathcal{H}$ -comodule algebra which is a relative injective right $\mathcal{H}_L$ -comodule. Set $B := A^{co\mathcal{H}}$, and assume that B is a T -ring (e.g. T is a k -subalgebra of B). Assume that the lifted canonical map (6.3) is a split epimorphism of L - L bimodules (with respect to the module structures (6.4) and (6.5)). Then A is a T -relative L -equivariantly projective right B -module.*

Proof. Let ζ_0^T be an L - L bimodule section of the lifted canonical map (6.3). By Lemma 6.6, the map (6.3) is split by the right $\mathcal{H}$ -comodule right A -module map ζ^T in (6.6). From the proof of implication (b) $\Rightarrow$ (a) in Theorem 6.7 we have that $(A \otimes_T A)^{co\mathcal{H}} = A \otimes_T B$. Hence taking the ‘coinvariants part’ of ζ^T , we obtain a right B -module section of the multiplication map $A \otimes_T B \rightarrow A$,

$$\chi_0^T : A \rightarrow A \otimes_T B, \quad a \mapsto \zeta_0^T(1_A \otimes_R S^{-1}(a_{[1]})(1))^{[0]} j \left(S(\zeta_0^T(1_A \otimes_R S^{-1}(a_{[1]})(1))^{[1]}) S^{-1}(a_{[1]})(2) \right) a_{[0]}.$$

Consider the left L -module structure (7.1) of A . The right $\mathcal{H}_L$ -coaction in A is left L -linear in the sense that $(a\pi_R \circ s_L(l))_{[0]} \otimes_L (a\pi_R \circ s_L(l))_{[1]} = a_{[0]} \otimes_L a_{[1]} s_R \circ \pi_R \circ s_L(l)$, for $l \in L$ and $a \in A$. The antipode satisfies $S^{-1}(hs_R \circ \pi_R \circ s_L(l)) = t_R \circ \pi_R \circ s_L(l) S^{-1}(h) = s_L(l) S^{-1}(h)$, for $l \in L$ and $h \in H$. The coproduct γ_L is left L -linear, i.e. $(s_L(l)h)_{(1)} \otimes_L (s_L(l)h)_{(2)} = s_L(l)h_{(1)} \otimes_L h_{(2)}$, for $l \in L$ and $h \in H$. The map ζ_0^T is left L -linear by assumption, with respect to the left L -module structures in (6.5). The right $\mathcal{H}_R$ -coaction in $A \otimes_T A$ is given via the second factor, so it is obviously left L -linear with respect to the left L -module structure in (6.5). All these considerations together verify the left L -linearity of χ_0^T with respect to the left L -module structure (7.1) in A . Hence χ_0^T is an L - B

bilinear section of the multiplication map $A \otimes_T B \rightarrow A$, which proves T -relative L -equivariant projectivity of the right B -module A . $\square$

Let A be a right comodule algebra of a Hopf algebroid $\mathcal{H}$ with a bijective antipode S . By Theorem 5.1 (c) $\Leftrightarrow$ (n), A is a relative injective right comodule for the constituent left bialgebroid $\mathcal{H}_L$ if and only if A^{op} is a relative injective right comodule for the left bialgebroid $(\mathcal{H}_L)_{cop}$, with coaction $a \mapsto a^{[0]} \otimes_{L^{op}} S^{-1}(a^{[1]})$. Hence Proposition 7.4 can be applied to the right $\mathcal{H}_{cop}$ -comodule algebra A^{op} , with $(\mathcal{H}_L)_{cop}$ coaction $a \mapsto a^{[0]} \otimes_{L^{op}} S^{-1}(a^{[1]})$. It yields a result about the equivariant projectivity of A as a B - L bimodule, with obvious left B -action, and right L -action related to the left R -action via

$$al = \pi_R \circ t_L(l)a, \quad \text{for } l \in L, a \in A.$$

COROLLARY 7.5. *In the setting of Proposition 7.4, A is a T -relative L -equivariantly projective left B -module.*

The following corollary is the main result of this section. It formulates sufficient conditions on a Galois extension $B \subseteq A$ by a Hopf algebroid $\mathcal{H}$ with a bijective antipode, under which A is a T -relative $\mathcal{H}$ -equivariantly projective left and right B -module, for an algebra T such that B is a T -ring.

COROLLARY 7.6. *Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$, and a bijective antipode S . Let A be a right $\mathcal{H}$ -comodule algebra which is a relative injective right $\mathcal{H}_L$ -comodule. Denote $B := A^{co\mathcal{H}}$ and assume that B is a T -ring (e.g. T is a subalgebra of B). Assume that the lifted canonical map (6.3) is a split epimorphism of L - L bimodules (with respect to the module structures (6.4) and (6.5)). Then $B \subseteq A$ is an $\mathcal{H}_R$ -Galois extension and A is a T -relative $\mathcal{H}$ -equivariantly projective left and right B -module with respect to the left and right $\mathcal{H}$ -comodule structures related via (2.26) and (2.27).*

Proof. By Theorem 5.1 (c) $\Rightarrow$ (f), A is a relative injective right $\mathcal{H}_R$ -comodule. Hence the Galois property, i.e. bijectivity of the canonical map (2.13), follows by virtue of Theorem 6.7 (b) $\Rightarrow$ (a).

A is a B -equivariantly injective right, and left $\mathcal{H}_L$ -comodule by Theorem 7.2, and its application to the co-opposite Hopf algebroid $\mathcal{H}_{cop}$ and the right $\mathcal{H}_{cop}$ -comodule algebra A^{op} (with $(\mathcal{H}_R)_{cop}$ -coaction $a \mapsto a_{[0]} \otimes_{R^{op}} S^{-1}(a_{[1]})$), respectively. By Proposition 7.4, A is a T -relative L -equivariantly projective right B -module. By Corollary 7.5, A is a T -relative L -equivariantly projective left B -module. Hence A is a T -relative $\mathcal{H}$ -equivariantly projective left and right B -module by Theorem 7.3, and its application to the co-opposite Hopf algebroid $\mathcal{H}_{cop}$ and the right $\mathcal{H}_{cop}$ -comodule algebra A^{op} (with $(\mathcal{H}_R)_{cop}$ -coaction $a \mapsto a_{[0]} \otimes_{R^{op}} S^{-1}(a_{[1]})$), respectively. $\square$

By [BB1, Theorem 3.7] we conclude that there exists a strong T -connection for an extension $B \subseteq A$ as in Corollary 7.6 whenever T is a k -subalgebra of B . In [BB1, Theorem 5.14] conditions are formulated for the independence of the corresponding relative Chern-Galois character of the choice of a strong T -connection. Note that in the case when in Corollary 7.6 the k -algebra T is equal to k , these conditions reduce to the assumption that A is a locally projective k -module.

EXAMPLE 7.7. *Cleft extensions by Hopf algebroids were introduced in [BB2, Definition 3.5], as follows. Let $\mathcal{H}$ be a Hopf algebroid with constituent left bialgebroid $\mathcal{H}_L = (H, L, s_L, t_L, \gamma_L, \pi_L)$, right bialgebroid $\mathcal{H}_R = (H, R, s_R, t_R, \gamma_R, \pi_R)$, and antipode S . Let A be a right $\mathcal{H}$ -comodule algebra with coinvariants $B := A^{co\mathcal{H}}$. Denote the unit map of the corresponding R -ring A by $\eta_R : R \rightarrow A$. The algebra extension $B \subseteq A$ is called $\mathcal{H}$ -cleft provided that the following conditions hold.*

- a) A is an L -ring (with some unit map $\eta_L : L \rightarrow A$) and B is an L -subring of A .
- b) There exist morphisms $j \in {}_L\text{Hom}^{\mathcal{H}}(H, A)$ and $\tilde{j} \in {}_R\text{Hom}_L(H, A)$, satisfying

$$\mu \circ (j \otimes_L j) \circ \gamma_L = \eta_R \circ \pi_R \quad \text{and} \quad \mu \circ (j \otimes_R \tilde{j}) \circ \gamma_R = \eta_L \circ \pi_L,$$

where μ denotes the multiplication in A , both as an L -ring and as an R -ring. The bimodule structures in H are given by

$$lhr := s_L(l)hs_R(r) \quad \text{and} \quad rhl = t_L(l)ht_R(r), \quad \text{for } l \in L, r \in R, h \in H.$$

The bimodule structures in A are given by

$$lar := \eta_L(l)a\eta_R(r) \quad \text{and} \quad ral = \eta_R(r)a\eta_L(l), \quad \text{for } l \in L, r \in R, a \in A.$$

In an $\mathcal{H}$ -cleft extension $B \subseteq A$ the map $\tilde{j}(1_H)j(-) : H \rightarrow A$ is right $\mathcal{H}$ -colinear and normalised. Hence, by Theorem 5.1 (a) $\Rightarrow$ (c), A is a relative injective right $\mathcal{H}_L$ -comodule. By definition B is an L -ring. The lifted canonical map

$$\widetilde{\text{can}}^L : A \otimes_L A \rightarrow A \otimes_R H, \quad a \otimes_L a' \mapsto aa'^{[0]} \otimes_R a'^{[1]}$$

possesses an L - L bilinear section (with respect to the module structures (6.4) and (6.5)):

$$(7.2) \quad \zeta_0^L : A \otimes_R H \rightarrow A \otimes_L A, \quad a \otimes_R h \mapsto a\tilde{j}(h_{(1)}) \otimes_L j(h_{(2)}).$$

The map (7.2) is well defined by the module map properties of j and $\tilde{j}$. It is left L -linear by the identity $\tilde{j}(t_R(r)h) = \tilde{j}(h)\eta_R(r)$, for $r \in R$ and $h \in H$, see [BB2, Lemma 3.7]. Right L -linearity of (7.2) follows by the left R -linearity of (the right $\mathcal{H}$ -comodule map) j , i.e. $j(s_R(r)h) = \eta_R(r)j(h)$, for $r \in R$ and $h \in H$. In view of Corollary 7.6, all these considerations together imply that a cleft extension $B \subseteq A$ by a Hopf algebroid $\mathcal{H}$ with a bijective antipode is an $\mathcal{H}$ -Galois extension which is an L -relative $\mathcal{H}$ -equivariantly projective left and right B -module, cf. [BB2, Lemma 5.1].

ACKNOWLEDGEMENTS

This paper was written while A. Ardizzoni and C. Menini were members of G.N.S.A.G.A. with partial financial support from M.I.U.R.. Work of G. Böhm is supported by the Hungarian Scientific Research Fund OTKA T043159 and the Bolyai János Fellowship. Her stay, as a visiting professor at University of Ferrara, was supported by I.N.D.A.M.. She would like to express her gratitude to the members of the Department of Mathematics at University of Ferrara for a very warm hospitality.

REFERENCES

- [Ar] A. Ardizzoni, *Separable Functors and formal smoothness*, Preprint arXiv:math.QA/0407095.
- [Bö1] G. Böhm, *Integral theory for Hopf algebroids*, Alg. Rep. Theory **8** (2005), 563–599.
- [Bö2] G. Böhm, *Galois theory for Hopf algebroids*, Ann. Univ. Ferrara - Sez. VII - Sc. Mat., Vol **LI** (2005), 233–262.
- [BB1] G. Böhm and T. Brzeziński, *Strong connections and the relative Chern-Galois character for corings*, Int. Math. Res. Not. 2005, **42** (2005), 2579–2625.
- [BB2] G. Böhm and T. Brzeziński, *Cleft extensions of Hopf algebroids*, Appl. Categorical Structures **14** (2006), 431–469.
- [BSz] G. Böhm and K. Szlachányi, *Hopf algebroids with bijective antipodes: axioms, integrals and duals*, J. Algebra **274** (2004), 708–750.
- [Bro] K. S. Brown, *Cohomology of groups*. Graduate Texts in Mathematics, **87**. Springer-Verlag, New York-Berlin, 1982.
- [Brz1] T. Brzeziński, *The structure of corings. Induction functors, Maschke-type theorem, and Frobenius and Galois-type properties*, Alg. Rep. Theory **5** (2002), 389–410.
- [Brz2] T. Brzeziński, *Galois comodules*, J. Algebra **290** (2005), 503–537.
- [Brz3] T. Brzeziński, *A note on coring extensions*, Ann. Univ. Ferrara - Sez. VII - Sc. Mat., Vol **LI** (2005), 15–27.
- [BM] T. Brzeziński and G. Militaru, *Bialgebroids, $\times_R$ -bialgebras and duality*, J. Algebra **251** (2002), 279–294.
- [BTW] T. Brzeziński, R.B. Turner and A.P. Wrightson, *The structure of weak coalgebra-Galois extensions*, Comm. Algebra **34** (2006), 1489–1509.
- [BW] T. Brzeziński and R. Wisbauer, *Corings and Comodules*. Cambridge University Press, Cambridge, 2003.
- [CM] S. Caenepeel and G. Militaru, *Maschke functors, semisimple functors and separable functors of the second kind. Applications*, J. Pure Appl. Algebra **178** (2003), 131–157.
- [CIMZ] S. Caenepeel, B. Ion, G. Militaru and S. Zhu, *Separable functors applied to Doi Hopf modules. Applications*, Adv. Math. **145** (1999), 239–290.
- [CMZ] S. Caenepeel, G. Militaru and Shenglin Zhu, *Frobenius Separable Functors for Generalized Module Categories and Nonlinear Equations*. LNM **1787** (2002), Springer-Verlag, Berlin - New York.
- [DGH] L. Dabrowski, H. Grosse and P.M. Hajac, *Strong connections and Chern-Connes pairing in the Hopf-Galois theory*, Comm. Math. Phys. **220** (2001), 301–331.
- [Doi] Y. Doi, *Algebras with total integrals*, Comm. Algebra **13** (1985), 2137–2159.
- [H] P.M. Hajac, *Strong connections on quantum principal bundles*, Comm. Math. Phys. **182** (1996), 579–617.
- [HM] P.M. Hajac and S. Majid, *Projective module description of the q -monopole*, Comm. Math. Phys. **206** (1999), 247–264.

- [HS] P.J. Hilton and U. Stambach, *A course in Homological algebra*. Graduate Text in Mathematics **4**, Springer, New York, 1971.
- [KSz] L. Kadison and K. Szlachányi, *Bialgebroid actions on depth two extensions and duality*, Adv. Math. **179** (2003), 75–121.
- [KT] H. F. Kreimer and M. Takeuchi, *Hopf algebras and Galois extensions of an algebra*, Indiana Univ. Math. J. **30** (1981), 675–692.
- [Lu] J.-H. Lu, *Hopf algebroids and quantum groupoids*, Int. J. Math. **7** (1996), 47–70.
- [MM1] C. Menini and G. Militaru, *Integrals, quantum Galois extensions and the affineness criterion for quantum Yetter-Drinfel'd modules*, J. Algebra **247** (2002), 467–508.
- [MM2] C. Menini and G. Militaru, *The affineness criterion for Doi-Koppinen modules*, in: "Hopf algebras in non-commutative geometry and physics", S. Caenepeel and F. Van Oystaeyen (eds.), Lecture Notes in Pure and Applied Mathematics **239** pp 215–228, Marcel Dekker, New York, 2004.
- [NVV] C. Năstăsescu, M. Van den Bergh and F. Van Oystaeyen, *Separable functors applied to graded rings*, J. Algebra **123** (1989), 397–413.
- [Raf] M. D. Rafael, *Separable Functors Revisited*, Comm. Algebra **18** (1990), 1445–1459.
- [Row] L.H. Rowen, *Ring theory. Vol. I*, Academic Press, Boston (1988).
- [Scha1] P. Schauenburg, *Hopf-Galois and bi-Galois extensions*, in: Galois theory, Hopf algebras, and semiabelian categories, Fields Inst. Commun. **43**, AMS 2004, pp. 469–515.
- [Scha2] P. Schauenburg, *Bialgebras over noncommutative rings and a structure theorem for Hopf bimodules*, Appl. Categorical Structures **6** (1998), 193–22.
- [SS] P. Schauenburg and H.-J. Schneider, *On generalized Hopf Galois extensions*, Journal of Pure and Applied Algebra **202** (2005), 168–194.
- [Schn] H.-J. Schneider, *Principal homogeneous spaces for arbitrary Hopf algebras*, Israel J. Math. **72** (1990), 167–195.
- [Ta] M. Takeuchi, *Groups of algebras over $A \otimes \bar{A}$* , J. Math. Soc. Japan **29** (1977), 459–492.
- [We] C. Weibel, *An introduction to homological algebra*. Cambridge Studies in Advanced Mathematics **38**, Cambridge University Press, 1994.

UNIVERSITY OF FERRARA, DEPARTMENT OF MATHEMATICS, VIA MACHIAVELLI 35, FERRARA, I-44100, ITALY

E-mail address: rdzlsn@unife.it

URL: <http://www.unife.it/utenti/alessandro.ardizzoni>

RESEARCH INSTITUTE FOR PARTICLE AND NUCLEAR PHYSICS, BUDAPEST, H-1525 BUDAPEST 114, P.O.B.49, HUNGARY

E-mail address: G.Bohm@rmki.kfki.hu

URL: <http://www.rmki.kfki.hu/~bgabr>

UNIVERSITY OF FERRARA, DEPARTMENT OF MATHEMATICS, VIA MACHIAVELLI 35, FERRARA, I-44100, ITALY

E-mail address: men@unife.it

URL: <http://www.unife.it/utenti/claudia.menini>