# INTEGRAL GROUP RING OF THE MATHIEU SIMPLE GROUP $M_{23}$

V.A. BOVDI AND A.B. KONOVALOV

Institute of Mathematics, University of Debrecen, Hungary School of Computer Science, University of St Andrews, Scotland

ABSTRACT. We investigate the classical Zassenhaus conjecture for the unit group of the integral group ring of Mathieu simple group  $M_{23}$  using the Luthar-Passi method. This work is a continuation of the research that we carried out for Mathieu groups  $M_{11}$  and  $M_{12}$ . As a consequence, for this group we confirm Kimmerle's conjecture on prime graphs.

### 1. INTRODUCTION AND MAIN RESULTS

Let  $V(\mathbb{Z}G)$  be the normalized unit group of the integral group ring  $\mathbb{Z}G$  of a finite group G. With respect to the structure of  $V(\mathbb{Z}G)$ , H. Zassenhaus in [Zas74] proposed the following conjecture:

(**ZC**) Every torsion unit  $u \in V(\mathbb{Z}G)$  is conjugate within the rational group algebra  $\mathbb{Q}G$  to an element of G.

In the general case the problem is still open, despite a lot of known results about (**ZC**) and methods developed for its investigation. In the present paper we will be focused on those of them which are relevant to finite simple groups.

The first result on (**ZC**) for simple groups was obtained by I.S. Luthar and I.B.S. Passi in [LP89] for the alternating group  $A_5$ , and the technique used in this paper is now known as the Luthar-Passi method. Later M. Hertweck in [Her07a] extended this method and applied it for the investigation of (**ZC**) for  $PSL(2, p^n)$ . This method proved to be useful for non-simple groups as well. For some recent results we refer to [BJK07, BHK04, BK07, Her07a, Her06, Her07b, HK06]. Also some related properties can be found in [AB89, LT91] and [BK00, Kim06]. In the latter papers some weakened variations of (**ZC**) have been made.

In [Kim06] W. Kimmerle proposed to relate (**ZC**) with some properties of graphs associated with groups. Let #(G) be the set of all primes dividing the order of G. The Gruenberg-Kegel graph (or the prime graph) of G is the graph  $\pi(G)$  with

<sup>1991</sup> Mathematics Subject Classification. Primary 16S34, 20C05; Secondary 20D08.

Key words and phrases. Zassenhaus conjecture, Kimmerle conjecture, torsion unit, partial augmentation, integral group ring.

The research was supported by OTKA grants No.T 037202, No.T 038059 and Francqui Stichting (Belgium) grant ADSI107.

V.A. Bovdi: Institute of Mathematics, University of Debrecen, P.O. Box 12, H-4010 Debrecen, Hungary; Institute of Mathematics and Informatics, College of Nyíregyháza, Sóstói út 31/b, H-4410 Nyíregyháza, Hungary; E-mail: vbovdi@math.klte.hu.

A.B. Konovalov: School of Computer Science, University of St Andrews, Jack Cole Building, North Haugh, St Andrews, Fife, KY16 9SX, Scotland; E-mail: konovalov@member.ams.org.

#### BOVDI AND KONOVALOV

vertices labeled by the primes in #(G) and with an edge from p to q if there is an element of order pq in the group G. The following conjecture was introduced by W. Kimmerle:

**(KC)** If G is a finite group then 
$$\pi(G) = \pi(V(\mathbb{Z}G))$$
.

In [Kim06] (**KC**) was solved for finite solvable groups and finite Frobenius groups. Note that with respect to the Zassenhaus conjecture the investigation of Frobenius groups was completed by M. Hertweck and the first author in [BH07]. In our previous papers [BJK07, BK07, BKS07] we confirmed (**KC**) for sporadic simple groups  $M_{11}$ ,  $M_{12}$  and some Janko simple groups.

In the present paper we continue these investigations for the Mathieu simple group  $M_{23}$  using the Luthar-Passi method. In our main result we obtain a lot of information on possible torsion units in  $V(\mathbb{Z}M_{23})$ , and as a consequence of part (i) of the main theorem we solve **(KC)** for  $M_{23}$ .

First we introduce some notation. Put  $G = M_{23}$ . Let

$$\mathcal{C} = \{C_1, C_{2a}, C_{3a}, C_{4a}, C_{5a}, C_{6a}, C_{7a}, C_{7b}, C_{8a}, \\ C_{11a}, C_{11b}, C_{14a}, C_{14b}, C_{15a}, C_{15b}, C_{23a}, C_{23b}\},\$$

be the collection of all conjugacy classes of G, where the first index denotes the order of the elements of this conjugacy class and  $C_1 = \{1\}$ . Suppose  $u = \sum \alpha_g g \in V(\mathbb{Z}G)$ has finite order k. Denote by  $\nu_{nt} = \nu_{nt}(u) = \varepsilon_{C_{nt}}(u) = \sum_{g \in C_{nt}} \alpha_g$  the partial augmentation of u with respect to  $C_{nt}$ . From S.D. Berman's Theorem [Ber55] one knows that  $\nu_1 = \alpha_1 = 0$  and

(1) 
$$\sum_{C_{nt}\in\mathcal{C}}\nu_{nt}=1.$$

Hence, for any character  $\chi$  of G, we get that  $\chi(u) = \sum \nu_{nt} \chi(h_{nt})$ , where  $h_{nt}$  is a representative of a conjugacy class  $C_{nt}$ .

The main result is the following.

**Theorem 1.** Let  $V(\mathbb{Z}G)$  be the normalized unit group of the integral group ring  $\mathbb{Z}G$ , where G is the sporadic simple Mathieu group  $M_{23}$ . Let u be a torsion unit of  $V(\mathbb{Z}G)$  of order |u|. Then:

- (i) if |u| ∉ {12, 24, 28, 56}, then |u| coincides with the order of some element g ∈ G. Equivalently, there is no elements of orders 10, 21, 22, 33, 35, 46, 55, 69, 77, 115, 161 and 253 in V(ZG).
- (ii) if  $|u| \in \{2, 3, 5, 23\}$ , then u is rationally conjugate to some  $g \in G$ ;
- (iii) if |u| = 4, then the tuple of partial augmentations of u belongs to the set

 $\Big\{ (\nu_{2a}, \nu_{3a}, \nu_{4a}, \nu_{5a}, \nu_{6a}, \nu_{7a}, \nu_{7b}, \nu_{8a}, \nu_{11a}, \nu_{11b}, \nu_{14a}, \nu_{14b}, \nu_{15a},$ 

$$\nu_{15b}, \nu_{23a}, \nu_{23b}) \in \mathbb{Z}^{16} \quad | \quad \nu_{kx} = 0,$$
  
$$kx \notin \{2a, 4a\}, \quad (\nu_{2a}, \nu_{4a}) \in \{ (0, 1), (-2, 3), (2, -1) \} \};$$

(iv) if |u| = 6, then the tuple of partial augmentations of u belongs to the set

$$\left\{ \begin{array}{ll} \left(\nu_{2a}, \nu_{3a}, \nu_{4a}, \nu_{5a}, \nu_{6a}, \nu_{7a}, \nu_{7b}, \nu_{8a}, \nu_{11a}, \nu_{11b}, \nu_{14a}, \nu_{14b}, \nu_{15a}, \\ \nu_{15b}, \nu_{23a}, \nu_{23b} \right) \in \mathbb{Z}^{16} \quad | \nu_{kx} = 0, \quad kx \notin \{2a, 3a, 6a\}, \\ \left(\nu_{2a}, \nu_{3a}, \nu_{6a}\right) \in \{ (0, 0, 1), \quad (4, -6, 3), \quad (-6, 6, 1), \quad (-6, 9, -2), \\ \left(-6, 12, -5\right), \quad (-4, 3, 2), \quad (-4, 6, -1), \quad (-4, 9, -4), \quad (-2, 0, 3), \\ \left(-2, 3, 0\right), \quad (-2, 6, -3), \quad (0, -3, 4), \quad (0, 3, -2), \quad (2, -6, 5), \\ \left(2, -3, 2\right), \quad \left(2, 0, -1\right), \quad (4, -9, 6), \quad (4, -3, 0), \quad (6, -12, 7), \\ \left(6, -9, 4\right), \quad \left(6, -6, 1\right) \right\} \right\};$$

- (v) if |u| = 7, then the tuple of partial augmentations of u belongs to the set
  - $\left\{ \begin{array}{l} (\nu_{2a}, \nu_{3a}, \nu_{4a}, \nu_{5a}, \nu_{6a}, \nu_{7a}, \nu_{7b}, \nu_{8a}, \nu_{11a}, \nu_{11b}, \nu_{14a}, \nu_{14b}, \nu_{15a}, \\ \nu_{15b}, \nu_{23a}, \nu_{23b}) \in \mathbb{Z}^{16} \quad | \nu_{kx} = 0, \\ kx \notin \{7a, 7b\}, \quad (\nu_{7a}, \nu_{7b}) \in \{ (0, 1), (2, -1), (1, 0), (-1, 2) \} \right\};$
- (vi) if |u| = 8, then the tuple of partial augmentations of u belongs to the set
  - $\left\{ \begin{array}{c} (\nu_{2a}, \nu_{3a}, \nu_{4a}, \nu_{5a}, \nu_{6a}, \nu_{7a}, \nu_{7b}, \nu_{8a}, \nu_{11a}, \nu_{11b}, \nu_{14a}, \nu_{14b}, \nu_{15a}, \end{array} \right.$

 $\nu_{15b}, \nu_{23a}, \nu_{23b}) \in \mathbb{Z}^{16} \quad | \nu_{kx} = 0,$  $kx \notin \{2a, 4a, 8a\}, \quad (\nu_{2a}, \nu_{4a}, \nu_{8a}) \in \{ (-2, 4, -1), (0, 4, -3), (2, 0, -1), (0, 0, 1), (0, 2, -1), (2, -2, 1), (-2, 2, 1), (0, -2, 3), (-2, 6, -3), (2, -4, 3)\} \};$ 

(vii) if |u| = 11, then the tuple of partial augmentations of u belongs to the set

 $\left\{ \left(\nu_{2a},\nu_{3a},\nu_{4a},\nu_{5a},\nu_{6a},\nu_{7a},\nu_{7b},\nu_{8a},\nu_{11a},\nu_{11b},\nu_{14a},\nu_{14b},\nu_{15a},\right.\right.$ 

 $\nu_{15b}, \nu_{23a}, \nu_{23b}) \in \mathbb{Z}^{16} | \nu_{kx} = 0,$  $kx \notin \{11a, 11b\}, \quad (\nu_{11a}, \nu_{11b}) \in \{ (-5, 6), (5, -4), (0, 1), (-2, 3), (-8, 9), (-9, 10), (-6, 7), (2, -1), (8, -7), (-3, 4), (6, -5), (-4, 5), (1, 0), (3, -2), (9, -8), (10, -9), (7, -6), (-1, 2), (-7, 8), (4, -3) \} \};$ 

(viii) if |u| = 15, then the tuple of partial augmentations of u belongs to the set

 $\Big\{ (\nu_{2a}, \nu_{3a}, \nu_{4a}, \nu_{5a}, \nu_{6a}, \nu_{7a}, \nu_{7b}, \nu_{8a}, \nu_{11a}, \nu_{11b}, \nu_{14a}, \nu_{14b}, \nu_{15a},$ 

 $\nu_{15b}, \nu_{23a}, \nu_{23b}) \in \mathbb{Z}^{16} \quad | \quad \nu_{kx} = 0,$   $kx \notin \{3a, 5a, 15a, 15b\}, \quad (\nu_{3a}, \nu_{5a}, \nu_{15a}, \nu_{15b}) \in \{ (-3, 5, -1, 0),$  $(-3, 5, 0, -1), (0, 0, 0, 1), (0, 0, 1, 0), (3, -5, 1, 2), (3, -5, 2, 1) \} \}.$ 

As an immediate consequence of the first part of the Theorem one obtains Kimmerle's conjecture for  $M_{23}$ .

**Corollary 1.** If  $G = M_{23}$  then  $\pi(G) = \pi(V(\mathbb{Z}G))$ .

## 2. Preliminaries

Throughout the paper we simply denote  $M_{23}$  by G. The *p*-Brauer character table of the group G will be denoted by  $\mathfrak{BCT}(p)$ .

The crucial restriction on partial augmentations is given by the next result, detailed explanation of which can be found in [LP89] and [Her07a].

**Proposition 1.** (see [LP89, Her07a]) Let either p = 0 or p is a prime divisor of |G|. Suppose that  $u \in V(\mathbb{Z}G)$  has finite order k and assume k and p are coprime in case  $p \neq 0$ . If z is a primitive k-th root of unity and  $\chi$  is either a classical character or a p-Brauer character of G then, for every integer l, the number

(2) 
$$\mu_l(u,\chi,p) = \frac{1}{k} \sum_{d|k} Tr_{\mathbb{Q}(z^d)/\mathbb{Q}} \{\chi(u^d) z^{-dl}\}$$

is a non-negative integer.

If p = 0, we will use the notation  $\mu_l(u, \chi, *)$  for  $\mu_l(u, \chi, 0)$ .

For the determination of possible orders of units we use the next bound.

**Proposition 2.** (see [CL65]) The order of a torsion element  $u \in V(\mathbb{Z}G)$  is a divisor of the exponent of G.

The next two results, the first of which is a special case of the second one, already yield that several partial augmentations are zero.

**Proposition 3.** (see [LP89] and Theorem 2.7 in [MRSW87]) Let u be a torsion unit of  $V(\mathbb{Z}G)$ . Let C be a conjugacy class of G. If  $a \in C$  and p is a prime dividing the order of a but not the order of u then  $\varepsilon_C(u) = 0$ .

**Proposition 4.** (see [Her06], Proposition 3.1; [Her07a], Proposition 2.2) Let G be a finite group and let u be a torsion unit in  $V(\mathbb{Z}G)$ . If x is an element of G whose p-part, for some prime p, has order strictly greater than the order of the p-part of u, then  $\varepsilon_x(u) = 0$ .

The following result relates the solution of the Zassenhaus conjecture to partial augmentations of torsion units.

**Proposition 5.** (see [LP89] and Theorem 2.5 in [MRSW87]) Let  $u \in V(\mathbb{Z}G)$  be of order k. Then u is conjugate in  $\mathbb{Q}G$  to an element  $g \in G$  if and only if for each d dividing k there is precisely one conjugacy class C with partial augmentation  $\varepsilon_C(u^d) \neq 0$ .

Finally, the next result is useful for the investigation of *p*-elements in  $V(\mathbb{Z}G)$ .

**Proposition 6.** (see [CL65]) Let p be a prime, and let u be a torsion unit of  $V(\mathbb{Z}G)$  of order  $p^n$ . Then for  $m \neq n$  the sum of all partial augmentations of u with respect to conjugacy classes of elements of order  $p^m$  is divisible by p.

### 3. Proof of the Theorem

It is well known [GAP06, Gor83] that  $|G| = 2^7 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 23$  and  $exp(G) = 2^3 \cdot 3 \cdot 5 \cdot 7 \cdot 11 \cdot 23$ . The character table of G, as well as the Brauer character tables  $\mathfrak{BCT}(p)$ , where  $p \in \{2, 3, 5, 7, 11, 23\}$ , can be found using the GAP system [GAP06]. Throughout the paper we will use the notation, indexation inclusive, for the characters and conjugacy classes as used in GAP.

Since the group G possesses elements of orders 2, 3, 4, 5, 6, 7, 8, 11, 14, 15 and 23, first of all we shall investigate units of these orders. After this, by Proposition 2, the order of each torsion unit divides the exponent of G, so it will be enough to consider units of orders 10, 12, 21, 22, 24, 28, 33, 35, 46, 55, 56, 69, 77, 115, 161 and 253, because if u will be a unit of another possible order, then there is  $t \in \mathbb{N}$ 

such that  $u^t$  has an order from this list. For all these orders except 12, 24, 28 and 56 we will prove that units of such order do not appear in  $V(\mathbb{Z}G)$ .

To reduce the volume of the paper, we omit from consideration units of orders 12, 14 and 28, because in these cases we obtain systems with enormous number of solutions. Note that with the LAGUNA package [BKRS07] we can investigate units of these orders and show that there are no units of order 28, and, consequently, of order 56 in  $V(\mathbb{Z}G)$ . Unfortunately, for units of order 12 the Luthar-Passi method is not enough to prove the same result.

Now we consider each case separately:

• Let u be a unit of order 2, 3 or 5. Then using Propositions 3 and 4 we obtain that all partial augmentations except one are zero.

• Let u be a unit of order 4. Then  $\nu_{2a} + \nu_{4a} = 1$  by (1) and Proposition 4. By (2), using  $\mathfrak{BCT}(23)$  we obtain the system of inequalities

$$\begin{aligned} \mu_0(u,\chi_{10},23) &= \frac{1}{4}(-16\nu_{2a}+272) \ge 0; \quad \mu_2(u,\chi_{10},23) = \frac{1}{4}(16\nu_{2a}+272) \ge 0; \\ \mu_0(u,\chi_2,23) &= \frac{1}{4}(10\nu_{2a}+2\nu_{4a}+26) \ge 0; \\ \mu_2(u,\chi_2,23) &= \frac{1}{4}(-10\nu_{2a}-2\nu_{4a}+26) \ge 0; \\ \mu_0(u,\chi_3,23) &= \frac{1}{4}(-6\nu_{2a}+2\nu_{4a}+42) \ge 0; \\ \mu_2(u,\chi_3,23) &= \frac{1}{4}(6\nu_{2a}-2\nu_{4a}+42) \ge 0, \end{aligned}$$

that has only three integral solutions  $(\nu_{2a}, \nu_{4a}) \in \{ (0, 1), (-2, 3), (2, -1) \}$  satisfying Proposition 6, such that all  $\mu_i(u, \chi_j, 23)$  are non-negative integers. • Let u be a unit of order 6. Clearly,  $\nu_{2a} + \nu_{3a} + \nu_{6a} = 1$  by (1) and Proposition 4.

Now from the system of inequalities

$$\begin{split} \mu_0(u,\chi_3,*) &= \frac{1}{6}(-6\nu_{2a}+42) \ge 0; \qquad \mu_3(u,\chi_3,*) = \frac{1}{6}(6\nu_{2a}+48) \ge 0; \\ \mu_0(u,\chi_{12},*) &= \frac{1}{6}(-8\nu_{3a}+888) \ge 0; \qquad \mu_3(u,\chi_{12},*) = \frac{1}{6}(8\nu_{3a}+888) \ge 0; \\ \mu_1(u,\chi_2,*) &= \frac{1}{6}(6\nu_{2a}+4\nu_{3a}+12) \ge 0; \\ \mu_2(u,\chi_2,*) &= \frac{1}{6}(-6\nu_{2a}-4\nu_{3a}+24) \ge 0; \\ \mu_1(u,\chi_5,*) &= \frac{1}{6}(-6\nu_{2a}+5\nu_{3a}+\nu_{6a}+203) \ge 0; \\ \mu_0(u,\chi_4,7) &= \frac{1}{6}(32\nu_{2a}+2\nu_{3a}+2\nu_{6a}+226) \ge 0; \\ \mu_3(u,\chi_2,23) &= \frac{1}{6}(-10\nu_{2a}-6\nu_{3a}+2\nu_{6a}+22) \ge 0, \end{split}$$

we obtain only those integral solutions that are listed in part (iv) of the Theorem. • Let u be a unit of order 7. By (1) and Proposition 4 we have  $\nu_{7a} + \nu_{7b} = 1$ . Now using (2) we obtain the system of inequalities

$$\mu_1(u, \chi_3, *) = \frac{1}{7}(4\nu_{7a} - 3\nu_{7b} + 45) \ge 0;$$
  

$$\mu_3(u, \chi_3, *) = \frac{1}{7}(-3\nu_{7a} + 4\nu_{7b} + 45) \ge 0;$$
  

$$\mu_1(u, \chi_2, 2) = \frac{1}{7}(-4\nu_{7a} + 3\nu_{7b} + 11) \ge 0;$$
  

$$\mu_3(u, \chi_2, 2) = \frac{1}{7}(3\nu_{7a} - 4\nu_{7b} + 11) \ge 0,$$

that has solutions  $(\nu_{7a}, \nu_{7b}) \in \{ (0, 1), (2, -1), (1, 0), (-1, 2) \}.$ 

• Let u be a unit of order 8. We get  $\nu_{2a} + \nu_{4a} + \nu_{8a} = 1$  by (1) and Proposition 4. It is necessary to consider three cases, defined by part (iii) of the Theorem. By (2)

we obtain the system of inequalities

$$\mu_0(u, \chi_4, 7) = \frac{1}{8}(64\nu_{2a} + \alpha_1) \ge 0; \quad \mu_4(u, \chi_4, 7) = \frac{1}{8}(-64\nu_{2a} + \alpha_1) \ge 0;$$

$$\mu_0(u, \chi_2, *) = \frac{1}{8}(24\nu_{2a} + 8\nu_{4a} + \alpha_2) \ge 0;$$

$$\mu_4(u, \chi_2, *) = \frac{1}{8}(-24\nu_{2a} - 8\nu_{4a} + \alpha_2) \ge 0;$$

$$\mu_0(u, \chi_3, *) = \frac{1}{8}(-12\nu_{2a} + 4\nu_{4a} - 4\nu_{8a} + \alpha_3) \ge 0;$$

$$\mu_4(u, \chi_3, *) = \frac{1}{8}(12\nu_{2a} - 4\nu_{4a} + 4\nu_{8a} + \alpha_3) \ge 0;$$

$$\mu_0(u, \chi_2, 23) = \frac{1}{8}(20\nu_{2a} + 4\nu_{4a} - 4\nu_{8a} + \alpha_4) \ge 0;$$

$$\mu_4(u, \chi_2, 23) = \frac{1}{8}(-20\nu_{2a} - 4\nu_{4a} + 4\nu_{8a} + \alpha_4) \ge 0;$$

$$\mu_4(u, \chi_2, 23) = \frac{1}{8}(-20\nu_{2a} - 4\nu_{4a} + 4\nu_{8a} + \alpha_4) \ge 0,$$
where  $(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = \begin{cases} (224, 32, 44, 28), & \text{if } \chi(u^2) = \chi(4a); \\ (160, 16, 60, 12), & \text{if } \chi(u^2) = -2\chi(2a) + 3\chi(4a); \end{cases}$ 

here  $(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = \begin{cases} (160, 16, 60, 12), & \text{if } \chi(u^2) = -2\chi(2a) + 3\chi(4a); \\ (288, 48, 28, 44), & \text{if } \chi(u^2) = 2\chi(2a) - \chi(4a). \end{cases}$ 

Union of solutions for all three cases, taking into account restrictions from Proposition 6, gives us part (vi) of the Theorem.

• Let u be a unit of order 11. Clearly,  $\nu_{11a} + \nu_{11b} = 1$  by (1) and Proposition 4. Using  $\mathfrak{BCT}(3)$  by (2) we obtain the system of inequalities

$$\mu_1(u, \chi_5, 3) = \frac{1}{11} (6\nu_{11a} - 5\nu_{11b} + 104) \ge 0;$$
  
$$\mu_2(u, \chi_5, 3) = \frac{1}{11} (-5\nu_{11a} + 6\nu_{11b} + 104) \ge 0,$$

that has only 20 integer solutions listed in part (vi) of the Theorem, such that are  $\mu_1(u, \chi_5, 3)$  and  $\mu_2(u, \chi_5, 3)$  are non-negative integers.

• Let u be a unit of order 15. Again,  $\nu_{3a} + \nu_{5a} + \nu_{15a} + \nu_{15b} = 1$  by (1) and Proposition 4. Using (2), we obtain the system of inequalities

$$\mu_0(u,\chi_7,2) = \frac{1}{15}(-40\nu_{3a}+210) \ge 0; \quad \mu_5(u,\chi_7,2) = \frac{1}{15}(20\nu_{3a}+225) \ge 0; \\ \mu_0(u,\chi_9,2) = \frac{1}{15}(-24\nu_{5a}+240) \ge 0; \quad \mu_5(u,\chi_9,2) = \frac{1}{15}(+12\nu_{5a}+240) \ge 0; \\ \mu_0(u,\chi_5,*) = \frac{1}{15}(40\nu_{3a}+240) \ge 0;$$

$$\begin{split} \mu_0(u,\chi_2,*) &= \frac{1}{15}(32\nu_{3a} + 16\nu_{5a} - 8\nu_{15a} - 8\nu_{15b} + 38) \geq 0; \\ \mu_3(u,\chi_2,*) &= \frac{1}{15}(-8\nu_{3a} - 4\nu_{5a} + 2\nu_{15a} + 2\nu_{15b} + 28) \geq 0; \\ \mu_5(u,\chi_2,*) &= \frac{1}{15}(-16\nu_{3a} - 8\nu_{5a} + 4\nu_{15a} + 4\nu_{15b} + 26) \geq 0; \\ \mu_1(u,\chi_7,*) &= \frac{1}{15}(-3\nu_{3a} + \nu_{5a} + 7\nu_{15a} - 8\nu_{15b} + 233) \geq 0; \\ \mu_7(u,\chi_7,*) &= \frac{1}{15}(-3\nu_{3a} + \nu_{5a} - 8\nu_{15a} + 7\nu_{15b} + 233) \geq 0; \\ \mu_1(u,\chi_2,2) &= \frac{1}{15}(2\nu_{3a} + \nu_{5a} - 8\nu_{15a} + 7\nu_{15b} + 8) \geq 0; \\ \mu_7(u,\chi_2,2) &= \frac{1}{15}(2\nu_{3a} + \nu_{5a} - 8\nu_{15a} + 7\nu_{15b} + 8) \geq 0; \\ \mu_0(u,\chi_4,7) &= \frac{1}{15}(8\nu_{3a} - 16\nu_{5a} + 8\nu_{15a} + 8\nu_{15b} + 202) \geq 0; \\ \mu_1(u,\chi_{10},23) &= \frac{1}{15}(-\nu_{3a} - 6\nu_{15a} - 9\nu_{15b} + 279) \geq 0; \\ \mu_1(u,\chi_{12},23) &= \frac{1}{15}(-\nu_{3a} - 6\nu_{15a} + 9\nu_{15b} + 666) \geq 0. \end{split}$$

This system has only six solutions such that all  $\mu_i(u, \chi_j, p)$  are non-negative integers, listed in part (viii) of the Theorem.

• Let u be a unit of order 23. By (1) and Proposition 4 we have  $\nu_{23a} + \nu_{23b} = 1$ . Using (2) we obtain the system of inequalities

$$\mu_1(u, \chi_2, 2) = \frac{1}{23}(12\nu_{23a} - 11\nu_{23b} + 11) \ge 0;$$
  
$$\mu_5(u, \chi_2, 2) = \frac{1}{23}(-11\nu_{23a} + 12\nu_{23b} + 11) \ge 0,$$

that has only two trivial solutions:  $(\nu_{23a}, \nu_{23b}) \in \{ (1,0), (0,1) \}.$ 

Using (2) we obtain the system of inequalities

It remains to prove part (i) of the Theorem. • Let u be a unit of order 10. By (1) and Proposition 4 we have  $\nu_{2a} + \nu_{5a} = 1$ .

$$\mu_0(u,\chi_5,*) = \frac{1}{10}(88\nu_{2a} + 252) \ge 0; \quad \mu_5(u,\chi_5,*) = \frac{1}{10}(-88\nu_{2a} + 208) \ge 0; \\ \mu_1(u,\chi_5,3) = \frac{1}{10}(8\nu_{2a} - \nu_{5a} + 97) \ge 0,$$

that has no integral solutions such that all  $\mu_i(u, \chi_j, p)$  are non-negative integers. • Let u be a unit of order 21. We have  $\nu_{3a} + \nu_{7a} + \nu_{7b} = 1$  by (1) and Proposition 4. We need to consider four cases given in part (v) of the Theorem:

Case 1. 
$$\chi(u^3) = \chi(7a)$$
.  
Case 2.  $\chi(u^3) = \chi(7a) - \chi(7b)$ .  
Case 2.  $\chi(u^3) = \chi(7b)$ .  
Case 4.  $\chi(u^3) = -\chi(7a) + 2\chi(7b)$ .

In all these cases using (2) we obtain the system of inequalities

$$\mu_0(u, \chi_2, 23) = \frac{1}{21}(36\nu_{3a} + 27) \ge 0; \quad \mu_7(u, \chi_2, 23) = \frac{1}{21}(-18\nu_{3a} + 18) \ge 0;$$
  
$$\mu_1(u, \chi_{10}, 23) = \frac{1}{21}(\nu_{3a} + 279) \ge 0,$$

that has no integral solutions such that all  $\mu_i(u, \chi_j, 23)$  are non-negative integers. • Let u be a unit of order 22. We have  $\nu_{2a} + \nu_{11a} + \nu_{11b} = 1$  by (1) and Proposition 4. We need to consider 20 cases given in part (vi) of the Theorem, but in all cases using (2) we obtain the system of inequalities

$$\mu_0(u,\chi_2,*) = \frac{1}{22}(60\nu_{2a} + 28) \ge 0; \quad \mu_{11}(u,\chi_2,*) = \frac{1}{22}(-60\nu_{2a} + 16) \ge 0,$$

that has no integral solutions such that all  $\mu_i(u, \chi_j, *)$  are non-negative integers. • Let u be a unit of order 33. We have  $\nu_{3a} + \nu_{11a} + \nu_{11b} = 1$  by (1) and Proposition 4. Again, we have 20 cases, and all of them give us the same system

$$\mu_0(u,\chi_2,*) = \frac{1}{33}(80\nu_{3a} + 30) \ge 0; \quad \mu_{11}(u,\chi_2,*) = \frac{1}{33}(-40\nu_{3a} + 18) \ge 0,$$

that has no integral solutions such that all  $\mu_i(u, \chi_j, *)$  are non-negative integers. • Let u be a unit of order 35. We have  $\nu_{5a} + \nu_{11a} + \nu_{11b} = 1$  by (1) and Proposition 4. We need to consider 4 cases, given by part (v) of the Theorem, and all of them give us the same system of inequalities

$$\mu_0(u,\chi_2,23) = \frac{1}{35}(24\nu_{5a}+25) \ge 0; \quad \mu_7(u,\chi_2,23) = \frac{1}{35}(-6\nu_{5a}+20) \ge 0,$$

that has no integral solutions such that all  $\mu_i(u, \chi_j, 23)$  are non-negative integers. • Let u be a unit of order 46. We have  $\nu_{2a} + \nu_{23a} + \nu_{23b} = 1$  by (1) and Proposition 4. We have two cases:  $\chi(u^2) \in \{\chi(23a), \chi(23b)\}$ , but in both of them

$$\mu_0(u,\chi_9,*) = \frac{1}{46}(286\nu_{2a} + 266) \ge 0; \quad \mu_{23}(u,\chi_9,*) = \frac{1}{46}(-286\nu_{2a} + 240) \ge 0,$$

that has no integral solutions such that all  $\mu_i(u, \chi_j, *)$  are non-negative integers. • Let u be a unit of order 55. We have  $\nu_{5a} + \nu_{11a} + \nu_{11b} = 1$  by (1) and Proposition 4. As before, we need to consider 20 cases given by part (vii) of the Theorem, and all of them give us the same system of inequalities

$$\mu_0(u,\chi_2,2) = \frac{1}{55}(40\nu_{5a} + 15) \ge 0; \quad \mu_5(u,\chi_2,2) = \frac{1}{55}(-4\nu_{5a} + 15) \ge 0,$$

that has no integral solutions such that all  $\mu_i(u, \chi_j, 2)$  are non-negative integers. • Let u be a unit of order 69. We have  $\nu_{3a} + \nu_{23a} + \nu_{23b} = 1$  by (1) and Proposition 4. We have two cases:  $\chi(u^3) \in \{\chi(23a), \chi(23b)\}$ , but both of them gives

$$\mu_0(u, \chi_3, *) = \frac{1}{69} (-44(\nu_{23a} + \nu_{23b}) + 23) \ge 0$$
  
$$\mu_{23}(u, \chi_3, *) = \frac{1}{69} (22(\nu_{23a} + \nu_{23b}) + 23) \ge 0,$$

that has no integral solutions such that all  $\mu_i(u, \chi_j, *)$  are non-negative integers. • Let u be a unit of order 77. Clearly,  $\nu_{7a} + \nu_{7b} + \nu_{11a} + \nu_{11b} = 1$  by (1) and Proposition 4. We need to consider 80 cases, determined by parts (v) and (vii) of the Theorem. All of them give us the same system of inequalities

$$\mu_0(u, \chi_2, 2) = \frac{1}{77} (30(\nu_{7a} + \nu_{7b}) + 14) \ge 0;$$
  
$$\mu_0(u, \chi_4, 2) = \frac{1}{77} (-90(\nu_{7a} + \nu_{7b}) + 35) \ge 0,$$

that has no integral solutions such that all  $\mu_0(u, \chi_j, 2)$  are non-negative integers. • Let u be a unit of order 115. We have  $\nu_{5a} + \nu_{23a} + \nu_{23b} = 1$  by (1) and Proposition 4. Consider two cases:  $\chi(u^5) \in \{\chi(23a), \chi(23b)\}$ . Both of them give

$$\mu_0(u,\chi_3,*) = \frac{1}{115} (-88(\nu_{23a} + \nu_{23b}) + 23) \ge 0$$
  
$$\mu_{23}(u,\chi_3,*) = \frac{1}{115} (22(\nu_{23a} + \nu_{23b}) + 23) \ge 0,$$

that has no integral solutions such that all  $\mu_i(u, \chi_j, *)$  are non-negative integers. • Let u be a unit of order 161. We have  $\nu_{7a} + \nu_{7b} + \nu_{23a} + \nu_{23b} = 1$  by (1) and Proposition 4. Consider 8 cases, determined by possible values of  $\chi(u^{23})$  and  $\chi(u^7)$ . All of them give us the same system of inequalities

$$\mu_0(u, \chi_5, *) = \frac{1}{161} (-132(\nu_{7a} + \nu_{7b}) + 224) \ge 0;$$
  
$$\mu_0(u, \chi_9, *) = \frac{1}{161} (132(\nu_{7a} + \nu_{7b}) + 259) \ge 0,$$

that has no integral solutions such that all  $\mu_0(u, \chi_j, *)$  are non-negative integers. • Let u be a unit of order 253. Again,  $\nu_{11a} + \nu_{11b} + \nu_{23a} + \nu_{23b} = 1$  by (1) and Proposition 4. We need to consider 40 cases, determined by possible values of  $\chi(u^{23})$  and  $\chi(u^{11})$ . All of them give us the same system of inequalities

$$\mu_0(u, \chi_2, *) = \frac{1}{253} (-220(\nu_{23a} + \nu_{23b})) \ge 0;$$
  

$$\mu_{23}(u, \chi_2, *) = \frac{1}{253} (22(\nu_{23a} + \nu_{23b})) \ge 0;$$
  

$$\mu_{11}(u, \chi_6, *) = \frac{1}{253} (-10(\nu_{23a} + \nu_{23b}) + 230) \ge 0,$$

that has no integral solutions such that all  $\mu_0(u, \chi_j, *)$  are non-negative integers.

Acknowledgment. The authors are grateful to Dr. Steve Linton and Dr. Tom Kelsey from the University of St Andrews for their advice with computational issues.

### References

- [AB89] V. A. Artamonov and A. A. Bovdi. Integral group rings: groups of invertible elements and classical K-theory. In Algebra. Topology. Geometry, Vol. 27 (Russian), Itogi Nauki i Tekhniki, pages 3–43, 232. Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989. Translated in J. Soviet Math. 57 (1991), no. 2, 2931–2958.
- [Ber55] S. D. Berman. On the equation  $x^m = 1$  in an integral group ring. Ukrain. Mat. Z., 7:253–261, 1955.
- [BH07] V. Bovdi and M. Hertweck. Zassenhaus conjecture for central extensions of S<sub>5</sub>. J. Group Theory, pages 1–11, to appear, 2007. (E-print arXiv:math.RA/0609435v1).

- [BHK04] V. Bovdi, C. Höfert, and W. Kimmerle. On the first Zassenhaus conjecture for integral group rings. Publ. Math. Debrecen, 65(3-4):291–303, 2004.
- [BJK07] V. Bovdi, E. Jespers, and A. Konovalov. Torsion units in integral group rings of Janko simple groups. *Preprint*, pages 1–30, submitted, 2007. (E-print arXiv:math/0608441v3).
- [BK00] F.M. Bleher and W. Kimmerle. On the structure of integral group rings of sporadic groups. LMS J. Comput. Math., 3:274–306 (electronic), 2000.
- [BK07] V. Bovdi and A. Konovalov. Integral group ring of the first Mathieu simple group. In Groups St. Andrews 2005. Vol. I, volume 339 of London Math. Soc. Lecture Note Ser., pages 237–245. Cambridge Univ. Press, Cambridge, 2007.
- [BKRS07] V. Bovdi, A. Konovalov, R. Rossmanith, and Cs. Schneider. LA-GUNA – Lie AlGebras and UNits of group Algebras, Version 3.4, 2007. (http://ukrgap.exponenta.ru/laguna.htm).
- [BKS07] V. Bovdi, A. Konovalov, and S. Siciliano. Integral group ring of the Mathieu simple group M<sub>12</sub>. Rend. Circ. Mat. Palermo (2), 56:125–136, 2007.
- [CL65] J.A. Cohn and D. Livingstone. On the structure of group algebras. I. Canad. J. Math., 17:583–593, 1965.
- [GAP06] The GAP Group. GAP Groups, Algorithms, and Programming, Version 4.4.9, 2006. (http://www.gap-system.org).
- [Gor83] D. Gorenstein. The classification of finite simple groups. Vol. 1. The University Series in Mathematics. Plenum Press, New York, 1983.
- [Her06] M. Hertweck. On the torsion units of some integral group rings. Algebra Colloq., 13(2):329–348, 2006.
- [Her07a] M. Hertweck. Partial augmentations and Brauer character values of torsion units in group rings. Comm. Algebra, pages 1–16, to appear, 2007. (E-print arXiv:math.RA/0612429v2).
- [Her07b] M. Hertweck. Torsion units in integral group rings or certain metabelian groups. Proc. Edinb. Math. Soc., pages 1–22, to appear, 2007.
- [HK06] C. Höfert and W. Kimmerle. On torsion units of integral group rings of groups of small order. In Groups, rings and group rings, volume 248 of Lect. Notes Pure Appl. Math., pages 243–252. Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [Kim06] W. Kimmerle. On the prime graph of the unit group of integral group rings of finite groups. In Groups, rings and algebras, volume 420 of Contemporary Mathematics, pages 215–228. AMS, 2006.
- [LP89] I. S. Luthar and I. B. S. Passi. Zassenhaus conjecture for A<sub>5</sub>. Proc. Indian Acad. Sci. Math. Sci., 99(1):1–5, 1989.
- [LT91] I. S. Luthar and P. Trama. Zassenhaus conjecture for S<sub>5</sub>. Comm. Algebra, 19(8):2353– 2362, 1991.
- [MRSW87] Z. Marciniak, J. Ritter, S. K. Sehgal, and A. Weiss. Torsion units in integral group rings of some metabelian groups. II. J. Number Theory, 25(3):340–352, 1987.
- [Zas74] H. Zassenhaus. On the torsion units of finite group rings. In Studies in mathematics (in honor of A. Almeida Costa) (Portuguese), pages 119–126. Instituto de Alta Cultura, Lisbon, 1974.