ON CLASSIFICATION OF FINITE DIMENSIONAL COMPLEX FILIFORM LEIBNIZ ALGEBRAS (PartI)

Bekbaev U.D.¹, Rakhimov I.S.²

Department of Mathematics & Institute for Mathematical Research,

FS,UPM, 43400, Serdang, Selangor Darul Ehsan, Malaysia.

Abstract

The paper is devoted to classification problem of finite dimensional complex none Lie filiform Leibniz algebras. The motivation to write this paper is an unpublished yet result of J.R.Gomez, B.A.Omirov [1] on necessary and sufficient conditions for two finite dimensional complex filiform Leibniz algebras to be isomorphic. We suggest another approach to this problem. The approach that we use for classification is in terms of invariants. In fact, utilizing this method for any given low dimensional case all filiform Leibniz algebras can be classified. Moreover, the results can be used for geometrical classification of orbits of such algebras.

2000 MSC: 17A32, 17B30.

Key-Words: filiform Leibniz algebra, invariant function, isomorphism.

1 Introduction

This paper aims to investigate a class of nonassociative algebras which generalizes the class of Lie algebras. These algebras satisfy certain identities that were suggested by J.-L.Loday [2]. When he used the tensor product instead of external product in the definition of the n-th cochain, in order to prove the differential property, that is defined on cochains, it sufficed to replace the anticommutativity and Jacoby identity by the Leibniz identity. This is an essential one of the motivation to appear for this class of algebras.

In this paper we suggest an algebraic approach to the classification problem for filiform Leibniz algebras. Utilizing this method for any fixed low dimensional case the corresponding classes of filiform Leibniz algebras can be classified completely. Moreover, the results may be used for geometric classification in the sense of geometric invariant theory [3]. It is assumed that it will be the subject of one of the next papers. For

¹e-mail: bekbaev@science.upm.edu.my

²e-mail: isamiddin@science.upm.edu.my

geometric classification of complex nilpotent Leibniz algebras of dimension at most four we refer to [4].

Let V be a vector space of dimension n over an algebraically closed field K (charK=0). The bilinear maps $V \times V \to V$ form a vector space $Hom(V \otimes V, V)$ of dimension n^3 , which can be considered together with its natural structure of an affine algebraic variety over K and denoted by $Alg_n(K) \cong K^{n^3}$. An n-dimensional algebra L over K may be considered as an element $\lambda(L)$ of $Alg_n(K)$ via the bilinear mapping $\lambda : L \otimes L \to L$ defining an binary algebraic operation on $L : \text{let } \{e_1, e_2, \ldots, e_n\}$ be a basis of the algebra L. Then the table of multiplication of L is represented by point (γ_{ij}^k) of this affine space as follow:

$$\lambda(e_i, e_j) = \sum_{k=1}^n \gamma_{ij}^k e_k.$$

 γ_{ij}^k are called *structural constants* of *L*. The linear reductive group $GL_n(K)$ acts on $Alg_n(K)$ by $(g * \lambda)(x, y) = g(\lambda(g^{-1}(x), g^{-1}(y)))$ ("transport of structure"). Two algebras λ_1 and λ_2 are isomorphic if and only if they belong to the same orbit under this action. The orbit of λ under this action is denoted by $O(\lambda)$. It is clear that elements of the given orbit are isomorphic to each other algebras. The classification means to specify the representatives of the orbits. A simple criterion, to decide if the given two algebras are isomorphic, is desired.

2 Preliminaries

Definition 1. An algebra L over a field K is called a *Leibniz algebra* if it satisfies the following Leibniz identity:

$$[x, [y, z]] = [[x, y], z] - [[x, z], y],$$

where $[\cdot, \cdot]$ denotes the multiplication in L. Let $Leib_n(K)$ be a subvariety of $Alg_n(K)$ consisting of all *n*-dimensional Leibniz algebras over K. It is invariant under the above mentioned action of $GL_n(K)$. As a subset of $Alg_n(K)$ the set $Leib_n(K)$ is specified by system of equations with respect to structural constants γ_{ij}^k :

$$\sum_{l=1}^{n} \left(\gamma_{jk}^{l} \gamma_{il}^{m} - \gamma_{ij}^{l} \gamma_{lk}^{m} + \gamma_{ik}^{l} \gamma_{lj}^{m} \right) = 0$$

It is easy to see that if the bracket in Leibniz algebra happens to be anticommutative then it is a Lie algebra. So Leibniz algebras are "noncommutative" generalization of Lie algebras. As to classifications of low dimensional Lie algebras they are well known. But unless simple Lie algebras the classification problem of all Lie algebras in common remains a big problem. Yu.I.Malcev [4] reduced the classification of solvable Lie algebras to the classification of nilpotent Lie algebras. Apparently the first non-trivial classification of some classes of low-dimensional nilpotent Lie algebra are due to Umlauf. In his thesis [6] he presented the redundant list of nilpotent Lie algebras of dimension at most seven. He gave also the list of nilpotent Lie algebras of dimension less than ten admitting so-called adapted basis (now, the nilpotent Lie algebras with this property are called *filiform Lie algebras*). It was shown by M.Vergne [7] the impotentness of filiform Lie algebras in the study of variety of nilpotent Lie algebras laws. Up to now the several classifications of low-dimensional nilpotent Lie algebras have been done. Unfortunately, many of these papers are based on direct computations (by hand) and the complexity of those computations leads frequently to errors. We refer the reader to [8] for comments and corrections of the classification errors.

Further if it is not asserted additionally all algebras assumed to be over the field of complex numbers.

Let L be a Leibniz algebra. We put:

$$L^1 = L, \quad L^{k+1} = [L^k, L], \ k \in N.$$

Definition 2. A Leibniz algebra L is said to be *nilpotent* if there exists an integer $s \in N$, such that $L^1 \supset L^2 \supset ... \supset L^s = \{0\}$. The smallest integer s for which $L^s = 0$ is called *the nilindex* of L.

Definition 3. An *n*-dimensional Leibniz algebra L is said to be *filiform* if $dimL^i = n - i$, where $2 \le i \le n$.

Theorem 1[9]. Arbitrary complex non-Lie filiform Leibniz algebra of dimension n + 1 is isomorphic to one of the following filiform Leibniz algebras:

a) (The first class):
$$\begin{cases} [e_0, e_0] = e_2, \\ [e_i, e_0] = e_{i+1}, & 1 \le i \le n-1 \\ [e_0, e_1] = \alpha_3 e_3 + \alpha_4 e_4 + \dots + \alpha_{n-1} e_{n-1} + \theta e_n, \\ [e_j, e_1] = \alpha_3 e_{j+2} + \alpha_4 e_{j+3} + \dots + \alpha_{n+1-j} e_n, & 1 \le j \le n-2 \end{cases}$$

b) (The second class):
$$\begin{cases} [e_0, e_0] = e_2, \\ [e_i, e_0] = e_{i+1}, \\ [e_0, e_1] = \beta_3 e_3 + \beta_4 e_4 + \dots + \beta_n e_n, \\ [e_1, e_1] = \gamma e_n, \\ [e_j, e_1] = \beta_3 e_{j+2} + \beta_4 e_{j+3} + \dots + \beta_{n+1-j} e_n, \\ 2 \le j \le n-2 \end{cases}$$

where [,] is the multiplication in a Leibniz algebra, $\{e_0, e_1, \ldots, e_n\}$ is a basis of the algebra and omitted products are assumed to be zero.

Note that the algebras from the first class and the second class never are isomorphic to each other.

In this paper we will deal with the first class of algebras, as for the second class it will be considered in our another paper.

Let us denote by $L(\alpha)$, the (n+1)-dimensional filiform non-Lie Leibniz algebra defined by parameters $\alpha = (\alpha_3, \alpha_4, ..., \alpha_n, \theta)$. The set of all (n+1)-dimensional complex filiform Leibniz algebras is denoted by FL_{n+1} . It is a closed and invariant subset of the variety of nilpotent Leibniz algebras.

Using the method of simplification of the basis transformations in [1] the following criterion on isomorphism of two (n + 1)-dimensional filiform Leibniz algebras was given. Namely: let $n \ge 3$.

Theorem 2.[1] Two algebras $L(\alpha)$ and $L(\alpha')$ from FL_{n+1} , where $\alpha = (\alpha_3, \alpha_4, ..., \alpha_n, \theta)$ and $\alpha' = (\alpha'_3, \alpha'_4, ..., \alpha'_n, \theta')$, are isomorphic if and only if there exist complex numbers A,B such that $A(A+B)\neq 0$ and the following conditions hold:

$$\begin{aligned} \alpha_3' &= \frac{A+B}{A^2} \alpha_3, \\ \alpha_t' &= \frac{1}{A^{t-1}} ((A+B)\alpha_t - \sum_{k=3}^{t-1} (C_{k-1}^{k-2}A^{k-2}B\alpha_{t+2-k} + C_{k-1}^{k-3}A^{k-3}B^2 \sum_{i_1=k+2}^{t} \alpha_{t+3-i_1} \cdot \alpha_{i_1+1-k} + \\ C_{k-1}^{k-4}A^{k-4}B^3 \sum_{i_2=k+3}^{t} \sum_{i_1=k+3}^{i_2} \alpha_{t+3-i_2} \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1-k} + \dots + \\ C_{k-1}^{1}AB^{k-2} \sum_{i_{k-3}=2k-2}^{t} \sum_{i_{k-4}=2k-2}^{i_{k-3}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{t+3-i_{k-3}} \cdot \alpha_{i_{k-3}+3-i_{k-4}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+5-2k} \\ &+ B^{k-1} \sum_{i_{k-2}=2k-1}^{t} \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} \alpha_{t+3-i_{k-2}} \cdot \alpha_{i_{k-2}+3-i_{k-3}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \alpha_{i_1+4-2k}) \cdot \alpha_k'), \end{aligned}$$
where $4 \leq t \leq n$.

$$\theta' &= \frac{1}{A^{n-1}} (A\theta + B\alpha_n - \sum_{k=3}^{n-1} (C_{k-1}^{k-2}A^{k-2}B\alpha_{n+2-k} + C_{k-1}^{k-3}A^{k-3}B^2 \sum_{i_1=k+2}^{n} \alpha_{n+3-i_1} \cdot \alpha_{i_1+1-k} + \\ C_{k-1}^{k-4}A^{k-4}B^3 \sum_{i_2=k+3}^{n} \sum_{i_1=k+3}^{i_2} \alpha_{n+3-i_2} \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1-k} + \dots + \\ C_{k-1}^{1}AB^{k-2} \sum_{i_{k-3}=2k-2}^{n} \sum_{i_{k-4}=2k-2}^{i_{k-4}} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{n+3-i_{k-3}} \cdot \alpha_{i_{k-3}+3-i_{k-4}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+5-2k} \\ &+ B^{k-1} \sum_{i_{k-2}=2k-1}^{n} \sum_{i_{k-3}=2k-2}^{n} \dots \sum_{i_1=2k-2}^{i_2} \alpha_{n+3-i_{k-3}} \cdot \alpha_{i_{k-3}+3-i_{k-4}} \cdot \dots \cdot \alpha_{i_2+3-i_1} \cdot \alpha_{i_1+4-2k}) \cdot \alpha_k'), \end{aligned}$$

Here are the above systems of equalities for some low dimensional cases:

Case of n = 4 i.e. **dimL=5**:

$$\begin{cases} \alpha'_{3} = \frac{1}{A}(1 + \frac{B}{A})\alpha_{3}, \\ \\ \alpha'_{4} = \frac{1}{A^{2}}(1 + \frac{B}{A})(\alpha_{4} - 2\frac{B}{A}\alpha_{3}^{2}) \\ \\ \\ \theta' = \frac{1}{A^{2}}[\theta + \frac{B}{A}\alpha_{4} - 2(1 + \frac{B}{A})\frac{B}{A}\alpha_{3}^{2}]. \end{cases}$$

Case of n = 5 i.e. **dimL=6**:

$$\begin{cases} \alpha_3' = \frac{1}{A} (1 + \frac{B}{A}) \alpha_3, \\ \alpha_4' = \frac{1}{A^2} (1 + \frac{B}{A}) (\alpha_4 - 2\frac{B}{A}\alpha_3^2) \\ \alpha_5' = \frac{1}{A^3} (1 + \frac{B}{A}) [\alpha_5 - 5\frac{B}{A} (\alpha_4 - \frac{B}{A}\alpha_3^2) \alpha_3] \\ \theta' = \frac{1}{A^3} [\theta + \frac{B}{A} \alpha_5 - 5(1 + \frac{B}{A})\frac{B}{A} (\alpha_4 - \frac{B}{A}\alpha_3^2) \alpha_3]. \end{cases}$$

Case of n = 6 i.e. **dimL=7**:

$$\begin{cases} \alpha_3' = \frac{1}{A} (1 + \frac{B}{A}) \alpha_3, \\ \alpha_4' = \frac{1}{A^2} (1 + \frac{B}{A}) (\alpha_4 - 2\frac{B}{A}\alpha_3^2) \\ \alpha_5' = \frac{1}{A^3} (1 + \frac{B}{A}) [\alpha_5 - 5\frac{B}{A} (\alpha_4 - \frac{B}{A}\alpha_3^2) \alpha_3] \\ \alpha_6' = \frac{1}{A^4} (1 + \frac{B}{A}) [\alpha_6 - 6\frac{B}{A} \alpha_3 \alpha_5 + 21(\frac{B}{A})^2 \alpha_3^2 \alpha_4 - 3\frac{B}{A} \alpha_4^2 - 14(\frac{B}{A})^3 \alpha_3^4] \\ \theta' = \frac{1}{A^4} \{\theta + \frac{B}{A} \alpha_6 - (1 + \frac{B}{A}) [6\frac{B}{A} \alpha_3 \alpha_5 - 21(\frac{B}{A})^2 \alpha_3^2 \alpha_4 + 3\frac{B}{A} \alpha_4^2 + 14(\frac{B}{A})^3 \alpha_3^4] \}.$$

Case of n = 7 i.e. **dimL=8**:

$$\begin{cases} \alpha'_{3} = \frac{1}{A} (1 + \frac{B}{A}) \alpha_{3}, \\ \alpha'_{4} = \frac{1}{A^{2}} (1 + \frac{B}{A}) (\alpha_{4} - 2\frac{B}{A}\alpha_{3}^{2}) \\ \alpha'_{5} = \frac{1}{A^{3}} (1 + \frac{B}{A}) [\alpha_{5} - 5\frac{B}{A}(\alpha_{4} - \frac{B}{A}\alpha_{3}^{2})\alpha_{3}] \\ \alpha'_{6} = \frac{1}{A^{4}} (1 + \frac{B}{A}) [\alpha_{6} - 6\frac{B}{A}\alpha_{3}\alpha_{5} + 21(\frac{B}{A})^{2}\alpha_{3}^{2}\alpha_{4} - 3\frac{B}{A}\alpha_{4}^{2} - 14(\frac{B}{A})^{3}\alpha_{3}^{4}] \\ \alpha'_{7} = \frac{1}{A^{5}} (1 + \frac{B}{A}) [\alpha_{7} - 7\frac{B}{A}\alpha_{3}\alpha_{6} + 28(\frac{B}{A})^{2}\alpha_{3}^{2}\alpha_{5} + 28(\frac{B}{A})^{2}\alpha_{3}\alpha_{4}^{2} - 7\frac{B}{A}\alpha_{4}\alpha_{5} \\ -84(\frac{B}{A})^{3}\alpha_{3}^{3}\alpha_{4} + 42(\frac{B}{A})^{4}\alpha_{5}^{5}] \\ \theta' = \frac{1}{A^{5}} \{\theta + \frac{B}{A}\alpha_{7} - (1 + \frac{B}{A})[7\frac{B}{A}\alpha_{3}\alpha_{6} - 28(\frac{B}{A})^{2}\alpha_{3}^{2}\alpha_{5} - 28(\frac{B}{A})^{2}\alpha_{3}\alpha_{4}^{2} + 7\frac{B}{A}\alpha_{4}\alpha_{5} \\ +84(\frac{B}{A})^{3}\alpha_{3}^{3}\alpha_{4} - 42(\frac{B}{A})^{4}\alpha_{5}^{5}] \}. \end{cases}$$

To deal with the classification of FL_{n+1} with respect to the above mentioned action we represent it as a disjoint union of an open and closed (with respect to the Zarisski topology) subsets. Moreover each of these subsets are invariant under the corresponding transformations presented in Theorem 2. Then we formulate the solution of the isomorphism problem for the corresponding algebras from the open subset. Similar approach can be used to solve isomorphism problem for the algebras from the corresponding closed subset. It is not difficult to notice that the expressions for α'_t , θ' in Theorem 2 can be represented in the following form:

$$\alpha_t' = \frac{1}{A^{t-2}}\varphi_t(\frac{B}{A};\alpha),$$

where $\alpha = (\alpha_3, \alpha_4, ..., \alpha_n, \theta)$ and $\varphi_t(y; z) = \varphi_t(y; z_3, z_4, ..., z_n, z_{n+1}) =$

$$\left((1+y)z_t - \sum_{k=3}^{t-1} (C_{k-1}^{k-2}yz_{t+2-k} + C_{k-1}^{k-3}y^2 \sum_{i_1=k+2}^{t} z_{t+3-i_1} \cdot z_{i_1+1-k} + \right)$$

$$C_{k-1}^{k-4}y^3 \sum_{i_2=k+3}^{t} \sum_{i_1=k+3}^{i_2} z_{t+3-i_2} \cdot z_{i_2+3-i_1} \cdot z_{i_1-k} + \dots +$$

$$C_{k-1}^{1}y^{k-2}\sum_{i_{k-3}=2k-2}^{t}\sum_{i_{k-4}=2k-2}^{i_{k-3}}\dots\sum_{i_{1}=2k-2}^{i_{2}}z_{t+3-i_{k-3}}\cdot z_{i_{k-3}+3-i_{k-4}}\cdot\dots\cdot z_{i_{2}+3-i_{1}}\cdot z_{i_{1}+5-2k}+$$

$$y^{k-1} \sum_{i_{k-2}=2k-1}^{t} \sum_{i_{k-3}=2k-1}^{i_{k-2}} \dots \sum_{i_1=2k-1}^{i_2} z_{t+3-i_{k-2}} \cdot z_{i_{k-2}+3-i_{k-3}} \cdot \dots \cdot z_{i_2+3-i_1} z_{i_1+4-2k}) \cdot \varphi_k(y;z)),$$

for
$$3 \le t \le n$$
.
 $\theta' = \frac{1}{A^{n-2}}\varphi_{n+1}(\frac{B}{A};\alpha)$, where $\varphi_{n+1}(y;z) = \varphi_{n+1}(y;z_3,z_4,...,z_n,z_{n+1}) = (z_{n+1} + yz_n - (1+y)\sum_{k=3}^{n-1} (C_{k-1}^{k-2}yz_{n+2-k} + C_{k-1}^{k-3}y^2 \sum_{i_1=k+2}^n z_{n+3-i_1} \cdot z_{i_1+1-k} + (z_{n+1} + yz_n) = (z_{n+1} + yz_n) =$

$$C_{k-1}^{k-4}y^3 \sum_{i_2=k+3}^n \sum_{i_1=k+3}^{i_2} z_{n+3-i_2} \cdot z_{i_2+3-i_1} \cdot z_{i_1-k} + \dots +$$

$$C_{k-1}^{1}y^{k-2}\sum_{i_{k-3}=2k-2}^{n}\sum_{i_{k-4}=2k-2}^{i_{k-3}}\dots\sum_{i_{1}=2k-2}^{i_{2}}z_{n+3-i_{k-3}}\cdot z_{i_{k-3}+3-i_{k-4}}\cdot\dots\cdot z_{i_{2}+3-i_{1}}\cdot z_{i_{1}+5-2k}$$

$$+y^{k-1}\sum_{i_{k-2}=2k-1}^{n}\sum_{i_{k-3}=2k-1}^{i_{k-2}}\dots\sum_{i_{1}=2k-1}^{i_{2}}z_{n+3-i_{k-2}}\cdot z_{i_{k-2}+3-i_{k-3}}\cdot\dots\cdot z_{i_{2}+3-i_{1}}\cdot z_{i_{1}+4-2k})\cdot\varphi_{k}(y;z))$$

To simplify notation let us agree that in the above case for transition from the (n+1)dimensional filiform Leibniz algebra $L(\alpha)$ to the (n + 1)-dimensional filiform Leibniz algebra $L(\alpha')$ we write $\alpha' = \rho(\frac{1}{A}, \frac{B}{A}; \alpha)$, where $\alpha = (\alpha_3, \alpha_4, ..., \alpha_n, \theta)$,

$$\rho(\frac{1}{A}, \frac{B}{A}; \alpha) = (\rho_1(\frac{1}{A}, \frac{B}{A}; \alpha), \rho_2(\frac{1}{A}, \frac{B}{A}; \alpha), \dots, \rho_{n-1}(\frac{1}{A}, \frac{B}{A}; \alpha)),$$
$$\rho_t(x, y; z) = x^t \varphi_{t+2}(y; z) \text{ for } 1 \le t \le n-2$$

and

$$\rho_{n-1}(x, y; z) = x^{n-2}\varphi_{n+1}(y, z)$$

Here are the main properties of the operator ρ , derived from the fact that $\rho(\frac{1}{A}, \frac{B}{A}; \cdot)$ is an action of a group, that will be used later on.

1⁰. $\rho(1,0;\cdot)$ is the identity operator.

2⁰.
$$\rho(\frac{1}{A_2}, \frac{B_2}{A_2}; \rho(\frac{1}{A_1}, \frac{B_1}{A_1}; \alpha)) = \rho(\frac{1}{A_1A_2}, \frac{A_1B_2 + A_2B_1 + B_1B_2}{A_1A_2}; \alpha).$$

3⁰. If $\alpha' = \rho(\frac{1}{A}, \frac{B}{A}; \alpha)$ then $\alpha = \rho(A, -\frac{B}{A+B}; \alpha').$

3 Classification theorems

Definition 4. An action of algebraic group G on a variety Z is a morphism

 $\sigma: G \times Z \longrightarrow Z$ with (i) $\sigma(e, z) = z$, where e is the unit element of G and $z \in Z$ (ii) $\sigma(g, \sigma(h, z)) = \sigma(gh, z)$, for any $g, h \in G$ and $z \in Z$. We shortly write gz for $\sigma(g, z)$, and call Z a G-variety.

Definition 5. A morphism $f: Z \longrightarrow K$ is said to be invariant if f(gz) = f(z) for any $g \in G$ and $z \in Z$.

The algebra of invariant morphisms on Z with respect to the action of the group G is denoted by $K[Z]^G$. Sometimes this algebra is a finitely generated K-algebra. This is referred to in [10] as the "first fundamental problem of invariant theory". If Z is an irreducible then the field of rational invariants can be defined as a quotient field of $K[Z]^G$. It is always finitely generated as a subalgebra of the finitely generated algebra K(Z). Description the field of rational invariants is an another important classical problem of the invariant theory [11].

Actually, we use some elements of the algebra of invariant morphisms under the above mentioned adapted action on the variety of filiform Leibniz algebras to solve isomorphism problem. From here on n is a positive integer. We assume that $n \ge 4$ since there are complete classifications of complex nilpotent Leibniz algebras of dimension at most four [12],[13].

We consider the following presentation of the first class of all n+1-dimensional filiform Leibniz algebras:

 $U \cup F$, where $U = \{L(\alpha) : \alpha_3(\alpha_4 + 2\alpha_3^2) \neq 0\}, F = \{L(\alpha) : \alpha_3(\alpha_4 + 2\alpha_3^2) = 0\}.$

Our main interest will be the cases of open sets i.e. "generic algebras" cases.

Theorem 3. i) Two algebras $L(\alpha)$ and $L(\alpha')$ from U are isomorphic if and only if

$$\rho_i(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = \rho_i(\frac{2\alpha_3'}{\alpha_4' + 2\alpha_3'^2}, \frac{\alpha_4'}{2\alpha_3'^2}; \alpha')$$

whenever $i = \overline{3, n-1}$.

ii) For any $(a_3, a_4, ..., a_{n-1}) \in C^{n-3}$ there is an algebra $L(\alpha)$ from U such that

$$\rho_i(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = a_i \text{ for all } i = \overline{3, n-1}$$

Proof. *i*). Part "if". Let two algebras $L(\alpha)$ and $L(\alpha')$ be isomorphic that is there exist $A, B \in \mathbb{C}$ such that $A(A + B) \neq 0$ and $\alpha' = \rho(\frac{1}{A}, \frac{B}{A}; \alpha)$. Consider algebra $L(\alpha^0)$, where $\alpha^0 = \rho(\frac{1}{A_0}, \frac{B_0}{A_0}; \alpha)$ and $A_0 = \frac{\alpha_4 + 2\alpha_3^2}{2\alpha_3}$, $B_0 = \frac{\alpha_4(\alpha_4 + 2\alpha_3^2)}{4\alpha_3^3}$. So $\alpha = \rho(A, \frac{-B}{A+B}; \alpha')$ and

$$\alpha^{0} = \rho(\frac{2\alpha_{3}}{\alpha_{4} + 2\alpha_{3}^{2}}, \frac{\alpha_{4}}{2\alpha_{3}^{2}}; \alpha) = \rho(\frac{1}{A_{0}}, \frac{B_{0}}{A_{0}}; \rho(A, \frac{-B}{A+B}; \alpha')) = \rho(\frac{A}{A_{0}}, \frac{B_{0}A - A_{0}B}{A(A+B)}; \alpha').$$

It is easy to check that $\frac{A}{A_0} = \frac{2\alpha'_3}{\alpha'_4 + 2\alpha'^2_3}$ and $\frac{B_0A - A_0B}{A(A+B)} = \frac{\alpha'_4}{2\alpha'^2_3}$. Therefore

$$\rho(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = \rho(\frac{2\alpha_3'}{\alpha_4' + 2\alpha_3'^2}, \frac{\alpha_4'}{2\alpha_3'^2}; \alpha')$$

and, in particular,

$$\rho_i(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = \rho_i(\frac{2\alpha_3'}{\alpha_4' + 2\alpha_3'^2}, \frac{\alpha_4'}{2\alpha_3'^2}; \alpha'),$$

for all $i = \overline{3, n-1}$. This procedure can be shown schematically by the following picture: $\alpha \xrightarrow{(A_0, B_0)} \alpha^0$

$$(A,B) \searrow \qquad \nearrow (A_0 A^{-1}, \frac{B_0 A - A_0 B}{A_0 (A+B)})$$
$$\alpha'$$

Part "Only if". Let the equalities

$$\rho_i(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = \rho_i(\frac{2\alpha_3'}{\alpha_4' + 2\alpha_3'^2}, \frac{\alpha_4'}{2\alpha_3'^2}; \alpha'), \quad i = \overline{3, n-1}$$

hold. Then it is easy to notice that

$$\rho_i(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = \rho_i(\frac{2\alpha_3'}{\alpha_4' + 2\alpha_3'^2}, \frac{\alpha_4'}{2\alpha_3'^2}; \alpha') \text{ for } i = \overline{1, 2}$$

as well and therefore $\rho(\frac{2\alpha_3}{\alpha_4+2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = \rho(\frac{2\alpha'_3}{\alpha'_4+2\alpha'_3^2}, \frac{\alpha'_4}{2\alpha'_3^2}; \alpha')$ that means the algebras $L(\alpha)$ and $L(\alpha')$ are isomorphic to the same algebra and therefore they are isomorphic to each other.

Proof of the part ii). The system of equations

$$\rho_i(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = a_i \quad 3 \le i \le n-1,$$

where $(a_3, a_4, ..., a_{n-1})$ is given and $\alpha = (\alpha_3, \alpha_4, ..., \alpha_{n-1}, \theta)$ is unknown, has a solution as far as for any $3 \leq i \leq n-1$ in $\rho_i(\frac{2\alpha_3}{\alpha_4+2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha)$ $(\rho_{n-1}(\frac{2\alpha_3}{\alpha_4+2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha))$ only variables $\alpha_3, \alpha_4, ..., \alpha_i$ (respectively, $\alpha_3, \alpha_4, ..., \alpha_{n-1}, \theta$) occur and each of these equations is a linear equation with respect to the last variable occurred in it. This is the end of the proof.

Here are the corresponding list of invariants for some low dimensional cases. Case of n = 4 i.e. dim L=5:

$$\left\{ \rho_3(\frac{2\alpha_3}{\alpha_4+2\alpha_3^2},\frac{\alpha_4}{2\alpha_3^2};\alpha) = (\frac{2\alpha_3}{\alpha_4+2\alpha_3^2})^2(\theta-\alpha_4). \right.$$

Case of n = 5 i.e. dim L=6:

$$\begin{cases} \rho_3(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = (\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2})^2 \frac{\alpha_5 + 5\alpha_3\alpha_4 + 5\alpha_3^3}{\alpha_3} - 5. \\ \\ \rho_4(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = (\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2})^3 (\theta - \alpha_5) + (\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2})^2 \frac{\alpha_5 + 5\alpha_3\alpha_4 + 5\alpha_3^3}{\alpha_3} - 5. \end{cases}$$

Case of n = 6 i.e. dim L=7:

$$\rho_{3}\left(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}},\frac{\alpha_{4}}{2\alpha_{3}^{2}};\alpha\right) = \left(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}}\right)^{2}\frac{\alpha_{5}+5\alpha_{3}\alpha_{4}+5\alpha_{3}^{3}}{\alpha_{3}} - 5.$$

$$\rho_{4}\left(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}},\frac{\alpha_{4}}{2\alpha_{3}^{2}};\alpha\right) = \left(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}}\right)^{3}\frac{\alpha_{6}+6\alpha_{3}\alpha_{5}+21\alpha_{3}^{2}\alpha_{4}+3\alpha_{4}^{2}+14\alpha_{3}^{4}}{\alpha_{3}} - \left(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}}\right)^{2}\frac{6\alpha_{3}\alpha_{5}+42\alpha_{3}^{2}\alpha_{4}+3\alpha_{4}^{2}+42\alpha_{3}^{4}}{\alpha_{3}^{2}} + 28.$$

$$\rho_{5}\left(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}},\frac{\alpha_{4}}{\alpha_{2}^{2}};\alpha\right) = \left(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}}\right)^{4}(\theta - \alpha_{6}) + \left(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}}\right)^{3}\left(\frac{\alpha_{6}+6\alpha_{3}\alpha_{5}+21\alpha_{3}^{2}\alpha_{4}+3\alpha_{4}^{2}+14\alpha_{3}^{4}}{\alpha_{3}^{2}}\right)^{3}$$

$$\rho_5(\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2}, \frac{\alpha_4}{2\alpha_3^2}; \alpha) = (\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2})^4(\theta - \alpha_6) + (\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2})^3(\frac{\alpha_6 + 6\alpha_3\alpha_5 + 21\alpha_3^2\alpha_4 + 3\alpha_4^2 + 14\alpha_3^4}{\alpha_3}) - (\frac{2\alpha_3}{\alpha_4 + 2\alpha_3^2})^2\frac{6\alpha_3\alpha_5 + 42\alpha_3^2\alpha_4 + 3\alpha_4^2 + 42\alpha_3^4}{\alpha_3^2} + 28.$$

Case of n = 7 i.e. dim L=8:

$$\begin{split} \rho_{3}(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}},\frac{\alpha_{4}}{2\alpha_{3}^{2}};\alpha) &= (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{2} \frac{\alpha_{5}+5\alpha_{3}\alpha_{4}+5\alpha_{3}^{3}}{\alpha_{3}} - 5. \\ \rho_{4}(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}},\frac{\alpha_{4}}{2\alpha_{3}^{2}};\alpha) &= (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{3} \frac{\alpha_{6}+6\alpha_{3}\alpha_{5}+21\alpha_{3}^{2}\alpha_{4}+3\alpha_{4}^{2}+14\alpha_{3}^{4}}{\alpha_{3}} \\ &- (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{2} \frac{6\alpha_{3}\alpha_{5}+42\alpha_{3}^{2}\alpha_{4}+3\alpha_{4}^{2}+42\alpha_{3}^{4}}{\alpha_{3}^{2}} + 28. \\ \rho_{5}(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}},\frac{\alpha_{4}}{2\alpha_{3}^{2}};\alpha) &= (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{4} \frac{\alpha_{7}+7\alpha_{3}\alpha_{6}+28\alpha_{3}\alpha_{4}^{2}+28\alpha_{3}^{2}\alpha_{5}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}\alpha_{4}+42\alpha_{5}^{5}}{\alpha_{3}} \\ &- (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{3} \frac{7\alpha_{3}\alpha_{6}+56\alpha_{3}\alpha_{4}^{2}+56\alpha_{3}^{2}\alpha_{5}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}\alpha_{4}+42\alpha_{5}^{5}}{\alpha_{3}^{2}} \\ &+ (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{2} \frac{28(\alpha_{4}^{2}+\alpha_{3}\alpha_{5}+9\alpha_{3}^{2}\alpha_{4}+9\alpha_{4}^{4})}{\alpha_{3}^{2}} - 126. \\ \rho_{6}(\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}},\frac{\alpha_{4}}{2\alpha_{3}^{2}};\alpha) &= (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{5}(\theta-\alpha_{7}) \\ &+ (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{4} \frac{\alpha_{7}+7\alpha_{3}\alpha_{6}+28\alpha_{3}\alpha_{4}^{2}+28\alpha_{3}^{2}\alpha_{5}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}\alpha_{4}+42\alpha_{3}^{5}}{\alpha_{3}} \\ &- (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{3} \frac{7\alpha_{3}\alpha_{6}+56\alpha_{3}\alpha_{4}^{2}+28\alpha_{3}^{2}\alpha_{5}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}\alpha_{4}+42\alpha_{3}^{5}}{\alpha_{3}} \\ &- (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{3} \frac{7\alpha_{3}\alpha_{6}+56\alpha_{3}\alpha_{4}^{2}+28\alpha_{3}^{2}\alpha_{5}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}\alpha_{4}+42\alpha_{3}^{5}}{\alpha_{3}} \\ &- (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{3} \frac{7\alpha_{3}\alpha_{6}+56\alpha_{3}\alpha_{4}^{2}+56\alpha_{3}^{2}\alpha_{5}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}\alpha_{4}+42\alpha_{3}^{5}}{\alpha_{3}} \\ &- (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{3} \frac{7\alpha_{3}\alpha_{6}+56\alpha_{3}\alpha_{4}^{2}+56\alpha_{3}^{2}\alpha_{5}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}\alpha_{4}+42\alpha_{3}^{5}}{\alpha_{3}} \\ &- (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{3} \frac{7\alpha_{3}\alpha_{6}+56\alpha_{3}\alpha_{4}^{2}+56\alpha_{3}^{2}\alpha_{5}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}\alpha_{4}+42\alpha_{3}^{5}}{\alpha_{3}} \\ &- (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{3} \frac{7\alpha_{3}\alpha_{6}+56\alpha_{3}\alpha_{4}^{2}+56\alpha_{3}^{2}\alpha_{5}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}\alpha_{4}+42\alpha_{3}^{5}}{\alpha_{3}} \\ &- (\frac{2\alpha_{3}}{\alpha_{4}+2\alpha_{3}^{2}})^{3} \frac{7\alpha_{3}\alpha_{6}+6\alpha_{6}\alpha_{4}+2\alpha_{3}^{2}}{\alpha_{5}}^{2}+7\alpha_{4}\alpha_{5}+84\alpha_{3}^{3}$$

The following two theorems deal with the isomorphism problem for elements from the closed set F. The proof of these theorems are similar to the proof of Theorem 3. The set F in it's turn can be represented as a union of two open and one closed subsets: $F = U'_1 \cup U'_2 \cup F'$, where $U'_1 = \{L(\alpha) \in F : \alpha_3 \neq 0 \text{ and } \alpha_4 + 2\alpha_3^2 = 0\}, U'_2 = \{L(\alpha) \in F : \alpha_3 = 0 \text{ and } \alpha_4 + 2\alpha_3^2 \neq 0\}$, and $F' = \{L(\alpha) \in F : \alpha_3 = 0 \text{ and } \alpha_4 + 2\alpha_3^2 = 0\}$.

Then represent U'_1 and U'_2 in the form

$$U_1' = U_1'' \bigcup F_1'',$$

where $U_1'' = \{ \alpha \in U_1' : (\alpha_5 - 5\alpha_3^3)(\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4) \neq 0 \}$, and $F_1'' = \{ \alpha \in U_1' : (\alpha_5 - 5\alpha_3^3)(\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4) = 0 \}$, and

$$U_2' = U_2'' \bigcup F_2'',$$

where

$$U_2'' = \{ \alpha \in U_2' : \alpha_5 \neq 0 \},\$$

$$F_2'' = \{ \alpha \in U_2' : \alpha_5 = 0 \},\$$

Theorem 4. Let $n \ge 6$. Then two algebras $L(\alpha)$ and $L(\alpha')$ from U''_1 are isomorphic if and only if

$$\rho_i(\frac{5\alpha_3^3 - \alpha_5}{\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4}, \frac{\alpha_6 + 7\alpha_3\alpha_5 - 21\alpha_3^4}{\alpha_3(5\alpha_3^3 - \alpha_5)}; \alpha) = \rho_i(\frac{5\alpha_3'^3 - \alpha_5'}{\alpha_6' + 6\alpha_3'\alpha_5' - 16\alpha_3'^4}, \frac{\alpha_6' + 7\alpha_3'\alpha_5' - 21\alpha_3'^4}{\alpha_3'(5\alpha_3'^3 - \alpha_5')}; \alpha')$$

for i = 4, n - 1. Moreover,

$$\rho_4(\frac{5\alpha_3^3 - \alpha_5}{\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4}, \frac{\alpha_6 + 7\alpha_3\alpha_5 - 21\alpha_3^4}{\alpha_3(5\alpha_3^3 - \alpha_5)}; \alpha) \neq -14.$$

and for any $(a_4, ..., a_{n-1}) \in C^{n-4}, a_4 \neq -14$ there is an algebra $L(\alpha)$ from U''_1 such that

$$\rho_i(\frac{5\alpha_3^3 - \alpha_5}{\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4}, \frac{\alpha_6 + 7\alpha_3\alpha_5 - 21\alpha_3^4}{\alpha_3(5\alpha_3^3 - \alpha_5)}; \alpha) = a_i \text{ for all } i = \overline{4, n-1}$$

Here are the list of invariants for n = 6 and n = 7 cases:

$$n=6$$

$$\begin{cases} \rho_4(\frac{5\alpha_3^3-\alpha_5}{\alpha_6+6\alpha_3\alpha_5-16\alpha_3^4},\frac{\alpha_6+7\alpha_3\alpha_5-21\alpha_3^4}{\alpha_3(5\alpha_3^3-\alpha_5)};\alpha) = 7\frac{(5\alpha_3^3-\alpha_5)^3}{\alpha_3(\alpha_6+6\alpha_3\alpha_5-16\alpha_3^4)^2} - 14.\\ \rho_5(\frac{5\alpha_3^3-\alpha_5}{\alpha_6+6\alpha_3\alpha_5-16\alpha_3^4},\frac{\alpha_6+7\alpha_3\alpha_5-21\alpha_3^4}{\alpha_3(5\alpha_3^3-\alpha_5)};\alpha) = (\frac{5\alpha_3^3-\alpha_5}{\alpha_6+6\alpha_3\alpha_5-16\alpha_3^4})^4(\theta-\alpha_6) + 7\frac{(5\alpha_3^3-\alpha_5)^3}{\alpha_3(\alpha_6+6\alpha_3\alpha_5-16\alpha_3^4)^2} - 14.\\ \mathbf{n=7:}\end{cases}$$

$$\begin{cases} \rho_4 \left(\frac{5\alpha_3^3 - \alpha_5}{\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4}, \frac{\alpha_6 + 7\alpha_3\alpha_5 - 21\alpha_3^4}{\alpha_3(5\alpha_3^3 - \alpha_5)}; \alpha\right) = 7 \frac{(5\alpha_3^3 - \alpha_5)^3}{\alpha_3(\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4)^2} - 14. \\ \rho_5 \left(\frac{5\alpha_3^3 - \alpha_5}{\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4}, \frac{\alpha_6 + 7\alpha_3\alpha_5 - 21\alpha_3^4}{\alpha_3(5\alpha_3^3 - \alpha_5)}; \alpha\right) = \left(\frac{5\alpha_3^3 - \alpha_5}{\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4}\right)^4 \frac{\alpha_7 + 7\alpha_3\alpha_6 - 14\alpha_3^2\alpha_5 - 14\alpha_3^5}{\alpha_3} \\ -35 \frac{(5\alpha_3^3 - \alpha_5)^3}{\alpha_3(\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4)^2} + 42. \\ \rho_6 \left(\frac{5\alpha_3^3 - \alpha_5}{\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4}, \frac{\alpha_6 + 7\alpha_3\alpha_5 - 21\alpha_3^4}{\alpha_3(5\alpha_3^3 - \alpha_5)}; \alpha\right) = \left(\frac{5\alpha_3^3 - \alpha_5}{\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4}\right)^5 (\theta - \alpha_7) \\ + \left(\frac{5\alpha_3^3 - \alpha_5}{\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4}\right)^4 \frac{\alpha_7 + 7\alpha_3\alpha_6 - 14\alpha_3^2\alpha_5 - 14\alpha_3^5}{\alpha_3} - 35 \frac{(5\alpha_3^3 - \alpha_5)^3}{\alpha_3(\alpha_6 + 6\alpha_3\alpha_5 - 16\alpha_3^4)^2} + 42. \end{cases}$$

Remark. An analog of Theorem 4 can be stated for n = 4 and n = 5 cases as well. But in these cases it is specific and therefore is not considered here.

Theorem 5. Let $n \ge 5$. Then two algebras $L(\alpha)$ and $L(\alpha')$ from U''_2 are isomorphic if and only if

$$\rho_i(\frac{\alpha_4}{\alpha_5}, \frac{\alpha_5^2}{\alpha_3^3} - 1; \alpha) = \rho_i(\frac{\alpha_4'}{\alpha_5'}, \frac{\alpha_5'^2}{\alpha_3'^3} - 1; \alpha')$$

for $\overline{i=4,n-1}$,

Moreover, for any $(a_4, ..., a_{n-1}) \in C^{n-4}$ there is an algebra $L(\alpha)$ from U''_2 such that

$$\rho_i(\frac{\alpha_4}{\alpha_5}, \frac{\alpha_5^2}{\alpha_3^3} - 1; \alpha) = a_i$$

for all $i = \overline{4, n-1}$.

Here are the corresponding list of invariants for n = 5, 6 and n = 7 cases: Case of n = 5 i.e. **dim L=6**:

$$\left\{ \rho_4\left(\frac{\alpha_4}{\alpha_5}, \frac{\alpha_5^2 - \alpha_4^3}{\alpha_4^3}; \alpha\right) = \frac{\alpha_4\left(\alpha_4^2\theta - \alpha_4^3\alpha_3 - 3\alpha_5^3\right)}{\alpha_5^3}.\right.$$

Case of n = 6 i.e. dim L=7:

$$\begin{cases} \rho_4(\frac{\alpha_4}{\alpha_5}, \frac{\alpha_5^2 - \alpha_4^3}{\alpha_4^3}; \alpha) = \frac{\alpha_4(\alpha_6 + 3\alpha_4^2)}{\alpha_5^2} - 3; \\ \\ \rho_5(\frac{\alpha_4}{\alpha_5}, \frac{\alpha_5^2 - \alpha_4^3}{\alpha_4^3}; \alpha) = (\frac{\alpha_4}{\alpha_5})^4(\theta - \alpha_6) + \frac{\alpha_4(\alpha_6 + 3\alpha_4^2)}{\alpha_5^2} - 3. \end{cases}$$

Case of n = 7 i.e. dim L=8:

$$\begin{cases} \rho_4(\frac{\alpha_4}{\alpha_5}, \frac{\alpha_5^2 - \alpha_4^3}{\alpha_4^3}; \alpha) = \frac{\alpha_4(\alpha_6 + 3\alpha_4^2)}{\alpha_5^2} - 3; \\ \\ \rho_5(\frac{\alpha_4}{\alpha_5}, \frac{\alpha_5^2 - \alpha_4^3}{\alpha_4^3}; \alpha) = \frac{\alpha_4^2(\alpha_7 + 7\alpha_4\alpha_5)}{\alpha_5^3} - 7; \\ \\ \\ \rho_6(\frac{\alpha_4}{\alpha_5}, \frac{\alpha_5^2 - \alpha_4^3}{\alpha_4^3}; \alpha) = (\frac{\alpha_4}{\alpha_5})^5(\theta - \alpha_7) + \frac{\alpha_4^2(\alpha_7 + 7\alpha_4\alpha_5)}{\alpha_5^3} - 7. \end{cases}$$

For any given low dimensional case the above suggested approach enable us to get the complete classification of filiform Leibniz algebras from the first class. It is hoped that we can present it in the near future.

Acknowledgement. We would like to thank J.R.Gomez and B.A.Omirov for their kind permission to use their unpublished result.

References

- J.R.Gomez, B.A.Omirov. On classification of complex filiform Leibniz Algebras (to appear).
- [2] J.-L.Loday. Une version non commutative des de Lie: les de Leibniz. L'Ens. Math., 39(1993), p.269-293.
- [3] H.Kraft, Geometrische Methoden in der Invarianten theorie, Aspekte der Mathematik, Vol. D1, Vieweg Verlag, Braunschweig-Wiesbaden, 1985. 2., durchgeschene Auflage.

- [4] S.Albeverio, B.A.Omirov, I.S.Rakhimov. Varieties of Nilpotent Complex Leibniz Algebras of Dimension Less than Five. Comm. in Algebra, vol. 33, 5, 2005, p. 1575-1585.
- [5] A.Malcev. On Solvable Lie algebras, Amer.Math.Soc.Transl.(1)9 (1962),228-262.
- [6] K.A.Umlauf, Uber die Zusammensetzung der endlichen continuierlichen Transformationsgrouppen insbesondere der Gruppen vom Range null, Thesis, Leipzig, 1891.
- [7] M.Vergne, Cohomologie des algebres de Lie nilpotentes. Application à l'étude de la variété des algèbres de Lie nilpotentes, Bull. Soc. Math. France 98(1970), 81-116.
- [8] J.R.Gomes, A.Jimenez-Merchan, Y.Khakimdjanov, Low-dimensional filiform Lie algebras, J. of Pure and Applied Algebra 130(1998), 133-158.
- Sh.A.Ayupov, B.A.Omirov, On some classes of nilpotent Leibniz algebras. // Sib. Math. J. (2001). V. 42, 1. 18-29.(in Russian)
- [10] H.Weyl, The classical groups, Princeton Univ. Press, Princeton, N.J., 1946.
- [11] V.L.Popov, E.B.Vinberg. Invariant theory. In: Encyclopaedia of Math.Sci.: Algebraic Geometry IV. Springer Verlag 55 (1994), 123-284.
- [12] Omirov B.A. Structural theory of nilpotent Leibniz algebras. PhD thesis, Institute of Mathematics of Uzbekistan Academy of Science, Uzbekistan (1999).
- [13] Albeverio S., Omirov B.A., Rakhimov I.S. Classification of four-dimensional nilpotent complex Leibniz algebras. Rheinische Friedrich-Wilhelms-Universitat Bonn, preprint, 2003, 12 p.(Appears in Extracta Mathematicae).