

ON COMPLETE SYSTEM OF COVARIANTS FOR THE BINARY FORM OF DEGREE 7

LEONID BEDRATYUK

ABSTRACT. A minimal system of 147 homogeneous generating elements of the algebra of covariants for the binary form of degree 7 is calculated.

1. INTRODUCTION

Let V_n be a vector $\mathbb{C}$ -space of the binary forms of degree d considered with natural action of the group $G = SL(2, \mathbb{C})$. Let us extend the action of the group G to the polynomial functions algebra $\mathbb{C}[V_d \oplus \mathbb{C}^2]$. Denote by $C_d = \mathbb{C}[V_d \oplus \mathbb{C}^2]^G$ the corresponding subalgebra of G -invariant functions. In the vocabulary of classical invariant theory the algebra C_d is called the algebra covariants of the binary form of d -th degree. Let C_d^+ be an ideal of C_d generated by all homogeneous elements of positive power. Denote by $\bar{C}_d$ a set of homogeneous elements of C_d^+ such that their images in $C_d^+/(C_d^+)^2$ form a basis of the vector space. The set $\bar{C}_d$ is called complete system of covariants of the d -th degree binary form. Elements of $\bar{C}_d$ form a minimal system of homogeneous generating elements of the invariants algebra C_d . Denote by c_d a number of elements of the set $\bar{C}_d$.

The complete systems of covariants was a topic of major research interest in classical invariant theory of the 19th century. It is easy to show that $c_1 = 0$, $c_2 = 2$, $c_3 = 4$. A complete system of covariants in the case $d = 4$ was calculated by Bool, Cayley, Eisenstein, see survey [1]. The complete systems of invariants and covariants in the cases $d = 5, 6$ were calculated by Gordan, see [2]. In particular, $c_4 = 5$, $c_5 = 23$, $c_6 = 26$.

Gall's attempt [3] to discover the complete system for the case $d = 7$ was unsuccessful. He did offered a system of 151 covariants but the system was not a minimal system, see [9], [16]. Also, Sylvester's attempts [6] to find the cardinality c_7 and covariant's degree-order distribution of the complete system were mistaken, see [9]. Therefore, the problem of finding a minimal system of homogeneous generating elements (or even a cardinality of the system) of the algebra of covariants for the binary form of degree 7 was open.

The case $d = 8$ was studied by Sylvester and Gall but they have obtained completely different results. By using Sylvester-Cayley technique, Sylvester in [6] got that $c_8 = 69$. Gall in [7], evolving the Gordan's constructive method, offered 68 covariants as a complete system of covariants for the case $d = 8$. In [17] the author have calculated the set $\bar{C}_8$.

For the cases $d = 9, 10$ Sylvester in [6] calculated the cardinalities c_9 and c_{10} but the present author, using a computer, found numerous mistakes in those computations.

Therefore, the complete systems of covariants for the binary form are so far known only for $d \leq 6$, see [1], [5], and for $d = 8$, see [17].

This article's purpose is calculating the set $\bar{C}_7$. Using a technique introduced in [16] we explicitly calculate a complete system of covariants of the 7th degree binary form. In particular, we prove that $c_7 = 147$.

All calculation were done with Maple.

2. PRELIMINARIES.

Before any calculation we try make a simplification of a covariant representation and their computation.

Let us indentify the algebra $\mathbb{C}[V_d]$ with the algebra $\mathbb{C}[X_d] := \mathbb{C}[t, x_1, x_2, \dots, x_d]$, and the algebra $\mathbb{C}[V_d \oplus \mathbb{C}^2]$ identify with the polynomial algebra $\mathbb{C}[t, x_1, x_2, \dots, x_n, Y_1, Y_2]$. Let

$$\kappa : C_d \longrightarrow \mathbb{C}[X_d]$$

be the $\mathbb{C}$ -linear map takes each homogeneous covariant of order k to his leading coefficient, i.e. a coefficient of Y_1^k . Follow by classical tradition an element of the algebra $\kappa(C_d)$ is called *semi-invariants*, a degree of a homogeneous covariant with respect to the variables set X_d is called *degree* of the covariant and its degree with respect to the variables set Y_1, Y_2 is called *order*.

In [16] showed that

$$\kappa(C_d) = k[t, z_2, z_3, \dots, z_d] \left[\frac{1}{t} \right] \cap \mathbb{C}[X_d],$$

$$z_i := \sum_{k=0}^{i-2} (-1)^k \binom{i}{k} x_{i-k} x_1^k t^{i-k-1} + (i-1)(-1)^{i+1} x_1^i, i = 2, \dots, d.$$

Let us consider the following derivation D of the algebra $k[t, x_1, z_2, z_3, \dots, z_d] \left[\frac{1}{t} \right]$:

$$\begin{aligned} D = & 7x_1 \frac{\partial}{\partial t} - \frac{(-15x_1 z_3 + 18z_2^2 - 4z_4)}{t} \frac{\partial}{\partial z_3} + \frac{(20x_1 z_4 - 24z_2 z_3 + 3z_5)}{t} \frac{\partial}{\partial z_4} + \\ & + \frac{(2z_6 + 25x_1 z_5 - 30z_2 z_4)}{t} \frac{\partial}{\partial z_5} + \frac{(z_7 + 30x_1 z_6 - 36z_2 z_5)}{t} \frac{\partial}{\partial z_6} + \\ & + \frac{7(5x_1 z_7 - 6z_2 z_6)}{t} \frac{\partial}{\partial z_7} + \frac{5(2x_1 z_2 + z_3)}{t} \frac{\partial}{\partial z_2}. \end{aligned}$$

Suppose $F = \sum_{i=0}^m f_i \binom{m}{i} Y_1^{m-i} Y_2^i$ be a covariant of order m , $\kappa(F) = f_0 \in \kappa(C_d)$. The classical Robert's theorem [10], states that the covariant F is completely and uniquely determined by its leading coefficient f_0 , namely, see [16] :

$$F = \sum_{i=0}^m \frac{D^i(f_0)}{i!} Y_1^{m-i} Y_2^i.$$

On the other hand, every semi-invariant is a leading coefficient of a covariant, see [13], [5]. This give us well defined explicit form of the inverse map

$$\kappa^{-1} : \kappa(C_d) \longrightarrow C_d,$$

namely

$$\kappa^{-1}(a) = \sum_{i=0}^{\text{ord}(a)} \frac{D^i(a)}{i!} Y_1^{\text{ord}(a)-i} Y_2^i,$$

here $a \in \kappa(C_d)$ and $\text{ord}(a)$ is an order of the element a with respect to the locally nilpotent (at $\kappa(C_d)$) derivation D , i.e. $\text{ord}(a) := \max\{s, D^s(a) \neq 0\}$. For example, since $\text{ord}(t) = d$, we have

$$\kappa^{-1}(t) = \sum_{i=0}^{\text{ord}(t)} \frac{D_2^i(t)}{i!} Y_1^{\text{ord}(t)-i} Y_2^i = t Y_1^d + \sum_{i=1}^d \binom{d}{i} x_i Y_1^{d-i} Y_2^i.$$

As we see the $\kappa^{-1}(t)$ is just the basic binary form. From polynomial functions point of view the covariant $\kappa^{-1}(t)$ is the evaluation map.

Thus, the problem of finding of complete system of the algebra $\overline{C}_d$ is equivalent to the problem of finding of complete system of semi-covariants's algebra $\kappa(C_d)$. It is well known classical results, see, for example, [4].

To calculate the semi-invariants we need have an analogue of the transvectants. Let f, g are two covariants of the degrees m and k respectively. The semi-invariant of the form

$$[f, g]^r := q_r(f, g) \cdot \sum_{i=0}^r (-1)^i \binom{r}{i} \frac{D^i(f)}{[m]_i} \Big|_{x_1=0, \dots, x_r=0} \frac{D^{r-i}(g)}{[k]_{r-i}} \Big|_{x_1=0, \dots, x_r=0}, \quad 0 \leq r \leq \min(m, k),$$

is called a r -th semitransvectant of the semi-invariants f and g . Here $q_r(f, g)$ is a normalization rational factor, see [16]. Note that up to multiplication on a rational number the semitransvectant $[f, g]^r$ is equal to $\kappa(\kappa^{-1}(f), \kappa^{-1}(g))^r$, where $(\cdot, \cdot)^r$ is usual transvectant, see [12], [5]. The following statements is direct consequences of corresponding transvectant properties, see [13]:

Lemma. *Let f, g be two semi-invariants. Then the following conditions hold*

- (i) *the semitransvectant $[t, f g]^i$ is reducible for $0 \leq i \leq \min(d, \max(\text{ord}(f), \text{ord}(g)))$;*
- (ii) *if $\text{ord}(f) = 0$, then $[t, f g]^i = f[t, g]^i$;*
- (iii) *$\text{ord}([f, g]^i) = \text{ord}(f) + \text{ord}(g) - 2i$;*
- (iv) *$\text{ord}(z_2^{i_1} z_3^{i_3} \cdots z_d^{i_d}) = d(i_2 + i_3 + \cdots + i_d) - 2(2i_2 + 3i_3 + \cdots + di_d)$.*

The representation of semi-invariants as elements of the algebra $\mathbb{C}[t, z_2, z_3, \dots, z_d][\frac{1}{t}]$ is more compact than their standard representation as elements of the algebra $\mathbb{C}[X_d]$. Very rough empire estimate is that a semi-invariant $\kappa(F)$ has terms number in $([\deg(F)/d] + 2)(\text{ord}(F) + 1)$ times less than the corresponding covariant F . Moreover, from computing point of view, the semitransvectant formula is more effective than the transvectant formula. These two favorable circumstances coupled with great Maple powers allow us to compute a complete system of covariants of the 7-th degree binary form.

3. IRREDUCIBLE COVARIANTS UP TO 13TH DEGREE

Let $\overline{C}_{7,i} := (C_7)_i$ be a subset of $\overline{C}_7$ whose elements has degree i and $\overline{C}_{7,i} := \overline{C}_7 \cap C_{7,i}$. By using an analogue of the Ω -process and the lemma, in [16] was found all irreducible covariants up to 13th degree. Let us present the lists of the generating elements.

The unique semi-invariant of the degree one obviously is t , $\text{ord}(t) = 7$.

The set $\overline{C}_{7,2}$, consists of the 3 irreducible semi-invariants

$$\begin{aligned} dv_1 &:= [t, t]^4 = \frac{3z_2^2 + z_4}{t^2} = x_4 t - 4x_1 x_3 + 3x_2^2, & \text{ord}(dv_1) &= 6, \\ dv_2 &:= [t, t]^6 = \frac{z_6 + 15z_2 z_4 - 10z_3^2}{t^4} = x_6 t - 6x_1 x_5 + 15x_2 x_4 - 10x_3^2, & \text{ord}(dv_2) &= 2, \\ dv_3 &:= [t, t]^2 = z_2 = x_2 t - x_1^2, & \text{ord}(dv_2) &= 10. \end{aligned}$$

The set $\overline{C}_{7,3}$, consists of the 6 irreducible semi-invariants

$$\begin{aligned} tr_1 &= [t, dv_1]^4, & \text{ord}(tr_1) &= 5, & tr_2 &= [t, dv_3], & \text{ord}(tr_2) &= 15, \\ tr_3 &= [t, dv_3]^3, & \text{ord}(tr_3) &= 11, & tr_4 &= [t, dv_3]^4, & \text{ord}(tr_4) &= 9, \\ tr_5 &= [t, dv_3]^5, & \text{ord}(tr_5) &= 7, & tr_6 &= [t, dv_3]^7, & \text{ord}(tr_6) &= 3. \end{aligned}$$

The set $\overline{C}_{7,4}$ consists of the following 8 irreducible semi-invariants:

$$\begin{aligned} ch_1 &= [t, tr_5]^7, & \text{ord}(ch_1) &= 0, & ch_2 &= [t, tr_3]^7, & \text{ord}(ch_2) &= 4, \\ ch_3 &= [t, tr_3]^2, & \text{ord}(ch_3) &= 14, & ch_4 &= [t, tr_3]^4, & \text{ord}(ch_4) &= 10, \\ ch_5 &= [t, tr_3]^5, & \text{ord}(tr_5) &= 8, & ch_6 &= [t, tr_1]^2, & \text{ord}(tr_6) &= 8, \\ ch_7 &= [t, tr_1]^3, & \text{ord}(ch_7) &= 6, & ch_8 &= [t, tr_1]^4, & \text{ord}(ch_6) &= 4. \end{aligned}$$

The set $\overline{C}_{7,5}$ consists of the following 10 irreducible semi-invariants:

$$\begin{aligned} pt_1 &= [t, ch_6]^5, & \text{ord}(pt_1) &= 5, & pt_2 &= [t, ch_6]^6, & \text{ord}(pt_2) &= 3, \\ pt_3 &= [t, ch_7]^2, & \text{ord}(pt_3) &= 9, & pt_4 &= [t, ch_7]^3, & \text{ord}(pt_4) &= 7, \\ pt_5 &= [t, ch_7]^5, & \text{ord}(pt_5) &= 3, & pt_6 &= [t, ch_6]^3, & \text{ord}(pt_6) &= 9, \\ pt_7 &= [t, ch_4]^2, & \text{ord}(pt_7) &= 13, & pt_8 &= [t, ch_4]^5, & \text{ord}(pt_8) &= 7, \\ pt_9 &= [t, dv_1^2]^7, & \text{ord}(pt_9) &= 5, & pt_{10} &= [t, dv_1 dv_2]^7, & \text{ord}(pt_{10}) &= 1. \end{aligned}$$

The set $\overline{C}_{7,6}$ consists of the following 10 irreducible semi-invariants:

$$\begin{aligned} sh_1 &= [t, pt_5]^5, & \text{ord}(sh_1) &= 6, & sh_2 &= [t, pt_7]^6, & \text{ord}(sh_2) &= 8, \\ sh_3 &= [t, pt_4]^5, & \text{ord}(sh_3) &= 4, & sh_4 &= [t, pt_4]^6, & \text{ord}(sh_4) &= 2, \\ sh_5 &= [t, pt_3]^2, & \text{ord}(sh_5) &= 12, & sh_6 &= [t, pt_3]^4, & \text{ord}(sh_6) &= 8, \\ sh_7 &= [t, pt_4]^4, & \text{ord}(sh_7) &= 6, & sh_8 &= [t, tr_1 dv_1]^7, & \text{ord}(sh_8) &= 4, \\ sh_9 &= [t, tr_1 dv_2]^6, & \text{ord}(sh_9) &= 2, & sh_{10} &= [t, tr_6 dv_1]^7, & \text{ord}(sh_{10}) &= 2. \end{aligned}$$

The set $\overline{C}_{7,7}$ consists of the following 12 irreducible semi-invariants:

$$\begin{aligned}
si_1 &= [t, sh_5]^4, & \text{ord}(si_1) &= 11, & si_2 &= [t, sh_7]^4, & \text{ord}(si_2) &= 5, \\
si_3 &= [t, tr_1^2]^7, & \text{ord}(si_3) &= 3, & si_4 &= [t, sh_1]^3, & \text{ord}(si_4) &= 7, \\
si_5 &= [t, ch_7 dv_1]^7, & \text{ord}(si_5) &= 5, & si_6 &= [t, ch_7 dv_2]^7, & \text{ord}(si_6) &= 1, \\
si_7 &= [t, tr_6^2]^4, & \text{ord}(si_7) &= 5, & si_8 &= [t, tr_6^2]^6, & \text{ord}(si_8) &= 1, \\
si_9 &= [t, tr_6 tr_1]^6, & \text{ord}(si_9) &= 3, & si_{10} &= [t, tr_6 tr_1]^7, & \text{ord}(si_{10}) &= 1, \\
si_{11} &= [t, tr_1^2]^6, & \text{ord}(si_{11}) &= 5, & si_{12} &= [t, sh_{10}], & \text{ord}(si_{12}) &= 7.
\end{aligned}$$

The set $\overline{C}_{7,8}$ consists of the following 13 irreducible semi-invariants:

$$\begin{aligned}
vi_1 &= [t, si_7]^3, & \text{ord}(vi_1) &= 6, & vi_2 &= [t, si_7]^4, & \text{ord}(vi_2) &= 4, \\
vi_3 &= [t, ch_8 tr_6]^7, & \text{ord}(vi_3) &= 0, & vi_4 &= [t, ch_8 tr_1]^6, & \text{ord}(vi_4) &= 4, \\
vi_5 &= [t, ch_8 tr_1]^7, & \text{ord}(vi_5) &= 2, & vi_6 &= [t, ch_7 tr_6]^7, & \text{ord}(vi_6) &= 2, \\
vi_7 &= [t, ch_7 tr_1]^7, & \text{ord}(vi_7) &= 4, & vi_8 &= [t, ch_8 tr_6]^6, & \text{ord}(vi_8) &= 2, \\
vi_9 &= [t, tr_6 dv_2]^7, & \text{ord}(vi_9) &= 0, & vi_{10} &= [t, si_4]^2, & \text{ord}(vi_{10}) &= 10, \\
vi_{11} &= [t, si_1^2]^4, & \text{ord}(vi_{11}) &= 6, & vi_{12} &= [t, si_{11}]^3, & \text{ord}(vi_{12}) &= 6, \\
& & & & vi_{13} &= [t, pt_9 dv_2]^7, & \text{ord}(vi_{13}) &= 0.
\end{aligned}$$

The set $\overline{C}_{7,9}$ consists of the following 11 irreducible semi-invariants:

$$\begin{aligned}
de_1 &= [t, sh_3 dv_1]^7, & \text{ord}(de_1) &= 3, & de_2 &= [t, ch_7 ch_8]^7, & \text{ord}(de_2) &= 3, \\
de_3 &= [t, pt_5 tr_6]^5, & \text{ord}(de_3) &= 3, & de_4 &= [t, pt_5 tr_1]^6, & \text{ord}(de_4) &= 3, \\
de_5 &= [t, pt_5 tr_1]^7, & \text{ord}(de_5) &= 1, & de_6 &= [t, sh_9 dv_1]^7, & \text{ord}(de_6) &= 1, \\
de_7 &= [t, sh_{10} dv_1]^7, & \text{ord}(de_7) &= 1, & de_8 &= [t, sh_{10} dv_2]^3, & \text{ord}(de_8) &= 5, \\
de_9 &= [t, vi_5]^2, & \text{ord}(de_9) &= 5, & de_{10} &= [t, vi_2]^4, & \text{ord}(de_{10}) &= 3, \\
& & & & de_{11} &= [t, vi_{11}]^2, & \text{ord}(de_{11}) &= 9.
\end{aligned}$$

The set $\overline{C}_{7,10}$ consists of the following 9 irreducible semi-invariants:

$$\begin{aligned}
des_1 &= [t, sh_9 tr_1]^6, & \text{ord}(des_1) &= 2, & des_2 &= [t, sh_4 tr_6]^4, & \text{ord}(des_2) &= 4, \\
des_3 &= [t, sh_4 tr_1]^6, & \text{ord}(des_3) &= 2, & des_4 &= [t, sh_1 tr_1]^7, & \text{ord}(des_4) &= 4, \\
des_5 &= [t, sh_3 tr_6]^5, & \text{ord}(des_5) &= 4, & des_6 &= [t, de_9]^2, & \text{ord}(des_6) &= 8, \\
des_7 &= [t, tr_6^3]^7, & \text{ord}(des_7) &= 2, & des_8 &= [t, sh_{10} tr_1]^6, & \text{ord}(des_8) &= 2, \\
& & & & des_9 &= [t, pt_1 ch_7]^7, & \text{ord}(des_9) &= 4.
\end{aligned}$$

The set $\overline{C}_{7,11}$ consists of the following 9 irreducible semi-invariants:

$$\begin{aligned}
odn_1 &= [t, vi_2 dv_1]^7, & \text{ord}(odn_1) &= 3, & odn_2 &= [t, vi_2, dv_2]^6, & \text{ord}(odn_2) &= 1, \\
odn_3 &= [t, vi_4 dv_2]^6, & \text{ord}(odn_3) &= 1, & odn_4 &= [t, vi_5 dv_1]^7, & \text{ord}(odn_4) &= 1, \\
odn_5 &= [t, vi_6 dv_1]^7, & \text{ord}(odn_5) &= 1, & odn_6 &= [t, vi_2 dv_2]^5, & \text{ord}(odn_6) &= 3, \\
odn_7 &= [t, des_6]^4, & \text{ord}(odn_7) &= 7, & odn_8 &= [t, des_6]^6, & \text{ord}(odn_8) &= 3, \\
& & & & odn_9 &= [t, vi_1 dv_2]^7, & \text{ord}(odn_9) &= 1.
\end{aligned}$$

The set $\overline{C}_{7,12}$ consists of the following 13 irreducible semi-invariants:

$$\begin{aligned} dvan_1 &= [t, sh_1 pt_2]^7, & \text{ord}(dvan_1) &= 2, & dvan_2 &= [t, sh_1 pt_5]^7, & \text{ord}(dvan_2) &= 2, \\ dvan_3 &= [sh_9, sh_{10}]^2, & \text{ord}(dvan_3) &= 0, & dvan_4 &= [t, odn_7]^6, & \text{ord}(dvan_4) &= 2, \\ dvan_5 &= [t, de_8 dv_2]^6, & \text{ord}(dvan_5) &= 2, & dvan_6 &= [sh_{10}, sh_{10}]^2, & \text{ord}(dvan_6) &= 0, \\ dvan_7 &= [t, de_9 dv_2]^6, & \text{ord}(dvan_7) &= 2, & dvan_8 &= [t, de_{10} dv_1]^7, & \text{ord}(dvan_8) &= 2, \\ dvan_9 &= [t, odn_7]^4, & \text{ord}(dvan_9) &= 6, & dvan_{10} &= [sh_1, sh_1]^2, & \text{ord}(dvan_{10}) &= 0, \\ dvan_{11} &= [sh_4, sh_4]^2, & \text{ord}(dvan_{11}) &= 0, & dvan_{12} &= [sh_4, sh_9]^2, & \text{ord}(dvan_{12}) &= 0, \\ & & dvan_{13} &= [sh_4, sh_2]^2, & \text{ord}(dvan_{13}) &= 0. \end{aligned}$$

The set $\overline{C}_{7,13}$ consists of the following 9 irreducible semi-invariants:

$$\begin{aligned} tryn_1 &= [t, dvan_9]^6, & \text{ord}(tryn_1) &= 1, & tryn_2 &= [t, vi_1 ch_7]^7, & \text{ord}(tryn_2) &= 5, \\ tryn_3 &= [t, vi_2 ch_8]^7, & \text{ord}(tryn_3) &= 1, & tryn_4 &= [t, vi_2 ch_2]^7, & \text{ord}(tryn_4) &= 1, \\ tryn_5 &= [t, vi_1 ch_8]^7, & \text{ord}(tryn_5) &= 3, & tryn_6 &= [t, vi_5 ch_2]^6, & \text{ord}(tryn_6) &= 1, \\ tryn_7 &= [t, vi_8 ch_8]^6, & \text{ord}(tryn_7) &= 1, & tryn_8 &= [t, vi_8 ch_7]^7, & \text{ord}(tryn_8) &= 1, \\ & & tryn_9 &= [t, vi_4 ch_8]^7, & \text{ord}(tryn_9) &= 1. \end{aligned}$$

The cardinalities of the sets $\overline{C}_{7,i}$ for $i = 1, \dots, 13$ and the covariant's degree-order distributions so far coincide completely with Gall's and Sylvester's results, see [3], [6].

4. COMPUTATION OF COVARIANTS OF DEGREE 14–30

Let $(C_+^2)_i$ be a subset of $(C_7^+)^2$ whose elements has degree i . Denote by $C_{i,j}$ a subspace of $C_{7,i}$ generated by elements of degree i and order j . Denote by δ_i a number of irreducible invariants of degree i and denote by $\delta_{i,j}$ a number of irreducible invariants of degree i and order j . It is evident that $\delta_i = \sum_j \delta_{i,j}$ and $c_7 = \sum_i \delta_i$. To compute the number $\delta_{i,j}$ we use the formula $\delta_{i,j} = \dim C_{i,j} - \dim (C_+^2)_{i,j}$. A dimension of the vector space $C_{i,j}$ is calculated by Cayley-Sylvester formula, see, for example [12], [14] and the dimension is equal to a coefficient of $T^{\frac{d+i-j}{2}}$ in the expansion of the series

$$\frac{(1 - T^{d+1})(1 - T^{d+2}) \dots (1 - T^{d+i})}{(1 - T^2) \dots (1 - T^i)}.$$

A dimension of the vector subspace $(C_+^2)_{i,j}$ of $(C_+^2)_i$ whose elements has order j is calculated by the formula $\dim(C_+^2)_{i,j} = \sigma_{i,j} - \dim S_{i,j}$. Here $\sigma_{i,j}$ is a number of monomial of the vector space $(C_+^2)_{i,j}$ and $S_{i,j}$ is a vector space of $(C_+^2)_{i,j}$ generated by syzygies. A dimension of the vector space $S_{i,j}$ is found by direct Maple computations. A calculation of the numbers $\sigma_{i,j}$ is a simple combinatorial problem. An explicit way of the irreducible semi-invariants is found by linear algebra technique, see details in [16], [17].

The set $\overline{C}_{14,0}$ we take from [16]:

$$chot_1 := [si_8, si_{10}], chot_2 := [si_6, si_{10}], chot_3 := [si_6, si_8], chot_4 := [si_3, si_9]^3.$$

For $j = 2$ we have $\dim C_{14,2} = 30$, $\sigma_{14,2} = 36$, $\dim S_{14,2} = 6$. Thus $\delta_{14,2} = 30 - (36 - 6) = 0$. For $j = 4$ we have $\dim C_{14,4} = 37$, $\sigma_{14,4} = 60$, $\dim S_{14,4} = 25$. Thus $\delta_{14,4} = 37 - (60 - 25) = 2$. After calculation we get the following irreducible semi-invariants of $\overline{C}_{14,2}$:

$$chot_5 := [sh_{10}, vi_2], chot_6 := [sh_9, vi_2].$$

The results of Sylvester's and Gall's for $\delta_{14,4}$ were 0 and 2 respectively. Taking into account the results of [3] we have $\delta_{14,j} = 0$ for $j > 4$. Therefore $\delta_{14} = 6$ and the set $C_{7,14}$ consists of the following 6 semi-invariants – $chot_1, \dots, chot_6$.

There exists only semi-invariants of orders 1 and 3 in the set $\overline{C}_{15}$, see [3]. We have $\dim C_{15,1} = 20$, $\sigma_{15,1} = 17$, $\dim S_{15,1} = 0$. Consequently $\delta_{15,1} = 3$. In the same way we get $\dim C_{15,3} = 42$, $\sigma_{15,3} = 61$, $\dim S_{15,3} = 20$. Thus $\delta_{15,3} = 1$ and $\delta_{15} = 4$. The results of Sylvester's and Gall's $\delta_{15,3}$ were 0 and 1 respectively. After calculation we obtain the 4 irreducible semi-invariants:

$$ptn_1 := [de_{10}, sh_9]^2, ptn_2 := [de_{10}, sh_4]^2, ptn_3 := [de_3, sh_9]^2, ptn_4 := [de_{10}, sh_{10}].$$

There exists only semi-invariants of orders 0 and 2 in the set $\overline{C}_{16}$. The two irreducible invariants were found in [16] -

$$shis_1 := [vi_2, vi_4]^4, shis_2 := [vi_4, vi_7]^4.$$

For $j = 2$ we have $\dim C_{16,2} = 33$, $\sigma_{16,2} = 39$, $\dim S_{16,2} = 9$. Thus $\delta_{16,2} = 3$ and we have $\delta_{16} = 5$. The results of Sylvester's and Gall's for $\delta_{16,2}$ were 0 and 3 respectively. After calculation we obtain the rest 3 semi-transvectants of the order 2:

$$shis_3 = [vi_5, vi_2]^2, shis_4 = [vi_8, vi_2]^2, shis_5 = [des_7, sh_{10}].$$

There exists only semi-invariants of order 1 in $\overline{C}_{17}$. We have $\dim C_{17,1} = 31$, $\sigma_{17,1} = 29$, $\dim S_{17,1} = 0$. Therefore $\delta_{17} = \delta_{17,1} = 2$. Sylvester and Gall have got the same value for $\delta_{17,1}$. After calculation we obtain the 2 irreducible semi-invariants:

$$simn_1 := [de_3, vi_5]^2, simn_2 := [si_8, des_7].$$

There exists only semi-invariants of orders 0 and 2 in the set $\overline{C}_{18}$. The nine irreducible invariants were found in [16] -

$$\begin{aligned} vis_1 &:= [de_4, de_3]^3, vis_2 := [de_4, de_{10}]^3, vis_3 := [de_5, de_6], vis_4 := [de_1, de_{10}]^3, vis_5 := [de_2, de_3]^3, \\ vis_6 &:= [de_2, de_{10}]^3, vis_7 := [de_3, de_{10}]^3, vis_8 := [de_6, de_7], vis_9 := [de_8, de_9]^5. \end{aligned}$$

For $\overline{C}_{18,2}$ we have $\dim C_{18,2} = 63$, $\sigma_{18,2} = 105$, $\dim S_{18,2} = 42$. Thus $\delta_{18,2} = 0$ i $\delta_{18} = 9$. The results of Sylvester's and Gall's for $\delta_{18,2}$ were 0 and 1 respectively.

There exists only semi-invariants of order 1 in $\overline{C}_{19}$. We have $\dim C_{19,1} = 46$, $\sigma_{19,1} = 57$, $\dim S_{19,1} = 12$. Thus $\delta_{19} = \delta_{19,1} = 1$. The results of Sylvester's and Gall's for $\delta_{19,1}$ were 0 and 2 respectively. After calculation we obtain the irreducible semi-invariant:

$$devn := [de_7, des_7].$$

There exists only one invariant in $\overline{C}_{20}$ which is calculated in [16] – $dvad_{20} := [des_7, des_7]^2$. The results of Sylvester's, Gall's and Diximier's for δ_{20} were 0, 2 and 1 respectively.

There exists only two invariants in $\overline{C}_{22}$ which are calculated in [16]:

$$dvdv_1 := [odn_6, odn_1]^3, dvdv_2 := [odn_8, odn_1]^3.$$

The results of Sylvester's, Gall's and Diximier's for δ_{22} were 1, 3 and 2 respectively.

There exists only semi-invariants of order 1 in $\overline{C}_{23}$. We have $\dim C_{23,1} = 85$, $\sigma_{23,1} = 142$, $\dim S_{23,1} = 58$. Thus $\delta_{23} = \delta_{23,1} = 1$. The results of Sylvester's and Gall's for δ_{23} were 0 and 1 respectively. After calculation we get the semi-invariant – $dvtr := [tryn_4, des_7]$.

In [3], [9], [16] proved that $\delta_{24} = 0$.

There exists only semi-invariants of order 1 in $\overline{C}_{25}$. We have $\dim C_{25,1} = 114$, $\sigma_{25,1} = 228$, $\dim S_{25,1} = 114$. $\delta_{25} = \delta_{25,1} = 0$. The results of Sylvester's and Gall's for δ_{25} were 0 and 1 respectively.

There exists only one covariant of degree 26 – $dvsh = [tryn_4, tryn_3]$, no covariants of degrees 27, 28, 29 and only one covariant of degree 30 – $trd := (h, \alpha)$ (in Gall's notations). By [3] it follows that $\delta_i = 0$ for $i > 30$.

Summarizing the above results we get

Theorem. *The system of the following 147 covariants*

$$\begin{aligned} & \kappa^{-1}(t) \\ & \kappa^{-1}(dv_i), i = 1, \dots, 3, \kappa^{-1}(tr_i), i = 1, \dots, 6, \kappa^{-1}(ch_i), i = 1, \dots, 8, \kappa^{-1}(pt_i), i = 1, \dots, 10 \\ & \kappa^{-1}(sh_i), i = 1, \dots, 10, \kappa^{-1}(si_i), i = 1, \dots, 12, \kappa^{-1}(vi_i), i = 1, \dots, 13, \kappa^{-1}(de_i), i = 1, \dots, 11 \\ & \kappa^{-1}(des_i), i = 1, \dots, 9, \kappa^{-1}(odn_i), i = 1, \dots, 8, \kappa^{-1}(dvn_i), i = 1, \dots, 13, \kappa^{-1}(tryn_i), i = 1, \dots, 9 \\ & \kappa^{-1}(chot_i), i = 1, \dots, 6, \kappa^{-1}(ptn_i), i = 1, \dots, 4, \kappa^{-1}(shis_i), i = 1, \dots, 5, \kappa^{-1}(simn_i), i = 1, \dots, 2 \\ & vis_i, i = 1, \dots, 9, \kappa^{-1}(devn), dvad, dvdv_1, dvdv_2, \kappa^{-1}(dvtr), dvsh, trd, \end{aligned}$$

is a complete system of the covariants for the binary form of degree 7.

REFERENCES

- [1] Dixmier, J., Quelques aspects de la théorie des invariants, *Gaz. Math., Soc. Math. Fr.*, 43, 39-64 (1990)
- [2] Gordan, P., Invariantentheorie, Teubner, Leipzig, reprinted by Chelsea Publ. Co., (1987).
- [3] von Gall, F., Das vollständige Formensystem der binären Form 7^{ter} Ordnung. *Math. Ann.* XXXI, 318-336, (1888).
- [4] Cayley, A. A Second Memoir upon Quantic. *Philosophical Transactions of the Royal Society of London*, Vol. 146, 101-126, (1856).
- [5] Olver, P., Classical invariant theory, Cambridge University Press, (1999).
- [6] Sylvester, J. J., Franklin, F. Tables of the generating functions and groundforms for the binary quantic of the first ten orders. *Am. J. II.* 223-251. (1879)
- [7] von Gall, F., Das vollständig Formensystem der binären Form achter Ordnung, *Math. Ann.*, 17, 31-52, 139-152, (1880).
- [8] Shioda, T. On the graded ring of invariants of binary octavics. *Am. J. Math.* 89, 1022-1046 (1967).
- [9] Dixmier, J., Lazard, D., Le nombre minimum d'invariants fondamentaux pour les formes binaires de degré 7. *Port. Math.*, 43,, 377–392, (1986). *J. Symb. Comput.*, 6, No.1, 113-115 (1988).
- [10] Roberts, M., The covariants of a binary quantic of the n-th degree, *Quarterly J. Math.* 4, 168–178, (1861).
- [11] van den Essen, A., Polynomial automorphisms and the Jacobian conjecture. *Progress in Mathematics* (Boston, Mass.). 190. Basel: Birkhauser, (2000).
- [12] Hilbert, D., Theory of algebraic invariants. Lectures. Cambridge University Press, (1993).
- [13] O. E. Glenn, reatise on theory of invariants, Boston, 1915.
- [14] Springer, T.A., Invariant theory, Lecture Notes in Mathematics, 585, Springer-Verlag, (1977).
- [15] Bedratyuk, L.P., On Differential equation of invariants of binary forms, math.AG/0602373.
- [16] Bedratyuk, L.P., On complete system of invariants for the binary form of degree 7, math.AG/0611122.
- [17] Bedratyuk, L.P., On complete system of covariants for the binary form of degree 8, math.AG/0612113.

5. APPENDIX

The degree-order distribution of $\overline{C}_8$.

		order															
		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
degree	1							1									
	2			1				1				1					
	3				1		1		1		1		1				1
	4	1				2		1		2		1				1	
	5		1		2		2		2		2				1		
	6			3		2		2		2				1			
	7		3		2		4		2				1				
	8	3		3		3		3				1					
	9		3		5		2				1						
	10			4		4				1							
	11		5		3				1								
	12	6		6				1									
	13		7		1		1										
	14	4				2											
	15		3		1												
	16	2		3													
	17		2														
	18	9															
	19		1														
	20	1															
	21																
	22	2															
	23		1														
	24																
	25		1														
	26	1															
	30	1															

KHMELNYTSKY NATIONAL UNIVERSITY, INSTYTUTS'KA ST. 11, KHMELNYTSKY , 29016, UKRAINE
E-mail address: bedratyuk@ief.tup.km.ua