Generating spaces of modular forms with η -quotients

L. J. P. Kilford

November 26, 2024

Abstract

In this note we consider a question of Ono, concerning which spaces of classical modular forms can be generated by sums of η -quotients. We give some new examples of spaces of modular forms which can be generated as sums of η -quotients, and show that we can write all modular forms of level $\Gamma_0(N)$ as rational functions of η -products.

1 Introduction

Let z be an element of the Poincaré upper half plane \mathcal{H} , and let $q := \exp(2\pi i z)$. We recall the definition of the Dedekind η -function:

$$\eta(q) := q^{1/24} \prod_{n=1}^{\infty} (1 - q^n).$$

It is well-known that this function plays an important role in the theory of modular forms; for instance, the unique normalised cusp form of level 1 and weight 12, the Δ function, can be written as $\Delta(q) = \eta(q)^{24}$.

Another classical application of the η -function is to the theory of partitions; a partition of n is a way of writing n as a sum of positive integers; for instance, 4 = 2 + 2 is a partition of 4. The number of partitions of n, p(n), grows quickly with n. For instance, in the early years of the $20^{\rm th}$ century, MacMahon computed that p(200) = 397299029388. The reciprocal of the η -function (with the $q^{1/24}$ removed) gives the following generating function for the p(n):

$$\sum_{n=0}^{\infty} p(n)q^n = \prod_{m=1}^{\infty} \left(\frac{1}{1-q^m}\right).$$

It can be shown that $\eta(q)^{24}$ is a modular form for $\mathrm{SL}_2(\mathbf{Z})$ In [1], an explicit transformation formula for $\eta(q)$ under the action of $\mathrm{SL}_2(\mathbf{Z})$ is given; it is of the form

$$\eta\left(\frac{az+b}{cz+d}\right) = \varepsilon \cdot (cz+d)^{1/2}\eta(z) \text{ for } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbf{Z}),$$
(1)

where $\varepsilon^{24} = 1$. (The explicit definition of ε , which depends on a, b, c, d, is complicated, but is given in Theorem 3.4 of [1]).

We see from (1) that $\eta(q)$ satisfies a "weight 1/2" transformation formula, which is consistent with the statement that Δ is a weight 12 modular form.

Definition 1. Let N be a positive integer, let $\{r_{\delta}\}$ be a set of integers, and let f be a meromorphic function from the Poincarè upper half plane to \mathbf{C} of the form

$$f(z) = \prod_{0 < \delta \mid N} \eta \left(q^{\delta} \right)^{r_{\delta}}$$
, where $q := \exp(2\pi i z)$.

We call f an η -quotient, and if all of the r_{δ} are non-negative we say that f is an η -product.

We see that Δ is an η -product; we will see some examples of η -quotients which are modular forms below.

There are some general theorems which tell us when an η -quotient is a classical modular form, by telling us how it transforms under elements of certain congruence subgroups.

Theorem 2 (Ligozat [11], quoted in [13]). Suppose that f(z) is an η -quotient which satisfies the following properties:

$$\sum_{0 < \delta | N} \delta \cdot r_{\delta} \equiv 0 \mod 24 \tag{2}$$

and

$$\sum_{0 < \delta | N} \frac{N}{\delta} \cdot r_{\delta} \equiv 0 \mod 24. \tag{3}$$

Then f(z) satisfies

$$f\left(\frac{az+b}{cz+d}\right) = \chi(d)(cz+d)^k f(z) \tag{4}$$

for every $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$, where $k := \frac{1}{2} \sum_{0 < \delta | N} r_{\delta}$, and

$$\chi(d) := \left(\frac{(-1)^k \cdot s}{d}\right), \text{ where } s := \prod_{0 < \delta \mid N} \delta^{r_\delta}.$$
(5)

In [15], the following question is posed:

Question 3 (Ono [15], Problem 1.68). Which spaces of classical modular forms can be generated by sums of η -quotients?

It is well-known (see for instance [16], Corollary 2 to Theorem VII.4) that every modular form of level 1 is a polynomial in the Eisenstein series E_4 and E_6 ,

so if we can write E_4 and E_6 in terms of η -quotients then we can write every modular form of level 1 in terms of η -quotients. In [15], we see that

$$E_4(z) = \frac{\eta(q)^{16}}{\eta(q^2)^8} + 2^8 \cdot \frac{\eta(q^2)^{16}}{\eta(q)^8}$$

and

$$E_6(z) = \frac{\eta(q)^{24}}{\eta(q^2)^{12}} - 2^5 \cdot 3 \cdot 5 \cdot \eta(q^2)^{12} - 2^9 \cdot 3 \cdot 11 \cdot \frac{\eta(q^2)^{12} \eta(q^4)^8}{\eta(q)^8} + 2^{13} \cdot \frac{\eta(q^4)^{24}}{\eta(q^2)^{12}}.$$

One can check that these are identities of modular forms by verifying that each of these η -quotients is a modular form on $\Gamma_0(4)$ and that the q-expansions of both sides agree up to a suitable bound, such as the Sturm bound [10] (also known as the Hecke bound); if a modular form has a zero at the cusp ∞ of large enough degree, then it must be the zero modular form. In this case, we find that the dimension of the space $M_4(\Gamma_0(4))$ is 3, and the Sturm bound here is 2. We can use a computer algebra package to check that these really are equalities.

In this paper we will give further examples of this form, and prove a theorem which says that we can write *every* modular form for a congruence subgroup $\Gamma_0(N)$ in terms of η -quotients of level at most $\Gamma_0(4N)$.

2 Writing eigenforms as η -products or η -quotients

We will briefly consider η -products, and show that there are only finitely many spaces of modular forms which can be generated by η -products. These do exist: for instance, $S_2(\Gamma_0(11)) = \mathbf{C} \cdot \eta(q)^2 \eta(q^{11})^2$, and $S_1(\Gamma_1(23)) = \mathbf{C} \cdot \eta(q)\eta(q^{23})$. However, it is clear that only finitely many spaces of modular forms can be completely generated by η -products, as there are only finitely many η -products with q-expansion of the form $q^a + O(q^{a+1})$ for any a; this follows from equation (2), because as the level increases the sum increases also.

Questions of this sort have been asked before; in [2], the question of when an η product is also a simultaneous eigenform for the Hecke operators T_p is addressed. In [5], a complete list of the 30 different η -products which are classical modular cuspidal eigenforms of integer weight with no zeroes outside the cusps is given (two of these have half-integral weight). Another proof of this classification was given by [9].

Similarly, in [12], a complete list of all η -quotients which are holomorphic new eigenforms of integral weight for the congruence subgroups $\Gamma_1(N)$ is derived. There are 74 of these, with levels ranging from $\mathrm{SL}_2(\mathbf{Z})$ (the Δ -function) up to $\Gamma_1(576)$ (the modular form $\eta(q^{12})^{-2} \cdot \eta(q^{24})^6 \cdot \eta(q^{48})^{-2} \in S_1(\Gamma_1(576))$). There have been other investigations; in [8], a list of the 65 η -products which have weight 1 and level $\Gamma^*(36)$ is given; the congruence subgroup $\Gamma^*(N)$ is generated by $\Gamma_0(36)$ and the Fricke involution $\begin{pmatrix} 0 & 1/\sqrt{N} \\ -1/\sqrt{N} & 0 \end{pmatrix}$. Another angle of inquiry about η -products is given in [6], which considers η -products which are lacunary modular forms; in other words, they have many zero Fourier coefficients.

3 Sums of η -products and η -quotients

We see that, from the results we have mentioned in the previous section, if we are to generate all spaces of modular forms for a given congruence subgroup Γ we will need to expand from η -products and η -quotients to consider more general objects. We will consider sums of η -products and η -quotients in this section.

From equation (4), we see that the only characters we can have are the Legendre symbols, so if we are considering congruence subgroups containing $\Gamma_1(N)$ we are restricted to the cases of $\Gamma_0(N)$ and $\Gamma_0(N)$ plus a character of order 2.

We will be using the standard formulae for the dimension of spaces of modular forms in this section; these can be found in many books on modular forms; one reference is Theorem 3.5.1 and Figure 3.3 of [4]. Let $g = \lfloor (p+1)/12 \rfloor$ if $p \not\equiv 1 \mod 12$ and $\lfloor (p+1)/12 \rfloor - 1$ otherwise (this is the genus of $X_0(p)$). The formulae (if N = p is prime) are (for $k \ge 4$ and $N \ge 5$):

$$\dim S_k(\Gamma_0(p)) = (k-1) \cdot (g-1) + \left\lfloor \frac{k}{4} \right\rfloor \cdot \left(1 + \left(\frac{-1}{p}\right)\right) + \left\lfloor \frac{k}{3} \right\rfloor \cdot \left(1 + \left(\frac{-3}{p}\right)\right) + k - 2.$$

The dimension of $S_2(\Gamma_0(p))$ is g, and the dimension of the space of Eisenstein series is 1 if k=2 and 2 if $k\geq 4$ (the Eisenstein series of weight $k\geq 4$ are the images of E_k under the two degeneracy maps from level 1 to level p, and the Eisenstein series of weight 2 is defined in terms of the Eisenstein series of weight 2 which is not a modular form). There are similar formulae for N composite and for other congruence subgroups.

We will first consider some more examples of spaces of modular forms which can be generated by η -quotients: for example, we can write every modular form of level $\Gamma_0(2)$ as a sum of η -quotients. It will suffice to show that $M_2(\Gamma_0(2))$, $M_4(\Gamma_0(2))$, $M_6(\Gamma_0(2))$ and $M_8(\Gamma_0(2))$ can be generated by sums of η -quotients, because the following relation holds:

$$S_{k+8}(\Gamma_0(2)) = f \cdot M_k(\Gamma_0(2)), \text{ for } k \in \mathbb{N}.$$

where $f = \lambda \cdot (\eta(z) \cdot \eta(2z))^8 \in S_8(\Gamma_0(2))$. This can be derived from the dimension formulae that we have given above.

Because we can write level 1 modular forms as sums of η -quotients, and we know (for instance, from Proposition III.3.19 of [7]) that the unique normalised cusp-form f of weight 8 and level $\Gamma_0(2)$ is an η -quotient, we need only to consider $M_2(\Gamma_0(2))$, which is 1-dimensional. Using MAGMA [3], and the fact that the Hecke bound for the space $M_2(\Gamma_0(8))$ is 3, we see that

$$\frac{\eta(q^2)^{20}}{\eta(q)^8 \cdot \eta(q^4)^8} + 16 \cdot \frac{\eta(q^8)}{\eta(q^2)^4} = E_{2,2} \in M_2(\Gamma_0(2)),$$

so every modular form of level $\Gamma_0(2)$ can be written as a sum of η -quotients. We need to consider the space of level $\Gamma_0(8)$ because the η -quotients have level $\Gamma_0(8)$.

Similarly, to show that every modular form for $\Gamma_0(3)$ is a sum of η -quotients, we need to check that $E_{3,2} \in M_2(\Gamma_0(3))$ is a sum of η -quotients. After some calculation, we find (using the fact that the Hecke bound for $M_2(\Gamma_0(27))$ is 6) that

$$36\frac{\eta(q^9)^6}{\eta(q^3)^2} + 12\frac{\eta(q^3)^2}{\eta(q^9)^2} + \frac{\eta(q^9)^{10}}{\eta(q^3)^3\eta(q^{27})^3} + 108\frac{\eta(q^9)^2\eta(q^{27})^3}{\eta(q^3)} + 9\frac{\eta(q^3)^3\eta(q^{27})^3}{\eta(q^9)^2} + 9\frac{\eta(q^{27})^6}{\eta(q^9)^2}$$

is equal to $E_{3,2}$; this, together with the knowledge that $S_6(\Gamma_0(3)) = \mathbf{C} \cdot \eta(q)^6 \eta(q^3)^6$ and that the Eisenstein series of weights 4 and 6 and level $\Gamma_0(3)$ are oldforms and therefore the image of forms of level 1 which can be written as η quotients, shows us that we can write modular forms of level $\Gamma_0(3)$ as sums of η quotients of level at most $4 \cdot 27$.

In fact, we can write E_4 and E_6 as sums of eta-quotients of level 27, so we can actually write every modular form for $\Gamma_0(3)$ in terms of η quotients of level 27.

In a similar fashion, we can write the elements of $M_3(\Gamma_1(3))$ as a linear combination of $\eta(q^3)^9/\eta(q)^3$ and $\eta(q)^9/\eta(q^3)^3$; these, together with the Eisenstein series of level 3 and weights 2,4 and 6 and the cusp form of level 3 and weight 6 generate all of the modular forms of level 3 and odd weight, so we can write every modular form of level $\Gamma_1(3)$ as a sum of η -quotients.

We can prove similar results for level $\Gamma_1(4)$ as well. Using Exercise III.3.17 of Koblitz's book on modular forms [7], we see that every $f \in M_k(\Gamma_1(4))$ can be written as a polynomial in the modular forms $F := \eta(q^4)^8/\eta(q^2)^4$, which has weight 2, and $\theta^2 := \eta(q^2)^{10}/(\eta(q)^4\eta(q^4)^4)$, which has weight 1. As both of these are η -quotients, we are done; every modular form of level $\Gamma_1(4)$ can be written as a sum of η -quotients of level at most 4.

A rather tedious MAGMA calculation shows us that the unique Eisenstein series $E_{5,2}$ of level $\Gamma_0(5)$ and weight 2 can be written as the following sum of η -quotients of level $\Gamma_0(20)$:

$$E_{5,2}(q) = 5 \cdot \frac{\eta(q^2)^3 \eta(q^4)^4 \eta(q^5)^3 \eta(q^{10})^5}{\eta(q)^7 \eta(q^{20})^4} - 2 \cdot \frac{\eta(q^2)^7 \eta(q^4)^2 \eta(q^5)^3 \eta(q^{10})}{\eta(q)^7 \eta(q^{20})^2}$$

$$- 4 \cdot \frac{\eta(q^2)^7 \eta(q^4)^7 \eta(q^{10})^5}{\eta(q)^7 \eta(q^5)^5 \eta(q^{20})^3} + \frac{\eta(q^4)^4 \eta(q^5)^6 \eta(q^{10})^4}{\eta(q)^6 \eta(q^{20})^4}$$

$$+ 2 \cdot \frac{\eta(q^2)^4 \eta(q^4)^2 \eta(q^5)^6}{\eta(q)^6 \eta(q^{20})^2} - \frac{\eta(q^2)^4 \eta(q^4)^7 \eta(q^{10})^4}{\eta(q)^6 \eta(q^5)^2 \eta(q^{20})^3}.$$

This follows because these six η -quotients generate $M_2(\Gamma_0(20))$, where $E_{5,2}$ appears as an oldform. We can find the coefficients of these using linear algebra.

In a very similar way, we can write both elements of $M_2(\Gamma_0(5), (\frac{\cdot}{5}))$ as η -quotients by considering them as oldforms of level 20 and finding a basis of η -quotients for this space of level 20 forms. This suffices to show that all modular forms with character $(\frac{\cdot}{5})$ can be written as sums of η -quotients.

At this point, we notice that if $p \ge 5$ is a prime then we cannot write every modular form for $\Gamma_0(p)$ as a sum of η -quotients of level p^r . We use the dimension

Group	Level of η -products needed
$\mathrm{SL}_2(\mathbf{Z})$	4 or 27
$\Gamma_1(2)$	8
$\Gamma_1(3)$	27
$\Gamma_1(4)$	4
$\Gamma_0(5)$	20
$\Gamma_0(7)$	28

Table 1: Levels of η -quotients needed to generate spaces of modular forms

formulae for $M_2(\Gamma_0(p))$ to show that there is at least one Eisenstein series of weight 2 with nonzero constant term in its Fourier expansion which we need to write as a sum of η quotients, each of which has the form $\eta(q)^a \eta(q^p)^b \cdots \eta(q^{p^r})^z$. Therefore, we need to solve the following two simultaneous equations:

$$a + p \cdot b + p^2 \cdot c + \dots + p^r \cdot z = 0 \tag{6}$$

$$a+b+c+\dots+z=4; (7)$$

putting these together, we find that

$$(p-1) \cdot b + (p^2 - 1) \cdot c + \dots + (p^r - 1) \cdot z = -4.$$
 (8)

We now reduce (8) modulo p-1; the left hand side vanishes, and the only way that the right hand side can vanish modulo p-1 is if $p \in \{2,3,5\}$.

Now we consider the case when p=5. We use the formula for the character of the η -quotient given in (5) to work on (8); after dividing both sides by 4, we see that the right hand side is -1, whereas the left hand side has only even entries (because the requirement that the character be trivial forces the sum $b+d+f+\cdots$ to be even). This is a contradiction.

We can tackle $\Gamma_0(7)$ in a similar fashion; in this case, we have to check that we can generate $M_2(\Gamma_0(7))$ and $M_4(\Gamma_0(7))$ by η -quotients, and again we prove this by finding a basis of η -quotients for $M_2(\Gamma_0(28))$ and $M_4(\Gamma_0(28))$ and then considering the level 7 spaces as oldforms in these spaces. Here we find that

$$E_{7,2} = 16 \cdot \frac{\eta(q^2)^6 \eta(q^4)^6 \eta(q^{14})^2}{\eta(q)^8 \eta(q^{28})^2} - 32 \cdot \frac{\eta(q^2)^4 \eta(q^4)^3 \eta(q^7)^6}{\eta(q)^6 \eta(q^{14})^2 \eta(q^{28})}$$

$$+ \frac{49}{2} \cdot \frac{\eta(q^4) \eta(q^7)^8 \eta(q^{14})^2}{\eta(q)^4 \eta(q^{28})^3} - 10 \cdot \frac{\eta(q^2)^4 \eta(q^4)^6 \eta(q^{14})^4}{\eta(q)^4 \eta(q^7)^4 \eta(q^{28})^2}$$

$$- \frac{16}{7} \cdot \frac{\eta(q^2)^5 \eta(q^7)^5}{\eta(q)^3 \eta(q^{14})} + 6 \cdot \frac{\eta(q^2)^7 \eta(q^4)^2 \eta(q^{28})^2}{\eta(q) \eta(q^7) \eta(q^{14})^5}$$

$$- \frac{17}{14} \cdot \frac{\eta(q) \eta(q^2)^3 \eta(q^7)^5 \eta(q^{14})^5}{\eta(q^4)^7 \eta(q^{28})^3};$$

the equations for the three cuspforms in $S_4(\Gamma_0(7))$ are similar.

4 Generating spaces of modular forms with rational functions of η -quotients

We now consider a larger set of functions than in the previous section; we will now consider rational functions of η -quotients and η -products. We will now prove the following theorem on the generation of spaces of modular forms by rational functions of η -quotients (and in fact by rational functions of η -products):

Theorem 4. Let N be a positive integer, let k be a positive even integer, and let $f \in M_k(\Gamma_0(N))$ be a modular form. Then we can write f as a rational function of η -quotients.

Proof. We first note that the classical j-invariant is an η -quotient, because we can write it as

$$j := \frac{E_4^3}{\Delta},$$

where E_4 is an η -quotient and $\Delta = \eta(q)^{24}$. This means that the modular function $j(q^N)$ is also an η -quotient, for any $N \in \mathbb{N}$.

We now quote a theorem about the field of modular functions of weight 0 for the congruence subgroup $\Gamma_0(N)$; this is proved in the online notes of Milne [14] on modular forms and modular functions:

Theorem 5 (Milne [14], Theorem 6.1). The field $C(X_0(N))$ of modular functions for $\Gamma_0(N)$ is generated over C by j(q) and $j(q^N)$. In particular, every modular function for $\Gamma_0(N)$ is a rational function of j(q) and $j(q^N)$.

By standard properties of modular forms, the quotient f/g of two modular forms of weight k for a congruence subgroup Γ is a modular function of weight 0 for that congruence subgroup, with possible poles at the zeroes of g.

We therefore take $f \in M_k(\Gamma_0(N))$ and consider the modular function

$$g := \frac{f \cdot E_4^a \cdot E_6^b}{\Delta^c} \in \mathcal{M}_0(\Gamma_0(N)), \tag{9}$$

where we choose $a \in \{0, 1, 2\}$, $b \in \{0, 1\}$ and $k \in \mathbb{N}$ such that k+4a+6b-12c=0. This has weight 0 by definition, so by Theorem 5 we can write it as a rational function of j(q) and $j(q^N)$, which are both rational functions of η -products. From before, we recall that E_4 , E_6 and Δ are all rational functions of η -products, so we can write f as a rational function of η -products, which is what we set out to do.

It is interesting to note that we only needed to use η -products of level 4N here (the N coming from the level of our form f, and the 4 coming from the η -quotient representations of E_4 and E_6). This corresponds to the figures in Table 1, where we needed to raise the level by a factor of at least 4, corresponding to the level 4 η -quotients that we used for E_4 and E_6 .

We see also that we have to have either the square of a prime or two distinct primes dividing the level, because we always need to write E_4 and E_6 in terms of η -products, and we need to find at least two different solutions to the set of simultaneous equations in the powers of the η^i (one for E_4 and one for E_6). This means that we need to have at least 3 i, so in particular we cannot write E_4 and E_6 just in terms of η -quotients of level p for any prime p.

If we are willing to raise the level then we can use $g := \eta(q)^2 \eta(q^{11})^2 \in S_2(\Gamma_0(11))$ as a multiplier in (9), so the only thing stopping our rational function being a polynomial is the numerator of the modular function of weight 0. However, we cannot assume that g only has a zero at ∞ , because while the modular form $\Delta(q) \in S_{12}(\operatorname{SL}_2(\mathbf{Z}))$ only has a simple zero at ∞ and no zeroes elsewhere, the modular form $\Delta(q) \in S_{12}(\Gamma_0(N))$ (viewed as an oldform) has many zeroes; it has one at each cusp of $\Gamma_0(N)$.

5 Acknowledgements

The author would like to thank Paul-Olivier Dehaye, Ken McMurdy and Alan Lauder for helpful conversations. The computations for this article were carried out on William Stein's computer MECCAH and on the Mathematics Institute in Oxford's computers; the author is grateful for this computer time.

References

- [1] Tom M. Apostol. Modular functions and Dirichlet series in number theory. Springer-Verlag, New York, 1976. Graduate Texts in Mathematics, No. 41.
- [2] Anthony J. F. Biagioli. Eta-products which are simultaneous eigenforms of Hecke operators. *Glasgow Math. J.*, 35(3):307–323, 1993.
- [3] W. Bosma, J. Cannon, and C. Playoust. The Magma algebra system I: The user language. *J. Symb. Comp.*, 24(3-4):235-265, 1997. See http://magma.maths.usyd.edu.au.
- [4] Fred Diamond and Jerry Shurman. A first course in modular forms, volume 228 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2005.
- [5] D. Dummit, H. Kisilevsky, and J. McKay. Multiplicative products of η-functions. In Finite groups—coming of age (Montreal, Que., 1982), volume 45 of Contemp. Math., pages 89–98. Amer. Math. Soc., Providence, RI, 1985.
- [6] Basil Gordon and Sinai Robins. Lacunarity of Dedekind η -products. Glasgow Math. J., 37(1):1–14, 1995.
- [7] N. Koblitz. Introduction to Elliptic Curves and Modular Forms. Springer, New York, 1993.

- [8] Günter Köhler. Eta products of weight 1 and level 36. *Arch. Math. (Basel)*, 76(3):202–214, 2001.
- [9] Masao Koike. On McKay's conjecture. Nagoya Math. J., 95:85–89, 1984.
- [10] Joan-C. Lario and René Schoof. Some computations with Hecke rings and deformation rings. *Experiment. Math.*, 11(2):303–311, 2002. With an appendix by Amod Agashe and William Stein.
- [11] Gérard Ligozat. Courbes modulaires de genre 1. Société Mathématique de France, Paris, 1975. Bull. Soc. Math. France, Mém. 43, Supplément au Bull. Soc. Math. France Tome 103, no. 3.
- [12] Yves Martin. Multiplicative η -quotients. Trans. Amer. Math. Soc., 348(12):4825-4856, 1996.
- [13] Ken McMurdy. A Splitting Criterion for Galois Representations Associated to Exceptional Modular Forms. PhD thesis, U.C. Berkeley, 2001. Available at http://www.rose-hulman.edu/~mcmurdy/research/.
- [14] J. Milne. Modular functions and modular forms. Available at http://www.jmilne.org/math/CourseNotes/math678.html, 1997.
- [15] Ken Ono. The web of modularity: arithmetic of the coefficients of modular forms and q-series, volume 102 of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2004.
- [16] J.-P. Serre. A course in arithmetic. Springer-Verlag, New York, 1973. Translated from the French, Graduate Texts in Mathematics, No. 7.