# The upper bound for the index of nilpotency for a matrix commuting with a given nilpotent matrix

#### Polona Oblak

Institute of M athematics, Physics and Mechanics, Department of Mathematics, Jadranska 19, SI-1000 Ljubljana, Slovenia

polona oblak@fm funi-ljsi

A bstract. We study the set P ( $N_B$ ) of all possible Jordan canonical form s of nilpotent matrices commuting with a given nilpotent matrix B. We describe P ( $N_B$ ) in the special case when B has only one Jordan block. In the general case, we not the maximal possible index of nilpotency in the set of all nilpotent matrices commuting with a given nilpotent matrix. We consider several examples.

M ath. Subj. Class (2000): 15A 04, 15A 21, 15A 27
K ey words: com muting nilpotent matrices, Jordan canonical form.

### 1 Introduction

We consider the following problem: What are possible sizes of Jordan blocks for a pair of commuting nilpotent matrices? Or equivalently, for which pairs of nilpotent orbits of matrices (under similarity) there exists a pair of matrices, one from each orbit, that commute. The answer to the question could be considered as a generalization of Gerstenhaber (Hesselink theorem on the partial order of nilpotent orbits [4].

The structure of the varieties of commuting pairs of matrices and of commuting pairs of nilpotent matrices is not yet well understood. It was proved by Motzkin and Taussky [12] (see also Guralnick [6]), that the variety of pairs of commuting matrices was irreducible. It was Guralnick [6] who showed that this is no longer the case for the variety of triples of commuting matrices (see also Guralnick and Sethuram an [7], Holbrook and Omladic [10], Omladic [13], Han [9]). Recently, it was proved that the variety of commuting pairs of nilpotent matrices was irreducible (Baranovsky [1], Basili [2]). Our motivation to study the problem is to contribute to better understanding of the structure of this variety and

which might also help in understanding the (ir) reducibility of the variety of triples of commuting matrices.

We are also motivated by the problems posed by Binding and Kosir [11] in the multiparam eter spectral theory and Gustafson [8] in the module theory over commutative rings. In both problems, certain pairs of commuting matrices appear. The matrices from the multiparam eter spectral theory generate an algebra that is a complete intersection, and the matrices from the theory of modules are both functions of another matrix.

Here we initiate the study of the problem. First we list all possible Jordan forms for nilpotent matrices commuting with a single Jordan block. In the Section 3 we recall the correspondence between directed graphs and generic matrices (also known as Gansner{ Saks Theorem), which is the main tool to prove our main result, i.e., to compute the maximal index of nilpotency of a nilpotent matrix commuting with a given nilpotent matrix with Jordan canonical form. In the last section we discuss some further examples.

## 2 The one Jordan block case and consequences

Let us denote by N = N (n; F) the variety of all n n nilpotent m atrices over a com m utative eld F and write  $N_2 = f(A; B) 2 N$  N; AB = BAg.

We follow the notations used in Basili [2] and write N<sub>B</sub> = fA 2 N; AB = BAg for someB 2 N. Suppose  $_1$   $_2$  :::  $_t > 0$  are the orders of Jordan blocks in the Jordan canonical form for B. We call the partition  $_=$  (  $_1$ ;  $_2$ ;:::;  $_t$ ) the shape of the matrix B and denote it by sh(B). We also write sh(B) = ( $_1$ ;  $_1$ ;  $_2$ ;  $_2$ ;  $_2$ ;  $_3$ ;  $_4$ ;  $_$ 

Let P (n) denote the set of all partitions of n 2 N and for a subset S N (n; F) write

$$P(S) = f(2P(n)); = sh(A) \text{ for som } eA 2 Sg:$$

D enote also

$$P(N_2) = f(sh(A); sh(B)); (A;B) 2 N_2g P(n) P(n):$$

Note that (\_; ) 2 P (N  $_2$ ) if and only if ( ;\_) 2 P (N  $_2$ ), i.e. P (N  $_2$ ) is sym m etric.

It is easy to see that for each t=1;2;:::;n there exists a uniquely de ned partition r(n;t):=(1;2;:::;t) 2 P (n), such that 1 there exists a uniquely defined partition r(n;t):=(1;2;:::;n) 2 P (n), such that 1 the same that r(n;t):=(1;2;:::;n) 3. It can be verified that r(n;t):=(n;t) 4. It can be verified that r(n;t):=(n;t) 5. It can be verified that r(n;t):=(n;t) 6. It can be verified that r(n;t):=(n;t) 8. It can be verified that

Take a matrix B 2 N (n;F) with dim kerB = 1 (i.e. sh(B) = (n)). Then it is well known that any matrix commuting with B is a polynomial in B. By computing the lengths of the Jordan chains of B, we observe that  $sh(B^k) = r(n;k)$  for k = 1;2;:::;n. Thus we have the following.

Proposition 1. For a matrix B with dim kerB = 1 it follows that P ( $N_B$ ) = R (n).

For a sequence  $(a_1; a_2; \dots; a_k)$ ,  $a_i \ge N$ , we write ord  $(a_1; a_2; \dots; a_k) = (a_{(1)}; a_{(2)}; \dots; a_{(k)})$ , where  $a_{(1)}$  a  $a_{(2)}$   $\dots$  a  $a_{(k)}$  and is a permutation of f1;2;...; kg. For  $a_{(1)} = (a_{(1)}; a_{(2)}; \dots; a_{(k)})$  we write

R (\_) = ford(r(
$$_1;s_1$$
);r( $_2;s_2$ );:::;r( $_t;s_t$ ));  $s_i = 1;2;:::;ig:$ 

So, for example, (3;3;2) 2 R ((5;3)), but  $(4;3;1) \ge R$  ((5;3)).

Proposition 2. For all 2P(n) it follows that f(:); 2R()g  $P(N_2)$ .

Proof. Take an arbitrary B 2 N with  $sh(B) = _ = (_1;_2;:::;_t)$  and pick \_ 2 R (\_). We want to nd a matrix A 2 N  $_B$  such that  $sh(A) = _ .$ 

By de nition, \_ is of the form \_ = ord(r( $_1;k_1$ );r( $_2;k_2$ );:::;r( $_t;k_t$ )). Let J  $_i$  be a  $_i$  \_ Jordan block corresponding to eigenvalue 0. Since sh( $_i$ ) = r( $_i;k_i$ ), it follows that the matrix \_ 3

$$A = \begin{cases} 2 & J_{1}^{k_{1}} & 0 & 0 & \dots & 0 \\ 6 & 0 & J_{2}^{k_{2}} & 0 & \dots & 0 \\ 7 & \vdots & \ddots & \vdots & 0 \\ 0 & 0 & 0 & \dots & J_{t}^{k_{t}} \end{cases}$$

has sh(A) =\_ and clearly  $A 2 N_B$ .

For a matrix A denote by (A) its index of nilpotency, i.e. (A) = minfi;  $A^i = 0$ g. For S N (n;F) write (S) = maxfi; i = (A) for some A 2 Sg. The following C orollaries can be easily obtained from P ropositions 1 and 2.

Corollary 3. If 2P(n), then 2R(n) if and only if  $((n); 2P(N_2)$ .

Corollary 4. For B 2 N it follows that sh(B) 2 R (n) if and only if  $(N_B) = n$ .

Exam ple 5. Note that Proposition 2 gives us a rather large subset of partitions in N<sub>B</sub>. Take sh(B) =  $(4;3;2^2;1)$ . First we directly observe that  $(4;3;2^2;1)$ ,  $(4;3;2;1^3)$ ,  $(4;3;1^5)$ ,  $(4;2^3;1^2)$ ,  $(4;2^2;1^4)$ ,  $(4;2;1^6)$ ,  $(4;1^8)$ ,  $(3;2^4;1)$ ,  $(3;2^3;1^3)$ ,  $(3;2^2;1^5)$ ,  $(3;2;1^7)$ ,  $(2^5;1^2)$ ,  $(2^4;1^4)$ ,  $(2^3;1^6)$ ,  $(2^2;1^8)$ ,  $(2;1^{10})$ ,  $(1^{12})$  are all in P(N<sub>B</sub>).

Next, we see that  $(4;3;2^2;1)$  is included in R (\_) if \_ is (7;5), (7;4;1), (7;3;2),  $(7;2^2;1)$ , (5;4;3), (5;4;2;1),  $(4;3^2;2)$  or  $(4;3;2^2;1)$ . Thus, by sym m etry, also all these partitions are in P (N  $_B$  ).

Corollary 3 shows that P (N  $_{\rm B}$ )  $\stackrel{\leftarrow}{\bullet}$  P (12). The natural question is whether the list above is the entire P (N  $_{\rm B}$ ). It can be veri ed with M athem atica that the answer is negative and that there are m ore partitions in P (N  $_{\rm B}$ ). However, it would take a lot of time to compute P (N  $_{\rm B}$ ) without computer. From results in Section 4 it follows that there exists a partition (  $_{1}$ ;  $_{2}$ ;:::;  $_{\rm k}$ ) 2 P (N  $_{\rm B}$ ) such that  $_{1}$  = 9.

# 3 A review of the Gansner-Saks theory

From now on we assume that F = C.

Let us recall the connection between generic matrices and directed graphs. (See also Brualdi, Ryser [3], Gansner [5], Saks [14].) We apply it to compute  $(N_B)$  in the next section.

A digraph is a directed graph (i.e. a graph each of whose edges are directed). A path of length k in a digraph is a sequence of vertices  $a_1; a_2; \ldots; a_k$  such that  $(a_i; a_{i+1})$  is an edge in for  $i=1;2;\ldots;k$  1. We do allow a path to consist of a single vertex. For a path P we denote by P jits length. We call  $a_1$  (resp.  $a_k$ ) the initial (resp. nal) vertex of P. A path  $a_1; \ldots; a_k$  with  $a_1 = a_k$  is called a cycle. A digraph without cycles is called an acyclic digraph.

Let be a nite acyclic digraph on n vertices, with the vertices labeled from 1 to n. A k{path in is a subset of the vertices that can be partitioned into k or fewer disjoint paths. Let  $\hat{d}_k = \hat{d}_k$ () be the largest cardinality of a k{path in and de ne  $\hat{d}_0 = 0$  and k = k() =  $\hat{d}_k$   $\hat{d}_k$  1. Since  $\hat{d}_k$   $\hat{d}_{k+1}$ , all the k0s are nonnegative, so we have the in nite sequence = () = (1; 2;:::) of nonnegative integers.

A matrix M 2 N (n;C) is called generic if all its nonzero entries are independent transcendentals over rational numbers Q. We say that matrices M =  $[m_{ij}]$  and N =  $[n_{ij}]$  of the same size have the same pattern if  $m_{ij} = 0$  ()  $n_{ij} = 0$  for i;j = 1;2;:::;n.

For a nilpotent matrix A and a generic M of the same pattern it is clear that (A) (M).

Given a nite acyclic digraph with n vertices, we assign to it a generic n n matrix  $M = [m_{ij}]$  such that  $m_{ij} = 0$  if (i;j) is not an edge in and such that the rest of the entries of M are nonzero complex numbers which are independent transcendentals. Since is acyclic, M is nilpotent. Conversely, given a nilpotent generic matrix, it corresponds, reversing the above assignment, to an acyclic digraph on n vertices.

The following theorem was independently proved by Gansner [5] and Saks [14].

Theorem 6. Let M be a generic nilpotent matrix. Then

$$(M_M) = \operatorname{sh}(M)$$
:

Corollary 7. The length of the longest path in a digraph is equal to (M).

# 4 M axim alindex of nilpotency in $N_B$

From now on, let B 2 N (n;C) be a xed matrix and write sh(B) =  $_{1}$  = (  $_{1}$ ;  $_{2}$ ;:::;  $_{t}$ ) = (m  $_{1}^{r_{1}}$ ; m  $_{2}^{r_{2}}$ ; :::; m  $_{1}^{r_{1}}$ ). By convention we have m  $_{1}$  > m  $_{2}$  > :::> m  $_{1}$  >  $_{0}$ .

For a pair of matrices (A;B) 2 N<sub>2</sub> denote sh (A;B) = (sh (A);sh (B)). Then there exists P 2 GL<sub>n</sub> (C) such that B = PJ<sub>B</sub> P  $^{-1}$ , where J<sub>B</sub> denotes the Jordan canonical form of matrix B. Thus sh (PAP  $^{-1}$ ;J<sub>B</sub>) = (sh (A);sh (B)) and (PAP  $^{-1}$ ;J<sub>B</sub>) 2 N<sub>2</sub>. Therefore we can assume that B is already in its upper triangular Jordan canonical form.

Write  $A = [A_{ij}]$  where  $A_{ij} 2$  M  $_{i}$   $_{j}$ . It is well known that if AB = BA, then  $A_{ij}$  are all upper triangular Toeplitz m atrices, i.e. for 1 i j twe have

$$A_{ij} = \begin{cases} 2 & \cdots & 0 & a_{ij}^{0} & a_{ij}^{1} & \cdots & a_{ij}^{i} & 1 \\ 6 & \cdots & 0 & a_{ij}^{0} & \ddots & \vdots & 7 \\ 6 & \cdots & 0 & a_{ij}^{0} & \ddots & \vdots & 7 \\ 7 & \cdots & 0 & \cdots & a_{ij}^{1} & 5 \end{cases} \text{ and } A_{ji} = \begin{cases} 2 & a_{ji}^{0} & a_{ji}^{1} & \cdots & a_{ji}^{i} & 7 \\ 6 & 0 & a_{ji}^{0} & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & a_{ji}^{1} & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \vdots & 7 \\ 6 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 7 & \vdots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots & \vdots \\ 8 & \vdots & \ddots & \ddots & \ddots$$

If i = j then we om it the rows or columns of zeros in A ij or A ji above.

 $A = [a_{ij}^0] \text{ where } q_{1} + 1 \text{ i; } j \text{ q.}$ 

We follow Basili [2] and introduce some further notation. For a matrix B with  $sh(B)=(1;2;\dots;t)$  we denote by  $r_B$  and  $k_i$ ,  $i=1;2;\dots;r_B$ , the numbers such that  $k_1=1$ ,  $k_i$   $k_{i+1}$  2,  $k_i$   $k_{i+1}$  1 for  $i=1;2;\dots;r_B$  1,  $k_{r_B}$  t 1. Note that  $k_i$  2 f1;2;\dots;tg. For example, if  $sh(A)=(5^3;3^2;1^3)$  and  $sh(B)=(5^3;4;3^2;2^4;1^3)$ , then  $r_A=r_B=3$ . If sh(C) 2 R (n), then  $r_C=1$ . Set  $q_0=0$ ,  $q_1=t$  and for  $q_1=t$  1;2;\dots;  $q_1=t$  2 f1;2;\dots;  $q_1=t$ 3, be such that  $q_1=t$ 4 if  $q_1=t$ 5 and  $q_1=t$ 5 and  $q_1=t$ 5. For a block matrix with blocks as in (1) we define

Lem m a 8. [2, Proposition 2.3] For an n n matrix A such that AB = BA it follows that  $A 2 N_B$  if and only if  $\overline{A}$  are nilpotent for  $= 1; 2; \ldots; 1$ .

M oreover, if  $A 2 N_B$  there exists a Jordan basis for B such that  $\overline{A}$  are all strictly upper triangular.

Therefore, when we study what are possible shapes of pairs from the set N  $_2$  (or in particular what are the indices of nilpotency in N  $_B$ ), we may consider only matrices A  $_2$  N  $_B$  such that we have  $a_{ij}^0=0$  for all 1  $_j$  i twhenever  $_i=_j$ . From now, we assume that the latter relations hold for A .

Suppose that a digraph corresponds to a generic matrix M such that M has the same pattern as a matrix A 2 N  $_{\rm B}$  . We call an (N  $_{\rm B}$  ; A )-digraph.

Let be an  $(N_B;A)$ -digraph that corresponds to  $A \ 2 \ N_B$ . Denote its vertices by (x;y), where x = 1;2;:::;t and y = 1;2;:::;t. Write blocks of A as in (1). If  $a_{ij}^k \in 0$  for some 1 i j t, then contains edges ((i;h);(j;h+k)) for h = 1;2;:::;t k. If

 $a_{ij}^k \in 0$  for some 1 j i t, then contains edges ((i;h); (j;h + k +  $_j$  i + 1)) for all h = 1;2;:::; i k.

We say that edges  $((i_1; j_1); (i_2; j_2))$  and  $((i_3; j_3); (i_4; j_4))$  are parallel if  $i_1 = i_3$ ,  $i_2 = i_4$  and  $j_1 + j_4 = j_2 + j_3$ . We call two paths  $P_1$  and  $P_2$  parallel if they consist of parallel edges.

The parts  $m_{i+1}^{r_i}$ ,  $m_{i+1}^{r_{i+1}}$  in sh (B) have to be treated dierently if  $m_{i+1} = 1$  or  $m_{i+1} = 1$ . We introduce some notation that will help us unify the treatments.

For k = 1; 2; ...; 1 1 de ne

$$s(k) = \begin{cases} r_k; & \text{ifm }_k & m_{k+1} & 2\\ r_k + r_{k+1}; & \text{ifm }_k & m_{k+1} & 1 \end{cases}$$

and  $s(1) = r_1$ .

We set  $_0=_1+1$ . Choose any k, 1 k t such that  $_{k=1}>_{k}$  and let s and z be such that  $_{k}=_{s}=_{s}1$ ,  $_{k}=_{s+1}=_{z}>_{z+1}$ . We denote by  $V_{B;k}$  the set of vertices

$$f(x;y)$$
;  $x = k;k + 1;...;z;y = 1;2;...; k and  $x = z + 1;z + 2;...;s;y = kg$ :$ 

W e call the path that contains vertices


$$V_1 = f(x;1) x = 1;2;...;kg; V_{B;k}$$
 and  $V_3 = f(x; x) x = 1;2;...;k$  1g [  $f(z; z)g$ 

the B  $_k$ -path (or B -path for short). We calls (k) the width of B  $_k$ -path (or of the set  $V_{B,k}$ ). The B  $_k$ -path consists of three subpaths P  $_i$ , i=1;2;3, such that P  $_1 \setminus P_2 = f(k;1)g$  and P  $_2 \setminus P_3 = f(z; _z)g$ . The vertices of P  $_1$  are from V  $_1$  and its edges are ((i;1); (i+1;1)) for  $i=1;2;\ldots;k$  1. The P  $_2$  consists of vertices V  $_B$   $_{jk}$  and edges ((i; j); (i+1; j)) for  $i=k;k+1;\ldots;s$  1,  $j=1;2;\ldots;k$  1 or  $i=k;k+1;\ldots;z$  1, j=k and ((s; j); (k; j+1)) for  $j=1;2;\ldots;k$  1. The subpath P  $_3$  on vertices V  $_3$  consists of edges ((i;  $_1$ ); (i+1;  $_{i+1}$ )) if  $_1=k+1$ , 1 i k 2 and ((j;  $_2$ ); (i;  $_3$ )) if  $_k=k+1$   $_3$  i <  $_3$  and if  $_3$  h j it follows that  $_3$  h =  $_3$  or  $_3$  or  $_3$  for a B  $_k$ -path P it follows that  $_3$  h =  $_3$  or  $_3$  or  $_3$  for a B  $_k$ -path P it follows that  $_3$  h =  $_3$  or  $_3$  for a B  $_k$ -path P it follows that  $_3$  h =  $_3$  or  $_3$  for a B  $_k$ -path P it follows that  $_3$  h =  $_3$  or  $_3$  for a B  $_k$ -path P it follows that  $_3$  h =  $_3$  or  $_3$  for a B  $_k$ -path P it follows that  $_3$ 

Recall that (A) (M) for A 2  $N_B$  and a generic matrix M which has the same pattern as A.ByCorollary 7 it follows that ( $N_B$ ) is less than or equal to the length of the longest path in a ( $N_B$ ; A)-digraph. The length of the B  $_k$ -path is equal to 2  $\stackrel{k}{i=1} r_i + r_k m_k + r_{k+1} m_{k+1}$  if s(k) =  $r_k + r_{k+1}$  and 2  $\stackrel{k}{i=1} r_i + r_k m_k$  otherwise.


In the next section we show that the longest path in an (N  $_{\rm B}$  ;A)-digraph  $\,\,$  is actually equal to the length of the longest B-path in  $\,$  .

Exam ple 9. Let sh (B) = (4;3 $^2$ ). By Corollary 4 it follows that (N<sub>B</sub>) = 10. The generic M , such that its digraph M equals to



and A 2 N<sub>B</sub> such that  $a_{12}^0$ ,  $a_{23}^0$ ,  $a_{31}^0$  are its only nonzero entries, have the same pattern. It is easy to verify that (M) = (A) = 10. The shown path is the only B-path in M.

Exam ple 10. In the case sh(B) =  $(4;3;2^2;1)$ , the (N<sub>B</sub>;A)-digraph has three B-paths of lengths 7 and 9, as shown below.



We will show that  $(N_B) = 9$ .

Lem m a 11. Let P be a path in an  $(N_B;A)$ -digraph that contains vertices (x;1) for x = 1;2;:::;t. Then there exists a B-path  $P_B$  such that  $P_j$   $P_B_j$ .

Proof. W rite sh(B) =  $(1; 2; ...; t) = (m_1^{r_1}; m_2^{r_2}; ...; m_1^{r_1})$ , where  $m_i = 1$  i+ 1 for i = 1; 2; ...; l and  $m_1 = t$ .

Let be an (N  $_{\rm B}$ ;A)-digraph and recall the de nitions of V  $_{\rm B}$ ;i and s(i). Let k be the maximal index such that s(k) = maxfs(i); 1 i lg. By P  $_{\rm B}$ ;k we denote the B  $_{\rm k}$ -path, i.e. the widest B -path in .

Suppose rst that  $s(k) = r_k + r_{k+1}$ , where  $r_{k+1} \in 0$ . Then

Let P be the path that contains vertices (x;1) for x=1;2;:::;t and P  $^0$  its subpath that contains all the vertices (x;y) in P , where y>1.

Exam ine the edge ((i;y<sub>i</sub>); (j;y<sub>j</sub>)) of P, where  $_{i}$  &  $_{j}$ . If i < j (and thus  $_{i} > _{j}$ ) then  $_{i}$  y<sub>i</sub> >  $_{j}$  y<sub>j</sub>. If i > j (and thus  $_{i} < _{j}$ ) then y<sub>i</sub> < y<sub>j</sub>. Since P<sub>B,k</sub> is the widest B-path in , it follows that  $_{i}$  P  $_{i=1}^{p-1}$  r<sub>i</sub> + (m<sub>1</sub> 1) (r<sub>k</sub> + r<sub>k+1</sub>). Therefore

$$\stackrel{\mbox{$\dot{x}$}^{1}$}{\mbox{$\dot{p}$}}$$
  $\stackrel{\mbox{$\dot{x}$}^{1}$}{\mbox{$\dot{z}$}}$   $\stackrel{\mbox{$\dot{x}$}}{\mbox{$\dot{z}$}}$   $\stackrel{\mbox{$\dot{x}$}}{\mbox{$\dot{z}$}}$   $\stackrel{\mbox{$\dot{z}$}}{\mbox{$\dot{z}$}}$   $\stackrel{\mbox{$\dot{z}$}}{\mbox{$\dot{z}$}}$ 

Since  $m_i = 1$  i+ 1 for  $i = 1; 2; \ldots; l$ , it follows that  $m_i = m_1 = l$  i for  $i = 1; 2; \ldots; l$ . In particular,  $m_l = m_k = l + k$ . Thus  $p \neq j = 2^{l+1} r_i + r_l + (m_k = l + k = l) (r_k + r_{k+1})$ . By de nition of k it follows that  $r_i + r_{i+1} p = r_k + r_{k+1} = l$  for all  $i = k+1; k+2; \ldots; l=1$ . Thus  $r_{k+1} + 2r_{k+2} + \ldots + 2r_{l+1} + r_l = l = l = l = l = l$  (l k l)  $(r_k + r_{k+1} = l)$ . Therefore

The case  $s(k) = r_k$  can be proved similarly, because  $P_{B;k}j = 2^{P_{k-1}} r_i + r_k m_k$  and  $P_{j} = 2^{P_{k-1}} r_i + r_l + (m_1 1) r_k$ .

Theorem 12. For A 2 N  $_{\rm B}$  it follows that

(A) 
$$\max_{1 \le t} 2i + \sum_{i+1} + \sum_{i+2} + \dots + \sum_{i+r} \sum_{i+1} \sum_{i+r} 1g$$
:

Proof. Write sh(B) = (1; 2; :::; t). We show by induction on t that the longest path in  $(N_B;A)$ -digraph is of the same length as a B-path.

Note rst that the longest path P in must contain (1;1) and  $(1; _1)$ . If it does not we can always add an edge from (1;1) to the rst vertex of P or an edge from the last vertex of P to  $(1; _1)$ .

If t = 1 then the longest path contains all vertices and therefore it is a B-path.

Suppose our claim holds for all partitions with at most t 1 parts and let P be the longest path in .

If P does not contain any of the vertices (x;y) for som e xed x and  $y=1;2;:::;_{x}$ , then P is a path in  $(N_B \circ; A)$ -digraph, where sh  $(B^0)=(_1;_2;:::;_{x}_1;_{x+1};_{x+2};:::;_{t})$ . By induction, there exists a  $B^0$ -path  $P_B \circ$  such that  $P_J : P_B \circ J$ . If a  $B^0$ -path is not already a B-path, it can be lengthened to a B-path. Therefore there exists a B-path  $P_B$  in , such that its length is greater than or equal to  $P_J$ . By assumption that  $P_J$  is the longest path in it follows that  $P_J : P_B \circ J$ .

Suppose now that for each x the path P contains a vertex  $(x;y_x)$  for som  $ey_x$ . The basic idea of the proof is to show the following claim: In there exists a path  $P^0$  of the same length as P such that its rst t vertices are  $(1;y_1)$ ,  $(2;y_2)$ ,...,  $(t;y_t)$ .

Since  $P^0$  is the longest path in it follows that  $y_i = 1$  for i = 1;2;:::;t. O therw ise the path that contains vertices (1;1), (2;1),..., (t;2),...,  $(t;y_t)$  and is after  $(t;y_t)$  equal to  $P^0$  is longer than  $P^0$ .

Let us prove the claim . Let  $y_t > 1$  be the sm allest integer such that P contains vertex  $(t;y_t)$ . Since P is the longest path, for x=1;:::;t there exists a vertex  $(x;y_x^3)$  that P visits after  $(t;y_t)$ , i.e.  $y_x^3 - y_t$ . Next, suppose that there exists some k, where 1-k < t and  $y_k^1 < y_k^2 - y_t$ , such that P contains vertices  $(k;y_k^1)$  and  $(k;y_k^2)$  and that  $y_k^2$  is such that there does not exist a vertex between  $(k;y_k^2)$  and  $(t;y_t)$  in P with k as its rst coordinate. Denote by  $P_1$  (resp.  $P_2$ ) the subpath of P with its initial vertex  $(k;y_k^1)$  (resp.  $(k;y_k^2)$ ) and its nal vertex  $(k;y_k^2)$  (resp.  $(k;y_k^3)$ ) and denote  $P_0 = P nfP_1; P_2g$ . In there exist the path  $P_1^0$  parallel to  $P_2$  starting at the  $(k;y_k^1)$  and the path  $P_2^0$  parallel to  $P_1$  ending at the  $(k;y_k^3)$ . Then, the initial vertex of  $P_2^0$  coincides with the nal vertex of  $P_1^0$ . Thus  $P_0 = P_1^0 =$ 

By repeating this swap for k=1;2;:::;t, we obtain a path  $P^0$  such that it does not contain vertices  $(x;y_x^1)$  and  $(x;y_x^2)$ , with  $y_x^1 < y_x^2$  yt for all x=1;2;:::;t 1. As argued above, this forces  $P^0$  to be a path in the  $(N_B;A)$ -digraph that contains vertices (x;1) for all x=1;2;:::;t. By Lem m a 11 there exists a B-path  $P_{B;k}$  such that  $P_{B;k}$  is  $P_{B;k}$  if  $P_{B;k}$  if  $P_{B;k}$  is  $P_{B;k}$  if  $P_$ 

Theorem 13. Let B be a nilpotent matrix such that sh(B) = (1; 2; :::; t). Then

$$(N_B) = \max_{1} \max_{i < t} f2i + \sum_{i+1} + \sum_{i+2} + \dots + \sum_{i+r} \sum_{i+1} \sum_{i+1} 1g:$$

Proof. We want to show that the maximum of Theorem 12 is attained. For the digraph with vertices (x;y),  $x=1;2;\ldots;t$ ,  $y=1;2;\ldots;t$ , recall the denition of a B<sub>k</sub>-path. Let k and s be such that maxf2i+  $_{i+1}+_{i+2}+\ldots+_{i+r};t_{i+1}-_{i+r}-1g=2k+_{k+1}+_{k+2}+\ldots+_{k+s}$ .

De ne path P as the B (k+1)-path in . It follows that  $P = 2k + k+1 + k+2 + \dots + k+s$ . The path P is de ned so that it can be completed (by drawing parallel edges) to an  $(N_B;A)$ -digraph. (For examples of such  $(N_B;A)$ -digraphs see Example 14.) By the proof of the Theorem 12 it follows that P is a longest path in .

For a generic matrix M , corresponding to , it follows that (M ) =  $\mathcal{P}$  j. Take A 2 N<sub>B</sub> to be a positive matrix with the same pattern as M . Thus  $\mathbb{A}^h|_{i,j}=0$  if and only if  $\mathbb{M}^h|_{i,j}=0$  for all h, i, j. Therefore (A) = (M) = 2k +  $_{k+1}$  +  $_{k+2}$  + :::+  $_{k+s}$  which proves the theorem .


### 5 Examples

Exam ple 14. As in Exam ple 10, we consider again the case  $sh(B) = (4;3;2^2;1)$ . By Theorem 13 it follows that  $(N_B) = m axf7;9;9g = 9$ . There exist matrices  $A_i \ge N_B$ ,

i = 1;2;3, with dierent shapes (and patterns) such that  $(A_i) = 9$ . Namely

and

The generic m atrices M  $_1$ , M  $_2$  and M  $_3$  w ith the same patterns as A  $_1$ , A  $_2$  and A  $_3$  have the following digraphs



It can be easily seen that  $\operatorname{sh}(A_1) = \operatorname{sh}(M_1) = (9;1^3)$ ,  $\operatorname{sh}(A_2) = \operatorname{sh}(M_2) = (9;2;1)$  and  $\operatorname{sh}(A_3) = \operatorname{sh}(M_3) = (9;3)$ .

Exam ple 15. Not all = ( $_1$ ;  $_2$ ;:::;  $_k$ ) 2 P (n), where  $_1$  = ( $N_B$ ), are in P ( $N_B$ ). For sh (B) = (6;4) it follows that ( $N_B$ ) = 6. We used M athematica to show that (6;3;1)  $\not\equiv$  P ( $N_B$ ).

Exam ple 16. Not all partitions in P (N  $_{\rm B}$ ) can be obtained from a generic matrix corresponding to an (N  $_{\rm B}$ ; A)-digraph. Let sh(B) = (5;3). We veri ed using M athematica that

One can check that there is no  $(N_B;A)$ -digraph with () equal to  $(4^2)$ ,  $(3^2;1^2)$  or  $(2^4)$ . However, there are matrices with these shapes in  $(N_B;A)$ . For example, for

we have  $\operatorname{sh}(A_1) = (2^4)$ ,  $\operatorname{sh}(A_2) = (3^2; 1^2)$ . Let  $M_i$ , i = 1; 2, be generic m atrices of the same shape as  $A_i$ . It can be proved, using G ansner-Saks theorem, that  $\operatorname{sh}(M_1) = \operatorname{sh}(M_2) = (5; 3)$ .

Note also that  $(4;3;1) \not\geq P$  (N<sub>B</sub>) and thus not all partitions  $(1;2;:::;k) \geq P$  (8) with 1 5 are in P (N<sub>B</sub>). Observe that (5;3) is a maximal partition in P (N<sub>B</sub>) with respect to the standard partition ordering. (See [4, Section 6.2] for more details.) By Gerstenhaber-Hesselink Theorem, the orbit corresponding to (4;3;1) is in the closure of the orbit corresponding to (5;3). Hence, it follows that N<sub>B</sub> need not intersect all the orbits in the closure of a maximal orbit that it does intersect.

### R eferences

- [1] V.Baranovsky: The variety of pairs of commuting nilpotent matrices is irreducible, Transform. Groups 6 (2001), no. 1, 3{8.
- [2] R.Basili: On the irreducibility of commuting varieties of nilpotent matrices, J.Algebra 268 (2003), 58-80.
- [3] R.A.Brualdi, H.J.Ryser: Combinatorial matrix theory, Cambridge University Press, 1991.
- [4] D.H.Collingwood, W.M.McGovern: Nilpotent orbits in semisimple Lie algebras, Van Nostrand Reinhold, 1993.
- [5] E.R. Gansner: Acyclic digraphs, Young tableaux and nilpotent matrices, SIAM J.A lgebraic Discrete Methods 2 (1981), no. 4, 429 (440.

- [6] R.Guralnick: A note on commuting pairs of matrices, Linear and Multilinear Algebra 31 (1992), no. 1-4, 71 (75.
- [7] R. Guralnick, B. A. Sethuram an: Commuting pairs and triples of matrices and related varieties, Linear Algebra Appl. 310 (2000) 139{148.
- [8] W . H . Gustafson: M odules and matrices, Linear Algebra Appl. 157 (1991), 3-19.
- [9] Y. Han: Commuting triples of matrices, Electron. J. Linear Algebra 13 (2005), 274-343.
- [10] J.Holbrook, M.Omladic: Approximating commuting operators, Linear Algebra Appl. 327 (2001), no. 1-3, 131 (149.
- [11] T.Kosir: The Cayley-Ham ilton theorem and inverse problems for multiparam eter systems, Linear Algebra Appl. 367 (2003), 155-163.
- [12] T.S.M otzkin, O. Taussky: Pairs of matrices with property L, Trans. Amer. Math. Soc. 73, (1952), 108{114.
- [13] M. Om ladic: A variety of commuting triples, Linear Algebra Appl. 383 (2004), 233{245.
- [14] M . Saks: Some sequences associated with combinatorial structures, D iscrete M ath.59 (1986), no.1-2, 135{166.