

PARAMETER TEST IDEALS OF COHEN MACAULAY RINGS

MORDECHAI KATZMAN

1. Introduction

The aim of this paper is to provide a description of parameter test ideals of local Cohen-Macaulay rings of prime characteristic p . The nature of this description will be such that it will allow us to give an algorithm for producing these ideals.

These descriptions will follow from the study of a certain Frobenius map on the injective hull of residue fields. In the complete case, which is the main case, Matlis duality will enable us to translate the Frobenius map above to a study of maps of quotients of regular rings to their images under the Frobenius functor.

2. Frobenius maps

Let S be a commutative ring of prime characteristic p and let M be an S -module. A Frobenius map on M is a \mathbb{Z} -linear map $\varphi: M \rightarrow M$ with the property that $\varphi(sm) = s^p \varphi(m)$ for all $s \in S$ and $m \in M$. The fundamental example of a Frobenius map is the Frobenius map $f: S \rightarrow S$ given by $f(s) = s^p$. The Frobenius map allows us to endow S with a structure of an S -bimodule: as a left S -module it has the usual S -module structure whereas S acts on itself on the right via the Frobenius map. We shall denote this bimodule $F_S(S)$ and so for all $a \in F_S(S)$ and $s \in S$, $sa = s^p a$ while $a \cdot s = a$, where \cdot denotes the action of S . We can extend this construction to obtain the Frobenius functor F sending any S -module M to $F_S(M) = F_S(S) \otimes_S M$ where S acts on $F_S(M)$ via its left-action on $F_S(S)$, so for $s \cdot m \in F_S(M)$ and $t \in S$ we have $t(s \cdot m) = ts \cdot m$ and $(s \cdot tm) = s \cdot t \cdot m = s^p t \cdot m$. We shall repeatedly (and tacitly) use the fact that the functor F_S is exact whenever S is regular (Theorem 2.1 in [K]).

Iterations $\varphi^e = \underbrace{\varphi \circ \dots \circ \varphi}_{e \text{ times}}$ of Frobenius maps $\varphi: M \rightarrow M$ result in maps $\varphi^e: M \rightarrow M$ which satisfy $\varphi^e(sm) = s^{p^e} \varphi^e(m)$ for all $s \in S$ and $m \in M$. More generally, we will consider the set $F^e(M)$ of all \mathbb{Z} -linear maps $\varphi: M \rightarrow M$ which satisfy $\varphi(sm) = s^{p^e} \varphi(m)$ for all $s \in S$ and $m \in M$. We can give $F^e(M)$ the structure of an S -module: for any $\varphi \in F^e(M)$ and $a \in S$ we simply let $a \cdot \varphi$ be the map sending $m \in M$ to $a \cdot \varphi(m)$. Furthermore, we can define a product in $F(M) := F^e(M)$ to be composition of \mathbb{Z} -linear maps and thus endow $F(M)$ with the structure of an S -algebra.

The iteration of the Frobenius map on R leads one to the iterated Frobenius functors $F_R^i(\cdot)$ which are defined for all $i \geq 1$ recursively by $F_R^1(\cdot) = F_R(\cdot)$ and $F_R^{i+1}(\cdot) = F_R \circ F_R^i(\cdot)$ for all $i \geq 1$. These higher Frobenius functors are also exact whenever S is regular.

In this paper we will be interested in studying Frobenius maps on injective hulls of residue fields and top local cohomology modules. An example of the latter when S is local and d -dimensional can be obtained as follows. The top local cohomology module $H_m^d(S)$ can be computed as the direct limit of

$$\frac{S}{(x_1, \dots, x_d)S} \xrightarrow{x_1} \frac{S}{(x_1^2, \dots, x_d)S} \xrightarrow{x_1} \frac{S}{(x_1^3, \dots, x_d)S} \xrightarrow{x_1} \dots$$

and we can define a Frobenius map $\varphi: H_m^d(S) \rightarrow H_m^d(S)$ on this direct limit by mapping the coset $a + (x_1^n, \dots, x_d^n)S$ in the n -th component of the direct limit to the coset $a^{p^e} + (x_1^{np^e}, \dots, x_d^{np^e})S$ in the np^e -th component of the direct limit. It is not hard to verify that this is indeed a well defined map from $H_m^d(S)$ to $H_m^d(S)$ and that it is a Frobenius map. An important observation used in this paper is the fact that when S is Cohen-Macaulay, the Frobenius map $\varphi: F^1(H_m^d(S)) \rightarrow F^1(H_m^d(S))$ described above generates the S -algebra $F(H_m^d(S))$ (cf. Example 3.7 in [LS]).

A different and fruitful way of thinking about Frobenius maps on M and their iterations is as left-module structures over certain skew-commutative rings. Given any commutative ring S we can construct a skew-commutative ring $S[T; f^e]$ as follows: as an S -module it will be the free module $\bigoplus_{i=0}^{\infty} S T^i$ and we extend the rule $Ts = s^{p^e} T$ for all $s \in S$ to a (non-commutative!) multiplicative structure on $S[T; f^e]$. Given a Frobenius map $\varphi: F^e(M) \rightarrow F^e(M)$ on an S -module M , we can then turn it into a left $S[T; f^e]$ -module by extending the rule $Tm = \varphi(m)$ for all $m \in M$. The fact that this gives M the structure of a left $S[T; f^e]$ -module is simply because for all $s \in S$ and $m \in M$,

$$T(sm) = (sm) = s^{p^e} \varphi(m) = s^{p^e} Tm = (Ts)m.$$

This approach has been taken in many previous papers, the most relevant to us being [S].

3. A duality

In this section we set up the main tool, based on Matlis Duality, which will enable us to explore given $R[T; f]$ -module structures of certain Artinian modules.

Henceforth in this paper $(R; m)$ will denote a complete local regular ring of characteristic p . We shall denote the injective hull of R/m with E and $(-)^\vee$ shall denote the functor $\text{Hom}_R(-; E)$.

Let M be any R -module and for all $m \in M$, let $e_m \in M^\vee$ be defined by $e_m(g) = g(m)$ for all $g \in M^\vee$. Matlis duality states that for all R -modules M which are either Noetherian or Artinian, the map $M \rightarrow M^\vee$ which sends $m \in M$ to e_m is an isomorphism of R -modules. If now M is an $R[T; f^e]$ -module, this map endows M^\vee with a structure of an $R[T; f^e]$ -module defined by $T e_m = e_{Tm}$ for all $m \in M$, so now we may identify M and M^\vee as $R[T; f^e]$ -modules.

Since R is complete, a straightforward modification of Lemma 4.1 in [L] provides us with a natural, functorial isomorphism $\varphi_M^e: F_R^e(M)^\vee \rightarrow F_R^e(M^\vee)$ defined for all Artinian R -modules. We shall use this isomorphism repeatedly in this section.

Fix now an ideal $I \subseteq R$ and write $S = R/I$. Let C^e be the category of Artinian $S[[T; f^e]]$ -modules. Let D^e be the category of R -linear maps $M \rightarrow F_R^e(M)$ where M is a finitely generated S -module and where a morphism between $M \xrightarrow{a} F_R^e(M)$ and $N \xrightarrow{b} F_R^e(N)$ is a commutative diagram of S -linear maps

$$\begin{array}{ccc} M & \xrightarrow{\quad} & N \\ \downarrow a & & \downarrow b \\ F_R^e(M) & \xrightarrow{F_R^e(\quad)} & F_R^e(N) \end{array}$$

In this section we construct a pair of functors $\tau^e : C^e \rightarrow D^e$ and $\tau_e^e : D^e \rightarrow C^e$ in a way that for all $M \in C^e$, the $S[[T; f^e]]$ -module $\tau_e^e \tau^e(M)$ is canonically isomorphic to M and for all $D = (N \xrightarrow{b} F_R^e(N)) \in D^e$, $\tau^e \tau_e(D)$ is canonically isomorphic to D .

The functor τ^e is the "first step" in the construction of Gennady Lyubeznik's functor $H_{R,S}$, i.e., for $M \in C^e$ we have an R -linear map $\tau_M^e : F_R^e(M) \rightarrow M$ given by $(r \cdot m) = rTm$ for all $r \in R$ and $m \in M$. Applying $(\quad)^{-}$ to the map τ_M^e one obtains an R -linear map $\bar{\tau}_M^e : M \rightarrow F_R^e(M)^{-}$. We now define $\tau(M)$ to be the map $M \xrightarrow{\bar{\tau}_M^e} F_R^e(M)^{-}$.

To define τ_e^e we retrace the steps above; given a finitely generated S -module N and a R -linear map $a : N \rightarrow F_R^e(N)$ we define $\tau_e^e(a)$ to coincide with the functor $(\quad)^{-}$ as a functor of S -modules giving $\tau_e^e(N)$ the additional structure of an $S[[T; f^e]]$ -module structure as follows.

We apply τ to the map a above to obtain a map $a^- : F_R^e(N)^{-} \rightarrow N^-$. We next obtain a map $\tau : F_R^e(N^-) \rightarrow F_R^e(N)^{-}$ as the following composition:

$$F_R^e(N^-) = F_R^e(N^-)^{-} \xrightarrow{(\bar{\tau}_N^-)^{-1}} F_R^e(N^{--})^- = F_R^e(N)^{-} :$$

We now obtain a functorial map $b = a^- : F_R^e(N^-) \rightarrow N^-$ and we define the action of T on N^- by defining $Tn = b(1 - n)$ for all $n \in N^-$.

Theorem 3.1. The functors $\tau^e : C^e \rightarrow D^e$ and $\tau_e^e : D^e \rightarrow C^e$ are exact. For all $M \in C^e$, the $S[[T; f]]$ -module $\tau_e^e \tau^e(M)$ is canonically isomorphic to M . For all $D = (N \xrightarrow{b} F_R^e(N)) \in D^e$, $\tau^e \tau_e(D)$ is canonically isomorphic to D .

Proof. The exactness of the functors follows from the exactness of the functors $\text{Hom}_R(\quad; E)$ and F_R^e .

To prove the second statement we notice that for all $M \in C^e$, $\tau_e^e \tau^e(M)$ is M^{--} which we identify as an S -module with M by identifying each $m \in M$ with the $e_m \in M^{--}$ which we defined at the beginning of this section. We want to show that this identification is an isomorphism of $S[[T; f^e]]$ -modules, and to do so we now describe Te_m for all $e_m \in \tau_e^e \tau^e(M)$. This will be the image of $1 - e_m$ under the map

$$(1) \quad F_R^e(M^{--}) \xrightarrow{i_1} F_R^e(M^{--})^{--} \xrightarrow{(\bar{\tau}_M^{--})^{-1}} F_R^e(M^{---})^- \xrightarrow{i_2} F_R^e(M^-)^- \xrightarrow{\bar{m}} F_R^e(M)^{-} \xrightarrow{\tau} M^{--}$$

where $i_1; i_2$ are the isomorphisms induced from the isomorphism of functors $(\) = (\)^{--}$. The functoriality of $(\)$ implies that we have a commutative diagram

$$\begin{array}{ccc} F_R^e(M^{---}) & \xrightarrow{i_2} & F_R^e(M^-) \\ \downarrow \bar{m}^{--} & & \downarrow \bar{m} \\ F_R^e(M^{--}) & \xrightarrow{i_1^{-1}} & F_R^e(M)^{--} \end{array}$$

We may now rewrite the composition in (1) as

$$F_R^e(M^{--}) \xrightarrow{i_1^{-1}} F_R^e(M^{--})^{--} (\bar{m}^{--})^1 \xrightarrow{i_2} F_R^e(M^{---}) \xrightarrow{\bar{m}^{-1}} F_R^e(M^{--}) \xrightarrow{i_1^{-1}} F_R^e(M)^{--} \xrightarrow{\bar{m}^{-1}} M^{--}$$

which simplifies into

$$F_R^e(M)^{--} \xrightarrow{\bar{m}^{-1}} M^{--} :$$

If we now start with $D = (N \xrightarrow{i_1} F_R^e(N)) \otimes D^e$

$$e \quad e(D) : N^{--} \rightarrow F^e(N^{--})$$

is given by the composition

(2)

$$N^{--} \xrightarrow{a^{-1}} F_R^e(N)^{--} \xrightarrow{i_2} F_R^e(N^{--})^{--} ((\bar{n}^{--})^1)^{-1} \xrightarrow{i_1} F_R^e(N^-)^{--} \xrightarrow{i_1^{-1}} F_R^e(N^-) \xrightarrow{n^{-1}} F_R^e(N^{--})$$

where $i_1; i_2$ are the isomorphisms induced from the isomorphism of functors $(\) = (\)^{--}$.

Now $(\bar{n}^{--})^1 = (\bar{n}^{--})^1$ and the functoriality of $(\)$ implies that we have a commutative diagram

$$\begin{array}{ccc} F_R^e(N^{--}) & \xrightarrow{(\bar{n}^{--})^1} & F_R^e(N^-)^{--} \\ \downarrow i_3^{-1} & & \downarrow i_4 \\ F_R^e(N^{--}) & \xrightarrow{n^{-1}} & F_R^e(N^-) \end{array}$$

We may now rewrite the composition in (2) as

$$N^{--} \xrightarrow{a^{-1}} F_R^e(N)^{--} \xrightarrow{i_2} F_R^e(N^{--})^{--} \xrightarrow{i_3^{-1}} F_R^e(N^{--}) \xrightarrow{n^{-1}} F_R^e(N^-) \xrightarrow{n^{-1}} F_R^e(N^{--})$$

which simplifies to a^{-1} .

Throughout this paper, when $e = 1$ we will drop the subscript e from our notation. Thus $C^1 = C$, $1^1 = 1$, etc.

As mentioned before, the functor F_R is a building block for another functor described in section 4 of [L]. This functor, denoted with $H_{R;S}$, is a functor from C to the category of F -finite F_R -modules (see sections 1,2 and 3 in [L] for definition and properties) and is obtained as follows. For $M \in C$ write $(M) = (N \xrightarrow{i_1} F_R(N))$. Now $H_{R;S}(M)$ is defined to be the direct limit of

$$(N \xrightarrow{i_1} F_R(N)) \xrightarrow{F_R(u)} F_R^2(N) \xrightarrow{F_R^2(u)} \dots$$

Various useful properties of Lyubeznik's functor can be found in section 4 of [L].

4. Frobenius maps on injective hulls

Henceforth in this paper we shall fix an ideal $I \subseteq R$ and denote R/I with S .

In this section we will first apply the tools developed in section 3 to yield a description of possible $S[T; f^e]$ -module structures of $E_S = E_S(S = mS)$, the injective hull of the residue field of S . This description is not new: it is contained in Proposition 5.2 in [LS]. Later in the section we shall use this description to describe explicitly the nilpotent elements of $E_S(S = mS)$.

Proposition 4.1. The $S[T; f^e]$ -module structures on E_S are given by

$${}^e R = I \rightarrow R = I^{[p^e]}$$

where the map above is given by multiplication by some $g \in I^{[p^e]} : I$.

Proof. Clearly, all R -linear maps $R/I \rightarrow R/I^{[p^e]}$ are given by multiplication by some $g \in I^{[p^e]} : I$. The proposition now follows from Theorem 3.1 and the fact that $E_S = S$.

The bijection between R -linear maps $R/I \rightarrow R/I^{[p^e]}$ and $S[T; f^e]$ -module structures on E_S has been described explicitly in Chapter 3 of [B] as follows. First, notice that E_S , thought of as the direct limit of

$$\frac{R}{(y_1, \dots, y_n)R} \xrightarrow{y_1 \cdots y^y} \frac{R}{(y_1^2, \dots, y_n^2)R} \xrightarrow{y_1 \cdots y^y} \cdots$$

where y_1, \dots, y_n is a system of parameters for R , has a natural Frobenius map given by

$$(x + (y_1^s, \dots, y_n^s)) \mapsto x^p + (y_1^{sp}, \dots, y_n^{sp}) \in \frac{R}{(y_1^{sp}, \dots, y_n^{sp})R} :$$

Now if $u \in (I^{[p]} :_R I)$, u , which is also a Frobenius map on E_S , will restrict to a Frobenius map on $E_S = \text{ann}_E I$ because for all $m \in \text{ann}_E I$,

$$Iu(m) \subseteq I^{[p]}(m) = (Im) = (0) = 0 :$$

In Chapter 3 of [B] it is shown that all Frobenius maps on E_S are obtained in this way.

Henceforth in this section we shall assume that E_S has a given $S[T; f]$ -module structure. Our next aim is to describe the $S[T; f]$ -submodules of E_S . Later in the section we shall use this description to describe explicitly the nilpotent elements of E_S . We start by recalling that the set of S -submodules of E_S is $\text{fann}_{E_S} J \mid J \subseteq S$ (cf. Theorem 5.21 in [SV].) If we now asked for a description of the $S[T; f]$ -submodules of E_S , the answer would obviously be "all $\text{ann}_{E_S} J$ which happen to be $S[T; f]$ -submodules of E_S ". With this in mind we define the following.

Definition 4.2. An ideal $J \subseteq S$ is called an E_S -ideal if $\text{ann}_{E_S} J$ is an $S[T; f]$ -submodule of E_S . An ideal $J \subseteq R$ is called an E_S -ideal if it contains I and its image in S is an E_S -ideal.

Notice that for an ideal $J \subseteq S$, being an E_S -ideal is equivalent to $\text{ann}_{E_S} J = \text{ann}_{E_S} JS[T; f]$. We also note that when S is Gorenstein the notion of E_S -ideals coincides with that of F -ideals studied in [Sm1].

Theorem 4.3. Let $u \in R$ be such that (E_S) is the map $\frac{R}{I} \xrightarrow{u} \frac{R}{I^{[p]}}$. The E_S -special ideals consist of all ideals $L \subseteq R$ containing I for which $uL \subseteq L^{[p]}$.

Proof. Assume first that L is E_S -special and hence an E -ideal. Apply the functor to the short exact sequence of $S[\Gamma; f]$ -modules

$$(3) \quad 0 \rightarrow \text{ann}_{E_S} L \rightarrow E_S \rightarrow E = \text{ann}_{E_S} L \rightarrow 0$$

to obtain the following short exact sequence in D

$$(4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \frac{L}{I} & \longrightarrow & \frac{R}{I} & \longrightarrow & \frac{R}{L} \longrightarrow 0 \\ & & \downarrow u & & \downarrow u & & \downarrow u \\ 0 & \longrightarrow & \frac{L^{[p]}}{I^{[p]}} & \longrightarrow & \frac{R}{I^{[p]}} & \longrightarrow & \frac{R}{L^{[p]}} \longrightarrow 0 \end{array}$$

and we must have $uL \subseteq L^{[p]}$.

On the other hand, if $uL \subseteq L^{[p]}$, we can construct the commutative diagram (4), and an application of the functor gives back the short exact sequence (3) and we deduce that L is an E -ideal and hence a special E_S ideal.

We now turn our attention to the nilpotent elements of E_S , i.e., to the $S[\Gamma; f]$ -submodule of E_S

$$\text{Nil}(E_S) = \{m \in E_S \mid T^e m = 0 \text{ for some } e \geq 0\}.$$

Recall that we can write $\text{Nil}(E_S)$ as $\text{ann}_{E_S} JS[\Gamma; f]$ for some E_S -ideal $J \subseteq R$. Also, it is known that there exists an $\ell \geq 1$ such that $T^\ell \text{Nil}(E_S) = 0$ (cf. [HS, Proposition 1.11] and [L, Proposition 4.4]). This invariant of S plays an important role in the study of the Frobenius closure (see [KS] and [KSY]). We now describe the ideal J and the index of nilpotency ℓ .

Definition 4.4. For all $e \geq 1$ write $e = 1 + p + \dots + p^{q-1}$.

Proposition 4.5. Let the map $(E_S) = (R=I \rightarrow R=I^{[p]})$ be given by multiplication by $u \in R$. Consider E_S as an $S[\Gamma; f^e]$ -module where for all $m \in E_S$, we define $T^e m = T^e m$. The map $(E_S)^e = (R=I \rightarrow R=I^{[p^e]})$ is given by multiplication by u^e .

Proof. For all $e \geq 1$ the R -linear map $\phi_e : F^e(E_S) \rightarrow E_S$ defined by $\phi_e(r \cdot m) = r T^e m$ can be factored as $\phi_e = \phi_{e-1} \circ \psi_e$ where for all $1 \leq i \leq e$, ψ_i is the R -linear map $\psi_i : F^i(M) \rightarrow F^{i-1}(M)$ defined by $\psi_i(r \cdot m) = r \cdot T m$. Also, it is not hard to see that $\psi_{i+1} = F(\psi_i)$ for all $1 \leq i \leq e$.

Now for all $e \geq 1$, the map $(E_S)^e = (R=I \rightarrow R=I^{[p^e]})$ is given by $\frac{R}{E_S} \xrightarrow{u^e} \frac{R}{I^{[p^e]}}$. It follows from the construction of $\frac{R}{E_S}$ that, if we identify E_S with $R=I$, $\frac{R}{E_S} \xrightarrow{u^e} R=I^{[p^e]}$ is the identity map. Now

$$\phi_e = \frac{R}{E_S} \quad \psi_1 = F^e(u) \quad F^{e-1}(u) \quad u^{[p^e]} u \quad u^e = u$$

Theorem 4.6. Let the map $(E_S) = (R=I \rightarrow R=I^{[p]})$ be given by multiplication by $u \in R$. For all $e \geq 1$ let J_e be the smallest ideal of R which contains I and such that $u^e \in J_e^{[p^e]}$. There exists an $e_0 \geq 1$ such that $J = J_{e_0+1}$ and for this e_0 , $\text{ann}_{E_S} J$ coincides with the $S[[f]]$ -module $\text{Nil}(E_S)$ of nilpotent elements of E_S . Furthermore, the index of nilpotency of $\text{Nil}(E_S)$, if not zero, is the smallest such e .

Proof. For all $e \geq 1$ let $N_e = \{m \in E_S \mid T^e m = 0\}$ and write $N_e = \text{ann}_{E_S} L_e$ for some E_S -ideal L_e .

Notice that $(N_e) = (R=L_e \rightarrow R=L_e^{[p^e]})$ and that the previous proposition implies that this map is given by multiplication by u^e . It follows from the construction of $(N_e) = (R=L_e \xrightarrow{u^e} R=L_e^{[p^e]})$ that this map is the zero map, i.e., $u^e \in L_e^{[p^e]}$; now the minimality of J_e implies that $J_e \subseteq L_e$ and $\text{ann}_{E_S} L_e \subseteq \text{ann}_{E_S} J_e$.

On the other hand, the map $R=J_e \xrightarrow{u^e} R=J_e^{[p^e]}$ is the zero map and so T^e kills

$$u^e R=J_e \xrightarrow{u^e} R=J_e^{[p^e]} = \text{ann}_{E_S} J_e$$

hence $\text{ann}_{E_S} J_e \subseteq N_e = \text{ann}_{E_S} L_e$ and we deduce that $\text{ann}_{E_S} J_e = N_e$.

Proposition 4.4 in [L] now implies that $\text{Nil}(E_S) = N$ for some $e \geq 1$, and since $\text{Nil}(E_S)$ is the union of the ascending chain $fN \subseteq g_1$, we see that $N = N$ for all e . Also, if $N = N_{e+1}$ but $\text{Nil}(E_S) \not\subseteq N$, pick any non-zero $m \in N \setminus \text{Nil}(E_S)$ and let $i \geq 0$ be minimal such that $T^i m \notin N$. Now $T^{e+1}(T^i m) = T^{i+e+1} m = 0$ so $T^i m \in N_{e+1} \subseteq N$, a contradiction.

We shall see in section 5 how to compute this smallest ideal $J = I$ for which $u^e \in J^{[p^e]}$.

We conclude this section by exhibiting another "naturally occurring" $S[[f]]$ -submodule of E_S .

Theorem 4.7. Let the map $(E_S) = (R=I \rightarrow R=I^{[p]})$ be given by multiplication by $u \in R$. For all $i \geq 0$ write $L_i = I^{[p^i]} :_R u$.

- (a) The sequence of ideals $fL_i \subseteq g_1$ is an ascending sequence.
- (b) If $L_A = L_{A+1}$ then $L_i = L_A$ for all $i \geq A$ and L_A is an E_S -ideal.
- (c) Let L be the stable value of $fL_i \subseteq g_1$. The quotient $E_S = \text{ann}_{E_S} L$ is nilpotent and for any E_S -ideal K , $E_S = \text{ann}_{E_S} K$ is nilpotent if and only if $K \subseteq L$.

Proof. For all $i \geq 1$ let the map $g_i : R=I \rightarrow R=I^{[p^i]}$ given by the composition

$$R=I \xrightarrow{u} R=I^{[p]} \xrightarrow{u^p} R=I^{[p^2]} \xrightarrow{u^{p^2}} \cdots \xrightarrow{u^{p^{i-1}}} R=I^{[p^i]}$$

is just the map $g_i : R=I \xrightarrow{u^i} R=I^{[p^i]}$ given by multiplication by u^i and whose kernel is L_i . These kernels form an ascending chain, so (a) follows.

The first statement in (b) follows from Proposition 2.3(b) in [L]. To prove the second statement, we first notice that since $uI \subseteq I^{[p]}$, $u^p I^{[p]} \subseteq I^{[p^{+1}]}$ for all $i \geq 1$ hence $u^i I \subseteq I^{[p^i]}$ and we deduce that $I \subseteq L$ for all $i \geq 1$. To show that L_A is an E_S -ideal it

remains to prove that $uL_A = L_A^{[p]}$:

$$\begin{aligned} uL_A &= uL_{A+1} \\ &= I^{[p^{A+1}]} \otimes_R u^{p^2 + \dots + p^A} \\ &= I^{[p^A]} \otimes_R u^{N_A \cdot p} \\ &= I^{[p^A]} \otimes_R u^{N_A \cdot [p]} \\ &= L_A^{[p]} \end{aligned}$$

where the penultimate equality is a consequence of the exactness of $F_R(\quad)$.

Let K be an E_S -ideal for which $E_S = \text{ann}_{E_S} K$ is nilpotent and choose some $e \geq 1$ for which $T^e(E_S = \text{ann}_{E_S} K) = 0$. An application of T^e to the short exact sequence

$$0 \rightarrow \text{ann}_{E_S} K \rightarrow E_S \rightarrow E_S / \text{ann}_{E_S} K \rightarrow 0$$

produces the following short exact sequence in D^e

$$\begin{array}{ccccccc} 0 & \longrightarrow & K=I & \longrightarrow & R=I & \longrightarrow & R=K \longrightarrow 0 \\ & & \downarrow u^e & & \downarrow u^e & & \downarrow u^e \\ 0 & \longrightarrow & K^{[p^e]}=I^{[p^e]} & \longrightarrow & R=I^{[p^e]} & \longrightarrow & R=K^{[p^e]} \longrightarrow 0 \end{array}$$

where the leftmost vertical map is the zero map, i.e., $u^e K = I^{[p^e]}$ and hence $K = L_e$.

5. The φ -closure

The statements of Theorems 4.6 and 4.7 in the previous section referred to certain smallest ideals $J \subseteq R$ with the property that $J^{[p^e]}$ contains a given ideal. The aim of this section is to describe an algorithm for computing these ideals.

Throughout this section T will denote a Noetherian regular ring of prime characteristic p .

Definition 5.1. Let $e \geq 1$. For any ideal $A \subseteq T$ we define

$$G^e(A) = \bigcap_{j=1}^n L_j L \quad \text{if } A \subseteq L \text{ is an ideal, } A \subseteq L^{[p^e]}$$

$$\text{and } L_e(A) = \bigcap_{L \subseteq G^e(A)} L.$$

Note that in general there is no reason why $L_e(A)$ should be in $G^e(A)$. Recall that an T -module M is φ -at if it is φ -at and if for all sets of T -modules $\{N_i\}_{i \in \mathbb{Z}_2}, M \subseteq T \setminus \bigcup_{i \in \mathbb{Z}_2} N_i = \bigcup_{i \in \mathbb{Z}_2} M \subseteq T \setminus N_i$ (cf. [HH2] pp. 41.) Notice that free modules are φ -at.

Proposition 5.2. Let $e \geq 1$ and assume that $T^{1=p^e}$ is a φ -at T -module. Let $A \subseteq T$ be an ideal.

- (a) $L_e(A) \subseteq G^e(A)$ and is the minimal element of $G^e(A)$.

(b) Let $B \subseteq T$ be any ideal. The smallest ideal $J \subseteq T$ which contains both $A^{[p^e]}$ and B is $I_e(A) + B$.

(c) If $A = A_1 + \dots + A_s$ then $I_e(A) = I_e(A_1) + \dots + I_e(A_s)$.

Proof. The first statement is an immediate consequence of the fact that the T -module $T^{1=p^e}$ is assumed to be flat. The second statement is straightforward.

An easy induction reduces the proof of (c) to the case $s = 2$. Now $A_1, A_2 \subseteq A$ so $A_1, A_2 \subseteq I_e(A)^{[p^e]}$, and the minimality of $I_e(A_1)$ and $I_e(A_2)$ now implies $I_e(A_1) + I_e(A_2) \subseteq I_e(A)$ hence $I_e(A_1) + I_e(A_2) = I_e(A)$. On the other hand $A = A_1 + A_2 \subseteq I_e(A_1)^{[p^e]} + I_e(A_2)^{[p^e]} = (I_e(A_1) + I_e(A_2))^{[p^e]}$ and the minimality of $I_e(A)$ implies $I_e(A) \subseteq I_e(A_1) + I_e(A_2)$.

Notice that if T is a polynomial ring $K[x_1, \dots, x_n]$ for some field K or a localization of it, then $T^{1=p^e}$ is a free T -module and hence flat. Throughout the remainder of this section we will assume that $T = K[x_1, \dots, x_n]$ or that $T = K[x_1, \dots, x_n]_{(p)}$ for some field K of prime characteristic p . We also assume $n \geq 1$.

The previous proposition reduces the calculation of $I_e(A)$ to the case where A is principal, and this is the content of the next proposition. This proposition has been proved in [ABL] and we reproduce the proof for the reader's convenience. We first define

$$f = (x_1, \dots, x_n)^{2N^n} \prod_{i=1}^n x_i^{p^e - 1}$$

and for each $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ let x^α denote the monomial $x_1^{\alpha_1} \dots x_n^{\alpha_n}$. Observe that, if \mathcal{B} is a basis of K as a K^{p^e} -vector space,

$$B = \{x^\alpha \mid \alpha \in \mathbb{N}^n, \alpha_i < p^e\}$$

is a free basis for the T^{p^e} -module T .

Proposition 5.3. Let $g \in T$; using the free basis B above write $g = \sum_{b \in B} g_b^{p^e} b$ where $g_b \in T$ for all $b \in B$. Then $I_e(gT)$ is the ideal generated by $fg_b \mid b \in B$.

Proof. If $L \subseteq T$ is such that $g = \sum_{b \in B} g_b^{p^e} b \in L^{[p^e]}$ then we can find $\alpha_1, \dots, \alpha_s \in \mathbb{N}^n$ and $r_1, \dots, r_s \in T$ such that $\sum_{b \in B} g_b^{p^e} b = \sum_{i=1}^s r_i x^{\alpha_i}$. For all $1 \leq i \leq s$ we can now write $r_i = \sum_{b \in B} r_{b,i}^{p^e} b$ where $r_{b,i} \in T$ for all $b \in B$ and we obtain

$$\sum_{b \in B} g_b^{p^e} b = \sum_{b \in B} \sum_{i=1}^s r_{b,i}^{p^e} x^{\alpha_i} b$$

Since these are direct sums, we may compare coefficients and deduce that for all $b \in B$, $g_b^{p^e} = \sum_{i=1}^s r_{b,i}^{p^e} x^{\alpha_i}$ hence $g_b = \sum_{i=1}^s r_{b,i} x^{\alpha_i}$ and $g_b \in L$. On the other hand, if $g_b \in L$ for all $b \in B$ we clearly have $g = \sum_{b \in B} g_b^{p^e} b \in L^{[p^e]}$, so we have shown that $I_e(gT)$ is the ideal generated by $fg_b \mid b \in B$.

We can now restate one of the statements of Theorem 4.6 as follows: The index of nilpotency of E_s , if not zero, is the index at which the ascending sequence of ideals $fI_e(u_e + 1) + Ig_{e-1}$ stabilizes.

Notice that when T is a polynomial ring and $W \subseteq T$ is a multiplicative set, Proposition 5.3 implies that for any ideal $A \subseteq T$, $I_e(W^{-1}A) = W^{-1}I_e(A)$. Similarly, if \mathfrak{m} is the irrelevant ideal of T and \hat{T} denotes the completion of T with respect to \mathfrak{m} , then $I_e(A\hat{T}) = I_e(A)\hat{T}$.

Definition 5.4. Fix any $u \in T$. For any ideal $A \subseteq T$ we define a sequence of ideals as follows: $A_0 = A$ and $A_{i+1} = I_e(uA_i) + A_i$ for all $i \geq 0$. Clearly this sequence is an ascending chain and as T is Noetherian it stabilizes to some ideal which we denote with $A^{?^e u}$.

Proposition 5.5. Fix any $u \in T$ and let $A \subseteq T$ be an ideal. If $B \subseteq T$ is an ideal containing A and if $uB \subseteq B^{[e]}$ then $A^{?^e u} \subseteq B$.

Proof. Let $fA_i g_{i=0}^1$ be the sequence of ideals as in Definition 5.4. We show by induction that $A_i \subseteq B$ for all $i \geq 0$. Since $A_0 = A$ and $A \subseteq B$ the claim is true for $i = 0$; assume that $i \geq 0$ and that $A_i \subseteq B$. Now $uA_i \subseteq uB \subseteq B^{[e]}$ and the minimality of $I_e(uA_i) + A_i$ now implies that $A_{i+1} = I_e(uA_i) + A_i \subseteq B$.

The regular local ring R at the focus of this paper is a power series hence it is of the form $K[[x_1; \dots; x_n]]$ for some field K of prime characteristic p . When A is expanded from the polynomial ring $T = K[x_1; \dots; x_n]$ and $u \in T$ we want to compute $A^{?^e u}$ by performing calculations in T rather than R . Proposition 5.6 below shows how to do that.

Proposition 5.6. Let A be an ideal of $T = K[x_1; \dots; x_n]$ and let $u \in T$. We have $(AR)^{?^e u} = A^{?^e u}R$.

Proof. Let $fB_i g_{i=0}$ and $fC_i g_{i=0}$ be the sequences introduced in Definition 5.4 whose stable values are $(AR)^{?^e u}$ and $A^{?^e u}$, respectively. We will show that $B_i = C_i R$ for all $i \geq 0$ using induction on i .

First, $C_0 R = AR = B_0$, so assume that $i > 0$ and that $B_{i-1} = C_{i-1} R$. Now notice that since $uC_{i-1} \subseteq C_i^{[e]}$ and $C_{i-1} \subseteq C_i$ we have $uB_{i-1} = uC_{i-1} R \subseteq C_i^{[e]} R = (C_i R)^{[e]}$ and $B_{i-1} = C_{i-1} R \subseteq C_i R$ so the minimality of B_i implies that $B_i \subseteq C_i R$. On the other hand, $uC_{i-1} R = uB_{i-1} \subseteq B_i^{[e]}$ implies that $uC_{i-1} = uC_{i-1} R \setminus T \subseteq B_i^{[e]} \setminus T = (B_i \setminus T)^{[e]}$ and $C_{i-1} R = B_{i-1} \subseteq B_i$ implies that $C_{i-1} = C_{i-1} R \setminus T \subseteq B_i \setminus T$ and the minimality of C_i implies that $C_i \subseteq B_i \setminus T$ hence $C_i R \subseteq (B_i \setminus T) R \subseteq B_i$.

6. E_S -ideals and special E_S -ideals

Following [S], we call an ideal $K \subseteq S[[T; f]]$ a graded two-sided ideal if $K = \sum_{i=0}^M K_i T^i$ for ideals K_0, K_1, \dots of S . An important example is $K = LS[[T; f]]$ for some ideal $L \subseteq S$. Let G be an $S[[T; f]]$ -module. An $S[[T; f]]$ -submodule $M \subseteq G$ is a graded annihilator submodule if $M = \text{ann}_G K$ for some graded two-sided ideal $K \subseteq S[[T; f]]$.

For any $S[[T; f]]$ -submodule $M \subseteq G$ we define the graded-annihilator of M , denoted $\text{gr-ann}_{S[[T; f]]} M$, to be the largest graded two-sided ideal contained in $\text{ann}_{S[[T; f]]} M$.

We call an ideal $L \subseteq S$ a G -special ideal whenever $LS[[T; f]]$ is the graded annihilator of some $S[[T; f]]$ -submodule $M \subseteq G$, in which case $LS[[T; f]] = \text{gr-ann}(\text{ann}_G LS[[T; f]])$ (cf. Lemma 1.7 in [S]; notice that we extended slightly the definition of special ideals to the case where G is not necessarily T -torsion-free.)

Proposition 6.1. Assume that R is complete and that E_S is T -torsion free. An ideal $L \subseteq R$ which contains I is an E_S -ideal if and only if LS is E -special.

Proof. Assume first that L is E_S -special, i.e., $LS[\mathbb{T};f] = \text{gr-ann } N$ for some $S[\mathbb{T};f]$ -submodule N of E_S , and since E_S is assumed to be T -torsion-free, we have $\text{gr-ann } N = (0 \neq N)S[\mathbb{T};f]$ (cf. Definition 1.10 in §). We can also write $N = \text{ann}_{E_S} L^0$ for some E_S -ideal L^0 and

$$\begin{aligned} LS[\mathbb{T};f] &= (0 \neq \text{ann}_{E_S} L^0)S[\mathbb{T};f] \\ &= (0 \neq (R=L^0)^-)S[\mathbb{T};f] \\ &= (0 \neq R=L^0)S[\mathbb{T};f] \\ &= L^0S[\mathbb{T};f] \end{aligned}$$

so $L = L^0$ and is an E_S -ideal.

If, on the other hand, L is an E_S -ideal, i.e., if $\text{ann}_{E_S} L = \text{ann}_{E_S} LS[\mathbb{T};f]$, then

$$\begin{aligned} \text{gr-ann ann}_{E_S} LS[\mathbb{T};f] &= (0 \neq \text{ann}_{E_S} LS[\mathbb{T};f])S[\mathbb{T};f] \\ &= (0 \neq \text{ann}_{E_S} LS)S[\mathbb{T};f] \\ &= (0 \neq (R=L)^-)S[\mathbb{T};f] \\ &= (0 \neq R=L)S[\mathbb{T};f] \\ &= LS[\mathbb{T};f] \end{aligned}$$

and so L is E_S -special.

7. The $S[\mathbb{T};f]$ -module structure of $H_{mS}^{\dim S}(S)$ and the induced structure on E_S

In what follows we describe a natural $S[\mathbb{T};f]$ -module structure on $H_{mS}^{\dim S}(S)$ and show how this induces an $S[\mathbb{T};f]$ -module structure on E_S ; the following section will describe its relevance to test-ideals.

We shall assume henceforth that S is Cohen-Macaulay with canonical module ω_S .

The short exact sequence $0 \rightarrow \omega_S \rightarrow S \rightarrow S/\omega_S \rightarrow 0$ yields a surjection $H_{mS}^{\dim S}(\omega_S) \rightarrow H_{mS}^{\dim S}(S)$; for any system of parameters x_1, \dots, x_d for S this map can also be described as the map

$$\lim_{i \rightarrow 0} \frac{\omega_S^i}{x_1^i \omega_S + \frac{1}{d}! x} \rightarrow \lim_{i \rightarrow 0} \frac{S^i}{x_1^i S + \frac{1}{d}! Sx}$$

induced by the maps

$$\frac{\omega_S^i}{x_1^i \omega_S + \frac{1}{d}! x} \rightarrow \frac{S^i}{x_1^i S + \frac{1}{d}! Sx}$$

given by $a + \frac{1}{d}! x \mapsto a + x_1^i S + \frac{1}{d}! Sx$. The natural action of Frobenius on $H_{mS}^{\dim S}(S)$ given by

$$f(a + x_1^i S + \frac{1}{d}! Sx) = a^p + x_1^{ip} S + \frac{1}{d}! x^p S$$

now lifts to an action on $E_S = \lim_{i \rightarrow 0} \frac{\omega_S^i}{x_1^i \omega_S + \frac{1}{d}! x}$ given by

$$f(a + x_1^i \omega_S + \frac{1}{d}! x) = a^p + x_1^{ip} \omega_S + \frac{1}{d}! x^p S$$

and this $S[[T;f]]$ module structure on E_S clearly makes the surjection $E_S \rightarrow H_m^{\dim S}(S)$ described above into a $S[[T;f]]$ -linear map.

If we apply ϕ to the $S[[T;f]]$ -linear surjection $E_S \rightarrow H_m^{\dim S}(S)$ and identify $E_{\bar{S}}$ with R/I we obtain the following commutative diagram with exact rows

$$(5) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_m^{\dim S}(S) & \longrightarrow & \frac{R}{I} & & \\ & & \downarrow u & & \downarrow u & & \\ 0 & \longrightarrow & F_R \oplus H_m^{\dim S}(S) & \longrightarrow & \frac{R}{I[[T]]} & & \end{array}$$

where $u \in R$, the second vertical map is multiplication by u and the first vertical map is given by restriction of the second, i.e., also by multiplication by u . We deduce that under the identification of $E_{\bar{S}}$ with R/I , $H_m^{\dim S}(S)$ is identified with J/I for some ideal $J \subset R$ containing I . This ideal J must then satisfy $uJ \subset J[[T]]$.

Our next step is to compute J and u effectively. Let ϕ be the full pre-image of ϕ in R . Working over R , the surjection $E_S = H_m^{\dim S}(\phi) \rightarrow H_m^{\dim S}(S)$ can be written as $E_S = H_m^{\dim S}(\phi = I) \rightarrow H_m^{\dim S}(R = I)$. Write $d = \dim R - \dim S$; recall that Local Duality states that the functors $H_m^{\dim S}(\phi)$ and $\text{Ext}_R(\phi; R)$ are isomorphic and the surjection above is induced by applying either of these functors to the inclusion $\phi \subset S$. Applying the latter and a further application of $(\phi)^\perp$ yields the injection $\text{Ext}_R(R = I; R) \hookrightarrow \text{Ext}_R(\phi = I; R)$ and so $J = I = \text{Ext}_R(R = I; R)$. This Ext-module can be computed effectively and J can be recovered by computing a minimal presentation of this module.

To find the map u in (5) we use the fact that $F = H_m^{\dim S}(S)$ is the R -algebra with one generator corresponding to the $S[[T;f]]$ -module structure defined above (cf. Example 3.7 in [LS]). Hence the S -linear maps $\text{Ext}_R(R = I; R) \rightarrow F_R \oplus \text{Ext}_R(R = I; R)$ form a rank-one S -free module and the generator u of this free module can be computed explicitly from the generator of

$$\frac{I[[T]] \otimes_R I \setminus J[[T]] \otimes_R J}{I[[T]]}$$

(cf. Chapter 3 of [B] and section 4.)

8. The computation of parameter-test-ideals

Throughout this section we will assume that $S = R/I$ is Cohen-Macaulay with canonical module ω_S . We shall write $H = H_m^{\dim S}(S)$ and we will assume that E_S is T -torsion-free. This last assumption implies that H , being a quotient of E_S by a graded annihilator submodule, is also T -torsion-free (cf. Lemma 3.1 in [S]). Recall now that Corollary 4.6 in [S] now states that the parameter-test-element of S is the smallest H -special ideal of S of positive height. In this section we relate the H -special ideals to E_S -special ideals and describe an algorithm for computing the parameter-test-element of S .

First we note that Proposition 6.1 and yields the following.

Theorem 8.1. Assume that E_S is T -torsion-free and write $H := H_{mS}^{\dim S}(S) = \frac{E_S}{\text{ann}_{E_S} J}$ where $J \subseteq R$ is an E_S -ideal. The H -special ideals are

$$f(L : J) \subseteq L \subseteq R \text{ is an } E_S\text{-special ideal contained in } Jg :$$

Proof. This follows from Proposition 3.3 in [5].

We are now ready for the main theorem in this section.

Theorem 8.2. Assume that E_S is T -torsion-free. Let $c \in R$ be such that its image in S is a parameter test element. The parameter test ideal τ of S is given by $((cJ + I)^{?u} :_R J)S$.

Proof. Notice that $(cJ + I)^{?u}$ is an E_S -ideal and that since $c \in ((cJ + I)^{?u} : J)$, we have $\text{ht}((cJ + I)^{?u} : J)S > 0$. Now

$$\begin{aligned} \tau &= \bigcap \{ f(K \subseteq J) \subseteq S \text{ is an } H\text{-special ideal; } \text{ht}(K) > 0 \} \\ &= \bigcap \{ f(L \subseteq J) \subseteq L \subseteq J \text{ is an } E_S\text{-ideal; } \text{ht}(L \subseteq J)S > 0 \} \\ &= \bigcap \{ f(L \subseteq J) \subseteq J \text{ is an } E_S\text{-ideal; } \text{ht}(L : J)S > 0 \} : J \end{aligned}$$

so we see that $\tau = ((cJ + I)^{?u} : J)$.

Also, $c \in \tau$ hence $cJ \subseteq L$ for all E_S -ideals L for which $\text{ht}(L : J)S > 0$ and Proposition 5.5 implies that $(cJ + I)^{?u} \subseteq L$ and hence that $((cJ + I)^{?u} : J) \subseteq (L : J)$ for all E_S -ideals L for which $\text{ht}(L : J) > 0$. We conclude that $((cJ + I)^{?u} : J) = \tau$.

In the case where E_S is T -torsion-free, if we are given one parameter test element, we can now compute the entire parameter test ideal of S as follows.

- (1) Find the element $u \in R$ as described in section 7 and use Theorem 4.6 to determine whether E_S is T -torsion-free. If E_S is T -torsion-free proceed as follows.
- (2) Find the ideal $I \subseteq J \subseteq R$ as described in section 7.
- (3) Given one parameter test element c , compute $L = (cJ + I)^{?u}$ as described in section 5.
- (4) The parameter test ideal of S is $(L \subseteq J)S$.

9. Applications and examples

A particularly simple instance of the results of the previous chapters is the case where S is a complete intersection, i.e., the case where I is generated by a regular sequence $u_1, \dots, u_s \in R$. Now $S = R/I$ is Gorenstein, $E_S = H_{mS}^{\dim S}(S)$ (so the surjection described in the beginning of section 7 is an equality) and $(E_S) = R/I \rightarrow R/I^{[p]}$ is given by multiplication by $u = (u_1 \dots u_s)^{p-1}$ whose image in $(I^{[p]} \subseteq I) = I^{[p]}$ generates it.

We first return to Theorem 4.6 and put it in its proper context and recover Richard Fedder's characterization of F -injective complete intersections (Proposition 2.1 in [F]). Assume now that $I \subseteq R$ is generated by a regular sequence u_1, \dots, u_s and notice that as $S = R/I$ is Gorenstein, $E_S = H_{mS}^{\dim(S)}(S)$. We call S F -injective if the natural Frobenius map on $E_S = H_{mS}^{\dim(S)}(S)$ is injective, i.e., if $\text{Nil}(E_S) = 0$. We now recover Fedder's Criterion which states that S is F -injective if and only if $u \notin m^{[p]}$ where $u = (u_1 \dots u_s)^{p-1}$. The crucial fact here is that (E_S) is the map $R/I \rightarrow R/I^{[p]}$. As in the proof of Theorem 4.6

- [HH1] M. Hochster and C. Huneke. Tight closure, invariant theory, and the Briançon-Skoda theorem. *Journal of the AMS* 3 (1990), no. 1, 31{116.
- [HH2] M. Hochster and C. Huneke. F -regularity, test elements, and smooth base change. *Transactions of the AMS*, 346 (1994), no. 1, pp. 1{62.
- [K] E. Kunz. Characterizations of regular local rings for characteristic p . *American Journal of Mathematics* 91 (1969), pp. 772{784.
- [KS] M. Katzman and R. Y. Sharp. Uniform behaviour of the Frobenius closures of ideals generated by regular sequences, *Journal of Algebra*, 295 (2006) 231{246.
- [L] G. Lyubeznik. F -modules: applications to local cohomology and D -modules in characteristic $p > 0$. *J. Reine Angew. Math.* 491 (1997), pp. 65{130.
- [LS] G. Lyubeznik and K. E. Smith. On the commutation of the test ideal with localization and completion. *Transactions of the AMS* 353 (2001), no. 8, pp. 3149{3180.
- [S] R. Y. Sharp. Graded annihilators of modules over the Frobenius skew polynomial ring, and tight closure. [arXiv.org/math/0605329](https://arxiv.org/math/0605329).
- [SV] D. W. Sharpe and P. Vámos. *Injective modules*. Cambridge Tracts in Mathematics and Mathematical Physics 62. Cambridge University Press, London-New York, 1972.
- [Sm1] K. E. Smith. Test ideals in local rings. *Transactions of the AMS* 347 (1995), no. 9, pp. 3453{3472.
- [Sm2] K. E. Smith. F -rational rings have rational singularities. *American Journal of Mathematics* 119 (1997), no. 1, pp. 159{180.

Department of Pure Mathematics, University of Sheffield, Hicks Building, Sheffield S3 7RH, United Kingdom, Fax number: +44-(0)114-222-3769

E-mail address: M.Katzman@sheffield.ac.uk