

# ON THE SHARPNESS OF FEW NOMIAL BOUND AND THE NUMBER OF COMPONENTS OF A FEW NOMIAL HYPERSURFACE

FREDERIC BIHAN, J. MAURICE ROJAS, AND FRANK SOTTILE

**Abstract.** We show the existence of systems of  $n$  polynomial equations in  $n$  variables, with a total of  $n + k + 1$  distinct monomial terms, possessing  $\frac{n+k}{k} 2^{\binom{n+k}{4}}$  nondegenerate positive solutions. This shows that the recent upper bound of  $\frac{e^2+3}{4} 2^{\binom{n+k}{4}} n^k$  for the number of nondegenerate positive solutions is asymptotically sharp for fixed  $k$  and large  $n$ . We also adapt a method of Perrucci to show that there are fewer than  $\frac{e^2+3}{4} 2^{\binom{n+k}{4}} 2^{n^k}$  connected components in a smooth hypersurface in the positive orthant of  $\mathbb{R}^n$  defined by a polynomial with  $n+k+1$  monomials. Our results hold for polynomials with real exponents.

Khovanskii [5] showed there are at most  $2^{\binom{n+k}{2}} (n+1)^{n+k}$  nondegenerate positive solutions to a system of  $n$  polynomial equations in  $n$  variables having a total of  $n+k+1$  distinct monomials. This [fewnomial bound](#) is also a bound for polynomials with real exponents, which are linear combinations of monomials with real-number exponents. The equivalence is discussed in [8] and the arguments of [3] require that we allow real exponents.

While this fewnomial bound was not believed to be sharp, only recently have smaller bounds been found. The first breakthrough was due to Li, Rojas, and Wang [7] who showed that a system of two trinomials in two variables has at most 5 positive solutions, which is significantly less than Khovanskii's bound of 5184. Bihan [2] showed that a system of  $n$  polynomials in  $n$  variables with  $n+2$  monomials has at most  $n+1$  nondegenerate positive solutions and proved the existence of such a system with  $n+1$  positive solutions. This gave the sharp bound of  $n+1$  for the case  $k=1$ . Bihan and Sottile [3] generalized this to all  $k$ , giving the upper bound of  $\frac{e^2+3}{4} 2^{\binom{n+k}{4}} n^k$  for the number of nondegenerate positive solutions, which was significantly smaller than Khovanskii's bound. We show that this upper bound is near-optimal for fixed  $k$  and large  $n$ .

**Theorem 1.** For any positive integers  $n, k$  with  $n > k$ , there exists a system of  $n$  polynomials in  $n$  variables involving  $n+k+1$  distinct monomials and having  $\frac{n+k}{k} 2^{\binom{n+k}{4}}$  nondegenerate positive solutions.

We believe that there is room for improvement in the dependence on  $k$ , both in the upper bound of [3] and in the lower bound of Theorem 1.

**Proof.** We will construct such a system when  $n = km$ , a multiple of  $k$ , from which we may deduce the general case as follows. Suppose that  $n = km + j$  with  $1 \leq j < k$  and we have a system of  $m$   $k$  equations in  $m$   $k$  variables involving  $m$   $k + k + 1$  monomials with  $(m+1)^k$  nondegenerate positive solutions. We add  $j$  new variables  $x_1, \dots, x_j$  and  $j$  new equations  $x_1 = 1, \dots, x_j = 1$ . Since the polynomials in the original system may be assumed to have

2000 Mathematics Subject Classification. 14P99.

Key words and phrases. Fewnomials, connected component.

Rojas supported by NSF CAREER grant DMS-0349309.

Sottile supported by NSF CAREER grant DMS-0538734.

constant term  $s$ , this gives a system with  $n$  polynomials in  $n$  variables having  $n + k + 1$  monomials and  $(m + 1)^k = \frac{n+k}{k}^k$  nondegenerate positive solutions.

Fix positive integers  $k, m$  and set  $n = km$ . Bihan [2] showed there exists a system of  $m$  polynomials in  $m$  variables

$$f_1(y_1; \dots; y_m) = f_2(y_1; \dots; y_m) = \dots = f_m(y_1; \dots; y_m) = 0$$

having  $m + 1$  solutions, and where each of polynomial has the same  $m + 2$  monomials, one of which we may take to be a constant.

For each  $j = 1; \dots; k$ , let  $y_{j1}; \dots; y_{jm}$  be  $m$  new variables and consider the system

$$f_1(y_{j1}; \dots; y_{jm}) = \dots = f_m(y_{j1}; \dots; y_{jm}) = 0;$$

which has  $m + 1$  positive solutions in  $(y_{j1}; \dots; y_{jm})$ . As the sets of variables are disjoint, the combined system consisting of all  $km$  polynomials in all  $km$  variables has  $(m + 1)^k$  positive solutions. Each subsystem has  $m + 2$  monomials, one of which is a constant. Thus the combined system has  $1 + k(m + 1) = km + k + 1 = n + k + 1$  monomials.

Remark 2. This proof of Theorem 1 is based on Bihan's non-constructive proof of the existence of a system of  $m$  polynomials in  $m$  variables having  $m + 2$  monomials and  $m + 1$  nondegenerate positive solutions. While finding explicit such systems is challenging in general, we give systems for  $n = 2$  and  $n = 3$ .

The system of  $n = 2$  equations with 2 variables

$$x^2y(1 + 4x^2) = xy(4 + x^2) = 0;$$

has  $4 = 2 + 1 + 1$  monomials and exactly 3 complex solutions, each of which are nondegenerate and lie in the positive quadrant. We give numerical approximations, computed with the computer algebra system SINGULAR [4, 10].

$$(2.618034; 4.1459); (1; 5); (0.381966; 10.854102) :$$

The system of  $n = 3$  equations with 3 variables

$$yz(1 + 8 + 2x^2) = xy(1 + 220 + x^2) = z(1 + x^2) = 0;$$

has  $5 = 3 + 1 + 1$  monomials and exactly 4 complex solutions, each of which are nondegenerate and lie in the positive octant.

$$(0.0766453; 0.13595; 1.005875); (0.0845135; 0.138297; 1.007143); \\ (0.540459; 0.548869; 1.2921); (1.298382; 1.301883; 2.6858) :$$

Khovanskii also considered smooth hypersurfaces in the positive orthant  $\mathbb{R}_{>}^n$  defined by polynomials with  $n + k + 1$  monomials. He showed that the total Betti number of such a fewnomial hypersurface is at most  $(2n^2 - n + 1)^{n+k} (2n)^{n-1} 2^{\binom{n+k}{2}}$  [6, x3.14, Cor. 4]. Li, Rojas, and Wang [7] bounded the number of connected components of such a hypersurfaces by  $n(n+1)^{n+k+1} 2^{n-1} 2^{\binom{n+k+1}{2}}$ . Perrucci [9] lowered this bound to  $(n+1)^{n+k} 2^{1+\binom{n+k}{2}}$ . His method was to bound the number of compact components and then use an argument based on the faces of the  $n$ -dimensional cube to bound the number of all components. We improve Perrucci's method, using the  $n$ -simplex and the bounds of Bihan and Sottile [3] to obtain a new, lower bound. This also extends an earlier bound of Benedetti, Loeser, and Risler to the context of sparse polynomial systems [1].

Theorem 3. A smooth hypersurface in  $\mathbb{R}_{>}^n$  defined by a polynomial with  $n + k + 1$  monomials having a line span  $\mathbb{R}^n$  has fewer than  $\frac{e^{2+3}}{4} \binom{k}{2} 2^n n^k$  connected components.

Remark 4. If a hypersurface  $f = 0$  is singular, we may bound its number of connected components by the number of connected components of the two hypersurfaces  $f = \pm \epsilon$ , for  $\epsilon$  small. (This is Lemma 13 in [7].) Since we can assume that  $f$  has a constant term, this gives the bound of  $2 \cdot \frac{e^2+3}{4} \binom{k}{e} 2^{n+k} 2^{n+k+1}$  for the number of connected components of an arbitrary (possibly singular) hypersurface in  $\mathbb{R}_{>}^n$  defined by a polynomial with  $n+k+1$  monomials.

Let  $(n; k)$  be the maximum number of compact connected components of a smooth hypersurface in  $\mathbb{R}_{>}^n$  defined by a polynomial having at most  $n+k+1$  monomials and let  $(n; k)$  be the maximal number of connected components of such a hypersurface.

Lemma 5. 
$$(n; k) \leq \sum_{i=0}^{n-1} \binom{n+i}{i} (n-i; k).$$

Proof of Theorem 3. Bihan and Sottile [3] proved that  $(n; k) \leq \frac{e^2+3}{8} 2^{\binom{k}{e}} n^k$ . Substituting this into Lemma 5 bounds  $(n; k)$  by

$$\sum_{i=0}^{n-1} \binom{n+i}{i} \left( \frac{e^2+3}{8} 2^{\binom{k}{e}} n^k \right)^{n-i} < \left( \frac{e^2+3}{8} 2^{\binom{k}{e}} n^k \right)^{\sum_{i=0}^{n-1} (n-i)} = \left( \frac{e^2+3}{8} 2^{\binom{k}{e}} n^k \right)^{n^2}.$$

Proof of Lemma 5. Let  $f$  be a polynomial in the variables  $x_1, \dots, x_n$  which has  $n+k+1$  distinct monomials and suppose that  $f(x) = 0$  defines a smooth hypersurface  $X$  in  $\mathbb{R}_{>}^n$ . We may apply a monomial change of coordinates to  $\mathbb{R}_{>}^n$  and assume that  $1, x_1, x_2, \dots, x_n$  are among the monomials appearing in  $f$ . Perturbing the exponents of  $f$  if necessary, we may assume that  $f$  also has a monomial  $x^a = x_1^{a_1} x_2^{a_2} \dots x_n^{a_n}$  with  $a_1 a_2 \dots a_n \neq 0$ . (Justifications for these assumptions are given in §1.) Applying the coordinate change  $x_i \mapsto x_i^{-1}$  whenever  $a_i < 0$ , we may further assume that the monomials of  $f$  include

$$1, x_i^{e_i}; \quad i = 1, \dots, n; \quad \text{and} \quad x^a;$$

where  $e_i \geq 1$  and  $a_i > 0$ , for  $i = 1, \dots, n$ .

Suppose that  $a = (a_1, \dots, a_n) \in \mathbb{R}_{>}^{1+n}$  with  $a_1 \dots a_n \neq 1$ . Define hypersurfaces  $H_0, H_1, \dots, H_n$  of  $\mathbb{R}_{>}^n$  by

$$H_0 = \{x \in \mathbb{R}_{>}^n \mid x^a = 1\} \quad \text{and} \quad H_i = \{x \in \mathbb{R}_{>}^n \mid x_i = 1\} \quad i = 1, \dots, n.$$

The transformation  $\text{Log}: \mathbb{R}_{>}^n \rightarrow \mathbb{R}^n$  defined on each coordinate by  $x_i \mapsto \log(x_i)$  sends the hypersurfaces  $H_0, H_1, \dots, H_n$  to hyperplanes in general position. That is, if  $S \subseteq \{0, 1, \dots, n\}$  and we define  $H_S = \bigcap_{i \in S} H_i$ , then  $\text{Log}(H_S)$  is an affine linear space of dimension  $n - |S|$ . Moreover, the complement of the hypersurfaces  $H_i$  has  $2^{n+1} - 1$  connected components, exactly one of which is bounded.

If we restrict  $f$  to some  $H_S$ , we will obtain a new polynomial  $f_S$  in  $n - |S|$  variables with at most  $1 + n - |S| + k$  monomials. (While this is obvious when  $0 \notin S$ , if  $0 \in S$ , then we pick an index  $j \notin S$  and use  $x^a = 1$  to eliminate  $x_j$ .) For almost all  $S$ , the polynomial  $f_S$  defines a smooth hypersurface  $X_S$  of  $H_S$ .

We may choose  $\epsilon$  small enough so that every compact connected component of  $X$  lies in the bounded region of the complement of the hypersurfaces  $H_i$ , and every noncompact connected component of  $X$  meets some hypersurface  $H_i$ . Shrinking  $\epsilon$  if necessary, we can insure that every bounded component of  $X_S$  lies in the bounded region of the complement of  $H_j \setminus H_S$  for  $j \notin S$ , and every unbounded component meets some  $H_j \setminus H_S$  for  $j \notin S$ .

Given a connected component  $C$  of  $X$ , the subsets  $S = \{f_0, \dots, f_n\}$  such that  $C$  meets  $H_S$  form a simplicial complex. If  $S$  represents a maximal simplex in this complex, then  $C \setminus H_S$  is a union of compact components of  $X_S$ , and  $|S| < n$  as  $H_S$  is not a point. Thus the number of connected components of  $X$  is bounded by the sum of the numbers of compact components of  $X_S$  for all  $S = \{f_0, \dots, f_n\}$  with  $n > |S|$ . Since each  $f_S$  has at most  $1 + n - |S|$  monomials, this sum is bounded by the sum in the statement of the lemma.

## References

- [1] R. Benedetti, F. Loeser, and J.-J. Risler, Bounding the number of connected components of a real algebraic set, *Discrete Comput. Geom.* 6 (1991), no. 3, 191{209.
- [2] F. Bihan, Polynomial systems supported on circuits and dessins d'enfants, 2005, *Journal of the London Mathematical Society*, to appear.
- [3] F. Bihan and F. Sottile, New fewnomial upper bounds from Gale dual polynomial systems, 2006, *Moscow Mathematical Journal*, to appear. [math.AG/0609544](http://math.AG/0609544).
- [4] G.-M. Greuel, G. Pfister, and H. Schönemann. Singular 3.0.2. A Computer Algebra System for Polynomial Computations. Centre for Computer Algebra, University of Kaiserslautern (2006). <http://www.singular.uni-kl.de>.
- [5] A. G. Khovanskii, A class of systems of transcendental equations, *Dokl. Akad. Nauk. SSSR* 255 (1980), no. 4, 804{807.
- [6] ———, Fewnomials, *Trans. of Math. Monographs*, 88, AMS, 1991.
- [7] T.-Y. Li, J. M. Rojas, and X. Wang, Counting real connected components of trinomial curve intersections and monomial hypersurfaces, *Discrete Comput. Geom.* 30 (2003), no. 3, 379{414.
- [8] D. Napolitano, A power function approach to Kouchnirenko's conjecture, *Symbolic computation: solving equations in algebra, geometry, and engineering* (South Hadley, MA, 2000), *Contemp. Math.*, vol. 286, Amer. Math. Soc., Providence, RI, 2001, pp. 99{106.
- [9] D. Perrucci, Some bounds for the number of components of real zero sets of sparse polynomials, *Discrete Comput. Geom.* 34 (2005), no. 3, 475{495.
- [10] M. Wenk and W. Pohl. normal.lib. A Singular 3.0.2 library for Complex Solving of Polynomial Systems (2006).

Laboratoire de Mathématiques, Université de Savoie, 73376 Le Bourget-du-Lac Cedex, France

E-mail address: Frederic.Bihan@univ-savoie.fr

URL: <http://www.lama.univ-savoie.fr/~bihan/>

Department of Mathematics, Texas A & M University, College Station, Texas 77843, USA

E-mail address: rojas@math.tamu.edu

URL: <http://www.math.tamu.edu/~rojas/>

Department of Mathematics, Texas A & M University, College Station, Texas 77843, USA

E-mail address: sottile@math.tamu.edu

URL: <http://www.math.tamu.edu/~sottile/>