

Stability of closed characteristics on compact convex hypersurfaces in \mathbf{R}^6

Wei Wang*

School of Mathematical Science
Peking University, Beijing 100871
PEOPLES REPUBLIC OF CHINA

Abstract

In this paper, let $\Sigma \subset \mathbf{R}^6$ be a compact convex hypersurface. We prove that if Σ carries only finitely many geometrically distinct closed characteristics, then at least two of them must possess irrational mean indices. Moreover, if Σ carries exactly three geometrically distinct closed characteristics, then at least two of them must be elliptic.

Key words: Compact convex hypersurfaces, closed characteristics, Hamiltonian systems, Morse theory, mean index identity, stability.

AMS Subject Classification: 58E05, 37J45, 37C75.

Running title: Stability of closed characteristics

1 Introduction and main results

In this paper, let Σ be a fixed C^3 compact convex hypersurface in \mathbf{R}^{2n} , i.e., Σ is the boundary of a compact and strictly convex region U in \mathbf{R}^{2n} . We denote the set of all such hypersurfaces by $\mathcal{H}(2n)$. Without loss of generality, we suppose U contains the origin. We consider closed characteristics (τ, y) on Σ , which are solutions of the following problem

$$\begin{cases} \dot{y} = JN_{\Sigma}(y), \\ y(\tau) = y(0), \end{cases} \quad (1.1)$$

where $J = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}$, I_n is the identity matrix in \mathbf{R}^n , $\tau > 0$, $N_{\Sigma}(y)$ is the outward normal vector of Σ at y normalized by the condition $N_{\Sigma}(y) \cdot y = 1$. Here $a \cdot b$ denotes the standard inner product

*Partially supported by NNSF, RFDP of MOE of China. E-mail: alexanderweiwang@yahoo.com.cn

of $a, b \in \mathbf{R}^{2n}$. A closed characteristic (τ, y) is *prime*, if τ is the minimal period of y . Two closed characteristics (τ, y) and (σ, z) are *geometrically distinct*, if $y(\mathbf{R}) \neq z(\mathbf{R})$. We denote by $\mathcal{J}(\Sigma)$ and $\tilde{\mathcal{J}}(\Sigma)$ the set of all closed characteristics (τ, y) on Σ with τ being the minimal period of y and the set of all geometrically distinct ones respectively. Note that $\mathcal{J}(\Sigma) = \{\theta \cdot y \mid \theta \in S^1, y \text{ is prime}\}$, while $\tilde{\mathcal{J}}(\Sigma) = \mathcal{J}(\Sigma)/S^1$, where the natural S^1 -action is defined by $\theta \cdot y(t) = y(t + \tau\theta)$, $\forall \theta \in S^1, t \in \mathbf{R}$.

Let $j : \mathbf{R}^{2n} \rightarrow \mathbf{R}$ be the gauge function of Σ , i.e., $j(\lambda x) = \lambda$ for $x \in \Sigma$ and $\lambda \geq 0$, then $j \in C^3(\mathbf{R}^{2n} \setminus \{0\}, \mathbf{R}) \cap C^0(\mathbf{R}^{2n}, \mathbf{R})$ and $\Sigma = j^{-1}(1)$. Fix a constant $\alpha \in (1, 2)$ and define the Hamiltonian function $H_\alpha : \mathbf{R}^{2n} \rightarrow [0, +\infty)$ by

$$H_\alpha(x) = j(x)^\alpha, \quad \forall x \in \mathbf{R}^{2n}. \quad (1.2)$$

Then $H_\alpha \in C^3(\mathbf{R}^{2n} \setminus \{0\}, \mathbf{R}) \cap C^1(\mathbf{R}^{2n}, \mathbf{R})$ is convex and $\Sigma = H_\alpha^{-1}(1)$. It is well known that the problem (1.1) is equivalent to the following given energy problem of the Hamiltonian system

$$\begin{cases} \dot{y}(t) = JH'_\alpha(y(t)), & H_\alpha(y(t)) = 1, & \forall t \in \mathbf{R}. \\ y(\tau) = y(0). \end{cases} \quad (1.3)$$

Denote by $\mathcal{J}(\Sigma, \alpha)$ the set of all solutions (τ, y) of (1.3) where τ is the minimal period of y and by $\tilde{\mathcal{J}}(\Sigma, \alpha)$ the set of all geometrically distinct solutions of (1.3). As above, $\tilde{\mathcal{J}}(\Sigma, \alpha)$ is obtained from $\mathcal{J}(\Sigma, \alpha)$ by dividing the natural S^1 -action. Note that elements in $\mathcal{J}(\Sigma)$ and $\mathcal{J}(\Sigma, \alpha)$ are one to one correspondent to each other, similarly for $\tilde{\mathcal{J}}(\Sigma)$ and $\tilde{\mathcal{J}}(\Sigma, \alpha)$.

Let $(\tau, y) \in \mathcal{J}(\Sigma, \alpha)$. The fundamental solution $\gamma_y : [0, \tau] \rightarrow \text{Sp}(2n)$ with $\gamma_y(0) = I_{2n}$ of the linearized Hamiltonian system

$$\dot{w}(t) = JH''_\alpha(y(t))w(t), \quad \forall t \in \mathbf{R}, \quad (1.4)$$

is called the *associate symplectic path* of (τ, y) . The eigenvalues of $\gamma_y(\tau)$ are called *Floquet multipliers* of (τ, y) . By Proposition 1.6.13 of [Eke3], the Floquet multipliers with their multiplicities of $(\tau, y) \in \mathcal{J}(\Sigma)$ do not depend on the particular choice of the Hamiltonian function in (1.3). For any $M \in \text{Sp}(2n)$, we define the *elliptic height* $e(M)$ of M to be the total algebraic multiplicity of all eigenvalues of M on the unit circle $\mathbf{U} = \{z \in \mathbf{C} \mid |z| = 1\}$ in the complex plane \mathbf{C} . Since M is symplectic, $e(M)$ is even and $0 \leq e(M) \leq 2n$. As usual a $(\tau, y) \in \mathcal{J}(\Sigma)$ is *elliptic*, if $e(\gamma_y(\tau)) = 2n$. It is *non-degenerate*, if 1 is a double Floquet multiplier of it. It is *hyperbolic*, if 1 is a double Floquet multiplier of it and $e(\gamma_y(\tau)) = 2$. It is well known that these concepts are independent of the choice of $\alpha > 1$.

For the existence and multiplicity of geometrically distinct closed characteristics on convex compact hypersurfaces in \mathbf{R}^{2n} we refer to [Rab1], [Wei1], [EkL1], [EkH1], [Szu1], [HWZ1], [LoZ1],

[LLZ1], and references therein. Note that recently in [WHL], Wang, Hu and Long proved $\#\tilde{\mathcal{J}}(\Sigma) \geq 3$ for every $\Sigma \in \mathcal{H}(6)$.

On the stability problem, in [Eke2] of Ekeland in 1986 and [Lon2] of Long in 1998, for any $\Sigma \in \mathcal{H}(2n)$ the existence of at least one non-hyperbolic closed characteristic on Σ was proved provided $\#\tilde{\mathcal{J}}(\Sigma) < +\infty$. Ekeland proved also in [Eke2] the existence of at least one elliptic closed characteristic on Σ provided $\Sigma \in \mathcal{H}(2n)$ is $\sqrt{2}$ -pinched. In [DDE1] of 1992, Dell'Antonio, D'Onofrio and Ekeland proved the existence of at least one elliptic closed characteristic on Σ provided $\Sigma \in \mathcal{H}(2n)$ satisfies $\Sigma = -\Sigma$. In [Lon3] of 2000, Long proved that $\Sigma \in \mathcal{H}(4)$ and $\#\tilde{\mathcal{J}}(\Sigma) = 2$ imply that both of the closed characteristics must be elliptic. In [LoZ1] of 2002, Long and Zhu further proved when $\#\tilde{\mathcal{J}}(\Sigma) < +\infty$, there exists at least one elliptic closed characteristic and there are at least $\lfloor \frac{n}{2} \rfloor$ geometrically distinct closed characteristics on Σ possessing irrational mean indices, which are then non-hyperbolic. In the recent paper [LoW1], Long and Wang proved that there exist at least two non-hyperbolic closed characteristic on $\Sigma \in \mathcal{H}(6)$ when $\#\tilde{\mathcal{J}}(\Sigma) < +\infty$. Motivated by these results, we prove the following results in this paper:

Theorem 1.1. *On every $\Sigma \in \mathcal{H}(6)$ satisfying $\#\tilde{\mathcal{J}}(\Sigma) < +\infty$, there exist at least two geometrically distinct closed characteristics possessing irrational mean indices.*

Theorem 1.2. *Suppose $\#\tilde{\mathcal{J}}(\Sigma) = 3$ for some $\Sigma \in \mathcal{H}(6)$. Then there exist at least two elliptic closed characteristics in $\tilde{\mathcal{J}}(\Sigma)$.*

The proofs of Theorems 1.1 and 1.2 are given in Section 3. Mainly ingredients in the proofs include: the mean index identity for closed characteristics established in [WHL] recently, Morse inequality and the index iteration theory developed by Long and his coworkers, specially the common index jump theorem of Long and Zhu (Theorem 4.3 of [LoZ1], cf. Theorem 11.2.1 of [Lon4]). In Section 2, we review briefly the equivariant Morse theory and the mean index identity for closed characteristics on compact convex hypersurfaces in \mathbf{R}^{2n} developed in the recent [WHL].

In this paper, let \mathbf{N} , \mathbf{N}_0 , \mathbf{Z} , \mathbf{Q} , \mathbf{R} , and \mathbf{R}^+ denote the sets of natural integers, non-negative integers, integers, rational numbers, real numbers, and positive real numbers respectively. Denote by $a \cdot b$ and $|a|$ the standard inner product and norm in \mathbf{R}^{2n} . Denote by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ the standard L^2 -inner product and L^2 -norm. For an S^1 -space X , we denote by X_{S^1} the homotopy quotient of X module the S^1 -action, i.e., $X_{S^1} = S^\infty \times_{S^1} X$. We define the functions

$$\begin{cases} [a] = \max\{k \in \mathbf{Z} \mid k \leq a\}, & E(a) = \min\{k \in \mathbf{Z} \mid k \geq a\}, \\ \varphi(a) = E(a) - [a], \end{cases} \quad (1.5)$$

Specially, $\varphi(a) = 0$ if $a \in \mathbf{Z}$, and $\varphi(a) = 1$ if $a \notin \mathbf{Z}$. In this paper we use only \mathbf{Q} -coefficients for all homological modules. For a \mathbf{Z}_m -space pair (A, B) , let $H_*(A, B)^{\pm \mathbf{Z}_m} = \{\sigma \in H_*(A, B) \mid L_*\sigma = \pm \sigma\}$,

where L is a generator of the \mathbf{Z}_m -action.

2 Equivariant Morse theory for closed characteristics

In the rest of this paper, we fix a $\Sigma \in \mathcal{H}(2n)$ and assume the following condition on Σ :

(F) **There exist only finitely many geometrically distinct closed characteristics**

$\{(\tau_j, y_j)\}_{1 \leq j \leq k}$ **on** Σ .

In this section, we review briefly the equivariant Morse theory for closed characteristics on Σ developed in [WHL] which will be needed in Section 3 of this paper. All the details of proofs can be found in [WHL].

Let $\hat{\tau} = \inf\{\tau_j \mid 1 \leq j \leq k\}$. Note that here τ_j 's are prime periods of y_j 's for $1 \leq j \leq k$. Then by §2 of [WHL], for any $a > \hat{\tau}$, we can construct a function $\varphi_a \in C^\infty(\mathbf{R}, \mathbf{R}^+)$ which has 0 as its unique critical point in $[0, +\infty)$ such that φ_a is strictly convex for $t \geq 0$. Moreover, $\frac{\varphi'_a(t)}{t}$ is strictly decreasing for $t > 0$ together with $\lim_{t \rightarrow 0^+} \frac{\varphi'_a(t)}{t} = 1$ and $\varphi_a(0) = 0 = \varphi'_a(0)$. More precisely, we define φ_a via Propositions 2.2 and 2.4 in [WHL]. The precise dependence of φ_a on a is explained in Remark 2.3 of [WHL].

Define the Hamiltonian function $H_a(x) = a\varphi_a(j(x))$ and consider the fixed period problem

$$\begin{cases} \dot{x}(t) = JH'_a(x(t)), \\ x(1) = x(0). \end{cases} \quad (2.1)$$

Then $H_a \in C^3(\mathbf{R}^{2n} \setminus \{0\}, \mathbf{R}) \cap C^1(\mathbf{R}^{2n}, \mathbf{R})$ is strictly convex. Solutions of (2.1) are $x \equiv 0$ and $x = \rho y(\tau t)$ with $\frac{\varphi'_a(\rho)}{\rho} = \frac{\tau}{a}$, where (τ, y) is a solution of (1.1). In particular, nonzero solutions of (2.1) are one to one correspondent to solutions of (1.1) with period $\tau < a$.

In the following, we use the Clarke-Ekeland dual action principle. As usual, let G_a be the Fenchel transform of H_a defined by $G_a(y) = \sup\{x \cdot y - H_a(x) \mid x \in \mathbf{R}^{2n}\}$. Then $G_a \in C^2(\mathbf{R}^{2n} \setminus \{0\}, \mathbf{R}) \cap C^1(\mathbf{R}^{2n}, \mathbf{R})$ is strictly convex. Let

$$L_0^2(S^1, \mathbf{R}^{2n}) = \left\{ u \in L^2([0, 1], \mathbf{R}^{2n}) \mid \int_0^1 u(t) dt = 0 \right\}. \quad (2.2)$$

Define a linear operator $M : L_0^2(S^1, \mathbf{R}^{2n}) \rightarrow L_0^2(S^1, \mathbf{R}^{2n})$ by $\frac{d}{dt}Mu(t) = u(t)$, $\int_0^1 Mu(t) dt = 0$. The dual action functional on $L_0^2(S^1, \mathbf{R}^{2n})$ is defined by

$$\Psi_a(u) = \int_0^1 \left(\frac{1}{2}Ju \cdot Mu + G_a(-Ju) \right) dt. \quad (2.3)$$

Then the functional $\Psi_a \in C^{1,1}(L_0^2(S^1, \mathbf{R}^{2n}), \mathbf{R})$ is bounded from below and satisfies the Palais-Smale condition. Suppose x is a solution of (2.1). Then $u = \dot{x}$ is a critical point of Ψ_a . Conversely,

suppose u is a critical point of Ψ_a . Then there exists a unique $\xi \in \mathbf{R}^{2n}$ such that $Mu - \xi$ is a solution of (2.1). In particular, solutions of (2.1) are in one to one correspondence with critical points of Ψ_a . Moreover, $\Psi_a(u) < 0$ for every critical point $u \neq 0$ of Ψ_a .

Suppose u is a nonzero critical point of Ψ_a . Then following [Eke3] the formal Hessian of Ψ_a at u is defined by

$$Q_a(v, v) = \int_0^1 (Jv \cdot Mv + G_a''(-Ju)Jv \cdot Jv)dt,$$

which defines an orthogonal splitting $L_0^2 = E_- \oplus E_0 \oplus E_+$ of $L_0^2(S^1, \mathbf{R}^{2n})$ into negative, zero and positive subspaces. The index of u is defined by $i(u) = \dim E_-$ and the nullity of u is defined by $\nu(u) = \dim E_0$. Let $u = \dot{x}$ be the critical point of Ψ_a such that x corresponds to the closed characteristic (τ, y) on Σ . Then the index $i(u)$ and the nullity $\nu(u)$ defined above coincide with the Ekeland indices defined by I. Ekeland in [Eke1] and [Eke3]. Specially $1 \leq \nu(u) \leq 2n - 1$ always holds.

We have a natural S^1 -action on $L_0^2(S^1, \mathbf{R}^{2n})$ defined by $\theta \cdot u(t) = u(\theta + t)$ for all $\theta \in S^1$ and $t \in \mathbf{R}$. Clearly Ψ_a is S^1 -invariant. For any $\kappa \in \mathbf{R}$, we denote by

$$\Lambda_a^\kappa = \{u \in L_0^2(S^1, \mathbf{R}^{2n}) \mid \Psi_a(u) \leq \kappa\}. \quad (2.4)$$

For a critical point u of Ψ_a , we denote by

$$\Lambda_a(u) = \Lambda_a^{\Psi_a(u)} = \{w \in L_0^2(S^1, \mathbf{R}^{2n}) \mid \Psi_a(w) \leq \Psi_a(u)\}. \quad (2.5)$$

Clearly, both sets are S^1 -invariant. Since the S^1 -action preserves Ψ_a , if u is a critical point of Ψ_a , then the whole orbit $S^1 \cdot u$ is formed by critical points of Ψ_a . Denote by $\text{crit}(\Psi_a)$ the set of critical points of Ψ_a . Note that by the condition (F), the number of critical orbits of Ψ_a is finite. Hence as usual we can make the following definition.

Definition 2.1. *Suppose u is a nonzero critical point of Ψ_a and \mathcal{N} is an S^1 -invariant open neighborhood of $S^1 \cdot u$ such that $\text{crit}(\Psi_a) \cap (\Lambda_a(u) \cap \mathcal{N}) = S^1 \cdot u$. Then the S^1 -critical modules of $S^1 \cdot u$ are defined by*

$$C_{S^1, q}(\Psi_a, S^1 \cdot u) = H_q((\Lambda_a(u) \cap \mathcal{N})_{S^1}, ((\Lambda_a(u) \setminus S^1 \cdot u) \cap \mathcal{N})_{S^1}).$$

We have the following proposition for critical modules.

Proposition 2.2. (Proposition 3.2 of [WHL]) *The critical module $C_{S^1, q}(\Psi_a, S^1 \cdot u)$ is independent of a in the sense that if x_i are solutions of (2.1) with Hamiltonian functions $H_{a_i}(x) \equiv$*

$a_i \varphi_{a_i}(j(x))$ for $i = 1$ and 2 respectively such that both x_1 and x_2 correspond to the same closed characteristic (τ, y) on Σ . Then we have

$$C_{S^1, q}(\Psi_{a_1}, S^1 \cdot \dot{x}_1) \cong C_{S^1, q}(\Psi_{a_2}, S^1 \cdot \dot{x}_2), \quad \forall q \in \mathbf{Z}.$$

Now let $u \neq 0$ be a critical point of Ψ_a with multiplicity $\text{mul}(u) = m$, i.e., u corresponds to a closed characteristic $(m\tau, y) \subset \Sigma$ with (τ, y) being prime. Hence $u(t + \frac{1}{m}) = u(t)$ holds for all $t \in \mathbf{R}$ and the orbit of u , namely, $S^1 \cdot u \cong S^1/\mathbf{Z}_m \cong S^1$. Let $f : N(S^1 \cdot u) \rightarrow S^1 \cdot u$ be the normal bundle of $S^1 \cdot u$ in $L_0^2(S^1, \mathbf{R}^{2n})$ and let $f^{-1}(\theta \cdot u) = N(\theta \cdot u)$ be the fibre over $\theta \cdot u$, where $\theta \in S^1$. Let $DN(S^1 \cdot u)$ be the ϱ -disk bundle of $N(S^1 \cdot u)$ for some $\varrho > 0$ sufficiently small, i.e., $DN(S^1 \cdot u) = \{\xi \in N(S^1 \cdot u) \mid \|\xi\| < \varrho\}$ and let $DN(\theta \cdot u) = f^{-1}(\theta \cdot u) \cap DN(S^1 \cdot u)$ be the disk over $\theta \cdot u$. Clearly, $DN(\theta \cdot u)$ is \mathbf{Z}_m -invariant and we have $DN(S^1 \cdot u) = DN(u) \times_{\mathbf{Z}_m} S^1$, where the \mathbf{Z}_m -action is given by

$$(\theta, v, t) \in \mathbf{Z}_m \times DN(u) \times S^1 \mapsto (\theta \cdot v, \theta^{-1}t) \in DN(u) \times S^1.$$

Hence for an S^1 -invariant subset Γ of $DN(S^1 \cdot u)$, we have $\Gamma/S^1 = (\Gamma_u \times_{\mathbf{Z}_m} S^1)/S^1 = \Gamma_u/\mathbf{Z}_m$, where $\Gamma_u = \Gamma \cap DN(u)$. Since Ψ_a is not C^2 on $L_0^2(S^1, \mathbf{R}^{2n})$, we need to use a finite dimensional approximation introduced by Ekeland in order to apply Morse theory. More precisely, we can construct a finite dimensional submanifold $\Gamma(\iota)$ of $L_0^2(S^1, \mathbf{R}^{2n})$ which admits a \mathbf{Z}_ι -action with $m|\iota$. Moreover Ψ_a and $\Psi_a|_{\Gamma(\iota)}$ have the same critical points. $\Psi_a|_{\Gamma(\iota)}$ is C^2 in a small tubular neighborhood of the critical orbit $S^1 \cdot u$ and the Morse index and nullity of its critical points coincide with those of the corresponding critical points of Ψ_a . Let

$$D_\iota N(S^1 \cdot u) = DN(S^1 \cdot u) \cap \Gamma(\iota), \quad D_\iota N(\theta \cdot u) = DN(\theta \cdot u) \cap \Gamma(\iota). \quad (2.6)$$

Then we have

$$C_{S^1, *}(\Psi_a, S^1 \cdot u) \cong H_*(\Lambda_a(u) \cap D_\iota N(u), (\Lambda_a(u) \setminus \{u\}) \cap D_\iota N(u))^{\mathbf{Z}_m}. \quad (2.7)$$

Now we can apply the results of Gromoll and Meyer in [GrM1] to the manifold $D_{p\iota}N(u^p)$ with u^p as its unique critical point, where $p \in \mathbf{N}$. Then $\text{mul}(u^p) = pm$ is the multiplicity of u^p and the isotropy group $\mathbf{Z}_{pm} \subseteq S^1$ of u^p acts on $D_{p\iota}N(u^p)$ by isometries. According to Lemma 1 of [GrM1], we have a \mathbf{Z}_{pm} -invariant decomposition of $T_{u^p}(D_{p\iota}N(u^p))$

$$T_{u^p}(D_{p\iota}N(u^p)) = V^+ \oplus V^- \oplus V^0 = \{(x_+, x_-, x_0)\}$$

with $\dim V^- = i(u^p)$, $\dim V^0 = \nu(u^p) - 1$ and a \mathbf{Z}_{pm} -invariant neighborhood $B = B_+ \times B_- \times B_0$ for 0 in $T_{u^p}(D_{p\iota}N(u^p))$ together with two \mathbf{Z}_{pm} -invariant diffeomorphisms

$$\Phi : B = B_+ \times B_- \times B_0 \rightarrow \Phi(B_+ \times B_- \times B_0) \subset D_{p\iota}N(u^p)$$

and

$$\eta : B_0 \rightarrow W(u^p) \equiv \eta(B_0) \subset D_{p\iota}N(u^p)$$

such that $\Phi(0) = \eta(0) = u^p$ and

$$\Psi_a \circ \Phi(x_+, x_-, x_0) = |x_+|^2 - |x_-|^2 + \Psi_a \circ \eta(x_0), \quad (2.8)$$

with $d(\Psi_a \circ \eta)(0) = d^2(\Psi_a \circ \eta)(0) = 0$. As [GrM1], we call $W(u^p)$ a local *characteristic manifold* and $U(u^p) = B_-$ a local *negative disk* at u^p . By the proof of Lemma 1 of [GrM1], $W(u^p)$ and $U(u^p)$ are \mathbf{Z}_{pm} -invariant. Then we have

$$\begin{aligned} & H_*(\Lambda_a(u^p) \cap D_{p\iota}N(u^p), (\Lambda_a(u^p) \setminus \{u^p\}) \cap D_{p\iota}N(u^p)) \\ &= H_*(U(u^p), U(u^p) \setminus \{u^p\}) \otimes H_*(W(u^p) \cap \Lambda_a(u^p), (W(u^p) \setminus \{u^p\}) \cap \Lambda_a(u^p)), \end{aligned} \quad (2.9)$$

where

$$H_q(U(u^p), U(u^p) \setminus \{u^p\}) = \begin{cases} \mathbf{Q}, & \text{if } q = i(u^p), \\ 0, & \text{otherwise.} \end{cases} \quad (2.10)$$

Now we have the following proposition.

Proposition 2.3. (Proposition 3.10 of [WHL]) *Let $u \neq 0$ be a critical point of Ψ_a with $\text{mul}(u) = 1$. Then for all $p \in \mathbf{N}$ and $q \in \mathbf{Z}$, we have*

$$C_{S^1, q}(\Psi_a, S^1 \cdot u^p) \cong \left(H_{q-i(u^p)}(W(u^p) \cap \Lambda_a(u^p), (W(u^p) \setminus \{u^p\}) \cap \Lambda_a(u^p)) \right)^{\beta(u^p)\mathbf{Z}_p}, \quad (2.11)$$

where $\beta(u^p) = (-1)^{i(u^p)-i(u)}$. Thus

$$C_{S^1, q}(\Psi_a, S^1 \cdot u^p) = 0, \quad \text{for } q < i(u^p) \text{ or } q > i(u^p) + \nu(u^p) - 1. \quad (2.12)$$

In particular, if u^p is non-degenerate, i.e., $\nu(u^p) = 1$, then

$$C_{S^1, q}(\Psi_a, S^1 \cdot u^p) = \begin{cases} \mathbf{Q}, & \text{if } q = i(u^p) \text{ and } \beta(u^p) = 1, \\ 0, & \text{otherwise.} \end{cases} \quad (2.13)$$

We make the following definition

Definition 2.4. *Let $u \neq 0$ be a critical point of Ψ_a with $\text{mul}(u) = 1$. Then for all $p \in \mathbf{N}$ and $l \in \mathbf{Z}$, let*

$$\begin{aligned} k_{l, \pm 1}(u^p) &= \dim \left(H_l(W(u^p) \cap \Lambda_a(u^p), (W(u^p) \setminus \{u^p\}) \cap \Lambda_a(u^p)) \right)^{\pm \mathbf{Z}_p}, \\ k_l(u^p) &= \dim \left(H_l(W(u^p) \cap \Lambda_a(u^p), (W(u^p) \setminus \{u^p\}) \cap \Lambda_a(u^p)) \right)^{\beta(u^p)\mathbf{Z}_p}. \end{aligned}$$

$k_l(u^p)$'s are called critical type numbers of u^p .

We have the following properties for critical type numbers

Proposition 2.5. (Proposition 3.13 of [WHL]) *Let $u \neq 0$ be a critical point of Ψ_a with $\text{mul}(u) = 1$. Then there exists a minimal $K(u) \in \mathbf{N}$ such that*

$$\nu(u^{p+K(u)}) = \nu(u^p), \quad i(u^{p+K(u)}) - i(u^p) \in 2\mathbf{Z},$$

and $k_l(u^{p+K(u)}) = k_l(u^p)$ for all $p \in \mathbf{N}$ and $l \in \mathbf{Z}$. We call $K(u)$ the minimal period of critical modules of iterations of the functional Ψ_a at u .

For a closed characteristic (τ, y) on Σ , we denote by $y^m \equiv (m\tau, y)$ the m -th iteration of y for $m \in \mathbf{N}$. Let $a > \tau$ and choose φ_a as above. Determine ρ uniquely by $\frac{\varphi'_a(\rho)}{\rho} = \frac{\tau}{a}$. Let $x = \rho y(\tau t)$ and $u = \dot{x}$. Then we define the index $i(y^m)$ and nullity $\nu(y^m)$ of $(m\tau, y)$ for $m \in \mathbf{N}$ by

$$i(y^m) = i(u^m), \quad \nu(y^m) = \nu(u^m).$$

These indices are independent of a when a tends to infinity. Now the mean index of (τ, y) is defined by

$$\hat{i}(y) = \lim_{m \rightarrow \infty} \frac{i(y^m)}{m}.$$

Note that $\hat{i}(y) > 2$ always holds which was proved by Ekeland and Hofer in [EkH1] of 1987 (cf. Corollary 8.3.2 and Lemma 15.3.2 of [Lon4] for a different proof).

By Proposition 2.2, we can define the critical type numbers $k_l(y^m)$ of y^m to be $k_l(u^m)$, where u^m is the critical point of Ψ_a corresponding to y^m . We also define $K(y) = K(u)$. Then we have

Proposition 2.6. *We have $k_l(y^m) = 0$ for $l \notin [0, \nu(y^m) - 1]$ and it can take only values 0 or 1 when $l = 0$ or $l = \nu(y^m) - 1$. Moreover, the following properties hold (cf. Lemma 3.10 of [BaL1], [Cha1] and [MaW1]):*

- (i) $k_0(y^m) = 1$ implies $k_l(y^m) = 0$ for $1 \leq l \leq \nu(y^m) - 1$.
- (ii) $k_{\nu(y^m)-1}(y^m) = 1$ implies $k_l(y^m) = 0$ for $0 \leq l \leq \nu(y^m) - 2$.
- (iii) $k_l(y^m) \geq 1$ for some $1 \leq l \leq \nu(y^m) - 2$ implies $k_0(y^m) = k_{\nu(y^m)-1}(y^m) = 0$.
- (iv) If $\nu(y^m) \leq 3$, then at most one of the $k_l(y^m)$'s for $0 \leq l \leq \nu(y^m) - 1$ can be non-zero.
- (v) If $i(y^m) - i(y) \in 2\mathbf{Z} + 1$ for some $m \in \mathbf{N}$, then $k_0(y^m) = 0$.

Proof. By Definition 2.4 we have

$$k_l(y^m) \leq \dim H_l(W(u^m) \cap \Lambda_a(u^m), (W(u^m) \setminus \{u^m\}) \cap \Lambda_a(u^m)) \equiv \eta_l(y^m).$$

Then from Corollary 1.5.1 of [Cha1] or Corollary 8.4 of [MaW1], (i)-(iv) hold.

For (v), if $\eta_0(y^m) = 0$, then (v) follows directly from Definition 2.4.

By Corollary 8.4 of [MaW1], $\eta_0(y^m) = 1$ if and only if u^m is a local minimum in the local characteristic manifold $W(u^m)$. Hence $(W(u^m) \cap \Lambda_a(u^m), (W(u^m) \setminus \{u^m\}) \cap \Lambda_a(u^m)) = (\{u^m\}, \emptyset)$. By Definition 2.4, we have:

$$\begin{aligned} k_{0,+1}(u^m) &= \dim H_0(W(u^m) \cap \Lambda_a(u^m), (W(u^m) \setminus \{u^m\}) \cap \Lambda_a(u^m))^{+\mathbf{Z}_m} \\ &= \dim H_0(\{u^m\})^{+\mathbf{Z}_m} \\ &= 1. \end{aligned}$$

This implies $k_0(u^m) = k_{0,-1}(u^m) = 0$. ■

For a closed characteristic (τ, y) on Σ , we define as in [WHL]

$$\hat{\chi}(y) = \frac{1}{K(y)} \sum_{\substack{1 \leq m \leq K(y) \\ 0 \leq l \leq 2n-2}} (-1)^{i(y^m)+l} k_l(y^m). \quad (2.14)$$

In particular, if all y^m 's are non-degenerate, then by Proposition 2.3 we have

$$\hat{\chi}(y) = \begin{cases} (-1)^{i(y)}, & \text{if } i(y^2) - i(y) \in 2\mathbf{Z}, \\ \frac{(-1)^{i(y)}}{2}, & \text{otherwise.} \end{cases} \quad (2.15)$$

We have the following mean index identity for closed characteristics.

Theorem 2.7. (Theorem 1.2 of [WHL]) *Suppose $\Sigma \in \mathcal{H}(2n)$ satisfies $\#\tilde{\mathcal{J}}(\Sigma) < +\infty$. Denote all the geometrically distinct closed characteristics by $\{(\tau_j, y_j)\}_{1 \leq j \leq k}$. Then the following identity holds*

$$\sum_{1 \leq j \leq k} \frac{\hat{\chi}(y_j)}{\hat{i}(y_j)} = \frac{1}{2}.$$

Let Ψ_a be the functional defined by (2.3) for some $a \in \mathbf{R}$ large enough and let $\varepsilon > 0$ be small enough such that $[-\varepsilon, +\infty) \setminus \{0\}$ contains no critical values of Ψ_a . Denote by I_a the greatest integer in \mathbf{N}_0 such that $I_a < i(\tau, y)$ hold for all closed characteristics (τ, y) on Σ with $\tau \geq a$. Then by Section 5 of [WHL], we have

$$H_{S^1, q}(\Lambda_a^{-\varepsilon}) \cong H_{S^1, q}(\Lambda_a^\infty) \cong H_q(CP^\infty), \quad \forall q < I_a. \quad (2.16)$$

For any $q \in \mathbf{Z}$, let

$$M_q(\Lambda_a^{-\varepsilon}) = \sum_{1 \leq j \leq k, 1 \leq m_j < a/\tau_j} \dim C_{S^1, q}(\Psi_a, S^1 \cdot u_j^{m_j}). \quad (2.17)$$

Then the equivariant Morse inequalities for the space $\Lambda_a^{-\varepsilon}$ yield

$$M_q(\Lambda_a^{-\varepsilon}) \geq b_q(\Lambda_a^{-\varepsilon}), \quad (2.18)$$

$$\begin{aligned} M_q(\Lambda_a^{-\varepsilon}) &= M_{q-1}(\Lambda_a^{-\varepsilon}) + \cdots + (-1)^q M_0(\Lambda_a^{-\varepsilon}) \\ &\geq b_q(\Lambda_a^{-\varepsilon}) - b_{q-1}(\Lambda_a^{-\varepsilon}) + \cdots + (-1)^q b_0(\Lambda_a^{-\varepsilon}), \end{aligned} \quad (2.19)$$

where $b_q(\Lambda_a^{-\varepsilon}) = \dim H_{S^1, q}(\Lambda_a^{-\varepsilon})$. Now we have the following Morse inequalities for closed characteristics

Theorem 2.8. *Let $\Sigma \in \mathcal{H}(2n)$ satisfy $\#\tilde{\mathcal{J}}(\Sigma) < +\infty$. Denote all the geometrically distinct closed characteristics by $\{(\tau_j, y_j)\}_{1 \leq j \leq k}$. Let*

$$M_q = \lim_{a \rightarrow +\infty} M_q(\Lambda_a^{-\varepsilon}), \quad \forall q \in \mathbf{Z}, \quad (2.20)$$

$$b_q = \lim_{a \rightarrow +\infty} b_q(\Lambda_a^{-\varepsilon}) = \begin{cases} 1, & \text{if } q \in 2\mathbf{N}_0, \\ 0, & \text{otherwise.} \end{cases} \quad (2.21)$$

Then we have

$$M_q \geq b_q, \quad (2.22)$$

$$M_q - M_{q-1} + \cdots + (-1)^q M_0 \geq b_q - b_{q-1} + \cdots + (-1)^q b_0, \quad \forall q \in \mathbf{Z}. \quad (2.23)$$

Proof. As we have mentioned before, $\hat{i}(y_j) > 2$ holds for $1 \leq j \leq k$. Hence the Ekeland index satisfies $i(y_j^m) = i(u_j^m) \rightarrow \infty$ as $m \rightarrow \infty$ for $1 \leq j \leq k$. Note that $I_a \rightarrow +\infty$ as $a \rightarrow +\infty$. Now fix a $q \in \mathbf{Z}$ and a sufficiently great $a > 0$. By Propositions 2.2, 2.3 and (2.17), $M_i(\Lambda_a^{-\varepsilon})$ is invariant for all $a > A_q$ and $0 \leq i \leq q$, where $A_q > 0$ is some constant. Hence (2.20) is meaningful. Now for any a such that $I_a > q$, (2.16)-(2.19) imply that (2.21)-(2.23) hold. \blacksquare

3 Proofs of the main theorems

In this section, we give proofs of Theorems 1.1 and 1.2 by using the mean index identity of [WHL], Morse inequality and the index iteration theory developed by Long and his coworkers.

As Definition 1.1 of [LoZ1], we define

Definition 3.1. For $\alpha \in (1, 2)$, we define a map $\varrho_n: \mathcal{H}(2n) \rightarrow \mathbf{N} \cup \{+\infty\}$

$$\varrho_n(\Sigma) = \begin{cases} +\infty, & \text{if } \#\mathcal{V}(\Sigma, \alpha) = +\infty, \\ \min \left\{ \left\lceil \frac{i(x, 1) + 2S^+(x) - \nu(x, 1) + n}{2} \right\rceil \mid (\tau, x) \in \mathcal{V}_\infty(\Sigma, \alpha) \right\}, & \text{if } \#\mathcal{V}(\Sigma, \alpha) < +\infty, \end{cases} \quad (3.1)$$

where $\mathcal{V}(\Sigma, \alpha)$ and $\mathcal{V}_\infty(\Sigma, \alpha)$ are variationally visible and infinite variationally visible sets respectively given by Definition 1.4 of [LoZ1] (cf. Definition 15.3.3 of [Lon4]).

Theorem 3.2. (cf. Theorem 15.1.1 of [Lon4]) *Suppose $(\tau, y) \in \mathcal{J}(\Sigma)$. Then we have*

$$i(y^m) \equiv i(m\tau, y) = i(y, m) - n, \quad \nu(y^m) \equiv \nu(m\tau, y) = \nu(y, m), \quad \forall m \in \mathbf{N}, \quad (3.2)$$

where $i(y, m)$ and $\nu(y, m)$ are the Maslov-type index and nullity of $(m\tau, y)$ defined by Conley, Zehnder and Long (cf. §5.4 of [Lon4]).

Recall that for a principal $U(1)$ -bundle $E \rightarrow B$, the Fadell-Rabinowitz index (cf. [FaR1]) of E is defined to be $\sup\{k \mid c_1(E)^{k-1} \neq 0\}$, where $c_1(E) \in H^2(B, \mathbf{Q})$ is the first rational Chern class. For a $U(1)$ -space, i.e., a topological space X with a $U(1)$ -action, the Fadell-Rabinowitz index is defined to be the index of the bundle $X \times S^\infty \rightarrow X \times_{U(1)} S^\infty$, where $S^\infty \rightarrow CP^\infty$ is the universal $U(1)$ -bundle.

As in P.199 of [Eke3], choose some $\alpha \in (1, 2)$ and associate with U a convex function H such that $H(\lambda x) = \lambda^\alpha H(x)$ for $\lambda \geq 0$. Consider the fixed period problem

$$\begin{cases} \dot{x}(t) = JH'(x(t)), \\ x(1) = x(0). \end{cases} \quad (3.3)$$

Define

$$L_0^{\frac{\alpha}{\alpha-1}}(S^1, \mathbf{R}^{2n}) = \{u \in L^{\frac{\alpha}{\alpha-1}}(S^1, \mathbf{R}^{2n}) \mid \int_0^1 u dt = 0\}. \quad (3.4)$$

The corresponding Clarke-Ekeland dual action functional is defined by

$$\Phi(u) = \int_0^1 \left(\frac{1}{2} Ju \cdot Mu + H^*(-Ju) \right) dt, \quad \forall u \in L_0^{\frac{\alpha}{\alpha-1}}(S^1, \mathbf{R}^{2n}), \quad (3.5)$$

where Mu is defined by $\frac{d}{dt}Mu(t) = u(t)$ and $\int_0^1 Mu(t)dt = 0$, H^* is the Fenchel transform of H defined in §2.

For any $\kappa \in \mathbf{R}$, we denote by

$$\Phi^{\kappa-} = \{u \in L_0^{\frac{\alpha}{\alpha-1}}(S^1, \mathbf{R}^{2n}) \mid \Phi(u) < \kappa\}. \quad (3.6)$$

Then as in P.218 of [Eke3], we define

$$c_i = \inf\{\delta \in \mathbf{R} \mid \hat{I}(\Phi^{\delta-}) \geq i\}, \quad (3.7)$$

where \hat{I} is the Fadell-Rabinowitz index given above. Then by Proposition 3 in P.218 of [Eke3], we have

Proposition 3.3. *Every c_i is a critical value of Φ . If $c_i = c_j$ for some $i < j$, then there are infinitely many geometrically distinct closed characteristics on Σ .*

As in Definition 2.1, we define the following

Definition 3.4. *Suppose u is a nonzero critical point of Φ , and \mathcal{N} is an S^1 -invariant open neighborhood of $S^1 \cdot u$ such that $\text{crit}(\Phi) \cap (\Lambda(u) \cap \mathcal{N}) = S^1 \cdot u$. Then the S^1 -critical modules of $S^1 \cdot u$ is defined by*

$$C_{S^1, q}(\Phi, S^1 \cdot u) = H_q((\Lambda(u) \cap \mathcal{N})_{S^1}, ((\Lambda(u) \setminus S^1 \cdot u) \cap \mathcal{N})_{S^1}), \quad (3.8)$$

where $\Lambda(u) = \{w \in L_0^{\frac{\alpha}{\alpha-1}}(S^1, \mathbf{R}^{2n}) \mid \Phi(w) \leq \Phi(u)\}$.

Comparing with Theorem 4 in P.219 of [Eke3], we have the following

Proposition 3.5. *For every $i \in \mathbf{N}$, there exists a point $u \in L_0^{\frac{\alpha}{\alpha-1}}(S^1, \mathbf{R}^{2n})$ such that*

$$\Phi'(u) = 0, \quad \Phi(u) = c_i, \quad (3.9)$$

$$C_{S^1, 2(i-1)}(\Phi, S^1 \cdot u) \neq 0. \quad (3.10)$$

Proof. By Lemma 8 in P.206 of [Eke3], we can use Theorem 1.4.2 of [Cha1] in the equivariant form to obtain

$$H_{S^1, *}(\Phi^{c_i+\epsilon}, \Phi^{c_i-\epsilon}) = \bigoplus_{\Phi(u)=c_i} C_{S^1, *}(\Phi, S^1 \cdot u), \quad (3.11)$$

for ϵ small enough such that the interval $(c_i - \epsilon, c_i + \epsilon)$ contains no critical values of Φ except c_i .

Similar to P.431 of [EkH1], we have

$$H^{2(i-1)}((\Phi^{c_i+\epsilon})_{S^1}, (\Phi^{c_i-\epsilon})_{S^1}) \xrightarrow{q^*} H^{2(i-1)}((\Phi^{c_i+\epsilon})_{S^1}) \xrightarrow{p^*} H^{2(i-1)}((\Phi^{c_i-\epsilon})_{S^1}), \quad (3.12)$$

where p and q are natural inclusions. Denote by $f : (\Phi^{c_i+\epsilon})_{S^1} \rightarrow CP^\infty$ a classifying map and let $f^\pm = f|_{(\Phi^{c_i \pm \epsilon})_{S^1}}$. Then clearly each $f^\pm : (\Phi^{c_i \pm \epsilon})_{S^1} \rightarrow CP^\infty$ is a classifying map on $(\Phi^{c_i \pm \epsilon})_{S^1}$. Let $\eta \in H^2(CP^\infty)$ be the first universal Chern class.

By definition of c_i , we have $\hat{I}(\Phi^{c_i-\epsilon}) < i$, hence $(f^-)^*(\eta^{i-1}) = 0$. Note that $p^*(f^+)^*(\eta^{i-1}) = (f^-)^*(\eta^{i-1})$. Hence the exactness of (3.12) yields a $\sigma \in H^{2(i-1)}((\Phi^{c_i+\epsilon})_{S^1}, (\Phi^{c_i-\epsilon})_{S^1})$ such that $q^*(\sigma) = (f^+)^*(\eta^{i-1})$. Since $\hat{I}(\Phi^{c_i+\epsilon}) \geq i$, we have $(f^+)^*(\eta^{i-1}) \neq 0$. Hence $\sigma \neq 0$, and then

$$H_{S^1}^{2(i-1)}(\Phi^{c_i+\epsilon}, \Phi^{c_i-\epsilon}) = H^{2(i-1)}((\Phi^{c_i+\epsilon})_{S^1}, (\Phi^{c_i-\epsilon})_{S^1}) \neq 0.$$

Now the proposition follows from (3.11) and the universal coefficient theorem. ■

Proposition 3.6. *Suppose u is the critical point of Φ found in Proposition 3.5. Then we have*

$$C_{S^1, 2(i-1)}(\Psi_a, S^1 \cdot u_a) \neq 0, \quad (3.13)$$

where Ψ_a is given by (2.3) and $u_a \in L_0^2(S^1, \mathbf{R}^{2n})$ is its critical point corresponding to u in the natural sense.

Proof. Fix this u , we modify the function H only in a small neighborhood Ω of 0 as in [Eke1] so that the corresponding orbit of u does not enter Ω and the resulted function \tilde{H} satisfies similar properties as Definition 1 in P. 26 of [Eke1] by just replacing $\frac{3}{2}$ there by α . Define the dual action functional $\tilde{\Phi} : L_0^{\frac{\alpha}{\alpha-1}}(S^1, \mathbf{R}^{2n}) \rightarrow \mathbf{R}$ by

$$\tilde{\Phi}(v) = \int_0^1 \left(\frac{1}{2} Jv \cdot Mv + \tilde{H}^*(-Jv) \right) dt, \quad (3.14)$$

since clearly Φ and $\tilde{\Phi}$ are C^1 close to each other. Then by the continuity of critical modules (cf. Theorem 8.8 of [MaW1] or Theorem 1.5.6 in P.53 of [Cha1], which can be easily generalized to the equivariant sense) for the u in the proposition, we have

$$C_{S^1, *}(\Phi, S^1 \cdot u) \cong C_{S^1, *}(\tilde{\Phi}, S^1 \cdot u). \quad (3.15)$$

Using a finite dimensional approximation as in Lemma 3.9 of [Eke1], we have

$$C_{S^1, *}(\tilde{\Phi}, S^1 \cdot u) \cong H_*(\tilde{\Lambda}(u) \cap D_\iota N(u), (\tilde{\Lambda}(u) \setminus \{u\}) \cap D_\iota N(u))^{\mathbf{Z}_m}, \quad (3.16)$$

where $\tilde{\Lambda}(u) = \{w \in L_0^{\frac{\alpha}{\alpha-1}}(S^1, \mathbf{R}^{2n}) \mid \tilde{\Phi}(w) \leq \tilde{\Phi}(u)\}$ and $D_\iota N(u)$ is a \mathbf{Z}_m -invariant finite dimensional disk transversal to $S^1 \cdot u$ at u (cf. Lemma 3.9 of [WHL]), m is the multiplicity of u .

By Lemma 3.9 of [WHL], we have

$$C_{S^1, *}(\Psi_a, S^1 \cdot u_a) \cong H_*(\Lambda_a(u_a) \cap D_\iota N(u_a), (\Lambda_a(u_a) \setminus \{u_a\}) \cap D_\iota N(u_a))^{\mathbf{Z}_m}. \quad (3.17)$$

By the construction of H_a in [WHL], $H_a = \tilde{H}$ in a L^∞ -neighborhood of $S^1 \cdot u$. We remark here that multiplying H by a constant will not affect the corresponding critical modules, i.e., the corresponding critical orbits have isomorphic critical modules. Hence we can assume $H_a = H$ in a L^∞ -neighborhood of $S^1 \cdot u$ and then the above conclusion. Hence Ψ_a and $\tilde{\Phi}$ coincide in a L^∞ -neighborhood of $S^1 \cdot u$. Note also by Lemma 3.9 of [Eke1], the two finite dimensional approximations are actually the same. Hence we have

$$\begin{aligned} & H_*(\tilde{\Lambda}(u) \cap D_\iota N(u), (\tilde{\Lambda}(u) \setminus \{u\}) \cap D_\iota N(u))^{\mathbf{Z}_m} \\ & \cong H_*(\Lambda_a(u_a) \cap D_\iota N(u_a), (\Lambda_a(u_a) \setminus \{u_a\}) \cap D_\iota N(u_a))^{\mathbf{Z}_m}. \end{aligned} \quad (3.18)$$

Now the proposition follows from Proposition 3.5 and (3.16)-(3.18). ■

Now we can give:

Proof of Theorem 1.1. By the assumption (F) at the beginning of Section 2, we denote by $\{(\tau_j, y_j)\}_{1 \leq j \leq k}$ all the geometrically distinct closed characteristics on Σ , and by $\gamma_j \equiv \gamma_{y_j}$ the associated symplectic path of (τ_j, y_j) on Σ for $1 \leq j \leq k$. Then by Lemma 15.2.4 of [Lon4], there exist $P_j \in \text{Sp}(6)$ and $M_j \in \text{Sp}(4)$ such that

$$\gamma_j(\tau_j) = P_j^{-1}(N_1(1, 1) \diamond M_j)P_j, \quad \forall 1 \leq j \leq k, \quad (3.19)$$

where recall $N_1(1, b) = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$ for $b \in \mathbf{R}$.

Without loss of generality, by Theorem 1.3 of [LoZ1] (cf. Theorem 15.5.2 of [Lon4]), we may assume that (τ_1, y_1) has irrational mean index. Hence by Theorem 8.3.1 and Corollary 8.3.2 of

[Lon4], $M_1 \in \text{Sp}(4)$ in (3.19) can be connected to $R(\theta_1) \diamond Q_1$ within $\Omega^0(M_1)$ for some $\frac{\theta_1}{\pi} \notin \mathbf{Q}$ and $Q_1 \in \text{Sp}(2)$, where $R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ for $\theta \in \mathbf{R}$. Here we use notations from Definition 1.8.5 and Theorem 1.8.10 of [Lon4]. By Theorem 2.7, the following identity holds

$$\frac{\hat{\chi}(y_1)}{\hat{i}(y_1)} + \sum_{2 \leq j \leq k} \frac{\hat{\chi}(y_j)}{\hat{i}(y_j)} = \frac{1}{2}. \quad (3.20)$$

Now we have the following four cases according to the classification of basic norm forms (cf. Definition 1.8.9 of [Lon4]).

Case 1. $Q_1 = R(\theta_2)$ with $\frac{\theta_2}{\pi} \notin \mathbf{Q}$ or $Q_1 = D(\pm 2) \equiv \begin{pmatrix} \pm 2 & 0 \\ 0 & \pm \frac{1}{2} \end{pmatrix}$.

In this case, by Theorems 8.1.6 and 8.1.7 of [Lon4], we have $\nu(y_1^m) \equiv 1$, i.e., y_1^m is non-degenerate for all $m \in \mathbf{N}$. Hence it follows from (2.15) that $\hat{\chi}(y_1) \neq 0$. Now (3.20) implies that at least one of the y_j 's for $2 \leq j \leq k$ must have irrational mean index. Hence the theorem holds.

Case 2. $Q_1 = N_1(1, b)$ with $b = \pm 1, 0$.

We have two subcases according to the value of $\hat{\chi}(y_1)$.

Subcase 2.1. $\hat{\chi}(y_1) \neq 0$.

In this case, (3.20) implies that at least one of the y_j 's for $2 \leq j \leq k$ must have irrational mean index. Hence the theorem holds.

Subcase 2.2. $\hat{\chi}(y_1) = 0$.

Note that by Theorems 8.1.4 and 8.1.7 of [Lon4] and our above Proposition 2.5, we have $K(y_1) =$

1. Since $\nu(y_1) \leq 3$, it follows from Proposition 2.6 and (2.14):

$$0 = \hat{\chi}(y_1) = (-1)^{i(y_1)}(k_0(y_1) - k_1(y_1) + k_2(y_1)). \quad (3.21)$$

By (iv) of Proposition 2.6, at most one of $k_l(y_1)$ for $l = 0, 1, 2$ can be nonzero. Then (3.21) yields $k_l(y_1) = 0$ for $l = 0, 1, 2$. Hence it follows from Proposition 2.3 and Definition 2.4 that

$$C_{S^1, q}(\Psi_a, S^1 \cdot u_1^p) = 0, \quad \forall p \in \mathbf{N}, q \in \mathbf{Z}, \quad (3.22)$$

where we denote by u_1 the critical point of Ψ_a corresponding to (τ_1, y_1) . In other words, u_1^m is homologically invisible for all $m \in \mathbf{N}$.

By Propositions 3.5 and 3.6, we can replace the term *infinite variationally visible* in Definition 1.4 of [LoZ1] (cf. Definition 15.3.3 of [Lon4]) by *homologically visible*, and it is easy to check that all the results in [LoZ1] remain true under this change. Hence by Theorem 1.3 of [LoZ1] (cf. Theorem 15.5.2 of [Lon4]), at least one of the y_j 's for $2 \leq j \leq k$ must have irrational mean index, i.e., we can forget y_1 and consider only y_j 's for $2 \leq j \leq k$, then apply that theorem. This proves our theorem.

Case 3. $Q_1 = N_1(-1, 1)$.

In this case, by Theorems 8.1.4, 8.1.5 and 8.1.7 of [Lon4], we have

$$i(y_1, m) = mi(y_1, 1) + 2E\left(\frac{m\theta_1}{2\pi}\right) - 2, \quad \nu(y_1, m) = 1 + \frac{1 + (-1)^m}{2}, \quad \forall m \in \mathbf{N},$$

with $i(y_1, 1) \in 2\mathbf{Z} + 1$. Hence $K(y_1) = 2$ by Proposition 2.5. Because y_1 is non-degenerate, we have $k_l(y_1) = \delta_0^l$ for all $l \in \mathbf{Z}$ by (2.11), (2.13) and Definition 2.4. By Theorem 3.2, we have $i(y_1) = i(y_1, 1) - 3 \in 2\mathbf{Z}$ and $i(y_1^2) - i(y_1) = i(y_1, 2) - i(y_1, 1) \in 2\mathbf{Z} + 1$. Hence $k_0(y_1^2) = 0$ by (v) of Proposition 2.6. Because $\nu(y_1^2) = 2$, we have $k_l(y_1^2) = 0$ for $l \geq 2$. Then (2.14) implies

$$\hat{\chi}(y_1) = \frac{1 + k_1(y_1^2)}{2} \neq 0.$$

Now (3.20) implies that at least one of the y_j 's for $2 \leq j \leq k$ must have irrational mean index. Hence the theorem holds.

Case 4. $Q_1 = N_1(-1, b)$ with $b = 0, -1$ or $Q_1 = R(\theta_2)$ with $\frac{\theta_2}{2\pi} = \frac{L}{N} \in \mathbf{Q} \cap (0, 1)$ with $N > 1$ and $(L, N) = 1$.

Note first that if $Q_1 = N_1(-1, b)$ with $b = 0, -1$, then Theorems 8.1.5 and 8.1.7 of [Lon4] imply that their index iteration formulae coincide with that of a rotational matrix $R(\theta)$ with $\theta = \pi$. Hence in the following we shall only consider the case $Q_1 = R(\theta_2)$ with $\theta_2/\pi \in (0, 2) \cap \mathbf{Q}$. The same argument also shows that the theorem is true for $Q_1 = N_1(-1, -1)$.

By Theorems 8.1.4 and 8.1.7 of [Lon4], we have

$$i(y_1, m) = m(i(y_1, 1) - 1) + 2E\left(\frac{m\theta_1}{2\pi}\right) + 2E\left(\frac{m\theta_2}{2\pi}\right) - 3, \quad (3.23)$$

$$\nu(y_1, m) = 3 - 2\varphi\left(\frac{m\theta_2}{2\pi}\right), \quad (3.24)$$

with $i(y_1, 1) \in 2\mathbf{Z} + 1$ and all $m \in \mathbf{N}$. By Proposition 2.5, we have $K(y_1) = N$. Note that because y_1^m is non-degenerate for $1 \leq m \leq N - 1$, $k_l(y_1^m) = \delta_0^l$ holds for $1 \leq m \leq N - 1$ by (2.11), (2.13) and Definition 2.4. By Theorem 3.2, we have $i(y_1) = i(y_1, 1) - 3 \in 2\mathbf{Z}$. Then (2.14) implies

$$\hat{\chi}(y_1) = \frac{N - 1 + k_0(y_1^N) - k_1(y_1^N) + k_2(y_1^N)}{N}. \quad (3.25)$$

This follows from $\nu(y_1^m) \leq 3$ for all $m \in \mathbf{N}$.

We have two subcases according to the value of $\hat{\chi}(y_1)$.

Subcase 4.1. $\hat{\chi}(y_1) \neq 0$.

In this subcase, (3.20) implies that at least one of the y_j 's for $2 \leq j \leq k$ must have irrational mean index. Hence the theorem holds.

Subcase 4.2. $\hat{\chi}(y_1) = 0$.

In this subcase, it follows from (3.25) and (iv) of Proposition 2.6 that

$$k_1(y_1^N) = N - 1 > 0. \quad (3.26)$$

Using the common index jump theorem (Theorems 4.3 and 4.4 of [LoZ1], Theorems 11.2.1 and 11.2.2 of [Lon4]), we obtain some $(T, m_1, \dots, m_k) \in \mathbf{N}^{k+1}$ such that $\frac{m_1 \theta_2}{\pi} \in \mathbf{Z}$ (cf. (11.2.18) of [Lon4]) and the following hold by (11.2.6), (11.2.7) and (11.2.26) of [Lon4]:

$$i(y_j, 2m_j) \geq 2T - \frac{e(\gamma_j(\tau_j))}{2}, \quad (3.27)$$

$$i(y_j, 2m_j) + \nu(y_j, 2m_j) \leq 2T + \frac{e(\gamma_j(\tau_j))}{2} - 1, \quad (3.28)$$

$$i(y_j, 2m_j + 1) = 2T + i(y_j, 1). \quad (3.29)$$

$$i(y_j, 2m_j - 1) + \nu(y_j, 2m_j - 1) = 2T - (i(y_j, 1) + 2S_{\gamma_j(\tau_j)}^+(1) - \nu(y_j, 1)). \quad (3.30)$$

By P. 340 of [Lon4], we have

$$\begin{aligned} & 2S_{\gamma_j(\tau_j)}^+(1) - \nu(y_j, 1) \\ &= 2S_{N_1(1, 1)}^+(1) - \nu_1(N_1(1, 1)) + 2S_{M_j}^+(1) - \nu_1(M_j) \\ &= 1 + 2S_{M_j}^+(1) - \nu_1(M_j) \\ &\geq -1, \quad 1 \leq j \leq k. \end{aligned} \quad (3.31)$$

In the last inequality, we have used the fact that the worst case for $2S_{M_j}^+(1) - \nu_1(M_j)$ happens when $M_j = N_1(1, -1)^{\circ 2}$ which gives the lower bound -2 .

By Corollary 15.1.4 of [Lon4], we have $i(y_j, 1) \geq 3$ for $1 \leq j \leq k$. Note that $e(\gamma_j(\tau_j)) \leq 6$ for $1 \leq j \leq k$. Hence Theorem 10.2.4 of [Lon4] yields

$$\begin{aligned} i(y_j, m) + \nu(y_j, m) &\leq i(y_j, m+1) - i(y_j, 1) + \frac{e(\gamma_j(\tau_j))}{2} - 1 \\ &\leq i(y_j, m+1) - 1, \quad \forall m \in \mathbf{N}, 1 \leq j \leq k. \end{aligned} \quad (3.32)$$

Specially, we have

$$i(y_j, m) < i(y_j, m+1), \quad \forall m \in \mathbf{N}, 1 \leq j \leq k.$$

Now (3.27)-(3.30) become

$$i(y_j, 2m_j) \geq 2T - 3, \quad (3.33)$$

$$i(y_j, 2m_j) + \nu(y_j, 2m_j) - 1 \leq 2T + 1, \quad (3.34)$$

$$i(y_j, 2m_j + m) \geq 2T + 3, \quad \forall m \geq 1, \quad (3.35)$$

$$i(y_j, 2m_j - m) + \nu(y_j, 2m_j - m) - 1 \leq 2T - 3, \quad \forall m \geq 1, \quad (3.36)$$

where $1 \leq j \leq k$. By Proposition 2.3, we have

$$C_{S^1, q}(\Psi_a, S^1 \cdot u_1^{2m_1}) = \delta_{i(u_1^{2m_1})+1}^q \mathbf{Q}^{k_1(y_1^N)} = \delta_{i(u_1^{2m_1})+1}^q \mathbf{Q}^{N-1}, \quad (3.37)$$

Note that by Theorem 3.2

$$i(y_j^m) = i(y_j, m) - 3, \quad \forall m \in \mathbf{N}, \quad 1 \leq j \leq k. \quad (3.38)$$

Hence (3.23) implies that $i(y_1^m)$ is even for all $m \in \mathbf{N}$. This together with (3.35)-(3.38) and Proposition 2.3 yield

$$C_{S^1, 2T-2}(\Psi_a, S^1 \cdot u_1^m) = 0, \quad \forall m \in \mathbf{N}, \quad (3.39)$$

$$C_{S^1, 2T-4}(\Psi_a, S^1 \cdot u_1^m) = 0, \quad \forall m \in \mathbf{N}, \quad (3.40)$$

$$C_{S^1, 2T-2}(\Psi_a, S^1 \cdot u_j^m) = 0, \quad \forall m \neq 2m_j, \quad 2 \leq j \leq k. \quad (3.41)$$

$$C_{S^1, 2T-4}(\Psi_a, S^1 \cdot u_j^m) = 0, \quad \forall m \neq 2m_j, \quad 2 \leq j \leq k. \quad (3.42)$$

In fact, by (3.35), (3.36) and (3.38) for $1 \leq j \leq k$, we have $i(u_j^m) = i(y_j^m) \geq 2T$ for all $m > 2m_j$ and $i(u_j^m) + \nu(u_j^m) - 1 = i(y_j^m) + \nu(y_j^m) - 1 \leq 2T - 6$ for all $m < 2m_j$. Thus (3.41)-(3.42) hold and (3.39)-(3.40) hold for $m \neq 2m_1$ by Proposition 2.3. Since $i(y_1^{2m_1})$ is even, by (3.37), (3.39)-(3.40) also hold for $m = 2m_1$.

Thus by Propositions 3.5 and 3.6 we can find $p, q \in \{2, \dots, k\}$ such that

$$\Phi'(u_p^{2m_p}) = 0, \quad \Phi(u_p^{2m_p}) = c_{T-1}, \quad C_{S^1, 2T-4}(\Psi_a, S^1 \cdot u_p^{2m_p}) \neq 0, \quad (3.43)$$

$$\Phi'(u_q^{2m_q}) = 0, \quad \Phi(u_q^{2m_q}) = c_T, \quad C_{S^1, 2T-2}(\Psi_a, S^1 \cdot u_q^{2m_q}) \neq 0, \quad (3.44)$$

where we denote also by $u_p^{2m_p}$ and $u_q^{2m_q}$ the corresponding critical points of Φ and which will not be confused.

Note that by assumption (F) and Proposition 3.3, we have $c_{T-1} < c_T$. Hence $p \neq q$ by (3.43) and (3.44). Then the proof of Lemma 3.1 in [LoZ1](cf. lemma 15.3.5 of [Lon4]) yields

$$\hat{i}(y_p, 2m_p) < \hat{i}(y_q, 2m_q). \quad (3.45)$$

Now if both $\hat{i}(y_p) \in \mathbf{Q}$ and $\hat{i}(y_q) \in \mathbf{Q}$ hold, then the proof of Theorem 5.3 in [LoZ1](cf. Theorem 15.5.2 of [Lon4]) yields

$$\hat{i}(y_p, 2m_p) = \hat{i}(y_q, 2m_q).$$

Note that we may choose T firstly such that $\frac{T}{M\hat{i}(y_j)} \in \mathbf{N}$ hold for all $\hat{i}(y_j) \in \mathbf{Q}$ then use the proof of Theorem 5.3 in [LoZ1]. Here M is the least integer in \mathbf{N} that satisfies $\frac{M\theta}{\pi} \in \mathbf{Z}$, whenever

$e^{\sqrt{-1}\theta} \in \sigma(\gamma_j(\tau_j))$ and $\frac{\theta}{\pi} \in \mathbf{Q}$ for some $1 \leq j \leq k$. Hence either $\hat{i}(y_p) \notin \mathbf{Q}$ or $\hat{i}(y_q) \notin \mathbf{Q}$ holds. This together with $\hat{i}(y_1) \notin \mathbf{Q}$ and $p, q \neq 1$ proves the theorem. \blacksquare

Proof of Theorem 1.2. We denote by $\{(\tau_j, y_j)\}_{1 \leq j \leq 3}$ the three geometrically distinct closed characteristics on Σ , and by $\gamma_j \equiv \gamma_{y_j}$ the associated symplectic path of (τ_j, y_j) on Σ for $1 \leq j \leq 3$. Then as in the proof of Theorem 1.1, there exist $P_j \in \text{Sp}(6)$ and $M_j \in \text{Sp}(4)$ such that

$$\gamma_j(\tau_j) = P_j^{-1}(N_1(1, 1) \diamond M_j)P_j, \quad \forall 1 \leq j \leq 3. \quad (3.46)$$

As in P.356 of [LoZ1], if there is no (τ_j, y_j) with $M_j = N_1(1, -1)^{\diamond 2}$ and $i(y_j, 1) = 3$ in $\mathcal{V}_\infty(\Sigma, \alpha)$, then $\varrho_n(\Sigma) = 3$. Hence we can use Theorem 1.4 of [LoZ1] (Theorem 15.5.2 of [Lon4]) to obtain the existence of at least two elliptic closed characteristics. This proves the theorem.

It remains to show that if there exists a (τ_j, y_j) with $M_j = N_1(1, -1)^{\diamond 2}$ and $i(y_j, 1) = 3$ in $\mathcal{V}_\infty(\Sigma, \alpha)$, we have at least two elliptic closed characteristics. We may assume $M_1 = N_1(1, -1)^{\diamond 2}$ and $i(y_1, 1) = 3$ without loss of generality. Note that (τ_1, y_1) has rational mean index by Theorem 8.3.1 of [Lon4] and Theorem 3.2.

By Theorem 1.3 of [LoZ1], we may assume that (τ_2, y_2) has irrational mean index. Hence by Theorem 8.3.1 and Corollary 8.3.2 of [Lon4], $M_2 \in \text{Sp}(4)$ in (3.46) can be connected to $R(\theta_2) \diamond Q_2$ within $\Omega^0(M_2)$ for some $\frac{\theta_2}{\pi} \in \mathbf{R} \setminus \mathbf{Q}$ and $Q_2 \in \text{Sp}(2)$, where $R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ for $\theta \in \mathbf{R}$. Here we use notations from Definition 1.8.5 and Theorem 1.8.10 of [Lon4]. By Theorem 2.7, the following identity holds

$$\frac{\hat{\chi}(y_1)}{\hat{i}(y_1)} + \frac{\hat{\chi}(y_2)}{\hat{i}(y_2)} + \frac{\hat{\chi}(y_3)}{\hat{i}(y_3)} = \frac{1}{2}. \quad (3.47)$$

Now if Q_2 is not hyperbolic, then both (τ_1, y_1) and (τ_2, y_2) are elliptic, so the theorem holds.

Hence it remains to consider the case that Q_2 is hyperbolic. Clearly (τ_2, y_2) is non-degenerate, then it follows from (2.15) that $\hat{\chi}(y_2) \neq 0$. Hence (3.47) implies that $\hat{i}(y_3) \in \mathbf{R} \setminus \mathbf{Q}$. Now by Theorem 8.3.1 and Corollary 8.3.2 of [Lon4], $M_3 \in \text{Sp}(4)$ in (3.46) can be connected to $R(\theta_3) \diamond Q_3$ within $\Omega^0(M_3)$ for some $\frac{\theta_3}{\pi} \in \mathbf{R} \setminus \mathbf{Q}$ and $Q_3 \in \text{Sp}(2)$. By the same reason as above, it suffices to consider the case that Q_3 is hyperbolic.

Combining all the above, the only case we need to kick off is that

$$M_1 = N_1(1, -1)^{\diamond 2}, \quad i(y_1, 1) = 3, \quad M_2 = R(\theta_2) \diamond Q_2, \quad M_3 = R(\theta_3) \diamond Q_3, \quad (3.48)$$

where both Q_2 and Q_3 are hyperbolic. Hence by Theorem 8.3.1 of [Lon4] and Theorem 3.2, we have

$$i(y_1^m) = m(i(y_1, 1) + 1) - 4 = 4m - 4, \quad \nu(y_1^m) = 3, \quad \forall m \in \mathbf{N}, \quad (3.49)$$

$$i(y_j^m) = m(i(y_j) + 3) + 2E\left(\frac{m\theta_j}{2\pi}\right) - 5, \quad \nu(y_j^m) = 1, \quad \forall m \in \mathbf{N}, \quad j = 2, 3. \quad (3.50)$$

By Proposition 2.5, we have $K(y_1) = 1$. Note that $i(y_1) = i(y_1, 1) - 3 = 0$ by Theorem 3.2. Hence Proposition 2.6, (2.14) and (2.15) imply

$$\hat{\chi}(y_1) \leq 1, \quad \hat{\chi}(y_1) \in \mathbf{Z}, \quad (3.51)$$

$$\hat{\chi}(y_j) = \begin{cases} -1, & \text{if } i(y_j) \in 2\mathbf{N}_0 + 1, \\ \frac{1}{2}, & \text{if } i(y_j) \in 2\mathbf{N}_0, \end{cases} \quad j = 1, 2. \quad (3.52)$$

By (3.49) and (3.50), we have

$$\hat{i}(y_1) = 4, \quad (3.53)$$

$$\hat{i}(y_j) = i(y_j) + 3 + \frac{\theta_j}{\pi} > 3, \quad j = 2, 3. \quad (3.54)$$

By (3.51)-(3.54), in order to make (3.47) hold, we must have

$$\hat{\chi}(y_1) = 1, \quad (3.55)$$

$$i(y_j) \in 2\mathbf{N}_0, \quad j = 2, 3. \quad (3.56)$$

In fact, by (3.52) and (3.54), we have

$$\frac{\hat{\chi}(y_2)}{\hat{i}(y_2)} + \frac{\hat{\chi}(y_3)}{\hat{i}(y_3)} < \frac{1}{6} + \frac{1}{6} < \frac{1}{2}.$$

Thus to make (3.47) hold, we must have $\frac{\hat{\chi}(y_1)}{\hat{i}(y_1)} > 0$. Hence (3.55) follows from (3.51). Now if $i(y_2) \in 2\mathbf{N}_0 + 1$ or $i(y_3) \in 2\mathbf{N}_0 + 1$ holds, then by (3.52), we have

$$\frac{\hat{\chi}(y_1)}{\hat{i}(y_1)} + \frac{\hat{\chi}(y_2)}{\hat{i}(y_2)} + \frac{\hat{\chi}(y_3)}{\hat{i}(y_3)} < \frac{1}{4} + \frac{1}{6} < \frac{1}{2}.$$

Hence (3.56) must hold.

By (2.14), (3.49) and (3.55), we have $1 = \hat{\chi}(y_1) = k_0(y_1) - k_1(y_1) + k_2(y_1)$. Since $\nu(y_1) = 3$, by Proposition 2.6, only one of $k_0(y_1)$, $k_1(y_1)$, $k_2(y_1)$ can be nonzero. Hence we obtain

$$k_1(y_1) = 0, \quad k_0(y_1) + k_2(y_1) = 1, \quad (3.57)$$

By Proposition 2.3, we have

$$C_{S^1, q}(\Psi_a, S^1 \cdot u_j^p) = 0, \quad \forall p \in \mathbf{N}, q \in 2\mathbf{Z} + 1, 1 \leq j \leq 3. \quad (3.58)$$

In fact, by (3.49), we have $i(y_1^m) \in 2\mathbf{N}$ for all $m \in \mathbf{N}$. Thus (3.58) holds for $j = 1$ by (2.11), (3.57) and Definition 2.4. By (3.50) and (3.56), for $j = 2, 3$, we have $i(y_j^m) \in 2\mathbf{N}$ when $m \in 2\mathbf{N}_0 + 1$ and $i(y_j^m) \in 2\mathbf{N}_0 + 1$ when $m \in 2\mathbf{N}$. In particular, all y_j^m are non-degenerate for $m \in \mathbf{N}$ and $j = 2, 3$. Thus (3.58) holds for $j = 2, 3$ by (2.13).

Note that (3.58) implies

$$M_q = 0, \quad \forall q \in 2\mathbf{Z} + 1. \quad (3.59)$$

Together with the Morse inequality Theorem 2.8, it yields

$$-M_{2k} - \cdots - M_2 - M_0 \geq -b_{2k} - \cdots - b_2 - b_0.$$

Thus together with the Morse inequality again, it yields

$$b_{2k} + \cdots + b_2 + b_0 \geq M_{2k} + \cdots + M_2 + M_0 \geq b_{2k} + \cdots + b_2 + b_0,$$

for all $k \geq 0$. Therefore we obtain

$$M_q = b_q, \quad \forall q \in \mathbf{Z}. \quad (3.60)$$

By (3.57), we have two cases according to the values of $k_l(y_1)$ s.

Case 1. $k_0(y_1) = 1$ and $k_2(y_1) = 0$.

In this case, by Propositions 2.3, 2.5 and Definition 2.4, we have

$$\dim C_{S^1, q}(\Psi_a, S^1 \cdot u_1^m) = \delta_{4m-4}^q, \quad \forall m \in \mathbf{N}, q \in \mathbf{Z}. \quad (3.61)$$

Then by (3.60) and (2.21), we must have

$$C_{S^1, 4m-4}(\Psi_a, S^1 \cdot u_j^p) = 0, \quad \forall p, m \in \mathbf{N}, j = 2, 3. \quad (3.62)$$

By (3.60) and (2.21) again, $M_2 = b_2 = 1$ implies

$$C \equiv C_{S^1, 2}(\Psi_a, S^1 \cdot u_j^p) = \mathbf{Q}, \quad (3.63)$$

for some $p \in \mathbf{N}$ and $j = 2$ or 3 . If $p \geq 2$, by (3.50), we have

$$i(y_j^p) \geq 3p + 2E\left(\frac{p\theta_j}{2\pi}\right) - 5 \geq 3. \quad (3.64)$$

Thus $C = 0$ by Proposition 2.3. Hence $p = 1$. Without loss of generality, we assume $j = 2$. Then by Proposition 2.3 and (3.63), we have

$$i(y_2) = 2. \quad (3.65)$$

Then by (3.50), we have

$$i(y_2^m) \geq 7, \quad \forall m \geq 2. \quad (3.66)$$

By (3.60) and (2.21), $M_6 = b_6 = 1$ implies

$$C_{S^1, 6}(\Psi_a, S^1 \cdot u_j^p) = \mathbf{Q}, \quad (3.67)$$

for some $p \in \mathbf{N}$ and $j = 2$ or 3 . By (3.65) and (3.66), we have $j \neq 2$, i.e., $j = 3$. We must have $p = 1$. In fact, by (3.61) and (3.63), y_1^m and y_2^n already contribute a 1 to M_q for $q = 0, 2, 4$. Hence by (2.21), (3.60) and (3.56), we have $i(y_3) \geq 6$, and then $i(y_3^m) \geq 15$ by (3.50) for $m \geq 2$. Thus $p = 1$ follows from Proposition 2.3. Now we have

$$i(y_3) = 6. \quad (3.68)$$

Hence by (3.53) and (3.55) for y_1 , (3.50), (3.52), (3.65) and (3.68) for y_2 and y_3 , we have

$$\frac{\hat{\chi}(y_1)}{\hat{i}(y_1)} + \frac{\hat{\chi}(y_2)}{\hat{i}(y_2)} + \frac{\hat{\chi}(y_3)}{\hat{i}(y_3)} = \frac{1}{4} + \frac{1}{2(5 + \frac{\theta_2}{\pi})} + \frac{1}{2(9 + \frac{\theta_3}{\pi})} < \frac{1}{2}.$$

This contradicts (3.47) and proves Case 1.

Case 2. $k_0(y_1) = 0$ and $k_2(y_1) = 1$.

The study for this case is similar to that of Case 1. Thus we are rather sketch here.

In this case, by Proposition 2.3 and Definition 2.4, we have

$$\dim C_{S^1, q}(\Psi_a, S^1 \cdot u_1^m) = \delta_{4m-2}^q, \quad \forall m \in \mathbf{N}, q \in \mathbf{Z}. \quad (3.69)$$

Then by (3.60) and (2.21), we must have

$$C_{S^1, 4m-2}(\Psi_a, S^1 \cdot u_j^p) = 0, \quad \forall p, m \in \mathbf{N}, j = 2, 3. \quad (3.70)$$

By (3.69), (3.60) and (2.21), $M_0 = b_0 = 1$ implies

$$C_{S^1, 0}(\Psi_a, S^1 \cdot u_j^p) = \mathbf{Q}, \quad (3.71)$$

for some $p \in \mathbf{N}$ and $j = 2$ or 3 . By (3.64), we have $p = 1$. Without loss of generality, we assume $j = 2$. Then by Proposition 2.3 and (3.50), we have

$$i(y_2) = 0, \quad i(y_2^m) \geq 6, \quad \forall m \geq 3. \quad (3.72)$$

By (3.60) and (2.21), $M_4 = b_4 = 1$ implies

$$C_{S^1, 4}(\Psi_a, S^1 \cdot u_j^p) = \mathbf{Q}, \quad (3.73)$$

for some $p \in \mathbf{N}$ and $j = 2$ or 3 . By (3.69) and (3.72), as in the verification of (3.68), we have $j = 3$ and $p = 1$. Then by Proposition 2.3, we have

$$i(y_3) = 4. \quad (3.74)$$

Hence by (3.53) and (3.55) for y_1 , (3.50), (3.52), (3.72) and (3.74) for y_2 and y_3 , we have

$$\frac{\hat{\chi}(y_1)}{\hat{i}(y_1)} + \frac{\hat{\chi}(y_2)}{\hat{i}(y_2)} + \frac{\hat{\chi}(y_3)}{\hat{i}(y_3)} = \frac{1}{4} + \frac{1}{2(3 + \frac{\theta_2}{\pi})} + \frac{1}{2(7 + \frac{\theta_3}{\pi})} < \frac{1}{2}.$$

This contradicts (3.47) and proves Case 2 and then the whole theorem. ■

Acknowledgements. I would like to sincerely thank my Ph. D. thesis advisor, Professor Yiming Long, for introducing me to Hamiltonian dynamics and for his valuable help and encouragement during the writing of this paper. I would like to say that how enjoyable it is to work with him. I would like to sincerely thank the referee for his/her careful reading and valuable comments and suggestions.

References

- [BaL1] V. Bangert and Y. Long, The existence of two closed geodesics on every Finsler 2-sphere. (2005) Preprint.
- [Chal] K. C. Chang, Infinite Dimensional Morse Theory and Multiple Solution Problems. Birkhäuser. Boston. 1993.
- [CoZ1] C. Conley and E. Zehnder, Morse-type index theory for flows and periodic solutions for Hamiltonian equations. *Comm. Pure. Appl. Math.* 37 (1984) 207-253.
- [DDE1] Dell’Antonio, G., B. D’Onofrio and I. Ekeland, Les systèm hamiltoniens convexes et pairs ne sont pas ergodiques en general. *C. R. Acad. Sci. Paris. Series I.* 315 (1992), 1413-1415.
- [Eke1] I. Ekeland, Une théorie de Morse pour les systèmes hamiltoniens convexes. *Ann. IHP. Anal. non Linéaire.* 1 (1984) 19-78.
- [Eke2] I. Ekeland, An index throphy for periodic solutions of convex Hamiltonian systems. *Proc. Symp. in Pure Math.* 45 (1986) 395-423.
- [Eke3] I. Ekeland, Convexity Methods in Hamiltonian Mechanics. Springer-Verlag. Berlin. 1990.
- [EkH1] I. Ekeland and H. Hofer, Convex Hamiltonian energy surfaces and their closed trajectories. *Comm. Math. Phys.* 113 (1987) 419-467.
- [EkL1] I. Ekeland and L. Lassoued, Multiplicité des trajectoires fermées d’un système hamiltonien sur une hypersurface d’énergie convexe. *Ann. IHP. Anal. non Linéaire.* 4 (1987) 1-29.

- [FaR1] E. Fadell and P. Rabinowitz, Generalized comological index theories for Lie group actions with an application to bifurcation equations for Hamiltonian systems.
- [GrM1] D. Gromoll and W. Meyer, On differentiable functions with isolated critical points. *Topology*. 8 (1969) 361-369.
- [HWZ1] H. Hofer, K. Wysocki and E. Zehnder, The dynamics on three-dimensional strictly convex energy surfaces. *Ann. of Math.* 148 (1998) 197-289.
- [LLZ1] C. Liu, Y. Long and C. Zhu, Multiplicity of closed characteristics on symmetric convex hypersurfaces in \mathbf{R}^{2n} . *Math. Ann.* 323 (2002), 201-215.
- [Lon1] Y. Long, Maslov-type index, degenerate critical points and asymptotically linear Hamiltonian systems. *Science in China. Series A.* 33(1990), 1409-1419.
- [Lon2] Y. Long, Hyperbolic closed characteristics on compact convex smooth hypersurfaces in \mathbf{R}^{2n} . *J. Diff. Equa.* 150 (1998), 227-249.
- [Lon3] Y. Long, Precise iteration formulae of the Maslov-type index theory and ellipticity of closed characteristics. *Advances in Math.* 154 (2000), 76-131.
- [Lon4] Y. Long, Index Theory for Symplectic Paths with Applications. Progress in Math. 207, Birkhäuser. Basel. 2002.
- [Lon5] Y. Long, Index iteration theory for symplectic paths with applications to nonlinear Hamiltonian systems. *Proc. of Inter. Congress of Math. 2002*. Vol.II, 303-313. Higher Edu. Press. Beijing. 2002.
- [Lon6] Y. Long, Index iteration theory for symplectic paths and multiple periodic solution orbits. *Frontiers of Math.* 8 (2006) 341-353.
- [LoW1] Y. Long and W. Wang, Stability of closed characteristics on compact convex hypersurfaces, *Memory Volume for Professor S. S. Chern*. Ed. by P. Griffiths. Nankai Tracts in Mathematics Vol. 11, World Scientific. 313-333..
- [LZe1] Y. Long and E. Zehnder, Morse theory for forced oscillations of asymptotically linear Hamiltonian systems. In *Stoc. Proc. Phys. and Geom.*, S. Albeverio et al. ed. World Sci. (1990) 528-563.

- [LoZ1] Y. Long and C. Zhu, Closed characteristics on compact convex hypersurfaces in \mathbf{R}^{2n} . *Ann. of Math.* 155 (2002) 317-368.
- [MaW1] J. Mawhin and M. Willem, Critical Point Theory and Hamiltonian Systems. Springer. New York. 1989.
- [Rab1] P. H. Rabinowitz, Periodic solutions of Hamiltonian systems. *Comm. Pure Appl. Math.* 31 (1978) 157-184.
- [Szu1] A. Szulkin, Morse theory and existence of periodic solutions of convex Hamiltonian systems. *Bull. Soc. Math. France.* 116 (1988) 171-197.
- [Vit1] C. Viterbo, Equivariant Morse theory for starshaped Hamiltonian systems. *Trans. Amer. Math. Soc.* 311 (1989) 621-655.
- [WHL] W. Wang, X. Hu and Y. Long, Resonance identity, stability and multiplicity of closed characteristics on compact convex hypersurfaces. math.SG/0701608, to appear in Duke Math. J.
- [Wei1] A. Weinstein, Periodic orbits for convex Hamiltonian systems. *Ann. of Math.* 108 (1978) 507-518.