

SOME OBSERVATIONS ON LOCAL AND PROJECTIVE HYPERSURFACES

HAILONG DAO

Abstract. Let R be a hypersurface in an equicharacteristic or unramified regular local ring. For a pair of modules $(M; N)$ over R we study applications of rigidity of $\mathrm{Tor}^R(M; N)$, based on ideas by Huneke, Wiegand and Jorgensen. We then focus on the hypersurfaces with isolated singularity and even dimension, and show that modules over such rings behave very much like those over regular local rings. Connections and applications to projective hypersurfaces such as intersection dimension of subvarieties and cohomological criterion for splitting of vector bundles are discussed.

1. Introduction

The purpose of this note is to continue our investigation of the rigidity and decent intersection of modules over a local hypersurface R done in [Dal]. There we showed that the vanishing of ${}^R(M; N)$, a function introduced by Hochster in [Hol], implies the rigidity of Tor for $(M; N)$. We will apply this to give various new results on modules over hypersurfaces. We also give supporting evidences for the following Conjecture made in [Dal]:

Conjecture 1.1. Let R be an hypersurface with isolated singularity. Assume that $\dim R$ is even. Then ${}^R(M; N)$ always vanishes.

In Section 2 we review the basic notations and preliminary results. In Section 3 we use the procedure invented by Huneke and Wiegand in [HW2] to show that $\mathrm{Hom}_R(M; M)$ rarely has good depth. The use of ${}^R(M; N)$ allows us to simplify and strengthen some results in [HW2] (see 3.2 and 3.3, also 4.4 and 4.5).

Section 4 is concerned with two consequences of Conjecture 1.1:

Conjecture 1.2. Let R be an admissible hypersurface (meaning \hat{R} is a quotient of an unramified or equicharacteristic regular local ring by a nonzero element) with isolated singularity. Assume that $\dim R$ is even. Let $M; N$ be R -modules such that $\mathrm{Tor}_1^R(M; N) = 0$ for some i , then $\mathrm{Tor}_j^R(M; N) = 0$ for all $j \geq i$.

Conjecture 1.3. Let R be an admissible hypersurface with isolated singularity. Assume that $\dim R$ is even. Let $M; N$ be R -modules such that $\mathrm{ld}(M \otimes_R N) < 1$. Then $\dim M + \dim N = \dim R$.

We will prove 1.1 when M is free on the punctured spectrum of R and $N = M$ (see 4.1). We also prove 1.3 when R is a standard graded local complete intersection and $M; N$ are graded modules (see 4.11). Many applications follow, such as generalizations of results on $\mathrm{Hom}_R(M; M)$ over regular local rings by Auslander and Auslander-Goldman (Corollary 4.8 and 4.9). In this section, the connection with geometry of projective hypersurfaces (which goes both ways) will be exploited.

For example, 4.1.1 will be proved using ℓ -adic cohomology, and from 4.1 we obtain the following (see 4.7):

Corollary 1.4. Let k be a field and n an even integer ≥ 4 . Let $X \subset \mathbb{P}_k^n$ be a smooth hypersurface. Let E be a vector bundle on X . If $H^1(X; (E \otimes \mathcal{O}_X(i))^{\otimes \ell}) = 0$ for all $i \in \mathbb{Z}$, then E is a direct sum of line bundles of the form $\mathcal{O}_X(i)$.

In Section 5, we give some further applications of rigidity. The first is a simple characterization of maximal Cohen-Macaulay modules. The second is a connection between vanishing of $\text{Ext}_R^n(M; M)$ with $\text{pd}_R M$, following a recent result by Jorgensen ([Jo]).

The author would like to thank Ragnar-Olaf Buchweitz, David Jorgensen, Craig Huneke, Mircea Mustata, and Paul Roberts for many useful conversations and communications regarding this note.

2. Notation and preliminary results

Unless otherwise specified, all rings are Noetherian, commutative and local, and all modules are finitely generated. A local ring $(R; \mathfrak{m}; k)$ is a hypersurface if its completion \hat{R} has the form $\hat{R} = (f)$, where T is a regular local ring and f is in the maximal ideal of T . We say that R is admissible (as a hypersurface) if T is a power series ring over a field or a discrete valuation ring.

For a ring R and a non-negative integer i , we set $X^{\leq i}(R) := \{p \in \text{Spec}(R) \mid \dim(R_p) \leq i\}$. We denote by $Y(R)$ the set $X^{\dim(R)-1}(R)$, the punctured spectrum of R . We say that R satisfies the condition (R_i) if R_p is regular for any $p \in X^{\leq i}(R)$. We denote by $G(R)$ the Grothendieck group of finitely generated modules over R and by $\bar{G}(R) := G(R)/[R]$, the reduced Grothendieck group. Also, we let $\text{Sing}(R) := \{p \in \text{Spec}(R) \mid R_p \text{ is not regular}\}$ be the singular locus of R . For an abelian group G , we let $G_{\mathbb{Q}} = G \otimes_{\mathbb{Z}} \mathbb{Q}$.

Let $M^* := \text{Hom}(M; R)$ be the dual of M . The module M is called reflexive provided the natural map $M \rightarrow M^{**}$ is an isomorphism. The module M is called maximal Cohen-Macaulay (MCM) if $\text{depth}_R M = \dim R$.

For a non-negative integer n , M is said to satisfy (S_n) if:

$$\text{depth}_{R_p} M_p \geq \min\{n, \dim(R_p)\} \text{ for } p \in \text{Spec}(R)$$

(The depth of the 0 module is set to be 1).

A pair of R -modules $(M; N)$ is called rigid if for any integer $i \geq 0$, $\text{Tor}_i^R(M; N) = 0$ implies $\text{Tor}_j^R(M; N) = 0$ for all $j \leq i$. Moreover, M is rigid if for all N , the pair $(M; N)$ is rigid.

One defines the finite length index of the pair $(M; N)$ as:

$$f_R(M; N) := \min\{i \mid \text{Tor}_i^R(M; N) \neq 0\}$$

The function $f_R^R(M; N)$

Let $R = (f)$ be an admissible local hypersurface. The function $f_R^R(M; N)$ was introduced by Hochster ([Ho1]) for any pair of finitely generated modules $M; N$ such that $f_R(M; N) < 1$ as:

$$f_R^R(M; N) = 1 - \text{length}_R(\text{Tor}_{2e+2}^R(M; N)) - \text{length}_R(\text{Tor}_{2e+1}^R(M; N)):$$

where e is any integer $(d+2)=2$. It is well known (see [1]) that $\text{Tor}^R(M; N)$ is periodic of period 2 after $d+1$ spots, so this function is well-defined. The theta function satisfies the following properties. First, if $M \otimes_R N$ has finite length, then:

$$\text{Tor}^R(M; N) = \text{Tor}^T(M; N):$$

Secondly, $\text{Tor}^R(M; N)$ is biadditive on short exact sequence, assuming it is defined. Specifically, for any short exact sequence:

$$0 \rightarrow N_1 \rightarrow N_2 \rightarrow N_3 \rightarrow 0$$

and any module M such that $f_R(M; N_i) < 1$ for all $i = 1, 2, 3$, we have $\text{Tor}^R(M; N_2) = \text{Tor}^R(M; N_1) + \text{Tor}^R(M; N_3)$. Similarly, $\text{Tor}^R(M; N)$ is additive on the first variable.

In [2], we show that when $\text{Tor}^R(M; N)$ can be defined and vanishes, then $(M; N)$ is rigid:

Proposition 2.1. Let R be an admissible hypersurface and $M; N$ be R -modules such that $f_R(M; N) < 1$ (so that $\text{Tor}^R(M; N)$ can be defined). Assume $\text{Tor}^R(M; N) = 0$. Then $(M; N)$ is rigid.

The Pushforward

Let R be a Gorenstein ring and M a torsion-free (equivalent to (S_1)) R -module. Then there is an exact sequence:

$$0 \rightarrow M \rightarrow R \rightarrow M_1 \rightarrow 0$$

Here $M_1 = (M)$. This exact sequence is called the pushforward of M . The following proposition is taken from [HJW]. As mentioned in section 2, only a tiny modification is needed for the proof to work with our definition of (S_k) .

Proposition 2.2. ([HJW], 1.6) Let $R; M; M_1$ as above. Then for any $p \in \text{Spec}(R)$:

- (1) M_p is free if and only if $(M_1)_p$ is free.
- (2) If M_p is a maximal Cohen-Macaulay R_p -module, then so is $(M_1)_p$.
- (3) $\text{depth}_{R_p}(M_1)_p = \text{depth}_{R_p} M_p - 1$.
- (4) If M satisfies (S_k) , then M_1 satisfies (S_{k-1}) .

The depth formula

A result by Huneke and Wiegand showed that when all the high Tor modules vanish, the depths of the modules satisfy a remarkable equation:

Proposition 2.3. ([HW1], 2.5) Let R be a complete intersection. Let $M; N$ be non-zero finitely generated modules over R such that $\text{Tor}_i^R(M; N) = 0$ for all $i \geq 1$. Then:

$$\text{depth}(M) + \text{depth}(N) = \text{depth}(R) + \text{depth}(M \otimes_R N)$$

3. On depth of $\text{Hom}_R(M; M)$

We will call a module M such that M is locally free over the punctured spectrum of R a vector bundle over $\text{Y}(R)$ (or just vector bundle). We observe that for a vector bundle M and any R -module N , $\text{Tor}^R(M; N)$ is always defined. In this section we will show that for certain modules over admissible hypersurfaces, $\text{Hom}(M; M)$ and $M \otimes M$ rarely have good depth. We will follow the same procedure as in [HW2], but with two essential additions: we focus on vector bundles from the beginning

(even if the results do not involve them) and we will exploit the function ${}^R(\mathcal{M}; N)$ rather heavily. These will allow us to simplify and strengthen some results in [HW 2].

Proposition 3.1. Let R be an admissible hypersurface. Let \mathcal{M} be a vector bundle over $Y(R)$ such that $\text{depth}(\mathcal{M}) \geq 1$. Let N be an R -module such that ${}^R(\mathcal{M}; N) = 0$ and $\text{depth}(\mathcal{M} \otimes_R N) \geq 1$. Then $\text{Tor}_i^R(\mathcal{M}; N) = 0$ for all $i > 0$.

Proof. The assumptions ensure that \mathcal{M} is S_1 . Hence we have the pushforward of \mathcal{M} :

$$0 \rightarrow \mathcal{M} \rightarrow R \rightarrow \mathcal{M}_1 \rightarrow 0$$

We can tensor with N to get:

$$0 \rightarrow \text{Tor}_1^R(\mathcal{M}_1; N) \rightarrow \mathcal{M} \otimes_R N \rightarrow N \rightarrow \mathcal{M}_1 \otimes_R N \rightarrow 0$$

By 2.2, \mathcal{M}_1 is also a vector bundle. So $\text{length}(\text{Tor}_1^R(\mathcal{M}_1; N)) < 1$. Since it embeds into a module of positive depth, $\text{Tor}_1^R(\mathcal{M}_1; N)$ must be 0. Clearly ${}^R(\mathcal{M}_1; N)$ is defined and equal to:

$${}^R(R; N) - {}^R(\mathcal{M}; N) = 0$$

So by 2.1, $\text{Tor}_i^R(\mathcal{M}_1; N) = 0$ for all $i > 0$. Since $\mathcal{M} = \text{syzy}_1^R(\mathcal{M}_1)$, we are done.

The next result is an analogue of Theorem 2.4 in [HW 2]

Theorem 3.2. Let $(R; \mathfrak{m})$ be an admissible hypersurface of dimension d . Let r be an integer such that $0 \leq r < d$. Let \mathcal{M} be a vector bundle over $Y(R)$ such that $\text{depth}(\mathcal{M}) \geq r$. Let N be an R -module such that: N satisfies (S_r) , ${}^R(\mathcal{M}; N) = 0$, and $H_m^r(\mathcal{M} \otimes_R N) = 0$. Then $\text{depth}(\mathcal{M} \otimes_R N) \geq r + 1$ and if $r > 0$, we have $\text{Tor}_i^R(\mathcal{M}; N) = 0$ for all $i > 0$.

Proof. We will use induction on r . If $r = 0$ the conclusion is trivial. Now we assume $r > 0$. Then N satisfies (S_1) and we have the pushforward:

$$0 \rightarrow N \rightarrow R \rightarrow N_1 \rightarrow 0$$

Tensoring with \mathcal{M} we get:

$$0 \rightarrow \text{Tor}_1^R(N_1; \mathcal{M}) \rightarrow N \otimes_R \mathcal{M} \rightarrow \mathcal{M} \rightarrow N_1 \otimes_R \mathcal{M} \rightarrow 0$$

which we break into two short exact sequences:

$$0 \rightarrow \text{Tor}_1^R(N_1; \mathcal{M}) \rightarrow N \otimes_R \mathcal{M} \rightarrow C \rightarrow 0$$

and:

$$0 \rightarrow C \rightarrow \mathcal{M} \rightarrow N_1 \otimes_R \mathcal{M} \rightarrow 0$$

Since \mathcal{M} is a vector bundle, $T = \text{Tor}_1^R(N_1; \mathcal{M})$ has finite length. In particular $H_m^{r+1}(T) = 0$. By applying $H_m^0(\)$ to the first exact sequence we get $H_m^r(C) = 0$. Now applying $H_m^0(\)$ to the second exact sequence and using $H_m^{r-1}(\mathcal{M}) = 0$ (because $\text{depth}(\mathcal{M}) \geq r$) and $H_m^r(\mathcal{M} \otimes_R N) = 0$ we get $H_m^{r-1}(N_1 \otimes_R \mathcal{M}) = 0$. We need to check the other inductive assumptions for N_1 . Clearly, N_1 is (S_{r-1}) by 2.2. Also, ${}^R(\mathcal{M}; N_1)$ is defined and equal to:

$${}^R(\mathcal{M}; R) - {}^R(\mathcal{M}; N) = 0$$

So by induction we have $\text{depth}(\mathcal{M} \otimes_R N_1) \geq r$. Then by 3.1 $\text{Tor}_i^R(N_1; \mathcal{M}) = 0$ for all $i > 0$, so we have the first assertion and:

$$0 \rightarrow N \otimes_R \mathcal{M} \rightarrow \mathcal{M} \rightarrow N_1 \otimes_R \mathcal{M} \rightarrow 0$$

which implies $\text{depth}(M \otimes_R N) \geq r + 1$.

Corollary 3.3. Let R be an admissible hypersurface satisfying condition (R_2) . Let M be a reflexive R -module such that $M = 0$ in $\overline{G}(R)_Q$. If $\text{Hom}(M; M)$ satisfy (S_3) then M is free.

Proof. We use induction on $d = \dim R$. If $d = 2$ then R is regular, and M , being reflexive, must be free. Assume $d \geq 3$ and (by induction hypothesis, since all the conditions localize) that M is free on the punctured spectrum. In other words, M is a vector bundle. Consider the natural map $\alpha: M \otimes_R M \rightarrow \text{Hom}(M; M)$. Since M is a vector bundle, the kernel and cokernel of α both have finite length. By considering the long exact sequence of local cohomology from the two short exact sequences:

$$0 \rightarrow \ker(\alpha) \rightarrow M \otimes_R M \rightarrow \text{im}(\alpha) \rightarrow 0$$

$$0 \rightarrow \text{im}(\alpha) \rightarrow \text{Hom}(M; M) \rightarrow \text{coker}(\alpha) \rightarrow 0$$

and using $H_m^i(\text{Hom}(M; M)) = 0$ for $i < 3$ (because $\text{Hom}(M; M)$ is (S_3) and $d \geq 3$) we can deduce that $H_m^2(M \otimes_R M) = 0$. Now M is also a reflexive vector bundle, hence ${}^R(M; M) = 0$. So Theorem 3.2 applies to give $\text{Tor}_i^R(M; M) = 0$ for all $i > 0$. By depth formula we get:

$$\text{depth}(M) + \text{depth}(M) = d + 3$$

so by [Va], 3.3.16) we must have $\text{depth}(M) = \text{depth}(M) = d$. On the other hand, the vanishing of all $\text{Tor}_i^R(M; M)$ forces one of these modules to have finite projective dimension (see [HW 2], 1.9). Either way, M must be free (since M is MCM).

Example 3.4. This will be our main example throughout this note. Let $R = k[[x; y; u; v]] = (x; y; u; v)$ with $m = (x; y; u; v)$ and $M = (x; y)$. Then $M = (x; v)$ and $M \otimes_R M = (x^2; xy; xv; yv) = (x^2; xy; xv; xu) = m$. Using the long exact sequence of local cohomology for the sequence $0 \rightarrow m \rightarrow R \rightarrow k \rightarrow 0$ we get $H_m^1(M \otimes_R M) = H_m^1(m) = k$ and $H_m^2(M \otimes_R M) = H_m^2(m) = 0$. Clearly $\text{depth}(M \otimes_R M) = 1$.

Note that both M and M are MCM modules over R . Since R has isolated singularity, they are also vector bundles over $Y(R)$. It can be easily computed that $\text{Tor}_1^R(M; M) = 0$, $\text{Tor}_2^R(M; M) = k$ and ${}^R(M; M) = 1$.

Consider Proposition 3.1 and Theorem 3.2. Let $N = M$. Then the example shows that the condition ${}^R(M; N) = 0$ can not be dropped.

Consider Corollary 3.3. Since $\overline{G}(R)_Q = Q[M]$, the condition $M = 0$ in $\overline{G}(R)_Q$ can not be dropped either.

4. Isolated hypersurface singularities of even dimensions

In this section we will show some supporting evidences for Conjecture 1.1. They indicate that modules over isolated hypersurface singularities of even dimensions behave very similar to those over regular local rings. We first prove:

Theorem 4.1. Let R be a hypersurface with isolated singularity. Assume that $\dim R$ is even. Then for any vector bundle M on $Y(R)$, ${}^R(M; M) = 0$.

We need to review the concept of stable (co)homology (for more details, see [Bu] or [AB]). Let R be a Noetherian ring, a complete resolution of an R -module M is a complex T such that $H_n(T) = H^n(T) = 0$ for all $n \geq 2$ and $T_r = P_r$ for some projective resolution of M and some integer r . It is known that the modules

$H^i(\text{Hom}_R(T; N))$ and $H_i(T \otimes_R N)$ are independent of the resolution, and one calls them $\mathbb{E}xt_R^i(M; N)$ and $\mathbb{F}or_1^R(M; N)$, respectively.

We will need the standard and easy results below, reproved for the reader's convenience (some of these could be found in [AB], but we could not find a full reference):

Lemma 4.2. Let R be a local hypersurface. Let $M; N$ be R -modules such that M is maximal Cohen-Macaulay (MCM). Then we have the following isomorphism:

- (1) $\text{Tor}_1^R(M; N) = \mathbb{F}or_1^R(M; N)$ for all $i > 0$.
- (2) $\text{Ext}_R^i(M; N) = \mathbb{E}xt_R^i(M; N)$ for all $i > 0$.
- (3) $\mathbb{F}or_1^R(M; N) = \mathbb{F}or_{i+2}^R(M; N)$ for all i .
- (4) $\mathbb{E}xt_R^i(M; N) = \mathbb{E}xt_{R^{i+2}}^R(M; N)$ for all i .
- (5) $\mathbb{F}or_1^R(M; N) = \mathbb{E}xt_{R^{i+1}}^R(M; N)$ for all i .
- (6) $\text{Tor}_1^R(M; N) = \mathbb{E}xt_R^{i+1}(M; N)$ for all $i > 0$.

Proof. Let $F : \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow M \rightarrow 0$ be a free resolution of M . Since M is MCM we have $M^\perp = M$ and $\text{Ext}^n(M; R) = 0$ for all $n \geq 1$. So dualizing F we get:

$$F^\perp : 0 \rightarrow M^\perp \rightarrow F_0^\perp \rightarrow F_1^\perp \rightarrow \cdots$$

We can splice the two complexes together to get:

$$T : \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow F_0 \rightarrow F_1 \rightarrow \cdots$$

It is obvious that T is a complete resolution of M . That proves (1) and (2). Also, since M is MCM and R is a hypersurface, F (and therefore T) is periodic of period 2. Hence (3) and (4) follow. From the construction, $T[-1]$ is a complete resolution of M , and the canonical isomorphism $\text{Hom}_R(T; N) = (T \otimes_R N)$ gives (5). Finally, (6) follows from combining (1), (5) and (4).

We will also need the following result by Buchweitz:

Proposition 4.3. ([Bu], 10.3.3) Let R be a local hypersurface with isolated singularity such that $\dim R$ is even. Then for any two MCM R -modules $M; N$, even integer i and odd integer j :

$$l(\mathbb{E}xt_R^i(M; N)) = l(\mathbb{E}xt_R^j(N; M))$$

Now we can prove 4.1:

Proof. (of Theorem 4.1) First, we will prove the theorem for the case when M is MCM. In this situation, for any integer $i > 0$ we have:

$$\text{Tor}_1^R(M; M) = \mathbb{E}xt_R^{i+1}(M; M)$$

by (6) of 4.2. Hence 4.3 shows that $\text{Tor}_1^R(M; M) = 0$.

Now, assume M is a vector bundle. Let $K = \text{syzy}_1^R(M)$. We claim that $\text{Tor}_1^R(M; M) = \text{Tor}_1^R(K; K)$. Dualizing the short exact sequence: $0 \rightarrow K \rightarrow F \rightarrow M \rightarrow 0$ we get:

$$0 \rightarrow M^\perp \rightarrow F^\perp \rightarrow K^\perp \rightarrow \text{Ext}_R^1(M; R) \rightarrow 0$$

So $M^\perp = K^\perp + \text{Ext}_R^1(M; R)$ in $\overline{G}(R)$. But $\text{Ext}_R^1(M; R)$ has finite length because M is free on the punctured spectrum of R , so it is equal to 0 in $\overline{G}(R)_0$. So $\text{Tor}_1^R(M; M) = \text{Tor}_1^R(K; K)$. Together with $\text{Tor}_1^R(M; M) = \text{Tor}_1^R(K; K)$, the claim follows.

Repeating the claim we get ${}^R(M; M) = {}^R(K; K)$ when $K = \text{syzy}_n^R(M)$ for any $n > 0$. But for $n = 0$ K is an MCM R -module, so ${}^R(K; K) = 0$.

Corollary 4.4. Let R be an admissible hypersurface such that R has isolated singularity and $\dim R > 2$ and is even. Let M be a reflexive R -module. If $\text{Hom}(M; M)$ satisfy (S_3) then M is free.

Proof. Examining the proof of 3.3 we can see that the only thing we need is ${}^R(M; M) = 0$. Since by induction we can assume M is a vector bundle, the Corollary follows from Theorem 4.1.

Corollary 4.5. Let R be an admissible hypersurface such that R has isolated singularity and $\dim R \geq 2$ and is even. Let M be a vector bundle over $Y(R)$.

- (1) If $H_m^1(M \otimes_R M) = 0$ then $M = F \oplus T$, where T is free and T has finite length.
- (2) If $H_m^2(M \otimes_R M) = 0$ and $\dim R > 2$ then M is free.

Proof. (1) By (3.3) of [HW 2] we may assume that M is torsion free. By 3.2 we have $\text{Tor}_i^R(M; M) = 0$ for all $i > 0$. An identical argument to the last part of 3.3 shows that M is free.

(2) By (3.3) of [HW 2] we have $H_m^2(M \otimes_R M) = 0$. Again, using 3.2 we have M is free.

Example 4.6. Let $R; M$ as in Example 3.4. Then $H_m^2(M \otimes_R M) = 0$ but M is not free. Note that $\dim R = 3$.

Corollary 4.7. Let k be a field and n an even integer ≥ 4 . Let $X \subset \mathbb{P}_k^n$ be a smooth hypersurface. Let E be a vector bundle on X . If $H^1(X; (E \otimes E)(l)) = 0$ for all $l \geq 2$, then E is a direct sum of line bundles of the form $\mathcal{O}_X(i)$.

Proof. This is a straightforward consequence of standard connections between a projective variety and its affine cone (see section 5 in [HW 2]).

It is worth noting that two obvious consequences of Corollary 4.4 which are generalizations of well-known results by Auslander and Goldman ([AG], Theorem 4.4) and Auslander ([Au], Theorem 1.3) for modules over regular local rings:

Corollary 4.8. Let R be an admissible hypersurface such that R has isolated singularity and $\dim R > 2$ and is even. Let M be a reflexive R -module. If $\text{Hom}(M; M) = R^n$ for some n then M is free.

Corollary 4.9. Let R be an admissible hypersurface such that R has isolated singularity and $\dim R > 2$ and is even. Let M be an R -module satisfying (S_3) . If $\text{Hom}(M; M) = M^n$ for some n then M is free.

We now show that Conjecture 1.3 is true in the standard graded case:

Theorem 4.10. Let k be an algebraically closed field. Let $X \subset \mathbb{P}_k^N$ be a smooth complete intersection such that $\dim X = n$ is odd. Let U, V be subvarieties of X such that $U \cap V = \emptyset$. Then $\dim U + \dim V < \dim X$.

Proof. We are going to use l-adic cohomology (for basic properties and notations, we refer to [Mil]). Let l be a prime number such that $l \nmid \text{char}(k)$. There is a class map:

$$\text{cl}: \text{CH}^r(X) \rightarrow H^{2r}(X; \mathbb{Q}_l(r))$$

This map gives a graded ring homomorphism $CH^*(X) \rightarrow H^{2*}(X; \mathbb{Q}_1(r))$ (with the intersection product on the left hand side and the cup product on the right hand side, see [M1], chapter 23). Let $a = \dim U$ and $b = \dim V$, and we may assume $a \leq b$. Suppose $a + b = \dim X = n$ (if $a + b > n$, we can always choose some subvariety of smaller dimension in U or V such that equality occurs). Then $2a \leq n$, but n is odd, so $2a > n$. Let $h \in CH^1(X)$ represents the hyperplane section. By Weak Lefschetz Theorem and the fact that $2(n - a) < n$, we have

$$H^{2(n-a)}(X; \mathbb{Q}_1(n-a)) = H^{2(n-a)}(\mathbb{P}_k^N; \mathbb{Q}_1(n-a))$$

the latter is generated by power of the class of the hyperplane section. Thus $cl(U) = cl(h)^{n-a} \in H^{2(n-a)}(X; \mathbb{Q}_1(n-a))$. We then have :

$$cl(U \cdot V) = cl(U) \cdot cl(V) = cl(h)^{n-a} \cdot cl(V) = cl(h)^n \cdot V$$

The last term is equal to the degree of $h^n \cdot V \in CH^n(X)$, so it is nonzero. But the first term has to be 0 by assumption. This contradiction proves the Theorem.

Theorem 4.11. Let k be a field. Let $A = k[x_0, \dots, x_N] / (I)$ be a homogeneous complete intersection. Let $(R; \mathfrak{m}; k)$ be the local ring at the origin of A . Suppose that R has isolated singularity and $\dim R$ is even. Let M, N be graded R -modules such that $l(M \otimes_R N) < 1$. Then $\dim M + \dim N = \dim R$.

Proof. Without affecting the statement we may assume k is algebraically closed. Since the minimal primes over a graded module are graded, we may replace M and N with R/P and R/Q , where P, Q are homogeneous primes in R . Now let $X = \text{Proj}(A)$, $U = \text{Proj}(A/P)$, $V = \text{Proj}(A/Q)$ and applying the previous Theorem.

5. Some other applications

We note a simple characterization of maximal Cohen-Macaulay modules, due to the fact that over an admissible hypersurface, a module of finite length is rigid (see 4.10 in [D1]):

Proposition 5.1. Let R be a admissible hypersurface and M an R -module. The following are equivalent:

- (1) There is a finite length module N such that $\text{Tor}_1^R(M; N) = 0$.
- (2) There is a finite length module N such that $\text{Ext}_R^1(M; N) = 0$.
- (3) M is a maximal Cohen-Macaulay R -module.

Proof. For equivalence of (1) and (2): we may assume R is complete. Then by Matlis duality $\text{Tor}_1^R(M; N) \cong \text{Ext}_R^1(M; N^\vee)$, so (1) and (2) are equivalent.

Assume (1). Then since N is rigid, we have $\text{Tor}_i^R(M; N) = 0$ for all $i \geq 1$. The depth formula shows:

$$\text{depth}(M) + \text{depth}(N) = \text{depth}(R) + \text{depth}(M \otimes_R N)$$

hence $\text{depth}(M) = \text{depth}(R)$ and (3) follows.

Finally, assume (3). Let $n = \dim R$, we can choose a regular sequence x_1, \dots, x_n on both M and R (by induction on n). Now, just take $N = R/(x_1, \dots, x_n)$. Then N is finite length and $\text{Tor}_1^R(M; N) = 0$.

It would be very interesting to know whether the previous Proposition is true for all local complete intersections. It is not hard to see that this question has intimate connection to the rigidity of modules of finite length over such rings:

Lemma 5.2. Let \mathcal{C} be the category of all local complete intersections. Considering the following properties:

- (1) For any $R \in \mathcal{C}$, an R -module of finite length is rigid.
- (2) For any $R \in \mathcal{C}$ and any R -module M , if there is a finite length R -module N such that $\text{Tor}_1^R(M; N) = 0$ then M is maximal Cohen-Macaulay.
- (3) For any $R \in \mathcal{C}$ such that $\dim R > 0$ and any R -module M , if there is a finite length R -module N such that $\text{Tor}_1^R(M; N) = 0$ then $\text{depth}(M) > 0$.
- (4) For any $R \in \mathcal{C}$, an R -module of finite length and finite projective dimension is rigid.

We have (1) \Rightarrow (2), (3) \Rightarrow (4).

Proof. (1) \Rightarrow (2) is the proof of 5.1.

(2) \Rightarrow (3). (2) clearly implies (3). Suppose (3) holds. Then since $\text{depth}(M) \leq \text{depth}(R) > 0$ we can find $x \in \text{Ann}_R N$ such that x is regular on both R and M . Then $\text{Tor}_1^{R=(x)}(M=xM; N) = \text{Tor}_1^R(M; N) = 0$. So we can replace $R; M$ by $R=(x); M=xM$ and repeat the process until $\dim R = 1$, at which point M is MCM by (3).

(2) \Rightarrow (4). Suppose $M; N$ are nonzero R -modules such that $\ell(N) < 1$, $\text{pd}_R N < 1$ and $\text{Tor}_i^R(M; N) = 0$ for some $i > 0$. Then let $M^0 = \text{syz}_{i-1}^R M$ and we get $\text{Tor}_1^R(M^0; N) = 0$. By (2) M^0 is MCM. Let q be the largest integer such that $\text{Tor}_q^R(M^0; N) \neq 0$. Then by Lemma 2.2 in [HW 2] we have:

$$\text{depth}(M^0) + \text{depth}(N) = \text{depth}(R) + \text{depth}(\text{Tor}_q^R(M^0; N)) - q$$

Since $\text{depth}(M^0) = \text{depth}(R)$ and $\text{depth}(N) = \text{depth}(\text{Tor}_q^R(M^0; N)) = 0$ we must have $q = 0$.

Remark. In [JS] a module M is constructed over a 0-dimensional Gorenstein ring such that M is not rigid. So property (1) fails for Gorenstein rings.

Finally, we want to use our approach to rigidity to prove a connection between vanishing of Ext and projective dimension, following a recent result by Jorgensen. In [Jo], the following was proved:

Theorem 5.3. ([Jo], Theorem 1.1) Let R be a local ring and M an R -module such that M is rigid. Then $\text{Ext}_R^n(M; M) = 0$ if and only if $\text{pd}_R(M) < n$.

We will modify Jorgensen's proof to show:

Proposition 5.4. Let R be an admissible hypersurface and M be an R -module such that $M = 0$ in $\overline{G}(R)_0$. Then $\text{Ext}_R^n(M; M) = 0$ if and only if $\text{pd}_R(M) < n$.

Remark. Over a hypersurfaces, the classes of rigid modules and of modules which are 0 in $\overline{G}(R)_0$ are incomparable (see [Dal]).

Proof. We first need to review some notation in [Jo]. Let N be an R -module and let

$$F = \cdots \rightarrow F_2 \rightarrow F_1 \rightarrow F_0 \rightarrow N \rightarrow 0$$

be a minimal free resolution of N . Then one can define for $i \geq 0$:

$$D^i(N) = \text{coker}(F_i \rightarrow F_{i+1})$$

Examining the proof of Lemma 2.2 and Theorem 1.1 in [Jo] shows that to prove $\text{Ext}_R^n(M; M) = 0$ implies $\text{pd}_R(M) < n$ one only needs that the pair $(D^n(\text{syz}_{n-1}^R M); M)$ is rigid. Let $L = D^n(\text{syz}_{n-1}^R M)$. We will show that $\text{Ext}_R^R(L; M) = 0$ using induction

on $\dim R$. If $\dim R = 0$ then $\overline{G}^R(L; M)$ is always defined and equals 0, since all modules have finite length and $\overline{G}^R(R)_Q = 0$. Take any prime $p \in Y(R)$. By induction $(\text{syz}_{n-1}^R M)_p$ is free, so L_p is a free R_p module. Thus $\overline{G}^R(L; M)$ is always defined and since $M = 0$ in $\overline{G}^R(R)_Q$ we must have $\overline{G}^R(L; M) = 0$. So $(L; M)$ is rigid and we are done.

Example 5.5. Let $R = k[[x; y]]/(xy)$ and $M = R/(x)$. Then $\text{Ext}^{2i+1}(M; M) = 0$ for all $i > 0$ but $\text{pd}_R M = 1$. Note that $\overline{G}^R(R) = \mathbb{Z}[M]$.

References

- [Au] M. Auslander, On the Purity of the Branch Locus, *Amer. J. Math.* 84 (1962), 116-125.
- [AB] L.L. Avramov, R.-O. Buchweitz, Support varieties and cohomology over complete intersections, *Invent. Math.* 142 (2000), 285-318.
- [AG] M. Auslander, O. Goldman, Maximal orders, *Trans. Amer. Math. Soc.* 97 (1960), 1-24.
- [BH] W. Bruns, J. Herzog, Cohen-Macaulay rings, Cambridge Univ. Press, Cambridge (1996).
- [Bu] R.-O. Buchweitz, Maximal Cohen-Macaulay modules and Tate cohomology over Gorenstein rings, Preprint, Univ. Hannover, (1986).
- [Dal] H. Dao, Decency and rigidity over hypersurfaces, *arXiv math.AC/0611568*, preprint.
- [Ei] D. Eisenbud, Homological algebra on a complete intersection, with an application to group representations, *Trans. Amer. Math. Soc.* 260 (1980), 35-64.
- [Fu] W. Fulton, Intersection Theory, Springer-Verlag, Berlin (1998).
- [Gr] A. Grothendieck, Elements de geometrie algebrique, Chapage IV Publ. Math. I.H.E.S. 24 (1965).
- [Ha] R. Hartshorne, Algebraic Geometry, Graduate Text in Mathematics, Springer-Verlag, New York, (1977).
- [Hol] M. Hochster, The dimension of an intersection in an ambient hypersurface, Proceedings of the First Midwest Algebraic Geometry Seminar (Chicago Circle, 1980), Lecture Notes in Mathematics 862, Springer-Verlag, 1981, 93-106.
- [HW 1] C. Huneke, R. Wiegand, Tensor products of modules and the rigidity of Tor, *Math. Ann.* 299 (1994), 449-476.
- [HW 2] C. Huneke, R. Wiegand, Tensor products of modules, rigidity and local cohomology, *Math. Scan.* 81 (1997), 161-183.
- [JW] C. Huneke, R. Wiegand, D. Jorgensen, Vanishing theorems for complete intersections, *J. Algebra* 238 (2001), 684-702.
- [Jo] D. Jorgensen, A note on the obstruction group to lifting module regular deformation, preprint.
- [JS] D. Jorgensen, L. Sega, Nonvanishing cohomology and classes of Gorenstein rings, *Adv. Math.* 188 (2004), 470-490.
- [Mi] J.S. Milne, Lectures on etale cohomology, available at www.jmilne.org/math.
- [PS] C. Peskine, L. Szpiro, Dimension projective et cohomologie locale. Applications a la demonstration de conjectures de M. Auslander, H. Bass et A. Grothendieck, *Inst. Hautes Etudes Sci. Publ. Math.* 42 (1973), 47-119.
- [Ro] P. Roberts, Multiplicities and Chern classes in Local Algebra, Cambridge Univ. Press, Cambridge (1998).
- [Va] W. Vasconcelos, Arithmetic of Blowup Algebra, London Math. Soc. Lecture Notes 195 (1994).

Department of Mathematics, University of Utah, 155 South 1400 East, Salt Lake City, UT 84112-0090, USA
 E-mail address: hdao@math.utah.edu