

Metric invariants of tropical conics and factorization of degree–two homogeneous tropical polynomials in three variables

M. Ansola, M.J. de la Puente^{*†}

March 23, 2019

Abstract

To every tropical conic we associate a certain non–negative symmetric 3×3 real matrix. The non–diagonal entries of this matrix are metric invariants of the given tropical conic, up to translation. There exist proper and improper tropical conics, pairs of lines being a particular sort of the latter. We characterize those tropical conics whose associated matrix is tropically singular. Again, pairs of lines are among the latter.

A tropical conic is defined by an arbitrary degree–two homogeneous tropical polynomial in three variables. The conic is a tree of a very particular kind. Given such a tree, we give a procedure to obtain a defining tropical polynomial.

Finally, we give criteria to decide whether a tropical degree–two homogeneous polynomial in three variables is reducible and, if so, we find a factorization.

2000 Math. Subj. Class.: 05C05; 12K99

Key words: Tropical conics, metric invariants, factorization of tropical polynomials, tropically singular matrix.

1 Introduction

In recent years, there has been a growing interest in *projective tropical geometry*, [4, 5, 8, 11, 13, 14, 15, 16, 17, 18, 21, 22, 23]. Some authors have searched for tropical versions of some classical theorems of plane projective geometry, see [19, 20]. Many of these old theorems involve conics and this has motivated us to make a thorough study of tropical conics.

This paper begins with the study of *tropical conics* from an euclidian point of view. It originated as an attempt to explain in full detail and give proofs for all statements given in example 3.4 in [16]. We introduce a triple of non–negative real numbers, (f_2^+, f_4^+, f_6^+) , which completely determines a tropical conic, up to translation. Given

^{*}Departamento de Algebra, Facultad de Matemáticas, Universidad Complutense, 28040–Madrid, Spain

[†]Partially supported by MTM 2005–02865 and UCM 910444.

a tropical degree–two homogeneous polynomial P in three variables, one can easily compute these numbers, and then draw the tropical conic \mathcal{C} associated to P (see procedure in p. 9). In the literature, one finds that the regular subdivision of the Newton polygon of P provides \mathcal{C} as a graph, by duality. But this method does not determine \mathcal{C} completely, since nothing is said about the precise coordinates of the vertices of \mathcal{C} nor the lengths of its edges.

Tropical conics can be classified, according to the values attained by (f_2^+, f_4^+, f_6^+) . In particular, we characterize degenerated (also called improper) tropical conics (corollary 4). We also characterize pairs of lines in terms of these invariants (corollary 7). Pairs of tropical lines are degenerated tropical conics, but the converse is not true.

The values (f_2^+, f_4^+, f_6^+) can be arranged into a 3×3 symmetric real matrix with non–negative entries. We characterize tropical conics whose associated matrix is tropically singular (lemma 4). These are pairs of tropical lines and one–point central conics.

On the other hand, it is known that every balanced weighted tree \mathcal{C}_d of a certain sort is a tropical curve of degree $d > 0$, see [12]. For $d = 2$, we give a procedure to find a defining polynomial P for such a graph \mathcal{C}_2 (see p. 14).

Having studied tropical conics thoroughly, we address the question of irreducibility of degree–two homogeneous tropical polynomials in three variables. We give a procedure to determine whether such a polynomial is reducible, and we explain how to obtain a factorization, if there exists one (see procedure p. 13).

Some of the results in this paper have already appeared in [1], while other are new. The idea of decomposing a matrix A as the sum of two matrices, $\text{trans}(A)$ and $\text{shp}(A)$ comes, somehow, from [9] and is developed in section 3. The factorization problem for tropical polynomials in one variable and the fundamental theorem of algebra have been investigated in [10, 6].

We would like to thank B. Bertrand, A. Díaz–Cano and P.D. González for some helpful discussions about the material presented here.

2 Notations

Tropical conics are degree–two plane tropical curves and they can be defined in the affine setting or in the projective one. *Tropical geometry* arises when one works over \mathbb{T} , the *tropical semi–field*. By definition, \mathbb{T} is the set $\mathbb{R} \cup \{-\infty\}$ endowed with two operations: \max and $+$. *Tropical addition* is \max and $+$ is taken as *tropical multiplication*. They are denoted \oplus and \odot , respectively. The neutral element for tropical addition is $-\infty$ and zero is the neutral element for tropical multiplication. It is noticeable that $a \oplus a = a$, for $a \in \mathbb{T}$, that is, tropical addition is *idempotent*. However, there does not exist an inverse element, with respect to \oplus , for $a \in \mathbb{T}$. This is all that \mathbb{T} lacks in order to be a field.

Here, \mathbb{R}^+ will denote the set of non–negative real numbers. For $a \in \mathbb{T}$, we will set $a^+ = \max\{a, 0\} = a \oplus 0$, the *positive part of a* (an exception to this notation is d_i^+ in p. 10). For a tropical polynomial P , the polynomial obtained by replacing every coefficient a of P by a^+ will be denoted P^+ . For a matrix A , the matrix obtained by replacing every entry a of A by a^+ will be denoted A^+ .

The tropical affine 2-space is \mathbb{T}^2 , with addition and multiplication are defined coordinatewise. It will be denoted \mathbb{TA}^2 . Here we can define *translations* and *homotheties* in the standard way. A translation depends on a vector $\tau \in \mathbb{R}^2$ and a homothety depends on a point in \mathbb{R}^2 , called *center*, and a *scaling factor* $\rho \in \mathbb{R}$, which we require to be positive.

In the space $\mathbb{T}^3 \setminus \{(-\infty, -\infty, -\infty)\}$ we define an equivalence relation \sim by letting $(b_1, b_2, b_3) \sim (c_1, c_2, c_3)$ if there exist $\lambda \in \mathbb{R}$ such that $(b_1 + \lambda, b_2 + \lambda, b_3 + \lambda) = (c_1, c_2, c_3)$. The equivalence class of (b_1, b_2, b_3) is denoted $[b_1, b_2, b_3]$ and its elements are obtained by adding multiples of the vector $(1, 1, 1)$ to the point (b_1, b_2, b_3) . The tropical projective 2-space, \mathbb{TP}^2 , is the set of such equivalence classes.

Points in \mathbb{TA}^2 (resp. \mathbb{TP}^2) without infinite coordinates will be called *interior points*. The rest of the points will be called *boundary points*. The *boundary* of \mathbb{TA}^2 (resp. \mathbb{TP}^2) is the union of its boundary points. Note that, at least, one of the coordinates of any point in \mathbb{TP}^2 must be finite. The plane \mathbb{TP}^2 is covered by three copies of \mathbb{TA}^2 as follows. There exist injective maps

$$\begin{aligned} j_3 : \mathbb{TA}^2 &\rightarrow \mathbb{TP}^2, & (x, y) &\mapsto [x, y, 0], & j_2 : \mathbb{TA}^2 &\rightarrow \mathbb{TP}^2, & (x, z) &\mapsto [x, 0, z], \\ j_1 : \mathbb{TA}^2 &\rightarrow \mathbb{TP}^2, & (y, z) &\mapsto [0, y, z] \end{aligned}$$

and $\mathbb{TP}^2 = \text{im } j_3 \cup \text{im } j_2 \cup \text{im } j_1$. The complementary set of $\text{im } j_3$ is the set

$$\mathbb{TP}^2 \setminus \text{im } j_3 = \{[x, y, -\infty] : x, y \in \mathbb{R} \cup \{-\infty\}\}.$$

Similarly for j_2, j_1 . It is easy to check that the set of interior points of \mathbb{TP}^2 is equal to the intersection $\text{im } j_3 \cap \text{im } j_2 \cap \text{im } j_1$.

We will use X, Y, Z as variables. Any exchange of the variables X, Y, Z provides a *change of projective tropical coordinates*. There might exist other types of changes of projective tropical coordinates but we will not consider them here.

Let us look at the map j_3 more closely. By j_3^{-1} , the directions given by the coordinate axes X, Y, Z in \mathbb{TP}^2 are transformed into the directions given by the coordinate axes X, Y in \mathbb{TA}^2 and the usual line $X = Y$. This is so because $j_3(x, x) = [x, x, 0] = [0, 0, -x]$, for $x \in \mathbb{T}$. Moreover, the negative Z half-axis in \mathbb{TP}^2 corresponds to the NE direction in \mathbb{TA}^2 . We consider the euclidian metric in \mathbb{TA}^2 . Notice that the composite maps $j_l^{-1} \circ j_k$ are NOT isometries, for $k, l \in \{1, 2, 3\}, k \neq l$.

The projective tropical coordinates of a point in \mathbb{TP}^2 are not unique. In order to avoid this inconvenience, we choose a *normalization*. For instance, making the last coordinate equal to zero is a normalization. We call it the $Z = 0$ *normalization* and say that *we work in $Z = 0$* . For any given normalization, there exist a few points in \mathbb{TP}^2 which do not admit it. This is the case for $[a, b, -\infty]$, for any $a, b \in \mathbb{T}$ and the $Z = 0$ normalization. To consider the $Z = 0$ normalization means to pass from the projective setting to the affine one, via j_3 . Other possible normalizations are $Y = 0$, or $X = 0$, or X, Y, Z all non-negative and, at least, one equal to zero, etc.

3 Matrices

All matrices considered in this paper are symmetric; therefore, we only write their lower triangular parts. We will arrange subscripts of matrices entries in an unusual way, taken from [16]. All subscripts will be taken modulo 6. To any symmetric matrix

$$A = \begin{pmatrix} a_1 & & & & & \\ a_2 & a_3 & & & & \\ a_6 & a_4 & a_5 & & & \end{pmatrix} \quad (1)$$

with $a_1, a_3, a_5 \in \mathbb{R}$ and $a_2, a_4, a_6 \in \mathbb{T}$ we associate the following values in \mathbb{T} :

$$f_2 = 2a_2 - a_1 - a_3, \quad f_4 = 2a_4 - a_3 - a_5, \quad f_6 = 2a_6 - a_5 - a_1. \quad (2)$$

Note that $a_2, a_4, a_6 \in \mathbb{R}$ if and only if $f_2, f_4, f_6 \in \mathbb{R}$. If this is the case, we can define three more values in \mathbb{R} :

$$d_1 = \frac{1}{2}(f_6 + f_2 - f_4), \quad d_3 = \frac{1}{2}(f_2 + f_4 - f_6), \quad d_5 = \frac{1}{2}(f_4 + f_6 - f_2), \quad (3)$$

so that

$$f_2 = d_1 + d_3, \quad f_4 = d_3 + d_5, \quad f_6 = d_5 + d_1, \quad (4)$$

$$d_1 = a_6 + a_2 - a_4 - a_1, \quad d_3 = a_2 + a_4 - a_6 - a_3, \quad d_5 = a_4 + a_6 - a_2 - a_5. \quad (5)$$

Note that the values $f_2, f_4, f_6, d_1, d_3, d_5$ (if defined) do not change if A is replaced by $A = \alpha + U$, for any $\alpha \in \mathbb{R}$, where U denotes the 3×3 matrix all whose entries are one. Now, let us prove some easy properties of such numbers.

Lemma 1. *If $f_2, f_4, f_6 \in \mathbb{R}$ and there exists $i \in \{1, 3, 5\}$ such that $d_i < 0$ and $f_{i-1} > 0, f_{i+1} > 0$. Then $d_{i-2} > 0, d_{i+2} > 0$ and $f_{i+3} > 0$.*

Proof. We just show one case, the others being similar. Suppose $d_1 < 0, f_6 > 0$ and $f_2 > 0$. Then $0 < d_1 + d_3$ and $0 < d_1 + d_5$, whence $0 < -d_1 < d_3$ and $0 < -d_1 < d_5$. Adding up, we obtain $0 < d_3 + d_5 = f_4$. \square

Note that the same assertion and proof are valid if all the inequalities are relaxed, i.e., if all occurrences of $<$ are replaced by \leq .

Lemma 2. *If f_2, f_4, f_6 are all non-negative, then d_i is negative for, at most, one $i \in \{1, 3, 5\}$. And if f_2, f_4, f_6 are all positive, then d_i is non-positive for, at most, one $i \in \{1, 3, 5\}$.* \square

Next, we decompose a matrix A into the sum of two symmetric matrices, $A = \text{trans}(A) + \text{shp}(A)$, where

$$\text{trans}(A) = \frac{1}{2} \begin{pmatrix} 2a_1 & & & & & \\ a_1 + a_3 & 2a_3 & & & & \\ a_1 + a_5 & a_3 + a_5 & 2a_5 & & & \end{pmatrix}, \quad \text{shp}(A) = \frac{1}{2} \begin{pmatrix} 0 & & & & & \\ f_2 & 0 & & & & \\ f_6 & f_4 & 0 & & & \end{pmatrix},$$

(see [9]). The entries of $\text{trans}(A)$, the *translation matrix of A*, are real, while the entries of $\text{shp}(A)$, the *shape matrix of A*, belong to \mathbb{T} , in general. The meaning of this names will become clear later. To A we also associate the following points in \mathbb{TP}^2 :

$$h(A) = [-a_1, -a_2, -a_6], \quad v(A) = [-a_2, -a_3, -a_4], \quad i(A) = [-a_6, -a_4, -a_5].$$

If not all a_k equal $-\infty$, for $k \in \{2, 4, 6\}$, then we can consider the point $c(A) = [a_4, a_6, a_2]$. We will simply write h, v, i, c if the matrix A is understood.

In \mathbb{R}^2 , the segment joining points a, b will be denoted $\text{conv}(ab)$.

Lemma 3. *Suppose that $a_2, a_4, a_6 \in \mathbb{R}$. In $Z = 0$, the length of the segment $\text{conv}(hc)$ is $|d_1|$ and it is parallel to the X axis. The length of the segment $\text{conv}(vc)$ is $|d_3|$ and it is parallel to the Y axis. The length of the segment $\text{conv}(ic)$ is $|d_5|\sqrt{2}$ and it is parallel to the line $X = Y$.*

Proof. In $Z = 0$, the coordinates of the points h, v, i, c are $h = (a_6 - a_1, a_6 - a_2)$, $v = (a_4 - a_2, a_4 - a_3)$, $i = (a_5 - a_6, a_5 - a_4)$, $c = (a_4 - a_2, a_6 - a_2)$ and now all the assertions easily follow from the equalities (5). \square

A shorter way to put the preceding lemma is

$$c = v + [0, -d_3, 0] = h + [-d_1, 0, 0] = i + [-d_5, -d_5, 0], \quad (6)$$

when c and d_i are defined, $i \in \{1, 3, 5\}$. We deduce that, in $Z = 0$, the points h, v, c determine a *usual right triangle*; the signs of d_1 and d_3 determine this triangle further. Similarly in $Z = 0$, the vertices h, i, c determine a usual triangle having at c an angle of $\frac{3\pi}{4}$ (resp. $\frac{\pi}{4}$) if $d_1 d_5 > 0$ (resp. $d_1 d_5 < 0$). These triangles become right triangles in $Y = 0$. Also the vertices v, i, c determine a usual triangle in $Z = 0$, having at c an angle of $\frac{3\pi}{4}$ (resp. $\frac{\pi}{4}$) if $d_3 d_5 > 0$ (resp. $d_3 d_5 < 0$). These triangles become right triangles in $X = 0$.

The *tropical determinant* of a 3×3 matrix is defined as follows:

$$\left| \begin{array}{ccc} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{array} \right|_t = \max_{\sigma \in S_3} \{a_{1\sigma(1)} + a_{2\sigma(2)} + a_{3\sigma(3)}\}$$

where S_3 denotes the permutation group in 3 symbols. This matrix is said *tropically singular* if the *maximum in the tropical determinant is attained, at least, twice*. If A equals (1) then

$$|\text{trans}(A)|_t = a_1 + a_3 + a_5, \quad |\text{shp}(A)|_t = \max\{0, s, s, f_2, f_4, f_6\}$$

$$|A|_t = |\text{trans}(A)|_t + |\text{shp}(A)|_t,$$

where $s = \frac{f_2 + f_4 + f_6}{2} = -a_1 + a_2 - a_3 + a_4 - a_5 + a_6$ and $s = d_1 + d_3 + d_5$, if d_1, d_3, d_5 are defined. The matrix $\text{trans}(A)$ is tropically singular, hence A is *tropically singular if and only if shp(A) is*.

Lemma 4. *Suppose that f_2, f_4, f_6 are all non-negative. The following are equivalent:*

1. $\text{shp}(A)$ is tropically singular,
2. the maximum of f_2, f_4, f_6 is no greater than the sum of the other two,
3. $d_1 \geq 0, d_3 \geq 0, d_5 \geq 0$.

Proof. Without loss of generality, we may suppose that $f_6 = \max\{f_2, f_4, f_6\}$. Then $f_6 \leq f_2 + f_4$ if and only if $f_6 \leq s$ if and only if $s = \max\{0, s, s, f_2, f_4, f_6\}$ if and only if the maximum $\{0, s, s, f_2, f_4, f_6\}$ is attained, at least, twice. Equivalence with $d_1 \geq 0, d_3 \geq 0, d_5 \geq 0$ follows from (3). \square

4 Tropical conics

Let $P(X, Y, Z)$

$$\begin{aligned} &= a_1 \odot X^{\odot 2} \oplus a_3 \odot Y^{\odot 2} \oplus a_5 \odot Z^{\odot 2} \oplus a_2 \odot X \odot Y \oplus a_4 \odot Y \odot Z \oplus a_6 \odot X \odot Z \\ &= \max\{a_1 + 2X, a_3 + 2Y, a_5 + 2Z, a_2 + X + Y, a_4 + Y + Z, a_6 + X + Z\} \end{aligned}$$

be a homogeneous tropical polynomial of degree two. This means that $a_1, a_3, a_5 \in \mathbb{R}$ and $a_2, a_4, a_6 \in \mathbb{T}$. The *tropical conic* given by P is, by definition, the set of points in \mathbb{TP}^2 where the *maximum is attained, at least, twice*. A simple computer program (in MAPLE, for instance) may be used in order to sketch this conic, say in $Z = 0$. Soon we will see that we can dispense with the computer!

It is well-known that \mathcal{C} is a tree, (see section 6). It has six pendant edges, counted with multiplicity. This multiplicity is one or two. If we draw \mathcal{C} in $Z = 0$, then \mathcal{C} has two pendant edges in the W direction, two in the S direction and two in the NE direction, all counted with multiplicity. Moreover, \mathcal{C} has, at most, four non-pendant vertices. The aim of this section is to give a *metric description of this tree*. Without loss of generality, we may work in $Z = 0$.

Let P be given. First, *we make the extra assumption that $a_2, a_4, a_6 \in \mathbb{R}$* , so that lemma 3 holds true. To P we associate a symmetric matrix A as in (1) —here we must recall that tropical addition is idempotent. From A we obtain the points h, v, i, c , the matrix $\text{shp}(A)$ and the real values d_1, d_3, d_5 .

- The simplest case occurs when $d_i > 0$ for all $i \in \{1, 3, 5\}$. Then, the points h, v, i, c are precisely the non-pendant vertices of the conic \mathcal{C} and, from the equalities (6), we conclude that in $Z = 0$ they are joined by edges of lengths $d_1, d_3, d_5\sqrt{2}$. To finish up with the sketch of \mathcal{C} , we must add six pendant edges, as explained above; see figure at the end (in a separate file) line 1, column 3. We will say that this is a **one-point central tropical conic**; see p. 14 for an explanation of this and coming names. If we work in $X = 0$, (resp. $Y = 0$) we obtain other representations of such a conic; see figure at the end (in a separate file) line 1, column 1 (resp. column 2).

Notice that if \mathcal{C} is a one-point central tropical conic, then its vertices h, v, i form a proper tropical triangle, as defined in [2]. This is so because the inequalities $d_i > 0$ for $i \in \{1, 3, 5\}$ yield the six inequalities on the coordinates of the vertices of such a triangle.

- The next case is given by the following condition: there exists $i \in \{1, 3, 5\}$ with $d_i < 0$ and $f_{i-1} > 0$, $f_{i+1} > 0$. Here we will talk about **two–point central tropical conics**. Three cases arise here, depending on the value of i :

type 1: $d_1 < 0$, $f_6 > 0$, $f_2 > 0$.

type 3: $d_3 < 0$, $f_2 > 0$, $f_4 > 0$.

type 5: $d_5 < 0$, $f_4 > 0$, $f_6 > 0$.

Associated to A , we have the points h, v, i, c and equalities (6) hold. In a tropical conic \mathcal{C} of **type 5**, we find that h, v are vertices, but i, c are NOT. Yet, the conic \mathcal{C} has two other non–pendant vertices, denoted ch, cv and determined as follows:

$$ch = h + [-f_6, 0, 0], \quad cv = v + [0, -f_4, 0].$$

The points i, cv, c, ch are the vertices of a usual square of side length equal to $-d_5$. From h, v, ch, cv hang six pendant edges of \mathcal{C} and such a tropical conic is represented in figure at the end (in a separate file) line 2, column 3.

In a tropical conic of **type 1**, we find the vertices v, i (from the matrix A) and two more non–pendant vertices, denoted by ci, cv and determined as follows:

$$ci = i + [f_6, f_6, 0], \quad cv = v + [0, -f_2, 0].$$

A tropical conic of type 1 is represented in figure at the end (in a separate file) line 2, column 1.

In a tropical **type 3** conic, we see the vertices h, i associated to the matrix A , but v, c do NOT appear. Two more non–pendant vertices exist in such a conic, denoted by ci, ch , and given by the following equalities:

$$ci = i + [f_4, f_4, 0], \quad ch = h + [-f_2, 0, 0].$$

A tropical conic of type 3 is represented in figure at the end (in a separate file) line 2, column 2.

Note that the exchange of the variables X (resp. Y) and Z takes a tropical conic of type 1 (resp. type 3) into a tropical conic of type 5. Therefore, in order to study all two–point central conics, it is enough to study type 5 conics. Note also that the factor $\sqrt{2}$ (resp. $\sqrt{5}$) appears in edges of finite length and slope ± 1 (resp. $\frac{1}{2}$ or 2); see p. 14.

As a result of the former discussion, we see that the condition

- (*) f_2, f_4, f_6 are all positive, (therefore d_1, d_3, d_5 are all defined) and, at least, two d_i are positive.

holds for one–point central conics (clearly) and for two–point central conics (by lemma 1). The preceding paragraphs can be summarized in the following result.

Theorem 1. *Let P be a homogeneous tropical degree–two polynomial in three variables with real coefficients (then the values $f_2, f_4, f_6, d_1, d_3, d_5$ are all defined and real). If condition (*) holds, then the tropical conic \mathcal{C} defined by P is either one–point central or two–point central. Furthermore, \mathcal{C} is completely determined by the values $f_2, f_4, f_6, d_1, d_3, d_5$ and any one point associated to the matrix A corresponding to P .*

Corollary 1. *Let P be a homogeneous tropical degree–two polynomial in three variables with real coefficients (then the values $f_2, f_4, f_6, d_1, d_3, d_5$ are all defined and real). If condition (*) holds, then in $Z = 0$, the tropical conic \mathcal{C} defined by P has two different pendant edges in the NE direction (resp. W direction) (resp. S direction) and they are $\sqrt{d_1^2 + d_3^2}$ (resp. f_4) (resp. f_6) apart.*

Proof. This follows from the equalities (4) and (6) and the analysis of one–point central and two–point central conics just made. \square

Notice that $\sqrt{d_1^2 + d_3^2}$ tends to zero if and only if $f_2 = d_1 + d_3$ tends to zero.

Suppose now that $\mathcal{C}, \mathcal{C}'$ are two tropical conics as in the preceding theorem and consider their affine representations, say in $Z = 0$. It is clear that there exists a translation taking \mathcal{C} to \mathcal{C}' if and only if $d_i = d'_i$, for $i \in \{1, 3, 5\}$ and $f_k = f'_k$, for $k \in \{2, 4, 6\}$. And if $d_i = \rho d'_i$, for $i \in \{1, 3, 5\}$ and $f_k = \rho f'_k$, for $k \in \{2, 4, 6\}$, for some positive ρ then a translation followed by a ρ –homothety takes \mathcal{C} into \mathcal{C}' .

Definition 1. *A tropical conic is proper if it is not the union of two tropical lines and all of its pendant edges have multiplicity one. A tropical conic is improper if it is not proper.*

This definition is, only apparently, different from that in [16]; see corollary 5.

Given a tropical conic \mathcal{C} , let us suppose that two consecutive non–pendant vertices a, b of \mathcal{C} collapse. Then another tropical conic \mathcal{C}' appears. When a, b collapse, then the edge (of finite length) \overline{ab} vanishes, i.e., \mathcal{C}' does not have an edge \overline{ab} . At the same time, if \mathcal{C} contains parallel pendant edges e, f such that e is incident to a and f is incident to b , then e, f also collapse. This means that in \mathcal{C}' , e is an pendant edge with multiplicity two.

Definition 2. *Let \mathcal{C}' be a tropical conic which can be obtained from \mathcal{C} by successively collapsing one or more pairs of consecutive non–pendant vertices. Then we will say that \mathcal{C}' is a degeneration of \mathcal{C} .*

The degenerations of one–point central or two–point central tropical conics are represented in figure at the end (in a separate file), lines 3 to 7, where a thick segment represents a multiplicity–two edge. These conics clearly satisfy

(**) f_2, f_4, f_6 are all non–negative, (therefore d_1, d_3, d_5 are all defined) and, at least, two d_i are non–negative.

Corollary 2. *If A is a matrix such that condition (**) holds, then the tropical conic \mathcal{C} given by A is a translation of the tropical conic \mathcal{C}' given by $\text{shp}(A)$.*

Proof. Let d_i, f_k be the values associated to A and d'_i, f'_k be the values associated to $\text{shp}(A)$. Then $d_i = d'_i, f_k = f'_k$, for $i \in \{1, 3, 5\}, k \in \{2, 4, 6\}$. Now, from theorem 1 and collapsing, we realize that these two tropical conics look identical, the only difference between \mathcal{C} and \mathcal{C}' being the position of one (and hence all) of their vertices. It is easy to compute the translation vector, say in $Z = 0$; it is $\tau = \left(\frac{a_5 - a_1}{2}, \frac{a_5 - a_3}{2} \right)$, i.e., $\mathcal{C} = \mathcal{C}' + \tau$. \square

As a consequence of the previous corollary, *up to a translation*, we can assume $A = \text{shp}(A)$, i.e., in P we can assume $a_i = 0$, for $i \in \{1, 3, 5\}$.

The next class of tropical conics that we will study are those having $f_k \leq 0$, for some $k \in \{2, 4, 6\}$, including the case $f_k = -\infty$. In particular, here we *remove the extra assumption made in p. 6* and will deal with arbitrary homogeneous degree–two polynomials P in three variables. This means that P has $a_2, a_4, a_6 \in \mathbb{T}$ —some d_i might not be defined. Here our goal is to show that *no further shapes of tropical conics do appear*.

Lemma 5. *Consider $P = X^{\odot 2} \oplus Y^{\odot 2} \oplus Z^{\odot 2} \oplus \frac{f_2}{2} \odot X \odot Y \oplus \frac{f_4}{2} \odot Y \odot Z \oplus \frac{f_6}{2} \odot X \odot Z$ and suppose that $-\infty \leq f_2 \leq 0, f_4 \geq 0, f_6 \geq 0$. Then the tropical conics associated to P and P^+ are equal.*

Proof. We have $P^+ = X^{\odot 2} \oplus Y^{\odot 2} \oplus Z^{\odot 2} \oplus X \odot Y \oplus \frac{f_4}{2} \odot Y \odot Z \oplus \frac{f_6}{2} \odot X \odot Z$.

Let $\mathcal{C}, \mathcal{C}^+$ denote the corresponding conics. We show that the term $X + Y + \frac{f_2}{2}$ is irrelevant in $P = \max \left\{ 2X, 2Y, 2Z, X + Y + \frac{f_2}{2}, Y + Z + \frac{f_4}{2}, X + Z + \frac{f_6}{2} \right\}$,

when $-\infty \leq f_2 < 0$. Let $(x, y) \in \mathbb{R}^2$. If $\frac{f_2}{2} < x - y$, then $x + y + \frac{f_2}{2} \leq 2x$. And if $\frac{f_2}{2} \geq x - y$, then $\frac{f_2}{2} \leq -\frac{f_2}{2} \leq y - x$, whence $x + y + \frac{f_2}{2} \leq 2y$. This implies that

$$\max\{2X, 2Y, X + Y + \frac{f_2}{2}, \dots\} = \max\{2X, 2Y, \dots\}.$$

Moreover, letting f_2 tend to zero, the argument still holds, proving that

$$\max\{2X, 2Y, X + Y, \dots\} = \max\{2X, 2Y, \dots\}.$$

□

Notice that $f_4 \geq 0, f_6 \geq 0$ play no role in the former proof; we have made such assumptions so that the conic \mathcal{C}^+ is known to us. Therefore, the following corollary is true.

Corollary 3. *If $f_2, f_4, f_6 \in \mathbb{T}$ and $P = X^{\odot 2} \oplus Y^{\odot 2} \oplus Z^{\odot 2} \oplus \frac{f_2}{2} \odot X \odot Y \oplus \frac{f_4}{2} \odot Y \odot Z \oplus \frac{f_6}{2} \odot X \odot Z$, then the tropical conics associated to P and P^+ are equal.* □

Now, we give a **procedure to sketch, say in $Z = 0$, the tropical conic \mathcal{C} defined by an arbitrary homogeneous degree–two polynomial P .**

- From P , compute the matrices A and $\text{shp}(A)^+$ and the translation vector $\tau = \left(\frac{a_5 - a_1}{2}, \frac{a_5 - a_3}{2} \right) \in \mathbb{R}^2$.

- Draw the tropical conic \mathcal{C}^+ given by $\text{shp}(A)^+$; it is either one–point central or two–point central conic or a degeneration of them. By corollary 3, the conic given by $\text{shp}(A)$ coincides with \mathcal{C}^+ .
- Translate \mathcal{C}^+ by τ to obtain the tropical conic \mathcal{C} .

In particular, *the conic \mathcal{C} is determined by the triple $(f_2^+, f_4^+, f_6^+) \in \mathbb{R}^+$ and any point associated to the matrix A .*

The triple (f_2^+, f_4^+, f_6^+) determines the number and multiplicity of the pendant edges of the conic \mathcal{C} , by corollary 1. Indeed:

- If f_2^+, f_4^+, f_6^+ are all positive, then \mathcal{C} has six pendant edges.
- If exactly two among f_2^+, f_4^+, f_6^+ are positive, then \mathcal{C} has five pendant edges, one of them of multiplicity two.
- If only one among f_2^+, f_4^+, f_6^+ is positive, then \mathcal{C} has four pendant edges, two of them of multiplicity two. Then \mathcal{C} is a double tropical line.
- If $f_2^+ = f_4^+ = f_6^+ = 0$, then \mathcal{C} has three pendant edges, all of them of multiplicity two.

Now let

$$d_1^+ = \frac{1}{2}(f_6^+ + f_2^+ - f_4^+), d_3^+ = \frac{1}{2}(f_2^+ + f_4^+ - f_6^+), d_5^+ = \frac{1}{2}(f_4^+ + f_6^+ - f_2^+) \in \mathbb{R}.$$

Note that, if d_i is defined, d_i^+ is NOT equal to the positive part of d_i , in general. Now, the triple (d_1^+, d_3^+, d_5^+) determines which non–pendant vertices does \mathcal{C} have, as well as the lengths of the edges joining them. Indeed:

- If d_1^+, d_3^+, d_5^+ are all non–negative, then \mathcal{C} is a one–point central conic or a degeneration of it. These are precisely the tropical conics having tropically singular matrix $\text{shp}(A)^+$.
- If $d_1^+ \geq 0, d_3^+ \geq 0, d_5^+ \leq 0$, then \mathcal{C} is a two–point central type 5 conic, or a degeneration of it. (If $d_1^+ \geq 0, d_3^+ \leq 0, d_5^+ \geq 0$ or $d_1^+ \leq 0, d_3^+ \geq 0, d_5^+ \geq 0$, the situation is similar).

Here, no further cases arise, by lemma 2.

The following are all direct consequences of our discussion.

Corollary 4. *A tropical conic is degenerated if and only if $-\infty \leq f_k \leq 0$, for some $k \in \{2, 4, 6\}$ or if f_2, f_4, f_6 are all positive and $d_i = 0$, for some $i \in \{1, 3, 5\}$.*

Proof. We can work in $Z = 0$. Let \mathcal{C} be a conic satisfying the hypothesis. Say $-\infty \leq f_6 \leq 0$. Then $f_6^+ = 0$, so that \mathcal{C} has one multiplicity–two pendant edge pointing S, and then \mathcal{C} is degenerated. On the other hand, suppose that $f_k > 0$, for all $k \in \{2, 4, 6\}$ and say $d_5 = 0$. Then \mathcal{C} is a degeneration of a type 5 conic. The converse is clear. \square

Corollary 5. *Every proper tropical conic is either one–point central or two–point central. And every improper tropical conic is a degeneration of a proper conic.* \square

Corollary 6. *The exchange of the variables X, Y, Z transforms proper (resp. one–point central) (resp. two–point central) conics into proper (resp. one–point central) (resp. two–point central) conics.* \square

To finish up this section, let us gather all we know about **pairs of tropical lines**. They are represented in figure at the end (in a separate file), lines 6 to 8.

Corollary 7. *A tropical conic is a pair of lines if and only if d_1^+, d_3^+, d_5^+ are all non–negative and, at least, one d_i^+ equals zero. The number of different pendant edges in a pair of tropical lines is six, five or three.* \square

Pairs of tropical lines are degenerated tropical conics but the converse is not true (take, for instance a tropical conic having f_k^+ positive, for only one k). Moreover, pairs of tropical lines are conics in one of the following cases:

- f_2, f_4, f_6 are all positive and $d_i = 0$ for just one $i \in \{1, 3, 5\}$. Equivalently, in the matrix $\text{shp}(A) = \text{shp}(A)^+$, the vertex c coincides with just one among h, v, i .
- There exists $k \in \{2, 4, 6\}$ such that $f_{k-2} = f_{k+2} > 0$ and $-\infty \leq f_k \leq 0$. Then we have $f_k^+ = 0 = d_{k-1}^+ = d_{k+1}^+$ and $d_{k+3}^+ > 0$. Equivalently, the vertex c equals to just two among h, v, i in the matrix $\text{shp}(A)^+$.
- f_2, f_4, f_6 are all non–positive. Then we have $0 = f_2^+ = f_4^+ = f_6^+ = d_1^+ = d_3^+ = d_5^+$. The matrix $\text{shp}(A)^+$ is null and all of its vertices h, v, i, c coincide. Here we have a double tropical line.

Corollary 8. *A tropical conic \mathcal{C} is a pair of lines if and only if $c \in \{h, v, i\}$ for the matrix $\text{shp}(A)^+$.*

Proof. Use equalities (6) and the preceding discussion. \square

Let us summarize. Consider the non–negative octant $\mathcal{O} = (\mathbb{R}^+)^3$ and let (f_2^+, f_4^+, f_6^+) be the coordinates of a point in \mathcal{O} . In figure 1 we see the plane section of \mathcal{O} given by $f_2^+ + f_4^+ + f_6^+ = k$, for some positive k . Tropical conics having tropically singular matrix $\text{shp}(A)^+$ correspond to the shaded closed triangle, the boundary of which corresponds to pairs of lines. Degenerated tropical conics which are not pairs of lines correspond to the boundary of the section.

5 Factorization

A tropical polynomial p (homogeneous or not) in any number of variables is called *reducible* if it is the tropical product of two non–constant tropical polynomials. A tropical hypersurface \mathcal{C} (affine or projective) is called *reducible* if it is the union of two hypersurfaces (affine or projective, accordingly) $\mathcal{C}_1, \mathcal{C}_2$ with $\mathcal{C}_1 \neq \mathcal{C} \neq \mathcal{C}_2$. It is




Figure 1: Section of octant \mathcal{O} given by $f_2^+ + f_4^+ + f_6^+ = k$.

clear that the reducibility of a polynomial causes the reducibility of the corresponding hypersurface but, in the tropical setting, the converse is NOT true, see example 3.

Let $P = a_1 \odot X^{\odot 2} \oplus a_3 \odot Y^{\odot 2} \oplus a_5 \odot Z^{\odot 2} \oplus a_2 \odot X \odot Y \oplus a_4 \odot Y \odot Z \oplus a_6 \odot X \odot Z$ be a homogeneous degree-two tropical polynomial in three variables. We have $a_1, a_3, a_5 \in \mathbb{R}$ and $a_2, a_4, a_6 \in \mathbb{T}$. Consider the associated matrix $A = \text{trans}(A) + \text{shp}(A)$. Let us look at the following extreme cases.

Example 1. If $A = \text{trans}(A)$, then

$$h = v = i = c = \left[\frac{-a_1}{2}, \frac{-a_3}{2}, \frac{-a_5}{2} \right] = \left[\frac{a_5 - a_1}{2}, \frac{a_5 - a_3}{2}, 0 \right].$$

From the results in the previous section we deduce that the tropical conic given by P is a double line with vertex at c . In particular, (recalling that \mathbb{T} is idempotent), we have

$$P = \left(\frac{a_1}{2} \odot X \oplus \frac{a_3}{2} \odot Y \oplus \frac{a_5}{2} \odot Z \right)^{\odot 2}.$$

Example 2. If $A = \text{shp}(A)$, i.e., $P = X^{\odot 2} \oplus Y^{\odot 2} \oplus Z^{\odot 2} \oplus \frac{f_2}{2} \odot X \odot Y \oplus \frac{f_4}{2} \odot Y \odot Z \oplus \frac{f_6}{2} \odot X \odot Z$, then, up to tropical multiplication by a real constant, a tropical factorization of P must have the form

$$(a \odot X \oplus b \odot Y \oplus c \odot Z) \odot ((-a) \odot X \oplus (-b) \odot Y \oplus (-c) \odot Z), \quad (7)$$

for $a, b, c \in \mathbb{R}$. Moreover, $\frac{f_2}{2} = |a - b|$, $\frac{f_4}{2} = |b - c|$, $\frac{f_6}{2} = |c - a| \in \mathbb{R}^+$.

Corollary 9. If $P = X^{\odot 2} \oplus Y^{\odot 2} \oplus Z^{\odot 2} \oplus \frac{f_2}{2} \odot X \odot Y \oplus \frac{f_4}{2} \odot Y \odot Z \oplus \frac{f_6}{2} \odot X \odot Z$ and $-\infty \leq f_k < 0$ for some $k \in \{2, 4, 6\}$, then P is irreducible.

Example 3. If $P = X^{\odot 2} \oplus Y^{\odot 2} \oplus Z^{\odot 2} \oplus \frac{f_2}{2} \odot X \odot Y \oplus \frac{f_4}{2} \odot Y \odot Z \oplus \frac{f_6}{2} \odot X \odot Z$ and $-\infty \leq f_k < 0$ for all $k \in \{2, 4, 6\}$, then P is irreducible, by the previous corollary, but the associated tropical conic is a double line, by the results at the end of the previous section.

Corollary 10. Suppose that $P = X^{\odot 2} \oplus Y^{\odot 2} \oplus Z^{\odot 2} \oplus \frac{f_2}{2} \odot X \odot Y \oplus \frac{f_4}{2} \odot Y \odot Z \oplus \frac{f_6}{2} \odot X \odot Z$ and f_2, f_4, f_6 are all non-negative. Then P is reducible if and only if the maximum of f_2, f_4, f_6 equals the sum of the other two.

Proof. Without loss of generality, we may assume that $f_6 = \max\{f_2, f_4, f_6\}$. Suppose that $f_6 = f_2 + f_4$ and take, for instance, $a = 0, b = \frac{f_2}{2}, c = \frac{f_4}{2}$. Then P equals the product (7). The converse is easy. \square

Given P , we compute the values f_2, f_4, f_6 and write $\text{shp}(P) = X^{\odot 2} \oplus Y^{\odot 2} \oplus Z^{\odot 2} \oplus \frac{f_2}{2} \odot X \odot Y \oplus \frac{f_4}{2} \odot Y \odot Z \oplus \frac{f_6}{2} \odot X \odot Z$. Arguing as in the previous section, we realize that P is irreducible if and only if $\text{shp}(P)$ is irreducible. Therefore, here is a **procedure to determine whether an arbitrary tropical homogeneous degree-two polynomial P in three variables is reducible, and to obtain a factorization, in such a case.**

- Compute the polynomial $\text{shp}(P)$.
- Decide whether $\text{shp}(P)$ is reducible or not, using corollaries 9 and 10.
- If $\text{shp}(P)$ is reducible, factor it, as in example 2. Then compute the vector $\tau = \left(\frac{a_5 - a_1}{2}, \frac{a_5 - a_3}{2} \right)$ and translate the factorization of $\text{shp}(P)$ by $-\tau$, to obtain a factorization of P .

Example 4. Let $P = X^{\odot 2} \oplus 12 \odot Y^{\odot 2} \oplus Z^{\odot 2} \oplus 7 \odot X \odot Y \oplus 6 \odot Y \odot Z \oplus 1 \odot X \odot Z$. The associated matrices are $A = \text{trans}(A) + \text{shp}(A)$, where

$$\text{trans}(A) = \begin{pmatrix} 0 & & \\ 6 & 12 & \\ 0 & 6 & 0 \end{pmatrix} \quad \text{shp}(A) = \begin{pmatrix} 0 & & \\ 1 & 0 & \\ 1 & 0 & 0 \end{pmatrix},$$

$f_2 = f_6 = 2, f_4 = 0$ and $\max\{f_2, f_4, f_6\} = f_6 = f_2 + f_4$. Therefore the polynomial $\text{shp}(P) = X^{\odot 2} \oplus Y^{\odot 2} \oplus Z^{\odot 2} \oplus 1 \odot X \odot Y \oplus Y \odot Z \oplus 1 \odot X \odot Z$ is reducible and a factorization is given by (7) with $a = 0, b = c = 1$. The translation vector is $\tau = (0, -6)$ and thus $P = (X \oplus 7 \odot Y \oplus 1 \odot Z) \odot (X \oplus 5 \odot Y \oplus (-1) \odot Z)$.

6 Trees

Let us recall some facts about *trees*; see [3, 7] for details. A *graph* G is an ordered pair (V, E) , where V is a finite set of points, called *vertices* of G , and E is a set of

cardinality–two subsets of V . The elements of E are called *edges* of G . The edge joining vertices u, v will be denoted \overline{uv} . The *degree* of a vertex v of G is the number of edges of G incident with v . Degree–one vertices are called *pendant vertices* and edges incident to pendant vertices are called *pendant edges*.

A *tree* is a connected graph without cycles. A tree $G = (V, E)$ naturally carries a *discrete metric*; it is the function $d : V \times V \rightarrow \mathbb{N}$, where $d(u, v)$ is the least number of edges to be passed through when going from u to v . If $d(u, v) = 1$, we say that u, v are *consecutive vertices*. The *eccentricity* of a vertex v is $e(v) = \max_{u \in V} d(u, v)$ and the *radius* of the graph G is $r(G) = \min_{v \in V} e(v)$. A vertex v in G is *central* in G if $e(v) = r(G)$ and the *center* of G is the set of all central points in G . It is known that *every tree has a center and it consists either of just one vertex or two consecutive vertices*. This explains the names one–point central and two–point central, given in p. 6 and 7. It is known that a tropical projective plane curve \mathcal{C} of degree $d > 0$ is a weighted tree of a very particular sort, see [5, 13, 14, 16]. Each vertex of \mathcal{C} is determined by its tropical projective coordinates. The pendant vertices of \mathcal{C} are precisely the points in \mathcal{C} which lay on the boundary of \mathbb{TP}^2 . There are $3d$ such vertices, counted with multiplicity. They are grouped in 3 families of d vertices each: d vertices have the X (resp. Y) (resp. Z) coordinate equal to $-\infty$. Every pendant edge in \mathcal{C} has infinite length. There are $3d$ such edges, counted with multiplicity, and they are grouped in 3 families of d edges each. The rest of the edges in \mathcal{C} have finite lengths. Edges in \mathcal{C} may carry a multiplicity, which is a natural number, no greater than d . The multiplicity of a vertex is deduced from the multiplicities of the edges incident to it.

A tropical projective plane curve \mathcal{C} can be represented in $Z = 0$ (or in $Y = 0$ or $X = 0$). More precisely, this means that we represent $j_3^{-1}(\mathcal{C})$ (and still denote it \mathcal{C}) (or $j_2^{-1}(\mathcal{C})$ or $j_1^{-1}(\mathcal{C})$) in \mathbb{TA}^2 . Say, we choose to work in $Z = 0$. Then the *slope* of every edge of finite length in \mathcal{C} is a rational number and at each non–pendant vertex v the *balance condition* holds. This means that $\sum_{j=1}^s \lambda_j e_j = 0$, where u_1, \dots, u_s are all the vertices in \mathcal{C} consecutive to v , $\lambda_1, \dots, \lambda_s \in \mathbb{N}$ are the weights of the edges $\overline{vu_1}, \dots, \overline{vu_s}$ and $e_1, \dots, e_s \in \mathbb{Z}^2$ are the primitive integral vectors at the point v in the directions of such edges.

It is known that every such balanced weighted tree is a tropical curve, see [5, 12, 13]. When $d = 2$, here is a **procedure to find a defining polynomial P for such a graph \mathcal{C}** .

- From \mathcal{C} , compute the values $f_2^+, f_4^+, f_6^+, d_1^+, d_3^+, d_5^+$ and the matrix $\text{shp}(A)^+$. Classify \mathcal{C} according to section 4.
- From \mathcal{C} , compute a symmetric matrix A , using unknowns if necessary. Then compute $\text{shp}(A)$.
- According to the classification, solve for the unknowns. Substitute the unknowns in A and then obtain P .

Example 5. In $Z = 0$, let the weighted tree \mathcal{C} in figure 2

- [9] Z. Izhakian, *Duality of tropical curves*, arXiv:math.AG/0503691, 2005.
- [10] K.H. Kim, F.W. Roush, *Factorization of polynomials in one variable over the tropical semiring*, arXiv:math.CO/0501167, 2005.
- [11] G.L. Litvinov, V.P. Maslov, (eds.) *Idempotent mathematics and mathematical physics*, Cont. Math, **377**, Proceedings Vienna 2003, American Mathematical Society, Contemp. Math. **377**, (2005).
- [12] G. Mikhalkin, *Decomposition into pairs-of-pants for complex algebraic hypersurfaces*, Topology, **43**, (2004), 10035–1065.
- [13] G. Mikhalkin, *Enumerative tropical algebraic geometry in \mathbb{R}^2* , J. Amer. Math. Soc. **18**, n.2, (2005), 313–378.
- [14] G. Mikhalkin, *Tropical geometry and its applications*, in Invited lectures, v. II, Proceedings of the ICM, Madrid, 2006, (M. Sanz–Solé et al. eds.) 827–852; arXiv:math.AG/0601041, 2006.
- [15] G. Mikhalkin, *What is a tropical curve?*, Notices AMS, April 2007, 511–513.
- [16] J. Richter–Gebert, B. Sturmfels, T. Theobald, *First steps in tropical geometry*, in [11], 289–317.
- [17] D. Speyer, B. Sturmfels, *The tropical grassmannian*, Adv. Geom. **4**, 389–411, (2004).
- [18] B. Sturmfels, *Solving systems of polynomial equations*, CBMS Regional Conference Series in Math. **97**, AMS, Providence, RI, 2002.
- [19] L.F. Tabera, *Tropical constructive Pappus’s theorem*, IMRN **39**, 2373–2389 (2005).
- [20] L.F. Tabera, *Tropical plane geometric constructions: a transfer technique in tropical geometry*, arXiv:math.AG/0511713, 2007.
- [21] M. D. Vigeland, *The group law on a tropical elliptic curve*, arXiv:math.AG/05411485, 2004.
- [22] O. Viro, *Dequantization of real algebraic geometry on logarithmic paper*, European Congress of Mathematics, Vol.I (Barcelona 2000), Prog. Math. 201, Birkhäuser, Basel, 2001, 135–146.
- [23] O. Viro, *What is an amoeba?*, Notices AMS, September 2002, 916–917.

This figure "mjp_conics.gif" is available in "gif" format from:

<http://arxiv.org/ps/math/0702143v2>