

**A SUFFICIENT CONDITION FOR FINITE TIME BLOW UP OF  
THE NONLINEAR KLEIN-GORDON EQUATIONS WITH  
ARBITRARILY POSITIVE INITIAL ENERGY**

YANJIN WANG

ABSTRACT. In this paper we consider the nonexistence of global solutions of a Klein-Gordon equation of the form

$$u_{tt} - \Delta u + m^2 u = f(u) \quad (t, x) \in [0, T) \times \mathbb{R}^n.$$

Here  $m \neq 0$  and the nonlinear power  $f(u)$  satisfies some assumptions which will be stated later. We give a sufficient condition on the initial datum with arbitrarily high initial energy such that the solution of the above Klein-Gordon equation blows up in a finite time.

1. INTRODUCTION

This paper studies the nonexistence of global solutions of the nonlinear Klein-Gordon equation

$$(1.1) \quad \begin{cases} u_{tt} - \Delta u + m^2 u = f(u), & (t, x) \in [0, T) \times \mathbb{R}^n \\ u(0, x) = u_0(x), & x \in \mathbb{R}^n \\ u_t(0, x) = u_1(x), & x \in \mathbb{R}^n \end{cases}$$

where  $\Delta$  is Laplacian operator on  $\mathbb{R}^n$  ( $n \geq 1$ ),  $u_0(x)$  and  $u_1(x)$  are real valued functions,  $m \neq 0$  is a real constant, the right hand side  $f(u)$  is a real-valued, nonlinear function of  $u$ . Without loss of generality, we may assume  $m = 1$  throughout the paper.

The above problem has various applications in the area of nonlinear optics, plasma physics, fluid mechanics, etc. There is a literature on the Cauchy problem for the equation (1.1) (see, for instance, [3][5][6][7][13][14][15][16] and the papers cited therein). The results about the blow up properties for the local solution of the equation (1.1) are investigated by [1][8][12][15][16]. For the typical form of the nonlinear power,  $f(u) = |u|^{p-1}u$  with  $1 < p < \frac{n+2}{n-2}$  ( $n \geq 2$ ), we refer the result [16] to readers, which established a sharp condition for the global existence and blow up of the solution of the Klein-Gordon equation (1.1). But in [16], the initial energy had an upper bound. This motivates us to consider the problem how about the solution when the initial energy is arbitrarily large.

In the paper we investigate the above problem for the equation (1.1) with the nonlinear power  $f(u)$  which satisfies that there exists  $\epsilon > 0$  such that for any  $s \in \mathbb{R}$ ,

$$(1.2) \quad f(s)s \geq (2 + \epsilon)F(s),$$

---

2000 *Mathematics Subject Classification.* Primary 35L05, 35L15.

*Key words and phrases.* Klein-Gordon equation, Blow-up, Positive initial energy.

The author wishes to express his deep gratitude to Prof. Hitoshi Kitada for his constant encouragement and kind guidance. The study is supported by Japanese Government Scholarship.

where  $F(s) = \int_0^s f(\xi)d\xi$ .

For the above nonlinear power, the nonexistence of the global solution of an abstract wave equation (including the Klein-Gordon equation) was established in [8] when the initial energy was negative. However until recently there has been very little work on the global existence and global nonexistence of the solutions of the initial value problem for Klein-Gordon equations when the initial energy is arbitrarily high.

The purpose of this paper is to establish a sufficient condition of the initial datum with arbitrarily high initial energy such that the corresponding local solution of the nonlinear Klein-Gordon equation (1.1) blows up in a finite time. As far as we know, this is the first blow up result for the Klein-Gordon equations with arbitrarily high initial energy on the whole space  $\mathbb{R}^n$ . Our proof is based on the concavity method which firstly introduced by Levine [8][9]. We note that our proof is very simple, and it can be applied to some other equations.

Here we also refer the result [10] to readers. Levine and Todorova [10] studied a damped wave equation in the following form

$$(1.3) \quad \begin{cases} u_{tt} + a|u_t|^{\rho-1}u_t - \Delta u = b|u|^{p-1}u - q(x)^2u \text{ in } [0, T) \times \mathbb{R}^n \\ u(0, x) = u_0(x) \text{ in } \mathbb{R}^n \\ u_t(0, x) = u_1(x) \text{ in } \mathbb{R}^n \end{cases}$$

where  $a, b > 0$ ,  $\rho \geq 1$ , and  $q(x)^2 \geq 0$ . They proved the existence of the initial datum with arbitrarily high initial energy such that the solution blows up in a finite time when  $1 \leq \rho < p$ . More recently, for the case,  $\rho = 1$  and  $q(x) = 0$ , when the initial energy is arbitrarily high, Gazzola and Squassina [4] have obtained the blow up result on an open bounded Lipschitz subset of  $\mathbb{R}^n$ . We note that their proof cannot be extended to the whole space  $\mathbb{R}^n$ . Our proof can be easily adapted to the damped wave equation (1.3) with  $\rho = 1$  and  $q(x) = m \neq 0$ . In the paper we will not discuss it in detail but will make some remarks.

The paper is organized as follows. Section 2 introduces some notations and known result, and states our main result. In Section 3 we prove the main result based on a concavity argument. We make some remarks in the last section.

## 2. PRELIMINARY AND MAIN RESULT

Before we state the main result, we introduce some notations. We denote by  $\|\cdot\|_q$  the  $L^q(\mathbb{R}^n)$  norm for  $1 \leq q \leq \infty$ , and we define the spaces:  $H^1(\mathbb{R}^n) = \{u \in L^2(\mathbb{R}^n); \|u\|_{H^1(\mathbb{R}^n)} = \|(1 - \Delta)^{1/2}u\| < \infty\}$ , and  $H_0^1(\mathbb{R}^n) = \{u \in H^1(\mathbb{R}^n); \text{supp}(u) \text{ is compact in } \mathbb{R}^n\}$ . For simplicity we will denote  $\int_{\mathbb{R}^n}$  by  $\int$ . The notation  $t \rightarrow T^-$  means  $t < T$  and  $t \rightarrow T$ .

We first state the result of the local existence established in [3]

**Theorem 2.1.** *Let the initial data  $(u_0, u_1) \in H_0^1(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$ , and let  $f$  satisfy the following conditions:  $f(0) = 0$  and*

$$(2.1) \quad |f(\lambda_1) - f(\lambda_2)| \leq c(|\lambda_1|^{p-1} + |\lambda_2|^{p-1})|\lambda_1 - \lambda_2|$$

for all  $\lambda_1, \lambda_2 \in \mathbb{R}$  and some constant  $c > 0$ , and

$$(2.2) \quad 1 < p < \frac{n}{n-2} \text{ if } n \geq 3; \quad 1 < p < \infty \text{ if } n = 1, 2.$$

Then there is a unique local solution  $u(t, x)$  of the equation (1.1) on a maximal time interval  $[0, T_{\max})$  satisfying  $u(0, x) = u_0(x)$  and  $u_t(0, x) = u_1(x)$ . In addition,  $u(t, x)$  satisfies

$$(2.3) \quad E(t) = E(0)$$

where

$$(2.4) \quad E(t) = \frac{1}{2} \int (|u_t(t, x)|^2 + |u(t, x)|^2 + |\nabla u(t, x)|^2 - 2F(u(t, x))) dx.$$

In order to state the main result, we next define a function,

$$(2.5) \quad I(u) = \int (|u(x)|^2 + |\nabla u(x)|^2) dx - \int f(u(x))u(x) dx.$$

Now we are in the position to state our main result:

**Theorem 2.2.** *Let  $f(s)$  satisfy the assumption (1.2). If the initial datum  $(u_0, u_1) \in H_0^1(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$  satisfies that*

$$(2.6) \quad E(0) > 0,$$

$$(2.7) \quad \|u_0\|^2 \geq \frac{2(2+\epsilon)}{\epsilon} E(0),$$

$$(2.8) \quad I(u_0) < 0,$$

$$(2.9) \quad \int u_0 u_1 dx > 0,$$

where  $\epsilon > 0$  is the constant in (1.2), then the corresponding local solution  $u(t, x)$  of the equation (1.1) will blow up in a finite time  $T_{\max} < \infty$ , that is to say,

$$(2.10) \quad \lim_{t \rightarrow T_{\max}^-} \|u(t, \cdot)\|^2 \rightarrow \infty.$$

### 3. PROOF OF THEOREM 2.2

The proof is split into two steps.

We first prove that

$$(3.1) \quad \|u(t, \cdot)\|^2 \geq \frac{2(2+\epsilon)}{\epsilon} E(0),$$

$$(3.2) \quad I(u(t, \cdot)) < 0$$

for every  $t \in [0, T_{\max})$ .

We prove  $I(u(t, \cdot)) < 0$  for every  $t \in [0, T_{\max})$ . Suppose to the contrary that there exists a time  $T > 0$  such that

$$T = \min\{t \in (0, T_{\max}); I(u(t, \cdot)) = 0\}.$$

We now define an auxiliary function:

$$(3.3) \quad G(t) = \int |u(t, x)|^2 dx.$$

By simple computations we have

$$(3.4) \quad G'(t) = 2 \int uu_t dx,$$

$$(3.5) \quad \frac{1}{2} G''(t) = \int |u_t|^2 dx + \int (f(u)u - |\nabla u|^2 - |u|^2) dx.$$

Noting the assumption  $I(u(t, \cdot)) < 0$  for every  $t \in [0, T)$ , we obtain  $G''(t) > 0$  for every  $t \in [0, T)$ , which implies that  $G'(t)$  is strictly increasing on  $[0, T)$ , then by (2.9) we see  $G'(t) > 0$  for every  $t \in [0, T)$ . In other words, we obtain that  $G(t)$  is also strictly increasing on  $[0, T)$ . Thus we obtain

$$G(t) > G(0) \geq \frac{2(2+\epsilon)}{\epsilon} E(0)$$

for every  $t \in (0, T)$ . From the continuity of  $u(t, x)$  at  $t = T$  it follows that

$$(3.6) \quad G(T) = \|u(T, \cdot)\|^2 > \frac{2(2+\epsilon)}{\epsilon} E(0).$$

But, by (2.4) it is obvious that

$$\|u(T, \cdot)\|^2 + \|\nabla u(T, \cdot)\|^2 - 2 \int F(u(T, x)) dx \leq 2E(T) = 2E(0).$$

Noting the assumption  $I(u(T, \cdot)) = 0$  and (1.2), we have

$$\|u(T, \cdot)\|^2 + \|\nabla u(T, \cdot)\|^2 \geq (2+\epsilon) \int F(u(T, x)) dx,$$

we now obtain

$$(3.7) \quad \|u(T, \cdot)\|^2 + \|\nabla u(T, \cdot)\|^2 \leq \frac{2(2+\epsilon)}{\epsilon} E(0).$$

Obviously there is a contradiction between (3.6) and (3.7). Thus we have proved that for every  $t \in [0, T_{\max})$

$$(3.8) \quad I(u(t, \cdot)) < 0.$$

By the proof above, we also see that, if  $I(u(t, \cdot)) < 0$  for every  $t \in [0, T_{\max})$  then  $G(t)$  is strictly increasing on  $[0, T_{\max})$ . Thus (3.8) implies

$$(3.9) \quad \|u(t, \cdot)\|^2 > \frac{2(2+\epsilon)}{\epsilon} E(0)$$

for every  $t \in (0, T_{\max})$ .

Therefore we have completed the proof of (3.1) and (3.2). We next prove the blow up result of the equation (1.1).

By (3.5) and (3.9), we see that

$$\begin{aligned} \frac{1}{2} G''(t) &= \int |u_t|^2 dx - \int (|u|^2 + |\nabla u|^2) dx + \int f(u) u dx \\ &\geq (2 + \frac{\epsilon}{2}) \int |u_t|^2 dx + \frac{\epsilon}{2} \int (|u|^2 + |\nabla u|^2) dx - (2 + \epsilon) E(0) \\ &\geq (2 + \frac{\epsilon}{2}) \int |u_t|^2 dx. \end{aligned}$$

By the Schwarz inequality, we therefore have

$$(3.10) \quad \begin{aligned} G''(t)G(t) - \frac{4+\epsilon}{4}(G'(t))^2 &\geq (4+\epsilon) \left\{ \int |u|^2 dx \int |u_t|^2 dx - \left( \int uu_t dx \right)^2 \right\} \\ &\geq 0 \end{aligned}$$

for every  $t \in [0, T_{\max})$ . As  $\frac{4+\epsilon}{4} > 1$ , letting  $\alpha = \frac{\epsilon}{4}$ , we have

$$(3.11) \quad (G^{-\alpha})' = -\alpha G^{-\alpha-1} G'(t) < 0,$$

$$(3.12) \quad (G^{-\alpha})'' = -\alpha G^{-\alpha-2} \left[ G''(t)G(t) - \frac{4+\epsilon}{4} (G'(t))^2 \right] \leq 0$$

for every  $t \in (0, T_{\max})$ , which means that the function  $G^{-\alpha}$  is concave. Obviously  $G(0) > 0$ , then from (3.12) it follows that the function  $G^{-\alpha} \rightarrow 0$  when  $t < T_{\max}$  and  $t \rightarrow T_{\max}$  ( $T_{\max} < \frac{\epsilon G(0)}{4G'(0)}$ ). Therefore, we see that there exists a finite time  $T_{\max} > 0$  such that

$$(3.13) \quad \lim_{t \rightarrow T_{\max}^-} \|u(t, \cdot)\|^2 \rightarrow \infty.$$

Thus, the proof of Theorem 2.2 is completed.

#### 4. SOME REMARKS

First we consider a special case of the nonlinear power  $f(u)$ , which is the typical form of the nonlinear power,  $f(u) = |u|^{p-1}u$  with  $1 < p < \frac{n+2}{n-2}$  and  $n \geq 2$ . We define the following action

$$(4.1) \quad J(u) = \frac{1}{2} \int \left( |u|^2 + |\nabla u|^2 - \frac{2}{p+1} |u|^{p+1} \right) dx.$$

As in [2], we have that

**Theorem 4.1.** *There exists  $\bar{u}(x) \in M$ , such that*

$$(4.2) \quad d = \inf_{u \in M} J(u) = J(\bar{u})$$

where  $M = \{u \in H^1(\mathbb{R}^n); I(u) = 0, u \neq 0\}$ . Moreover,  $\bar{u}$  is the ground state of the following elliptic equation

$$-\Delta u + u = u^p, \quad u \in H^1(\mathbb{R}^n).$$

Thus, when  $E(0) < d$ , as in the first part of the proof of Theorem 2.2, we can prove that if there exists  $T$  such that  $I(u(T, \cdot)) = 0$ , the function  $u(T, x) \in H_0^1(\mathbb{R}^n) \setminus \{0\}$  satisfies

$$(4.3) \quad I(u(T, \cdot)) = 0,$$

$$(4.4) \quad J(u(T, \cdot)) \leq E(T) = E(0) < d.$$

This contradicts Theorem 4.1. Thus for the case  $E(0) < d$  the blow-up result holds. This reproduces the result in [16][11].

Secondly, we consider an example to which our method is applicable. For the damped wave equation (1.3) with  $\rho = 1$  and  $q(x)^2 = m^2 = 1$ , we define the auxiliary function  $G(t)$  corresponding to (3.3) in the following form

$$G(t) = \int |u(t, x)|^2 dx + \int_0^t \|u(\tau, \cdot)\|^2 d\tau + (T_0 - t) \|u_0\|^2,$$

where  $T_0 > 0$  is some constant. Then the argument in [4] proves  $I(u(t)) < 0$ . Thus similarly to the proof of our Theorem 2.2, we get the blow up result for the damped wave equation.

## REFERENCES

1. J.M. Ball, *Finite time blow up in nonlinear problem*, in: M.G. Grandall(Ed.), *Nonlinear Evolution equations*, Academic Press, New York, 1978, 189-205.
2. H. Berestycki and T. Cazenave, *Instabilité des états stationnaires dans les équations de Schrödinger et de Klein-Gordon non linéaires*, C.R. Acad. Sci. Paris 293(1981) 489-492.
3. T. Cazenave, *Uniform estimates for solutions of nonlinear Klein-Gordon equations*, Journal of Functional Analysis 60(1985) 36-55.
4. F. Gazzola and M. Squassina, *Global solutions and finite time blow up for damped semilinear wave equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire 23(2006) 185-207.
5. J. Ginibre and G., Velo, *The global Cauchy problem for the nonlinear Klein-Gordon equation*, Math. Z. 189(1985) 487-505.
6. J. Ginibre and G., Velo, *The global Cauchy problem for the nonlinear Klein-Gordon equation*, Ann. Inst. H. Poincaré Anal. Non Linéaire 6(1989) 15-35.
7. S. Klainerman, *Global existence of small amplitude solutions to nonlinear Klein-Gordon equations in four space-time dimensions* Commun. Pure Appl. Math. 38(1985) 631-641.
8. H.A. Levine, *Instability and nonexistence of global solutions of nonlinear wave equations of the form  $Pu_{tt} = -Au + F(u)$* , Trans. Amer. Math. Soc. 192(1974) 1-21
9. H. A. Levine, *Some additional remarks on the nonexistence of global solutions to nonlinear wave equations*, SIAM J. Math. Anal. 5(1974), 138-146.
10. H.A. Levine and G. Todorova, *Blow up of solutions of the Cauchy problem for a wave equation with nonlinear damping and source terms and positive initial energy*, Proc. Amer. Math. Soc. 129(2003), 793-805.
11. Y. Liu, M. Ohta and M. Todorova, *Strong instability of solitary waves for nonlinear Klein-Gordon equations and generalized Boussinesq equations*, Ann. I. H. Poincaré C AN (2006), doi:10.1016/j.anihpc.2006.03.005
12. L.E. Payne and D.H. Sattinger, *Saddle points and instability of nonlinear hyperbolic equations*, Israel J. Math. 22(1975) 273-303.
13. H. Peter,  *$L^p$ -Abschätzungen und klassische Lösungen für nichtlineare Wellengleichungen, I.*, Math. Z. 150(1976) 159-183.
14. J. Simon and E. Taffin *The Cauchy problem for non-linear Klein-Gordon equations*, Commun. Math. Phys. 152(1993), 433-478.
15. W. Strauss, *Nonlinear wave equations* Am. Math. Soc. 73(1989).
16. J. Zhang, *Sharp conditions of global existence for nonlinear Schrödinger and Klein-Gordon equations*, Nonlinear Analysis, 48(2002), 191-207.

GRADUATE SCHOOL OF MATHEMATICS, UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO, TOKYO, 153-8914, JAPAN.

*E-mail address:* wangyj@ms.u-tokyo.ac.jp