

ASSOCIATION SCHEMES AND CLASSICAL RCFT'S AS "GRAPHIC" FOURIER TRANSFORMATIONS

BRIAN J. DAY

ABSTRACT. Association schemes and rational conformal field theories (RCFT's) can each collectively be regarded as a type of Fourier transformation (pseudo-functor). Mathematically this is a fairly straightforward observation, but may be worth pursuing from a physical viewpoint.

1. INTRODUCTION

The (enriched) probicategories that we begin with provide the local structure constants for more complete biclosed bicategories of functors into a ground category such as $\mathcal{V} = \mathbf{Set}$ or $\mathcal{V} = \mathbf{Vect}$; they were introduced in [4] for precisely this purpose. Such complete bicategories of functors behave a bit like metric function algebras, both being familiar manifestations of "form" and "function".

Actually there is more $*$ -autonomous bicategory structure associated with the second example (cf. [5, Example 7] for the RCFT case) but we leave the formulation of this to the reader's initiative. One can find an introduction to bicategories in [2], [7], or [9].

2. THE PSEUDOFUNCTORIAL TRANSFORMATION

A *procategory* (\mathcal{A}, p, j) over $\mathcal{V} = \mathbf{Set}$ or $\mathcal{V} = \mathbf{Vect}$ (say) [4] consists of 0-cells $\{x, y, \dots\}$, and hom-categories $\mathcal{A}(x, y)$ over \mathcal{V} , related by a promultiplication and prounit

$$\begin{aligned} p &: \mathcal{A}(x, y)^{\text{op}} \otimes \mathcal{A}(y, z)^{\text{op}} \otimes \mathcal{A}(x, z) \longrightarrow \mathcal{V} \\ j &: \mathcal{A}(x, x) \longrightarrow \mathcal{V} \end{aligned}$$

designed to provide part of a biclosed bicategory structure on $\mathcal{P}(\mathcal{A}) = [\mathcal{A}, \mathcal{V}]$ via the Kan extension (convolution [3]) of the promultiplication and prounit to the respective hom-categories $[\mathcal{A}(x, y), \mathcal{V}]$ of $\mathcal{P}(\mathcal{A})$ (which we suppose has the same 0-cells as \mathcal{A}).

Next let \mathcal{B} be a biclosed bicategory over \mathcal{V} whose hom-categories $\mathcal{B}(u, v)$ are small \mathcal{V} -cocomplete. Then a given set map

$$F : \text{ob}(\mathcal{A}) \longrightarrow \text{ob}(\mathcal{B})$$

with functors $F_{x,y} : \mathcal{A}(x, y)^{\text{op}} \rightarrow \mathcal{B}(Fx, Fy)$ and natural isomorphisms

$$\begin{aligned} \int^c p(a, b, c) \otimes F(c) &\xrightarrow{\cong} F(a) \circ F(b) \\ \int^c j(c) \otimes F(c) &\xrightarrow{\cong} 1, \end{aligned}$$

yields 2-cells (which are isomorphisms)

$$\begin{array}{ccc} \mathcal{A}(x, y)^{\text{op}} \otimes \mathcal{A}(y, z)^{\text{op}} & \xrightarrow{p} & [\mathcal{A}(x, z), \mathcal{V}] \\ F \otimes F \downarrow & \cong & \downarrow \hat{F} \\ \mathcal{B}(Fx, Fy) \otimes \mathcal{B}(Fy, Fz) & \xrightarrow{\circ} & \mathcal{B}(Fx, Fz) \end{array}$$

where

$$\hat{F}(f) = \int^c f(c) \otimes F(c).$$

These extend to 2-cells (isomorphisms)

$$\begin{array}{ccc} [\mathcal{A}(x, y), \mathcal{V}] \otimes [\mathcal{A}(y, z), \mathcal{V}] & \xrightarrow{\circ} & [\mathcal{A}(x, z), \mathcal{V}] \\ \hat{F} \otimes \hat{F} \downarrow & \cong & \downarrow \hat{F} \\ \mathcal{B}(Fx, Fy) \otimes \mathcal{B}(Fy, Fz) & \xrightarrow{\circ} & \mathcal{B}(Fx, Fz); \end{array}$$

that is, isomorphisms

$$\begin{aligned} \hat{F}(f \circ g) &\cong \hat{F}(f) \circ \hat{F}(g) \\ \hat{F}(j) &\cong 1, \end{aligned}$$

given by

$$\begin{aligned} \int^c \int^{ab} f(a) \otimes g(b) \otimes p(a, b, c) \otimes F(c) &\xrightarrow{\cong} \int^{ab} f(a) \otimes g(b) \otimes (F(a) \circ F(b)) \\ \text{and } \int^c j(c) \otimes F(c) &\xrightarrow{\cong} 1, \end{aligned}$$

which we shall suppose to satisfy the axioms for \hat{F} to be a pseudofunctor (see [9, §9, Example 7]) from $\mathcal{P}(\mathcal{A})$ to \mathcal{B} .

3. ASSOCIATION SCHEMES

Suppose the **Set**-procategory \mathcal{A} is defined to have as 0-cells certain finite sets x, y, \dots , and discrete hom-categories given by $\mathcal{A}(x, y)$ is the set of members of a given partition of the cartesian product $x \times y$, together with an associative promultiplication p and prounit j (where each $\mathcal{A}(x, x)$ is assumed to contain the diagonal of $x \times x$ as representing object (1-cell) for the functor j).

A *realisation* of (\mathcal{A}, p, j) is a (locally) coproduct preserving pseudofunctor

$$\hat{F} : [\mathcal{A}, \mathbf{Set}_f] \longrightarrow \mathcal{B}$$

where \mathcal{B} is a biclosed bicategory over **Set**. The main point is that the data

$$\begin{aligned} \hat{F}(f \circ g) &\cong \hat{F}(f) \circ \hat{F}(g) \\ \hat{F}(j) &\cong 1 \end{aligned}$$

for pseudofunctoriality of \hat{F} is determined completely as the left Kan extension of the corresponding natural isomorphisms (2-cells)

$$\begin{aligned} F(a) \circ F(b) &\cong \sum_c p(a, b, c) \otimes F(c) \\ 1 &\cong F(1) \end{aligned}$$

in \mathcal{B} (as noted in §2) because \hat{F} is the left Kan extension of F . If \mathcal{B} is chosen to be the evident bicategory given by the objects all finite sets (u, v, \dots) and with hom-categories $\mathcal{B}(u, v) = \mathbf{Set}_f^{u \times v}$ with matrix multiplication as the composition, then the assignment

$$F : \mathcal{A}^{\text{op}} \longrightarrow \mathcal{B}$$

given by $F(x) = x$ and

$$\begin{aligned} F_{x,y} : \mathcal{A}(x, y) &\longrightarrow \mathcal{B}(x, y) \\ a &\longmapsto \text{the } 0\text{-}1 \text{ matrix determined by } a, \end{aligned}$$

extends to a pseudofunctor

$$\hat{F} : [\mathcal{A}, \mathbf{Set}_f] \longrightarrow \mathcal{B}$$

if (\mathcal{A}, p, j) consists of association schemes (see [1] and [10]).

4. CLASSICAL RCFT

A similar interpretation is available for RCFT based partly on the observation that each solution to the Moore-Seiberg polynomial equations yields a semisimple rigid braided tensor category (see [8, Appendix C] and [6]).

For example, let K be a field and let (\mathcal{A}, p, j) be a \mathbf{Vect}_f -procategory over K with 0-cells x, y, \dots , and hom-categories $\mathcal{A}(x, y) = 0$ if $x \neq y$, and $\mathcal{A}(x, x)$ a finite (discrete) promonoid with an identity element (usually denoted by 0) to represent the functor j . Let

$$p : \mathcal{A}(x, x)^{\text{op}} \otimes \mathcal{A}(x, x)^{\text{op}} \otimes \mathcal{A}(x, x) \longrightarrow \mathbf{Vect}_f$$

be the braided associative \mathbf{Vect} -promultiplication, usually denoted by

$$p(i, j, k) = V_{i,j}^k = N_{i,j}^k \cdot K \text{ for } i, j, k \in \mathcal{A}(x, x)$$

(which is only non-trivial on each endohom $\mathcal{A}(x, x)$). Then, if \mathcal{B} is a locally cocomplete biclosed \mathbf{Vect} -bicategory, one can view a pseudofunctor

$$\hat{F} : \mathcal{P}(\mathcal{A}) = [\mathcal{A}, \mathbf{Vect}_f] \longrightarrow \mathcal{B}$$

which is the left Kan extension of the data

$$F : \mathcal{A}^{\text{op}} \longrightarrow \mathcal{B}$$

and

$$\begin{aligned} F(i) \otimes F(j) &\cong \bigoplus_k V_{ij}^k \otimes F(k) \\ I &\cong F(0) \end{aligned}$$

in the monoidal endocategory $\mathcal{B}(Fx, Fx)$, as a realisation of the given conformal field theories (\mathcal{A}, p, j) . Of course, we could have allowed a bit more scope and not supposed that $\mathcal{A}(x, y) = 0$ for $x \neq y$ in $\text{ob}(\mathcal{A})$.

Remark. Admittedly, the probicategories \mathcal{A} featuring in §3 and §4 are rather “trivial”, however the general process (§2 etc.) promises interesting examples in more substantial cases.

REFERENCES

- [1] R. A. Bailey, Association schemes: Designed experiments, algebra and combinatorics, Cambridge University Press, 2004.
- [2] J. Bénabou, Introduction to bicategories, Lecture Notes in Math. 47 (Springer 1967) pp 1–77.
- [3] B. J. Day, On closed categories of functors, Lecture Notes in Mathematics 137 (Springer 1970), pp. 1–38.
- [4] B. J. Day, Biclosed bicategories; localisation of convolution, Macquarie Math. Reports, Report 81-0030, 1981, arXiv:math.CT/07053485, 2007.
- [5] B. J. Day, *-Autonomous categories in quantum theory, arXiv:math.CT/0605037, 2006.
- [6] R. Häring-Oldenburg, Reconstruction of weak quasi-Hopf algebras, J. Alg. 194 (1997) pp. 14–35.
- [7] S. Mac Lane, Categories for the working mathematician, Graduate Texts in Math. 5 (Springer 1998).
- [8] G. Moore and N. Seiberg, Classical and quantum conformal field theory, Comm. Math. Phys. 123 (1989) pp. 177–254.
- [9] R. Street, Categorical structures, Handbook of Algebra, Vol 1 (Edited by M. Hazewinkel), Elsevier Science B.V. (1996) pp. 529–577.
- [10] P.-H. Zieschang, An algebraic approach to association schemes, Lecture Notes in Math. 1628 (Springer 1996).

DEPARTMENT OF MATHEMATICS, MACQUARIE UNIVERSITY, NSW, 2109, AUSTRALIA