

Bounds of some real (complex) solution of a finite system
of polynomial equations with rational coefficients

Apoloniusz Tyszk

Abstract. We discuss two conjectures. **(I)** For each $x_1, \dots, x_n \in \mathbb{R}$ ($\mathbb{C}$) there exist $y_1, \dots, y_n \in \mathbb{R}$ ($\mathbb{C}$) such that

$$\begin{aligned} \forall i \in \{1, \dots, n\} \quad |y_i| &\leq 2^{2^{n-2}} \\ \forall i \in \{1, \dots, n\} \quad (x_i = 1 &\Rightarrow y_i = 1) \\ \forall i, j, k \in \{1, \dots, n\} \quad (x_i + x_j &= x_k \Rightarrow y_i + y_j = y_k) \\ \forall i, j, k \in \{1, \dots, n\} \quad (x_i \cdot x_j &= x_k \Rightarrow y_i \cdot y_j = y_k) \end{aligned}$$

(II) Let $\mathbf{G}$ be an additive subgroup of $\mathbb{C}$. Then for each $x_1, \dots, x_n \in \mathbf{G}$ there exist $y_1, \dots, y_n \in \mathbf{G} \cap \mathbb{Q}$ such that

$$\begin{aligned} \forall i \in \{1, \dots, n\} \quad |y_i| &\leq 2^{n-1} \\ \forall i \in \{1, \dots, n\} \quad (x_i = 1 &\Rightarrow y_i = 1) \\ \forall i, j, k \in \{1, \dots, n\} \quad (x_i + x_j &= x_k \Rightarrow y_i + y_j = y_k) \end{aligned}$$

For a positive integer n we define the set of equations E_n by

$$E_n = \{x_i = 1 : 1 \leq i \leq n\} \cup \{x_i + x_j = x_k : 1 \leq i \leq j \leq n, 1 \leq k \leq n\} \cup \{x_i \cdot x_j = x_k : 1 \leq i \leq j \leq n, 1 \leq k \leq n\}$$

Since there is a finite number of non-empty subsets of E_n , we get:

- (1) There is a function $\chi : \{1, 2, 3, \dots\} \rightarrow \{1, 2, 3, \dots\}$ with the property: for each positive integer n , if a non-empty subset of E_n forms a system of equations that is consistent over $\mathbb{Z}$, then this system has a solution being a sequence of integers whose absolute values are not greater than $\chi(n)$.
- (2) There is a function $\gamma : \{1, 2, 3, \dots\} \rightarrow \{1, 2, 3, \dots\}$ with the property: for each positive integer n , if a non-empty subset of E_n forms a system of equations that is consistent over $\mathbb{R}$, then this system has a solution being a sequence of real numbers whose absolute values are not greater than $\gamma(n)$.
- (3) There is a function $\theta : \{1, 2, 3, \dots\} \rightarrow \{1, 2, 3, \dots\}$ with the property: for each positive integer n , if a non-empty subset of E_n forms a system of equations that is consistent over $\mathbb{C}$, then this system has a solution being a sequence of complex numbers whose absolute values are not greater than $\theta(n)$.

2000 Mathematics Subject Classification: 03B25, 12D10, 12L05, 14P05, 15A06, 26C10.

Key words and phrases: system of polynomial equations, small solutions.

Remark. Let us consider the problem of finding a complex solution of a polynomial system with n variables and integer coefficients. This problem reduces to the problem of finding a real solution of some polynomial system with $2n$ variables and integer coefficients. Therefore, the problem of consistency over $\mathbb{C}$ of a polynomial system with n variables and integer coefficients reduces to the problem of consistency over $\mathbb{R}$ of some polynomial system with $2n$ variables and integer coefficients.

Let us note three facts:

- (4) There is a finite number of non-empty subsets of E_n .
- (5) There is an algorithm for quantifier elimination for $(\mathbb{R}, +, \cdot, 0, 1, =, \leq)$ (A. Tarski and A. Seidenberg, and later G. E. Collins with his cylindrical algebraic decomposition algorithm, see [1]). In particular, there is an algorithm checking the consistency over $\mathbb{R}$ of each finite system of polynomial equations with integer coefficients.
- (6) Applying the cylindrical algebraic decomposition algorithm, for each consistent finite system of polynomial equations with n variables and integer coefficients, we can determine $a > 0$ such that $[-a, a]^n$ contains a solution.

By the Remark and facts (4), (5), (6), we can find computable γ and θ .

There are known direct estimates which enable us to find computable γ . Let $V \subseteq \mathbb{R}^n$ be a real algebraic variety given by the system of equations $f_1 = \dots = f_m = 0$, where $f_i \in \mathbb{Q}[x_1, \dots, x_n]$ ($i = 1, \dots, m$). We denote by L the maximum of the bit-sizes of the coefficients of the system and set $d = \sum_{i=1}^m \deg(f_i)$, $r = \binom{n+2d}{n}$. We recall ([1, p. 245]) that the bit-size of a non-zero integer is the number of bits in its binary representation. More precisely, the bit-size of $k \in \mathbb{Z} \setminus \{0\}$ is τ if and only if $2^{\tau-1} \leq |k| < 2^\tau$. The bit-size of a rational number is the sum of the bit-sizes of its numerator and denominator. N. N. Vorobjov Jr. proved ([17]) that there exists $(x_1, \dots, x_n) \in V$ such that $|x_i| < 2^{H(r,L)}$ ($i = 1, \dots, n$), where H is some polynomial not depending on the initial system; for a simplified proof see [7, Lemma 9, p. 56]. A more general bound follows from [1, Theorem 13.15, p. 476].

Let M be the maximum of the absolute values of the coefficients of the polynomials $f_1(x_1, \dots, x_n), \dots, f_m(x_1, \dots, x_n) \in \mathbb{Z}[x_1, \dots, x_n]$. Assume that the system

$$f_1(x_1, \dots, x_n) = \dots = f_m(x_1, \dots, x_n) = 0$$

is consistent over $\mathbb{R}$ ($\mathbb{C}$). Let $d(i, j)$ be the degree of variable x_i in the polynomial $f_j(x_1, \dots, x_n)$. Assume that $d_i = \max\{d(i, j) : 1 \leq j \leq m\} \geq 1$ for each $i \in \{1, \dots, n\}$. Let $\mathcal{T}$ denote the family of all polynomials $W(x_1, \dots, x_n) \in \mathbb{Z}[x_1, \dots, x_n]$ for which all coefficients belong to the interval $[-M, M]$ and each variable x_i has degree less than or equal to d_i . Then, $\{x_1, \dots, x_n\} \subseteq \mathcal{T}$ and

$$\text{card } \mathcal{T} = (2M + 1)^{(d_1+1) \cdots (d_n+1)}$$

To each polynomial that belongs to $\mathcal{T} \setminus \{x_1, \dots, x_n\}$ we assign a new variable x_i with $i \in \{n+1, \dots, (2M+1)^{(d_1+1) \cdots (d_n+1)}\}$. Let $\mathcal{K}$ denote the family of all equations of the form

$$x_i = 1, x_i + x_j = x_k, x_i \cdot x_j = x_k \quad (i, j, k \in \{1, \dots, (2M+1)^{(d_1+1) \cdots (d_n+1)}\})$$

which are polynomial identities in $\mathbb{Z}[x_1, \dots, x_n]$. Let $f_j(x_1, \dots, x_n) = x_{q(j)}$, where $j \in \{1, \dots, m\}$ and $q(j) \in \{1, \dots, (2M+1)^{(d_1+1) \cdots (d_n+1)}\}$. The system

$$f_1(x_1, \dots, x_n) = \dots = f_m(x_1, \dots, x_n) = 0$$

can be equivalently write down as

$$\mathcal{K} \cup \{x_{q(1)} + x_{q(1)} = x_{q(1)}, \dots, x_{q(m)} + x_{q(m)} = x_{q(m)}\}$$

We have proved that introducing additional variables we can equivalently write down the system

$$f_1(x_1, \dots, x_n) = \dots = f_m(x_1, \dots, x_n) = 0$$

as a system of equations of the form $x_i = 1, x_i + x_j = x_k, x_i \cdot x_j = x_k$, where $i, j, k \in \{1, \dots, (2M+1)^{(d_1+1) \cdots (d_n+1)}\}$ and the variables $x_{n+1}, \dots, x_{(2M+1)^{(d_1+1) \cdots (d_n+1)}}$ are new.

Conjecture 1. Let S be a consistent system of equations in real (complex) numbers $x_1, x_2, \dots, x_n$, where each equation in S is one of the following three forms: $x_i = 1$ or $x_i + x_j = x_k$ or $x_i \cdot x_j = x_k$. Then S has a real (complex) solution $(x_1, \dots, x_n)$ in which $|x_i| \leq 2^{2^{n-2}}$ for each i .

Conjecture 1 implies that the system

$$f_1(x_1, \dots, x_n) = \dots = f_m(x_1, \dots, x_n) = 0$$

has a real (complex) solution $(x_1, \dots, x_n)$ in which $|x_i| \leq 2^{2^{(2M+1)^{(d_1+1) \cdots (d_n+1)} - 2}}$ for each i . This upper bound is rough because does not depend on the number of equations. We describe a better bound that depends on m . Let

$$L = \{(s_1, \dots, s_n) \in \mathbb{Z}^n : (0 \leq s_1 \leq d_1) \wedge \dots \wedge (0 \leq s_n \leq d_n) \wedge (1 \leq s_1 + \dots + s_n)\}$$

$$f_j(x_1, \dots, x_n) = a_j + \sum_{(s_1, \dots, s_n) \in L} a_j(s_1, \dots, s_n) \cdot x_1^{s_1} \dots x_n^{s_n}$$

where $j \in \{1, \dots, m\}$, $a_j \in \mathbb{Z}$, $a_j(s_1, \dots, s_n) \in \mathbb{Z}$. Let $\mathcal{L} = \{x_1^{s_1} \dots x_n^{s_n} : (s_1, \dots, s_n) \in L\}$. Of course, $\{x_1, \dots, x_n\} \subseteq \mathcal{L}$. We define the lexicographic order $\preceq$ on $\mathcal{L}$. We will define new variables x_i .

Step 1. To each integer in $[-M, M]$ we assign a separate variable x_i .

In this step we introduce $2M + 1$ new variables.

Step 2. To each monomial in $\mathcal{L} \setminus \{x_1, \dots, x_n\}$ we assign a separate variable x_i .

In this step we introduce $(d_1 + 1) \dots (d_n + 1) - 1 - n$ new variables.

Step 3. To each monomial $a_j(s_1, \dots, s_n) \cdot x_1^{s_1} \dots x_n^{s_n}$ ($j \in \{1, \dots, m\}$, $(s_1, \dots, s_n) \in L$) we assign a separate variable x_i .

In this step we introduce $m \cdot ((d_1 + 1) \dots (d_n + 1) - 1)$ new variables.

Step 4. To each polynomial $a_j + \sum_{(t_1, \dots, t_n) \preceq (s_1, \dots, s_n)} a_j(t_1, \dots, t_n) \cdot x_1^{t_1} \dots x_n^{t_n}$ ($j \in \{1, \dots, m\}$, $(s_1, \dots, s_n) \in L$) we assign a separate variable x_i .

In this step we introduce $m \cdot ((d_1 + 1) \dots (d_n + 1) - 1)$ new variables.

The total number of new variables is equal to

$$p = 2(M - m) - n + (2m + 1) \cdot (d_1 + 1) \dots (d_n + 1)$$

Without loss of generality we can assume that we have introduced the variables $x_{n+1}, \dots, x_{n+p}$. Let $\mathcal{H}$ denote the family of all equations of the form

$$x_i = 1, x_i + x_j = x_k, x_i \cdot x_j = x_k \quad (i, j, k \in \{1, \dots, n + p\})$$

which are polynomial identities in $\mathbb{Z}[x_1, \dots, x_n]$. Let $f_j(x_1, \dots, x_n) = x_{q(j)}$, where $j \in \{1, \dots, m\}$ and $q(j) \in \{1, \dots, n + p\}$. The system

$$f_1(x_1, \dots, x_n) = \dots = f_m(x_1, \dots, x_n) = 0$$

can be equivalently write down as

$$\mathcal{H} \cup \{x_{q(1)} + x_{q(1)} = x_{q(1)}, \dots, x_{q(m)} + x_{q(m)} = x_{q(m)}\}$$

Conjecture 1 implies that the system

$$f_1(x_1, \dots, x_n) = \dots = f_m(x_1, \dots, x_n) = 0$$

has a real (complex) solution $(x_1, \dots, x_n)$ in which $|x_i| \leq 2^{2^{2(M-m)-2+(2m+1) \cdot (d_1+1) \dots (d_n+1)}}$ for each i .

Concerning Conjecture 1, for $n = 1$ estimation by $2^{2^{n-2}}$ can be replaced by estimation by 1. For $n > 1$ estimation by $2^{2^{n-2}}$ is the best estimation. Indeed, let $n > 1$ and $\widetilde{x}_1 = 1, \widetilde{x}_2 = 2^{2^0}, \widetilde{x}_3 = 2^{2^1}, \dots, \widetilde{x}_n = 2^{2^{n-2}}$. In any ring $\mathbf{K}$ of characteristic 0, from the system of all equations belonging to E_n and are satisfied under the substitution $[x_1 \rightarrow \widetilde{x}_1, \dots, x_n \rightarrow \widetilde{x}_n]$, it follows that $x_1 = \widetilde{x}_1, \dots, x_n = \widetilde{x}_n$.

If a system $S \subseteq E_1$ is consistent over $\mathbb{C}$, then S has a solution $\widehat{x}_1 \in \{0, 1\}$. This proves Conjecture 1 for $n = 1$. If a system $S \subseteq E_2$ is consistent over $\mathbb{C}$, then S has a solution $(\widehat{x}_1, \widehat{x}_2) \in \{(0, 0), (0, 1), (1, 0), (\frac{1}{2}, 1), (1, \frac{1}{2}), (1, 1), (1, 2), (2, 1)\}$. This proves Conjecture 1 for $n = 2$. Let

$$\begin{aligned} \mathcal{W} = & \{\{1\}, \{0\}, \{1, 0\}, \{1, 2\}, \left\{1, \frac{1}{2}\right\}, \left\{1, 2, \frac{1}{2}\right\}, \{1, 0, 2\}, \left\{1, 0, \frac{1}{2}\right\}, \\ & \{1, 0, -1\}, \{1, 2, -1\}, \{1, 2, 3\}, \{1, 2, 4\}, \left\{1, \frac{1}{2}, -\frac{1}{2}\right\}, \left\{1, \frac{1}{2}, \frac{1}{4}\right\}, \left\{1, \frac{1}{2}, \frac{3}{2}\right\}, \\ & \{1, -1, -2\}, \left\{1, \frac{1}{3}, \frac{2}{3}\right\}, \{1, 2, \sqrt{2}\}, \left\{1, \frac{1}{2}, \frac{1}{\sqrt{2}}\right\}, \left\{1, \sqrt{2}, \frac{1}{\sqrt{2}}\right\}, \\ & \left\{1, \frac{\sqrt{5}-1}{2}, \frac{\sqrt{5}+1}{2}\right\}, \left\{1, \frac{\sqrt{5}+1}{2}, \frac{\sqrt{5}+3}{2}\right\}, \left\{1, \frac{-\sqrt{5}-1}{2}, \frac{\sqrt{5}+3}{2}\right\} \} \end{aligned}$$

If a system $S \subseteq E_3$ is consistent over $\mathbb{R}$, then S has a real solution $(\widehat{x}_1, \widehat{x}_2, \widehat{x}_3)$ with $\{\widehat{x}_1\} \cup \{\widehat{x}_2\} \cup \{\widehat{x}_3\} \in \mathcal{W}$. This proves Conjecture 1 for $\mathbb{R}$ and $n = 3$. If a system $S \subseteq E_3$ is consistent over $\mathbb{C}$, then S has a solution $(\widehat{x}_1, \widehat{x}_2, \widehat{x}_3)$ with $\{\widehat{x}_1\} \cup \{\widehat{x}_2\} \cup \{\widehat{x}_3\} \in \mathcal{W} \cup \left\{\left\{1, \frac{-1+\sqrt{-3}}{2}, \frac{1+\sqrt{-3}}{2}\right\}, \left\{1, \frac{1-\sqrt{-3}}{2}, \frac{1+\sqrt{-3}}{2}\right\}\right\}$. This proves Conjecture 1 for $\mathbb{C}$ and $n = 3$.

Now we demonstrate the use of the *Mathematica* software for checking Conjecture 1 for $n = 3$. Without loss of generality, we can adopt the following assumptions which allow for reducing the number of studied systems $S \subseteq E_3$.

- (7) The equation $x_1 = 1$ belongs to system S , as when all equations $x_1 = 1, x_2 = 1, x_3 = 1$ do not belong to system S , then system S has the solution $(0, 0, 0)$.
- (8) Equations $x_1 + x_1 = x_1, x_1 + x_2 = x_2, x_1 + x_3 = x_3$ do not belong to system S , as each of these equations is contradictory when $x_1 = 1$.
- (9) We only consider these systems S where each real solution $(1, x_2, x_3)$ has pairwise different 1, x_2, x_3 , as Conjecture 1 is proven for $n = 2$. Therefore, all equations $x_1 \cdot x_i = x_j$ ($i \neq j$) do not belong to system S .

- (10) Instead of each equation $x_1 + x_1 = x_i$ ($i = 2, 3$) we consider the equation $x_i = 2$.
- (11) Instead of each equation $x_i + x_i = x_1$ ($i = 2, 3$) we consider the equation $x_i = \frac{1}{2}$.
- (12) Instead of each equation $x_i + x_j = x_j$ ($2 \leq i \leq j \leq 3$) we consider the equation $x_i = 0$.
- (13) Instead of each equation $x_i + x_j = x_i$ ($1 \leq i \leq j \leq 3, 2 \leq j$) we consider the equation $x_j = 0$.
- (14) All equations $x_i \cdot x_j = x_i$ ($1 \leq i \leq j \leq 3$) do not belong to system S , because they are met when $j = 1$, and when $j > 1$ they are equivalent to equation $x_i = 0$.
- (15) All equations $x_i \cdot x_j = x_j$ ($1 \leq i \leq j \leq 3$) do not belong to system S , because they are met when $i = 1$, and when $i > 1$ they are equivalent to equation $x_j = 0$.

After replacement of variables x_1, x_2, x_3 with 1 and variables x, y , instead of set E_3 we receive 16 equations:

$$\begin{array}{cccccc}
 x = 2 & y = 2 & x = \frac{1}{2} & y = \frac{1}{2} & x = 0 & y = 0 \\
 x \cdot x = y & x \cdot x = 1 & x + x = y & y \cdot y = x & y \cdot y = 1 & y + y = x \\
 x \cdot y = 1 & x + y = 1 & x + 1 = y & y + 1 = x & &
 \end{array}$$

If $n = 3$, the following code in *Mathematica* verifies Conjecture 1 for $\mathbb{R}$ and $\mathbb{C}$:

```

Clear[x, y, i, j]
A := {x == 2, y == 2, x == 1/2, y == 1/2, x == 0, y == 0, x*x == y,
      x*x == 1, x + x == y, y*y == x, y*y == 1, y + y == x, x*y == 1,
      x + y == 1, x + 1 == y, y + 1 == x}
f[i_, j_] :=
  Reduce[Exists[{x, y},
    A[[i]] && A[[j]] && (Abs[x] > 4 || Abs[y] > 4)], Complexes] /;
  i < j
f[i_, j_] := {} /; i >= j
Union[Flatten[Table[f[i, j], {i, 1, 16}, {j, 1, 16}]]]

```

The output is {False}.

Multiple execution of the algorithm described in items (16)-(19) yields partial (as probabilistic) resolution of Conjecture 1 for $\mathbb{R}$ and $n \geq 4$.

(16) From the set E_n we remove the equations:

$$\begin{aligned}
x_i &= 1 & (1 \leq i \leq n), \\
x_1 + x_i &= x_i & (1 \leq i \leq n), \\
x_1 + x_1 &= x_i & (i \neq 2), \\
x_i + x_i &= x_1 & (i \neq 3), \\
x_i + x_j &= x_i & (1 \leq i \leq j \leq n, (i, j) \neq (4, 4)), \\
x_i + x_j &= x_j & (1 \leq i \leq j \leq n, (i, j) \neq (4, 4)), \\
x_i \cdot x_j &= x_i & (1 \leq i \leq j \leq n), \\
x_i \cdot x_j &= x_j & (1 \leq i \leq j \leq n), \\
x_1 \cdot x_i &= x_j & (i, j \in \{1, 2, \dots, n\}),
\end{aligned}$$

and in other equations we replace x_1 by 1. We obtain a non-empty set H_n .

(17) We introduce a random linear order on H_n , but with a reservation that the first equation is to be among equations involving the digit 1.

(18) We define by induction a finite sequence $(s_1, \dots, s_m)$ of equations belonging to H_n . As s_1 we put the first equation in H_n . When the sequence $(s_1, \dots, s_i)$ is defined, and there exists $h \in H_n \setminus \{s_1, \dots, s_i\}$ for which the system $\{s_1, \dots, s_i, h\}$ has a real solution $(x_2, \dots, x_n)$ with pairwise different $1, x_2, \dots, x_n$, then as s_{i+1} we put the smallest such h . If such h does not exist, then $m = i$ and the construction of the sequence $(s_1, \dots, s_m)$ is finished. If any of the systems $\{s_1, \dots, s_m, x_i = 1\}$ ($i = 2, \dots, n$), $\{s_1, \dots, s_m, x_i = x_j\}$ ($2 \leq i < j \leq n$) has a real solution, then we return to item (17).

(19) We resolve Conjecture 1 for $\mathbb{R}$ for the system $\{s_1, \dots, s_m\}$.

The condition "with pairwise different $1, x_2, \dots, x_n$ " may be removed from item (18), but this will increase the average number of executions of item (18). An intermediate possibility is that the sentence in item (18) be formulated as follows:

When the sequence $(s_1, \dots, s_i)$ is defined, and there exists $h \in H_n \setminus \{s_1, \dots, s_i\}$ for which the system $\{s_1, \dots, s_i, h\}$ has a real solution and the left side of equation h is different from the left side of each of the equations $s_1, \dots, s_i$, then as s_{i+1} we put the smallest such h .

A more complicated variant of the algorithm leads to a shorter execution time. In this variant, we first check whether the left side of equation h is different from the left side of each of the equations $s_1, \dots, s_i$ (which is not time-consuming, and preliminarily many equations h are eliminated), and then we check whether the system $\{s_1, \dots, s_i, h\}$ has a real solution with pairwise different $1, x_2, \dots, x_n$.

In the computer execution of items (18) and (19) one may use *Mathematica* software and the *CylindricalDecomposition* or *Reduce* procedure. The algorithm for Conjecture 1 for $\mathbb{C}$ is analogical, but we only apply the *Reduce* procedure. Neither for $\mathbb{R}$ nor for $\mathbb{C}$ can we apply the *Resolve* procedure in Mathematica 6.0, as it yields wrong results, as presented below:

```
Resolve[Exists[{x}, x == 0 && x*x == 1], Reals]
True
Resolve[Exists[{x}, x == 0 && x*x == 1], Complexes]
True
```

Theorem 1. If $n \in \{1, 2, 3\}$, then Conjecture 1 holds true for each subring $\mathbf{K} \subseteq \mathbb{C}$.

Proof. If a system $S \subseteq E_1$ is consistent over $\mathbf{K}$, then S has a solution $\hat{x}_1 \in \{0, 1\}$. If a system $S \subseteq E_2$ is consistent over $\mathbf{K}$ and $\frac{1}{2} \notin \mathbf{K}$, then S has a solution $(\hat{x}_1, \hat{x}_2) \in \{(0, 0), (0, 1), (1, 0), (1, 1), (1, 2), (2, 1)\}$. If a system $S \subseteq E_2$ is consistent over $\mathbf{K}$ and $\frac{1}{2} \in \mathbf{K}$, then S has a solution $(\hat{x}_1, \hat{x}_2) \in \{(0, 0), (0, 1), (1, 0), (\frac{1}{2}, 1), (1, \frac{1}{2}), (1, 1), (1, 2), (2, 1)\}$. For reducing the number of studied systems $S \subseteq E_3$, we may assume that the equation $x_1 = 1$ belongs to S , as when all equations $x_1 = 1, x_2 = 1, x_3 = 1$ do not belong to S , then S has the solution $(0, 0, 0) \in \mathbf{K}^3$. Let

$$A_2 = \{\hat{x}_2 \in \mathbb{C} : \text{there exists } \hat{x}_3 \in \mathbb{C} \text{ for which } (1, \hat{x}_2, \hat{x}_3) \text{ solves } S\}$$

$$A_3 = \{\hat{x}_3 \in \mathbb{C} : \text{there exists } \hat{x}_2 \in \mathbb{C} \text{ for which } (1, \hat{x}_2, \hat{x}_3) \text{ solves } S\}$$

We may assume that $A_2 \not\subseteq \{z \in \mathbb{C} : |z| \leq 4\}$ or $A_3 \not\subseteq \{z \in \mathbb{C} : |z| \leq 4\}$.

Case 1: $A_2 \not\subseteq \{z \in \mathbb{C} : |z| \leq 4\}$ and $A_3 \subseteq \{z \in \mathbb{C} : |z| \leq 4\}$. If $(1, \hat{x}_2, \hat{x}_3) \in \mathbf{K}^3$ solves S , then $(1, 1, \hat{x}_3) \in \mathbf{K}^3$ solves S .

Case 2: $A_2 \subseteq \{z \in \mathbb{C} : |z| \leq 4\}$ and $A_3 \not\subseteq \{z \in \mathbb{C} : |z| \leq 4\}$. If $(1, \hat{x}_2, \hat{x}_3) \in \mathbf{K}^3$ solves S , then $(1, \hat{x}_2, 1) \in \mathbf{K}^3$ solves S .

Case 3: $A_2 \not\subseteq \{z \in \mathbb{C} : |z| \leq 4\}$ and $A_3 \not\subseteq \{z \in \mathbb{C} : |z| \leq 4\}$. If $(1, \hat{x}_2, \hat{x}_3) \in \mathbf{K}^3$ solves S , then $(1, 0, 1) \in \mathbf{K}^3$ solves S or $(1, 1, 0) \in \mathbf{K}^3$ solves S or $(1, 1, 1) \in \mathbf{K}^3$ solves S .

□

The number $2 + 273^2$ is prime.

Theorem 2. If $k \in \mathbb{Z} \cap [273, \infty)$ and $2 + k^2$ is prime, then Conjecture 1 fails for $n = 6$ and the ring $\mathbb{Z}[\frac{1}{2+k^2}] = \{\frac{x}{(2+k^2)^m} : x \in \mathbb{Z}, m \in \mathbb{Z} \cap [0, \infty)\}$.

Proof. $(1, 2, k, k^2, 2 + k^2, \frac{1}{2+k^2})$ solves the system

$$\begin{cases} x_1 &= 1 \\ x_1 + x_1 &= x_2 \\ x_3 \cdot x_3 &= x_4 \\ x_2 + x_4 &= x_5 \\ x_5 \cdot x_6 &= x_1 \end{cases}$$

Assume that $(x_1, x_2, x_3, x_4, x_5, x_6) \in (\mathbb{Z}[\frac{1}{2+k^2}])^6$ solves the system. Let $x_5 = \frac{a}{(2+k^2)^p}$, $x_6 = \frac{b}{(2+k^2)^q}$, $a, b \in \mathbb{Z}$, $p, q \in \mathbb{Z} \cap [0, \infty)$. Since $2 + k^2$ is prime and $1 = |x_1| = |x_5 \cdot x_6| = \frac{|a| \cdot |b|}{(2+k^2)^{p+q}}$, we conclude that $|a| = (2 + k^2)^{\tilde{p}}$ for some $\tilde{p} \in \mathbb{Z} \cap [0, \infty)$. Hence $|x_5| = (2 + k^2)^{\tilde{p}-p}$. On the other hand, $|x_5| = |x_2 + x_4| = |x_1 + x_1 + x_3 \cdot x_3| = |1 + 1 + x_3^2| \geq 2$. Therefore, $\tilde{p} - p \geq 1$. Consequently, $|x_5| = (2 + k^2)^{\tilde{p}-p} \geq 2 + k^2 > 2^{2^{6-2}}$. $\square$

The number $-2^{32} - 2^{16} - 1$ is square-free, because $-3 \cdot 7 \cdot 13 \cdot 97 \cdot 241 \cdot 673$ is the factorization of $-2^{32} - 2^{16} - 1$ into prime numbers.

Theorem 3. If $n = 6$, then Conjecture 1 fails for the ring $\mathbb{Z}[\sqrt{-2^{32} - 2^{16} - 1}] = \{x + y \cdot \sqrt{-2^{32} - 2^{16} - 1} : x, y \in \mathbb{Z}\}$.

Proof. $(1, 2^{16} + 1, -2^{16}, -2^{32} - 2^{16}, \sqrt{-2^{32} - 2^{16} - 1}, -2^{32} - 2^{16} - 1)$ solves the system

$$\begin{cases} x_1 &= 1 \\ x_2 + x_3 &= x_1 \\ x_2 \cdot x_3 &= x_4 \\ x_5 \cdot x_5 &= x_6 \\ x_1 + x_6 &= x_4 \end{cases}$$

which has no integer solutions. For each $z \in \mathbb{Z}[\sqrt{-2^{32} - 2^{16} - 1}]$, if $|z| \leq 2^{2^{6-2}}$ then $z \in \mathbb{Z}$. $\square$

Observation 1. If $q, a, b, c, d \in \mathbb{Z}$, $b \neq 0$ or $d \neq 0$, $q \geq 2$, q is square-free, and $(a + b\sqrt{q}) \cdot (c + d\sqrt{q}) = 1$, then

$$(a \geq 1 \wedge b \geq 1) \vee (a \leq -1 \wedge b \leq -1) \vee (c \geq 1 \wedge d \geq 1) \vee (c \leq -1 \wedge d \leq -1)$$

The number $4 \cdot 13^4 - 1$ is square-free, because $3 \cdot 113 \cdot 337$ is the factorization of $4 \cdot 13^4 - 1$ into prime numbers.

Theorem 4. If $p \in \mathbb{Z} \cap [13, \infty)$ and $4p^4 - 1$ is square-free, then Conjecture 1 fails for $n = 5$ and the ring $\mathbb{Z}[\sqrt{4p^4 - 1}] = \{x + y \cdot \sqrt{4p^4 - 1} : x, y \in \mathbb{Z}\}$.

Proof. $(1, 2p^2 + \sqrt{4p^4 - 1}, 2p^2 - \sqrt{4p^4 - 1}, 4p^2, 2p)$ solves the system

$$\begin{cases} x_1 &= 1 \\ x_2 \cdot x_3 &= x_1 \\ x_2 + x_3 &= x_4 \\ x_5 \cdot x_5 &= x_4 \end{cases}$$

Assume that $(x_1, x_2, x_3, x_4, x_5) \in (\mathbb{Z}[\sqrt{4p^4 - 1}])^5$ solves the system. Let $x_2 = a + b\sqrt{4p^4 - 1}$, $x_3 = c + d\sqrt{4p^4 - 1}$, $a, b, c, d \in \mathbb{Z}$. Since the system has no integer solutions, $b \neq 0$ or $d \neq 0$. Since $x_2 \cdot x_3 = x_1 = 1$, Observation 1 implies that $|x_2| \geq 1 + \sqrt{4p^4 - 1} > 2^{2^{5-2}}$ or $|x_3| \geq 1 + \sqrt{4p^4 - 1} > 2^{2^{5-2}}$. □

For each $a, b, c \in \mathbb{R}$ ($\mathbb{C}$) we define $S(a, b, c)$ as

$$\{\mathcal{E} \in E_3 : \mathcal{E} \text{ is satisfied under the substitution } [x_1 \rightarrow a, x_2 \rightarrow b, x_3 \rightarrow c]\}$$

If $a, b, c \in \mathbb{R}$ and $\{a\} \cup \{b\} \cup \{c\} \in \mathcal{W}$, then the system $S(a, b, c)$ is consistent over $\mathbb{R}$, has a finite number of real solutions, and each real solution of $S(a, b, c)$ belongs to $[-4, 4]^3$. The family

$$\{S(a, b, c) : a, b, c \in \mathbb{R} \wedge \{a\} \cup \{b\} \cup \{c\} \in \mathcal{W}\}$$

equals to the family of all systems $S \subseteq E_3$ which are consistent over $\mathbb{R}$ and maximal with respect to inclusion.

If $a, b, c \in \mathbb{C}$ and $\{a\} \cup \{b\} \cup \{c\} \in \mathcal{W} \cup \left\{ \left\{ 1, \frac{-1+\sqrt{-3}}{2}, \frac{1+\sqrt{-3}}{2} \right\}, \left\{ 1, \frac{1-\sqrt{-3}}{2}, \frac{1+\sqrt{-3}}{2} \right\} \right\}$, then the system $S(a, b, c)$ is consistent over $\mathbb{C}$, has a finite number of solutions, and each solution of $S(a, b, c)$ belongs to $\{(z_1, z_2, z_3) \in \mathbb{C}^3 : |z_1| \leq 4 \wedge |z_2| \leq 4 \wedge |z_3| \leq 4\}$. The family

$$\begin{aligned} &\{S(a, b, c) : a, b, c \in \mathbb{C} \wedge \{a\} \cup \{b\} \cup \{c\} \in \\ &\mathcal{W} \cup \left\{ \left\{ 1, \frac{-1+\sqrt{-3}}{2}, \frac{1+\sqrt{-3}}{2} \right\}, \left\{ 1, \frac{1-\sqrt{-3}}{2}, \frac{1+\sqrt{-3}}{2} \right\} \right\} \} \end{aligned}$$

equals to the family of all systems $S \subseteq E_3$ which are consistent over $\mathbb{C}$ and maximal with respect to inclusion.

A stronger form of Conjecture 1 for $\mathbb{R}$ is:

- (20) If a system $S \subseteq E_n$ is consistent over $\mathbb{R}$ and maximal with respect to inclusion, then S has a finite number of real solutions and each real solution of S belongs to $[-2^{2^{n-2}}, 2^{2^{n-2}}]^n$.

A stronger form of Conjecture 1 for $\mathbb{C}$ is:

- (21) If a system $S \subseteq E_n$ is consistent over $\mathbb{C}$ and maximal with respect to inclusion, then S has a finite number of solutions and each solution of S belongs to $\{(z_1, \dots, z_n) \in \mathbb{C}^n : |z_1| \leq 2^{2^{n-2}} \wedge \dots \wedge |z_n| \leq 2^{2^{n-2}}\}$.

We can formulate Conjecture 1 as follows: for each $x_1, \dots, x_n \in \mathbb{R}$ ($\mathbb{C}$) there exist $y_1, \dots, y_n \in \mathbb{R}$ ($\mathbb{C}$) such that

$$\begin{aligned} \forall i \in \{1, \dots, n\} \quad & |y_i| \leq 2^{2^{n-2}} \\ \forall i \in \{1, \dots, n\} \quad & (x_i = 1 \Rightarrow y_i = 1) \\ \forall i, j, k \in \{1, \dots, n\} \quad & (x_i + x_j = x_k \Rightarrow y_i + y_j = y_k) \\ \forall i, j, k \in \{1, \dots, n\} \quad & (x_i \cdot x_j = x_k \Rightarrow y_i \cdot y_j = y_k) \end{aligned}$$

Conjecture 2 strengthens Conjecture 1.

Conjecture 2. For each $x_1, \dots, x_n \in \mathbb{R}$ ($\mathbb{C}$) there exist $y_1, \dots, y_n \in \mathbb{R}$ ($\mathbb{C}$) such that

$$\begin{aligned} \forall i \in \{1, \dots, n\} \quad & |y_i| \leq 2^{2^{n-2}} \\ \forall i \in \{1, \dots, n\} \quad & |y_i - 1| \leq |x_i - 1| \\ \forall i, j, k \in \{1, \dots, n\} \quad & |y_i + y_j - y_k| \leq |x_i + x_j - x_k| \\ \forall i, j, k \in \{1, \dots, n\} \quad & |y_i \cdot y_j - y_k| \leq |x_i \cdot x_j - x_k| \end{aligned}$$

We say that $X \subseteq \mathbb{R}$ ($\mathbb{C}$) has a property $\mathcal{B}$, if for each $x_1, \dots, x_n \in X$ there exist $y_1, \dots, y_n \in X$ with the above four properties. We define:

$$\mathcal{F}_{\mathbb{R}} = \{X \subseteq \mathbb{R} : X \text{ has property } \mathcal{B}\}$$

$$\mathcal{F}_{\mathbb{C}} = \{X \subseteq \mathbb{C} : X \text{ has property } \mathcal{B}\}$$

If $X \subseteq [-\sqrt{2}, \sqrt{2}]$ then $X \in \mathcal{F}_{\mathbb{R}}$. If $X \subseteq \{z \in \mathbb{C} : |z| \leq \sqrt{2}\}$ then $X \in \mathcal{F}_{\mathbb{C}}$.

Theorem 5. The family $\mathcal{F}_{\mathbb{R}}$ ($\mathcal{F}_{\mathbb{C}}$) has a maximal element.

We prove: if $\mathcal{C} \subseteq \mathcal{F}_{\mathbb{R}}$ is a chain, then $\bigcup_{X \in \mathcal{C}} X \in \mathcal{F}_{\mathbb{R}}$. Since $\mathcal{C}$ is a chain, for each $x_1, \dots, x_n \in \bigcup_{X \in \mathcal{C}} X$ there exists $X \in \mathcal{C}$ with $x_1, \dots, x_n \in X$. Since X has property $\mathcal{B}$,

we obtain suitable $y_1, \dots, y_n \in X \cap [-2^{2^{n-2}}, 2^{2^{n-2}}] \subseteq \left(\bigcup_{X \in \mathcal{C}} X\right) \cap [-2^{2^{n-2}}, 2^{2^{n-2}}]$.

We have proved that $\bigcup_{X \in \mathcal{C}} X \in \mathcal{F}_{\mathbb{R}}$. By Zorn's lemma, the family $\mathcal{F}_{\mathbb{R}}$ has a maximal element. The proof for $\mathcal{F}_{\mathbb{C}}$ is analogical.

□

It is hardly to decide whether Theorem 5 may help prove that $\mathbb{R} \in \mathcal{F}_{\mathbb{R}}$ and $\mathbb{C} \in \mathcal{F}_{\mathbb{C}}$.

Since $(\mathbb{R}, +, \cdot, 0, 1, =, \leq)$ is decidable, Conjectures 1 and 2 for $\mathbb{R}$ are decidable for each fixed n . By the same reason and the Remark, Conjectures 1 and 2 for $\mathbb{C}$ are decidable for each fixed n .

Hilbert's tenth problem is to give a computing algorithm which will tell of a given polynomial equation with integer coefficients whether or not it has a solution in integers. Yu. V. Matijasevič proved ([10]) that there is no such algorithm, see also [11], [4], [5], [8]. It implies that Conjecture 1 is false for $\mathbb{Z}$ instead of $\mathbb{R}$ ($\mathbb{C}$). Moreover, Matijasevič's theorem implies that Conjecture 1 for $\mathbb{Z}$ is false with any other computable estimation instead of $2^{2^{n-2}}$, so each χ in item (1) is not computable.

As we have proved Conjecture 1 for $\mathbb{Z}$ is false. We describe a counterexample showing it directly.

Lemma 1. In $\mathbb{Z}$, the formula $x < y$ is equivalent to

$$\begin{aligned} & \exists w \exists s_1 \exists s_2 \exists s_3 \exists s_4 \exists u_1 \exists u_2 \exists u_3 \exists u_4 \exists z_1 \exists z_2 \exists z_3 \exists z_4 \\ & ((w = 1) \wedge (s_1 \cdot s_1 = u_1) \wedge (s_2 \cdot s_2 = u_2) \wedge (s_3 \cdot s_3 = u_3) \wedge (s_4 \cdot s_4 = u_4) \wedge \\ & (x + w = z_1) \wedge (z_1 + u_1 = z_2) \wedge (z_2 + u_2 = z_3) \wedge (z_3 + u_3 = z_4) \wedge (z_4 + u_4 = y)) \end{aligned}$$

Proof. It follows from the Lagrange four square theorem ([13]) which states that each non-negative integer is a sum of four squares of integers.

□

Let the sequence $\{a_n\}$ be defined inductively by $a_1 = 2$, $a_{n+1} = a_n^{a_n}$. Let $A = \{a_1, a_2, a_3, \dots\}$.

Lemma 2. In $\mathbb{Z}$, for some integer $m \geq 2$ the condition $t_1 \in A$ can be equivalently formulated as $\exists t_2 \dots \exists t_m \Phi(t_1, t_2, \dots, t_m)$, where $\Phi(t_1, t_2, \dots, t_m)$ is a conjunction of formulae of the form $t_i = 1$, $t_i + t_j = t_k$, $t_i \cdot t_j = t_k$.

Proof. A is recursive, so, in particular, A is recursively enumerable. By Matijasevič's theorem ([10],[11],[4],[5],[8]) there exists a Diophantine definition of A .

□

Applying Lemma 1 $n - 1$ times and applying Lemma 2 n times, we obtain: in $\mathbb{Z}$, the condition

$$(x_1 < x_2) \wedge \dots \wedge (x_{n-1} < x_n) \wedge (x_1 \in A) \wedge \dots \wedge (x_n \in A)$$

can be equivalently formulated as $\exists x_{n+1} \dots \exists x_{p(n)} \Psi(x_1, \dots, x_n, x_{n+1}, \dots, x_{p(n)})$, where $p(n) = n + (n - 1) \cdot 13 + n \cdot (m - 1)$ and $\Psi(x_1, \dots, x_n, x_{n+1}, \dots, x_{p(n)})$ is a conjunction of formulae of the form $x_i = 1$, $x_i + x_j = x_k$, $x_i \cdot x_j = x_k$. We find a positive integer r for which $a_r > 2^{2^{p(r)-2}}$. There exist integers $x_1, \dots, x_r, x_{r+1}, \dots, x_{p(r)}$ satisfying $\Psi(x_1, \dots, x_r, x_{r+1}, \dots, x_{p(r)})$. If integers $x_1, \dots, x_r, x_{r+1}, \dots, x_{p(r)}$ satisfy $\Psi(x_1, \dots, x_r, x_{r+1}, \dots, x_{p(r)})$, then $x_r \geq a_r > 2^{2^{p(r)-2}}$.

A similar counterexample, with a more direct construction, is presented now.

Yu. V. Matijasevič proved ([10]) that the relation $z = x^y$ is Diophantine, see also [3], [4], [12], [8]. Applying it $n - 1$ times, we find the Diophantine definition of the relation

$$\left\{ (x_1, \dots, x_n) \in \mathbb{Z}^n : (x_1 = 2) \wedge (x_2 = x_1^{x_1}) \wedge \dots \wedge (x_{i+1} = x_i^{x_i}) \wedge \dots \wedge (x_n = x_{n-1}^{x_{n-1}}) \right\}$$

The rest of the proof goes as before.

Question. For which $n \in \{1, 2, 3, \dots\}$ there exists a continuous function

$$\mathbb{R}^n \ni (x_1, \dots, x_n) \xrightarrow{f_n} (f_{(n,1)}(x_1, \dots, x_n), \dots, f_{(n,n)}(x_1, \dots, x_n)) \in [-2^{2^{n-2}}, 2^{2^{n-2}}]^n$$

such that

$$\forall (x_1, \dots, x_n) \in \mathbb{R}^n \forall i \in \{1, \dots, n\} (x_i = 1 \Rightarrow f_{(n,i)}(x_1, \dots, x_n) = 1)$$

$$\forall (x_1, \dots, x_n) \in \mathbb{R}^n \forall i, j, k \in \{1, \dots, n\} (x_i + x_j = x_k \Rightarrow$$

$$f_{(n,i)}(x_1, \dots, x_n) + f_{(n,j)}(x_1, \dots, x_n) = f_{(n,k)}(x_1, \dots, x_n))$$

$$\forall (x_1, \dots, x_n) \in \mathbb{R}^n \forall i, j, k \in \{1, \dots, n\} (x_i \cdot x_j = x_k \Rightarrow$$

$$f_{(n,i)}(x_1, \dots, x_n) \cdot f_{(n,j)}(x_1, \dots, x_n) = f_{(n,k)}(x_1, \dots, x_n))$$

Theorem 6. Such functions exist for $n = 1$ and $n = 2$.

Proof. Case $n = 1$. We define $f_1 : \mathbb{R} \rightarrow [0, 1]$ by

$$f_1(x) = \begin{cases} 0 & \text{if } x \in (-\infty, 0) \\ x & \text{if } x \in [0, 1] \\ 1 & \text{if } x \in (1, \infty) \end{cases}$$

Case $n = 2$. Let A be a closed subset of a metric space X and let X be a locally convex topological linear space. The Dugundji theorem ([6]) states that every continuous map $f : A \rightarrow X$ can be extended continuously to all of X in such a way that the range of

the extension lies in the convex hull of $f(A)$. Applying the Dugundji theorem we find a continuous function $f_2 : \mathbb{R}^2 \rightarrow [-2, 2]^2$ with the following properties:

$$\begin{aligned} \forall x, y \in [-2, 2] \quad f_2(x, y) &= (x, y) \\ \forall x \in (-\infty, -2) \quad (f_2(x, 1) &= (-2, 1) \quad \wedge \quad f_2(1, x) = (1, -2)) \\ \forall x \in (2, \infty) \quad (f_2(x, 1) &= (2, 1) \quad \wedge \quad f_2(1, x) = (1, 2)) \\ \forall x \in (-\infty, -2) \quad (f_2(x, 0) &= (-2, 0) \quad \wedge \quad f_2(0, x) = (0, -2)) \\ \forall x \in (2, \infty) \quad (f_2(x, 0) &= (2, 0) \quad \wedge \quad f_2(0, x) = (0, 2)) \\ \forall x \in (-\infty, -1) \quad (f_2(x, 2x) &= (-1, -2) \quad \wedge \quad f_2(2x, x) = (-2, -1)) \\ \forall x \in (1, 2] \quad (f_2(x, 2x) &= (2 - x, 4 - 2x) \quad \wedge \quad f_2(2x, x) = (4 - 2x, 2 - x)) \\ \forall x \in (2, \infty) \quad (f_2(x, 2x) &= (0, 0) \quad \wedge \quad f_2(2x, x) = (0, 0)) \\ \forall x \in (-\infty, -\sqrt{2}) \quad (f_2(x, x^2) &= (-\sqrt{2}, 2) \quad \wedge \quad f_2(x^2, x) = (2, -\sqrt{2})) \\ \forall x \in (\sqrt{2}, 2] \quad (f_2(x, x^2) &= (\sqrt{4 - x^2}, 4 - x^2) \quad \wedge \quad f_2(x^2, x) = (4 - x^2, \sqrt{4 - x^2})) \\ \forall x \in (2, \infty) \quad (f_2(x, x^2) &= (0, 0) \quad \wedge \quad f_2(x^2, x) = (0, 0)) \end{aligned}$$

We propose an effective description of a continuous $f_2 : \mathbb{R}^2 \rightarrow [-2, 2]^2$. We define $\sigma : \mathbb{R} \rightarrow [-2, 2]$ by

$$\sigma(x) = \begin{cases} -2 & \text{if } x \in (-\infty, -2) \\ x & \text{if } x \in [-2, 2] \\ 2 & \text{if } x \in (2, \infty) \end{cases}$$

Let

$$T = [-2, 2]^2 \cup$$

$$\{(x, y) \in \mathbb{R}^2 : y = 1 \vee x = 1 \vee y = 0 \vee x = 0 \vee y = 2x \vee x = 2y \vee y = x^2 \vee x = y^2\}$$

Let $\rho : \mathbb{R}^2 \setminus T \rightarrow (0, \infty)$ be defined by

$$\begin{aligned} \rho(x, y) &= \frac{1}{|x - \sigma(x)| + |y - \sigma(y)|} + \\ &\frac{1}{|y - 1|} + \frac{1}{|x - 1|} + \frac{1}{|y - 0|} + \frac{1}{|x - 0|} + \frac{1}{|y - 2x|} + \frac{1}{|x - 2y|} + \frac{1}{|y - x^2|} + \frac{1}{|x - y^2|} \end{aligned}$$

and let $g : \mathbb{R}^2 \setminus T \rightarrow [-2, 2]^2$ be defined by

$$\begin{aligned} g(x, y) &= \frac{1}{\rho(x, y)} \cdot \left(\frac{f_2(\sigma(x), \sigma(y))}{|x - \sigma(x)| + |y - \sigma(y)|} + \right. \\ &\left. \frac{f_2(x, 1)}{|y - 1|} + \frac{f_2(1, y)}{|x - 1|} + \frac{f_2(x, 0)}{|y - 0|} + \frac{f_2(0, y)}{|x - 0|} + \frac{f_2(x, 2x)}{|y - 2x|} + \frac{f_2(2y, y)}{|x - 2y|} + \frac{f_2(x, x^2)}{|y - x^2|} + \frac{f_2(y^2, y)}{|x - y^2|} \right) \end{aligned}$$

Let $f_2|T$ denote f_2 restricted to T . The function f_2 has an exact definition on T , and $(f_2|T) \cup g : \mathbb{R}^2 \rightarrow [-2, 2]^2$ is continuous.

□

Let $\mathbf{K}$ be a ring and let $A \subseteq \mathbf{K}$. We say that a map $f : A \rightarrow \mathbf{K}$ is *arithmetic* if it satisfies the following conditions:

if $1 \in A$ then $f(1) = 1$,

if $a, b \in A$ and $a + b \in A$ then $f(a + b) = f(a) + f(b)$,

if $a, b \in A$ and $a \cdot b \in A$ then $f(a \cdot b) = f(a) \cdot f(b)$.

We call an element $r \in \mathbf{K}$ *arithmetically fixed* if there is a finite set $A \subseteq \mathbf{K}$ (an *arithmetic neighbourhood* of r inside $\mathbf{K}$) with $r \in A$ such that each arithmetic map $f : A \rightarrow \mathbf{K}$ fixes r , i.e. $f(r) = r$. If $\mathbf{K}$ is a field, then any $r \in \mathbf{K}$ is arithmetically fixed if and only if $\{r\}$ is existentially first-order definable in the language of rings without parameters, see [15]. Articles [14], [9], [15] dealt with a description of a situation where for an element in a field there exists an arithmetic neighbourhood. Article [16] describes various types of arithmetic neighbourhoods inside $\mathbb{Z}$ and $\mathbb{Q}$.

Let $\widetilde{\mathbf{K}}$ denote the set of all $r \in \mathbf{K}$ that are arithmetically fixed. Let $\widetilde{\mathbf{K}}_n$ ($n = 1, 2, 3, \dots$) denote the set of all $r \in \mathbf{K}$ for which there exists an arithmetic neighbourhood A of r such that $\text{card}(A) \leq n$. Obviously, $\widetilde{\mathbf{K}}_1 = \{0, 1\}$ and $\widetilde{\mathbf{K}}_2 \in \{\{0, 1\}, \{0, 1, 2\}, \{0, 1, 2, \frac{1}{2}\}\}$.

By Theorem 3 in [14] $\widetilde{\mathbb{R}}_n \subseteq \mathbb{R}^{\text{alg}} = \{x \in \mathbb{R} : x \text{ is algebraic over } \mathbb{Q}\}$. By this, Conjecture 1 implies $\widetilde{\mathbb{R}}_n \subseteq \mathbb{R}^{\text{alg}} \cap [-2^{2^{n-2}}, 2^{2^{n-2}}]$. By Corollary 2 in [14] $\widetilde{\mathbb{C}}_n \subseteq \mathbb{Q}$. By this, Conjecture 1 implies $\widetilde{\mathbb{C}}_n \subseteq \mathbb{Q} \cap [-2^{2^{n-2}}, 2^{2^{n-2}}]$.

Theorem 7 (cf. [14]). For each $n \in \{3, 4, 5, \dots\}$ we have $\text{card}(\widetilde{\mathbf{K}}_n) \leq (n+1)^{n^2+n} + 2$.

Proof. If $\text{card}(\mathbf{K}) < n$ then $\text{card}(\widetilde{\mathbf{K}}_n) \leq \text{card}(\mathbf{K}) < n < (n+1)^{n^2+n} + 2$. In the rest of the proof we assume that $\text{card}(\mathbf{K}) \geq n$. Let $r \in \widetilde{\mathbf{K}}_n \setminus \{0, 1\}$ and A is a neighbourhood of r such that $\text{card}(A) \leq n$. Then each set B with $A \subseteq B \subseteq \mathbf{K}$ and $\text{card}(B) = n$ is a neighbourhood of r . Observe that $1 \in B$, because in the opposite case the arithmetic map $B \rightarrow \{0\}$ moves $r \neq 0$, which is impossible. Since $r \neq 1$, we can choose B with $\mathbf{K} \supseteq B = \{x_1, \dots, x_n\} \supseteq A$, where $x_1 = r$, $x_n = 1$, and $x_i \neq x_j$ if $i \neq j$. We choose all formulae $x_i + x_j = x_k$, $x_i \cdot x_j = x_k$ ($1 \leq i \leq j \leq n$, $1 \leq k \leq n$) that are satisfied in B . Joining these formulae with conjunctions we get some formula Φ . Let V denote the set of variables in $\Phi \wedge (x_n = 1)$. Observe that $x_1 \in V$, since otherwise for any $s \in \mathbf{K} \setminus \{r\}$ the mapping $f = \text{id}(B \setminus \{r\}) \cup \{(r, s)\}$ satisfies conditions (1)-(3) and

$f(r) \neq r$. The formula $\underbrace{\dots \exists x_i \dots}_{x_i \in V, i \neq 1} (\Phi \wedge (x_n = 1))$ is satisfied in $\mathbf{K}$ if and only if $x_1 = r$.

For each $(i, j) \in \{(i, j) : 1 \leq i \leq j \leq n\}$ there are $n + 1$ possibilities:

$$x_i + x_j = x_1, \dots, x_i + x_j = x_n, \quad x_i + x_j \notin \{x_1, \dots, x_n\}.$$

For each $(i, j) \in \{(i, j) : 1 \leq i \leq j \leq n\}$ there are $n + 1$ possibilities:

$$x_i \cdot x_j = x_1, \dots, x_i \cdot x_j = x_n, \quad x_i \cdot x_j \notin \{x_1, \dots, x_n\}.$$

Since $\text{card}(\{(i, j) : 1 \leq i \leq j \leq n\}) = \frac{n^2+n}{2}$, the number of possible formulae $\Phi \wedge (x_n = 1)$ does not exceed $(n + 1)^{\frac{n^2+n}{2}} \cdot (n + 1)^{\frac{n^2+n}{2}} = (n + 1)^{n^2+n}$. Thus $\text{card}(\widetilde{\mathbf{K}}_n \setminus \{0, 1\}) \leq (n + 1)^{n^2+n}$, so $\text{card}(\widetilde{\mathbf{K}}_n) \leq (n + 1)^{n^2+n} + 2$. □

As we have seen in the proof of Theorem 7, each n -element arithmetic neighbourhood of $r \in \mathbf{K}$ determines a system of equations belonging to some non-empty subset of E_n . In the ring $\mathbf{K}$, for each solution of this system the value of variable x_1 is r .

Considering all systems $S \subseteq E_i$ ($i = 1, 2, 3$) we get: $\widetilde{\mathbb{Q}}_1 = \widetilde{\mathbb{R}}_1 = \widetilde{\mathbb{C}}_1 = \{0, 1\}$, $\widetilde{\mathbb{Q}}_2 = \widetilde{\mathbb{R}}_2 = \widetilde{\mathbb{C}}_2 = \{0, 1, 2, \frac{1}{2}\}$, $\widetilde{\mathbb{Q}}_3 = \widetilde{\mathbb{R}}_3 = \widetilde{\mathbb{C}}_3 = \{0, 1, 2, \frac{1}{2}, -1, 3, 4, -\frac{1}{2}, \frac{1}{4}, \frac{3}{2}, -2, \frac{1}{3}, \frac{2}{3}\}$.

For any ring $\mathbf{K}$ and any $r \in \widetilde{\mathbf{K}}$ we define $\omega(r) \in \{1, 2, 3, \dots\}$ as

$$\min \{\text{card}(A) : \{r\} \subseteq A \subseteq \mathbf{K} \wedge A \text{ is an arithmetic neighbourhood of } r \text{ inside } \mathbf{K}\}$$

As a corollary of Theorem 7 we obtain

$$\forall n \in \{3, 4, 5, \dots\} \forall B \subseteq \widetilde{\mathbf{K}} \left(\text{card}(B) > (n + 1)^{n^2+n} + 2 \implies \exists r \in B \omega(r) > n \right)$$

Obviously, each $\widetilde{\mathbf{K}}_n$ is finite (Theorem 7 gives a concrete upper bound for $\text{card}(\widetilde{\mathbf{K}}_n)$), so for any subring $\mathbf{K} \subseteq \mathbb{C}$ there exists $\lambda : \{1, 2, 3, \dots\} \rightarrow \{1, 2, 3, \dots\}$ such that

$$\forall n \in \{1, 2, 3, \dots\} \forall z \in \widetilde{\mathbf{K}}_n |z| \leq \lambda(n)$$

The author does not know whether for $\mathbf{K} = \mathbb{Z}$ there exists a computable $\lambda : \{1, 2, 3, \dots\} \rightarrow \{1, 2, 3, \dots\}$ with the above property.

Conjecture 3. Let $\mathbf{G}$ be an additive subgroup of $\mathbb{C}$. Let S be a consistent system of equations in $x_1, x_2, \dots, x_n \in \mathbf{G}$, where each equation in S is one of the following two forms: $x_i = 1$ or $x_i + x_j = x_k$. Then S has a solution $(x_1, x_2, \dots, x_n) \in (\mathbf{G} \cap \mathbb{Q})^n$ in which $|x_j| \leq 2^{n-1}$ for each j .

In case when $\mathbf{G} \supseteq \mathbb{Q}$ or $\mathbf{G} = \mathbb{Z}$, we will prove a weaker version of Conjecture 3 with the estimation given by $(\sqrt{5})^{n-1}$.

Observation 2. If $\mathcal{A} \subseteq \mathbb{C}^k$ is an affine subspace and $\text{card } \mathcal{A} > 1$, then there exists $m \in \{1, 2, \dots, k\}$ with

$$\emptyset \neq \mathcal{A} \cap \{(x_1, x_2, \dots, x_k) \in \mathbb{C}^k : x_m + x_m = x_m\} \subsetneq \mathcal{A}$$

Theorem 8. Let S be a consistent system of equations in complex numbers $x_1, x_2, \dots, x_n$, where each equation in S is one of the following two forms: $x_i = 1$ or $x_i + x_j = x_k$. Then S has a rational solution $(x_1, x_2, \dots, x_n)$ in which $|x_j| \leq (\sqrt{5})^{n-1}$ for each j .

Proof. We shall describe how to find a solution $(x_1, x_2, \dots, x_n) \in \mathbb{Q}^n$ in which $|x_j| \leq (\sqrt{5})^{n-1}$ for each j . We can assume that for a certain $i \in \{1, 2, \dots, n\}$ the equation $x_i = 1$ belongs to S , as otherwise $(0, 0, \dots, 0)$ is a solution. Without loss of generality we can assume that the equation $x_1 = 1$ belongs to S . Each equation belonging to S has a form

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = b,$$

where $a_1, a_2, \dots, a_n, b \in \mathbb{Z}$. Since $x_1 = 1$, we can equivalently write this equation as

$$a_2x_2 + a_3x_3 + \dots + a_nx_n = b - a_1$$

We receive a system of equations whose set of solutions is a non-empty affine subspace $\mathcal{A} \subseteq \mathbb{C}^{n-1}$. If $\text{card } \mathcal{A} > 1$, then by Observation 2 we find $m \in \{2, 3, \dots, n\}$ for which

$$\emptyset \neq \mathcal{A} \cap \{(x_2, x_3, \dots, x_n) \in \mathbb{C}^{n-1} : x_m + x_m = x_m\} \subsetneq \mathcal{A}$$

The procedure described in the last sentence is applied to the affine subspace

$$\mathcal{A} \cap \{(x_2, x_3, \dots, x_n) \in \mathbb{C}^{n-1} : x_m + x_m = x_m\}$$

and repeated until one point is achieved. The maximum number of procedure executions is $n - 1$. The received one-point affine subspace is described by equations belonging to a certain set

$$\mathcal{U} \subseteq \{x_i = 1 : i \in \{2, 3, \dots, n\}\} \cup \{x_i + x_j = x_k : i, j, k \in \{1, 2, \dots, n\}, i + j + k > 3\}$$

Each equation belonging to $\mathcal{U}$ has a form

$$a_2x_2 + a_3x_3 + \dots + a_nx_n = c,$$

where $a_2, a_3, \dots, a_n, c \in \mathbb{Z}$. Among these equations, we choose $n - 1$ linearly independent equations. Let $\mathbf{A}$ be the matrix of the system, and the system of equations has the following form

$$\mathbf{A} \cdot \begin{bmatrix} x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} c_2 \\ c_3 \\ \vdots \\ c_n \end{bmatrix}$$

Let $\mathbf{A}_j$ be the matrix formed by replacing the j -th column of $\mathbf{A}$ by the column vector $c_2, c_3, \dots, c_n$. Obviously, $\det(\mathbf{A}) \in \mathbb{Z}$, and $\det(\mathbf{A}_j) \in \mathbb{Z}$ for each $j \in \{1, 2, \dots, n - 1\}$. By Cramer's rule $x_j = \frac{\det(\mathbf{A}_{j-1})}{\det(\mathbf{A})} \in \mathbb{Q}$ for each $j \in \{2, 3, \dots, n\}$.

When the row of matrix $\mathbf{A}$ corresponds to the equation $x_i = 1$ ($i > 1$), then the entries in the row are 1, 0 ($n - 2$ times), while the right side of the equation is 1.

When the row of matrix $\mathbf{A}$ corresponds to the equation $x_1 + x_i = x_i$ ($i > 1$), then the entries in the row are 1, 0 ($n - 2$ times), while the right side of the equation is 2.

When the row of matrix $\mathbf{A}$ corresponds to one of the equations: $x_1 + x_i = x_1$ or $x_i + x_1 = x_1$ ($i > 1$), then the entries in the row are 1, 0 ($n - 2$ times), while the right side of the equation is 0.

When the row of matrix $\mathbf{A}$ corresponds to one of the equations: $x_1 + x_i = x_j$ or $x_i + x_1 = x_j$ ($i > 1, j > 1, i \neq j$), then the entries in the row are 1, -1 , 0 ($n - 3$ times), while the right side of the equation is 1.

When the row of matrix $\mathbf{A}$ corresponds to the equation $x_i + x_i = x_1$ ($i > 1$), then the entries in the row are 2, 0 ($n - 2$ times), while the right side of the equation is 1.

When the row of matrix $\mathbf{A}$ corresponds to the equation $x_i + x_j = x_1$ ($i > 1, j > 1, i \neq j$), then the entries in the row are 1, 1, 0 ($n - 3$ times), while the right side of the equation is 1.

From now on we assume that $i, j, k \in \{2, 3, \dots, n\}$.

When the row of matrix $\mathbf{A}$ corresponds to the equation $x_i + x_j = x_k$ ($i \neq j, i \neq k, j \neq k$), then the entries in the row are 1, 1, -1 , 0 ($n - 4$ times), while the right side of the equation is 0.

When the row of matrix $\mathbf{A}$ corresponds to the equation $x_i + x_i = x_k$ ($i \neq k$), then the entries in the row are 2, -1 , 0 ($n - 3$ times), while the right side of the equation is 0.

When the row of matrix $\mathbf{A}$ corresponds to the equation $x_i + x_j = x_k$ ($k = i$ or $k = j$),

then the entries in the row are 1, 0 ($n - 2$ times), while the right side of the equation is 0.

Contradictory equations, e.g. $x_1 + x_i = x_i$ do not belong to $\mathcal{U}$, therefore their description has been disregarded. The presented description shows that each row of matrix $\mathbf{A}_j$ ($j \in \{1, 2, \dots, n - 1\}$) has the length less than or equal to $\sqrt{5}$. By Hadamard's inequality $|\det(\mathbf{A}_j)| \leq (\sqrt{5})^{n-1}$ for each $j \in \{1, 2, \dots, n - 1\}$. Hence, $|x_j| = \frac{|\det(\mathbf{A}_{j-1})|}{|\det(\mathbf{A})|} \leq |\det(\mathbf{A}_{j-1})| \leq (\sqrt{5})^{n-1}$ for each $j \in \{2, 3, \dots, n\}$. □

Lemma 3 ([2]). Let $\mathbf{A}$ be a matrix with m rows, n columns, and integer entries. Let $b_1, b_2, \dots, b_m \in \mathbb{Z}$, and the matrix equation

$$\mathbf{A} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

defines the system of linear equations with rank m . Denote by δ the maximum of the absolute values of the $m \times m$ minors of the augmented matrix $(\mathbf{A}, b)$. We claim that if the system is consistent over $\mathbb{Z}$, then it has a solution in $(\mathbb{Z} \cap [-\delta, \delta])^n$.

Theorem 9. Let S be a consistent system of equations in integers $x_1, x_2, \dots, x_n$, where each equation in S is one of the following two forms: $x_i = 1$ or $x_i + x_j = x_k$. Then S has an integer solution $(x_1, x_2, \dots, x_n)$ in which $|x_j| \leq (\sqrt{5})^{n-1}$ for each j .

Proof. We shall describe how to find a solution $(x_1, x_2, \dots, x_n) \in \mathbb{Z}^n$ in which $|x_j| \leq (\sqrt{5})^{n-1}$ for each j . We can assume that for a certain $i \in \{1, 2, \dots, n\}$ the equation $x_i = 1$ belongs to S , as otherwise $(0, 0, \dots, 0)$ is a solution. Without loss of generality we can assume that the equation $x_1 = 1$ belongs to S . Analogously as in the proof of Theorem 8, we construct a system of linear equations with variables $x_2, \dots, x_n$. For the augmented matrix of this system, the length of each row is not greater than $\sqrt{5}$. We finish the proof by applying Hadamard's inequality and Lemma 3. □

References

- [1] S. BASU, R. POLLACK, M. F. ROY, *Algorithms in real algebraic geometry*, 2nd ed., Springer-Verlag, Berlin, 2006.

- [2] I. BOROSH, M. FLAHIVE, D. RUBIN, B. TREYBIG, *A sharp bound for solutions of linear Diophantine equations*, Proc. Amer. Math. Soc. 105 (1989), 844–846.
- [3] M. DAVIS, *An explicit Diophantine definition of the exponential function*, Comm. Pure Appl. Math. 24 (1971), 137–145.
- [4] M. DAVIS, *Hilbert’s tenth problem is unsolvable*, Amer. Math. Monthly 80 (1973), 233–269.
- [5] M. DAVIS, *Computability and unsolvability*, Dover Publications, New York, 1982.
- [6] J. DUGUNDJI, *An extension of Tietze’s theorem*, Pacific J. Math. 1 (1951), 353–367.
- [7] D. YU. GRIGOR’EV AND N. N. VOROBYOV, JR., *Solving systems of polynomial inequalities in subexponential time*, J. Symbolic Comput. 5 (1988), 37–64.
- [8] J. P. JONES AND YU. V. MATIJASEVIČ, *Proof of recursive unsolvability of Hilbert’s tenth problem*, Amer. Math. Monthly 98 (1991), 689–709.
- [9] G. LETTL, *Finitely arithmetically fixed elements of a field*, Arch. Math. (Basel) 87 (2006), 530–538.
- [10] YU. V. MATIJASEVIČ, *The Diophantineness of enumerable sets*, Soviet Math. Dokl. 11 (1970), 354–358.
- [11] YU. V. MATIJASEVIČ, *Hilbert’s tenth problem*, MIT Press, Cambridge, MA, 1993.
- [12] YU. V. MATIJASEVIČ AND J. ROBINSON, *Reduction of an arbitrary Diophantine equation to one in 13 unknowns*, Acta Arith. 27 (1975), 521–553.
- [13] W. SIERPIŃSKI, *Elementary theory of numbers*, 2nd ed. (ed. A. Schinzel), PWN (Polish Scientific Publishers) and North-Holland, Warsaw-Amsterdam, 1987.
- [14] A. TYŚKA, *A discrete form of the theorem that each field endomorphism of $\mathbb{R}(\mathbb{Q}_p)$ is the identity*, Aequationes Math. 71 (2006), 100–108.
- [15] A. TYŚKA, *On $\emptyset$ -definable elements in a field*, Collect. Math. 58 (2007), 73–84.
- [16] A. TYŚKA, *Arithmetic neighbourhoods of numbers*, Int. Math. Forum 3 (2008), no. 8, 349–362, to appear.

- [17] N. N. VOROBJOV, JR., *Estimates of real roots of a system of algebraic equations*,
J. Sov. Math. 34 (1986), 1754–1762.

Apoloniusz Tyszk

Technical Faculty

Hugo Kołłątaj University

Balicka 116B, 30-149 Kraków, Poland

E-mail: *rttyszka@cyf-kr.edu.pl*