BRAIDED BIALGEBRAS OF TYPE ONE

A. ARDIZZONI AND C. MENINI

ABSTRACT. Braided bialgebras of type one in abelian braided monoidal categories are characterized as braided graded bialgebras which are strongly \mathbb{N} -graded both as an algebra and as a coalgebra.

Contents

Introduction		1
1.	Preliminaries and Notations	6
2.	Graded and Strongly N-graded Coalgebras	Ę
3.	Graded and Strongly N-graded Algebras	13
4.	Braided Bialgebra Structure of the Cotensor Coalgebra	17
5.	Graded Braided Bialgebra Structure of the Tensor Algebra	20
6.	Braided Bialgebras of Type One	21
7.	Radford-Majid bosonization	25
References		28

Introduction

Bialgebras of type one were introduced by Nichols in [Ni]. They came out to play a relevant role in the theory of Hopf Algebras. In particular, their "coinvariant" part, called Nichols algebra, has been deeply investigated, see e.g. [Ro], [AS] and the references therein.

The main purpose of this paper is to study some relevant properties of bialgebras of type one in the framework of abelian braided monoidal categories.

Let H be a braided bialgebra in a cocomplete and complete abelian braided monoidal category (\mathcal{M},c) satisfying AB5. Assume that the tensor product commutes with direct sums and is two-sided exact. Let M be in ${}^H_H\mathcal{M}^H_H$. Let $T=T_H(M)$ be the relative tensor algebra and let $T^c=T^c_H(M)$ be the relative cotensor coalgebra as introduced in [AMS1]. We prove that both T and T^c have a natural structure of graded braided bialgebra and that the natural algebra morphism $F:T\to T^c$, which coincide with the canonical injections on H and M, is a graded bialgebra homomorphism. Thus its image is a graded braided bialgebra which we denote by H[M] and call, accordingly to [Ni], the braided bialgebra of type one associated to H and M.

We would like to outline that a treatment of bialgebras of type one in the case H=1 can be found in [Sch]. Following a different approach, that traces back to some Nichols' original ideas, in order to characterize braided bialgebras of type one, we develop some relevant properties of graded algebras and coalgebras. We introduce the definition of strongly N-graded algebras and coalgebras which is inspired in the second case to [NT].

Given a graded coalgebra $(C = \bigoplus_{n \in \mathbb{N}} C_n, \Delta, \varepsilon)$ in \mathcal{M} , we can write $\Delta_{|C_n}$ as the sum of unique components $\Delta_{i,j} : C_{i+j} \to C_i \otimes C_j$ where i+j=n. The coalgebra C is defined to be a *strongly*

¹⁹⁹¹ $Mathematics\ Subject\ Classification.$ Primary 18D10; Secondary 16W30.

 $[\]it Key\ words\ and\ phrases.$ Braided bialgebras, Monoidal categories, Cotensor Coalgebras.

This paper was written while both authors were members of G.N.S.A.G.A. with partial financial support from M.I.U.R..

 \mathbb{N} -graded coalgebra when $\Delta_{i,j}^C: C_{i+j} \to C_i \otimes C_j$ is a monomorphism for every $i, j \in \mathbb{N}$. In Theorem 2.22, we prove that the following assertions are equivalent.

- (a) C is a strongly \mathbb{N} -graded coalgebra.
- (b) the canonical morphism $\psi: C \to T^c_{C_0}(C_1)$ is a monomorphism.
- (c) $C_0 \oplus C_1 = C_0 \wedge_C C_0$.

Also the "dual" notion of $strongly \ \mathbb{N}$ -graded algebra is introduced and a similar result is achieved. Braided bialgebras of type one are then characterized as braided graded bialgebras which are strongly \mathbb{N} -graded both as an algebra and as a coalgebra.

Assume that H is a braided Hopf algebra in (\mathcal{M}, c) with bijective antipode. Using general results by Bespalov and Drabant (see [Bes] and [BD]), we prove that H[M] is the Radford-Majid bosonization

$$H[M] = \mathbf{1} \left[M^{co(H)} \right] \rtimes H$$

where $\mathbf{1}\left[M^{co(H)}\right]$ is the braided bialgebra of type one associated to $\mathbf{1}$ and $M^{co(H)}$ in ${}^{H}_{H}\mathcal{YD}\left(\mathcal{M}\right)$.

Finally we would like to outline that many results are firstly stated and proved in the coalgebra case. Our choice is motivated by the special feature of this case, due to the lack of the existence of the coradical in the monoidal context. As far as the algebra case is concerned, we would like to point out that here we can drop the assumptions of completeness and AB5-condition essentially because the analogue of Lemma 2.2 doesn't require these extra conditions. Also proofs in the algebra case are not given whenever they would have been an easy adaptation of the coalgebra ones.

Elsewhere we will apply the results of the present paper to study strictly graded bialgebras and graded bialgebras associated to a (co)algebra homomorphism.

1. Preliminaries and Notations

Let $[(X, i_X)]$ be a subobject of an object E in an abelian category C, where $i_X = i_X^E : X \hookrightarrow E$ is a monomorphism and $[(X, i_X)]$ is the associated equivalence class. By abuse of language, we will say that (X, i_X) is a subobject of E and we will write $(X, i_X) = (Y, i_Y)$ to mean that $(Y, i_Y) \in [(X, i_X)]$. The same convention applies to cokernels. If (X, i_X) is a subobject of E then we will write $(E/X, p_X) = \operatorname{Coker}(i_X)$, where $p_X = p_X^E : E \to E/X$.

1.1. **Monoidal Categories.** Recall that (see [Ka, Chap. XI]) a monoidal category is a category \mathcal{M} endowed with an object $\mathbf{1} \in \mathcal{M}$ (called unit), a functor $\otimes : \mathcal{M} \times \mathcal{M} \to \mathcal{M}$ (called tensor product), and functorial isomorphisms $a_{X,Y,Z} : (X \otimes Y) \otimes Z \to X \otimes (Y \otimes Z), l_X : \mathbf{1} \otimes X \to X, r_X : X \otimes \mathbf{1} \to X$, for every X,Y,Z in \mathcal{M} . The functorial morphism a is called the associativity constraint and satisfies the Pentagon Axiom, that is the following relation

$$(U \otimes a_{V,W,X}) \circ a_{U,V \otimes W,X} \circ (a_{U,V,W} \otimes X) = a_{U,V,W \otimes X} \circ a_{U \otimes V,W,X}$$

holds true, for every U, V, W, X in \mathcal{M} . The morphisms l and r are called the *unit constraints* and they obey the *Triangle Axiom*, that is $(V \otimes l_W) \circ a_{V,1,W} = r_V \otimes W$, for every V, W in \mathcal{M} .

It is well known that the Pentagon Axiom completely solves the consistency problem arising out of the possibility of going from $((U \otimes V) \otimes W) \otimes X$ to $U \otimes (V \otimes (W \otimes X))$ in two different ways (see [Mj1, page 420]). This allows the notation $X_1 \otimes \cdots \otimes X_n$ forgetting the brackets for any object obtained from $X_1, \cdots X_n$ using \otimes . Also, as a consequence of the coherence theorem, the constraints take care of themselves and can then be omitted in any computation involving morphisms in \mathcal{M} .

Thus, for sake of simplicity, from now on, we will omit the associativity constraints.

The notions of algebra, module over an algebra, coalgebra and comodule over a coalgebra can be introduced in the general setting of monoidal categories. Given an algebra A in \mathcal{M} on can define the categories ${}_{A}\mathcal{M}$, \mathcal{M}_{A} and ${}_{A}\mathcal{M}_{A}$ of left, right and two-sided modules over A respectively. Similarly, given a coalgebra C in \mathcal{M} , one can define the categories of C-comodules ${}^{C}\mathcal{M}$, \mathcal{M}^{C} , ${}^{C}\mathcal{M}^{C}$. For more details, the reader is referred to [AMS2].

DEFINITIONS 1.2. Let \mathcal{M} be a monoidal category.

We say that \mathcal{M} is an **abelian monoidal category** if \mathcal{M} is abelian and both the functors $X \otimes (-)$: $\mathcal{M} \to \mathcal{M}$ and $(-) \otimes X : \mathcal{M} \to \mathcal{M}$ are additive and right exact, for any $X \in \mathcal{M}$.

We say that \mathcal{M} is an **coabelian monoidal category** if \mathcal{M}^o is an abelian monoidal category, where \mathcal{M}^o denotes the dual monoidal category of \mathcal{M} . Recall that \mathcal{M}^o and \mathcal{M} have the same objects but $\mathcal{M}^o(X,Y) = \mathcal{M}(Y,X)$ for any X,Y in \mathcal{M} .

Given an algebra A in an abelian monoidal category \mathcal{M} , there exist a suitable functor \otimes_A : ${}_A\mathcal{M}_A\times{}_A\mathcal{M}_A\to{}_A\mathcal{M}_A$ and constraints that make the category $({}_A\mathcal{M}_A,\otimes_A,A)$ abelian monoidal, see [AMS2, 1.11]. The tensor product over A in \mathcal{M} of a right A-module V and a left A-module W is defined to be the coequalizer:

$$(V \otimes A) \otimes W \xrightarrow{AXV,W} V \otimes_A W \otimes_A W \xrightarrow{AXV,W} V \otimes_A W \otimes_A$$

Note that, since \otimes preserves coequalizers, then $V \otimes_A W$ is also an A-bimodule, whenever V and W are A-bimodules.

Dually, given a coalgebra (C, Δ, ε) in a coabelian monoidal category \mathcal{M} , there exist a suitable functor $\Box_C : {}^C \mathcal{M}^C \times {}^C \mathcal{M}^C \to {}^C \mathcal{M}^C$ and constraints that make the category $({}^C \mathcal{M}^C, \Box_C, C)$ coabelian monoidal. The cotensor product over C in \mathcal{M} of a right C-bicomodule V and a left C-comodule W is defined to be the equalizer:

$$0 \longrightarrow V \square_C W \xrightarrow{C \leq V, W} V \otimes W \Longrightarrow V \otimes (C \otimes W)$$

Note that, since \otimes preserves equalizers, then $V \square_C W$ is also a C-bicomodule, whenever V and W are C-bicomodules.

1.3. **Graded Objects.** Let $(X_n)_{n\in\mathbb{N}}$ be a sequence of objects a cocomplete abelian monoidal category \mathcal{M} and let

$$X = \bigoplus_{n \in \mathbb{N}} X_n$$

be their coproduct in \mathcal{M} . In this case we also say that X is a graded object of \mathcal{M} and that the sequence $(X_n)_{n\in\mathbb{N}}$ defines a graduation on X. A morphism

$$f: X = \bigoplus_{n \in \mathbb{N}} X_n \to Y = \bigoplus_{n \in \mathbb{N}} Y_n$$

is called a graded homomorphism whenever there exists a family of morphisms $(f_n: X_n \to Y_n)_{n \in \mathbb{N}}$ such that $f = \bigoplus_{n \in \mathbb{N}} f_n$ i.e. such that

$$f \circ i_{X_n}^X = i_{Y_n}^Y \circ f_n$$
, for every $n \in \mathbb{N}$.

We fix the following notations:

$$X\left(n\right) = \left\{ \begin{array}{ll} 0, & \text{for } n=0 \\ \underset{0 \leq i \leq n-1}{\oplus} X_i, & \text{for } n \geq 1 \end{array} \right. \quad \text{and} \quad X\left[n\right] = \underset{i \geq n}{\oplus} X_i, \text{ for } n \in \mathbb{N}.$$

Let $\sigma_i^{i+1}: X(i) \to X(i+1)$ be the canonical inclusion and for any j > i, define:

$$\sigma_{i}^{j} = \sigma_{j-1}^{j} \sigma_{j-2}^{j-1} \cdots \sigma_{i+1}^{i+2} \sigma_{i}^{i+1} : X(i) \to X(j)$$

Throughout let

$$\pi_n^m : X(n) \to X(m) \ (m \le n), \qquad \pi_n : X \to X(n),$$

$$p_n^m : X(n) \to X_m \ (m < n), \qquad p_n : X \to X_n,$$

be the canonical projections and let

$$\begin{split} \sigma_{m}^{n} & : & X\left(m\right) \to X\left(n\right) \ \left(m \leq n\right), \qquad \sigma_{n} : X\left(n\right) \to X, \\ i_{m}^{n} & : & X_{m} \to X\left(n\right) \ \left(m < n\right), \qquad i_{n} : X_{n} \to X, \end{split}$$

be the canonical injections for any $m, n \in \mathbb{N}$. For technical reasons we set $\pi_n^m = 0$, $\sigma_m^n = 0$ for any n < m and $p_n^m = 0$, $i_m^n = 0$ for any $n \le m$. Then, we have the following relations:

$$p_n \sigma_k = p_k^n$$
, $p_n i_k = \delta_{n,k} \operatorname{Id}_{X_k}$, $\pi_n i_k = i_k^n$.

Moreover, we have:

$$\begin{array}{lll} \pi_n^m\sigma_k^n=\sigma_k^m, \text{ if } k\leq m\leq n, & \text{and} & \pi_n^m\sigma_k^n=\pi_k^m, \text{ if } m\leq k\leq n, \\ p_n^m\pi_k^n=p_k^m, \text{ if } m< n\leq k, & \text{and} & \sigma_n^mi_k^n=i_k^m, \text{ if } k< n\leq m, \\ p_n^m\sigma_k^n=p_k^m, \text{ if } m< k\leq n, & \text{and} & \pi_n^mi_k^n=i_k^m, \text{ if } k< m\leq n, \\ p_n^m\pi_n=p_m, \text{ if } m< n, & \text{and} & \sigma_ni_m^n=i_m, \text{ if } m< n, \\ \pi_n\sigma_k=\sigma_k^n, \text{ if } k\leq n, & \text{and} & \sigma_ni_m^n=i_m, \text{ if } m< n, \\ p_n^mi_m^n=\mathrm{Id}_{X_m}, \text{ if } m< n. & & & & & \\ \end{array}$$

In the other cases, these compositions are zero. Similarly let

$$\tau_n^m : X[n] \to X[m] \ (n \le m), \qquad \tau_n : X \to X[n],$$

 $\zeta_n^m : X[n] \to X_m \ (n \le m),$

be the canonical projections and let

$$u_m^n : X[m] \to X[n] \ (n \le m), \qquad \nu_n : X[n] \to X,$$

$$\lambda_m^n : X_m \to X[n] \ (n \le m),$$

be the canonical injection for any $m,n\in\mathbb{N}$. For technical reasons we set $\tau_n^m=0,\ \nu_m^n=0$ for any m< n and $\zeta_n^m=0,\ \lambda_m^n=0$ for any m< n. Then, we have the following relations:

$$p_n \nu_k = \zeta_k^n, \qquad \tau_n i_k = \lambda_k^n.$$

Moreover, we have:

$$\begin{array}{lll} \tau_n^m \nu_k^n = \nu_k^m, \text{ if } n \leq m \leq k, & \text{and} & \tau_n^m \nu_k^n = \tau_k^m, \text{ if } n \leq k \leq m, \\ \zeta_n^m \tau_n^n = \zeta_k^m, \text{ if } k \leq n \leq m, & \text{and} & \nu_n^m \lambda_k^n = \lambda_k^m, \text{ if } m \leq n \leq k, \\ \zeta_n^m \nu_k^n = \zeta_k^m, \text{ if } n \leq k \leq m, & \text{and} & \tau_n^m \lambda_k^n = \lambda_k^m, \text{ if } n \leq m \leq k, \\ \zeta_n^m \tau_n = p_m, \text{ if } n \leq m, & \text{and} & \nu_n \lambda_m^n = i_m, \text{ if } n \leq m, \\ \tau_n \nu_k = \nu_k^n, \text{ if } n \leq k, & \text{and} & \tau_n \nu_k = \tau_k^n, \text{ if } k \leq n, \\ \zeta_n^m \lambda_n^m = \text{Id}_{X_m}, \text{ if } n \leq m. & & & & & \\ \end{array}$$

In the other cases, these compositions are zero. Given graded objects X, Y in \mathcal{M} we set

$$(X \otimes Y)_n = \bigoplus_{a+b=n} (X_a \otimes Y_b).$$

Then this defines a graduation on $X \otimes Y$ whenever the tensor product commutes with direct sums. We denote by

$$X_a \otimes Y_b \overset{\gamma^{X,Y}_{a,b}}{\overset{\sim}{\rightarrow}} (X \otimes Y)_{a+b}$$
 and $(X \otimes Y)_{a+b} \overset{\omega^{X,Y}_{a,b}}{\overset{\sim}{\rightarrow}} X_a \otimes Y_b$

the canonical injection and projection respectively. We have

(1)
$$\sum_{a+b=n} \left(i_a^X \otimes i_b^Y \right) \omega_{a,b}^{X,Y} = \nabla \left[\left(i_a^X \otimes i_b^Y \right)_{a+b=n} \right]$$

(2)
$$\sum_{a,b=n} \gamma_{a,b}^{X,Y} \left(p_a^X \otimes p_b^Y \right) = \Delta \left[\left(p_a^X \otimes p_b^Y \right)_{a+b=n} \right]$$

where $\nabla \left[\left(i_a^X \otimes i_b^Y \right)_{a+b=n} \right]$ denotes the codiagonal morphism associated to the family $\left(i_a^X \otimes i_b^Y \right)_{a+b=n}$ and $\Delta \left[\left(p_a^X \otimes p_b^Y \right)_{a+b=n} \right]$ denotes the diagonal morphism associated to the family $\left(p_a^X \otimes p_b^Y \right)_{a+b=n}$.

LEMMA 1.4. Let $X = \bigoplus_{n \in \mathbb{N}} X_n$ be a graded object in a cocomplete abelian category \mathcal{M} . Then, for every $n \in \mathbb{N}$

$$0 \to X[n] \xrightarrow{\nu_n} X \xrightarrow{\pi_n} X(n) \to 0$$
 and $0 \to X(n) \xrightarrow{\sigma_n} X \xrightarrow{\tau_n} X[n] \to 0$

are splitting exact sequences.

2. Graded and Strongly N-graded Coalgebras

In this section $(\mathcal{M}, \otimes, \mathbf{1})$ will denote a cocomplete coabelian monoidal category such that the tensor product commutes with direct sums.

2.1. Recall that a graded coalgebra in \mathcal{M} is a coalgebra (C, Δ, ε) where

$$C = \bigoplus_{n \in \mathbb{N}} C_n$$

is a graded object of \mathcal{M} such that $\Delta: C \to C \otimes C$ is a graded homomorphism i.e. there exists a family $(\Delta_n)_{n \in \mathbb{N}}$ of morphisms

$$\Delta_n^C = \Delta_n : C_n \to (C \otimes C)_n = \bigoplus_{a+b=n} (C_a \otimes C_b) \text{ such that } \Delta = \bigoplus_{n \in \mathbb{N}} \Delta_n.$$

We set

$$\Delta_{a,b}^{C} = \Delta_{a,b} := \left(C_{a+b} \stackrel{\Delta_{a+b}}{\to} (C \otimes C)_{a+b} \stackrel{\omega_{a,b}^{C,C}}{\to} C_{a} \otimes C_{b} \right).$$

A homomorphism $f:(C,\Delta_C,\varepsilon_C)\to(D,\Delta_D,\varepsilon_D)$ of coalgebras is a graded coalgebra homomorphism if it is a graded homomorphism too.

LEMMA 2.2. Let I be a set and let $(X_i)_{i\in I}$ be a family of objects in a cocomplete and complete abelian category C satisfying AB5. Let Y be an object in C and let $f: Y \to \bigoplus_{i\in I} X_i$ be a morphism such that

$$p_k f = 0$$
 for any $k \in I$,

where $p_k: \bigoplus_{i\in I} X_i \to X_k$ denotes the canonical projection. Then f=0.

PROPOSITION 2.3. Let (C, Δ, ε) be a coalgebra in \mathcal{M} which is a graded object with graduation defined by $(C_k)_{k\in\mathbb{N}}$. Fix $n\in\mathbb{N}$. Consider the following assertions.

(i) There exists a morphism

$$\Delta_n: C_n \to (C \otimes C)_n = \bigoplus_{a+b=n} (C_a \otimes C_b) \text{ such that } \Delta i_n = \nabla \left[(i_a \otimes i_b)_{a+b=n} \right] \Delta_n.$$

(ii) There exists a family $(\Delta_{a,b})_{a+b=n}$ of morphisms $\Delta_{a,b}: C_n \to C_a \otimes C_b$ such that

(3)
$$\Delta i_n = \sum_{a+b=n} (i_a \otimes i_b) \, \Delta_{a,b}.$$

(iii) $(p_a \otimes p_b)\Delta i_n = 0$, for every $a, b \in \mathbb{N}$, $a + b \neq n$.

Then $(i) \Leftrightarrow (ii) \Rightarrow (iii)$. Furthermore, if \mathcal{M} is also complete and satisfies AB5, then $(ii) \Leftrightarrow (iii)$.

Proof. $(i) \Rightarrow (ii)$ Set

$$\Delta_{a,b} = \left(C_n \stackrel{\Delta_n}{\to} (C \otimes C)_n \stackrel{\omega_{a,b}^{C,C}}{\to} C_a \otimes C_b \right)$$

for every $a, b \in \mathbb{N}$ such that a + b = n. We have

$$\sum_{a+b=n} \left(i_a \otimes i_b\right) \Delta_{a,b} = \sum_{a+b=n} \left(i_a \otimes i_b\right) \omega_{a,b}^{C,C} \Delta_n \stackrel{(1)}{=} \nabla \left[\left(i_a \otimes i_b\right)_{a+b=n} \right] \Delta_n = \Delta i_n.$$

 $(ii) \Rightarrow (iii)$ We have

(4)
$$(p_a \otimes p_b) \sum_{u+v=n} (i_u \otimes i_v) \Delta_{u,v} = \delta_{a+b,n} \Delta_{a,b} = \Delta_{a,b} p_{a+b} i_n,$$

so that, in view of (3), we obtain

$$(p_a \otimes p_b) \Delta i_n = \delta_{a+b,n} \Delta_{a,b}$$
, for every $a, b \in \mathbb{N}$.

 $(ii) \Rightarrow (i)$ Set

$$\Delta_n = \sum_{a+b=n} \left(C_n \stackrel{\Delta_{a,b}}{\to} C_a \otimes C_b \stackrel{\gamma_{a,b}^{C,C}}{\to} (C \otimes C)_n \right).$$

Then

$$\nabla \left[(i_u \otimes i_v)_{u+v=n} \right] \Delta_n = \sum_{a+b=n} \nabla \left[(i_u \otimes i_v)_{u+v=n} \right] \gamma_{a,b}^{C,C} \Delta_{a,b} = \sum_{a+b=n} (i_a \otimes i_b) \Delta_{a,b} = \Delta i_n.$$

Assume \mathcal{M} is also complete and satisfies AB5.

 $(iii) \Rightarrow (ii)$ Set $\Delta_{a,b} = (p_a \otimes p_b)\Delta i_{a+b}$ for every $a,b \in \mathbb{N}$. For every $u,v \in \mathbb{N}$, we have

$$(p_u \otimes p_v) \sum_{a+b-n} (i_a \otimes i_b) \Delta_{a,b} \stackrel{(4)}{=} \delta_{u+v,n} \Delta_{u,v} = (p_u \otimes p_v) \Delta i_n.$$

since the codomain of Δi_n is $C \otimes C = \bigoplus_{u,v \in \mathbb{N}} C_u \otimes C_v$ and \mathcal{M} satisfies AB5, by Lemma 2.2, we conclude.

PROPOSITION 2.4. 1) If $(C = \bigoplus_{n \in \mathbb{N}} C_n, \Delta, \varepsilon)$ is a graded coalgebra in \mathcal{M} , then

(5)
$$(p_a \otimes p_b)\Delta = \Delta_{a,b}p_n$$
, for every $a, b \in \mathbb{N}$.

2) Assume that \mathcal{M} is also complete and satisfies AB5. Let (C, Δ, ε) be a coalgebra in \mathcal{M} which is a graded object with graduation defined by $(C_n)_{n\in\mathbb{N}}$. If there exists a family $(\Delta_{a,b})_{a,b\in\mathbb{N}}$ of morphisms $\Delta_{a,b}: C_{a+b} \to C_a \otimes C_b$ such that (5) holds, then $(C = \bigoplus_{n\in\mathbb{N}} C_n, \Delta, \varepsilon)$ is a graded coalgebra in \mathcal{M} .

Proof. 1) For every $a, b, t \in \mathbb{N}$ we have

$$(p_a \otimes p_b) \Delta i_t \stackrel{\text{(3)}}{=} (p_a \otimes p_b) \sum_{u+v=n} (i_u \otimes i_v) \Delta_{u,v} \stackrel{\text{(4)}}{=} \Delta_{a,b} p_{a+b} i_n.$$

Since this equality holds for an arbitrary $n \in \mathbb{N}$, we get (5).

2) By (5) we have $(p_a \otimes p_b)\Delta i_n = 0$, for every $a, b \in \mathbb{N}, a+b \neq n$. Thus, by Proposition 2.3, C is a graded coalgebra.

PROPOSITION 2.5. 1) Let $C = \bigoplus_{n \in \mathbb{N}} C_n$ be a graded object of \mathcal{M} such that there exists a family $(\Delta_{a,b})_{a,b \in \mathbb{N}}$

$$\Delta_{a,b}: C_{a+b} \to C_a \otimes C_b$$
,

of morphisms and a morphism $\varepsilon_0: C_0 \to \mathbf{1}$ which satisfy

(6)
$$(\Delta_{a,b} \otimes C_c) \, \Delta_{a+b,c} = (C_a \otimes \Delta_{b,c}) \, \Delta_{a,b+c},$$

(7)
$$(C_d \otimes \varepsilon_0) \Delta_{d,0} = r_{C_d}^{-1}, \qquad (\varepsilon_0 \otimes C_d) \Delta_{0,d} = l_{C_d}^{-1},$$

for every $a, b, c \in \mathbb{N}$. Then there exists a unique morphism $\Delta : C \to C \otimes C$ such that (3) holds. Moreover $(C = \bigoplus_{n \in \mathbb{N}} C_n, \Delta, \varepsilon = \varepsilon_0 p_0)$ is a graded coalgebra.

2) If C is a graded coalgebra, then $\varepsilon = \varepsilon i_0 p_0$ so that ε is a graded homomorphism, and we have that (6) and (7) hold for every $a, b, c \in \mathbb{N}$, where $\varepsilon_0 = \varepsilon i_0$.

Moreover $(C_0, \Delta_0 = \Delta_{0,0}, \varepsilon_0 = \varepsilon i_0)$ is a coalgebra in \mathcal{M} , i_0 is a coalgebra homomorphism and, for every $n \in \mathbb{N}$, $(C_n, \Delta_{0,n}, \Delta_{n,0})$ is a C_0 -bicomodule such that $p_n : C \to C_n$ is a morphism of C_0 -bicomodules $(C \text{ is a } C_0\text{-bicomodule through } p_0)$.

Proof. 1) Since the tensor product commutes with direct sums, we have that $C \otimes C = \bigoplus_{a,b \in \mathbb{N}} (C_a \otimes C_b)$. Thus, by the universal property of coproduct there exists a unique morphism $\Delta : C \to C \otimes C$ such that (3) holds. By using (3) and (6) we get $(\Delta \otimes C) \Delta i_n = (C \otimes \Delta) \Delta i_n$, for every $n \in \mathbb{N}$ and hence Δ is coassociative. Similarly, by using (3) and (7), we get $(C \otimes \varepsilon) \Delta = r_C^{-1}$ and $(\varepsilon \otimes C) \Delta = l_C^{-1}$. In view of (3), by Proposition 2.3, C is a graded coalgebra.

2) By using (5), we obtain

$$(p_a \otimes p_b \otimes p_c) (\Delta \otimes C) \Delta i_{a+b+c} = (\Delta_{a,b} \otimes C_c) \Delta_{a+b,c}$$

$$(p_a \otimes p_b \otimes p_c) (C \otimes \Delta) \Delta i_{a+b+c} = (C_a \otimes \Delta_{b,c}) \Delta_{a,b+c}$$

so that (6) holds true.

From $(C \otimes \varepsilon_C) \Delta = r_C^{-1}$, we deduce

$$(p_b \otimes \mathbf{1}) (C \otimes \varepsilon_C) \Delta i_a = (p_b \otimes \mathbf{1}) r_C^{-1} i_a.$$

We compute the first term:

$$(p_b \otimes \mathbf{1}) (C \otimes \varepsilon) \Delta i_a \stackrel{(3)}{=} (p_b \otimes \mathbf{1}) (C \otimes \varepsilon) \sum_{u+v=a} (i_u \otimes i_v) \Delta_{u,v} = (C_b \otimes \varepsilon i_{a-b}) \Delta_{b,a-b}$$

and the second one:

$$(p_b \otimes \mathbf{1}) r_C^{-1} i_a = r_{C_b}^{-1} p_b i_a = \delta_{b,a} r_{C_b}^{-1}.$$

Thus we obtain

$$(C_b \otimes \varepsilon i_{a-b}) \, \Delta_{b,a-b} = \delta_{b,a} r_{C_b}^{-1}.$$

Similarly one gets $(\varepsilon_C i_{a-b} \otimes C_b) \Delta_{a-b,b} = \delta_{a,b} l_{C_b}^{-1}$. In particular one has (7).

Let us prove that ε is a graded homomorphism. Let us check that

$$\varepsilon = \varepsilon i_0 p_0$$
.

Set $\varepsilon' := \varepsilon i_0 p_0$. Then

$$r_{C}\left(A\otimes\varepsilon'\right)\Delta i_{n}\overset{(3)}{=}r_{C}\left(A\otimes\varepsilon i_{0}p_{0}\right)\sum_{a+b=n}\left(i_{a}\otimes i_{b}\right)\Delta_{a,b}=r_{C}\left(i_{n}\otimes\varepsilon i_{0}\right)\Delta_{n,0}\overset{(7)}{=}r_{C}\left(i_{n}\otimes\mathbf{1}\right)r_{C_{n}}^{-1}=i_{n}.$$

Since this holds for every $n \in \mathbb{N}$, we deduce that $r_C(A \otimes \varepsilon') \Delta = \operatorname{Id}_C$ so that ε' is a right counit for C. Since ε is a counit for C, one gets $\varepsilon = \varepsilon'$.

Hence ε is a graded homomorphism.

By applying (6) and (7) to the cases

$$(a, b, c) = (0, 0, 0), d = 0,$$

we get that $(C_0, \Delta_0 = \Delta_{0,0}, \varepsilon_0 = \varepsilon i_0)$ is a coalgebra in \mathcal{M} .

By applying (6) and (7) to the cases

$$(a,b,c) = (n,0,0), (0,n,0), (0,0,n), d = n,$$

we get that $(C_n, \Delta_{0,n}, \Delta_{n,0})$ is a C_0 -bicomodule for every $n \in \mathbb{N}$. By applying (5) to the cases (a,b) = (n,0) and (a,b) = (0,n), we get the last assertion.

LEMMA 2.6. Let $f:(C,\Delta_C,\varepsilon_C)\to (D,\Delta_D,\varepsilon_D)$ be a homomorphism of graded coalgebras in \mathcal{M} . Then

(8)
$$\Delta_{a,b}^{D} \circ f_{a+b} = (f_a \otimes f_b) \circ \Delta_{a,b}^{C}, \text{ for every } n, a, b \in \mathbb{N}.$$

Proof. By definition of graded homomorphism, $f \circ i_t^C = i_t^D \circ f$ and hence $p_n^D \circ f \circ i_t^C = f \circ p_n^C \circ i_t^C$ for every $n, t \in \mathbb{N}$. Therefore $p_n^D \circ f = f_n \circ p_n^C$ for every $n \in \mathbb{N}$. Using this relation and (5), for every $a, b \in \mathbb{N}$, the equality

$$(p_a^D \otimes p_b^D) \circ \Delta_D \circ f \circ i_{a+b}^C = (p_a^D \otimes p_b^D) \circ (f \otimes f) \circ \Delta_C \circ i_{a+b}^C$$

rewrites as $\Delta_{a,b}^D \circ f_{a+b} = (f_a \otimes f_b) \circ \Delta_{a,b}^C$.

2.7. Let C be a coalgebra in \mathcal{M} . As in the case of vector spaces, we can introduce the wedge product of two subobjects X, Y of C in \mathcal{M} :

$$(X \wedge_C Y, i_{X \wedge Y}^C) := Ker[(p_X \otimes p_Y) \circ \triangle_C],$$

where $p_X: C \to C/X$ and $p_Y: C \to C/Y$ are the canonical quotient maps. In particular we have the following exact sequence:

$$0 \longrightarrow X \wedge_C Y \xrightarrow{i_{X \wedge Y}^C} C \xrightarrow{(p_X \otimes p_Y) \circ \triangle_C} C/X \otimes C/Y.$$

Let $\delta: D \to C$ be a monomorphism which is a homomorphism of coalgebras in \mathcal{M} . Denote by (L, p) the cokernel of δ in \mathcal{M} . Regard D as a C-bicomodule via δ and observe that L is a C-bicomodule and p is a morphism of bicomodules. Let

$$(D^{\wedge_C^n}, \delta_n) := \ker(p^{\otimes n} \Delta_C^{n-1})$$

for any $n \in \mathbb{N} \setminus \{0\}$. Note that $(D^{\wedge_C^1}, \delta_1) = (D, \delta)$ and $(D^{\wedge_C^2}, \delta_2) = D \wedge_C D$.

In order to simplify the notations we set $(D^{\wedge_C^0}, \delta_0) = (0, 0)$.

Now, since \mathcal{M} has left exact tensor functors and since $p^{\otimes n}\Delta_C^{n-1}$ is a morphism of C-bicomodules (as a composition of morphisms of C-bicomodules), we get that $D^{\wedge_C^n}$ is a coalgebra and $\delta_n: D^{\wedge_C^n} \to C$ is a coalgebra homomorphism for any n > 0 and hence for any $n \in \mathbb{N}$.

PROPOSITION 2.8. [AMS1, Proposition 2.17] Let $\delta: D \to C$ be a monomorphism which is a coalgebra homomorphism in \mathcal{M} . Then we have

$$(9) D^{\wedge_C^m} \wedge_C D^{\wedge_C^n} = D^{\wedge_C^{m+n}}.$$

DEFINITION 2.9. Let $(C = \bigoplus_{n \in \mathbb{N}} C_n, \Delta, \varepsilon)$ be a graded coalgebra in \mathcal{M} . In analogy with the group graded case (see [NT]), we say that C is a *strongly* \mathbb{N} -graded coalgebra whenever

 $\Delta_{i,j}^C: C_{i+j} \to C_i \otimes C_j$ is a monomorphism for every $i, j \in \mathbb{N}$, where $\Delta_{i,j}^C$ is the morphism defined in Definition 2.1.

THEOREM 2.10. Let $(C = \bigoplus_{n \in \mathbb{N}} C_n, \Delta, \varepsilon)$ be a graded coalgebra in a cocomplete and complete coabelian monoidal category \mathcal{M} satisfying AB5. Then:

1) Let $n \geq 2$ be such that

(10)
$$\left\{ \begin{array}{c} \text{there exist } 0 \leq a, b \leq n-1 \text{ such that } a+b=n \\ \text{and } \Delta_{u,v} : C_{u+v} \to C_u \otimes C_v \text{ is a monomorphism for every } a \leq u \text{ and } b \leq v \end{array} \right\}$$

$$0 \to C\left(n\right) \xrightarrow{\sigma_{n}} C \xrightarrow{(\tau_{a} \otimes \tau_{b})^{\Delta}} C\left[a\right] \otimes C\left[b\right]$$

is an exact sequence and

$$C(n) = C(a) \wedge_C C(b)$$
.

2) Assume that C is a strongly \mathbb{N} -graded coalgebra. Then

$$(C(n), \sigma_n) = C_0^{\wedge_C^n}, \text{ for every } n \in \mathbb{N}.$$

Proof. 1) Let $n \geq 2$ be such that condition (10) holds true. Let us prove that the following sequence

$$0 \to C(n) \xrightarrow{\sigma_n} C \xrightarrow{(\tau_a \otimes \tau_b)\Delta} C[a] \otimes C[b]$$

is exact. For every $0 \le t \le n-1$, we compute

$$(\tau_a \otimes \tau_b) \, \Delta \sigma_n i_t^n = (\tau_a \otimes \tau_b) \, \Delta i_t \stackrel{(3)}{=} \sum_{u+v=t} (\tau_a i_u \otimes \tau_b i_v) \, \Delta_{u,v} = \sum_{u+v=t} (\lambda_u^a \otimes \lambda_v^b) \, \Delta_{u,v} = 0.$$

Let $g: Y \to C$ be a morphism such that $(\tau_a \otimes \tau_b) \Delta g = 0$. Now, for every $a \leq u$ and $b \leq v$, we have

$$0 = (\zeta_a^u \tau_a \otimes \zeta_b^v \tau_b) \Delta g = (p_u \otimes p_v) \Delta g \stackrel{(5)}{=} \Delta_{u,v} p_{u+v} g.$$

By hypothesis $\Delta_{u,v}$ is a monomorphism so that $p_{u+v}g = 0$. We conclude that

$$p_c g = 0$$
, for every $c \ge n$.

Set $\overline{g} = \pi_n g$ and let us prove that $g = \sigma_n \overline{g}$. By Lemma 2.2 this is the case if and only if

$$p_c g = p_c \sigma_n \overline{g}$$
, for every $c \in \mathbb{N}$.

We calculate

$$p_c \sigma_n \overline{g} = p_n^c \overline{g} = p_n^c \pi_n g = \begin{cases} p_c g & \text{for every } c < n \\ 0 = p_c g & \text{otherwise.} \end{cases}$$

Thus the required sequence is exact.

By Lemma 1.4, the following sequence

$$0 \to C\left(n\right) \xrightarrow{\sigma_n} C \xrightarrow{\tau_n} C\left[n\right] \to 0$$

is split exact. Then we have

$$(C(n), \sigma_n) = \ker [(\tau_a \otimes \tau_b) \Delta] = \ker (\tau_a) \wedge_C \ker (\tau_b) = C(a) \wedge_C C(b).$$

2) Assume that C is a strongly \mathbb{N} -graded coalgebra. Let us prove that $C(n) = C_0^{\wedge_C^n}$, for every $n \in \mathbb{N}$.

The case n=0 is trivial. Let us prove the equality above for every $n \ge 1$ by induction on n.

If n = 1, by definition, we have $C(1) = C_0 = C_0^{\wedge_C^1}$.

Let $n \ge 2$ and assume that $C(t) = C_0^{\wedge t}$ for every $0 \le t \le n-1$. By hypothesis we get that condition (10) holds true for n and hence by 2) we have $C(n) = C(a) \wedge_C C(b)$ so that, by Proposition 2.8, we have

$$C_0^{\wedge_C^n} = C_0^{\wedge_C^a} \wedge_C C_0^{\wedge_C^b} = C\left(a\right) \wedge_C C\left(b\right) = C\left(n\right).$$

PROPOSITION 2.11. Assume that \mathcal{M} is also complete and satisfies AB5. Let $(C = \bigoplus_{n \in \mathbb{N}} C_n, \Delta_C, \varepsilon_C)$ and $(D = \bigoplus_{n \in \mathbb{N}} D_n, \Delta_D, \varepsilon_D)$ be graded coalgebras in \mathcal{M} . Let $f : C \to D$ be a graded coalgebra homomorphism. Assume that D is a strongly \mathbb{N} -graded coalgebra. Then the following assertions are equivalent

- (a) f_0 and f_1 are monomorphisms and C is a strongly \mathbb{N} -graded coalgebra.
- (a') f_0 and f_1 are monomorphisms and $\Delta_{a,1}^C: C_{a+1} \to C_a \otimes C_1$ is a monomorphism for every $a \in \mathbb{N}$.
 - (b) f_n is a monomorphism for every $n \in \mathbb{N}$.
 - (c) f is a monomorphism.
 - (d) f_0 and f_1 are monomorphisms and $(C(n), \sigma_n) = C_0^{\wedge c}$, for every $n \in \mathbb{N}$.
 - (e) f_0 and f_1 are monomorphisms and $(C(2), \sigma_2) = C_0^{\wedge_C^2}$.

Proof. Recall that, by Lemma 2.6, we have (8).

- $(a) \Rightarrow (a')$ It is trivial.
- $(a') \Rightarrow (b)$ We proceed by induction on $a \geq 1$. For a = 1 there is nothing to prove. Let $a \geq 2$ and assume that f_{a-1} is a monomorphism. Since $\Delta_{a-1,1}^C : C_{a+1} \to C_a \otimes C_1$ is a monomorphism, in view of (8), we get that $f_a = f_{(a-1)+1}$ is a monomorphism.
 - $(b) \Rightarrow (c)$ Since $f = \bigoplus_{n \in \mathbb{N}} f_n$ and \mathcal{M} satisfies AB5, it is trivial (see e.g. [Po, page 53]).
- $(c) \Rightarrow (b)$ Since f is a monomorphism and $f \circ i_n^C = i_n^D \circ f_n$ for every $n \in \mathbb{N}$, it is clear that each f_n is a monomorphism too.
 - $(b) \Rightarrow (a)$ Let $a, b \geq 0$. In view of (8), $\Delta_{a,b}^C$ is a monomorphism as $\Delta_{a,b}^D$ is a monomorphism.
 - $(a) \Rightarrow (d)$ It follows by Theorem 2.10.
- $(d) \Rightarrow (e)$ It is trivial.
- $(e) \Rightarrow (a)$ By hypothesis,

$$\ker\left[\left(\tau_{1}^{C}\otimes\tau_{1}^{C}\right)\Delta_{C}\right]=\left(C_{0}^{\wedge_{C}^{2}},\delta_{2}^{C}\right)=\left(C\left(2\right),\sigma_{2}^{C}\right)=\ker\left(\tau_{2}^{C}\right)$$

so that

$$\operatorname{Im}\left[\left(\tau_{1}^{C}\otimes\tau_{1}^{C}\right)\Delta_{C}\right]=\operatorname{Coker}\left[\ker\left(\tau_{2}^{C}\right)\right]=\operatorname{Im}\left(\tau_{2}^{C}\right)=\left(C\left[2\right],\tau_{2}^{C}\right).$$

Hence there is a monomorphism $\beta:C\left[2\right]\to C\left[1\right]\otimes C\left[1\right]$ such that

$$\beta \circ \tau_2^C = \left(\tau_1^C \otimes \tau_1^C\right) \circ \Delta_C.$$

For every $n \in \mathbb{N}$, we have

$$\beta \circ \lambda_n^2 = \beta \circ \tau_2^C \circ i_n^C = \left(\tau_1^C \otimes \tau_1^C\right) \circ \Delta_C \circ i_n^C \stackrel{(3)}{=} \left(\tau_1^C \otimes \tau_1^C\right) \sum_{a+b=n} \left(i_a^C \otimes i_b^C\right) \circ \Delta_{a,b}^C = \sum_{a+b=n} \left(\lambda_a^1 \otimes \lambda_b^1\right) \circ \Delta_{a,b}^C$$

so that

(11)
$$\beta \circ \lambda_n^2 = \sum_{a+b=n} \left(\lambda_a^1 \otimes \lambda_b^1 \right) \circ \Delta_{a,b}^C.$$

Let us prove that $\Delta_{a,b}^C: C_{a+b} \to C_a \otimes C_b$ is a monomorphism for every $0 \leq a, b$.

For a=0 or b=0 there is nothing to prove as $(C_n, \Delta_{0,n}^C, \Delta_{n,0}^C)$ is a C_0 -bicomodule in view of Proposition 2.5. Let us check that $\Delta_{a,b}^C: C_{a+b} \to C_a \otimes C_b$ is a monomorphism for every a, b by induction on n=a+b.

For n = 1 there is nothing to prove.

For n=2, since $\lambda_0^1=0$, we have

$$\beta \circ \lambda_2^2 \stackrel{(11)}{=} \sum_{a+b=2} \left(\lambda_a^1 \otimes \lambda_b^1 \right) \circ \Delta_{a,b}^C = \left(\lambda_1^1 \otimes \lambda_1^1 \right) \circ \Delta_{1,1}^C.$$

So that, since β and λ_i^j are monomorphisms for every $j \leq i$, we have

$$\ker\left(\Delta_{1,1}^C\right) = \ker\left[\left(\lambda_1^1 \otimes \lambda_1^1\right) \circ \Delta_{1,1}^C\right] = \ker\left(\beta \circ \lambda_2^2\right) = 0.$$

Let $n \geq 3$ and assume that $\Delta_{u,v}^C : C_{u+v} \to C_u \otimes C_v$ is a monomorphism for every $u, v \in \mathbb{N}$ with u+v < n. Fix $a, b \in \mathbb{N}$ such that n = a+b and let h be a morphism in \mathcal{M} such that $\Delta_{a,b}^C \circ h = 0$. Let us prove that h = 0.

By (6), for every i, j > 0 with n = i + j, we have

$$\left(\Delta_{i,a-i}^{C} \otimes C_b \right) \Delta_{a,b}^{C} = \left(C_i \otimes \Delta_{j-b,b}^{C} \right) \Delta_{i,j}^{C}, \text{ if } a \geq i \text{ and hence } j \geq b$$

$$\left(\Delta_{a,i-a}^{C} \otimes C_j \right) \Delta_{i,j}^{C} = \left(C_a \otimes \Delta_{b-j,j}^{C} \right) \Delta_{a,b}^{C}, \text{ if } i \geq a \text{ and hence } b \geq j.$$

In the first case (the other one is analogous) we have

$$0 = \left(\Delta_{i,a-i}^C \otimes C_b\right) \Delta_{a,b}^C h = \left(C_i \otimes \Delta_{j-b,b}^C\right) \Delta_{i,j}^C h.$$

Since (j-b)+b=j < n, we have that $\Delta_{j-b,b}^C$ is a monomorphism and hence $C_i \otimes \Delta_{j-b,b}^C$ is a monomorphism too. Thus $\Delta_{i,j}^C h = 0$ for every i,j>0 with n=i+j.

Then, since $\lambda_0^1 = 0$, we have

$$\beta \circ \lambda_n^2 \circ h \stackrel{(11)}{=} \sum_{i+j=n} \left(\lambda_i^1 \otimes \lambda_j^1 \right) \circ \Delta_{i,j}^C \circ h = \sum_{\substack{i+j=n\\i \neq 0, j \neq 0}} \left(\lambda_i^1 \otimes \lambda_j^1 \right) \circ \Delta_{i,j}^C \circ h = 0$$

Since β and λ_n^2 are monomorphisms, we get h = 0.

2.12. The Cotensor Coalgebra. For any object $X \in \mathcal{M}$, we can define the cotensor coalgebra $T^c(X)$ as follows. As an object in \mathcal{M} it is defined by setting

$$T^{c} = T^{c}(X) := \bigoplus_{p \in \mathbb{N}} X^{\otimes p}.$$

Let $p_a: T(X) \to X^{\otimes a}$ be the canonical projection and let $\varepsilon_0 := \mathrm{Id}_1$. Let

$$\Delta_{p,q}: X^{\otimes p+q} \to X^{\otimes p} \otimes X^{\otimes q}, \text{ for every } p,q \in \mathbb{N}$$

be the canonical isomorphism, which is unique by Coherence Theorem. Still by Coherence Theorem and the definitions of $\Delta_{d,0}$ and $\Delta_{0,d}$, one has

$$[\Delta_{p,q} \otimes X^{\otimes r}] \Delta_{p+q,r} = [X^{\otimes p} \otimes \Delta_{q,r}] \Delta_{p,q+r}.$$
$$(X^{\otimes d} \otimes \varepsilon_0) \Delta_{d,0} = r_{X^{\otimes d}}^{-1}, \qquad (u_0 \otimes X^{\otimes d}) \Delta_{0,d} = l_{X^{\otimes d}}^{-1}.$$

so that (6) and (7) hold.

By Proposition 2.5, there there exists a unique morphism $\Delta_{T^c}: T^c \to T^c \otimes T^c$ such that (3) holds i.e.

(12)
$$\Delta \circ i_n = \sum_{a+b=n} (i_a \otimes i_b) \circ \Delta_{a,b}.$$

Moreover $(T^c, \Delta_{T^c}, \varepsilon_{T^c} = \varepsilon_0 p_0 = p_0)$ is a graded coalgebra with graduation defined by $(X^{\otimes p})_{p \in \mathbb{N}}$. Let $(C, \Delta_C, \varepsilon_C)$ be a coalgebra in \mathcal{M} .

As explained in Section 1, we have that $({}^{C}\mathcal{M}^{C}, \square_{C}, C)$ is a coabelian monoidal category. Furthermore one can see that ${}^{C}\mathcal{M}^{C}$ has also arbitrary direct sums which commute with \square_{C} .

Therefore we can consider, in the monoidal category $({}^{C}\mathcal{M}^{C}, \square_{C}, C)$, the cotensor coalgebra of an arbitrary C-bimodule M. We will denote it by

$$(T^c = T_C^c(M), \overline{\Delta}_{T^c}, \overline{\varepsilon_{T^c}}).$$

Note that

$$T^c = \bigoplus_{p \in \mathbb{N}} M^{\square_{C}p}, \qquad \overline{\Delta}_{T^c} : T^c \to T^c \square_C T^c \qquad \text{and} \qquad \overline{\varepsilon_{T^c}} : T^c \to C.$$

Set

$$\Delta_T = {}_{C}\varsigma_{V,W} \circ \overline{\Delta}_T : T^c \to T^c \otimes T^c \quad \text{and} \quad \varepsilon_{T^c} = \varepsilon_C \circ \overline{\varepsilon_{T^c}} : T^c \to \mathbf{1},$$

where $_{CSV,W}: T^c \square_C T^c \longrightarrow T^c \otimes T^c$ is the canonical morphism introduced in Section 1.

PROPOSITION 2.13. Let $(C, \Delta_C, \varepsilon_C)$ be a coalgebra in \mathcal{M} and let (M, ρ_M^r, ρ_M^l) be a C-bicomodule. Let $T^c = T_C^c(M)$ be the cotensor coalgebra. Then $T^c = \bigoplus_{n \in \mathbb{N}} T_n^c$ is a graded coalgebra where $T_n^c = M^{\square_C n}$. Moreover we have

$$\Delta_{m,n} = \begin{cases} (M^{\square m-1} \square \varsigma_M \square M^{\square n-1}), & \text{for any } m, n \ge 1; \\ (M^{\square m-1} \square \rho_M^r), & \text{for any } m \ge 1, n = 0; \\ (\rho_M^l \square M^{\square n-1}), & \text{for any } m = 0, n \ge 1; \\ \Delta_C, & \text{for any } m = 0, n = 0. \end{cases}$$

In particular T^c is a strongly \mathbb{N} -graded coalgebra.

Proof. It follows by (2.12) and by Proposition 2.3 as (3) is clearly satisfied.


Remark 2.14. Note that the coalgebra structure $T_C^c(M)$ in Proposition 2.13 is the same introduced in [AMS1, Theorem 2.9].

We have the following result.

NOTATION 2.15. Let $\delta: D \to C$ be a homomorphism of coalgebras in \mathcal{M} . By Proposition [AMS1, Proposition 1.10], $((D^{\wedge_{C}^{i}})_{i\in\mathbb{N}}, (\xi_{i}^{j})_{i,j\in\mathbb{N}})$ is a direct system in \mathcal{M} whose direct limit carries a natural coalgebra structure that makes it the direct limit of $((D^{\wedge_{C}^{i}})_{i\in\mathbb{N}}, (\xi_{i}^{j})_{i,j\in\mathbb{N}})$ as a direct system of coalgebras.

From now on we set: $(\widetilde{D}_C, (\xi_i)_{i \in \mathbb{N}}) = \varinjlim (D^{\wedge_C^i})_{i \in \mathbb{N}}$, where $\xi_i : D^{\wedge_C^i} \to \widetilde{D}_C$ denotes the structural morphism of the direct limit. We simply write \widetilde{D} if there is no danger of confusion. We note that, since \widetilde{D} is a direct limit of coalgebras, the canonical (coalgebra) homomorphisms $(\delta_i : D^{\wedge_C^i} \to C)_{i \in \mathbb{N}}$, which are compatible, factorize to a unique coalgebra homomorphism $\widetilde{\delta} : \widetilde{D} \to C$ such that $\widetilde{\delta}\xi_i = \delta_i$ for any $i \in \mathbb{N}$.

THEOREM 2.16. [AMS1, Theorems 2.13 and 2.15] Let (C, Δ, ε) be a coalgebra in a cocomplete and complete coabelian monoidal category \mathcal{M} satisfying AB5. Let (M, ρ_M^r, ρ_M^l) be a C-bicomodule. Let $\delta: D \to E$ be a monomorphism which is a homomorphism of coalgebras. Let $f_C: \widetilde{D} \to C$ be a coalgebra homomorphism and let $f_M: \widetilde{D} \to M$ be a morphism of C-bicomodules such that $f_M \xi_1 = 0$, where \widetilde{D} is a bicomodule via f_C . Then there is a unique coalgebra homomorphism $f: \widetilde{D} \to T_C^c(M)$ such that $p_0 f = f_C$ and $p_1 f = f_M$, where $p_n: T_C^c(M) \to M^{\square n}$ denotes the canonical projection.



Moreover

(13)
$$p_n \circ f = f_M^{\square_C n} \circ \overline{\Delta}_{\widetilde{D}}^{n-1} \text{ for any } n \in \mathbb{N}.$$

PROPOSITION 2.17. [AMS1, Proposition 2.11] Let (C, Δ, ε) be a coalgebra in \mathcal{M} and let (M, ρ_M^r, ρ_M^l) be a C-bicomodule. Let $T^c := T_C^c(M)$. Let E be a coalgebra and let $\alpha : E \to T^c$ and $\beta : E \to T^c$ be coalgebra homomorphisms. If $p_1\alpha = p_1\beta$, then $p_n\alpha = p_n\beta$ for any $n \ge 1$.

COROLLARY 2.18. Assume that \mathcal{M} is also complete and satisfies AB5. Let (C, Δ, ε) be a coalgebra in \mathcal{M} and let (M, ρ_M^r, ρ_M^l) be a C-bicomodule. Let $T^c := T_C^c(M)$. Let E be a coalgebra and let $\alpha : E \to T^c$ and $\beta : E \to T^c$ be coalgebra homomorphisms.

Then $\alpha = \beta$ whenever $p_n \alpha = p_n \beta$, for n = 0, 1.

Proof. It follows by Proposition 2.17 and Lemma 2.2.

PROPOSITION 2.19. [AMS1, Proposition 3.3] Let $C = \bigoplus_{t \in \mathbb{N}} C_t$ be a graded coalgebra in \mathcal{M} . Then

(14)
$$\tau_1^{\otimes n+1} \Delta_C^n i_b = 0, \text{ for every } 0 \le b \le n.$$

Moreover

$$C = \underline{\lim} (C_0^{\wedge_C^t})_{t \in \mathbb{N}}.$$

THEOREM 2.20. Let $(\mathcal{M}, \otimes, \mathbf{1})$ be a cocomplete and complete coabelian monoidal category satisfying AB5 and such that the tensor product commutes with direct sums.

Let (C, Δ, ε) be a coalgebra in \mathcal{M} and let (M, ρ_M^r, ρ_M^l) be a C-bicomodule.

Let B be a graded coalgebra, let $g_C: B_0 \to C$ be a coalgebra homomorphism and let $g_M: B_1 \to M$ be a morphism in ${}^C\mathcal{M}^C$, where $B_1 \in {}^C\mathcal{M}^C$ via g_C .

Then there is a unique coalgebra homomorphism $f: B \to T_C^c(M)$ such that

$$p_0^{T^c} \circ f = g_C \circ p_0^B$$
 and $p_1^{T^c} \circ f = g_M \circ p_1^B$.

Moreover f is a morphism of graded coalgebras where

$$f_t = \left(g_M \circ p_1^B\right)^{\square_C t} \circ \overline{\Delta}_B^{t-1} \circ i_t^B \text{ for every } t \in \mathbb{N}$$

where $\overline{\Delta}_B^{t-1}: B \to B^{\square_C t}$ is the t-th iterated comultiplication of B.

Proof. In view of Proposition 2.5, $(B_0, \Delta_0 = \Delta_{0,0}, \varepsilon_0 = \varepsilon i_0)$ is a coalgebra in \mathcal{M} , i_0^B is a coalgebra homomorphism and $(B_1, \Delta_{0,1}, \Delta_{1,0})$ is a B_0 -bicomodule. Moreover, $p_0^B : B \to B_0$ is a coalgebra homomorphism (see 5) and $p_1^B : B \to B_1$ is a morphism of B_0 -bicomodules (B is a B_0 -bicomodule through p_0).

By Theorem 2.16 applied in the case when δ is the morphism $i_0^B: B_0 \to B$ and by Proposition 2.19, there is a unique coalgebra homomorphism $f: B \to T_C^c(M)$ such that

$$p_0^{T^c} \circ f = g_C \circ p_0^B$$
 and $p_1^{T^c} \circ f = g_M \circ p_1^B$.

By (13), we have

$$p_t^{T^c} \circ f = \left(g_M \circ p_1^B\right)^{\square_C t} \circ \overline{\Delta}_B^{t-1} \text{ for any } t \in \mathbb{N}.$$

Let us prove there is a family $(f_t)_{t\in\mathbb{N}}$ of morphisms such that

$$p_t^{T^c} \circ f = f_t \circ p_t^B.$$

By (5) we have

$$(p_a^B \otimes p_b^B)\Delta_B = \Delta_{a,b}^B p_{a+b}^B$$

By Proposition 2.5, for every $n \in \mathbb{N}$, $p_n^B : B \to B_n$ is a morphism of B_0 -bicomodules and hence of C-bicomodules via g_C . Thus

$$(p_a^B \square_C p_b^B) \overline{\Delta}_B = \overline{\Delta}_{a,b}^B p_{a+b}^B$$

Hence there is a morphism $\alpha_t: (B_1)^{\otimes_A t} \to B_t$ such that

$$(p_1^B)^{\square_C t} \circ \overline{\Delta}_B^{t-1} = \alpha_t \circ p_t^B.$$

Then $\alpha_t = \left(p_1^B\right)^{\square_C t} \circ \overline{\Delta}_B^{t-1} \circ i_t^B$. We have

$$p_t^{T^c} \circ f = \left(g_M \circ p_1^B\right)^{\square_C t} \circ \overline{\Delta}_B^{t-1} = \left(g_M\right)^{\square_C t} \circ \left(p_1^B\right)^{\square_C t} \circ \overline{\Delta}_B^{t-1} = \left(g_M\right)^{\square_C t} \circ \alpha_t \circ p_t^B.$$

so that (15) holds true for $f_t := (g_M)^{\square_C t} \circ \alpha_t = (g_M)^{\square_C t} \circ (p_1^B)^{\square_C t} \circ \overline{\Delta}_B^{t-1} \circ i_t^B$.

Now, f is a morphism of graded coalgebras if and only if $f \circ i_t^B = i_t^{T^c} \circ f_t$, for every $t \in \mathbb{N}$. In view of Lemma 2.2, this is equivalent to prove that $p_s^{T^c} \circ f \circ i_t^B = p_s^{T^c} \circ i_t^{T^c} \circ f_t$, for every $s, t \in \mathbb{N}$ which is trivially true in view of (15).

REMARK 2.21. In the case C = 1, the universal property of the cotensor coalgebra stated in Theorem 2.20, differs from [Sch, Remark 2.3] where it is proved that there exists a unique graded coalgebra homomorphism $f: B \to T_c^c(M)$ such that $f_1 = g_M$. In fact we need extra assumptions as AB5 condition in order to establish the uniqueness of f as a coalgebra homomorphism. This uniqueness allows us to prove that f is graded whenever B is graded.

Theorem 2.22. Let $(C = \bigoplus_{n \in \mathbb{N}} C_n, \Delta_C, \varepsilon_C)$ be a graded coalgebra in a cocomplete and complete coabelian monoidal category M satisfying AB5. Assume that that the tensor product commutes with direct sums.

Let $T^c := T^c_{C_0}(C_1)$ be the cotensor coalgebra. Then there is a unique coalgebra homomorphism

$$\psi: C \to T_{C_0}^c(C_1),$$

such that $p_0^{T^c} \circ \psi = p_0^C$ and $p_1^{T^c} \circ \psi = p_1^C$.

Moreover ψ is a graded coalgebra homomorphism with

$$\psi_m = \left(p_1^C\right)^{\square m} \circ \overline{\Delta}_C^{m-1} \circ i_m^C \text{ for every } m \in \mathbb{N}$$

and the following assertions are equivalent.

- (a) C is a strongly \mathbb{N} -graded coalgebra.
- (a') $\Delta_{a,1}^C: C_{a+1} \to C_a \otimes C_1$ is a monomorphism for every $a \in \mathbb{N}$.
- (b) ψ_n is a monomorphism for every $n \in \mathbb{N}$.
- (c) ψ is a monomorphism.
- $(d) \ (C(n), \sigma_n^C) = C_0^{\wedge_C^n}, \text{ for every } n \in \mathbb{N}.$ $(e) \ (C(2), \sigma_2^C) = C_0^{\wedge_C}.$

Proof. In view of Proposition 2.5, C_1 is a bicomodule over the coalgebra C_0 so that we can consider $T^c_{C_0}(C_1)$. By Theorem 2.20, there is a unique coalgebra homomorphism $\psi: C \to T^c_{C_0}(C_1)$ such that $p_0^{T^c} \circ \psi = p_0^C$ and $p_1^{T^c} \circ \psi = p_1^C$. Moreover ψ is a morphism of graded coalgebras where

$$\psi_t = \left(p_1^C\right)^{\Box t} \circ \overline{\Delta}_C^{t-1} \circ i_t^C \text{ for every } t \in \mathbb{N}$$

where $\overline{\Delta}_C^{t-1}: C \to C^{\square_{C_0}t}$ is the t-th iterated comultiplication of C.

Since, by Proposition 2.13, T^c is a strongly N-graded coalgebra, we can apply Proposition 2.11. We conclude by observing that

$$\psi_0 = (p_1^C)^{\Box 0} \circ \overline{\Delta}_C^{0-1} \circ i_0^C = p_0^C \circ i_0^C = \operatorname{Id}_{C_0} \quad \text{and} \quad \psi_1 = (p_1^C)^{\Box 1} \circ \overline{\Delta}_C^{1-1} \circ i_1^C = p_1^C \circ i_1^C = \operatorname{Id}_{C_1}.$$

3. Graded and Strongly N-graded Algebras

In this section $(\mathcal{M}, \otimes, \mathbf{1})$ will denote a cocomplete abelian monoidal category such that the tensor product commutes with direct sums. Analogous results to those included in Section 2 can be obtained for graded algebras by a careful "dualization". Still we would like to point out that here we don't need to assume that \mathcal{M} is also complete and satisfies AB5 essentially because the analogue of Lemma 2.2 with the injections doesn't require these extra conditions.

DEFINITION 3.1. Recall that a graded algebra in \mathcal{M} is an algebra (A, m, u) where

$$A = \bigoplus_{n \in \mathbb{N}} A_n$$

is a graded object of \mathcal{M} such that $m:A\otimes A\to A$ is a graded homomorphism i.e. there exists a family $(m_n)_{n\in\mathbb{N}}$ of morphisms

$$m_n^A = m_n : \bigoplus_{a+b=n} (A_a \otimes A_b) = (A \otimes A)_n \to A_n$$
 such that $m = \bigoplus_{n \in \mathbb{N}} m_n$.

We set

$$m_{a,b}^A := \left(A_a \otimes A_b \overset{\gamma_{a,b}^{A,A}}{\to} (A \otimes A)_{a+b} \overset{m_{a+b}}{\to} A_{a+b} \right).$$

A homomorphism $f:(A, m_A, u_A) \to (B, m_B, u_B)$ of algebras is a graded algebra homomorphism if it is a graded homomorphism too.

PROPOSITION 3.2. Let (A, m, u) be an algebra in \mathcal{M} which is a graded object with graduation defined by $(A_k)_{k \in \mathbb{N}}$. Fix $n \in \mathbb{N}$. The following assertions are equivalent.

(i) There exists a morphism

$$m_n: \bigoplus_{a+b=n} (A_a \otimes A_b) = (A \otimes A)_n \to A_n \text{ such that } p_n m = m_n \Delta \left[(p_a \otimes p_b)_{a+b=n} \right].$$

(ii) There exists a family $(m_{a,b})_{a+b=n}$ of morphisms $m_{a,b}: A_a \otimes A_b \to A_n$ such that

$$(16) p_n m = \sum_{a+b=n} m_{a,b} \left(p_a \otimes p_b \right).$$

(iii) $p_n m(i_a \otimes i_b) = 0$, for every $a, b \in \mathbb{N}$, $a + b \neq n$.

Proof. It is analogous to that of Proposition 2.3.

PROPOSITION 3.3. 1) If $(A = \bigoplus_{n \in \mathbb{N}} A_n, m, u)$ is a graded algebra in \mathcal{M} , then

(17)
$$m(i_a \otimes i_b) = i_{a+b} m_{a,b}, \text{ for every } a, b \in \mathbb{N}.$$

2) Let (A, m, u) be an algebra in \mathcal{M} which is a graded object with graduation defined by $(A_k)_{k \in \mathbb{N}}$. If there exists a family $(m_{a,b})_{a,b \in \mathbb{N}}$ of morphisms $m_{a,b} : A_a \otimes A_b \to A_n$ such that (17) holds, then $(A = \bigoplus_{n \in \mathbb{N}} A_n, m, u)$ is a graded algebra in \mathcal{M} .

Proof. 1) By definition, we have

$$i_{a+b}m_{a,b} = i_{a+b}m_{a+b}\gamma_{a,b}^{A,A} = m\nabla \left[(i_u \otimes i_v)_{u+v=a+b} \right] \gamma_{a,b}^{A,A} = m(i_a \otimes i_b).$$

2) By the universal property of coproducts there is a unique

$$m_n: \bigoplus_{a+b=n} (A_a \otimes A_b) = (A \otimes A)_n \to A_n$$

such that $m_n \gamma_{a,b}^{A,A} = m_{a,b}$ for every $a, b \in \mathbb{N}$ such that a+b=n. For every $n, a, b \in \mathbb{N}$ such that a+b=n, we have

$$m\nabla \left[(i_u \otimes i_v)_{u+v=n} \right] \gamma_{a,b}^{A,A}$$

$$= m\nabla \left[(i_u \otimes i_v)_{u+v=a+b} \right] \gamma_{a,b}^{A,A} = m(i_a \otimes i_b) \stackrel{(17)}{=} i_{a+b} m_{a,b} = i_n m_n \gamma_{a,b}^{A,A}$$

so that $m\nabla \left[(i_u \otimes i_v)_{u+v=n} \right] = i_n m_n$.

PROPOSITION 3.4. 1) Let $A = \bigoplus_{n \in \mathbb{N}} A_n$ be a graded object of \mathcal{M} such that there exists a family $(m_{a,b})_{a,b \in \mathbb{N}}$

$$m_{a,b}: A_a \otimes A_b \to A_{a+b},$$

of morphisms and a morphism $u_0: \mathbf{1} \to A_0$ which satisfy

(18)
$$m_{a+b,c} \left(m_{a,b} \otimes A_c \right) = m_{a,b+c} \left(A_a \otimes m_{b,c} \right),$$

(19)
$$m_{d,0}(A_d \otimes u_0) = r_{A_d}, \qquad m_{0,d}(u_0 \otimes A_d) = l_{A_d},$$

for every $a, b, c \in \mathbb{N}$. Then there exists a unique morphism $m : A \otimes A \to A$ such that (17) holds. Moreover $(A \oplus_{n \in \mathbb{N}} A_n, m, u = i_0 u_0)$ is a graded algebra.

2) If A is a graded algebra then (16) holds, $u = i_0 p_0 u$ so that u is a graded homomorphism, and we have that (18) and (19) hold for every $a, b, c \in \mathbb{N}$, where $u_0 = p_0 u$.

Moreover $(A_0, m_0 = m_{0.0}, u_0 = p_0 u)$ is an algebra in \mathcal{M} , p_0 is an algebra homomorphism and, for every $n \in \mathbb{N}$, $(A_n, m_{0,n}, m_{n,0})$ is an A_0 -bimodule such that $i_n : A_n \to A$ is a morphism of A_0 -bimodules (A is an A_0 -bimodule through i_0).

Proof. 1) Since the tensor product commutes with direct sums, we have that $A \otimes A = \bigoplus_{a,b \in \mathbb{N}} (A_a \otimes A_b)$. Thus, by the universal property of coproduct there exists a unique morphism $m:A\otimes A\to A$ such that (17) holds.

The remaining part of the proof is similar to that of Proposition 2.5.

2) It is analogous to Proposition 2.5.

Given an ideal (I, i_I) of an algebra A in \mathcal{M} , one can define as for ordinary algebras, the iterated n-th power I^n of I in A (see e.g. [AMS2, Example 3.2]).

DEFINITION 3.5. Let $(A = \bigoplus_{n \in \mathbb{N}} A_n, m, u)$ be a graded algebra in \mathcal{M} . In analogy with the group graded case, we say that A is a strongly \mathbb{N} -graded algebra whenever

 $m_{i,j}^A: A_i \otimes A_j \to A_{i+j}$ is an epimorphism for every $i, j \in \mathbb{N}$,

where $m_{i,j}^A$ is the morphism of Definition 3.1.

PROPOSITION 3.6. Let $(A = \bigoplus_{n \in \mathbb{N}} A_n, m_A, u_A)$ and $(B = \bigoplus_{n \in \mathbb{N}} B_n, m_B, u_B)$ be graded algebras in \mathcal{M} . Let $f:A\to B$ be a graded algebra homomorphism. Assume that A is a strongly \mathbb{N} -graded algebra.

Then the following assertions are equivalent

- (a) f_0 and f_1 are epimorphisms and B is is a strongly N-graded algebra.
- (a') f_0 and f_1 are epimorphisms and $m_{a,1}^B: B_a \otimes B_1 \to B_{a+1}$ is an epimorphism for every $a \in \mathbb{N}$.
- (b) f_n is an epimorphism for every $n \in \mathbb{N}$.
- (c) f is an epimorphism.
- (d) f_0 and f_1 are epimorphisms and $(B[n], \nu_n^B) = B[1]^n$, for every $n \in \mathbb{N}$. (e) f_0 and f_1 are epimorphisms and $(B[2], \nu_2^B) = B[1]^2$.

Proof. It is analogous to that of Proposition 2.11.

3.7. The tensor algebra. For any object $X \in \mathcal{M}$, we can define the tensor algebra T(X) as follows. As an object in \mathcal{M} it is defined by setting

$$T = T(X) := \bigoplus_{p \in \mathbb{N}} X^{\otimes p}$$
.

Let $i_p: X^{\otimes p} \to T(X)$ be the canonical injection and let $u_0 := \mathrm{Id}_1$.

$$m_{p,q}: X^{\otimes p} \otimes X^{\otimes q} \to X^{\otimes p+q}$$
, for every $p, q \in \mathbb{N}$

be the canonical isomorphism, which is unique by Coherence Theorem. Still by Coherence Theorem and the definitions of $m_{d,0}$ and $m_{0,d}$, one has

(20)
$$m_{p+q,r}[m_{p,q} \otimes X^{\otimes r}] = m_{p,q+r}[X^{\otimes p} \otimes m_{q,r}].$$
$$m_{d,0} \left(X^{\otimes d} \otimes u_0\right) = r_{X^{\otimes d}}, \qquad m_{0,d} \left(u_0 \otimes X^{\otimes d}\right) = l_{X^{\otimes d}}$$

so that (18) and (19) hold.

By Proposition 3.4, there exists a unique morphism $m_T: T \otimes T \to T$ such that (17) holds, i.e.

(21)
$$m_T \circ (i_p \otimes i_q) = i_{p+q} \circ m_{p,q}, \text{ for every } p, q \in \mathbb{N}.$$

Moreover $(T, m_T, u_T = i_0 u_0 = i_0)$ is a graded algebra with graduation defined by $(X^{\otimes p})_{n \in \mathbb{N}}$. Note that

$$m_T := \bigoplus_{p \in \mathbb{N}} (\nabla [(m_{i,j})_{i+j=p}]) : T(X) \otimes T(X) \to T(X),$$

where $\nabla[(m_{i,j})_{i+j=p}]: \bigoplus_{i+j=p} (X^{\otimes i} \otimes X^{\otimes j}) \longrightarrow X^{\otimes p}$ denotes the codiagonal morphism associated to the family $(m_{i,j})_{i+j=p}$. Let (A, m_A, u_A) be an algebra in \mathcal{M} .

As explained in Section 1, we have that $({}_{A}\mathcal{M}_{A}, \otimes_{A}, A)$ is an abelian monoidal category. Furthermore one can see that ${}_{A}\mathcal{M}_{A}$ has also arbitrary direct sums which commute with \otimes_{A} .

Therefore we can consider, in the monoidal category $({}_{A}\mathcal{M}_{A}, \otimes_{A}, A)$, the tensor algebra of an arbitrary A-bimodule M. We will denote it by

$$(T = T_A(M), \overline{m}_T, \overline{u}_T)$$
.

Note that

$$T = \bigoplus_{p \in \mathbb{N}} M^{\otimes_A p}, \quad \overline{m}_T : T \otimes_A T \to T \quad \text{and} \quad \overline{u}_T : A \to T.$$

Set

$$m_T = \overline{m}_T \circ {}_A \chi_{T,T} : T \otimes T \to T$$
 and $u_T = \overline{u_T} \circ u_A : \mathbf{1} \to T$,

where $A\chi_{T,T}: T \otimes T \longrightarrow T \otimes_A T$ is the canonical morphism introduced in Section 1.

We have the following result.

PROPOSITION 3.8. Let (A, m_A, u_A) be an algebra in \mathcal{M} and let (M, μ_M^r, μ_M^l) be an A-bimodule. Let $T = T_A(M)$ be the tensor algebra. Then $(T = \bigoplus_{n \in \mathbb{N}} T_n, m_T, u_T)$ is a graded algebra in $(\mathcal{M}, \otimes, \mathbf{1})$, where $T_n = M^{\otimes_A n}$. Moreover we have

$$m_{u,v} = \begin{cases} (M^{\otimes_A u - 1} \otimes_A \chi_M \otimes_A M^{\otimes_A v - 1}), & \text{for any } u, v \ge 1; \\ (M^{\otimes_A u - 1} \otimes_A \mu_M^r), & \text{for any } u \ge 1, v = 0; \\ (\mu_M^l \otimes_A M^{\otimes_A v - 1}), & \text{for any } u = 0, v \ge 1; \\ m_A, & \text{for any } u = 0, v = 0. \end{cases}$$

In particular T is a strongly \mathbb{N} -graded algebra.

Proof. Follows by (3.7) and by Proposition 3.3 as (17) is clearly satisfied.

We are now able to state the Universal property of the relative tensor algebra.

THEOREM 3.9 (Universal property of the relative tensor algebra). Let $(\mathcal{M}, \otimes, \mathbf{1})$ be a cocomplete abelian monoidal category such that the tensor product commutes with direct sums.

Let A, B be algebras in M and let $f_A: A \to B$ be an algebra homomorphism.

Let $M \in {}_{A}\mathcal{M}_{A}$, and let $f_{M} : M \to B$ be a morphism in ${}_{A}\mathcal{M}_{A}$, where $B \in {}_{A}\mathcal{M}_{A}$ via f_{0} .

Then there is a unique algebra homomorphism $f: T_A(M) \to B$ such that

$$M \xrightarrow{i_1} T_A(M)$$

$$f_M \downarrow \qquad \qquad f_{i_0}$$

$$A$$

where $i_0: A \to T_A(M)$ and $i_1: M \to T_A(M)$ are the canonical injections. Moreover

$$f \circ i_t^T = \overline{m}_B^{t-1} \circ (f_M)^{\otimes_A t}$$
 for every $t \in \mathbb{N}$,

where $\overline{m}_B^{t-1}: B^{\otimes_A t} \to B$ is the t-th iterated multiplication of B.

When B is a graded algebra the universal property of $T_A(M)$ has a graded version which is the following.

THEOREM 3.10. Let $(\mathcal{M}, \otimes, \mathbf{1})$ be a cocomplete abelian monoidal category such that the tensor product commutes with direct sums. Let A be an algebra in \mathcal{M} and let $M \in {}_{A}\mathcal{M}_{A}$.

Let B be a graded algebra, let $g_A: A \to B_0$ be an algebra homomorphism and let $g_M: M \to B_1$ be a morphism in ${}_A\mathcal{M}_A$, where $B_1 \in {}_A\mathcal{M}_A$ via g_A .

Then there is a unique algebra homomorphism $f: T_A(M) \to B$ such that

$$f \circ i_0^T = i_0^B \circ g_A$$
 and $f \circ i_1^T = i_1^B \circ g_M$.

Moreover f is a morphism of graded algebras where

$$f_t = p_t^B \circ \overline{m}_B^{t-1} \circ (i_1^B \circ g_M)^{\otimes_A t}$$
 for every $t \in \mathbb{N}$

and $\overline{m}_B^{t-1}: B^{\otimes_A t} \to B$ is the t-th iterated multiplication of B.

Proof. It is analogous to that of Theorem 2.20.

Theorem 3.11. Let $(A = \bigoplus_{n \in \mathbb{N}} A_n, m_A, u_A)$ be a graded algebra in a cocomplete abelian monoidal category \mathcal{M} . Assume that the tensor product commutes with direct sums.

Let $T:=T_{A_0}(A_1)$ be the tensor algebra. Then there is a unique algebra homomorphism

$$\varphi: T_{A_0}(A_1) \to A,$$

 $such\ that\ \varphi\circ i_0^T=i_0^A\ and\ \varphi\circ i_1^T=i_1^A.$

Moreover φ is a graded algebra homomorphism with

$$\varphi_t = p_t^A \circ \overline{m}_A^{t-1} \circ (i_1^A)^{\otimes_{A_0} t} \text{ for every } t \in \mathbb{N}$$

and the following assertions are equivalent.

- (a) A is a strongly \mathbb{N} -graded algebra.
- (a') $m_{a,1}^A: A_a \otimes A_1 \to A_{a+1}$ is an epimorphism for every $a \in \mathbb{N}$.
- (b) φ_n is an epimorphism for every $n \in \mathbb{N}$.
- (c) φ is an epimorphism.
- (d) $(A[n], \nu_n^A) = A[1]^n$, for every $n \in \mathbb{N}$. (e) $(A[2], \nu_2^A) = A[1]^2$.

Proof. It is analogous to that of Theorem 2.22.

4. Braided Bialgebra Structure of the Cotensor Coalgebra

The main aim of this section is to provide a braided bialgebra structure for the cotensor coalgebra inside a braided monoidal category. This structure is used to extend the notion of bialgebra of type one, introduced in the classical case by Nichols in [Ni], to the wider context of a braided monoidal category (see Definition 6.7). A universal property for the cotensor bialgebra is also proven (see Theorem 4.7).

4.1. Braided bialgebras. A braided bialgebra in (\mathcal{M}, c) , is a sextuple $(H, m, u, \Delta, \varepsilon)$ such that (H, m, u) is an algebra in \mathcal{M} , (H, Δ, ε) is a coalgebra in \mathcal{M} and this two structure are compatible in the sense that

(22)
$$\Delta \circ m = (m \otimes m) \circ (H \otimes c \otimes H) \circ (\Delta \otimes \Delta)$$
 and $m_1 \circ (\varepsilon \otimes \varepsilon) = \varepsilon \circ m$.

DEFINITIONS 4.2. Let $(B, m_B, u_B, \Delta_B, \varepsilon_B)$ be a braided bialgebra in a cocomplete braided monoidal category (\mathcal{M}, c) . Then B is called a graded braided bialgebra if B is a graded object with graduation defined by $(B_k)_{k\in\mathbb{N}}$ and if, with respect to this graduation, (B, m_B, u_B) is a graded algebra and $(B, \Delta_B, \varepsilon_B)$ is a graded coalgebra. A morphism $f: A \to B$ between graded braided bialgebras is called a morphism of graded (braided) bialgebras whenever it is both a morphism of graded algebras and a morphism of graded coalgebras i.e. if it is a bialgebra homomorphism which is also a graded homomorphism.

DEFINITION 4.3. Let H be a braided bialgebra in (\mathcal{M}, c) .

An object in ${}^H_H\mathcal{M}^H_H$ is a 5-tuple $(M,\mu^r_M,\mu^l_M,\rho^r_M,\rho^l_M)$ such that

- (M, μ_M^r, μ_M^l) is an *H*-bimodule;
- (M, ρ_M^r, ρ_M^l) is an H-bicomodule;
- the following compatibility conditions are fulfilled:

(23)
$$\rho_M^l \mu_M^l = (m_H \otimes \mu_M^l) (H \otimes c_{H,H} \otimes M) (\Delta_H \otimes \rho_M^l),$$

(24)
$$\rho_M^l \mu_M^r = (m_H \otimes \mu_M^r) (H \otimes c_{M,H} \otimes H) (\rho_M^l \otimes \Delta_H),$$

(25)
$$\rho_M^r \mu_M^l = (\mu_M^l \otimes m_H) (H \otimes c_{H,M} \otimes H) (\Delta_H \otimes \rho_M^r),$$

(26)
$$\rho_M^r \mu_M^r = (\mu_M^r \otimes m_H) (M \otimes c_{H,H} \otimes H) (\rho_M^r \otimes \Delta_H).$$

PROPOSITION 4.4. Let (\mathcal{M}, c) be a cocomplete coabelian braided monoidal category and let C = $\bigoplus_{n\in\mathbb{N}} C_n$ be a graded coalgebra in \mathcal{M} . Assume that the tensor product commutes with direct sums. Then $(C \otimes C, \Delta_{C \otimes C}, \varepsilon_{C \otimes C})$ is a graded coalgebra where

$$\Delta_{C \otimes C} : C \otimes C \xrightarrow{\Delta_C \otimes \Delta_C} C \otimes C \otimes C \otimes C \otimes C \xrightarrow{C \otimes c_{C,C} \otimes C} C \otimes C \otimes C \otimes C, \quad \varepsilon_{C \otimes C} : C \otimes C \xrightarrow{\varepsilon_C \otimes \varepsilon_C} \mathbf{1} \otimes \mathbf{1} \xrightarrow{m_1} \mathbf{1}.$$

and with graduation given by $(C \otimes C)_n = \bigoplus_{a+b=n} C_a \otimes C_b$. Moreover, for every $s, t \in \mathbb{N}$ we have

(27)
$$\Delta_{s,t}^{C\otimes C} := \nabla \left[\sum_{\substack{i+j=a, u+v=b\\i+u=s, j+v=t}} \left(\gamma_{i,u}^{C,C} \otimes \gamma_{j,v}^{C,C} \right) \left(C_i \otimes c_{C_j,C_u} \otimes C_v \right) \left(\Delta_{i,j} \otimes \Delta_{u,v} \right) \right]_{\substack{a,b,\\a+b=s+t}}$$

Proof. It is well known that $(C \otimes C, \Delta_{C \otimes C}, \varepsilon_{C \otimes C})$ is a coalgebra (it is dual to [Mj1, Lemma 9.2.12, page 438]). Let us check the part of the statement concerning the graduation. In view of Proposition 2.3, since C is a graded coalgebra, for every $n \in \mathbb{N}$ there exists a family $(\Delta_{a,b})_{a+b=n}$ of morphisms $\Delta_{a,b}: C_n \to C_a \otimes C_b$, such that (3) holds true. For every $s,t \in \mathbb{N}$ define $\Delta_{s,t}^{C \otimes C}: (C \otimes C)_{s+t} \to (C \otimes C)_s \otimes (C \otimes C)_t$ accordingly to (27). It is straightforward to prove that

$$\Delta_{C \otimes C} \circ i_n^{C \otimes C} = \sum_{s+t=n} \left(i_s^{C \otimes C} \otimes i_t^{C \otimes C} \right) \Delta_{s,t}^{C \otimes C}.$$

THEOREM 4.5. Let H be a braided bialgebra in a cocomplete and complete coabelian braided monoidal category (\mathcal{M}, c) satisfying AB5. Assume that the tensor product commutes with direct sums.

Let $(M, \mu_M^r, \mu_M^l, \rho_M^r, \rho_M^l)$ be in ${}^H_H \mathcal{M}^H_H$. Let $T^c = T^c_H(M)$ be the cotensor coalgebra. Then there are unique coalgebra homomorphisms

$$m_{T^c}: T^c \otimes T^c \to T^c$$
 and $u_{T^c}: \mathbf{1} \to T^c$

such that the diagrams

$$(28) \qquad T^{c} \xrightarrow{p_{1}} M \qquad T^{c} \xrightarrow{p_{1}} M$$

$$\downarrow p_{0} \qquad m_{T^{c}} \qquad \uparrow \mu_{M}^{l}(p_{0} \otimes p_{1}) + \mu_{M}^{r}(p_{1} \otimes p_{0}) \qquad p_{0} \qquad \downarrow m_{T^{c}} \qquad \uparrow 0$$

$$H \xleftarrow{m_{H}(p_{0} \otimes p_{0})} T^{c} \otimes T^{c} \qquad H \xleftarrow{u_{H}} 1$$

are commutative, where $p_n: T^c \to M^{\square_H n}$ denotes the canonical projection. Moreover $(T^c, m_{T^c}, u_{T^c}, \Delta_{T^c}, \varepsilon_{T^c})$ is a graded braided bialgebra in \mathcal{M} .

Proof. First of all, by Proposition 4.4, we have that $(E = T^c \otimes T^c, \Delta_E, \varepsilon_E)$ is a graded coalgebra where

$$\Delta_E : T^c \otimes T^c \overset{\Delta_{T^c} \otimes \Delta_{T^c}}{\longrightarrow} T^c \otimes T^c \otimes T^c \otimes T^c \overset{T^c \otimes c_{T^c, T^c} \otimes T^c}{\longrightarrow} T^c \otimes T^c \otimes T^c \otimes T^c,$$

$$\varepsilon_E : T^c \otimes T^c \overset{\varepsilon_{T^c} \otimes \varepsilon_{T^c}}{\longrightarrow} \mathbf{1} \otimes \mathbf{1} \overset{m_1}{\longrightarrow} \mathbf{1}.$$

Since $T^c = \bigoplus_{n \in \mathbb{N}} T_n^c$ is a graded coalgebra, we have that (5) holds so that, by Proposition 2.13, we have

$$(p_0 \otimes p_0)\Delta_{T^c} = \Delta_H p_0$$
 and $(p_0 \otimes p_1)\Delta_{T^c} = \rho_M^l p_1$.

Set

 $g_H:=m_H:E_0=H\otimes H\to H$ and $g_M:=\nabla\left[\left(\mu_M^l,\mu_M^r\right)\right]:E_1=(H\otimes M)\oplus (M\otimes H)\to M.$ Then g_H is a coalgebra homomorphism:

$$\Delta_H g_H = \Delta_H m_H = (m_H \otimes m_H) (H \otimes c_{H,H} \otimes H) (\Delta_H \otimes \Delta_H) = (g_H \otimes g_H) \Delta_{H \otimes H}.$$

Moreover g_M is a morphism of left H-comodules

$$(g_{H} \otimes g_{M}) \Delta_{0,1}^{T^{c} \otimes T^{c}} \stackrel{(27)}{=} (g_{H} \otimes g_{M}) \nabla \begin{bmatrix} \left(\gamma_{0,0}^{T^{c},T^{c}} \otimes \gamma_{0,1}^{T^{c},T^{c}}\right) \left(T_{0}^{c} \otimes c_{T_{0}^{c},T_{0}^{c}} \otimes T_{1}^{c}\right) \left(\Delta_{0,0} \otimes \Delta_{0,1}\right), \\ \left(\gamma_{0,0}^{T^{c},T^{c}} \otimes \gamma_{1,0}^{T^{c},T^{c}}\right) \left(T_{0}^{c} \otimes c_{T_{1}^{c},T_{0}^{c}} \otimes T_{0}^{c}\right) \left(\Delta_{0,1} \otimes \Delta_{0,0}\right) \end{bmatrix}$$

$$= \nabla \begin{bmatrix} \left(m_{H} \otimes \mu_{M}^{l}\right) \left(H \otimes c_{H,H} \otimes M\right) \left(\Delta_{H} \otimes \rho_{M}^{l}\right), \\ \left(m_{H} \otimes \mu_{M}^{r}\right) \left(H \otimes c_{M,H} \otimes H\right) \left(\rho_{M}^{r} \otimes \Delta_{H}\right) \end{bmatrix} \stackrel{(23),(24)}{=} \nabla \left[\rho_{M}^{l} \mu_{M}^{l}, \rho_{M}^{l} \mu_{M}^{r}\right] = \rho_{M}^{l} g_{M}$$

Analogously $\rho_M^r g_M = (g_M \otimes g_H) \Delta_{1,0}^{T^c \otimes T^c}$, i.e. g_M is a morphism of right H-comodules and hence a morphism of H-bicomodules. By applying Theorem 2.20 to the case "C" = H and "B" = $H \otimes H$, using the maps g_H and g_M above, we get a unique coalgebra homomorphism $m_{T^c} = f : E \to T^c$ such that the left side of (28) is commutative so that

$$p_0 m_{T^c} (m_{T^c} \otimes T^c) = p_0 m_{T^c} (T^c \otimes m_{T^c})$$
 and $p_1 m_{T^c} (m_{T^c} \otimes T^c) = p_1 m_{T^c} (T^c \otimes m_{T^c})$.

Since $T^c \otimes T^c \otimes T^c$ is a coalgebra and $m_{T^c} (m_{T^c} \otimes T^c)$ and $m_{T^c} (T^c \otimes m_{T^c})$ are coalgebra homomorphisms, then by Corollary 2.18, we obtain

$$m_{T^c} (m_{T^c} \otimes T^c) = m_{T^c} (T^c \otimes m_{T^c}).$$

Set

$$g'_H := u_H : \mathbf{1} \to H$$
 and $g'_M := 0 : 0 \to M$.

Then g'_H is a coalgebra homomorphism and g'_M is a morphism of H-bicomodules. By applying Theorem 2.20 to the case $B = \mathbf{1}$, we get a unique coalgebra homomorphism $u_{T^c} = f' : \mathbf{1} \to T^c$ such that the right side of 28 is commutative so that

$$p_0 m_{T^c} (u_{T^c} \otimes T^c) = m_H (u_H \otimes H) (\mathbf{1} \otimes p_0) = l_H (\mathbf{1} \otimes p_0) = p_0 l_{T^c}$$

$$p_1 m_{T^c} (u_{T^c} \otimes T^c) = \mu_M^l (u_H \otimes M) (\mathbf{1} \otimes p_1) = l_M (\mathbf{1} \otimes p_1) = p_1 l_{T^c}.$$

Since $\mathbf{1} \otimes T^c$ is a coalgebra and $m_{T^c} (u_{T^c} \otimes T^c)$ and r_{T^c} are coalgebra homomorphisms, then by Corollary 2.18, we have $m_{T^c} (u_{T^c} \otimes T^c) = l_{T^c}$. Analogously $m_{T^c} (T^c \otimes u_{T^c}) = r_{T^c}$. Now, in view of Theorem 2.20, m_{T^c} and u_{T^c} are graded homomorphisms so that $(T^c, m_{T^c}, u_{T^c}, \Delta_{T^c}, \varepsilon_{T^c})$ is a graded braided bialgebra in \mathcal{M} .

REMARK 4.6. Let M be an object in a cocomplete and complete coabelian braided monoidal category (\mathcal{M},c) satisfying AB5. Assume that the tensor product commutes with direct sums. By applying Theorem 4.5 to the case $H=\mathbf{1}$ we endow the cotensor coalgebra $T^c=T^c_H(M)$ with an algebra structure such that T^c becomes a braided bialgebra. This algebra structure is the braided analogue of the so called "Shuffle Algebra" in the category of vector spaces.

Theorem 4.7. Let H be a braided bialgebra in a cocomplete and complete coabelian braided monoidal category (\mathcal{M}, c) satisfying AB5. Assume that the tensor product commutes with direct sums

Let $(M, \mu_M^r, \mu_M^l, \rho_M^r, \rho_M^l)$ be in ${}^H_H\mathcal{M}^H_H$. Let $T^c = T^c_H(M)$ be the cotensor coalgebra.

Let $\delta: D \to E$ be a monomorphism which is a homomorphism of coalgebras. Assume that there exist morphisms


$$m_{\widetilde{D}}:\widetilde{D}\otimes\widetilde{D} o\widetilde{D} \qquad and \qquad u_{\widetilde{D}}:\mathbf{1} o\widetilde{D}$$

such that $\left(\widetilde{D}, m_{\widetilde{D}}, u_{\widetilde{D}}, \Delta_{\widetilde{D}}, \varepsilon_{\widetilde{D}}, \right)$ is a braided bialgebra in \mathcal{M} .

Let $f_H: \widetilde{\widetilde{D}} \to H$ be a bialgebra homomorphism and let $f_M: \widetilde{D} \to M$ be a morphism of H-bicomodules such that $f_M \xi_1 = 0$, where \widetilde{D} is a bicomodule via f_H . Assume that

(29)
$$f_{M}m_{\widetilde{D}} = \mu_{M}^{l} \left(f_{H} \otimes f_{M} \right) + \mu_{M}^{r} \left(f_{M} \otimes f_{H} \right)$$

(i.e. f_M is a derivation of \widetilde{D} with values in the \widetilde{D} -bimodule M, where M is regarded as a bimodule via f_H). Then there is a unique coalgebra homomorphism $f: \widetilde{D} \to T_H^c(M)$ such that $p_0 f = f_H$ and $p_1 f = f_M$, where $p_n: T_C^c(M) \to M^{\square n}$ denotes the canonical projection.



Moreover f is a bialgebra homomorphism.

Proof. By Theorem 2.16, there is a unique coalgebra homomorphism $f: \widetilde{D} \to T_H^c(M)$ such that $p_0 f = f_H$ and $p_1 f = f_M$. We have

$$p_{0}m_{T^{c}}\left(f\otimes f\right) \stackrel{(28)}{=} m_{H}\left(f_{H}\otimes f_{H}\right) = f_{H}m_{\widetilde{D}} = p_{0}fm_{\widetilde{D}},$$

$$p_{1}m_{T^{c}}\left(f\otimes f\right) \stackrel{(28)}{=} \mu_{M}^{l}\left(f_{H}\otimes f_{M}\right) + \mu_{M}^{r}\left(f_{M}\otimes f_{H}\right) = f_{M}m_{\widetilde{D}} = p_{1}fm_{\widetilde{D}}.$$

We also obtain

$$f_{M}u_{\widetilde{D}} = f_{M}m_{\widetilde{D}}\left(u_{\widetilde{D}} \otimes u_{\widetilde{D}}\right)m_{\mathbf{1}}^{-1} \stackrel{(29)}{=} \mu_{M}^{l}\left(f_{H}u_{\widetilde{D}} \otimes f_{M}u_{\widetilde{D}}\right)m_{\mathbf{1}}^{-1} + \mu_{M}^{r}\left(f_{M}u_{\widetilde{D}} \otimes f_{H}u_{\widetilde{D}}\right)m_{\mathbf{1}}^{-1}$$
$$= \mu_{M}^{l}\left(u_{H} \otimes f_{M}u_{\widetilde{D}}\right)m_{\mathbf{1}}^{-1} + \mu_{M}^{r}\left(f_{M}u_{\widetilde{D}} \otimes u_{H}\right)m_{\mathbf{1}}^{-1} = f_{M}u_{\widetilde{D}} + f_{M}u_{\widetilde{D}}$$

so that $f_M u_{\widetilde{D}} = 0$ and hence, by Theorem 2.16,

$$p_0 f u_{\widetilde{D}} = f_H u_{\widetilde{D}} = u_H = p_0 u_{T^c} \qquad \text{and} \qquad p_1 f u_{\widetilde{D}} = f_M u_{\widetilde{D}} = 0 = p_1 u_{T^c}.$$

Since $m_{T^c}(f \otimes f)$, $fm_{\widetilde{D}}: \widetilde{D} \otimes \widetilde{D} \to T^c$ and $fu_{\widetilde{D}}, u_{T^c}: \mathbf{1} \to T^c$ are coalgebra homomorphisms and since, for i = 0, 1,

$$p_i m_{T^c} (f \otimes f) = p_i f m_{\widetilde{D}}$$
 and $p_i f u_{\widetilde{D}} = p_i u_{T^c}$,

then, by Corollary 2.18, we get that $m_{T^c}(f \otimes f) = f m_{\widetilde{D}}$ and $f u_{\widetilde{D}} = u_{T^c}$ i.e. that f is an algebra homomorphism.

5. Graded Braided Bialgebra Structure of the Tensor Algebra

In this section $((\mathcal{M}, \otimes, \mathbf{1}), c)$ will denote a cocomplete abelian braided monoidal category such that the tensor product commutes with direct sums.

Next aim is to provide a braided bialgebra structure (see 4.1) for the tensor algebra inside a braided monoidal category.

PROPOSITION 5.1. Let $A = \bigoplus_{n \in \mathbb{N}} A_n$ be a graded algebra in \mathcal{M} . Then $(A \otimes A, m_{A \otimes A}, u_{A \otimes A})$ is a graded algebra where

$$m_{A\otimes A}: A\otimes A\otimes A\otimes A \overset{A\otimes c_{A,A}\otimes A}{\longrightarrow} A\otimes A\otimes A\otimes A \overset{m\otimes m}{\longrightarrow} A\otimes A, \quad u_{A\otimes A}: \mathbf{1} \overset{m_{\mathbf{1}}^{-1}}{\longrightarrow} \mathbf{1}\otimes \mathbf{1} \overset{u\otimes u}{\longrightarrow} A\otimes A.$$
and with graduation given by $(A\otimes A)_n = \bigoplus_{a+b=n} A_a\otimes A_b.$

THEOREM 5.2. Let H be a braided bialgebra in a cocomplete abelian braided monoidal category (\mathcal{M}, c) . Assume that the tensor product commutes with direct sums.

Let $(M, \mu_M^r, \mu_M^l, \rho_M^r, \rho_M^l)$ be in ${}^H_H \mathcal{M}^H_H$. Let $T = T_H(M)$ be the tensor algebra. Then there are unique coalgebra homomorphisms

$$\Delta_T: T \to T \otimes T$$
 and $\varepsilon_T: T \to \mathbf{1}$

such that the diagrams

are commutative, where $i_n: M^{\otimes_H n} \to T$ denotes the canonical injection. Moreover $(T, m_T, u_T, \Delta_T, \varepsilon_T)$ is a graded braided bialgebra in \mathcal{M} .

We are now able to state the universal property of the tensor bialgebra.

THEOREM 5.3. Let H be a braided bialgebra in a cocomplete abelian braided monoidal category (\mathcal{M},c) . Assume that the tensor product commutes with direct sums.

Let $(M, \mu_M^r, \mu_M^l, \rho_M^r, \rho_M^l)$ be in ${}^H_H \mathcal{M}^H_H$. Let $T = T_H(M)$ be the tensor algebra.

Let E be a braided bialgebra in \mathcal{M} .

Let $f_H: H \to E$ be a bialgebra homomorphism and let $f_M: M \to E$ be a morphism of H-bimodules, where E is a bimodule via f_H . Assume that

$$\Delta_E f_M = (f_H \otimes f_M) \, \rho_M^l + (f_M \otimes f_H) \, \rho_M^r$$

(i.e. f_M is a coderivation of E with domain the E-bicomodule M, where M is regarded as a bicomodule via f_H). Then there is a unique algebra homomorphism $f: T_H(M) \to E$ such that $fi_0 = f_H$ and $fi_1 = f_M$, where $i_n: M^{\otimes_H n} \to T_H(M)$ denotes the canonical injection.

$$M \xrightarrow{i_1} T_H(M)$$

$$f_M \downarrow \qquad \qquad \downarrow i_0$$

$$E \xleftarrow{f_H} H$$

 $Moreover\ f\ is\ a\ bialgebra\ homomorphism.$

Proof. By Theorem 3.9, there is a unique algebra homomorphism $f: T \to E$ such that $fi_0 = f_H$ and $fi_1 = f_M$. The rest of the proof is similar to that of Theorem 4.7.

6. Braided Bialgebras of Type One

Even if not all the assumptions are always needed, in this section $((\mathcal{M}, \otimes, \mathbf{1}), c)$ will denote a cocomplete and complete abelian coabelian braided monoidal category satisfying AB5 and such that the tensor product commutes with direct sums.

LEMMA 6.1. Let $(B, m_B, u_B, \Delta_B, \varepsilon_B)$ be a graded braided bialgebra in (\mathcal{M}, c) . Then

- 1) $(B_0, m_{0,0}, u_0, \Delta_{0,0}, \varepsilon_0)$ is a braided bialgebra. Moreover both p_0^B and i_0^B are bialgebra homomorphisms.
 - 2) $(B_1, \mu_{B_1}^r = m_{1,0}, \mu_{B_1}^l = m_{0,1}, \rho_{B_1}^r = \Delta_{1,0}, \rho_{B_1}^l = \Delta_{0,1})$ is an object in $^{B_0}_{B_0}\mathcal{M}^{B_0}_{B_0}$. 3) i_1^B is a morphism of B_0 -bimodules, where B is a B_0 -bimodule via i_0^B , and we have

$$\Delta_B i_1^B = \left(i_0^B \otimes i_1^B\right) \Delta_{0,1} + \left(i_1^B \otimes i_0^B\right) \Delta_{1,0}$$

(i.e. i_1^B is a coderivation of B with domain the B-bicomodule B_1 , where B_1 is regarded as a bicomodule via i_0^B).

4) p_1^B is a morphism of B_0 -bicomodules, where B is a B_0 -bicomodule via p_0^B , and we have

$$p_{1}^{B}m_{B}=m_{0,1}\left(p_{0}^{B}\otimes p_{1}^{B}\right)+m_{1,0}\left(p_{1}^{B}\otimes p_{0}^{B}\right)$$

(i.e. p_1^B is a derivation of B with values in the B-bimodule B_1 , where B_1 is regarded as a bimodule $via p_0^B$).

Proof. We prove the statements by means of (22).

1) By Proposition 2.5, $(B_0, \Delta_0 = \Delta_{0,0}, \varepsilon_0 = \varepsilon_B i_0^B)$ is a coalgebra in \mathcal{M} and i_0^B is a coalgebra homomorphism. By Proposition 3.4 $(B_0, m_0 = m_{0,0}, u_0 = p_0^B u_B)$ is an algebra in \mathcal{M} and p_0^B is an algebra homomorphism.

By (16), (5) and by naturality of the braiding we obtain

$$(m_{0,0} \otimes m_{0,0}) (B_0 \otimes c_{B_0,B_0} \otimes B_0) (\Delta_{0,0} \otimes \Delta_{0,0})$$

$$= (p_0^B \otimes p_0^B) (m_B \otimes m_B) (B \otimes c_{B,B} \otimes B) (\Delta_B \otimes \Delta_B) (i_0^B \otimes i_0^B)$$

$$\stackrel{(22)}{=} (p_0^B \otimes p_0^B) \Delta_B m_B (i_0^B \otimes i_0^B) = \Delta_{0,0} m_{0,0}.$$

Moreover

$$m_1(\varepsilon_0 \otimes \varepsilon_0) = m_1(\varepsilon_B \otimes \varepsilon_B) (i_0^B \otimes i_0^B) \stackrel{(22)}{=} \varepsilon_B m_B (i_0^B \otimes i_0^B) \stackrel{(17)}{=} \varepsilon_B i_0^B m_{0,0} = \varepsilon_0 m_{0,0}$$

so that $(B_0, m_{0,0}, u_0, \Delta_{0,0}, \varepsilon_0)$ is a braided bialgebra. Let us check that i_0^B is an algebra homomorphism. We have

$$m_B (i_0^B \otimes i_0^B) \stackrel{(17)}{=} i_0^B m_{0.0}.$$

Moreover, by Proposition 3.4, $u = i_0^B p_0 u = i_0^B u_0$. Thus i_0^B is an algebra homomorphism.

Similarly, by means of (5) and of Proposition 2.5, we get that p_0^B is also a coalgebra homomorphism.

2) By (16), (5) and by naturality of the braiding we obtain

$$(31) \qquad (p_0^B \otimes p_1^B) (m_B \otimes m_B) (B \otimes c_{B,B} \otimes B) (\Delta_B \otimes \Delta_B)$$

$$= \begin{bmatrix} (m_{0,0} \otimes m_{1,0}) (B_0 \otimes c_{B_1,B_0} \otimes B_0) [\Delta_{0,1} p_1^B \otimes \Delta_{0,0} p_0^B] + \\ + (m_{0,0} \otimes m_{0,1}) (B_0 \otimes c_{B_0,B_0} \otimes B_1) [\Delta_{0,0} p_0^B \otimes \Delta_{0,1} p_1^B] \end{bmatrix}$$

Hence, by (5), (17) and by naturality of the braiding we obtain

$$\Delta_{0,1}m_{0,1} = \left(p_0^B \otimes p_1^B\right) \Delta_B m_B \left(i_0^B \otimes i_1^B\right)$$

$$\stackrel{(22)}{=} \left(p_0^B \otimes p_1^B\right) \left(m_B \otimes m_B\right) \left(B \otimes c_{B,B} \otimes B\right) \left(\Delta_B \otimes \Delta_B\right) \left(i_0^B \otimes i_1^B\right)$$

$$\stackrel{(31)}{=} \left(m_{0,0} \otimes m_{0,1}\right) \left(B_0 \otimes c_{B_0,B_0} \otimes B_1\right) \left[\Delta_{0,0} \otimes \Delta_{0,1}\right]$$

Similarly we get (24), (25) and (26).

THEOREM 6.2. Let H be a braided bialgebra in a cocomplete and complete abelian and coabelian braided monoidal category (\mathcal{M}, c) satisfying AB5. Assume that the tensor product commutes with direct sums.

Let $(M, \mu_M^r, \mu_M^l, \rho_M^r, \rho_M^l)$ be in ${}^H_H\mathcal{M}^H_H$. Let $T^c = T^c_H(M)$ be the cotensor coalgebra.

Let B be a graded bialgebra, let $g_H: B_0 \to H$ be a bialgebra homomorphism and let $g_M: B_1 \to M$ be a morphism of H-bicomodules, where B_1 is an H-bicomodule via g_H , and a morphism of B_0 -bimodules, where M is a B_0 -bimodule via g_H .

Then there is a unique coalgebra homomorphism $f: B \to T^c_H(M)$ such that

$$p_0^{T^c} \circ f = g_H \circ p_0^B \qquad \text{and} \qquad p_1^{T^c} \circ f = g_M \circ p_1^B.$$

Moreover f is a graded bialgebra homomorphism.

Proof. By Theorem 2.20, there is a unique coalgebra homomorphism $f: B \to T_H^c(M)$ such that $p_0^C \circ f = g_H \circ p_0^B$ and $p_1^{T^c} \circ f = g_M \circ p_1^B$. Moreover f is a morphism of graded coalgebras. On the other hand, set

$$f_H := g_H \circ p_0^B$$
 and $f_M := g_M \circ p_1^B$.

By Lemma 6.1, we have

$$p_1^B m_B = m_{0,1} \left(p_0^B \otimes p_1^B \right) + m_{1,0} \left(p_1^B \otimes p_0^B \right).$$

Since $g_M: B_1 \to M$ is a morphism of B_0 -bimodules, where M is a B_0 -bimodule via g_H , we get

$$f_{M} \circ m_{B} = g_{M} \circ p_{1}^{B} \circ m_{B}$$

$$= \mu_{M}^{l} \circ (g_{H} \otimes g_{M}) \circ (p_{0}^{B} \otimes p_{1}^{B}) + \mu_{M}^{r} \circ (g_{M} \otimes g_{H}) \circ (p_{1}^{B} \otimes p_{0}^{B})$$

$$= \mu_{M}^{l} \circ (f_{H} \otimes f_{M}) + \mu_{M}^{r} \circ (f_{M} \otimes f_{H}).$$

Hence, by Proposition 2.19 and Theorem 4.7, there is a unique coalgebra homomorphism $f': B \to T_H^c(M)$ such that $p_0f' = f_H$ and $p_1f' = f_M$. Moreover f' is a bialgebra homomorphism. By uniqueness f = f'.

THEOREM 6.3. Let H be a braided bialgebra in a cocomplete and complete abelian and coabelian braided monoidal category (\mathcal{M}, c) satisfying AB5. Assume that the tensor product commutes with direct sums.

Let $(M, \mu_M^r, \mu_M^l, \rho_M^r, \rho_M^l)$ be in ${}^H_H \mathcal{M}^H_H$. Let $T = T_H(M)$ be the tensor algebra. Let B be a graded bialgebra, let $g_H : H \to B_0$ be a bialgebra homomorphism and let $g_M : M \to B_1$

be a morphism of H-bimodules, where B_1 is an H-bimodule via g_H , and a morphism of B_0 -bicomodules, where M is a B_0 -bicomodule via g_H .

Then there is a unique algebra homomorphism $f: T_H(M) \to B$ such that

$$f \circ i_0^T = i_0^B \circ g_H$$
 and $f \circ i_1^T = i_1^B \circ g_M$.

Moreover f is a graded bialgebra homomorphism.

Proof. It is analogous to the proof of Theorem 6.2.

6.4. Let $f: X \to X'$ be a graded homomorphism in \mathcal{M} . Hence $f = \bigoplus_{n \in \mathbb{N}} f_n$ for suitable morphisms $f_n: X_n \to X'_n$, for every $n \in \mathbb{N}$. Write

$$f_n = \psi_n \circ \varphi_n$$

where $\varphi_n: X_n \to \operatorname{Im}(f_n)$ is an epimorphism and $\psi_n: \operatorname{Im}(f_n) \to X'_n$ is a monomorphism. Since \mathcal{M} satisfies AB5, it is in particular an AB4 category. Hence coproducts are both left and right exact. Thus

$$\varphi_f := \bigoplus_{n \in \mathbb{N}} \varphi_n : X \to \bigoplus_{n \in \mathbb{N}} \operatorname{Im}(f_n)$$
 and $\psi_f := \bigoplus_{n \in \mathbb{N}} \psi_n : \bigoplus_{n \in \mathbb{N}} \operatorname{Im}(f_n) \to X'$

are an epimorphism and a monomorphism respectively. Moreover

$$f = \bigoplus_{n \in \mathbb{N}} f_n = (\bigoplus_{n \in \mathbb{N}} \psi_n) \circ (\bigoplus_{n \in \mathbb{N}} \varphi_n) = \psi_f \circ \varphi_f.$$

In particular we obtain $\operatorname{Im}(f) = \bigoplus_{n \in \mathbb{N}} \operatorname{Im}(f_n)$ so that $\operatorname{Im}(f)_n = \operatorname{Im}(f_n)$ for every $n \in \mathbb{N}$. We would like to outline that the last statements do not in general hold without assuming AB5. Clearly if $f_n = \operatorname{Id}_{X_n}$ we can choose $\psi_n = \varphi_n = \operatorname{Id}_{X_n}$.

PROPOSITION 6.5. Let $f: A \to A'$ be a graded braided bialgebra homomorphism in (\mathcal{M}, c) . Then $\operatorname{Im}(f)$ can be endowed with a unique graded braided bialgebra structure such that $\varphi_f: A \to \operatorname{Im}(f)$ and $\psi_f: \operatorname{Im}(f) \to A'$ are graded braided bialgebra homomorphisms.

- 1) φ_f is epimorphism and ψ_f is a monomorphism in \mathcal{M} .
- 2) $\operatorname{Im}(f)_n = \operatorname{Im}(f_n)$ for every $n \in \mathbb{N}$.

Proof. Set $B = \operatorname{Im}(f)$, $B_n = \operatorname{Im}(f_n)$, $\varphi = \varphi_f$ and $\psi = \psi_f$.

Since $(\operatorname{Im}(f), \varphi) = \operatorname{Coker}(\ker(f))$, it is clear that $\operatorname{Im}(f)$ can be endowed with a unique algebra structure such that φ is an algebra homomorphism.

Since $(\operatorname{Im}(f), \psi) = \ker(\operatorname{Coker}(f))$, it is clear that $\operatorname{Im}(f)$ can be endowed with a unique coalgebra structure such that ψ is a coalgebra homomorphism.

Let us check that φ is also a coalgebra homomorphism. We have

$$(\psi \otimes \psi) \circ \Delta_B \circ \varphi = \Delta_{A'} \circ \psi \circ \varphi = \Delta_{A'} \circ f = (f \otimes f) \circ \Delta_A = (\psi \otimes \psi) \circ (\varphi \otimes \varphi) \circ \Delta_A,$$

$$\varepsilon_B \circ \varphi = \varepsilon_{A'} \circ \psi \circ \varphi = \varepsilon_{A'} \circ f = \varepsilon_A.$$

Since ψ is a monomorphism and \mathcal{M} is a coabelian monoidal category, we get that $\psi \otimes \psi$ is a monomorphism so that

$$\Delta_B \circ \varphi = (\varphi \otimes \varphi) \circ \Delta_A$$
 and $\varepsilon_B \circ \varphi = \varepsilon_A$

and hence φ is also a coalgebra homomorphism.

Similarly it can be proven that ψ is also an algebra homomorphism.

Let us check that B is a braided bialgebra. Since φ is a bialgebra homomorphism, A is a braided bialgebra and by naturality of the braiding one computes

$$(m_B \otimes m_B) \circ (B \otimes c_{B,B} \otimes B) \circ (\Delta_B \otimes \Delta_B) \circ (\varphi \otimes \varphi) = \Delta_B \circ m_B \circ (\varphi \otimes \varphi)$$
$$\varepsilon_B \circ m_B \circ (\varphi \otimes \varphi) = m_1 \circ (\varepsilon_B \otimes \varepsilon_B) \circ (\varphi \otimes \varphi).$$

Since $\varphi \otimes \varphi$ is an epimorphism we get (22) for B.

By Proposition 2.4 in order to prove that $B = \bigoplus_{n \in \mathbb{N}} B_n$ is a graded coalgebra we have to check that

there are morphisms $\Delta_{a,b}^B: B_{a+b} \to B_a \otimes B_b$, such that $(p_a^B \otimes p_b^B) \circ \Delta_B = \Delta_{a,b}^B \circ p_{a+b}^B$, for every $a,b \in \mathbb{N}$. Set

$$\Delta_{a,b}^B := \left(B_{a+b} \overset{i_{a+b}^B}{\to} B \overset{\Delta_B}{\to} B \otimes B \overset{p_a^B \otimes p_b^B}{\to} B_a \otimes B_b \right).$$

Since φ is a graded coalgebra homomorphism, by (5), one easily checks that

$$\Delta_{a,b}^{B} \circ p_{a+b}^{B} \circ \varphi = (p_{a}^{B} \otimes p_{b}^{B}) \circ \Delta_{B} \circ \varphi$$

Since φ is an epimorphism, we get $\Delta_{a,b}^B \circ p_{a+b}^B = (p_a^B \otimes p_b^B) \circ \Delta_B$ that is B is a graded coalgebra. Similarly one proves that B is also a graded algebra and hence a graded braided bialgebra.

Clearly both φ and ψ are graded braided bialgebra homomorphism.

1) and 2) follow by
$$(6.4)$$
.

THEOREM 6.6. Let H be a braided bialgebra in a cocomplete and complete abelian coabelian braided monoidal category (\mathcal{M},c) satisfying AB5. Assume that the tensor product commutes with direct sums.

Let $(M, \mu_M^r, \mu_M^l, \rho_M^r, \rho_M^l)$ be in ${}^H_H \mathcal{M}_H^H$. Let $T = T_H(M)$ and $T^c = T_H^c(M)$. Then there is a unique algebra homomorphism $F = F_{H,M} : T \to T^c$ such that $Fi_0^T = i_0^{T^c}$ and $Fi_1^T = i_1^{T^c}$:

$$M \xrightarrow{i_1^{T^c}} T_H(M)$$

$$\downarrow i_1^{T^c} \downarrow \qquad \qquad \downarrow i_0^{T^c}$$

$$T_H^c(M) \xleftarrow{i_0^{T^c}} H$$

Moreover F is a graded bialgebra homomorphism such that $F_0 = \operatorname{Id}_H$ and $F_1 = \operatorname{Id}_M$. Write $F = \psi_F \circ \varphi_F$ as in Proposition 6.5, where $\varphi_F : T \to \operatorname{Im}(F)$ and $\psi_F : \operatorname{Im}(F) \to T^c$ are graded bialgebra homomorphisms, φ_F is epimorphism and ψ_F is a monomorphism in M. We have:

- 1) $\operatorname{Im}(F)$ is a graded braided bialgebra such that $\operatorname{Im}(F)_0 = H$ and $\operatorname{Im}(F)_1 = M$.
- 2) φ_F is the unique algebra homomorphism such that $\varphi_F \circ i_0^T = i_0^{\operatorname{Im}(F)}$ and $\varphi_F \circ i_1^T = i_1^{\operatorname{Im}(F)}$ as in Theorem 3.11.
- 3) ψ_F is the unique coalgebra homomorphism such that $p_0^{T^c} \circ \psi_F = p_0^{\operatorname{Im}(F)}$ and $p_1^{T^c} \circ \psi_F = p_1^{\operatorname{Im}(F)}$ as in Theorem 2.22.

Proof. Set B = Im(F). In view of Theorem 4.5 $T^c = T_c^c(M)$ is a graded braided bialgebra.

By Corollary 6.3 there is a unique algebra homomorphism $F: T \to T^c$ such that $F \circ i_0^T = i_0^{T^c}$ and $F \circ i_1^T = i_1^{T^c}$. Moreover F is a graded bialgebra homomorphism.

By Proposition 6.5, we have that $\operatorname{Im}(F)$ is a graded braided bialgebra such that $\operatorname{Im}(F)_n = \operatorname{Im}(F_n)$ for every $n \in \mathbb{N}$. Moreover we can write $F = \psi \circ \varphi$ where $\varphi = \varphi_F : T \to \operatorname{Im}(F)$ and $\psi = \psi_F : \operatorname{Im}(F) \to T^c$ are graded braided bialgebra homomorphisms, φ is an epimorphism and ψ is a monomorphism in \mathcal{M} . Since F is graded we have

$$F_0 = p_0^{T^c} \circ F \circ i_0^T = p_0^{T^c} \circ i_0^{T^C} = \mathrm{Id}_H \qquad \text{and} \qquad F_1 = p_1^{T^c} \circ F \circ i_1^T = p_1^{T^c} \circ i_1^{T^C} = \mathrm{Id}_M.$$

Then $\operatorname{Im}(F)_0 = \operatorname{Im}(F_0) = H$ and $\operatorname{Im}(F)_1 = \operatorname{Im}(F_1) = M$. Now, since $F_0 = \operatorname{Id}_H$ and $F_1 = \operatorname{Id}_M$ we can choose $\varphi_0 = \psi_0 = \operatorname{Id}_H$ and $\varphi_1 = \psi_1 = \operatorname{Id}_M$ so that $\varphi \circ i_n^T = i_n^B$ and $p_n^B = p_n^{T^c} \circ \psi$ for n = 0, 1.

Definition 6.7. Take the notations and assumptions of Theorem 6.6. Following [Ni, page 1533], let

$$(H[M], i_{H[M]}) = (\operatorname{Im}(F), \psi_F).$$

Then H[M] is a graded braided bialgebra such that $H[M]_0 = H$ and $H[M]_1 = M$. This will be called the braided bialgebra of type one associated to H and M. THEOREM 6.8. Let $(B, m_B, u_B, \Delta_B, \varepsilon_B)$ be a braided graded bialgebra in a cocomplete and complete abelian coabelian braided monoidal category (\mathcal{M}, c) satisfying AB5. Assume that the tensor product commutes with direct sums.

1) There is a unique algebra homomorphism

$$\varphi: T_{B_0}(B_1) \to B$$

such that $\varphi \circ i_0^T = i_0^B$ and $\varphi \circ i_1^T = i_1^B$. Moreover φ is a graded bialgebra homomorphism.

2) There is a unique coalgebra homomorphism

$$\psi: B \to T^c_{B_0}(B_1)$$

such that $p_0^{T^c} \circ \psi = p_0^B$ and $p_1^{T^c} \circ \psi = p_1^B$. Moreover ψ is a graded bialgebra homomorphism. 3) $\psi \circ \varphi = F$ where $F: T_{B_0}(B_1) \to T_{B_0}^c(B_1)$ is the morphism defined in Theorem 6.6. In particular F is a graded bialgebra homomorphism.

Furthermore the following assertions are equivalent

- (a) B is strongly \mathbb{N} -graded both as an algebra and a coalgebra.
- (b) φ is an epimorphism and ψ is a monomorphism.
- (c) $\varphi = \varphi_F$ and $\psi = \psi_F$.
- (d) $B = B_0[B_1]$ is the braided bialgebra of type one associated to B_0 and B_1 .

Proof. 1)-2) By Lemma 6.1, we can apply Theorems 6.3 and 6.2.

3) By Theorem 6.6, there is a unique algebra homomorphism $F: T_{B_0}(B_1) \to T_{B_0}^c(B_1)$ such that $F \circ i_0^T = i_0^{T^c}$ and $F \circ i_1^T = i_1^{T^c}$. Moreover F is a graded bialgebra homomorphism such that $F_0 = \operatorname{Id}_H$ and $F_1 = \operatorname{Id}_M$. Since

$$\psi \circ \varphi \circ i_0^T = \psi \circ i_0^B \circ \varphi_0 = i_0^{T^c} \circ \psi_0 \circ \varphi_0 = i_0^{T^c}, \qquad \psi \circ \varphi \circ i_1^T = \psi \circ i_1^B \circ \varphi_1 = i_1^{T^c} \circ \psi_1 \circ \varphi_1 = i_1^{T^c},$$

we infer that $F = \psi \circ \varphi$. By Proposition 6.5, $\operatorname{Im}(F)$ can be endowed with a unique graded braided bialgebra structure such that $\varphi_F : T \to \operatorname{Im}(F)$ and $\psi_F : \operatorname{Im}(F) \to T^c$ are graded braided bialgebra homomorphisms. Furthermore $\operatorname{Im}(F)_n = \operatorname{Im}(F_n)$ for every $n \in \mathbb{N}$, we can write $F = \psi_F \circ \varphi_F$, φ_F is an epimorphism and ψ_F is a monomorphism in \mathcal{M} .

- $(a) \Leftrightarrow (b)$ It follows by Theorems 3.11 and 2.22.
- $(b) \Rightarrow (c)$ By hypothesis, since the category \mathcal{M} is abelian, B identifies as an object with Im $(F) = B_0[B_1]$. By uniqueness in 2) and 3) of Theorem 6.6 and in view of 1) and 2) above, we get $\varphi = \varphi_F$ and $\psi = \psi_F$.
- $(c) \Rightarrow (d)$ Since the graded braided bialgebra structure of B and of Im (F) make $\varphi_F : T \to B$ and $\psi_F : B \to T^c$ graded braided bialgebra homomorphisms, by Proposition 6.5, the two structures coincide.
- $(d) \Rightarrow (c)$ Since $B = \operatorname{Im}(F)$, it follows by 2) and 3) in Theorem 6.6.

$$(c) \Rightarrow (b)$$
 It is trivial.

REMARK 6.9. Recall that φ is an epimorphism if and only if it fulfills one of the equivalent conditions in Theorem 3.11 and ψ is a monomorphism if and only if it fulfills one of the equivalent conditions in Theorem 2.22. In particular we have the following result.

THEOREM 6.10. Let $(B, m_B, u_B, \Delta_B, \varepsilon_B)$ be a braided graded bialgebra in a cocomplete and complete abelian coabelian braided monoidal category (\mathcal{M}, c) satisfying AB5. Assume that the tensor product commutes with direct sums.

Then B is the braided bialgebra of type one $B_0[B_1]$ associated to B_0 and B_1 if and only if

$$\left(B\left[2\right],\nu_{2}^{B}\right)=B\left[1\right]^{2}\qquad and \qquad \left(B\left(2\right),\sigma_{2}^{B}\right)=B_{0}^{\wedge_{B}^{2}}.$$

7. Radford-Majid Bosonization

In this section $((\mathcal{M}, \otimes, \mathbf{1}), c)$ will denote a cocomplete and complete abelian coabelian braided monoidal category satisfying AB5 and such that the tensor product commutes with direct sums.

7.1. Let $(H, m_H, u_H, \Delta_H, \varepsilon_H, S_H)$ be a braided graded Hopf algebra with invertible antipode S_H . By [Bes, Theorem 3.4.3], $(H\mathcal{YD}(\mathcal{M}), \otimes, \mathbf{1}), \Psi$ is a braided monoidal category, where, for every $(U, \mu_U^l, \rho_U^l), (V, \mu_V^l, \rho_V^l) \in {}_H^H \mathcal{YD}(\mathcal{M}),$ the braiding is defined by

$$\Psi_{U,V} := \left(\mu_V^l \otimes U\right) \circ \left(H \otimes c_{U,V}\right) \circ \left(\rho_U^l \otimes V\right)$$

with inverse given by

$$\Psi_{U,V}^{-1} := \left(U \otimes \mu_V^l \right) \circ \left(U \otimes c_{H,V}^{-1} \right) \circ \left(c_{U,V}^{-1} \otimes S_H^{-1} \right) \circ \left(V \otimes c_{U,H}^{-1} \right) \circ \left(V \otimes \rho_U^l \right).$$

7.2. In view of [Bes, Lemma 3.9.4], recall that given an object (V, μ_V^l, ρ_V^l) in the monoidal category $({}^{H}_{H}\mathcal{YD}(\mathcal{M}), \otimes, \mathbf{1}), \text{ then }$

$$\left(V \otimes H, \mu_{V \otimes H}^{l}, \mu_{V \otimes H}^{r}, \rho_{V \otimes H}^{l}, \rho_{V \otimes H}^{r}\right)$$

is an object in ${}^H_H\mathcal{M}^H_H$ where

$$\mu_{V\otimes H}^{l} = (\mu_{V}^{l}\otimes m_{H})\circ (H\otimes c_{H,V}\otimes H)\circ (\Delta_{H}\otimes V\otimes H), \qquad \mu_{V\otimes H}^{r} = V\otimes m_{H},$$

$$\rho_{V\otimes H}^{l} = (m_{H}\otimes V\otimes H)\circ (H\otimes c_{V,H}\otimes H)\circ (\rho_{V}^{l}\otimes \Delta_{H}), \qquad \rho_{V\otimes H}^{r} = V\otimes \Delta_{H}.$$

Given a braided bialgebra $(Q, \mu_Q^l, \rho_Q^l), m_Q, u_Q, \Delta_Q, \varepsilon_Q)$ in the braided monoidal category $\left(\left({}_{H}^{H}\mathcal{YD}\left(\mathcal{M}\right),\otimes,\mathbf{1}\right),\Psi\right)$ then the Radford-Majid bosonization of Q

$$Q \rtimes H$$

is the object $Q \otimes H$ endowed with the following braided bialgebra structure:

$$m_{Q\rtimes H} = (m_Q\otimes m_H)\circ (Q\otimes \mu_Q^l\otimes H\otimes H)\circ (Q\otimes H\otimes c_{H,Q}\otimes H)\circ (Q\otimes \Delta_H\otimes Q\otimes H),$$

$$u_{Q \rtimes H} = (u_Q \otimes u_H) \circ \Delta_1,$$

$$\Delta_{Q \rtimes H} = (Q \otimes m_H \otimes Q \otimes H) \circ (Q \otimes H \otimes c_{Q,H} \otimes H) \circ (Q \otimes \rho_Q^l \otimes H \otimes H) \circ (\Delta_Q \otimes \Delta_H),$$

$$\varepsilon_{Q \rtimes H} = m_1 \circ (\varepsilon_Q \otimes \varepsilon_H).$$

and the above structures of object in ${}^H_H\mathcal{M}^H_H$.

LEMMA 7.3. Let $(H, m_H, u_H, \Delta_H, \varepsilon_H, S_H)$ be a braided graded Hopf algebra with invertible antipode S_H in $((\mathcal{M}, \otimes, \mathbf{1}), c)$.

Let $(Q = \bigoplus_{n \in \mathbb{N}} Q_n, \mu^l, \rho^l)$, $m, u, \Delta, \varepsilon)$ be a graded braided bialgebra in the braided monoidal category $(H\mathcal{YD}(\mathcal{M}), \otimes, \mathbf{1}), \Psi$. Then

$$Q \rtimes H = \bigoplus_{n \in \mathbb{N}} (Q_n \otimes H)$$

is a graded braided bialgebra in the monoidal category $(\mathcal{M}, \otimes, \mathbf{1})$ and

$$m_{a,b}^{Q \rtimes H} = (m_{a,b} \otimes m_H) \circ (Q_a \otimes \mu_{Q_b}^l \otimes H \otimes H) \circ (Q_a \otimes H \otimes c_{H,Q_b} \otimes H) \circ (Q_a \otimes \Delta_H \otimes Q_b \otimes H),$$

$$\Delta_{a,b}^{Q \rtimes H} = (Q_a \otimes m_H \otimes Q_b \otimes H) \circ (Q_a \otimes H \otimes c_{Q_b,H} \otimes H) \circ (Q_a \otimes \rho_{Q_b}^l \otimes H \otimes H) \circ (\Delta_{a,b} \otimes \Delta_H).$$

- Moreover, for every $a,b \in \mathbb{N}$, we have that

 1) $m_{a,b}^{Q \rtimes H}$ is an epimorphism whenever $m_{a,b}$ is an epimorphism

 2) $\Delta_{a,b}^{Q \rtimes H}$ is a monomorphism whenever $\Delta_{a,b}$ is a monomorphism.

Proof. Set $B = Q \times H$ and $B_n = Q_n \otimes H$ for every $n \in \mathbb{N}$. Then we have

$$i_n^B = i_n^Q \otimes H$$
 and $p_n^B = p_n^Q \otimes H$

for every $n \in \mathbb{N}$.

By Proposition 3.3, in order to prove that $Q \rtimes H$ is a graded algebra and that $m_{a,b}^{Q \rtimes H}$ is the required morphism, it is enough to prove that (17) holds for every $a, b \in \mathbb{N}$. By naturality of the braiding, the fact that i_b^Q is left *H*-linear and *Q* fulfills (17), we obtain $m_B \circ (i_a^B \otimes i_b^B) = i_{a+b}^B \circ m_{a,b}^B$. Similarly, by Proposition 2.4, $Q \times H$ is a graded coalgebra and hence a graded braided bialgebra.

Now, in view of 7.1, for every (U, μ_U^l, ρ_U^l) , $(V, \mu_V^l, \rho_V^l) \in ^H_H \mathcal{YD}(\mathcal{M})$ we have that the morphism

$$\Psi_{U,V} := \left(\mu_V^l \otimes U\right) \circ \left(H \otimes c_{U,V}\right) \circ \left(\rho_U^l \otimes V\right)$$

is invertible. Recall that, by [BD, Example 4.1.2], using the left adjoint action

$$ad := m_H \circ (m_H \otimes H) \circ (H \otimes c_{H,H}) \circ (H \otimes S \otimes H) \circ (\Delta_H \otimes H)$$

for H and the regular coaction, we have that $H_{ad} := (H, ad, \Delta_H)$ is an object in ${}^H_H \mathcal{YD}(\mathcal{M})$. Dually using the left coadjoint coaction

$$coad := (m_H \otimes H) \circ (H \otimes S \otimes H) \circ (H \otimes c_{H,H}) \circ (\Delta_H \otimes H) \circ \Delta_H$$

for H and the regular action, we have that $H^{coad} := (H, m_H, coad)$ is an object in ${}^H_H \mathcal{YD}(\mathcal{M})$. Then we have

$$m_{a,b}^{Q\rtimes H} = (m_{a,b}\otimes m_H)\circ (Q_a\otimes \Psi_{Q_b,H_{ad}}\otimes H) \quad \text{and} \quad \Delta_{a,b}^{Q\rtimes H} = (Q_a\otimes \Psi_{H^{coad},Q_b}\otimes H)\circ (\Delta_{a,b}\otimes \Delta_H).$$

Since $\Psi_{Q_b,H_{ad}}$ is an isomorphism we get that $m_{a,b}^{Q\rtimes H}$ is an epimorphism whenever $m_{a,b}$ is an epimorphism and $\Delta_{a,b}^{Q\rtimes H}$ is a monomorphism whenever $\Delta_{a,b}$ is a monomorphism.

THEOREM 7.4. Let $(H, m_H, u_H, \Delta_H, \varepsilon_H, S_H)$ be a braided graded Hopf algebra with invertible antipode S_H in a cocomplete and complete abelian coabelian braided monoidal category $((\mathcal{M}, \otimes, \mathbf{1}), c)$ satisfying AB5. Assume that the tensor product commutes with direct sums.

Let (V, μ_V^l, ρ_V^l) be in ${}^H_H \mathcal{YD}(\mathcal{M})$ and let $\mathbf{1}[V]$ be the braided bialgebra of type one associated to $\mathbf{1}$ and V in ${}^H_H \mathcal{YD}(\mathcal{M})$. Then $V \otimes H$ is an object in ${}^H_H \mathcal{M}_H^H$ and

$$\mathbf{1}\left[V\right]\rtimes H=H\left[V\otimes H\right]$$

is the braided bialgebra of type one associated to H and $V \otimes H$ in \mathcal{M} .

Proof. $V \otimes H$ is an object in ${}^H_H \mathcal{M}^H_H$ as in 7.2. Let $T_1(V)$ be the tensor algebra of V in the monoidal category $({}^H_H \mathcal{YD}(\mathcal{M}), \otimes, \mathbf{1})$. Clearly the isomorphisms

$$g_H := l_H^{-1} : H \longrightarrow \mathbf{1} \otimes H$$
 and $g_{V \otimes H} := \mathrm{Id}_{V \otimes H} : V \otimes H \longrightarrow V \otimes H$

are a braided bialgebra homomorphism and a morphism in ${}^H_H\mathcal{M}^H_H$ respectively. By Lemma 7.3, $T_1(V) \rtimes H = \bigoplus_{n \in \mathbb{N}} (V^{\otimes n} \otimes H)$ is a graded braided bialgebra in the monoidal category $(\mathcal{M}, \otimes, \mathbf{1})$. By Theorem 6.3, there is a unique algebra homomorphism $f: T_H(V \otimes H) \to T_1(V) \rtimes H$ such that

$$f\circ i_0^{T_H(V\otimes H)} = \left(i_0^{T_1(V)}\otimes H\right)\circ l_H^{-1} \qquad \text{and} \qquad f\circ i_1^{T_H(V\otimes H)} = i_1^{T_1(V)}\otimes H.$$

Moreover f is a graded bialgebra homomorphism. By Theorem 3.11, f is an epimorphism. In fact, by Proposition 3.8, $m_{a,b}^{T_1(V)}$ is an epimorphism for every $a,b\in\mathbb{N}$ and hence, in view of Lemma 7.3, $m_{a,b}^{T_1(V)\rtimes H}$ is an epimorphism too.

The isomorphisms

$$g_H^c := l_H : \mathbf{1} \otimes H \longrightarrow H$$
 and $g_{V \otimes H}^c := \mathrm{Id}_{V \otimes H} : V \otimes H \longrightarrow V \otimes H$

are a braided bialgebra homomorphism and a morphism in ${}^H_H\mathcal{M}^H_H$ respectively. By Lemma 7.3, $T^c_1(V) \rtimes H = \bigoplus_{n \in \mathbb{N}} (V^{\otimes n} \otimes H)$ is a graded braided bialgebra in the monoidal category $(\mathcal{M}, \otimes, \mathbf{1})$. Then, by Theorem 6.2, there is a unique coalgebra homomorphism $f^c: T^c_1(V) \rtimes H \to T^c_H(V \otimes H)$ such that

$$p_0^{T_H^c(V\otimes H)}\circ f^c=l_H\circ \left(p_0^{T_1^c(V)}\otimes H\right)\qquad\text{and}\qquad p_1^{T_c^c(V\otimes H)}\circ f^c=\left(p_1^{T_1^c(V)}\otimes H\right).$$

Moreover f^c is a graded bialgebra homomorphism. By Theorem 2.22 , f^c is a monomorphism. In fact, by Proposition 2.13, $\Delta_{a,b}^{T_1^c(V)}$ is a monomorphism for every $a,b\in\mathbb{N}$ and hence, in view of Lemma 7.3, $\Delta_{a,b}^{T_1^c(V)\rtimes H}$ is a monomorphism too. Using the same notations of Theorem 6.6, let us check that

$$F_{H,V\otimes H} = f^c \circ (F_{1,V} \rtimes H) \circ f.$$

By Theorem 6.6 it is enough to check that

$$f^c \circ (F_{\mathbf{1},V} \rtimes H) \circ f \circ i_0^{T_H(V \otimes H)} = i_0^{T_H^c(V \otimes H)} \qquad \text{and} \qquad f^c \circ (F_{\mathbf{1},V} \rtimes H) \circ f \circ i_1^{T_H(V \otimes H)} = i_1^{T_H^c(V \otimes H)} = i_1^{T_$$

We have

$$\begin{split} f^c \circ (F_{\mathbf{1},V} \rtimes H) \circ f \circ i_0^{T_H(V \otimes H)} &= f^c \circ (F_{\mathbf{1},V} \otimes H) \circ \left(i_0^{T_1(V)} \otimes H\right) \circ l_H^{-1} = f^c \circ \left(i_0^{T_1^c(V)} \otimes H\right) \circ l_H^{-1} \\ f^c \circ (F_{\mathbf{1},V} \rtimes H) \circ f \circ i_1^{T_H(V \otimes H)} &= f^c \circ (F_{\mathbf{1},V} \otimes H) \circ \left(i_1^{T_1(V)} \otimes H\right) = f^c \circ \left(i_1^{T_1^c(V)} \otimes H\right). \end{split}$$

Therefore we have to prove that

$$(32) \qquad f^c \circ \left(i_0^{T_1^c(V)} \otimes H\right) \circ l_H^{-1} = i_0^{T_1^c(V \otimes H)} \qquad \text{and} \qquad f^c \circ \left(i_1^{T_1^c(V)} \otimes H\right) = i_1^{T_1^c(V \otimes H)}.$$

By construction of f^c we get

$$\begin{split} p_0^{T_H^c(V \otimes H)} \circ f^c \circ \left(i_0^{T_1^c(V)} \otimes H \right) \circ l_H^{-1} &= l_H \circ \left(p_0^{T_1^c(V)} i_0^{T_1^c(V)} \otimes H \right) \circ l_H^{-1} = \operatorname{Id}_H = p_0^{T_H^c(V \otimes H)} \circ i_0^{T_H^c(V \otimes H)} \\ p_0^{T_H^c(V \otimes H)} \circ f^c \circ \left(i_1^{T_1^c(V)} \otimes H \right) &= l_H \circ \left(p_0^{T_1^c(V)} i_1^{T_1^c(V)} \otimes H \right) = 0 = p_0^{T_H^c(V \otimes H)} \circ i_1^{T_H^c(V \otimes H)}, \\ p_1^{T_H^c(V \otimes H)} \circ f^c \circ \left(i_0^{T_1^c(V)} \otimes H \right) \circ l_H^{-1} &= \left(p_1^{T_1^c(V)} i_0^{T_1^c(V)} \otimes H \right) \circ l_H^{-1} = 0 = p_1^{T_H^c(V \otimes H)} \circ i_0^{T_H^c(V \otimes H)}, \\ p_1^{T_H^c(V \otimes H)} \circ f^c \circ \left(i_1^{T_1^c(V)} \otimes H \right) &= \left(p_1^{T_1^c(V)} i_1^{T_1^c(V)} \otimes H \right) = \operatorname{Id}_V = p_1^{T_H^c(V \otimes H)} \circ i_1^{T_H^c(V \otimes H)}, \end{split}$$

so that, by Corollary 2.18, we have proved (32). Thus $F_{H,V\otimes H} = f^c \circ (F_{1,V} \rtimes H) \circ f$. Now, since f is an epimorphism and f^c is a monomorphism we get

$$H\left[V\otimes H\right] = \operatorname{Im}\left(F_{H,V\otimes H}\right) = \operatorname{Im}\left(f^{c}\circ\left(F_{\mathbf{1},V}\rtimes H\right)\circ f\right) = \operatorname{Im}\left(F_{\mathbf{1},V}\rtimes H\right) = \operatorname{Im}\left(F_{\mathbf{1},V}\right)\rtimes H = \mathbf{1}\left[V\right]\rtimes H.$$

7.5. Let $(H, m_H, u_H, \Delta_H, \varepsilon_H, S_H)$ be a braided graded Hopf algebra with invertible antipode S_H in $((\mathcal{M}, \otimes, \mathbf{1}), c)$.

We have an equivalence of braided monoidal categories

$$F: ({}_{H}^{H}\mathcal{M}_{H}^{H}, \otimes_{H}, H) \to ({}_{H}^{H}\mathcal{YD}(\mathcal{M}), \otimes, \mathbf{1}),$$

given by

$$F\left(M\right) = M^{co(H)} = \operatorname{Eq}\left[\rho_{M}^{r}, \left(M \otimes u_{H}\right) \circ r_{M}^{-1}\right]$$

for every $(M, \mu_M^r, \mu_M^l, \rho_M^r, \rho_M^l) \in {}_H^H \mathcal{M}_H^H$ (see [Bes, Theorem 3.9.6]). The inverse of F is the functor F^{-1} defined by

$$F^{-1}\left(V\right) = \left(V \otimes H, \mu_{V \otimes H}^{l}, \mu_{V \otimes H}^{r}, \rho_{V \otimes H}^{l}, \rho_{V \otimes H}^{r}\right)$$

as in 7.2 for every $V \in {}^{H}_{H} \mathcal{YD}(\mathcal{M})$.

THEOREM 7.6. Let $(H, m_H, u_H, \Delta_H, \varepsilon_H, S_H)$ be a braided graded Hopf algebra with invertible antipode S_H in a cocomplete and complete abelian coabelian braided monoidal category $((\mathcal{M}, \otimes, \mathbf{1}), c)$ satisfying AB5. Assume that the tensor product commutes with direct sums.

Let $(M, \mu_M^r, \mu_M^l, \rho_M^r, \rho_M^l)$ be in ${}^H_H \mathcal{M}^H_H$ and let $\mathbf{1} \left[M^{co(H)} \right]$ be the braided bialgebra of type one associated to $\mathbf{1}$ and $M^{co(H)}$ in ${}^H_H \mathcal{YD} (\mathcal{M})$. Then

$$\mathbf{1}\left[M^{co(H)}\right] \rtimes H = H[M]$$

is the braided bialgebra of type one associated to H and M in \mathcal{M} .

Proof. By Theorem 7.4 **1** $[M^{coH}] \rtimes H = H[M^{coH} \otimes H] \simeq H[M]$ is the braided bialgebra of type one associated to H and M in \mathcal{M} .

References

[AS] N. Andruskiewitsch and H-J. Schneider, *Pointed Hopf algebras*. New directions in Hopf algebras, 1–68, Math. Sci. Res. Inst. Publ., **43**, Cambridge Univ. Press, Cambridge, 2002.

[AMS1] A. Ardizzoni, C. Menini and D. Ştefan, Cotensor Coalgebras in Monoidal Categories, Comm. Algebra., to appear. (arXiv:math.QA/0507334)

[AMS2] A. Ardizzoni, C. Menini and D. Ştefan, Hochschild Cohomology And 'Smoothness' In Monoidal Categories, J. Pure Appl. Algebra, 208 (2007), 297-330.

[Bes] Yu. N. Bespalov, Crossed modules and quantum groups in braided categories, Appl. Categ. Structures 5 (1997), 155–204.

[BD] Y. Bespalov, B. Drabant, Hopf (bi-)modules and crossed modules in braided monoidal categories, J. Pure Appl. Algebra 123 (1998), no. 1-3, 105–129.

[Ka] Kassel, Quantum Groups, Graduate Text in Mathematics 155, Springer, 1995.

[Mj1] S. Majid, Foundations of quantum group theory, Cambridge University Press, 1995.

[NT] C. Năstăsescu, B. Torrecillas, Graded coalgebras, Tsukuba J. Math. 17 (1993), 461-479.

[Ni] Nichols, W. D. Bialgebras of type one. Comm. Algebra 6 (1978), no. 15, 1521–1552.

[Po] N. Popescu, Abelian Categories with Application to Rings and Modules, Academic Press, London & New York, (1973).

[Ro] M. Rosso, Quantum groups and quantum shuffles, Invent. Math. 133 (1998), no. 2, 399-416.

[Sch] P. Schauenburg, A characterization of the Borel-like subalgebras of quantum enveloping algebras Comm. Algebra 24 (1996), no. 9, 2811–2823.

University of Ferrara, Department of Mathematics, Via Machiavelli 35, Ferrara, I-44100, Italy $E\text{-}mail\ address$: rdzlsn@unife.it

 URL : http://www.unife.it/utenti/alessandro.ardizzoni

University of Ferrara, Department of Mathematics, Via Machiavelli 35, Ferrara, I-44100, Italy

E-mail address: men@unife.it

 $\mathit{URL} : \mathtt{http://www.unife.it/utenti/claudia.menini}$