

ALGEBRAIC CYCLES AND COMPLETIONS OF EQUIVARIANT K -THEORY

DAN EDIDIN AND WILLIAM GRAHAM

ABSTRACT. Let G be a complex, linear algebraic group acting on an algebraic space X . The purpose of this paper is to prove a Riemann-Roch theorem (Theorem 5.5) which gives a description of the completion of the equivariant Grothendieck group $G_0(G, X) \otimes \mathbb{C}$ at any maximal ideal of the representation ring $R(G) \otimes \mathbb{C}$ in terms of equivariant cycles. The main new technique for proving this theorem is our non-abelian completion theorem (Theorem 4.3) for equivariant K -theory. Theorem 4.3 generalizes the classical localization theorems for diagonalizable group actions to arbitrary groups.

CONTENTS

1. Introduction	2
1.1. Background	5
2. Induced representations in algebraic K -theory	6
2.1. Merkurjev's theorem and Weyl group actions	7
2.2. Twisted induction	8
2.3. Induction for Levi factors of parabolic subgroups	10
3. Induction, restriction and completion	12
3.1. Some facts about completions and invariants	12
3.2. Induction and restriction	14
4. The non-abelian completion theorem	16
4.1. The completion theorem for a central element	16
4.2. Statement of the non-abelian completion theorem	18
4.3. Proof of Theorem 4.3(a),(b)	19
4.4. The case of finite stabilizer	22
4.5. More on the finite stabilizer case	23
5. The twisted equivariant Riemann-Roch theorem	26
5.1. The equivariant Riemann-Roch theorem	26
5.2. Twisting by a central element	27
5.3. The twisted equivariant Riemann-Roch theorem	28
6. Equivariant envelopes and proofs of uniqueness	30

The second author was supported by N.S.F.

6.1. Equivariant envelopes	30
6.2. Proofs of Theorem 4.3(c), Corollary 4.5(b) and Theorem 5.5(d)	33
References	34

1. INTRODUCTION

The now-classic Riemann-Roch theorem of Baum, Fulton and MacPherson [BFM, Ful] states that for any separated algebraic space X there is a natural isomorphism between the Grothendieck group $G_0(X)$ of coherent sheaves on X and the Chow group CH^*X of cycles on X .¹ When a linear algebraic group G acts on X , the equivariant Riemann-Roch problem is to relate the equivariant Grothendieck group $G_0(G, X)$ of G -equivariant coherent sheaves and the G -equivariant Chow group CH_G^*X . In contrast to the non-equivariant case, elements of $G_0(G, X)$ cannot in general have unique representations by equivariant algebraic cycles. For example, if G is a finite group and X is a point then $G_0(G, X) = \mathbb{C}^r$, where r is the number of conjugacy classes in G , while $CH_G^*(X) = \mathbb{C}$. To obtain precise results we must use the fact that the equivariant Grothendieck group is a module for the representation ring $R(G)$.

The main theorem of [EG2] is an equivariant Riemann-Roch isomorphism between the completion of $G_0(G, X)$ at the ideal in $R(G)$ of virtual representations of rank 0 (the augmentation ideal) and the infinite product of equivariant Chow groups $\prod_{n=0}^{\infty} CH_G^n(X)$. Hence elements of the augmentation completion of $G_0(G, X)$ may be represented by equivariant cycles. An obvious problem is to determine whether other completions of $G_0(G, X)$ admit geometric descriptions.

One of the goals of this paper is to solve this problem for completions of $G_0(G, X)$ at maximal ideals in $R(G)$ when G is a complex algebraic group. Because we are using complex coefficients, maximal ideals in $R(G)$ correspond bijectively to semisimple conjugacy classes in G . The correspondence takes a semisimple conjugacy class Ψ to \mathfrak{m}_{Ψ} , the maximal ideal of virtual representations whose characters vanish at Ψ ; the augmentation ideal corresponds to the conjugacy class of the identity element of G . In this paper we generalize the results of [EG2] by proving that if h is an element of Ψ with centralizer Z , then elements of the \mathfrak{m}_{Ψ} -adic completion of $G_0(G, X)$ can be represented by Z -equivariant

¹Throughout this paper all K -groups and Chow groups are taken with complex coefficients.

cycles on the fixed subspace X^h . Precisely, we prove (Theorem 5.5) that for any G -space X , there is a natural Riemann-Roch isomorphism τ_X^Ψ from the \mathfrak{m}_Ψ -adic completion of $G_0(G, X)$ to the infinite product $\prod_{n=0}^{\infty} CH_Z^n(X^h)$. When X is a smooth scheme, there is an explicit formula for τ_X^Ψ in terms of Chern characters and Todd classes.

The equivariant Riemann-Roch theorem proved here follows from our non-abelian completion theorem (Theorem 4.3), a result of independent interest. The progenitor for Theorem 4.3 is the classical localization theorem originally proved by Segal [Seg] for Lie group actions, and extended to algebraic K -theory by Nielsen [Nie] and Thomason [Tho2]. It states that if G is a diagonalizable group (e.g., a torus) and h is an element of G , then the pushforward $G(G, X^h) \xrightarrow{i_*} G(G, X)$ becomes an isomorphism after localizing at the maximal ideal \mathfrak{m}_h of $R(G)$. (Here $G(G, X)$ denotes the infinite direct sum of equivariant K -groups $\bigoplus_{n=0}^{\infty} G_n(G, X)$.) Moreover, if X is smooth then there is an explicit localization formula

$$(1) \quad \alpha = i_* \left(\frac{i^* \alpha}{\lambda_{-1}(N_i^*)} \right)$$

where $\alpha \in G(G, X)_{\mathfrak{m}_h}$ and N_i^* is the conormal bundle of the regular embedding $i: X^h \rightarrow X$. Many applications of the localization theorem, such as a simple proof of the Weyl character formula, come from this explicit localization formula. In [EG3], we extended this result to the case where G is an arbitrary algebraic group, but with the assumption that G acts with finite stabilizer, and used our version of the explicit localization formula to give a Riemann-Roch formula for orbifolds. Related results were also obtained in that case by Toen [Toe] and Vezzosi-Vistoli [VV].

In this paper, we remove the restriction on the group action, but to do so, we must complete rather than localize. (In the finite stabilizer case these two operations coincide because the equivariant K -theory is supported at a finite number of maximal ideals in $R(G)$.) The nonabelian completion theorem states that there is a pushforward isomorphism

$$i_! : \widehat{G(Z, X^h)}_h \rightarrow \widehat{G(G, X)},$$

where $\widehat{G(Z, X^h)}_h$ is the completion of $G(Z, X^h)$ at the maximal ideal \mathfrak{m}_h of $R(Z)$, and $\widehat{G(G, X)}$ is the completion of $G(G, X)$ at \mathfrak{m}_Ψ . Moreover, when X is smooth a formula analogous to (1) above holds. This formula allows us to obtain our explicit formula for the Riemann-Roch isomorphism τ_X^Ψ when X is smooth.

The proof of the nonabelian completion theorem rests on the construction of a “twisted induction” functor in equivariant K -theory. If G is a connected reductive group and $Z \subset G$ is a connected subgroup of maximal rank such that both groups have simply connected commutator subgroups then we can use a theorem of Merkurjev to define an induction map $\text{ind}: G(Z, X) \rightarrow G(G, X)$ satisfying several natural properties, including what we call a “twisted reciprocity” formula relating ind to the usual restriction functor $\text{res}: G(G, X) \rightarrow G(Z, X)$. When Z is the centralizer of a semi-simple element $h \in G$, generalities about completions imply that the \mathfrak{m}_h -completion $\widehat{G(Z, X)}_h$ of $G(Z, X)$ is naturally identified as a summand in the \mathfrak{m}_Ψ -adic completion $\widehat{G(G, X)}$. By composing the inclusion as a summand with the map ind , we obtain a map

$$\text{ind}_h: \widehat{G(Z, X)}_h \rightarrow \widehat{G(G, X)}.$$

Using some basic facts about invariants and completion, we prove that ind_h and the natural map

$$\text{res}_h: \widehat{G(G, X)} \rightarrow \widehat{G(Z, X)}_h$$

are inverse isomorphisms (Proposition 3.5). (This proposition illustrates the necessity of working with completions rather than localizations, as the natural restriction map of localizations $G(G, X)_{\mathfrak{m}_\Psi} \rightarrow G(Z, X)_{\mathfrak{m}_h}$ is not in general an isomorphism (Example 3.7).) In the case that Z and G are both connected and reductive and G has simply connected commutator subgroup, the map i_i can be defined as a composition

$$\widehat{G(Z, X^h)}_h \xrightarrow{i_*} \widehat{G(Z, X)}_h \xrightarrow{\text{ind}_h} \widehat{G(G, X)}.$$

Because h is in the center of Z , we can argue as in the proof of the localization theorem for tori to show that i_* is an isomorphism; hence i_i is an isomorphism as well. For general G and Z a change of groups argument similar to that of [EG3] can be used to define i_i .

There are a number of natural questions arising from this work. If the element h is defined over a subfield $L \subset \mathbb{C}$ then our techniques imply that Theorem 4.3 holds for completions of equivariant K -theory tensored with L . An interesting problem for further study is to prove a version of Theorem 4.3 for completions of integral or rational equivariant K -theory at maximal (or prime) ideals of the representation ring $R(G)$. It would also be interesting to extend these results to algebraic groups defined over arbitrary fields. Similarly, if Ψ is the conjugacy class of an element $h \in G$ which is defined over a subfield $L \subset \mathbb{C}$ then

our techniques show that there is an isomorphism of completions of equivariant K -theory tensored with L and equivariant Chow groups with coefficients in L . Given an arbitrary maximal (or prime) ideal \mathfrak{m} in $R(G) \otimes \mathbb{Q}$ a very interesting open problem is to represent elements of the \mathfrak{m} -adic completion of $G_0(G, X)$ as formal equivariant algebraic cycles on an appropriate subspace of X . Another natural question is whether a version of Theorem 5.5 holds for higher K -theory. Specifically one can ask if there is a natural isomorphism between the \mathfrak{m}_Ψ -adic completion of $G_n(G, X)$ and the infinite product of higher equivariant Chow groups $\prod_{i=0}^{\infty} CH_G^i(X, n)$. Such a result would follow from the methods of this paper and the higher K -theory version of the Riemann-Roch isomorphism of [EG2]. Unfortunately we have not been able to construct such an isomorphism due to the difficulties in comparing different completions of higher equivariant K -theory (cf [EG2, Remark 2.2]).

Acknowledgement: The authors are very grateful to the referees for their careful reading as well as for many comments which helped to clarify the exposition.

1.1. Background. We work entirely over the ground field \mathbb{C} of complex numbers. All algebraic spaces are assumed to be separated and of finite type over \mathbb{C} . For a reference on the theory of algebraic spaces see the book [Knu]. As in [EG1] we will refer to an integral subspace of an algebraic space X as a *subvariety* of X . Note that if h is an automorphism of a separated algebraic space X , then the fixed point subspace X^h is closed in X , since it is the inverse image of the diagonal under the morphism $X \rightarrow X \times X$ given by the graph of h .

All algebraic groups are assumed to be linear. A basic reference for the theory of algebraic groups is [Bor]. If G is an algebraic group and $h \in G$, then $\mathcal{Z}_G(h)$ denotes the centralizer of h in G , and $C_G(h)$ the conjugacy class of h in G .

Throughout the paper we take the complex numbers as our coefficients for K -groups and Chow groups, e.g., we write $G_0(G, X)$ for $G_0(G, X) \otimes \mathbb{C}$.

1.1.1. Representation rings. We recall here some facts about representation rings proved in [EG3]. If G is an algebraic group then $R(G)$ denotes the representation ring of G tensored with \mathbb{C} . By [EG3, Proposition 2.5] there is a bijective correspondence between semisimple conjugacy classes in G and maximal ideals in $R(G)$. If Ψ is the conjugacy class of a semisimple element, then the corresponding maximal ideal \mathfrak{m}_Ψ consists of virtual representations whose characters vanish at Ψ .

If G is an algebraic group and $H \subset G$ is a closed subgroup then the restriction map $R(G) \rightarrow R(H)$ is a finite morphism ([EG3, Proposition 2.3]). As a result, if Ψ is a semisimple conjugacy class in G , then $\Psi \cap H$ decomposes into a finite number of conjugacy classes Ψ_1, \dots, Ψ_l , and a maximal ideal $\mathfrak{m}_{\Psi'} \subset R(H)$ lies over $\mathfrak{m}_{\Psi} \subset R(G)$ if and only if $\Psi' = \Psi_k$ for some k ([EG3, Proposition 2.6]).

1.1.2. *Equivariant K -theory.* As in [EG3], $G(G, X)$ denotes the infinite direct sum $\bigoplus_{i=0}^{\infty} G_i(G, X)$, where $G_i(G, X)$ is the i -th Quillen K -group of the category of G -equivariant coherent sheaves on X , tensored with \mathbb{C} . Thus, $G_0(G, X)$ is the Grothendieck group of G -equivariant coherent sheaves, tensored with \mathbb{C} . Likewise, $K_0(G, X)$ denotes the Grothendieck ring of G -equivariant vector bundles, also tensored with \mathbb{C} . When X is a smooth scheme then $K_0(G, X)$ and $G_0(G, X)$ may be identified thanks to Thomason's equivariant resolution theorem [Tho1]. If $X = \text{Spec } \mathbb{C}$ then $K_0(G, X) = G_0(G, X) = R(G)$.

If N is a G -equivariant vector bundle of rank n on X then $\lambda_{-1}(N)$ is the formal sum $\sum_{i=0}^n (-1)^i [\Lambda^i N]$.

If Z is a closed subgroup of G and X is a Z -space, then there is a ‘‘Morita equivalence’’ identification of $G(Z, X)$ with $G(G, G \times_Z X)$ (see Section 3.1 of [EG3] for more details). In particular, an element of $R(Z)$ such as $\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*)$ can be viewed as operating on $G(G, G \times_Z X)$. If X is a G -space, then $G \times_Z X$ is isomorphic to $G/Z \times X$, and the relative tangent bundle of the morphism $G \times_Z X \rightarrow X$ is identified under this equivalence with the pullback of the element $\mathfrak{g}/\mathfrak{z}$ of $R(Z)$ to $G(Z, X)$. In this case we will often simply identify the relative tangent bundle of the morphism as $\mathfrak{g}/\mathfrak{z}$ without further comment.

For a sketch of some basic properties of equivariant K -theory, most of which are due to Thomason, we refer the reader to Section 3.1 of our paper [EG3]. Facts from that section will be used in this paper without further reference.

2. INDUCED REPRESENTATIONS IN ALGEBRAIC K -THEORY

Let G be a connected reductive group and let $H \subset G$ be a connected reductive subgroup of the same rank. Assume that G and H have simply connected commutator subgroups. The goal of this section is to define a natural twisted induction map $G(H, X) \rightarrow G(G, X)$ satisfying an appropriate reciprocity formula (2.4). A theorem of Merkurjev is crucial for proving properties of the twisted induction map.

2.1. Merkurjev's theorem and Weyl group actions. Suppose that H is a subgroup of G and X is a G -space. There is a natural map

$$(2) \quad \theta : R(H) \otimes_{R(G)} G(G, X) \rightarrow G(H, X),$$

given by $\theta([V] \otimes \alpha) = [V] \text{res } \alpha$. This map can be realized another way (cf. [EG2, Prop. 3.2]). We can identify $G \times_H X$ with $G/H \times X$ via the map taking $(g, x)H$ to (gH, gx) . This identification is G -equivariant, where G acts on $G \times_H X$ by left multiplication on G , and on $G/H \times X$ by the direct product of the actions on G/H and X . Consequently the category \mathbf{coh}_X^H of H -equivariant coherent sheaves on X is equivalent to the category $\mathbf{coh}_{G/H \times X}^G$ of G -equivariant coherent sheaves on $G/H \times X$. In terms of this equivalence the map θ is given as follows. Let π_1 and π_2 denote the projections of $G/H \times X$ to G/H and X , respectively, and given a representation V of H , let \mathcal{V} denote the vector bundle $G \times_H V$ on G/H . Then θ is the map $R(H) \otimes_{R(G)} G(G, X) \rightarrow G(H, X)$ induced by the family of functors $f_V : \mathbf{coh}_G X \rightarrow \mathbf{coh}_H X$, $\mathcal{F} \mapsto \pi_1^* \mathcal{V} \otimes \pi_2^* \mathcal{F}$, defined for each H -module V . We may write θ_H^G for θ if we wish to make explicit the groups involved.

If G is a connected reductive group with simply connected commutator subgroup and T is a maximal torus of G , then Merkurjev proves ([Mer, Proposition 31]) that the map $\theta_T^G : R(T) \otimes_{R(G)} G(G, X) \rightarrow G(T, X)$ is an isomorphism. (Note that Merkurjev's result holds for K -theory with integral coefficients.) The following extension of Merkurjev's result will be useful for proving properties of the twisted induction map.

Proposition 2.1. *Suppose that $H \subset G$ are connected reductive groups of the same rank both with simply connected commutator subgroups. Then the natural map $\theta_H^G : R(H) \otimes_{R(G)} G(G, X) \rightarrow G(H, X)$ taking $[V] \otimes \alpha$ to $\text{res}(\alpha)[V]$ is an isomorphism.*

Proof. Let T be a maximal torus of H (hence also of G). By Merkurjev's theorem,

$$G(T, X) = R(T) \otimes_{R(G)} G(G, X) = R(T) \otimes_{R(H)} G(H, X).$$

Hence the map

$$R(T) \otimes_{R(H)} (R(H) \otimes_{R(G)} G(G, X)) \rightarrow R(T) \otimes_{R(H)} G(H, X)$$

is an isomorphism, since both sides are identified with $G(T, X)$. Since H is connected and reductive, the restriction map $R(H) \rightarrow R(T)$ is a split injection. Thus tensoring with $R(T)$ is a fully faithful functor. Therefore $G(G, X) \otimes_{R(G)} R(H) = G(H, X)$ as claimed. \square

Let G be as above; and let T be a maximal torus of G and W the corresponding Weyl group. If X is a G -space, then W acts on $G/T \times X$ by the rule $w \cdot (gT, x) = (gw^{-1}T, x)$. This action commutes with the action of G and hence induces an action of W on $G(G, G/T \times X)$, which we can identify with $G(T, X)$. This action of W is natural with respect to flat pullbacks and proper pushforwards arising from morphisms of G -spaces. The isomorphism

$$\theta_T^G : R(T) \otimes_{R(G)} G(G, X) \rightarrow G(T, X)$$

is W -equivariant, where W acts on the source by its action on $R(T)$. By a theorem of Serre [Ser, Theorem 4] we may identify $R(G) = R(T)^W$, so $G(T, X)^W = G(G, X)$. (Note that Serre's result holds even if the commutator subgroup is not simply connected.)

2.2. Twisted induction.

Definition 2.2. Let $H \subseteq G$ be connected reductive groups of equal rank with simply connected commutator subgroups, and let T be a maximal torus of H (hence also of G). Let $W_1 \subseteq W$ denote the Weyl groups of T in H and G , respectively. Let X be a G -space. We define

$$\text{ind}_H^G : G(H, X) \rightarrow G(G, X)$$

as follows. Identify $G(G, X)$ (resp. $G(H, X)$) as the W -invariants (resp. W_1 -invariants) in $G(T, X)$, and for any $\alpha \in G(H, X)$, set

$$\text{ind}_H^G(\alpha) = \sum_{w \in W/W_1} w\alpha.$$

(Note that we use the same notation for elements of W/W_1 and lifts to W ; sometimes we also use the same notation for lifts of elements of W to G .)

Definition 2.3. If G is a reductive group, define a linear map $R(G) \rightarrow \mathbb{C}$, $\alpha \mapsto \alpha^G$, by setting $[V]^G = \dim V^G$ for any G -module V and extending by linearity. Likewise, define a symmetric bilinear map $\text{Hom}_G(-, -) : R(G) \otimes R(G) \rightarrow \mathbb{C}$ by setting

$$\text{Hom}_G([V], [W]) = [V^* \otimes W]^G$$

for G -modules V, W .

The twisted induction map has properties given in the following proposition. Part (d) is analogous to the Frobenius reciprocity formula for induced representations, and justifies our use of the term twisted induction map.

Proposition 2.4. *Let $H \subseteq G$ be connected reductive groups of equal rank. Assume that both H and G have simply connected commutator subgroups. The map*

$$\mathrm{ind}_H^G : G(H, X) \rightarrow G(G, X)$$

has the following properties.

(a) ind_H^G is natural with respect to flat pullbacks and proper pushforwards arising from morphisms of G -spaces.

(b) If $K \subset H \subset G$ are connected reductive groups of equal rank with simply connected commutator subgroups, then

$$\mathrm{ind}_H^G \circ \mathrm{ind}_K^H = \mathrm{ind}_K^G.$$

(c) (Projection formula) If $\alpha \in R(H)$ and $\beta \in G(G, X)$ then

$$\mathrm{ind}_H^G(\alpha \mathrm{res}(\beta)) = \mathrm{ind}_H^G(\alpha)\beta.$$

(d) (Reciprocity formula) When $X = \mathrm{Spec} \mathbb{C}$, the map $\mathrm{ind}_H^G : R(H) \rightarrow R(G)$ satisfies

$$(3) \quad \mathrm{Hom}_G(\mathrm{ind}_H^G \alpha, \beta) = \mathrm{Hom}_H(\lambda_{-1}(\mathfrak{g}^*/\mathfrak{h}^*)\alpha, \mathrm{res} \beta),$$

where \mathfrak{g} and \mathfrak{h} are the Lie algebras of G and H respectively.

Moreover, the map ind_H^G is uniquely determined by properties (c) and (d).

Proof. Naturality of the map ind_H^G follows from naturality of the action of the Weyl group described in Section 2.1. Assertion (b) is a straightforward calculation. For the projection formula, if $\alpha \in R(H)$ and $\beta \in G(G, X)$, then $\alpha \mathrm{res}(\beta)$ is identified with the element $\alpha\beta \in G(T, X)$. Since β is W -invariant, $w(\alpha\beta) = w(\alpha)\beta$; the projection formula follows. We now turn to the reciprocity formula. Using the projection formula we reduce to the case where β is trivial; then we want to show

$$(\mathrm{ind}_H^G(\alpha))^G = (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{h}^*)\alpha)^H.$$

First assume that $H = T$ so by Serre's theorem we may identify $R(G)$ with $R(T)^W$. The groups H and G are complexifications of their corresponding maximal compact Lie subgroups [OV, Theorem 8, p. 244], and a compact Lie group has the same representation ring as its complexification [BtD, Ch. III, Prop. 8.6]. The Weyl integration formula [BtD, Ch. IV, Theorem 1.11] for compact Lie groups implies that if $\nu \in R(G)$, then

$$(4) \quad \nu^G = \frac{1}{|W|}((\lambda_{-1}(\mathfrak{g}^*/\mathfrak{t}^*)\nu)^T).$$

If $\nu = \text{ind}_T^G(\alpha)$, then, since $\lambda_{-1}(\mathfrak{g}^*/\mathfrak{t}^*)$ is W -invariant, the right hand side of (4) becomes

$$\frac{1}{|W|} \sum_{w \in W} (w(\lambda_{-1}(\mathfrak{g}^*/\mathfrak{t}^*)\alpha))^T.$$

But if $r \in R(T)$ then $r^T = (wr)^T$ for all $w \in W$; the reciprocity formula follows for $H = T$. For general H , we suppose $T \subset H \subset G$. We may assume $\alpha = \text{ind}_T^H(\beta)$. Then, applying (b) and what we have just proved,

$$(\text{ind}_H^G(\alpha))^G = (\text{ind}_T^G(\beta))^G = (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{t}^*)\alpha)^T.$$

On the other hand, applying the projection formula and what we have just proved with H in place of G , we obtain

$$(\lambda_{-1}(\mathfrak{g}^*/\mathfrak{h}^*)\alpha)^H = (\text{ind}_T^H(\lambda_{-1}(\mathfrak{g}^*/\mathfrak{h}^*)\beta))^H = (\lambda_{-1}(\mathfrak{h}^*/\mathfrak{t}^*)\lambda_{-1}(\mathfrak{g}^*/\mathfrak{h}^*)\beta)^T.$$

Since $\lambda_{-1}(\mathfrak{g}^*/\mathfrak{t}^*) = \lambda_{-1}(\mathfrak{h}^*/\mathfrak{t}^*)\lambda_{-1}(\mathfrak{g}^*/\mathfrak{h}^*)$, the reciprocity formula follows. Finally, the projection formula and Proposition 2.1 imply that the map ind_H^G is uniquely determined by its value when X is a point, in which case it is determined by the reciprocity formula. \square

2.3. Induction for Levi factors of parabolic subgroups. If H is a Levi factor of a parabolic subgroup P of G , and X is a G -space, then

$$G(H, X) \simeq G(G, G/H \times X) \simeq G(G, G/P \times X).$$

Here the first isomorphism was explained in Section 2.1, and the second is because the projection $G/H \times X \rightarrow G/P \times X$ has fibers isomorphic to affine space.

We include a proof of the following lemma because of the lack of a reference.

Lemma 2.5. *Let G be a connected reductive group. The commutator subgroup G' of G is simply connected if and only if $\pi_1(G)$ is torsion-free.*

Proof. The group G is the product of the semisimple group G' and a torus S , with $G' \cap S$ finite (see [Hum, Section 0.9]). Therefore G/G' is isomorphic to the torus $S/(G' \cap S)$. Since $\pi_2(G/G') = 0$, the long exact homotopy sequence of the fibration $G \rightarrow G/G'$ yields a short exact sequence $0 \rightarrow \pi_1(G') \rightarrow \pi_1(G) \rightarrow \pi_1(G/G') \rightarrow 0$. Since G/G' is a torus, $\pi_1(G/G')$ is free abelian. Since G' is semisimple, $\pi_1(G')$ is finite [Hel, Chapter VII, Theorem 6.1]. The lemma follows. \square

Proposition 2.6. *Let H be a Levi factor of a parabolic subgroup of G . If G has simply connected commutator subgroup, then so does H .*

Proof. The quotient G/H has the homotopy type of G/P , so G/H is simply connected and by the Hurewicz theorem, $\pi_2(G/H)$ is isomorphic to $H_2(G/H) \simeq H_2(G/P)$, which is torsion-free. Since $\pi_2(G) = 0$, the long exact homotopy sequence of the fibration $G \rightarrow G/H$ implies that $\pi_1(H)$ is torsion-free. \square

In this situation, the following proposition gives another construction of the twisted induction map.

Proposition 2.7. *Let G and H be as in Proposition 2.6. Identify $G(H, X)$ with $G(G, G/P \times X)$, and let $q : G/P \times X \rightarrow X$ denote the projection. If $\alpha \in G(H, X)$, then*

$$(5) \quad \text{ind}_H^G(\alpha) = q_*(\lambda_{-1}(\mathfrak{g}^*/\mathfrak{p}^*)\alpha).$$

Proof. The map defined by the right hand side of (5) satisfies the projection formula. Hence by Propositions 2.1 and 2.4 it suffices to show (5) hold for $\alpha \in R(H)$. When computing the right hand side of (5) we view $\lambda_{-1}(\mathfrak{g}^*/\mathfrak{p}^*)\alpha$ as inducing a virtual vector bundle on G/P , and q_* as taking the G -equivariant Euler characteristic of that bundle. Here, if V is a representation of H , then we may view it as a P -module by making the unipotent radical act trivially. The induced vector bundle on G/P is $G \times^P V$. The vector bundle on G/P induced by $\mathfrak{g}^*/\mathfrak{p}^*$ is the cotangent bundle $T^* = T_{G/P}^*$, so what we need to compute is $q_*(\lambda_{-1}(T^*)\alpha)$. The latter computation may be done using the same technique as in the proof Proposition 3.10 of [EG3], as we now explain.

The restriction map $G_0(G, G/P) \rightarrow G_0(T, G/P)$ is injective, so we may restrict to T -equivariant K -theory, and make the calculation in $G_0(T, G/P)$. In particular if $a \in T$ is any element such that $(G/P)^a = (G/P)^T$, then the localization theorem in T -equivariant K -theory [EG3, Theorem 3.3] implies that

$$(6) \quad \lambda_{-1}(T^*)\alpha = \sum_{x \in (G/P)^T} i_{x*} \frac{i_x^*(\lambda_{-1}(T^*)\alpha)}{\lambda_{-1}(T_x^*)} = \sum_{x \in (G/P)^T} i_{x*} i_x^* \alpha$$

in $G_0(T, G/P)_{\mathfrak{m}_a}$, where i_x is the inclusion of x in G/P . Since G/P has a cellular decomposition, [CG, Lemma 5.5.1] implies that $G_0(T, G/P)$ is a free $R(T)$ -module of rank equal to the number of T -fixed points. Since $R(T)$ is an integral domain, it follows that the equality

$$(7) \quad \lambda_{-1}(T^*)\alpha = \sum_{x \in (G/P)^T} i_{x*} i_x^* \alpha$$

also holds in $G_0(T, G/P)$. The T -fixed points in G/P are the points wP , for $w \in W/W_1$. If x corresponds to the coset wP then, identifying

$G(T, x)$ with $R(T)$, the map $q_* \circ i_x$ is the identity, and for $\alpha \in R(H) = R(T)^{W_1}$, we have $i_x^*(\alpha) = w\alpha$. Applying q_* to both sides of (7) yields

$$q_*(\lambda_{-1}(T^*)\alpha) = \sum_{w \in W/W_1} w\alpha = \text{ind}_H^G \alpha,$$

as desired. \square

3. INDUCTION, RESTRICTION AND COMPLETION

Let G be a connected reductive group with simply connected commutator subgroup and let $\Psi \subset G$ be a semisimple conjugacy class. Let h be an element of Ψ , and let $Z = \mathcal{Z}_G(h)$ be the centralizer of h . Assume that Z is also connected and reductive and has simply connected commutator subgroup.² In this section we prove that, after appropriate completions, the induction functor ind_Z^G is in fact the inverse of the restriction functor.

3.1. Some facts about completions and invariants. In this section we prove some basic commutative algebra results about invariants and completions which will be necessary for proving properties of the induction map in equivariant K -theory.

3.1.1. Completions and finite morphisms. Let $B \rightarrow A$ be a finite morphism of Noetherian rings. Let \mathfrak{m} be a maximal ideal of B , let $\mathfrak{a} = \mathfrak{m}A$, and let $\mathfrak{m}_1, \dots, \mathfrak{m}_r$ be the maximal ideals of A lying over \mathfrak{m} . Let M be an A -module and let \widehat{M} be the \mathfrak{a} -adic completion of M . Likewise, let \widehat{M}_i denote the \mathfrak{m}_i -adic completion of M . Since $\mathfrak{m}M \subset \mathfrak{m}_i M$ there is an induced map of completions $v_i: \widehat{M} \rightarrow \widehat{M}_i$. The following proposition will be needed. It is a generalization of [Bou, Chapter III, Section 2, no. 13, Proposition 18].

Lemma 3.1. *The map*

$$(v_i): \widehat{M} \rightarrow \prod_{i=1}^r \widehat{M}_i$$

is an isomorphism of \widehat{A} -modules.

Proof. Let N denote the product $\prod_{i=1}^r M$ with the topology given by the product of the \mathfrak{m}_i -adic topologies, $i = 1, \dots, r$. The completion of N in this topology is the product $\prod_{i=1}^r \widehat{M}_i$.

²If G has simply connected commutator subgroup then Z is automatically connected (c.f. [Hum, Theorem 2.11]).

We claim that the \mathfrak{a} -adic topology on M is the same as the topology induced by the inclusion by the diagonal map $M \xrightarrow{\Delta} N$. To see this argue as follows. Let $N_k = \prod_{i=1}^r \mathfrak{m}_i^k M \subset N$. The filtration $N \supset N_1 \supset \dots \supset N_k \supset \dots$ is a fundamental system of neighborhoods of 0 so the completion of N can be viewed as the completion with respect to this filtration. Now $\Delta^{-1}(N_k) = \cap \mathfrak{m}_i^k M = \mathfrak{m}_1^k \dots \mathfrak{m}_r^k M$ so the induced topology on M is the $\mathfrak{m}_1 \dots \mathfrak{m}_r$ -adic topology. But $\sqrt{\mathfrak{a}} = \mathfrak{m}_1 \dots \mathfrak{m}_r$ so this coincides with the \mathfrak{a} -adic topology as A is Noetherian.

If $n = (m_1, \dots, m_r) \in N$ then by the Chinese remainder theorem for modules [Bou, Chapter II, Section 1, no. 2, Proposition 6] there exists $m \in M$ such that $m \equiv m_i \pmod{\mathfrak{m}_i^n M}$ so $n \in \Delta(M) + N_n$ for any n . Hence $\Delta(M)$ is dense in N .

Let $N/\Delta(M)$ have the topology induced from the topology on N . Since the \mathfrak{a} -adic topology on M is induced from the topology on N , the sequence of completions

$$0 \rightarrow \widehat{M} \xrightarrow{\widehat{\Delta}} \widehat{N} \rightarrow \widehat{N/\Delta(M)} \rightarrow 0$$

is exact [Bou, Chapter III, Section 12, no. 12 Lemma 2]. But $\Delta(M)$ is dense, so $\widehat{N/\Delta(M)} = 0$. Identifying the completion \widehat{N} with $\prod_{i=1}^r \widehat{M}_i$ and $\widehat{\Delta}$ with $\prod_{i=1}^r v_i$ yields the proposition. \square

3.1.2. Completions and invariants. Let W be a finite group acting on a ring A . Assume that $|W|$ is invertible in A . Let $B = A^W$ denote the subring of invariants. Let $I \subset B$ be an ideal, and \widehat{B} the I -adic completion of B .

Let M be an A -module with a W -action compatible with the module structure, and let M^W be the B -submodule of W -invariants. Since $I \subset B$ is W -invariant, there is an action of W on the I -adic completion \widehat{M} of M .

Lemma 3.2. *Let \widehat{M}^W be the I -adic completion of M^W . There is a natural isomorphism of \widehat{B} -modules $\widehat{M}^W \rightarrow (\widehat{M})^W$.*

Proof. First observe that since W acts trivially on \widehat{B} , $(\widehat{M})^W$ is in fact a \widehat{B} -module. The hypothesis that $1/|W| \in A$ allows us to define a projection $M \rightarrow M^W$ by the formula $m \mapsto \frac{1}{|W|} \sum_{w \in W} wm$. Thus, for any integer $k > 0$, the exact sequence of A -modules

$$0 \rightarrow I^k M \rightarrow M \rightarrow M/I^k M \rightarrow 0$$

induces an exact sequence of B -modules

$$0 \rightarrow (I^k M)^W \rightarrow M^W \rightarrow (M/I^k M)^W \rightarrow 0.$$

Moreover, $(I^k M)^W = I^k M^W$, so the exact sequence induces isomorphisms

$$(8) \quad M^W / I^k M^W \rightarrow (M / I^k M)^W$$

for all k . By definition,

$$\lim_{\leftarrow} M^W / I^k M^W = \widehat{M^W}.$$

Also, an element $(m_k) \in \widehat{M}$ is W -invariant if and only if each m_k is W -invariant, i.e.,

$$(\widehat{M})^W = \lim_{\leftarrow} (M / I^k M)^W \subset \lim_{\leftarrow} M / I^k M = \widehat{M}.$$

Thus the desired isomorphism $\widehat{M^W} \rightarrow (\widehat{M})^W$ follows from (8) by taking inverse limits. \square

3.1.3. Invariants and direct products. Let A be a ring, let $M_1 \dots, M_r$ be A -modules, and let $M = \prod_{i=1}^r M_i$. Suppose that a finite group W acts compatibly on A and M . Assume that for all $w \in W$, $wM_i = M_j$ for some j , and conversely that for any i and j there exists $w \in W$ such that $wM_i = M_j$. Let $W_i = \{w \in W \mid wM_i = M_i\}$. The transitivity of the action on the components of M implies that the W_i are conjugate subgroups of W . Let $\pi_i: M \rightarrow M_i$ denote the projection. If $m \in M^W$ then the i -th component of m must be in $M_i^{W_i}$, so π_i restricts to a map $\text{res}_i: M^W \rightarrow M_i^{W_i}$.

Let w_1, \dots, w_r be a set of representatives for the cosets of W/W_i such that $w_j M_i = M_j$. We define a map $\text{ind}_i: M_i^{W_i} \rightarrow M^W$ by $\text{ind}_i(m_i) \mapsto (w_1 m_i, \dots, w_r m_i)$.

Lemma 3.3. *With the above assumptions and notation, the maps res_i and ind_i are inverse isomorphisms of the A^W -modules M^W and $M_i^{W_i}$.*

Proof. Let $m \in M^W$. Since W acts transitively on the components of M , $\text{res}_i(m) = 0$ if and only if $m = 0$, so res_i is injective. Since $\text{res}_i \circ \text{ind}_i$ is the identity, res_i and ind_i are inverse isomorphisms. \square

Remark 3.4. Our hypothesis on the W action on M implies that $M = \text{Ind}_{W_i}^W(M_i)$ for any i , where Ind is the standard induction functor from W_i -modules to W -modules. Viewed this way, Lemma 3.3 is simply the statement that $(\text{Ind}_{W_i}^W(M_i))^W = M_i^{W_i}$.

3.2. Induction and restriction. Let G be a complex algebraic group and $\Psi \subset G$ a semisimple conjugacy class. Fix $h \in \Psi$ and set $Z = \mathcal{Z}_G(h)$. Let $\mathfrak{m}_\Psi \subset R(G)$ be the maximal ideal of virtual representations whose character vanishes on Ψ and $\mathfrak{m}_h \subset R(Z)$ the maximal ideal of virtual representations whose character vanishes at the central element

h . By [EG3, Propositions 2.3, 2.6] the restriction map $R(\widehat{G}) \rightarrow R(\widehat{Z})$ is finite and \mathfrak{m}_h lies over \mathfrak{m}_Ψ . If M is an $R(\widehat{G})$ -module, let \widehat{M} denote the \mathfrak{m}_Ψ -adic completion of M . If M is also an $R(\widehat{Z})$ -module, such that the $R(\widehat{G})$ -module structure is obtained from the $R(\widehat{Z})$ -module structure by the map $R(\widehat{G}) \rightarrow R(\widehat{Z})$, then Lemma 3.1 implies that the \mathfrak{m}_h -adic completion \widehat{M}_h of M is canonically identified as a summand in the \mathfrak{m}_Ψ -adic completion \widehat{M} . The natural map $\widehat{M} \rightarrow \widehat{M}_h$ corresponds to projection onto this summand.

If X is a G -space then, in the context of equivariant K -theory, we have maps

$$\text{res}_h : \widehat{G(G, X)} \rightarrow \widehat{G(Z, X)} \rightarrow \widehat{G(Z, X)}_h$$

and

$$\text{res}_h : \widehat{K_0(G, X)} \rightarrow \widehat{K_0(Z, X)} \rightarrow \widehat{K_0(Z, X)}_h$$

corresponding to the composition of the restriction maps $\widehat{G(G, X)} \rightarrow \widehat{G(Z, X)}$ (resp. $\widehat{K_0(G, X)} \rightarrow \widehat{K_0(Z, X)}$) with projection on the summand $\widehat{G(Z, X)}_h$ (resp. $\widehat{K_0(Z, X)}_h$). If in addition we assume that both G and Z are connected and reductive and have simply connected commutator subgroups, then the induction map $\text{ind} : G(Z, X) \rightarrow G(G, X)$ induces a map on completions at the ideal \mathfrak{m}_Ψ . Composing with the inclusion of $\widehat{G(Z, X)}_h$ as a summand in $\widehat{G(Z, X)}$ gives a map

$$\text{ind}_h : \widehat{G(Z, X)}_h \rightarrow \widehat{G(G, X)}.$$

Proposition 3.5. *If G and Z are connected, reductive and have simply connected commutator subgroups, then the maps res_h and ind_h are inverse isomorphisms of $\widehat{R(\widehat{G})}$ -modules.*

Before we prove Proposition 3.5 we need a lemma about Weyl groups of centralizers of semisimple elements.

Lemma 3.6. *Let G be a connected reductive group and let h be a semisimple element of G such that the centralizer $Z = Z_G(h)$ is connected. Let Ψ denote the conjugacy class of h in G . Let T be a maximal torus of Z (hence also of G). Let $W = W(G, T)$ and $W_1 = W(Z, T)$. Then $h \in T$, $\Psi \cap T = W \cdot h$, and the stabilizer in W of h is W_1 .*

Proof of Lemma 3.6. First, h is contained in some maximal torus of Z . The conjugacy of maximal tori in Z implies that the Z -conjugacy class of h meets T , but this class consists of $\{h\}$. Hence $h \in T$. Suppose that $h' = ghg^{-1} \in T$. Since $g^{-1}Tg = T_1$ and T are two maximal tori containing h , they are both contained in Z . Therefore there is an element $z \in Z$ such that $zTz^{-1} = T_1$. Then $h' = gzh(gz)^{-1}$.

The element gz of $N(T)$ represents an element of $W = N(T)/T$, so $h' \in W \cdot h$. Hence $\Psi \cap T \subseteq W \cdot h$; the reverse inclusion holds as well since elements of W are represented by elements of $N(T) \subset G$. This proves the first statement. For the second, suppose $w \in W$ fixes h . Then w is represented by an element of Z , so w lies in the Weyl group of Z , namely W_1 . \square

Proof of Proposition 3.5. Choose a maximal torus T of Z (and G). Let $W = W(G, T)$ and $W_1 = W(Z, T)$. By Lemma 3.6, the Weyl group W acts transitively on the set $\Psi \cap T$ and the stabilizer of h is W_1 . Thus, $\Psi \cap T = \{h_1, h_2, \dots, h_r\}$ where $r = \frac{|W|}{|W_1|}$ (we take $h = h_1$). Set $M = G(T, X)$ and let \widehat{M}_i be the completion of M at the maximal ideal \mathfrak{m}_{h_i} of $R(T)$. By Lemma 3.1, $\widehat{M} \simeq \bigoplus_{i=1}^r \widehat{M}_i$. By Lemma 3.2, $G(\widehat{G}, X) = (\widehat{M})^W$ and $G(\widehat{Z}, X)_h = (\widehat{M}_1)^{W_1}$. The map ind_h corresponds to the induction map $(\widehat{M}_1)^{W_1} \rightarrow (\widehat{M})^W$ defined in Section 3.1.3 above, and the map res_h corresponds to the projection $(\widehat{M})^W \rightarrow (\widehat{M}_1)^{W_1}$. Hence Lemma 3.3 implies that ind_h and res_h are inverse isomorphisms. \square

Example 3.7. Proposition 3.5 illustrates the need to work with completions rather than localizations, because the restriction map of localizations $G(G, X)_{\mathfrak{m}_\Psi} \rightarrow G(Z, X)_{\mathfrak{m}_h}$ will not generally be an isomorphism. For example, if $G = \text{GL}_n$ and Ψ is a regular conjugacy class (i.e. elements of Ψ have distinct eigenvalues), then $Z = T$ is a maximal torus. The restriction map $R(\text{GL}_n) \rightarrow R(T)$ has degree $n!$, so the map of local rings $R(G)_{\mathfrak{m}_\Psi} \rightarrow R(T)_{\mathfrak{m}_h}$ cannot be an isomorphism. However, this map of local rings is étale so the corresponding map of completions is indeed an isomorphism.

4. THE NON-ABELIAN COMPLETION THEOREM

In this section we state and prove Theorem 4.3, our general non-abelian completion theorem for equivariant K -theory. This theorem extends the non-abelian localization theorem of [EG3, Theorem 5.1 and Corollary 5.2] which was proved for actions with finite stabilizer.

4.1. The completion theorem for a central element. Let Z be an algebraic group and let h be an element in the center of Z . If M is an $R(Z)$ -module, we denote by \widehat{M}_h the \mathfrak{m}_h -adic completion of M , where, as usual, \mathfrak{m}_h is the ideal in $R(H)$ corresponding to virtual representations whose characters vanish at h .

If X is a Z -space, then since h is central, the fixed locus X^h is Z invariant. When X is smooth, then so is X^h , and thus the inclusion

$i_*: X^h \rightarrow X$ is a regular embedding. As a tool in proving our general completion theorem, we need the following result.

Theorem 4.1. *Let Z be an algebraic group and h a central element of Z . Let X be a Z -space and let $i: X^h \rightarrow X$ denote the inclusion of the fixed point locus of h .*

(a) *The proper pushforward $i_*: G(Z, X^h) \rightarrow G(Z, X)$ is an isomorphism after completing at \mathfrak{m}_h .*

(b) *If X is smooth, then the map of $R(Z)$ -modules*

$$\cap \lambda_{-1}(N_i^*): G(Z, X^h) \rightarrow G(Z, X^h)$$

is invertible after completing at \mathfrak{m}_h , and if $\alpha \in \widehat{G(Z, X)}_h$ then

$$(9) \quad \alpha = i_*(\lambda_{-1}(N_i^*)^{-1} \cap i^* \alpha),$$

where the notation $\lambda_{-1}(N_i^)^{-1} \cap i^* \alpha$ means inverse image of $i^* \alpha$ under the isomorphism $\cap \lambda_{-1}(N_i^*)$.*

Proof. The proof of Theorem 4.1 is essentially the same as the proof of the central localization theorem ([EG3, Theorem 3.3]). Details may be found in that paper; here is a brief sketch. By a change of groups argument the result for a general group reduces to the case of a product of general linear groups, and this in turn reduces to the case of a maximal torus. Therefore we may assume that $G = T$ is a torus. Using the localization long exact sequence for an open set, to prove part (a) it suffices to show that if X^h is empty then $\widehat{G(T, X)}_h = 0$. By Thomason's generic slice theorem we may reduce to the case that $X = T/T' \times Y$ where $T' \subset T$ is a closed subgroup not containing h , and the action of T on Y is trivial. In this case $G(T, X) = R(T') \otimes G(Y)$ and $\widehat{G(T, X)}_h = \widehat{R(T')}_h \otimes G(Y)$. The crucial point is that if $h \notin T'$ then $R(T')_{\mathfrak{m}_h} = 0$ (which gives the localization result) and $\widehat{R(T')}_h = 0$ (which gives the completion result). This proves (a). To prove (b), we again use the generic slice theorem to reduce to the case where $X^h = T/T' \times X^h/T$, but now $T' \subset T$ is a closed subgroup containing h . Then $G(T, X^h) = R(T') \otimes G(X^h/T)$ so $\widehat{G(T, X^h)}_h = \widehat{R(T')}_h \otimes G(X^h/T)$. Let \mathcal{N}_x denote the fiber of N_i at a point $x \in X^h$. The action of $\lambda_{-1}(N_i^*)$ on $\widehat{G(T, X^h)}_h$ is invertible if and only if $\lambda_{-1}(\mathcal{N}_x)$ is invertible in $\widehat{R(T)}_h$ for some closed point x in each connected component of X^h . This follows from the fact (proved in [EG3]) that $\lambda_{-1}(\mathcal{N}_i) \notin \mathfrak{m}_h$. Finally, to deduce (9), if $\alpha \in \widehat{G(Z, X)}_h$, then by (a) we can write $\alpha = i_* \beta$ for $\beta \in \widehat{G(Z, X^h)}_h$. By the equivariant self-intersection formula, $\lambda_{-1}(N_i^*) \beta = i^* \alpha$, so $\beta = \lambda_{-1}(N_i^*)^{-1} \cap i^* \alpha$. \square

Remark 4.2. In the proof of the central localization theorem of [EG3], the change of groups step proceeded by embedding G into a product of general linear groups Q such that the conjugacy class of h in Q intersects G in the single point h . We asserted that h central in G implies h central in Q . Unfortunately this assertion can fail for the embeddings we constructed in that paper. However, we can arrange it to be true if G is reductive, since then we can take Q to be $\prod GL(V_i)$, where V_i is an irreducible G -module; in this case the assertion follows from Schur's lemma, and the proof of the central localization theorem goes through. To deal with nonreductive G we replace G by a Levi factor L and use the identification of $G(G, X)$ with $G(L, X)$; then the argument works because L is reductive.

4.2. Statement of the non-abelian completion theorem. Let X be a G -space and Ψ a semisimple conjugacy class in G . Let h be an element of Ψ and $Z = \mathcal{Z}_G(h)$. As usual, let $i: X^h \rightarrow X$ be the inclusion of the fixed point locus of h ; as noted above, if X is smooth then so is X^h , and the inclusion is a regular embedding. Since Z acts on X^h , if X is smooth we may define a map $i^!: \widehat{G(G, X)} \rightarrow \widehat{G(Z, X^h)}_h$ as the composition

$$\widehat{G(G, X)} \xrightarrow{\text{res}_h} \widehat{G(Z, X)}_h \xrightarrow{i^*} \widehat{G(Z, X^h)}_h,$$

where res_h is as in Section 3.2.

For arbitrary X we may also define $i^!: K_0(\widehat{G(G, X)}) \rightarrow K_0(\widehat{G(Z, X^h)}_h)$ in the same way. This pullback will be used in the statement of Theorem 5.5.

Theorem 4.3. *Let X be a G -space, and use the notation above. There is an isomorphism $i_!: \widehat{G(Z, X^h)}_h \rightarrow \widehat{G(G, X)}$ with the following properties.*

(a) (covariance) *Let $Y \xrightarrow{p} X$ be a proper G -morphism of algebraic spaces, and let $Y^h \xrightarrow{q} X^h$ be the corresponding proper map of h -fixed loci. Then $i_! q_* = p_* i_!$ as maps $\widehat{G(Z, Y^h)}_h \rightarrow \widehat{G(G, X)}$.*

(b) *If X is smooth, then $i^!$ is also an isomorphism and we have the following analogue of the explicit localization formula:*

$$(10) \quad \alpha = i_! (\lambda_{-1}(N_i^*)^{-1} \cap i^! \alpha)$$

(c) *The map $i_!: \widehat{G_0(Z, X)}_h \rightarrow \widehat{G_0(G, X)}$ is uniquely determined by properties (a) and (b).*

Remark 4.4. We conjecture that the uniqueness statement of part (c) holds for higher K -theory as well; this would follow if we had a

Riemann-Roch theorem relating higher equivariant G -theory to higher equivariant Chow groups.

When G and Z are connected and reductive with simply connected commutator subgroup, then Proposition 3.5 states that for any G -space X , the map $\text{res}_h: \widehat{G(G, X)} \rightarrow \widehat{G(Z, X)}_h$ is an isomorphism. Using Theorem 4.3, we can partially generalize this to arbitrary G and Z :

Corollary 4.5. (a) *If X is a smooth G -space, then $\text{res}_h: \widehat{G(G, X)} \rightarrow \widehat{G(Z, X)}_h$ is an isomorphism. In particular, the restriction map $R(G) \rightarrow R(Z)$ induces an isomorphism between the \mathfrak{m}_Ψ -adic completion of $R(G)$ and the \mathfrak{m}_h -adic completion of $R(Z)$.*

(b) *For any G -space X , the restriction map of completed equivariant Grothendieck groups $\text{res}_h: \widehat{G_0(G, X)} \rightarrow \widehat{G_0(Z, X)}_h$ is an isomorphism.*

Proof. If X is smooth, then the central completion theorem implies that $i^*: \widehat{G(Z, X)}_h \rightarrow \widehat{G(Z, X^h)}_h$ is an isomorphism. Since $i_!$ is an isomorphism, formula (10) of Theorem 4.3 implies that $i^!$ is an isomorphism. Hence res_h must also be an isomorphism, proving (a). The proof of (b) uses envelopes and Riemann-Roch, and will be deferred to Section 6.2. \square

We conjecture that the conclusion of part (a) holds for arbitrary X .

4.3. Proof of Theorem 4.3(a),(b). In this section we prove parts (a) and (b) of Theorem 4.3. Part (c) uses a Chow envelopes argument and our generalized equivariant Riemann-Roch theorem; it will be proved in Section 6.2.

If G and Z are connected and reductive with simply connected commutator subgroups, we define $i_!$ as $\text{ind}_h \circ i_*$. By the central completion theorem (Theorem 4.1) i_* is an isomorphism, and by Proposition 3.5, ind_h is an isomorphism. If $Y \rightarrow X$ is proper, then the diagram

$$\begin{array}{ccc} Y^h & \rightarrow & Y \\ \downarrow & & \downarrow \\ X^h & \rightarrow & X \end{array}$$

is a commutative diagram of proper morphisms. Covariance of $i_!$ then follows from covariance of both pushforwards and ind_h for proper G -morphisms. Part (b) follows from Theorem 4.1(b) and the fact that $\text{ind}_h \circ \text{res}_h = \text{id}$.

To prove the general case we use a change of groups argument similar to that used in [EG3]. By [EG3, Proposition 2.8], we may embed G into a product of general linear groups Q such that, writing $\Psi = C_G(h)$ and $\Psi_Q = C_Q(h)$, we have $\Psi_Q \cap G = \Psi$.

Write $Z = \mathcal{Z}_G(h)$ and $Z_Q = \mathcal{Z}_Q(h)$. The groups Q and Z_Q are both connected and reductive and have simply connected commutator subgroups because they are both products of general linear groups. Let $Y = Q \times_G X$, and let $j : Y^h \rightarrow Y$ be the inclusion of the fixed point locus. By what we have proved, there is an isomorphism of completions $j_! : \widehat{G(Z_Q, Y^h)}_h \rightarrow \widehat{G(Q, Y)}$. Here $\widehat{G(Z_Q, Y^h)}_h$ denotes the completion of $G(Z_Q, Y^h)$ at the maximal ideal \mathfrak{m}'_h of $R(Z_Q)$ corresponding to the central element h , and $\widehat{G(Q, Y)}$ is the \mathfrak{m}_{Ψ_Q} -adic completion of $G(Q, Y)$.

We want to use $j_!$ to define $i_!$. First observe that the only maximal ideal of $R(G)$ containing $\mathfrak{m}_{\Psi_Q}R(G)$ is \mathfrak{m}_{Ψ} ([EG3, Proposition 2.6]), so $\mathfrak{m}_{\Psi_Q}R(G)$ is an \mathfrak{m}_{Ψ} -primary ideal. Thus, under the Morita equivalence which identifies $G(Q, Y)$ with $G(G, X)$, the \mathfrak{m}_{Ψ_Q} -adic and \mathfrak{m}_{Ψ} -adic topologies coincide. Hence we may identify the \mathfrak{m}_{Ψ_Q} -adic completion of $G(Q, Y)$ with the \mathfrak{m}_{Ψ} -adic completion of $G(G, X)$. Similarly, under the Morita equivalence identification of $G(Z, X^h)$ with $G(Z_Q, Z_Q \times_Z X^h)$, the \mathfrak{m}_h and \mathfrak{m}'_h -adic topologies coincide.

The following lemma is a consequence of an extension of [EG3, Lemma 5.5] to non-smooth spaces.

Lemma 4.6. *The natural map $Z_Q \times_Z X^h \rightarrow Y^h$ is an isomorphism of Z_Q -spaces.*

Proof of Lemma 4.6. Let $S = \{(g, x) \mid g \in \Psi, gx = x\} \subset G \times X$ and $S_Q = \{(q, y) \mid q \in \Psi_Q, qy = y\} \subset Q \times Y$. As in the proof of [EG3, Lemma 5.5], consider the map

$$T : Q \times S \rightarrow \Psi_Q \times Q \times X, \quad (q, g, x) \mapsto (qqq^{-1}, q, x).$$

This map induces a map of quotients $\tilde{\Phi} : Q \times_G S \rightarrow \Psi_Q \times Y$ which factors through a map $\Phi : Q \times_G S \rightarrow S_Q \subset \Psi_Q \times Y$. In view of the identifications $S = G \times_Z X^h$ and $S_Q = Q \times_{Z_Q} Y^h$ ([EG3, Lemma 4.3]), to prove the lemma, it suffices to show that $\Phi : Q \times_G S \rightarrow S_Q$ is an isomorphism of Q -spaces. This was proved for smooth X in [EG3, Lemma 5.5]; to extend to arbitrary X we argue as follows. To check that Φ is an isomorphism, we may work locally in the smooth topology on X and assume that X is affine.³ By local linearizability of group actions [Bor], there is a G -equivariant embedding $X \subset V$ for some finite dimensional representation V of G . Let $W = Q \times_G V$. Define $S' = \{(g, v) \mid g \in \Psi, gv = v\} \subset \Psi \times V$ and $S'_Q = \{(q, w) \mid q \in \Psi_Q, qw =$

³Every G -space has a smooth G -cover by an affine G -scheme. To see this, let $U \xrightarrow{\pi} X$ be any étale cover of X by a scheme. Replacing U by an affine Zariski open cover, we may assume that U is also affine. Then the map $G \times U \rightarrow X$, $(g, u) \mapsto g\pi(u)$, is a smooth, affine G -cover of X .

$w\} \subset Q \times W$. Since V is smooth, the map $\Phi': Q \times_G S' \rightarrow S'_Q$ is an isomorphism. Now, $X^h = X \cap V^h$, so $S = S' \cap (\Psi \times X)$. Hence the isomorphism Φ' restricts to the map $\Phi: Q \times_G S \rightarrow S_Q$. Therefore, Φ is an isomorphism. \square

Hence we may identify the \mathfrak{m}_h -adic completion of $G(Z, X^h)$ with the \mathfrak{m}'_h -adic completion of $G(Z_Q, Y^h)$.

Applying Lemma 4.6 to the case $X = \text{Spec } \mathbb{C}$, we see that Z_Q/Z can be identified with $(Q/G)^h$. Let $e: Z_Q/Z \rightarrow Q/G$ be the corresponding Z_Q -equivariant regular embedding. Under the identification of $G(Z_Q, Z_Q/Z)$ with the representation ring $R(Z)$, the class of the conormal bundle $[N_e^*]$ corresponds to the virtual Z -module $\mathfrak{q}^*/\mathfrak{g}^* - \mathfrak{z}_Q^*/\mathfrak{z}^* = \mathfrak{q}^*/\mathfrak{z}_Q^* - \mathfrak{g}^*/\mathfrak{z}^*$, where \mathfrak{q} , \mathfrak{z}_Q , \mathfrak{g} , and \mathfrak{z} denote the Lie algebras of the groups Q , Z_Q , G , and Z , respectively. The central completion theorem implies that $\lambda_{-1}(N_e^*)$ is invertible in $G(\widehat{Z_Q, Z_Q/Z})_h = \widehat{R(Z)}_h$.

To complete the proof of the theorem for general G , we use the identifications of $G(\widehat{Z, X^h})_h$ with $G(\widehat{Z_Q, Y^h})_h$ and $G(\widehat{G, X})$ with $G(\widehat{Q, Y})$ to define

$$i_! : G(\widehat{Z, X^h})_h \rightarrow G(\widehat{G, X})$$

by the formula

$$(11) \quad i_! \beta = j_! (\lambda_{-1}(N_e^*)^{-1} \cap \beta)$$

Because $j_!$ is an isomorphism, $i_!$ is as well, and the covariance of $j_!$ implies covariance of $i_!$.

Now suppose that X is smooth. To verify formula (10) of Theorem 4.3, observe that $N_j^* = N_i^* + N_e^*$ in $G(Z, X^h) = G(Z_Q, Y^h)$, because j factors as the composition of embeddings

$$Y^h = (Q \times_G X)^h = (Z_Q \times_Z X^h) \rightarrow Z_Q \times_Z X \rightarrow Y = Q \times_G X.$$

Moreover, under our identifications, the maps $j^!$ and $i^!$ coincide. Thus, formula (10) for $i_!$ follows from the corresponding formula for $j_!$.

Remark 4.7. A priori, our definition of $i_!$ depends on the choice of the embedding of G into Q . However, if X is smooth, then $i_!$ is independent of the embedding, because equation (10) implies that composition of $i_!$ with the map $\cap \lambda_{-1}(N_i^*)^{-1}$ is the inverse of $i^!$. Since $i^!$ is defined intrinsically, so is $i_!$ in this case. For arbitrary X , the uniqueness statement Theorem 4.3(c) implies that $i_!$ is independent of Q as a map of completed Grothendieck groups. As noted above, we conjecture that Theorem 4.3(c) extends to higher K -theory; this would imply that $i_!$ is independent of Q in the higher K -theory case as well.

4.4. The case of finite stabilizer. In this section we show that if G acts with finite stabilizer on a smooth space X then the nonabelian completion theorem is equivalent to the nonabelian localization theorem of [EG3]. In the finite stabilizer case, the equivariant K -theory is supported at a finite number of maximal ideals, so localization and completion coincide. Therefore the issue is to compare the map used in this paper with the one of [EG3].

Let G be an algebraic group acting with finite stabilizer on a smooth algebraic space X . Following the notation of [EG3], let Ψ be a semisimple conjugacy class in G and let $S_\Psi = \{(g, x) | g \in \Psi, gx = x\}$. Choose $h \in \Psi$, and let $Z = \mathcal{Z}_G(h)$. Then $S_\Psi = G \times_Z X^h$, so by Morita equivalence we may identify $G(G, S_\Psi)$ with $G(Z, X^h)$, which is an $R(Z)$ -module. By [EG3, Lemma 4.6] the localization $G(G, S_\Psi)_{\mathfrak{m}_h} = G(Z, X^h)_{\mathfrak{m}_h}$ at \mathfrak{m}_h is a summand in the localization $G(G, S_\Psi)_{\mathfrak{m}_\Psi}$, and is independent of the choice of $h \in \Psi$. This summand is called the *central summand* and is denoted by $G(G, S_\Psi)_{c_\Psi}$ [EG3, Definition 4.10]. Because G acts with finite stabilizer, the G -equivariant and Z -equivariant Grothendieck groups are supported at a finite number of maximal ideals, so completion and localization coincide. Thus,

$$G(G, S_\Psi)_{c_\Psi} = G(Z, X^h)_{\mathfrak{m}_h} = \widehat{G(Z, X^h)}_{\mathfrak{m}_h}.$$

The hypothesis that G acts with finite stabilizer implies that the projection $f: S_\Psi \rightarrow X$ is finite so it induces a pushforward $f_*: G(Z, X^h) \rightarrow G(G, X)$. The nonabelian localization theorem for actions with finite stabilizer [EG3, Theorem 5.1] states that if $\alpha \in G(G, X)_{\mathfrak{m}_\Psi}$ then

$$(12) \quad \alpha = f_* (\lambda_{-1}(N_f^*)^{-1} \cap (f^*\alpha)_{c_\Psi})$$

where $(f^*\alpha)_{c_\Psi}$ denotes the projection of $f^*\alpha$ to the central summand and N_f is the relative tangent bundle of the l.c.i. morphism $S_\Psi \rightarrow X$.

Remark 4.8. Because the map $f: S_\Psi \rightarrow X$ is not an embedding, N_f^* is only a virtual bundle, and the class $\lambda_{-1}(N_f^*)$ is not defined in $K_0(G, S_\Psi)$. Nevertheless, we can define the operation $\cap \lambda_{-1}(N_f^*)^{-1}$ on $G(G, S_\Psi)_{c_\Psi}$, as follows. Since f factors as the composition of the smooth projection $G \times_Z X \rightarrow X$ with the regular embedding $i: S_\Psi \rightarrow G \times_Z X$, we have $[N_f^*] = [N_i^*] - \mathfrak{g}^*/\mathfrak{z}^*$ (here, as usual, we follow the conventions of Section 1.1 in identifying relative tangent bundles). After localizing $G(G, S_\Psi)$ at \mathfrak{m}_h , the endomorphism $\cap \lambda_{-1}(N_i^*)$ of $G(G, S_\Psi)_{c_\Psi}$ is invertible. Hence we can define $\cap \lambda_{-1}(N_f^*)^{-1}$ as the composition of the inverse of the operation $\cap \lambda_{-1}(N_i^*)$ with multiplication by $\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*)$. This point was obscured in [EG3].

By abuse of notation use i to denote both the Z -equivariant embedding $i: X^h \rightarrow X$ and the G -equivariant embedding $i: S_\Psi \rightarrow G \times_Z X$; this convention has the advantage that under our Morita equivalence identifications the same symbol i_* denotes the proper pushforwards $G(Z, X^h) \rightarrow G(Z, X)$ and $G(G, S_\Psi) \rightarrow G(G, G \times_Z X)$. Likewise, the conormal bundles to both of these maps may be denoted N_i^* . With this convention we can rewrite (12) as

$$(13) \quad \alpha = f_* (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*)(\lambda_{-1}(N_i^*)^{-1} \cap (f^*\alpha)_{c_\Psi})).$$

Under our identifications, if $\alpha \in \widehat{G(G, X)}$, then $f^*(\alpha)_{c_\Psi} = i^!(\alpha)$. The following result implies that when G acts on X with finite stabilizers then the nonabelian localization theorem of [EG3] is in fact equivalent to Theorem 4.3.

Theorem 4.9. *Assume that G acts on the smooth space X with finite stabilizer. If $\beta \in G(G, S_\Psi)_{c_\Psi}$, then*

$$(14) \quad f_* (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*) \cap \beta) = i_!\beta$$

(as elements of $G(G, X)_{\mathfrak{m}_\Psi}$).

Proof. Since the map $\alpha \mapsto f^*(\alpha)_{c_\Psi} = i^!(\alpha)$ is an isomorphism and $G(G, S_\Psi)_{c_\Psi} = \widehat{G(Z, X^h)}_h$, we can compare formulas (10) and (12) to obtain

$$(15) \quad f_*(\lambda_{-1}(N_f^*)^{-1} \cap \beta) = i_!(\lambda_{-1}((N_i^*)^{-1} \cap \beta)).$$

But as noted in the remark above,

$$\lambda_{-1}(N_f^*)^{-1} \cap \beta = \lambda_{-1}(N_i^*)^{-1} \cap (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*) \cap \beta).$$

Replacing β by $\lambda_{-1}(N_i^*) \cap \beta$ in (15) yields the formula of equation (14). \square

4.5. More on the finite stabilizer case. The proof of Theorem 4.9 used the localization and completion theorems. However, if G is connected and reductive with simply connected commutator subgroup and Z is a Levi factor of parabolic subgroup P of G then, by Proposition 2.6, Z also has a simply connected commutator. In this case we can prove a stronger result (Theorem 4.10) namely, that the maps agree even before localizing (or completing). This proof does not make use of the localization or completion theorems. Using Theorem 4.10 one could replace some constructions from [EG3] with constructions from this paper and obtain somewhat different proofs of the main results in [EG3]. However, Theorem 4.10 is not needed for the main results of this paper.

Assume that we are still in the finite stabilizer case. The proper pushforward f_* is defined without localizing (or completing); that is, f_* maps $G(G, S_\Psi)$ to $G(G, X)$. The map $i_!$ can also be defined without localizing or completing; we identify $G(G, S_\Psi)$ with $G(Z, X^h)$ and then define $i_! = \text{ind} \circ i_*$. In this case, the equality of Theorem 4.9 holds even without localizing:

Theorem 4.10. *Suppose that the connected reductive group G acts with finite stabilizer on a smooth space X . Let $h \in G$ be an element of the semisimple conjugacy class Ψ , and assume that $Z = \mathcal{Z}_G(h)$ is a Levi factor of a parabolic subgroup P of G . If $\beta \in G(G, S_\Psi)$, then*

$$(16) \quad f_* (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*) \cap \beta) = i_! \beta$$

as elements of $G(G, X)$.

Proof. We use the factorization of the map $S_\Psi \rightarrow G \times_P X$ from Section 4.1 of [EG3]. Consider the projections

$$G \times_Z X \xrightarrow{p} G \times_P X \xrightarrow{q} X.$$

By [EG3, Lemma 4.5] the composition $j = p \circ i: S_\Psi \rightarrow G \times_P X$ is also a regular embedding and the finite map $f: S_\Psi \rightarrow X$ is the composition of j with the smooth proper projection map q . Thus,

$$(17) \quad f_* (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*) \cap \beta) = q_* j_* (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*) \cap \beta).$$

On the other hand, $i_! \beta = \text{ind} \circ i_* \beta$. By Proposition 2.7 the right hand side of (17) equals

$$(18) \quad q_* (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{p}^*) \cap (p^*)^{-1} i_* \beta)$$

(Note that the symbol $(p^*)^{-1}$ was suppressed in the statement of Proposition 2.7.)

The projection p is a bundle map with fiber isomorphic to the affine space P/Z . The tangent bundle T_p to p is the bundle induced by Morita equivalence from the Z -module $\mathfrak{p}/\mathfrak{z}$. By abuse of notation we will write $T_p = \mathfrak{p}/\mathfrak{z}$. Letting the unipotent radical of P act trivially we may also view $\mathfrak{p}/\mathfrak{z}$ as a P -module. It follows that T_p is the pullback of the G -bundle on $G \times_P X$ induced by Morita equivalence from $\mathfrak{p}/\mathfrak{z}$ (viewed as Z -module). Continuing our abuse of notation we will also refer to this bundle on $G \times_P X$ as $\mathfrak{p}/\mathfrak{z}$.

Claim: If $\beta \in G(G, S_\Psi)$ then

$$(19) \quad i_* \beta = \lambda_{-1}(\mathfrak{p}^*/\mathfrak{z}^*) \cap p^* j_* \beta$$

We prove the claim. Let $\rho: W \rightarrow S_\Psi$ be the P/Z -bundle obtained by base change from the bundle $p: G \times_Z X \rightarrow G \times_P X$ along the morphism $j: S_\Psi \rightarrow G \times_P X$. Let $k: W \rightarrow G \times_Z X$ be the map obtained by base

change from j . The map $i: S_\Psi \rightarrow G \times_Z X$ induces a section $s: S_\Psi \rightarrow W$ of the projection ρ . In particular we have a cartesian diagram

$$\begin{array}{ccc} W & \xrightarrow{k} & G \times_Z X \\ \downarrow \rho & & \downarrow p \\ S_\Psi & \xrightarrow{j} & G \times_P X \end{array}$$

where the horizontal maps are regular embeddings.

Since direct image for finite morphisms commutes with flat pullback of coherent sheaves, the compositions $p^* \circ j_*$ and $k_* \circ \rho^*$ are equal as maps on K -theory. Thus the right hand side of (19) equals

$$(20) \quad \lambda_{-1}(\mathfrak{p}^*/\mathfrak{z}^*) \cap k_* \rho^* \beta$$

Since the bundle $\lambda_{-1}(\mathfrak{p}^*/\mathfrak{z}^*)$ on W is the pullback of the corresponding bundle on $G \times_Z X$, the the projection formula implies that the right hand side of (20) equals

$$(21) \quad k_*(\lambda_{-1}(\mathfrak{p}^*/\mathfrak{z}^*) \cap \rho^* \beta)$$

The normal bundle to the section $s: S_\Psi \rightarrow W$ is the restriction of T_ρ to s . Now, T_ρ is the pullback of T_p , which is the pullback to $G \times^Z X$ of the G -bundle on $G \times_P X$ induced by Morita equivalence from $\mathfrak{p}/\mathfrak{z}$ (viewed as P -module). By abuse of notation we will refer to this bundle on $G \times_P X$, as well as the various pullbacks of this bundle, as $\mathfrak{p}/\mathfrak{z}$. Thus, by the self intersection formula $s^* s_* \beta = \lambda_{-1}(\mathfrak{p}^*/\mathfrak{z}^*) \cap \beta$. Substituting into (21) yields

$$(22) \quad k_*(\rho^* s^* s_* \beta)$$

Since s is a section of the smooth morphism ρ , the composition $\rho^* \circ s^*$ is the identity. Hence we conclude that $\lambda_{-1}(\mathfrak{p}^*/\mathfrak{z}^*) \cap \rho^* j_* \beta = k_* s_* \beta = \beta$, proving our claim.

Substituting the expression for $i_* \beta$ for (19) into the right hand side of (18) we obtain

$$(23) \quad q_*(\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*) \cap (\lambda_{-1}(\mathfrak{p}^*/\mathfrak{z}^*) \cap j_* \beta))$$

Since all of the terms involving Lie algebras correspond to bundles which are pulled back from $G \times_P X$, the projection formula implies that (23) can be rewritten as

$$(24) \quad q_* j_* (\lambda_{-1}(\mathfrak{g}^*/\mathfrak{p}^*) \lambda_{-1}(\mathfrak{p}^*/\mathfrak{z}^*) \cap \beta).$$

Since $f = q \circ j$ and $\mathfrak{g}^*/\mathfrak{z}^* = \mathfrak{g}^*/\mathfrak{p}^* + \mathfrak{p}^*/\mathfrak{z}^*$ in $R(Z)$ we may rewrite (24) as $f_*(\lambda_{-1}(\mathfrak{g}^*/\mathfrak{z}^*) \cap \beta)$, completing the proof of the theorem. \square

5. THE TWISTED EQUIVARIANT RIEMANN-ROCH THEOREM

As above, let $\Psi = C_G(h)$ be a semi-simple conjugacy class. In this section we prove a Riemann-Roch isomorphism generalizing the main theorem of [EG2], which dealt with the case $h = 1$. This theorem gives a geometric description of the completion of equivariant K -theory at any maximal ideal in $R(G)$.

5.1. The equivariant Riemann-Roch theorem. We recall here the equivariant Riemann-Roch theorem of [EG2]. We begin by recalling some basic facts about equivariant Chow groups. Most of these facts are contained in the paper [EG1], but here we continue to use some notation from [EG2]. If X is a G -space then as in [EG2] we denote $CH_G^n(X)$ the “codimension n ” equivariant Chow groups. An element of $CH_G^n(X)$ is represented by a codimension n cycle on a quotient $X \times_G U$; here U is an open set in a representation V such that G acts freely on U , and such that the complement of U has codimension more than n in V . Less precisely, but more intuitively, we may view an element of $CH_G^n(X)$ as being represented by a G -invariant cycle of codimension n on a product $X \times V$ where V is a representation of G . We denote by $A_G^k(X)$ the “codimension n ” operational Chow ring. An element $x \in A_G^k(X)$ is an operation on $CH_G^*(X)$ which increases degree by k and satisfies basic naturality properties with respect to equivariant morphisms. The direct sum $A_G^*(X) = \bigoplus_{k=0}^{\infty} A_G^k(X)$ is a graded, commutative ring with multiplication defined by composition of operations. When X is a smooth algebraic space, then a basic result is that the map $A_G^*(X) \rightarrow CH_G^*(X)$, $x \mapsto x \cap [X]$ is an isomorphism, where $[X] \in CH_G^n(X)$ is the fundamental class. Thus, when X is smooth we may view the product of operations as an intersection product.

The infinite direct product $\prod_{k=0}^{\infty} A_G^k(X)$ is a completion of the equivariant operational Chow ring; it acts on the infinite product of equivariant Chow groups $\prod_{n=0}^{\infty} CH_G^n(X)$. If E is an equivariant vector bundle then we may define the equivariant Chern character $\text{ch}^G(E)$ and equivariant Todd class $\text{Td}^G(E)$ as the appropriate formal power series in the Chern classes of E , [EG2, Definition 3.1]. As such they may be viewed as elements in the infinite product $\prod_{n=0}^{\infty} A_G^n(X)$. The Chern character and Todd class extend to maps ch^G and Td^G from $K_0(G, X)$ to $\prod_{n=0}^{\infty} A_G^n(X)$.

We paraphrase from [EG2] the Riemann-Roch theorem for equivariant Chow groups. Let X be a G -space and let $\widehat{G_0(G, X)}$ denote the completion of $G_0(G, X)$ at the augmentation ideal in $R(G)$ —in other words, the completion at the maximal ideal $\mathfrak{m}_1 \subset R(G)$.

Theorem 5.1. *For all separated G -spaces X there is a map $\tau_X^G: G_0(G, X) \rightarrow \prod_{n=0}^{\infty} CH_G^n(X)$ which factors through an isomorphism $G_0(G, X) \rightarrow \prod_{n=0}^{\infty} CH_G^n(X)$ and satisfies the following properties.*

- (a) τ_X^G is covariant for proper morphisms.
- (b) If $\epsilon \in K_0(G, X)$ and $\alpha \in G_0(G, X)$ then $\tau_X^G(\epsilon\alpha) = \text{ch}^G(\epsilon)\tau^G(\alpha)$.
- (c) If X is a smooth scheme and either G is connected or X has a G -equivariant ample line bundle, then $\tau^G(\alpha) = \text{ch}(\alpha) \text{Td}^G(X)$, where $\text{Td}^G(X)$ is the equivariant Todd class of the virtual bundle $T_X - \mathfrak{g}$. (Here we implicitly identify $K_0(G, X)$ and $G_0(G, X)$ as X is a smooth scheme.

Remark 5.2. The notation used here differs somewhat from that of [EG2]: higher K -theory is not used in that paper, and the groups $G^G(X)$ and $K^G(X)$ of that paper correspond to the groups denoted here by $G_0(G, X)$ and $K_0(G, X)$, respectively.

Remark 5.3. The need to subtract \mathfrak{g} follows from the fact that if G acts freely on X and $X \xrightarrow{\pi} X/G$ is the quotient map, then, by definition, the Todd class map is the pullback of the Todd class map on X/G to the corresponding equivariant theories on X . However as an element of $K_0^G(X)$, $\pi^*(T_{X/G}) = T_X - \mathfrak{g}$.

5.2. Twisting by a central element. We recall the following construction from [EG3]. Let Z be an algebraic group, and let h be a semisimple element of the center of Z . If V is any representation of Z , then V decomposes as a direct sum of H -eigenspaces V_χ , where $\chi \in \mathbb{C}^*$. The assignment $[V] \rightarrow \sum_\chi \chi[V_\chi]$ defines an automorphism of $R(Z)$ which we call twisting by h . It is clear from the definition that this automorphism takes \mathfrak{m}_h to the augmentation ideal \mathfrak{m}_1 .

More generally, let X be a Z -space on which h acts trivially. Then any Z -equivariant coherent sheaf \mathcal{F} decomposes into a direct sum of eigensheaves \mathcal{F}_χ for the action of h . Define the twist of \mathcal{F} by h to be the virtual coherent sheaf $\mathcal{F}(h) = \sum_\chi \chi \mathcal{F}_\chi$. It is easy to see that twisting by a central element induces a natural automorphism of equivariant Grothendieck groups. The basic properties of this automorphism are as follows.

Proposition 5.4. *Let X be a Z -space on which the central element h acts trivially. Then the assignment $\mathcal{F} \mapsto \mathcal{F}(h)$ induces automorphisms of equivariant Grothendieck rings $t_h: K_0(Z, X) \rightarrow K_0(Z, X)$ and of equivariant Grothendieck groups $t_h: G_0(Z, X) \rightarrow G_0(Z, X)$ with the following properties.*

- (a) If $\epsilon \in K_0(Z, X)$ and $\alpha \in G_0(Z, X)$, then $t_h(\epsilon\alpha) = t_h(\epsilon)t_h(\alpha)$.

(b) If $f: Y \rightarrow X$ is an morphism of Z -spaces and h acts trivially on Y , then $f^* \circ t_h = t_h \circ f^*$ as maps $K_0(Z, X) \rightarrow K_0(Z, Y)$.

(c) t_h commutes with flat pullbacks and proper pushforwards of Grothendieck groups of equivariant coherent sheaves, provided the element h acts trivially on the spaces involved.

(d) t_h induces isomorphisms of completions $\widehat{K_0(Z, X)}_h \rightarrow \widehat{K_0(Z, X)}_1$ and $\widehat{G_0(Z, X)}_h \rightarrow \widehat{G_0(Z, X)}_1$.

5.3. The twisted equivariant Riemann-Roch theorem. We can now state the twisted equivariant Riemann-Roch theorem. For each algebraic G -space X , and each semisimple conjugacy class Ψ in G , this theorem gives a way to represent elements of the \mathfrak{m}_Ψ -adic completion of $G_0(G, X)$ by Z -equivariant cycles on X^h .

As usual, we fix Ψ and let $\widehat{G_0(G, X)}$ denote the \mathfrak{m}_Ψ -adic completion of $G_0(G, X)$.

Theorem 5.5. *For all separated G -spaces X , there is an isomorphism $\tau_X^\Psi: \widehat{G_0(G, X)} \rightarrow \prod_{i=0}^\infty CH_Z^i(X^h)$ with the following properties.*

- (a) τ_X^Ψ is covariant for proper morphisms.
- (b) If $\epsilon \in K_0(G, X)$ then

$$\tau_X^\Psi(\epsilon\alpha) = \text{ch}^Z(t_h(i^!\epsilon)) \cap \tau_X^\Psi(\alpha).$$

(c) If X is smooth, then $\text{ch}^Z(t_h(N_i^*))$ is invertible in the infinite product $\prod_{n=0}^\infty CH_Z^n(X^h)$. If in addition X^h is a scheme, and either Z is connected or X^h has a Z -equivariant ample line bundle, then

$$\tau_X^\Psi(\alpha) = \text{ch}^Z(t_h(i^!\alpha)) \text{ch}^Z(t_h(\lambda_{-1}(N_i^*)))^{-1} \text{Td}^Z(X^h).$$

(d) The map τ_X^Ψ is uniquely determined for G -schemes by properties (a) and (c).

Remark 5.6. When X and X^h are smooth schemes then both $G_0(G, X) = K_0(G, X)$ and $\prod_{i=0}^\infty CH_Z^i(X^h)$ have products. The map τ_X^Ψ is not a ring isomorphism but standard properties of the Chern character imply that the map $K_0(G, X) \rightarrow \prod_{i=0}^\infty CH_Z^i(X^h)$, $\epsilon \mapsto \text{ch}(t_h(i^!\epsilon))$ induces a ring isomorphism $\widehat{G_0(G, X)} \rightarrow \prod_{i=0}^\infty CH_Z^i(X^h)$.

Remark 5.7. We conjecture that τ_X^Ψ is also uniquely determined by properties (a) and (c) for all algebraic G -spaces. However, the proof of uniqueness uses an equivariant version of Chow's lemma for schemes (Proposition 6.3), and we do not know if the corresponding result holds for algebraic spaces.

Proof. Define τ_X^Ψ to be the composition $\tau_{X^h}^Z \circ t_h \circ (i_!)^{-1}$. The fact that τ_X^Ψ is an isomorphism, as well as the covariance for proper morphisms, follows because these properties hold for $i_!$ (Theorem 4.3), t_h (Proposition 5.4), and $\tau_{X^h}^Z$ (Theorem 5.1).

We now prove part (b). Let $G \subset Q$ be an embedding of G into a connected reductive group such that $Z_Q = \mathcal{Z}_Q(h)$ is also connected and reductive, and such that $\Psi_Q \cap G = \Psi$. As in the proof of Theorem 4.3, let $Y = Q \times_G X$. In the notation of that proof, the map of completions $i_!: \widehat{G(Z, X)}_h \rightarrow \widehat{G(G, X)}$ is defined as the composition of maps $\text{ind}_h \circ j_* \circ (\cap \lambda_{-1}(N_e^*)^{-1})$. Here

$$\text{ind}_h: \widehat{G(Z_Q, Y)} \rightarrow \widehat{G(Q, Y)} = \widehat{G(G, X)}$$

is the induction map corresponding to the inclusion $Z_Q \subset Q$, $j_*: \widehat{G(Z, X^h)}_h = \widehat{G(Z_Q, Y^h)}_h \rightarrow \widehat{G(Z_Q, Y)}_h$ is the map induced by the inclusion $j: Y^h \rightarrow Y$, and N_e is the normal bundle to Z_Q/Z in Q/G . Thus, if $\epsilon \in K_0(G, X)$, then

$$\begin{aligned} (i_!)^{-1}(\epsilon\alpha) &= \lambda_{-1}(N_e^*) \cap (j_*)^{-1}(\text{res}_h(\epsilon\alpha)) \\ &= \lambda_{-1}(N_e^*) \cap j^* \text{res}_h(\epsilon) \cap (j_*)^{-1}(\text{res}_h \alpha) \\ &= j^! \epsilon \cap \lambda_{-1}(N_e^*) \cap (j_*)^{-1}(\text{res}_h \alpha) \\ &= i^! \epsilon \cap (i_!)^{-1}(\alpha). \end{aligned}$$

The first equality holds since by Proposition 3.5, ind_h and res_h are inverse isomorphisms of $\widehat{G(Q, Y)}$ and $\widehat{G(Z_Q, Y)}_h$. The second equality is a K -theoretic consequence of the projection formula for the direct image functor j_* . The third equality follows from the definition of $j^!$. Finally, the fourth equality follows from the fact that $i^!$ and $j^!$ are equal as maps $K_0(G, X) = K_0(Q, Y) \rightarrow K_0(Z_Q, Y^h) = K_0(Z, X^h)$. Applying Theorem 5.1(b), we see that

$$\tau_X^\Psi(\epsilon\alpha) = \text{ch}(t_h(i^! \epsilon)) \cap \tau_X^\Psi(\alpha),$$

proving (b).

We now prove (c). Since multiplication by $\lambda_{-1}(N_i^*)$ is an automorphism of $G_0(Z, X^h)_h$, it follows that multiplication by $t_h(\lambda_{-1}(N_i^*))$ is an automorphism of $G_0(\widehat{Z, X^h})_1$. Since $\tau_{X^h}^Z: G_0(\widehat{Z, X^h})_1 \rightarrow \prod_{i=0}^\infty CH_Z^i(X^h)$ is an isomorphism, and

$$\tau_{X^h}^Z(t_h(\lambda_{-1}(N_i^*)) \cap \alpha) = \text{ch}(t_h(\lambda_{-1}(N_i^*))) \tau_{X^h}(\alpha),$$

it follows that multiplication by $\text{ch}(t_h(\lambda_{-1}(N_i^*)))$ is an isomorphism, proving the first assertion. The formula for $\tau_X^\Psi(\alpha)$ now follows from Theorem 4.3 and Theorem 5.1(c).

The proof of Part (d) uses equivariant Chow envelopes and will be deferred to Section 6.2. \square

By taking X to be a point, we obtain the following corollary relating completions of representation rings and Chow rings of classifying stacks. (By definition, the Chow ring of the classifying stack BG is the G -equivariant Chow ring of a point.)

Corollary 5.8. *Let G be a complex algebraic group, Ψ the conjugacy class of a semisimple element $h \in G$, and $Z = \mathcal{Z}_G(h)$. The map $R(G) \rightarrow \prod_{i=0}^{\infty} CH^i(BZ)$ which takes $[V]$ to $\text{ch}(t_h([\text{res } V]))$ induces an isomorphism $\widehat{R(G)} \rightarrow \prod_{i=0}^{\infty} CH^i(BZ)$.*

6. EQUIVARIANT ENVELOPES AND PROOFS OF UNIQUENESS

In this section we use envelope arguments to prove the uniqueness statements Theorem 4.3(c) and Theorem 5.5(d), as well as the isomorphism of completions stated in Corollary 4.5(b).

6.1. Equivariant envelopes.

Definition 6.1. Let X be a G -space. A proper G -equivariant map $\tilde{X} \rightarrow X$ is an *equivariant envelope* if for every G -invariant subspace $W \subset X$ whose connected components are integral, there is a corresponding G -invariant subspace $\tilde{W} \subset \tilde{X}$, with integral connected components, such that the map $\tilde{W} \rightarrow W$ is an isomorphism over a dense open subspace of W .

If G is connected then every component of an invariant subspace is invariant, so it would suffice to require the existence of \tilde{W} for every G -invariant subvariety W .

By canonical resolution of singularities for algebraic spaces (cf. [BM]), every separated algebraic G -space over a field of characteristic 0 has a nonsingular equivariant envelope [EG2, Corollary 2].

The following lemma will be used in the proofs of Theorem 4.3(c) and Corollary 4.5(b).

Lemma 6.2. *Let X be a separated G -space, and let $\pi: \tilde{X} \rightarrow X$ be an equivariant envelope. If $h \in Z(G)$ then the induced map $\tilde{X}^h \rightarrow X^h$ is also an equivariant envelope.*

Proof. Since h is central in G , the fixed loci X^h and \tilde{X}^h are G -invariant, so the map $\pi: \tilde{X}^h \rightarrow X^h$ is G -equivariant. Let $W \subset X^h$ be any G -invariant subspace with integral connected components, and let \tilde{W} be a subspace of \tilde{X} as in Definition 6.1. To prove that \tilde{X}^h is an equivariant envelope it suffices to prove that we can take $\tilde{W} \subset \tilde{X}^h$. The restricted

map $\pi: \tilde{W} \rightarrow W$ is a G -equivariant isomorphism over a dense open subspace of U of W . Replace \tilde{W} by the closure of $\pi^{-1}(U)$. Since h acts trivially on $\pi^{-1}(U)$ and \tilde{W}^h is closed, $\tilde{W} \subset \tilde{X}^h$ as desired. \square

6.1.1. *Equivariant Chow envelopes.* Let X be an algebraic space with a G -action. An equivariant envelope $\tilde{X} \rightarrow X$ is an *equivariant Chow envelope* if \tilde{X} has a G -linearized ample line bundle. In this section we prove that nonsingular Chow envelopes exist in the case where X is a separated scheme (Proposition 6.5). As a first step, we need an equivariant version of Chow's lemma, which extends a result of Sumihiro to disconnected groups. Once we establish this version of Chow's lemma, Proposition 6.5 will follow from equivariant resolution of singularities and Noetherian induction.

Proposition 6.3 (Equivariant Chow's Lemma). *Let X be a reduced and separated G -scheme. There exists a proper G -map $X' \xrightarrow{\pi} X$ such that X' has a G -linearized ample line bundle, and such that there is a dense open G -invariant subspace $U \subset X$ such that $\pi^{-1}(U) \rightarrow U$ is an isomorphism.*

Proof of Proposition 6.3. When G is connected this is proved by Sumihiro in [Sum]. In general, let G_0 be the identity component of G . Let $X_1 \xrightarrow{\pi} X$ be a proper G_0 -map with X_1 normal and quasi-projective and π an isomorphism over a dense G_0 -invariant open set U in X . By replacing U by the intersection of all σU , where σ runs over a set of coset representatives of G_0 in G , we may assume that U is G -invariant. Then the inverse image U_1 of U admits a G -action (since it is isomorphic to U). To produce a G -equivariant Chow cover we apply a construction used in the proof of [Sum, Theorem 3].

If Y is any G_0 -space, let IY denote the set of all functions $f: G \rightarrow Y$ satisfying $f(gg_0) = g_0^{-1}f(g)$ for all $g \in G$ and $g_0 \in G_0$. As a space, IY is isomorphic to Y^r by the map taking f to $(f(\sigma_1), \dots, f(\sigma_r))$, where $\sigma_1, \dots, \sigma_r$ is a collection of coset representatives of G_0 in G . (This isomorphism depends on the choice of coset representatives.) The group G acts on IY by the rule $(g \cdot f)(h) = f(g^{-1}h)$.

If Y has a G_0 -equivariant ample line bundle, then IY has a G -equivariant ample line bundle. This follows because a G_0 -equivariant embedding of Y into \mathbb{P}^N induces a G -equivariant embedding of IY into $I\mathbb{P}^N \simeq (\mathbb{P}^N)^r$. By Kambyashi's theorem [Kam], $(\mathbb{P}^N)^r$ has a G -linearized ample line bundle and this restricts to a G -equivariant ample line bundle on IY .

The projection $P: IY \rightarrow Y$, which takes f to $f(e)$ is G_0 -equivariant. If in addition Y is a G -space then the map $\Delta: Y \rightarrow IY$ which takes

$y \in Y$ to the function $f: G \rightarrow Y; g \mapsto g^{-1}y$ is a G -invariant section of P . In this case we denote the image of the section by Δ_Y .

We now return to the proof the proposition.

The proper G_0 -equivariant map $\pi: X_1 \rightarrow X$ induces a proper G -equivariant map $I\pi: IX_1 \rightarrow IX$. Let X' denote the inverse image of Δ_X under this map. Then the map $X' \rightarrow \Delta_X \simeq X$ is proper and G -equivariant. Since U_1 is a G -space, $\Delta_{U_1} \simeq U_1$ embeds in X' and maps isomorphically to $\Delta_U \simeq U$. It follows that $X' \rightarrow X$ is an isomorphism over the dense open set U . Finally, there is a G -equivariant ample line bundle on IX_1 , and this restricts to a G -equivariant ample bundle on X' . \square

To prove Proposition 6.5, we also need the following lemma, which is stated without proof in [PV].

Lemma 6.4. *Let G be an algebraic group acting on a normal quasi-projective variety Y . Then there exists a G -equivariant embedding $Y \subset \mathbb{P}^N$ such that the G -action is the restriction of the action of PGL_{N+1} on \mathbb{P}^N ; i.e., Y has a G -linearized ample line bundle.*

Proof of Lemma 6.4. If G is connected then our statement is [Sum, Theorem 1]. Otherwise let G_0 be the identity component of G , and let $Y \hookrightarrow \mathbb{P}^N$ be an G_0 -equivariant projective embedding. As in the proof of Proposition 6.3, since Y is a G -space we have $Y \simeq \Delta_Y \subset IY$. The composite embedding of Y into $I\mathbb{P}^N \simeq (\mathbb{P}^N)^r$ is G -equivariant. Since $(\mathbb{P}^N)^r$ is projective, it has a G -linearized ample line bundle by Kambayashi's theorem [Kam]. Pulling this ample bundle back to X gives us our desired G -linearized ample line bundle. \square

Proposition 6.5. *If X is a separated G -scheme, then there exists an equivariant Chow envelope $\tilde{X} \rightarrow X$ with \tilde{X} non-singular.*

Proof. The map $X_{red} \rightarrow X$ is an equivariant envelope, as is the normalization map. Thus we may assume that X is normal. We will show that there is a non-singular quasi-projective variety X' and a proper G -equivariant morphism $X' \xrightarrow{\pi} X$ such that π is an isomorphism over a dense G -equivariant open subspace U of X . This implies the proposition, since by Noetherian induction we may assume there is an equivariant Chow envelope \tilde{Y} of $Y = X - U$, so $\tilde{X} = X' \cup \tilde{Y} \rightarrow X$ is the desired envelope.

By Proposition 6.3 there is a quasi-projective scheme X_1 and a proper G -equivariant morphism $X_1 \rightarrow X$ which is an isomorphism over a dense open subspace of X . By canonical resolution of singularities there is a canonical (hence G -equivariant) resolution of singularities $X' \rightarrow X_1$.

Since X' is obtained by a sequence of blowups of the quasi-projective scheme X_1 , X' is quasi-projective. The composite map $X' \rightarrow X$ is an isomorphism over a dense open subspace of X . Finally, since X' is non-singular it has a G -linearized ample line bundle by Lemma 6.4. \square

6.1.2. *Envelopes and completions of equivariant K -theory.* A crucial property of (equivariant) envelopes is the following:

Lemma 6.6. *Let X be a G -space and $\pi: \tilde{X} \rightarrow X$ a birational envelope. Then $\pi_*: CH_G^*(\tilde{X}) \rightarrow CH_G^*(X)$ is surjective.*

Proof. This follows from [EG1, Lemma 3] and the corresponding fact for non-equivariant envelopes [Ful, Lemma 18.3(6)] \square

Using Riemann-Roch and this lemma, we can prove an analogous result on completions of equivariant Grothendieck groups:

Proposition 6.7. *Let $\tilde{X} \xrightarrow{\pi} X$ be a G -equivariant envelope. Then for any $h \in Z(G)$, the proper pushforward $\pi_*: G_0(\widehat{G, \tilde{X}^h})_h \rightarrow G_0(\widehat{G, X^h})_h$ is surjective.*

Proof. By Theorem 5.1 and Proposition 5.4 the diagram

$$\begin{array}{ccc} \widehat{G_0(G, \tilde{X}^h)}_h & \xrightarrow{\tau^{G, \text{ot}_h}} & \prod_{n=0}^{\infty} CH_G^n(\tilde{X}^h) \\ \pi_* \downarrow & & \pi_* \downarrow \\ \widehat{G_0(G, X^h)}_h & \xrightarrow{\tau^{G, \text{ot}_h}} & \prod_{n=0}^{\infty} CH_G^n(X^h) \end{array}$$

commutes and the horizontal arrows are isomorphisms. By Lemma 6.6, π_* is surjective as a map of equivariant Chow groups. Hence π_* must also be surjective as a map of completed equivariant Grothendieck groups. \square

6.2. **Proofs of Theorem 4.3(c), Corollary 4.5(b) and Theorem 5.5(d).** Suppose that there is an assignment $j_X: G_0(\widehat{Z, X^h})_h \rightarrow G_0(\widehat{G, X})$ for every G -space X which is covariant for proper morphisms and satisfies equation (10) for smooth X . Equation (10) implies that

$$j_X \alpha = (i^!)^{-1} (\lambda_{-1}(N_i^*)^{-1} \cap \alpha) = i_! \alpha$$

when X is smooth. We wish to show that $j_X = i_!$ for arbitrary X . Let $\pi: \tilde{X} \rightarrow X$ be a G -equivariant envelope with \tilde{X} non-singular. By Proposition 6.7, $\pi_*: G_0(\widehat{Z, \tilde{X}^h})_h \rightarrow G_0(\widehat{Z, X^h})_h$ is surjective. By covariance, $j_X(\pi_* \alpha) = \pi_* j_{\tilde{X}} \alpha$. Since $j_{\tilde{X}} = i_!$ as maps $G_0(\widehat{Z, \tilde{X}^h})_h \rightarrow$

$\widehat{G_0(G, \tilde{X})}$, it follows from covariance of $i_!$ that $\pi_* j_{\tilde{X}} \alpha = i_! \pi_* \alpha$. Hence $j_X = i_!$ completing the proof of Theorem 4.3(c).

To prove part (b) of Corollary 4.5 we use a similar argument. First observe that since $i_!$ is an isomorphism which is covariant for proper morphisms, Proposition 6.7 implies that the pushforward $\pi_*: \widehat{G_0(G, \tilde{X})} \rightarrow \widehat{G_0(G, X)}$ is surjective. Consider, for each G -space X , the map $(i_*)^{-1} \circ \text{res}_h: \widehat{G_0(G, X)} \rightarrow \widehat{G_0(Z, X^h)}_h$. This map is covariant for proper morphisms. When X is smooth, equation (10) implies that $(i_!)^{-1}$ is the map

$$\alpha \mapsto \lambda_{-1}(N_i^*) \cap i^*(\text{res}_h \alpha) = (i_*)^{-1} \circ \text{res}_h(\alpha),$$

i.e., that $(i_*)^{-1} \circ \text{res}_h = (i_!)^{-1}$. Using the same argument as in the previous paragraph, we conclude that $(i_*)^{-1} \circ \text{res}_h = (i_!)^{-1}$ as maps $\widehat{G_0(G, X)} \rightarrow \widehat{G_0(Z, X^h)}_h$ for arbitrary X . Since i_* and $i_!$ are isomorphisms, it follows that $\text{res}_h: \widehat{G_0(G, X)} \rightarrow \widehat{G_0(Z, X)}_h$ is also an isomorphism.

We conclude with a proof of Theorem 5.5(d), the uniqueness of the functor τ^Ψ . If X is a G -scheme, then by Proposition 6.3 there is a non-singular equivariant Chow envelope $\tilde{X} \rightarrow X$. If there is a functor τ' such that τ' satisfies properties (a) and (c), then $\tau' = \tau^\Psi$ on the smooth quasi-projective scheme \tilde{X} . Arguing as above shows that $\tau' = \tau^\Psi$ for arbitrary X .

REFERENCES

- [BFM] Paul Baum, William Fulton, and Robert MacPherson, *Riemann-Roch for singular varieties*, Inst. Hautes Études Sci. Publ. Math. (1975), no. 45, 101–145.
- [BM] Edward Bierstone and Pierre D. Milman, *Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant*, Invent. Math. **128** (1997), no. 2, 207–302.
- [Bor] Armand Borel, *Linear algebraic groups*, second ed., Springer-Verlag, New York, 1991.
- [BtD] Theodor Bröcker and Tammo tom Dieck, *Representations of compact Lie groups*, Graduate Texts in Mathematics, vol. 98, Springer-Verlag, New York, 1995.
- [Bou] Nicolas Bourbaki, *Commutative algebra. Chapters 1–7*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 1989, Translated from the French, Reprint of the 1972 edition.
- [CG] Neil Chriss and Victor Ginzburg, *Representation theory and complex geometry*, Birkhäuser Boston Inc., Boston, MA, 1997.
- [EG1] Dan Edidin and William Graham, *Equivariant intersection theory*, Invent. Math. **131** (1998), no. 3, 595–634.

- [EG2] ———, *Riemann-Roch for equivariant Chow groups*, Duke Math. J. **102** (2000), no. 3, 567–594.
- [EG3] ———, *Nonabelian localization in equivariant K -theory and Riemann-Roch for quotients*, Adv. Math. **198** (2005), no. 2, 547–582.
- [Ful] William Fulton, *Intersection theory*, Springer-Verlag, Berlin, 1984.
- [Hel] Sigurdur Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Pure and Applied Mathematics, vol. 80, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1978.
- [Hum] James E. Humphreys, *Conjugacy classes in semisimple algebraic groups*, Mathematical Surveys and Monographs, vol. 43, American Mathematical Society, Providence, RI, 1995.
- [Kam] T. Kambayashi, *Projective representation of algebraic linear groups of transformations*, Amer. J. Math. **88** (1966), 199–205.
- [Knu] Donald Knutson, *Algebraic spaces*, Springer-Verlag, Berlin, 1971, Lecture Notes in Mathematics, Vol. 203.
- [Mer] Alexander S. Merkurjev, *Equivariant K -theory*, Handbook of K -theory. Vol. 1, 2, Springer, Berlin, 2005, pp. 925–954.
- [Nie] H. Andreas Nielsen, *Diagonalizably linearized coherent sheaves*, Bull. Soc. Math. France **102** (1974), 85–97.
- [OV] A. L. Onishchik and È. B. Vinberg, *Lie groups and algebraic groups*, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1990, Translated from the Russian and with a preface by D. A. Leites.
- [PV] V. L. Popov and È. B. Vinberg, *Invariant theory*, Algebraic geometry. IV (I. R. Shafarevich, ed.), Encyclopedia of Mathematical Sciences, vol. 55, Springer-Verlag, Berlin, 1994, pp. vi+284.
- [Seg] Graeme Segal, *Equivariant K -theory*, Inst. Hautes Études Sci. Publ. Math. (1968), no. 34, 129–151.
- [Ser] Jean-Pierre Serre, *Groupes de Grothendieck des schémas en groupes réductifs déployés*, Inst. Hautes Études Sci. Publ. Math. (1968), no. 34, 37–52.
- [Sum] Hideyasu Sumihiro, *Equivariant completion*, J. Math. Kyoto Univ. **14** (1974), 1–28.
- [Tho1] R. W. Thomason, *Algebraic K -theory of group scheme actions*, Algebraic topology and algebraic K -theory (Princeton, N.J., 1983), Ann. of Math. Stud., vol. 113, Princeton Univ. Press, Princeton, NJ, 1987, pp. 539–563.
- [Tho2] ———, *Une formule de Lefschetz en K -théorie équivariante algébrique*, Duke Math. J. **68** (1992), no. 3, 447–462.
- [Toe] B. Toen, *Théorèmes de Riemann-Roch pour les champs de Deligne-Mumford*, K -Theory **18** (1999), no. 1, 33–76.
- [VV] Gabriele Vezzosi and Angelo Vistoli, *Higher algebraic K -theory of group actions with finite stabilizers*, Duke Math. J. **113** (2002), no. 1, 1–55.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211

E-mail address: edidin@math.missouri.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, BOYD GRADUATE STUDIES RESEARCH CENTER, ATHENS, GA 30602

E-mail address: wag@math.uga.edu