Matrix Norms and Rapid Mixing for Spin Systems

M artin D yer School of C om puting U niversity of Læds Læds LS2 9JT, UK Leslie Ann Goldberg Department of Computer Science, University of Liverpool, Liverpool L69 3BX, UK

M ark Jerrum SchoolofM athem atical Sciences, Queen M ary, University of London M ile End Road, London E1 4N S, UK

21st February 2007

A bstract

We give a system atic development of the application of matrix norms to rapid mixing in spin systems. We show that rapid mixing of both random update G lauber dynamics and system atic scan G lauber dynamics occurs if any matrix norm of the associated dependency matrix is less than 1. We give improved analysis for the case in which the diagonal of the dependency matrix is 0 (as in heat bath dynamics). We apply the matrix norm methods to random update and system atic scan G lauber dynamics for colouring various classes of graphs. We give a general method for estimating a norm of a symmetric non-regular matrix. This leads to improved mixing times for any class of graphs which is hereditary and su ciently sparse including several classes of degree-bounded graphs such as non-regular graphs, trees, planar graphs and graphs with given tree-width and genus.

1 Introduction

A spin system consists of a nite set of sites and a nite set of spins. A con guration is an assignment of a spin to each site. Sites interact locally, and these interactions specify the relative likelihood of possible (local) subcon gurations. Taken together, these give a well-de ned probability distribution on the set of con gurations.

G lauber dynamics is a M arkov chain whose states are con gurations. In the transitions of the M arkov chain, the spins are updated one at a time. The M arkov chain converges to the stationary distribution . During each transition of random update G lauber dynamics, a site is chosen uniform ly at random and a new spin is chosen from an appropriate probability distribution (based on the local subcon guration around the chosen site). During a transition of system atic scan G lauber dynamics, the sites are updated in a (determ inistic) system atic order, one after another. Again, the updates are from an appropriate probability distribution guration.

It is well-known that the mixing times of random update G lauber dynamics and system atic scan G lauber dynamics can be bounded in terms of the in uences of sites on each other. A dependency matrix for a spin system with n sites is an n n matrix R in which $R_{i;j}$ is an upper bound on the in uence (de ned below) of site i on site j.

An easy application of the path coupling m ethod of Bubley and D yer shows that if the L_1 norm of R (which is its maximum row sum, and is written $kR k_1$) is less than 1 then random update G lauber dynamics is rapidly mixing. The same is true if the L_1 norm (the maximum column sum of R, written $kR k_1$) is less than 1. The latter condition is known as the D obrushin condition. D obrushin [7] showed that, if $kR k_1 < 1$, then the corresponding countable spin system has a unique G ibbs measure. As we now know (see W eitz [27]), there is a very close connection between rapid mixing of G lauber dynamics for nite spin system s, and uniqueness of G ibbs measure for the corresponding countable system s. D obrushin and Shlosm an [8] were the rst to establish uniqueness when $kR k_1 < 1$. Their analysis extends to block dynamics but we will stick to G lauber dynamics in this paper. For an extension of som e of our ideas to block dynamics, see [21].

The D obrushin condition kR $k_1 < 1$ implies that system atic scan is rapidly mixing. A proof follows easily from the account of D obrushin uniqueness in Sim on's book [24], some of which is derived from the account of Follm er [13]. In [10], we showed that kR $k_1 < 1$ also implies rapid mixing of system atic scan G lauber dynam ics. [10, Section 3.5] notes that it is possible to prove rapid mixing by observing a contraction in other norm s besides the L₁ norm and the L₁ norm. This idea was developed by H ayes [16], who showed that rapid mixing occurs when the spectral norm kR k_2 is less than one. For symmetric matrices, the spectral norm is equal to the largest eigenvalue of R, (R). So, for symmetric matrices, [16] gives rapid mixing when (R) < 1. In general, kR₂k= (R) can be arbitrarily large, see Section 2.1.

In this paper, we give a system atic development of the application of matrix norm s to rapid mixing. We rst show that rapid mixing of random update G lauber dynamics occurs if any matrix norm is less than 1. Formally, we prove the following, where J_n is the norm of the all 1's matrix. All de nitions are given in Section 2.

Lem m a 1. Let R be a dependency matrix for a spin system, and let k = k be any matrix norm such that $kRk = \langle 1. Then the mixing time of random update G lauber dynamics is bounded by$

$$r'_{r}$$
 (") n(1) 1 ln (1) 1 J_n=":

W e prove a sim ilar result for system atic scan G lauber dynam ics.

Lem m a 2. Let R be a dependency matrix for a spin system and k k any matrix norm such that kRk < 1. Then the mixing time of system atic scan G lauber dynamics is bounded by

$$f_{s}(")$$
 (1)¹ ln (1)¹ J_n=":

The proofs of Lemma 1 and 2 in Section 3.1 use path coupling. Despite historical di erences, this approach is essentially equivalent to Dobrushin uniqueness, and we show how to prove the same lemmas using Dobrushin uniqueness in Section 3.2. We also give an improved analysis for the case in which the diagonal of R is 0, which is the case for heat bath dynamics. We prove the following.

Lem m a 3. Let R be symmetric with zero diagonal and $kR k_2 = (R) = < 1$. Then the mixing time of systematic scan is at most

$$_{s}^{(")}$$
 (1 $\frac{1}{2}$)(1)¹ ln (1)¹ n=":

An interesting observation is that when (R) is close to 1, the number of G lauber steps given in the upper bound from Lemma 3 is close to half the number that we get in our best estimate for random update G lauber dynamics (see Remark 6) | perhaps this can be interpreted as weak evidence in support of the conjecture that systematic scan mixes faster than random update for G lauber dynamics.

1.1 Applications

Hayes [16] gives applications of conditions of the D obrushin type to problem s on graphs, using the norm $k \ k$ In [10], we observed that the dependency matrix for the G lauber dynam ics on graph coburings can be bounded by a multiple of the adjacency matrix of the graph. This was applied to analysing the systematic scan dynam ics for colouring near-regular graphs and hence to regular graphs. Hayes extends the observation of [10] to the G lauber dynam ics for the Ising and hard core models. He applies these observations with a new estimate of the mixing time of these chains on planar graphs with given maximum degree. He also applies them to bounded-degree trees, using an eigenvalue estimate due to Stevanovic [25], for which he provides a di erent proof. He extends these results to the systematic scan chain for each problem, using ideas taken from [10].

In Section 4, we apply the matrix norm methods developed here to the random update G lauber dynamics and system atic scan dynamics for colouring various classes of graphs. We give a general method for estimating the norm k_{2} kof a symmetric non-negative matrix R.Ourmethod is again based on matrix norms. We show that there exists a \decom position" $R = B + B^{T}$, for some matrix B, where kB k₁, kB k₁ can be bounded in terms of kR k₁ and the maximum density of R. The bounds on kB k₁, kB k₁ can then be combined to bound kR k₂. In particular, our methods allow us to give a common generalisation of results of Hayes [16], Stevanovic [25] and others for the maximum eigenvalue of certain graphs. In most cases we are also able to strengthen the previous results. We show that k _kcan be bounded analogously to [16, 25] for any class of graphs which is hereditary and su ciently sparse. We apply this to give in proved mixing results for the G lauber and system atic scan dynamics for colouring several classes of degree-bounded graph: near-regular graphs, trees, planar graphs and, more generally, graphs of given tree-width or genus. These examples do not exhaust the classes of graphs to which our methods apply. Our methods could be used equally well to provide new mixing results for the Ising and hard-core models, as is done in [16], but we do not explore other applications here.

2 Prelim inaries

Let $[n] = f1;2;:::;ng, N = f1;2;3;:::g, and N_0 = N$ [f0g. W e use Z;R for the integers and reals, and R₊ for the non-negative reals. Let jcjdenote the absolute value of c.

2.1 Matrix norms

Let $M_{mn} = R^{m-n}$ be the set of realm n matrices. We denote M_{nn} , the set of square matrices, by M_n . The set of non-negative matrices will be denoted by M_{mn}^+ , and the set of square non-negative matrices by M_n^+ . We will write 0 for the zero m n matrix and I for the n n identity matrix. The dimensions of these matrices can usually be inferred from the context, but where am biguity is possible, or emphasis required, we will write $0_{m,n}$, I_n etc. Whether vectors are row or column will be determined either by context or explicit statement. The ith component of a vector v will be written both as v_i and v(i), whichever is more convenient. If R is a matrix and v a vector, R v(i) will mean $(R v)_i$. We will use J for the n n matrix of 1's, 1 for the column n-vector of 1's, and 1^T for the row n-vector of 1's. Again the dimensions can be inferred from context.

- A matrix norm (see [17]) is a function k k: M_n ! R₊ for each m; n 2 N such that
 - (i) kR k = 0 and $R 2 M_{mn}$ if and only if $R = 0 2 M_{mn}$;
 - (ii) k R k = j j k R k for all 2 R and R 2 M m n;
 - (iii) kR + Sk kR k + kSk for all R; S 2 M_{m n};
 - (iv) kRSk kRkkSk for all R 2 M_{mk} , S 2 M kn (k 2 N).

Note that property (iv) (submultiplicativity) is sometimes not required for a matrix norm, but we will require it here. The condition that k k be de ned for all m; n is, in fact, a m ild requirement. Suppose k k is initially de ned only on M_A , for any large enough n, then we can de ne kR k for R 2 M_{mk} (m; k 2 [n]) by \embedding" R in M_n, i.e.

$$kR k \stackrel{\text{def}}{=} \left\| \begin{array}{c|c} & & & \\ \hline R & O_{m,m-k} \\ \hline & O_{n-m,k} & O_{n-m,m-k} \end{array} \right\| :$$

It is straightforward to check that this de nition gives the required properties. For m any m atrix norm s, this embedding norm coincides with the actual norm for all m; k 2 [n].

Examples of matrix norms are operator norms, dened by $kRk = max_{x \in 0} jRx j = jk j$ for any vector norm jkjdened on R^n for all n 2 N. Observe that we denote a matrix norm by k k and a vector norm by j j. Since vector norms occur only in this section, this should not cause confusion. In fact, their meanings will also be very close, as we discuss below.

For any operator norm we clearly have kIk = 1. The norm $k_1k_2k_3$ and k_1k_4 , are in portant examples, derived from the corresponding vector norms. The norm kRk_1 is the maximum column sum of R, kRk_1 is the maximum row sum, and the spectral norm $kRk_2 = {p - \atop p}$, where is the largest eigenvalue of R^TR . (See [17, pp.294 [5], but observe that jj jj is used for what we denote here by k k.) The Frobenius norm $kR_Fk = {p - \atop ij} R_{ij}^2$ (see [17, p.291]) is an example of a matrix norm which is not an operator norm. Note that $kIk = \frac{p}{n}$ for the Frobenius norm, so it cannot be dened as an operator norm.

New matrix norms can also be created easily from existing ones. If W_n 2 M_n is a xed nonsingular matrix for each n, then k $_{W}k = kW_{m}$ $W_{n}^{-1}k$ is a matrix norm. (See [17, p.296].) Note that k $_{W}k$ is an operator norm whenever k k is, since it is induced by the vector norm \mathbf{j}_{M} \mathbf{j}_{n} .

The following relate matrix norms to absolute values and corresponding vector norms.

Lemma 4. Suppose c2R. Letk kbe a matrix norm on 1 1 matrices. Then jcj kck.

Proof. This follows from the axioms for a matrix norm. First, kck = kc 1k = jcklk by (ii). A lso, kck = kc 1k kck klk by (iv). Finally, $klk \in 0$ by (i).

Lem m a 5. Suppose x is a column vector, j = j a vector norm, and k k the corresponding operator norm. Then j k j = j l j k x k.

Proof. Let x be a length-'column vector. $j_k j$ is the vector norm applied to x, $j_l j$ is the same norm applied to the length-1 column vector containing a single 1. kxk is the operator norm applied to the '1 matrix containing the single column x. Then kxk = $\max_{0} j_k j = j$ jwhere is a non-negative real number. Pulling constants out of the vector norm, $\max_{0} j_k j = j j = j_k j = j_l j$.

The dual (or adjoint [17, p.309]) nom k kofa matrix nom k kwillbe de ned by kR k= kR^T k. Thusk 1kand k 1k are dual, and k 2kis self-dual. Note that, for any colum n vector x, kx^T k = kxk so, for example, kx^T k₁ = kxk₁. C learly, any matrix nom k k induces a vector norm j jon colum n vectors. Then the dual matrix nom, as de ned here, is closely related to the dual vector norm.

Lem m a 6. Suppose x is a column vector, j = j l j kxk.

Proof. By de nition, $jlj = m ax_{\neq 0} j \neq j j = 1 = jlj$, after pulling out constants, and

$$\mathbf{j}\mathbf{l}\mathbf{j}\mathbf{k}\mathbf{j} = \mathbf{j}\mathbf{l}\mathbf{j}\mathbf{m} \underset{y \in 0}{\text{max}} \frac{\mathbf{\dot{x}}^{\mathrm{T}} \mathbf{y}\mathbf{j}}{\mathbf{j}\mathbf{y}\mathbf{j}} = \max_{y \in 0} \frac{\mathbf{\dot{x}}^{\mathrm{T}} \mathbf{y}\mathbf{j}\mathbf{j}\mathbf{l}\mathbf{j}}{\mathbf{j}\mathbf{y}\mathbf{j}} = \max_{y \in 0} \frac{\mathbf{j}\mathbf{x}^{\mathrm{T}} \mathbf{y}\mathbf{j}}{\mathbf{j}\mathbf{y}\mathbf{j}} = \mathbf{k}\mathbf{x}^{\mathrm{T}}\mathbf{k} = \mathbf{k}\mathbf{x}\mathbf{k} : \square$$

W ith any matrix $R = (R_{ij}) 2 M_n$ we can associate a weighted digraph G (R) with vertex set [n], edge set $E = (i; j) 2 [n]^2 : R_{ij} \in 0$, and (i; j) 2 E has weight R_{ij} . The (zero-one) adjacency matrix of G (R) will be denoted by A (R). If G (R) is labelled so that each component has consecutive numbers, then R is block diagonal, and the (principal) blocks correspond to the components of G (R). A block is irreducible if the corresponding component of G (R) is strongly connected. Note, in particular, that R is irreducible if R > 0. If R is symmetric, G (R) is an undirected graph, and R is irreducible when G (R) is connected. For i; j 2 V, d(i; j) will denote the number of edges in a shortest directed path from i to j. If there is no such path, d(i; j) = 1. The diameter of G, D (G) = max_{i;j2V} d(i; j). Thus G is strongly connected when D (G) < 1. For R 2 M $_n^+$, let (R) denote the largest eigenvalue (the spectral radius). We know that (R) 2 R $_+$ from Perron-Frobenius theory [23, Ch.1]. We use the following facts about (R). The rst is a restatement of [23, Thm .1.6], a version of the Perron-Frobenius Theorem.

Lem m a 7. If R 2 M⁺_n is irreducible, there exists a row vector w > 0 satisfying wR w if and only if (R). If = (R), then w is the unique left eigenvector of R for the eigenvalue . Lem m a 8. If R 2 M⁺_n has blocks R₁, R₂, ..., R_k, then (R) = m ax₁ i k (R_i). Lem m a 9 (See [23], Thm .1.1). If R; R⁰2 M⁺_n and R R⁰, then (R) (R⁰).

() is not a matrix norm. For example,

$$\begin{array}{ccc}
0 & 1 \\
0 & 0
\end{array} = 0$$

so axiom (i) in the denition of a matrix norm is violated by (). Nevertheless, (R) is a lower bound on the value of any norm of R.

Lem m a 10 (See [17], Thm .5.6.9). If R 2 M $_{n}^{+}$, then (R) kRk for any matrix norm k k.

Furtherm ore, for every R 2 M $_n^+$ there is a norm k k, depending on R, such that the value of this norm coincides with () when evaluated at R.

Lem m a 11. For any irreducible R 2 M $_{n}^{+}$, there exists a matrix norm k k such that (R) = kR k.

Proof. Let w > 0 be a left eigenvector for = (R), and let $W = \text{diag}(w) 2 M_n^+$. Then $k_w k = kW$ () $W^{-1}k_1$ is the required norm, since $kR k_w = kW R W^{-1}k_1 = kw R W^{-1}k_1 = kw W^{-1}k_1 = k1^T k_1 = k1k_1 = .$

The norm $k = \sqrt{k} de$ ned in the proof of Lemma 11 is the minimum matrix norm for R, but this norm is clearly dependent upon R since w is.

The numerical radius [17] of R 2 M $_{n}^{+}$ is dened as (R) = maxfx^T R x : x^T x = 1g. () is not submultiplicative since

 $\begin{array}{cccccccc} 0 & 1 \\ 0 & 0 \end{array} = \begin{array}{cccccccc} 0 & 0 \\ 1 & 0 \end{array} = \frac{1}{2};$

but applying to the product gives

 $\begin{array}{cccc}
1 & 0 \\
0 & 0
\end{array} = 1:$

Thus () is not a matrix norm in our sense. Nevertheless, (R) provides a lower bound on the norm $kR\,k_2$.

Lem m a 12. (R) (R) kR_{k} , with equality throughout if R is symmetric.

Proof. Let w, w ith $kwk_2 = 1$, be an eigenvector for = (R). Then (R) $w^T R w = w^T w =$ (R). Also (R) = $x^T R x$ kR kg for some x w ith $kxk_2 = 1$, and $x^T R x = kx^T R xk_2$ kR kg since k gk is submultiplicative. If R is a symmetric matrix, then $R = Q^T Q$, for Q orthonormal and a diagonal matrix of eigenvalues. Then $kRk_2^2 = (R^T R) = (2^{-1}) = (R)^2$.

Thus, when R is symmetric we have $(R) = kRk_2$, and hence k_2k is the minimum matrix norm, uniform by for all symmetric R. However, when R is not symmetric, $kRk_2 = (R)$ can be arbitrarily large, even though 0 < R < J. Consider, for example,

$$R = "1 2"$$

for any $0 < " < \frac{1}{2}$. Then (R) $< p_{"}$ + ", and $kRk_2 > 1$ 2", so $\lim_{n!} kRk_2 = (R) = 1$. Also k $_2k$ is not necessarily the minimum norm for asymmetric R. We always have kRk_2 [17, p.314], but this bound can almost be achieved for 0 < R < J. Consider

for any $0 < " < \frac{1}{n}$. Then $kRk_1 = 1$ ", but $kRk_2 > (1 n")^{p}\overline{n}$, so $\lim_{n \to \infty} w_{n+1} = kRk_1 = p\overline{n}$. On the other hand, k _2kdoes have the follow ing m in in ality property.

Lem m a 13. For any matrix norm k k, $kR_2k = \frac{p}{kR kkR k}$.

Proof. $kRk_2^2 = (R^T R) kR^T R k kR^T kkR k = kR kkR k$, using Lem m as 10 and 12.

For a matrix norm k k, the quantities $_{n}J = kJk$, for $J \ge M_{n}$, and $C_{n} = k1kk1k$, will be used below. We collect some of its properties here. In particular, $J_{n} = n$ for k $_{1}k$, k $_{2}k$, k $_{1}k$ and the Frobenius norm, by direct calculation. M ore generally,

Lem m a 14. Let k k be a matrix norm . Then

(i) if
$$J_n = kJk$$
, then $J_n = J_n$;

- (ii) n J_h C_n ;
- (iii) if k = k is an operator norm, then $_{n}J = C_{n}$.
- (iv) if k k is induced by a vector norm which is symmetric in the coordinates, then $_n J = n$;
- (v) if k_{p} k is induced by the vector p-norm (1 p 1), then J = n.
- (vi) if $k = kW = W^{-1}k_1$, where W = diag(w) for a column vector w > 0 with $kw k_1 = 1$, then $J_n = 1 = w_{\min}$, where $w_{\min} = \min_i w_i$.

Proof. W e have

(i) $J_n = kJk = kJ^T k = kJk = J_n$;

- (ii) $J = 11^T$ so n1 = J1. Thus nk1k kJkk1k. Now k1k \notin 0, so cancellation gives the rst inequality. The second follows by submultiplicativity and duality.
- (iii) $J_n = kJk = k11^T k = max_{x \in 0} j l 1^T x j j x j$ where x is a length-n vector. Pulling scalar multiples out of the vector norm in the numerator, this is equal to j l j max_{x \in 0} j T x j j x j. Now, by Lemma 5, j l j = j l j k1k, and hence $J_n = k1kmax_{x \in 0} j T x j j x j = k1kk1^T k = C_n$.
- (iv) Let x be any column vector such that $1^{T} x = n$. Let x be x after a coordinate permutation , and x = x = n!. Clearly x = 1. Also kxk kxk, and $1^{T} x = 1^{T} x = n$ by subadditivity of k k and symmetry, so klk= $\max_{x \in 0} 1^{T} x = kxk$ $\max_{x \in 0} 1^{T} x = kxk = n = klk$.
- (v) This follows directly from (iv).

(vi)
$$J_n = kJk_w = kZk_1$$
, where $Z_{ij} = w_i = w_j$. Thus $J_n = \prod_{i=1}^{P} w_i = m in_i w_i = 1 = w_{min}$.

Remark 1. For an arbitrary matrix norm we can have $C_n > J_n$. This is true even if the norm is invariant under row and column permutations. For example k $k = maxfk_1$; k $_1kg$ is a matrix norm, with kJk = k1k = k1k = n, which even satisfies kIk = 1 (see [17, p.308]). For this norm $C_n = J_n = n$. In general, the ratio is unbounded, even for a xed n. Consider, for example, k k = maxfkW W^1k_1 ; kW W^1k_1 g, where W = diag(v) for a column vector v > 0 with $kvk_1 = 1$. It is easy to show that this is a matrix norm with $C_n = J_n = max_iv_i = min_iv_i$, which can be arbitrarily large.

W e will use the following technical Lemma, which appears as Lemma 9 in [10] for the norm k $_{1}k$. W e show that, for any non-negative matrix R with kRk < 1, there is a row vector w which approximately satisfies the condition of Lemma 7, and has w_{min} not too small.

Lem m a 15. Let R 2 M $_{n}^{+}$, and let k k be a matrix norm such that kR k < 1. Then, for any 0 < < 1, there is a matrix R⁰ R and a row vector w > 0 such that w R⁰ $_{w}^{0}$, kw k₁ = 1 and w_{min} = m in_iw_i = J, where 0 = + < 1.

Proof. Let $J^0 = J = J_n$, and $R^0 = R + J^0$. Then R^0 is irreducible, and kR^0k kRk + . Then, by Lem ma 10, $(R^0) + = 0$. Thus, by Lem ma 7, there exists w > 0 such that $wR^0 = \sqrt[6]{w}$. We normalise so that $kwk_1 = 1$. Then $w = \sqrt[6]{w} = wR^0 = \sqrt[6]{r} = J_n$; and hence $w_{min} = J_n$.

2.2 Random update and system atic scan G lauber dynam ics

The fram ework and notation is from [10, 11]. The set of sites of the spin system will be V = [n] = f1;2;:::;ng, and the set of spins will be = [q]. A con guration (or state) is an assignment of a spin to each site, and $^{+} = ^{n}$ denotes the set of all such con gurations. Let $M = q^{n} = jj^{n} = j^{+}j$, and we will suppose $^{+} = [M]$.

Local interaction between sites speci es the relative likelihood of possible (local) sub-con gurations. Taken together, these give a well-de ned probability distribution on the set of con gurations $^+$. G lauber dynamics is a M arkov chain (x_t) on con gurations that updates spins one site at a time, and converges to . We measure the convergence of this chain by the total variation distance

 $d_{TV}(;)$. We will abuse notation to write, for exam $pl_{FV}d(x_t;)$ rather than $d_{TV}(L(x_t);)$. The mixing time (") is then de ned by (") = mix_f d_{TV}(x_t;) "g. In our setting, n measures the size of of con gurations in +, and we presume it to be large. Thus, for convenience, we also use asymptotic bounds $^(")$, which have the property that $\lim_{t \to 0} sup_{n!,1}$ (")= $^(")$ 1.

We use the following notation. If x is a conguration and j is a site then x_j denotes the spin at site j in x. For each site j, S_j denotes the set of pairs of congurations that agree o of site j. That is, S_j is the set of pairs $(x;y) 2^{+} + x_j$ such that, for all $i \notin j$, $x_i = y_i$. For any state x and spin c, we use $x \rightarrow jc$ for the state y such that $y_i = x_i$ ($i \notin j$) and $y_j = c$. For each site j, we have a transition matrix P^[j] on the state space + which satis es two properties:

- (i) P ^[j] changes one con guration to another by updating only the spin at site j. That is, if P ^[j](x;y) > 0 then (x;y) 2 S_j.
- (ii) The equilibrium distribution is invariant with respect to $P^{[j]}$. That is, $P^{[j]} =$

Random update G lauber dynamics corresponds to a Markov chain M with state space ⁺ and transition matrix P = $(1=n)^{P} {n \atop j=1} P^{[j]}$. Systematic scan G lauber dynamics corresponds to a Markov chain M \rightarrow with state space ⁺ and transition matrix $P_{\rightarrow} = {Q \atop j=1}^{n} P^{[j]}$.

It is well-known that the mixing times r(") of M and s(") of M \rightarrow can be bounded in terms of the in uences of sites on each other. To be more precise, let j(x;) be the distribution on spins at site j induced by P ^[j](x;). Thus $j(x;c) = P^{[j]}(x;x \rightarrow jc)$. Now let \mathfrak{F}_{ij} be the in uence of site i on site j, which is given by $\mathfrak{F}_{ij} = \max_{(x,y) \geq S_i} d_{TV} (j(x;); j(y;))$. A dependency matrix for the spin system is any n n matrix $R = (\mathfrak{F}_{ij})$ such that $\mathfrak{F}_{ij} = \mathfrak{F}_{ij}$. Clearly we may assume $\mathfrak{F}_{ij} = 1$.

G iven a dependency matrix R, let $*_j$ denote the jth column of R, for j 2 [n]. Now let R_j 2 M $_n^+$ be the matrix which is an identity except for column j, which is $*_j$, i.e.

$$(R_{j})_{ik} = \begin{cases} 8 \\ < 1; & \text{if } i = k \notin j; \\ \$_{ij}; & \text{if } k = j; \\ 0; & \text{otherw ise.} \end{cases}$$
(1)

Let $R = \frac{1}{n} \prod_{j=1}^{P} R_j = \frac{n-1}{n} I + \frac{1}{n} R$ de nether random update matrix for R, and let $\mathcal{R} = R_1 R_2 ::: R_n$ de nether scan update matrix for R.

3 M ixing conditions for G lauber dynam ics

There are two approaches to proving mixing results based on the dependency matrix, path coupling and D obrushin uniqueness. These are, in a certain sense, dual to each other. All the results given here can be derived equally well using either approach, as we will show.

3.1 Path coupling

F irst consider applying path coupling to the random update G lauber dynam ics. W e will begin by proving a simple property of R $\,$.

Lem m a 16. Let R be a dependency matrix for a spin system, and k k any operator norm such that kR k < 1. Then kR k where = $1 \frac{1}{n} (1) < 1$.

Proof. kR k
$$\frac{n-1}{n}$$
 kIk + $\frac{1}{n}$ kR k = $\frac{n-1}{n}$ + $\frac{1}{n}$ kR k 1 (1) = n = .

W e can use this to bound the mixing time of the random update G lauber dynam ics.

Lem m a 17. Suppose R is a dependency matrix for a spin system, and let k k be any matrix norm. If kR k < 1, then the mixing time $_{r}$ (") of random update G lauber dynamics is at most n (1) 1 ln (C $_{n}$ =").

Proof. We will use path coupling. See, for example, [12]. Let $x_0; y_0 2^{-+}$ be the initial congurations of the coupled chains, and $x_t; y_t$ the states after t steps of the coupling. The path $z_0; \ldots; z_n$ from x_t to y_t has states $z_0 = x_t$, and $z_i = (z_{i-1} \rightarrow^i y_t(i))$ (i2 [n]), so $z_n = y_t$. We use a path metric d (;) determined by $(x_i; y) = _i$ if $(x; y) 2 S_i$, for given constants $0 < _i = 1$ (i2 [n]). The $_i$ (i2 [n]) make up a column vector > 0. Note that d_i (;) is the usual Hamming distance.

The coupling will be to make the same vertex choice for all $(x_t; y_t) \ge S_i$ and then maximally couple the spin choices. W ith this coupling, \hat{s}_{ij} bounds the probability of creating a disagreement at site j for any $(x_t; y_t) \ge S_i$ and time t.

Let $_{t}(i) = Pr(x_{t}(i) \in y_{t}(i))$ determine a row vector $_{t}$, so $E[d(x_{t};y_{t})] = _{t}$. Clearly 0 $_{t}$ 1^T. Since $_{t}(i)$ and $Pr(x_{t+1} = x; y_{t+1} = y jx_{t};y_{t})$ are independent, it follows that

(The ith term in the sum comes from considering how the distance between $z_{i 1}$ and z_i changes under the coupling. A ssum ing $z_{i 1}$ and z_i di er (at site i) then $_i$ is the reduction in distance that comes about by updating site i and removing the disagreement there, while $_j$ %_{ij} is the expected increase in distance that arises when site j is updated and a disagreement is created there.) Now Equation (2) holds for all with $0 < _i$ 1. In particular, for any ", it holds for any vector in which one component is 1 and the other components are ". Taking the limit, as "! 0, we nd that, componentwise,

$$t+1$$
 tR : (3)

Now, using (3) and induction on t, we nd that

$$t+1 0 R t+1$$
: (4)

Equation (4) implies $t+1 = 0R^{t+1}$. Using the Coupling Lemma [9, 20],

$$d_{TV}(x_t; y_t) \quad \Pr(x_t \in y_t) \qquad \begin{array}{c} X^n \\ \Pr(x_t(i) \in y_t(i)) = t^1 \\ i = 1 \end{array} \quad 0^{R^{t}} 1 \qquad 1^{T} R^{t} 1:$$

Now applying Lemma 4 with $c = 1^{T} R^{t} 1$ and using submultiplicativity (property (iv) of matrix norms),

$$1^{T}R$$
^t 1 k1kkR k^tk $1^{T}k = C_{n}kR$ k^t:

But kR k = 1 (1)=n by Lem m al6. Thus, when t n (1)¹ ln ($C_n = "$),

$$d_{TV}(x_t; y_t)$$
 $C_n = C_n 1 (1) = n^t C_n e^{t(1) = n}$ ":

C orollary 18. Let R be a dependency matrix for a spin system. Then the mixing time r(") of random update G lauber dynamics is at most n (1) $^{1} \ln (n=")$ if R satis es any of the following:

- (i) the D obrushin condition $= kR k_1 < 1;$
- (ii) the D obrushin-Shlosm an condition $^{0} = kR k_{1} < 1$;
- (iii) a p-norm condition $kR\,k_p \qquad < 1$ for any 1 .

Proof. This follows directly from Lemma 17, since $C_n = J_n = n$ for these norms, by Lemma 14. \Box

C orollary 19. Let R be a dependency matrix for a spin system . Suppose either of the following conditions holds

- (i) w > 0 is a row vector such that w R w, $kw k_1 = 1$ and $w_{min} = m in_i w_i$,
- (ii) w > 0 is a column vector such that Rw = w, $kw k_i = 1$ and $w_{min} = m in_i w_i$.

Then the mixing time r(") of random update G lauber dynamics is at most n (1) 1 ln (1= w_{min} ").

Proof. Both are proved similarly, using Lemma 17 with a suitable operator norm, so $C_n = J_n$.

- (i) Let W = diag(w) de ne the norm $kR k_w = kW RW^{-1}k_1$. Then $kR k_w$, and $J_h = 1 = w_{min}$ by Lem m a 14.
- (ii) Let W = diag(w) dene the norm $kR k_w = kW^{-1}RW k_1 = kW R^TW^{-1}k_1$. Then $kR k_w$, and $J_n = 1 = w_{min}$ by Lemma 14.

In the setting of C orollary 19(i), we can also show contraction of the associated metric d_w (;). Lem m a 20. Suppose R is a dependency matrix for a spin system, and let w > 0 be a column vector such that Rw w. Then $E[d_w(x_{t+1};y_{t+1})] = E[d_w(x_t;y_t)]$ for all t 0.

Proof. Note that
$$\mathbb{R} = \frac{n}{n} \mathbb{W} + \frac{1}{n} \mathbb{R} \mathbb{W}$$
 $(\frac{n}{n} + \frac{1}{n}) \mathbb{W} = \mathbb{W}$. Putting $= \mathbb{W}$ in (\mathbb{Q}) ,
 $\mathbb{E} [d_{\mathbb{W}} (x_{t+1}; y_{t+1})] = t_{t+1} \mathbb{W}$ $t_{\mathbb{R}} \mathbb{W}$ $t_{\mathbb{W}} = \mathbb{E} [d_{\mathbb{W}} (x_t; y_t)]$:

Remark 2. We may be able to use Lemma 20 obtain a polynom ialm ixing time in the \equality case" = 1 of path coupling. However, it is dicult to give a general result other than in \soft core" systems, where all spins can be used to update all sites in every con guration. See [2] for details. We will not pursue this here, however. Note that mixing for the equality case apparently cannot be obtained from the D obrushin analysis of Section 3.2. This is perhaps the most signi cant di erence between the two approaches.




Figure 1: Exam ple 1

We would like to use an eigenvector in C orollary 19, since then = (R) kR k for any norm . An important observation is that we cannot necessarily do this, because R m ay not be irreducible (so $w_{m in}$ m ay be 0), or $w_{m in}$ m ay simply be too sm all.

Example 1. Consider the matrix of Figure 1. Here R is irreducible, with (R) = 0.4 and left eigenvector w such that $w_i / 2^i$ (i 2 [n]). Thus $w_{min} < w_n = w_1 = 2^{1 n}$ is exponentially small, and Corollary 19(i) would give a mixing time estimate of (n²) site updates. In fact, R satisfies the Dobrushin condition with = 0.8 and the Dobrushin-Shlosm an condition with ⁰ = 0.9, so we know mixing actually occurs in 0 (n logn) updates.

However, if we know kRk < 1 for any norm k = k, we can use Lem m 15 to create a better lower bound on $w_{m in}$. We apply this observation as follows.

C orollary 21. Let R be a dependency matrix for a spin system, and let k k be any matrix norm. Suppose kR k < 1. Then, for any 0 < < 1 , the mixing time of random update G lauber dynamics is bounded by

r(") n (1)¹ ln (J_n = "):

Proof. Choose 0 < < 1 . Let \mathbb{R}^0 be the matrix from Lem ma 15. Since \mathbb{R}^0 R, it is a dependency matrix for the spin system. Let w be the vector from Lem ma 15. Now by Corollary 19, the mixing time is bounded by r(") n $(1 \ ^0)^{-1} \ln (1=w_{\min}")$: where $w_{\min} = J_1$ and $^0 = +$.

From this we can now prove Lemma 1, which is a strengthening of Lemma 17 for an arbitrary norm.

Lem m a 1. Let R be a dependency m atrix for a spin system, and let k = k be any m atrix norm such that kRk = < 1. Then the m ixing time of random update G lauber dynam ics is bounded by

$$r'_{r}$$
(") n(1) 1 ln (1) 1 J_n=":

Proof. Choose = (1) = ln n. Substituting this into the mixing time from Corollary 21 now in plies the conclusion, since J_n n.

Remark 3. The mixing time estimate is $r(n) n(1)^{1} \ln (1)^{1} J_{n} = n$. If (1) is not too small, e.g. if $(1) = (\log^{k} n)$ for any constant k = 0, we have $r(n) n(1)^{1} \ln (J_{n} = n)$. Thus we lose little asymptotically using Lemma 1, which holds for an arbitrary matrix norm, from the mixing time estimate $r(n) = n(1)^{1} \ln (J_{n} = n)$, which results from applying C orollary 17 with an operator norm k = k. The condition $(1) = (\log^{k} n)$ holds, for example, when (1) is a small positive constant, which is the case in many applications.

We can easily extend the analysis above to deal with system atic scan. Here the mixing time $_{s}$ (") will be bounded as a num ber of complete scans. The num ber of individual G lauber steps is then n times this quantity. The following lemma modiles the proof technique of [10, Section 7].

Lem m a 22. Let R be a dependency matrix for a spin system, and k k any matrix norm. If \mathbb{R}^{k} < 1, the mixing time $_{s}$ (") of system atic scan G lauber dynamics is at most (1) $^{1} \ln (C_{n} = ")$. If k k is an operator norm, the mixing time is at most (1 $^{\frac{1}{2}} \ln (J_{n} = ")$.

Proof. We use the same notation and proof method as in Lemma 17. Consider an application of P^[j], with associated matrix R_{j} , as de ned in (1). Then it follows that

$$E [d(x_1;y_1)] \qquad X^n \\ 0 (i) (i + j *_{ij}) = 0 R_j ;$$

i=1

If, as before, i = 1 and j ! 0 for $j \in i$, we have $Pr(x_1(i) \in y_1(i)) = {}_{0}R_j$. Now it follows that $E[d(x_n; y_n)] = {}_{0}C_{i=1}^{n}R_j = {}_{0}C$ and $E[d(x_{nt}; y_{nt})] = {}_{0}C^{t}$. Thus $Pr(x_{nt}(i) \in y_{nt}(i)) = {}_{0}C^{t}(i)$. Hence

$$d_{TV}(\mathbf{x}_{nt}; \mathbf{y}_{nt}) \quad \Pr(\mathbf{x}_{nt} \notin \mathbf{y}_{nt}) \qquad \Pr(\mathbf{x}_{nt}(\mathbf{i}) \notin \mathbf{y}_{nt}(\mathbf{i})) \quad {}_{0}\mathbb{R}^{t}\mathbf{1} \quad \mathbf{1}^{T}\mathbb{R}^{t}\mathbf{1} \quad \mathbf{k}\mathbb{R}^{t}\mathbf{k}^{t}\mathbf{k}\mathbf{1}^{T}\mathbf{k}\mathbf{k}\mathbf{1}\mathbf{k}:$$

$$\stackrel{i=1}{=} \mathbf{1}$$

The remainder of the proof is now similar to Lemma 17.

The following lemma was proved, in a slightly different form, in [10, Lemma 11]. It establishes the key relationship between R and R.

Lem m a 23. Let R be a dependency matrix for a spin system. Suppose w > 0 is a row vector such that w R w for some 1. Then wR w.

Proof. Note that for any row vector z, $zR_i = [z_1 \ i \ 1 \ \mathbf{z} \otimes_i z_{i+1} \ n]$. Since wR w w, w \otimes_i w_i. Now we can show by induction on ithat w $R_1 ::: R_i$ [w $\otimes_i ::: w \otimes_i w_{i+1} ::: w_n$]. For the inductive step, w $R_1 ::: R_i$ $zR_i = [z_1 \ i \ 1 \ \mathbf{z} \otimes_i z_{i+1} \ n]$ where $z = [w \otimes_1 ::: w \otimes_i \ 1 \ w_i ::: w_n]$. But then z w, so $z \otimes_i$ w \otimes_i so zR_i [w $\otimes_i ::: w \otimes_i w_{i+1} ::: w_n$]. Taking i = n, we have w \mathbb{R} [w \otimes_1 n] w $\otimes R$ w.

Corollary 24. (R) (R) and, if $kRk_1 = 1$, $kRk_1 = kRk_1$.

Proof. The rst statement follows directly from Lemmas 7 and 23. For the second, note that $1^{T}R \quad kR k_{1}1^{T}$, so $1^{T}R \quad kR k_{1}1^{T}$ by Lemma 23. But this implies $kR k_{1} \quad kR k_{1}$.

We can now apply this to the mixing of systematic scan. First we show, as in [24], that the Dobrushin criterion implies rapid mixing.

C orollary 25. Let R be a dependency matrix for a spin system. Then, if R satis es the D obrushin condition = kRk_1 < 1, the mixing time of system atic scan G lauber dynamics is at most $(1)^{1} \ln(n=")$.

Proof. This follows from Lemma 22 and Corollary 24, since $J_n = n$ for the norm k_{1k} .

Next we show, as in [10, Section 3.3], that a weighted Dobrushin criterion im plies rapid mixing.

C orollary 26. Let R be a dependency matrix for a spin system. Suppose w > 0 is a row vector satisfying $kw k_1 = 1$ and w R w for some < 1. Let $w_{m in} = m in_i w_i$. Then the mixing time s (") of system atic scan G lauber dynam ics is bounded by $(1)^{1} \ln (1 = w_{m in})$.

Proof. By Lemma 23, w \hat{R} w. We use the norm $k_w k = k W$ W¹ k_1 , where W = diag(w). Then apply Lemma 22 with $k \hat{R}' k_w$.

Once again, we cannot necessarily apply C orollary 26 directly since w_{min} m ay be too sm all (or even 0). Applying C orollary 26 to E xam ple 1 would give a mixing time estimate of (n) scans. However, R satisfies the D obrushin condition with = 0.8 so we know mixing actually occurs in 0 (log n) scans. Once again, our solution is to perturb R using Lemma 15.

C orollary 27. Let R be a dependency matrix for a spin system and k $\,$ k a matrix norm such that $kR\,k\,$ $\,$ < 1. Then, for any 0 < $\,$ < 1 $\,$, the mixing time of systematic scan G lauber dynamics is bounded by

 $_{\rm s}$ (") (1)¹ ln (J_n = "):

Proof. Let \mathbb{R}^0 be the matrix and w the vector from Lemma 15. Since \mathbb{R}^0 R, it is a dependency matrix for the spin system. Now by Corollary 26, the mixing time satisfies $_{s}(")$ (1 0) $^{1} \ln (1=w_{\min}")$, where $w_{\min} = \min_{i} w_{i} = J$ and $^{0} = +$.

We can now use this to prove Lemma 2.

Lemma 2.

Let R be a dependency matrix for a spin system and k k any matrix norm such that kRk < 1. Then the mixing time of system atic scan G lauber dynamics is bounded by

$$_{s}(")$$
 (1)¹ ln (1)¹ J_n=":

Proof. This follows from Corollary 27 exactly as Lemma 1 follows from Corollary 21. \Box

Remark 4. If, for example, k $k = k_p$, kfor any $1 < p_1$, $J_h = n$, and we obtain a mixing time $\hat{s}(")$ (1)¹ ln (1)¹ n=". If, in addition, (1) = (log ^k n) for any k 0 (as in Remark 3), we have $\hat{s}(")$ (1)¹ ln (n="), which matches the bound from C orollary 25 for the norm k $_1k$. Note that there is a di erence from the random update case, since here we do not have a result like Lemma 17 which we can apply directly with any operator norm.

3.2 Dobrushin uniqueness

The natural view of path coupling in this setting corresponds to multiplying R on the left by a row vector , as in Lemma 17. The Dobrushin uniqueness approach corresponds to multiplying R on the right by a column vector . A swe showed in [0, Section 7], these two approaches are essentially equivalent. However, for historical reasons, the Dobrushin uniqueness approach is frequently used in the statistical physics literature. See, for example, [22, 24]. Therefore, for completeness, we will now describe the Dobrushin uniqueness fram ework, using the notation of [10].

Recall that $^{+} = M$]. For any column vector f 2 R^M, let $_{i}(f) = \max_{(x,y)2S_{i}} f(x)$ f(y)j. Let (f) be the column vector given by (f) = $(_{1}(f); _{2}(f); ...; _{n}(f))$. Thus $: \mathbb{R}^{M} ! \mathbb{R}^{n}$. The following Lemma gives the key property of this function.

Lem m a 28. ([10, Lem m a 10]) The function satisfies $(\mathbb{P}^{[j]}f) = \mathbb{R}_{j}$ (f).

Proof. Suppose $(x;y) \ge S_i \mod x \lim ses P^{[j]}f(x) = P^{[j]}f(y)$. Then

$$\begin{array}{rcl} {}_{i}(\mathbb{P}^{[j]}f) &=& \mathbb{P}^{[j]}f(x) & \mathbb{P}^{[j]}f(y) \\ &=& f(x \rightarrow^{j}c)\mathbb{P}^{[j]}(x;x \rightarrow^{j}c) & f(y \rightarrow^{j}c)\mathbb{P}^{[j]}(y;y \rightarrow^{j}c) \\ &=& x^{c} & x & c \\ &=& f(x \rightarrow^{j}c) \ _{j}(x;c) & f(y \rightarrow^{j}c) \ _{j}(y;c) \\ &=& x^{c} & x \\ &=& f(x \rightarrow^{j}c) & f(y \rightarrow^{j}c) \ _{j}(x;c) + & f(y \rightarrow^{j}c)(\ _{j}(x;c) \ _{j}(y;c)) \\ && x^{c} & x \\ &=& c \end{array}$$

W e will bound the two terms in the last expression separately. First

$$\begin{array}{c} X \\ f(x \rightarrow^{j} c) \quad f(y \rightarrow^{j} c) \quad j(x;c) \\ c \end{array} \qquad \begin{array}{c} \max_{c} f(x \rightarrow^{j} c) \quad f(y \rightarrow^{j} c) \\ \end{array} \qquad \begin{array}{c} 1_{i \neq j i}(f): \\ \end{array} \qquad (5)$$

For the second, let $f^+ = \max_c f(y \to^j c)$, $f = \min_c f(y \to^j c)$ and $f^0 = \frac{1}{2}(f^+ + f)$. Note that $f^+ = \frac{1}{2}(f^+ - f) = \frac{1}{2}(f^+ - f)$. Then, since $\int_{c} \int_{c} (x;c) \int_{c} (y;c) = 0$,

$$\begin{array}{ccc} X & f(y \rightarrow^{j} c) (j(x; c) & j(y; c)) &= & X & f(y \rightarrow^{j} c) & f^{0} (j(x; c) & j(y; c)) \\ c & & c & \\ & & 2d_{TV} (j(x;); j(y;)) & \max f(y \rightarrow^{j} c) & f^{0} \\ & & = & 2d_{TV} (j(x;); j(y;)) & f & f^{0} \end{array}$$

The conclusion now follows by adding (5) and (6).

The following lemma allows us to apply Lemma 28 to bound mixing times.

Lem m a 29. Let $M = (X_t)$ be a M arkov chain with transition m atrix P, and k k a m atrix norm. Suppose there is a matrix R such that, for any column vector f 2 R^M, (Pf) R (f), and kRk < 1. Then the mixing time of M is bounded by

(")
$$(1)^{1} \ln (C_{n} = ");$$

Proof. For a column vector f_0 , let f_t be the column vector $f_t = P^t f_0$. Let be the row vector corresponding to the stationary distribution of M. Note that $f_t = P^t f_0$, which is f_0 since is a left eigenvector of P with eigenvalue 1.

Now let f_0 be the indicator vector for an arbitrary subset A of M = +. That is, let $f_0(z) = 1$ if z 2 A and $f_0(z) = 0$ otherwise. Then, since $P^t(x;y) = Pr(X_t = y jX_0 = x)$, we have $f_t(x) = Pr(X_t 2 A jx_0 = x)$. Also $f_t = f_0 = (A)$ for all t. Let $f_t = m in_z f_t(z)$ and $f_t^+ = m ax_z f_t(z)$. Since is a probability distribution, $f_t = f_t f_t^+$, so $f_t = (A) f_t^+$.

By induction on t, using the condition in the statem ent of the lem ma, we have $(f_t) = R^t (f_0)$. But $R^t (f_0) = R^t (f_$

$$f_{t}^{+} \quad f_{t} = f_{t}(y) \quad f_{t}(x) \qquad \begin{array}{c} X^{n} & X^{n} \\ jf_{t}(z_{i}) & f_{t}(z_{i-1})j \\ & i = 1 \end{array} \quad (f_{t}) = 1^{T} \quad (f_{t}) \quad 1^{T} R^{t} 1:$$

This implies that $\max_{x} jPr(x_t 2 A jx_0 = x)$ (A) j $\hat{I}R^{t}1$. Since A is arbitrary, for all t $(1)^{1} \ln (C_n = ")$ we have

$$d_{TV}(x_t;)$$
 $1^T R^t 1 k R k^t k 1 k k 1 k = C_n k R k^t C_n^t C_n e^{(1)t} ": \square$

The following lemma and Lemma 17, whose proof follows, enable us to use Lemma 29 to bound the mixing time of random update G lauber dynamics.

Lem m a 30. Let R be a dependency matrix for a spin system Let R be the random update matrix for R. Then, for f 2 R^M , (P f) R (f).

Proof. For each i 2 [n], from the denition of i, i(f) 0 and, for any c 2 R and f 2 R^M, $i(cf) = jcj_i(f)$. Also, $i(f_1 + f_2) = i(f_1) + i(f_2)$ for any $f_1; f_2 2 R^M$. Now,

$$(P f) = \frac{1}{n} \prod_{j=1}^{P} P^{[j]}f = \frac{1}{n} \prod_{j=1}^{P} P^{[j]}f \qquad \frac{1}{n} \prod_{j=1}^{P} P^{[j]}f):$$

By Lemma 28, this is at most $\frac{1}{n} \prod_{j=1}^{P} R_j$ (f) = R (f).

Remark 5. The proof shows that $_{i}(f)$ is a (vector) seminorm for all i2 [n]. It fails to be a norm because $_{i}(f) = 0$ does not imply f = 0. For example, $_{i}(1) = 0$ for all i2 [n].

We can now give a proof of Lem m a 17 using this approach.

Lem m a 17. Suppose R is a dependency matrix for a spin system, and let k k be any matrix norm. If kR k < 1, then the mixing time $_{r}$ (") of random update G lauber dynamics is at most n (1) 1 ln (C $_{n}$ =").

Proof. By Lemma 16, kR k = $1 \frac{1}{n} (1)$ and, by Lemma 30, (P f) R (f). Then, by Lemma 29, r(") (1) $1 \ln (C_n = ") = n (1)$ $1 \ln (C_n = ")$.

C orollaries 18 and 19 and the rest of that section now follow exactly as before. A similar analysis applies to system atic scan, though it is slightly easier. It relies on the analogue of Lemma 30, which in this case is immediate from Lemma 28.

Lem m a 31. Let R be a dependency matrix for a spin system. Let \hat{R} be the scan update matrix for R. Then, for any f 2 R^M, \hat{F} f) \hat{R} (f).

We can now give a proof of Lem m a 22 using this approach.

Lem m a 22 Let R be a dependency matrix for a spin system, and k k any matrix norm. If \mathbb{R}^{k} < 1, the mixing time s(") of system atic scan G lauber dynamics is at most (1) $^{1} \ln (C_{n} = ")$. If k k is an operator norm, the mixing time is at most (1) $^{\frac{1}{2}} \ln (J_{n} = ")$.

Proof. By Lemma 31, for any $f \ge \mathbb{R}^{M}$, $\mathfrak{P} f$) \mathfrak{R} (f). Then, by assumption, $\mathfrak{k} \mathfrak{R} \mathfrak{k} < 1$. Now apply Lemma 29.

The results following Lemma 22 in section 3.1 can then be obtained identically to the proofs given there.

3.3 Im proved analysis of system atic scan

W e m ay improve the analysis of C orollary 27 for the case in which the diagonal of R is 0, which is the case for the heat bath dynamics. For 0, de ne R by

so R has its upper triangle scaled by . Let ${}^{\circ}_{j}$ denote the jth column of R, for j 2 [n]. We can now prove the following strengthening of Lemma 23.

Lemma 32. If wR w, for some w 0 and 0 1, then $\tilde{\mathbf{W}}$ wR.

Proof. W e prove by induction that

 $wR_1R_2 \quad i \quad R \quad [w_1^{\circ}] \quad i \quad w_{1+1}^{\circ} \quad n] \quad w \quad [w_1 \quad i \quad w_{1+1}^{\circ} \quad n]: w$

The second inequality follows by assumption. The hypothesis is clearly true for i = 0. For i > 0,

where $we = [w \aleph_1 \quad i \aleph w i \quad n] w [w_1 \quad i \mid ww i \quad n]$. We follows that $we \aleph_i \quad w \aleph_i$, continuing the induction. Putting i = n gives the conclusion

Lemma 33. If R is symmetric and = (R) < 1 then (R) if = = (2).

Proof. We have = (R) = kRk_2 by Lemma 12. Since R is symmetric with zero diagonal, $x^TR x = \frac{1}{2}(1 +)x^TRx$. It follows that (R) (R) = $\frac{1}{2}(1 +)$ (R) = $\frac{1}{2}(1 +)$. Therefore (R) if 2 = (1 +). This holds if = (2).

Lem m a 34. Let R be symmetric with zero diagonal and $kR k_2 = (R) = < 1$, and 0 < < 1Let = + < 1. Then the mixing time of system atic scan is at most

$$s(") \quad \frac{2}{2 \quad 2} \ln (n = "):$$

Proof. Let $n^0 = n$ 1, and S = R + (J I) = h. Since S R, S is a dependency matrix for the original spin system . Also, S is symmetric and its diagonal is 0. Now

$$(S) = kSk_2 = kR + (J I) = nk_2 kRk_2 + kJ Ik_2 = n^0 + = :$$

Denote by $S = S_1S_2$ n the scan matrix. Let = = (2). Now by Lemma 33, we have (S) . Furthermore, S is irreducible, so by Lemma 7, there exists a row vector w > 0satisfying wS w. We can assume without loss of generality that w is normalised so that $kw k_1 = 1$. Finally, we can conclude from Lemma 32 that wS wS.

Since w S w S w, we have established that convergence is geometric with ratio, but we need a lower bound on $w_{min} = w_{min}$ in order to obtain an upper bound on mixing time via Lemma 29. Now

$$w \ w \ S \ w \ R + (J \ I) = n^{0} \ w \ (J \ I) = n^{0} = (=n^{0}) (1 \ w):$$
So w $(1 + = n^{0})$ $(=n^{0})1$, and $w_{m \ in} = (n^{0} +) = n \ By \ Corollary 26$, the m ixing time satisfies
$$s^{(")} (1 \)^{1} \ln (1 = w_{m \ in} ") (1 \)^{1} \ln (n = "):$$

We can now prove Lemma 3.

Lem m a 3. Let R be symmetric with zero diagonal and $kRk_2 = (R) = < 1$. Then the mixing time of systematic scan is at most

$$f_{s}(")$$
 (1 $\frac{1}{2}$)(1)¹ ln (1)¹ n=":

Proof. We apply Lemma 34 with $= (1) = \ln n$, and hence

Remark 6. If, as in Remark 3, if $(1) = (\log^{k} n)$ for some k 0, then we have mixing time $\binom{s}{s}$ (") $(1 \frac{1}{2})(1)^{1} \ln (n=")$ for system atic scan. We may compare the number of G lauber steps n_{s} (") with the estimate $\binom{r}{r}$ (") = $(1)^{1} n \ln (n=")$ for random update G lauber dynamics obtained from C orollary 18 using the minimum norm k $_{2}k$. The ratio is $(1 \frac{1}{2}) < 1$. This is close to $\frac{1}{2}$ when (R) is close to 1, as in many applications.

Example 2. Consider colouring a -regular graph with (2 + 1) colours [18, 22] using heat bath G lauber dynamics, we have (R) = = (+1). (See section 4). Then (1) = 1 = (+1) = (1), if = 0 (1), and the above ratio is $(1 \frac{1}{2}) = (+2) = (2 + 2)$. This is close to $\frac{1}{2}$ for large.

A lthough the improvement in the mixing time bound is a modest constant factor, this provides some evidence in support of the conjecture that systematic scan mixes faster than random update, for G lauber dynamics at least. The improvement is because we know, later in the scan, that most vertices have already been updated. In random update, some vertices are updated many times before others are updated at all. Lemma 34 suggests that this may be wasteful.

4 Colouring sparse graphs

In this section, we consider an application of the methods developed above to graph colouring problems, particularly in sparse graphs. By sparse, we will mean here that the number of edges of the graph is at most linear in its number of vertices.

Let G = (V; E), with V = [n], be an undirected (simple) graph or multigraph, without self-boops. Then d_v will denote degree of vertex $v \ge V$. If S = V, we will denote the induced subgraph by $G_S = (S; E_S)$. The (symmetric) adjacency matrix A(G) is a non-negative integer matrix, with zero diagonal, giving the number of edges between each pair of vertices. We write A for A(G) and (G) for (A(G)). Thus the adjacency matrix of a graph is a $O\{1 \text{ matrix}$. We also consider digraphs and directed multigraphs G = (V; E). We denote the indegree and outdegree of $v \ge V$ by $d_v; d_v^+$ respectively.

If G is a graph with maximum degree , we consider the heat bath G lauber dynam ics for properly colouring V with q > colours. The dependency matrix R for this chain satis es i_{j} 1=(q d_j) (i; j 2 [h]) (see Section 5.2 of [10]). Thus R = AD, where D = diag(1=(q d_j)). Let $D^{1=2} = diag(1= q d_j)$ and $\hat{A} = D^{1=2}AD^{1=2}$. Note that \hat{A} is symmetric. Also, $(\hat{A}) = (AD)$, since $(D^{1=2}AD^{1=2})(D^{1=2}x) = (D^{1=2}x)$ if and only if AD x = x. If (i; j) 2 E, we have $\hat{A}_{ij} = 1= q (q d_j)(q d_j)$. Since $\hat{A} = 1 q A$, we have $(\hat{A}) = 1 q A$ from Lemma 9. So if q > 1= A, we have $(\hat{A}) = 1 q A$. However, (\hat{A}) seems more discussion to estimate than (A), since it depends more on the detailed structure of G. Therefore we will use the bound $\frac{1}{q}$ (A) in the remainder of this section, and restrict most of the discussion to (G). The follow ing is well known.

Lem m a 35. If G has maximum degree and average degree d, then d (G) . If either bound is attained, there is equality throughout and G is -regular.

Proof. The vertex degrees of G are the row or column sum sof A (G). The upper bound then follows from (G) $kAk_{i} = m ax_{v2V} d_{v} = using Lemma 10$. For the lower bound, since G is undirected, (G) = (A) $\frac{1^{T}}{P_{n}}A\frac{p_{n}}{P_{n}} = 2jE j = n = d$, using Lemma 12. If the lower bound is attained, then the inequalities in the previous line are equalities, so 1 is an eigenvector of A. Thus A1 = d1, and every vertex has degree d = . When the upper bound is attained, since the columns sum s of A are at most , 1A = 1, so 1 is an eigenvector from Lemma 7, and 1A = 1. Then every vertex has degree = d.

Thus the resulting bound for colouring will be q > 2 when G is -regular, as already shown by Jerrum [18] or Salas and Sokal [22]. Thus we can only achieve mixing for q = 2 by this approach if the degree sequence of G is nonregular.

We now derive a bound on (R) for symmetric R which is very simple, but nonetheless can be used to provide good estimates in some applications.

Lem m a 36. Suppose R 2 M $_{n}^{+}$, and we have R = B + B^T, for som e B 2 M $_{n}$. If k k is any matrix norm, then (R) 2 kB kkB k.

Proof. (R) = $kB + B^T k_2$ $kB k_2 + kB^T k_2 = 2kB k_2$ 2 kB kkB k, using the self-duality of $k_2 k$ and Lem m as 12 and 13.

Corollary 37. If
$$R = B + B^T$$
, then (R) $2 kB k_1 kB k_1$.

We can use Corollary 37 as follows. If $R \ge M_n^+$, let $(R) = \max_{[n]} \sum_{i;j \ge 1} \sum_{ij=2}^{n} \sum_{ij=2}^{n}$

$$(\mathbf{R}) = \max_{x \ge f0; \lg^n} \frac{x^T \mathbf{R} x}{x^T x} \qquad \max_{x \ge R^n} \frac{x^T \mathbf{R} x}{x^T x} = (\mathbf{R}) = (\mathbf{R}):$$

Also, (R) = kRk_1 2 (R), since X(R) = $\max_{I \ [n]} \sum_{i,j \ge I} \sum_{I \ [n]} \sum_{j \ge [n]; j \ge I} \sum_{I \ [n]} \sum_{j \ge [n]; j \ge I} \sum_{I \ [n]} \sum_{I \ [n]} \sum_{i \ge [n]; j \ge I} \sum_{I \ [n]} \sum_{I \ [n]}$

W e m ay easily bound the maximum density for some some classes of graphs. For any $a;b \ge 2$, let us de ne G (a;b) to be the maxim al class of graphs such that

(i) G (a;b) is hereditary (closed under taking induced subgraphs);

(ii) for all G = (V; E) 2 G(a; b) with j V j = n, we have j E j an b.

Lem m a 38. Let G 2 G (a;b) with j j = n. If

(i) b 0, then (G) a b=n. (ii) b 0, let k = $a + \frac{p}{(a + \frac{1}{2})^2}$ 2b, then (G) = maxf(bk c 1)=2; a b=dkeg. Proof. In case (i), clearly f = f j a b=n. If S V, f = f j a b=f j a b=n. In case(i),

note that (G) $\frac{1}{n} \frac{n}{2} = \frac{1}{2}$ (n 1) for any simple graph G on n vertices. Thus,

Note that (s 1)=2 is increasing in s and a b=s is decreasing in s. Also, s = k is the positive solution to (s 1)=2 = a b=s. The other solution is not positive since b 0. Thus

Remark 7. We could consider a more general class G (a_n ; b_n), where $b_n j = o(na_n)$. This includes, for example, subgraphs of the d-dimensional hypercubic grid with vertex set $V = [k]^d$ in which each interior vertex has 2d neighbours. Then f_{ij} dn $dn^{1-1=d}$, so $a_n = d$ and $b_n = dn^{1-1=d}$. However, we will not pursue this further here.

W e can apply Lem m a 38 directly to som e classes of sparse graphs.

For the de nition of the tree-width t(G) of a graph G, see [6]. We say that a graph G has genus g if it can be embedded into a surface of genus g. See [5] for details, but note that that text (and several others) de ne the genus of the graph to be the sm allest genus of all surfaces in which G can be embedded. We use our de nition because it is appropriate for hereditary classes. Thus, for us a planar graph has genus 0, and a graph which can be embedded in the torus has genus 1 (whether or not it is planar).

Lem m a 39. If a graph G = (V; E) is

- (i) a nonregular connected graph with maximum degree , then G 2 G (=2;1);
- (ii) a forest, then $G \ 2 \ G \ (1;1);$
- (iii) a graph of tree-width t, then G 2 G (t;t(t + 1)=2);
- (iv) a planar graph, then $G \ 2 \ G \ (3; 6)$.
- (v) a graph of genus g, then $G \ 2 \ G \ (3; 6(1 g))$.

Proof. Note that (ii) is a special case of (iii), and (iv) is a special case of (v). For (i), if $G_s = (S; E_s)$ is an induced subgraph of G, then G_s cannot be -regular, and $E_s j = \frac{1}{2}$ f j 1. For (iii) and (v), the graph properties of having tree-width at most t, or genus at most g, are hereditary. A lso, if j' j = n, a graph of tree-width thas at most tn t(t+1)=2 edges (See, for example, 1, Theorem 1, Theorem 34].), and a graph of genus g at most 3n 6(1 - g) edges (See, for example, 5, Theorem 7.5, C orollary 7.9].).

Remark 8. In (i) { (iv) of Lemma 39 we have b > 0, but observe that in (v) we have b > 0 if g = 0 (planar), b = 0 if g = 1 (toroidal) and b < 0 if g > 1.




Figure 2: Upper and lower bounds on maximum density for small genus g

Corollary 40. If a graph G = (V; E) on n vertices is

(i) a nonregular connected graph with maximum degree, then (G) $\frac{1}{2}$ $\frac{1}{n}$; (ii) a forest, then (G) 1 $\frac{1}{n}$; (iii) a graph of tree-width t, then (G) t $\frac{t(t+1)}{2n}$; (iv) a planar graph, then (G) 3 $\frac{6}{n}$. (v) a graph of genus g > 0, let $k_g = 7=2 + p \frac{1}{12g + 1=4}$, then (G) $g = m \exp(bk_g c - 1)=2;3 + 6(g - 1)=dkgeg$.

Proof. Follows directly from Lemmas 38 and 39.

Remark 9. Suppose that g is chosen so that k_g is an integer. The bound in Corollary 40 (v) gives $g = (k_g \quad 1)=2$ (because k_g is the point at which the two arguments to the maximum are equal). The bound says that for every graph G with genus g, (G) g. This bound is tight because there is a graph G with density (G) = g and genus g. In particular, the complete graph K k_g has density g. If $k_g \quad 3$, it also has genus g. The sm allest genus of a surface in which it can be embedded is $g = d(k_g \quad 3)(k_g \quad 4)=12e$ (see, for example, 5, Thm .7.10]). This is at least¹ g since

$$\frac{k_g^2 - 7k_g + 12}{12} = g;$$

so the genus of G is g as required. The bound in Corollary 40 (v) m ay not be tight for those g for which k_g is not integral. However, the bound is not greatly in error. Consider any g > 0. The graph $G = K_{bk_gc}$ can be embedded in a surface of genus g so it has genus g. A lso, as noted above, $(G) = \frac{1}{2}$ (bk_gc 1). If the bound is not tight for this g and G then

(G)
$$_{g} = 3 + \frac{6(g-1)}{dk_{g}e} - 3 + \frac{6(g-1)}{k_{g}} = \frac{k_{g}-1}{2} = \frac{bk_{g}c-1}{2} + \frac{k_{g}-bk_{g}c}{2}$$
 (G) $+ \frac{1}{2}$: (7)

so g cannot be too much bigger than (G). It is easy to see that $g = \frac{P}{3g}$ for large g. For small g, a plot of the upper bound g on maximum density is shown in Figure 2, together with the lower bound l=2 ($bk_{g}c = 1$).

We now show that there exists a suitable B for applying C orollary 37.

¹ In fact, = g, though we don't use this fact here.

Lem m a 41. Let R 2 M $_n^+$ be symmetric with maximum density and let = kR k₁. Then there exists B 2 M $_n^+$ such that R = B + B^T and kB k₁ = , kB k₁ = .

Proof. It will be su cient to show that $kB k_1$, $kB k_1$, since then we have

$$= kR k_{1} = kB + B^{T} k_{1} \quad kB k_{1} + kB^{T} k_{1} = kB k_{1} + kB k_{1} + () = ;$$
(8)

First suppose R is rational. Note that is then also rational. Let $R^0 = R$ D, where D = diag $({}^{\circ}_{Si})$. Thus, for some large enough integer N > 0, A (G) = N R⁰ is the adjacency matrix of an undirected multigraph G = (V; E) with V = [n], N D is a matrix of even integers, and N is an integer. Thus, provided B is eventually rescaled to B=N, we may assume these properties hold for R⁰, D and . An orientation of G is a directed multigraph G = (V; E') such that exactly one of $e^+ = (v; w)$, e = (w; v) is in E for every e = fv; wg 2 E. Clearly A (G) = A (G) + A (G)^T, so we may take $B = A (G) + \frac{1}{2}D$. Note that kB k₁ = max_{v2V} (d_v + $\frac{1}{2}$ %_{vv}) and kB k₁ = max_{v2V} (d_v⁺ + $\frac{1}{2}$ %_{vv}). We now apply the following (slightly restated) theorem of Frank and G yarfas [14].

Theorem 42 (Frank and Gyarfas). Suppose v_v u_v for all $v \ge V$ in an undirected multigraph G = (V; E). Then G has an orientation G satisfying v_v d_v u_v if and only if, for all S V, we have $E_S j$ max $v_{2S} u_v$; $v_{2S} d_v$ v_v .

We will take $u_v = \frac{1}{2} \aleph_{vv}$, $\aleph_v = d_v + \frac{1}{2} \aleph_{vv}$. Then $\aleph_v = u_v$, since $d_v = (\aleph_v)$, and $(d_v \ \aleph_v)$ u_v , since 2. The conditions of Theorem 42 are satisfied since, for all S V, $\mathfrak{F}_S j = \frac{1}{2} \qquad \aleph_{vw} \qquad \frac{1}{2} \qquad \aleph_{vv} \qquad \mathfrak{F}_j \qquad \frac{1}{2} \qquad \aleph_{vv} = \begin{array}{c} X & X \\ \mathfrak{F}_S j = \frac{1}{2} \qquad \aleph_{vw} \qquad \frac{1}{2} \qquad \aleph_{vv} \qquad \mathfrak{F}_j \qquad \mathfrak{F}_j \qquad \mathfrak{F}_j \qquad \mathfrak{F}_v = \begin{array}{c} X & X \\ \mathfrak{F}_v = \chi & \chi \\ \mathfrak{$

The result now follows for rational R, since we have

 $kB k_{1} = \max_{v \ge V} (d_{v} + \frac{1}{2} \aleph_{vv}) \qquad ; \quad kB k_{1} = \max_{v \ge V} (d_{v}^{+} + \frac{1}{2} \aleph_{vv}) \qquad \max_{v \ge V} (d_{v}^{-} + \frac{1}{2} \aleph_{vv}) = :$

If R is irrational, standard continuity arguments now give the conclusion.

Remark 10. The use of Theorem 42 in the proof can be replaced by an application of the max-ow min-cut theorem, as in [15], but Theorem 42 seems more easily applicable here.

We can show that Lemma 41 is best possible, in the following sense.

Lem m a 43. Let R 2 M⁺_n be symmetric with maximum density and let = $kR k_1$. If R = B + B^T for any B 2 M_n, then $kB k_1$ and $kB k_1$ kB k.

Proof. Let I be any set achieving the maximum density of R. Then

$$2 \operatorname{JI} j = \begin{cases} X & X & X & X \\ & & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ & & \\ &$$

so kB k_1 . The second assertion follows from $\ensuremath{\left. \boldsymbol{\beta} \right)}$.

Theorem $p_{(R)} = \frac{44 \cdot 16R}{2} \times M_n^+$ is a symmetric matrix with maximum density and $k = kRk_1$, then

Proof. Follows directly from Corollary 37 and Lemma 41.

Remark 11. Since () is increasing for =2, an upper bound 0 can be used, as long as we ensure that 0 =2.

Rem ark 12. We can adapt this for asymmetric R by considering the \symmetrisation" $\frac{1}{2}$ (R + R^T). Note that (R) = $\frac{1}{2}$ (R + R^T). Let ~(R) = $k\frac{1}{2}$ (R + R^T) k_1 $p\frac{1}{2}$ (kR k_1 + kR k_1). We also have (R) (R) = $\frac{1}{2}$ (R + R^T) = $\frac{1}{2}$ (R + R^T). Then (R) $2\frac{1}{2}$ (~).

The following application, used together with Lemma 2, strengthens [10, Thm .15].

Theorem 45. Suppose R is a symmetric and irreducible dependency matrix with row sums at most 1, and suppose $0 < min_{j;2[n]}f_{ij} : s_{ij} > 0$ g. If there is any row with sum at most 1 , then (R) $p = \frac{1}{1} \frac{2}{2} = n^2$ 1 $2^2 = 2n^2$.

Proof. Since R is irreducible, for any I [n], $\stackrel{P}{}_{i;j^{2}I} \stackrel{*}{}_{ij}$ jIj . This also holds for I = [n] by assumption. Thus $\frac{1}{2}$ $\frac{1}{2n}$. Since kR k₁ 1, we have (R) $2 \frac{1}{2} \frac{1}{2n} \frac{1}{2n} \frac{1}{2n} = \frac{1}{2} \frac{1}{2n} \frac{1}{2n} \frac{1}{2} \frac{1}{2n} \frac{1}{2n} \frac{1}{2} \frac{1}{2}$

We can also apply Theorem 44 straightforwardly to (simple) graphs.

Corollary 46. If G has maximum density and maximum degree, then (G) 2^{p} (). Proof. In Theorem 44, we have = .

Theorem 47. If G = (V; E) 2 G(a; b), with b 0, 2a and j V j = n, then

(G)
$$2 = a \frac{b}{n} = a + \frac{b}{n}$$

 $a = b + \frac{b}{n}$
 $a = b + \frac{b}{a}$
 $a = b + \frac{b}{a$

Proof. The rst inequality follows directly from Lemma 38 and Corollary 46. Note that the condition 2a 2b=n is required in view of Remark 11. For the second, squaring gives

4a (a)
$$\frac{4b(2a)}{n} \frac{4b^2}{n^2}$$
 4a (a) $\frac{4b(2a)}{n} + \frac{b^2(2a)^2}{a(a)n^2}$;

which holds for all b and 2a. W hen = 2a, using $p_{1-x} = 1 \frac{1}{2}x$,

(G)
$$2 a^2 \frac{b^2}{n^2} = 2a 1 \frac{b^2}{a^2n^2}$$
 $2a 1 \frac{b^2}{2a^2n^2} = a 2 \frac{b^2}{a^2n^2}$:

Theorem 48. If G = (V; E) 2 G(a; b), with b 0 and j V j = n, let $k = a + \frac{1}{2} + \frac{p}{(a + \frac{1}{2})^2} 2b$ and = maxf(bk c 1)=2; a b=dkeg. Then, if 2, (G)

W e can apply this to the exam ples from Lemma 39.

Corollary 49. If G = (V; E), with maximum degree and j V j = n, is

(i) a connected nonregular graph, then

(G)
$$\frac{1}{2} \frac{4}{n^2} < \frac{2}{n^2}$$
:

(ii) a tree with 2, then

(G)
$$2 \ 1 \ \frac{1}{n} \qquad 1 + \frac{1}{n} < \frac{p}{1} \ 2 \ \frac{2}{(1)n}$$
:

If = 2 then (G) < 2 $1=n^2$, and if < 2 then (G) = . (iii) a graph with tree-width at most t and 2t, then

(G) 2 t
$$\frac{t(t+1)}{2n}$$
 t $\frac{t(t+1)}{2n}$ < $p = \frac{t(t+1)(2t)}{2(t)n}$:

If =
$$2t$$
 then (G) < $2t$ t(t + 1)²= $4n^2$

(iv) a planar graph with 6, then

r

(G) 2 3 6=n 3+6=n <
$$2^{p} \frac{1}{3(3)} 1 \frac{6}{(3)n}$$
:

If = 6, (G) 6
$$12=n^2$$
. If 5, (G) is best possible.
(v) a graph of genus g > 0, let $k_g = 7=2+p^2$ $12g + 1=4$ and $g = m \exp(bk_g c 1)=2;3+6(g 1)=dkgeg$.
If 2 g, then q

Proof. Using Lemma 39, these follow using Theorem 47 and Theorem 48 with

(i)
$$a = -2; b = 1 and = 2a$$
.

(ii) a = 1; b = 1, if > 2. If = 2, the result follows from the = 2a case. = 1, G is a single edge and, if = 0, an isolated vertex.

(iii)
$$a = t; b = t(t + 1)=2.$$

- (iv) a = 3; b = 6. If 5, regular planar graphs with degree exist, and we use Lemma 35. (v) a = 3; b = 6 (g 1).
- Rem ark 13. If G is a disconnected graph, the component having the largest eigenvalue determ ines (G), using Lemma 8. This can be applied to a forest.

Remark 14. Corollary 49(i) in proves on a result of Stevanovic [26], who showed that

(G) <
$$\frac{1}{2n(n-1)^{-2}}$$
:

This was improved by Zhang [28] to (approximately) $\frac{1}{2}$ (n) ², which is still inferior to (i). But recently the bound has been improved further by Cioaba, G regory and N ikiforov [4], who showed

(G) <
$$\frac{1}{n(D + 1)}$$
;

where D is the diam eter of G. This gives (G) $1=n^2$ even in the worst case, which signi – cantly in proves on (i). However, C orollary 49 is an easy consequence of the general C orollary 46, whereas [4] uses a calculation carefully tailored for this application.

Remark 15. When G is a degree-bounded forest, Corollary 49(ii) strengthens another result of Stevanovic [25], who showed (G) < 2^{p} 1.

Remark 16. When G is a planar graph, Theorem 47 (iv) im proves a result of Hayes [16].

We can now apply these results to the mixing of G lauber dynamics for proper colourings in the classes of sparse graphs G (a;b).

Theorem 50. Let G = (V; E) 2 G (a; b), with b > 0, have maximum degree 2a, where jV j = n. Let = $2^{p} \overline{a(a)}$, = 2a and = = (q). Then, if

(i) q > + , the random update and system atic scan G lauber dynam ics m ix in time

r(") (1) $\ln \ln (n=");$ $\hat{s}(")$ (1) $\frac{1}{2}$) (1) $\ln (n="):$

(ii) q = + and > 0, the random update and system atic scan G lauber dynam ics m ix in time

(")
$$(^{2}=2b)n^{2}\ln(n=");$$
 $^{s}(")$ $(^{2}=2b)n\ln(n="):$

(iii) q = + and = 0, the random update and system atic scan G lauber dynam ics m ix in time

$$r(")$$
 2 (a=b)² n³ ln (n="); $s(")$ 3 (a=b)² n² ln (n="):

Proof. Recall from the beginning of Section 4 that (R) (G)=(q) where (G) denotes (A (G)). Note also that, if is not an integer, then q = (1). By Theorem 47, for (i) we have $kR k_2 = (R)$ (G)=(q) = (q) = < 1. For (ii), we have (R) 1 ($2b = {}^2n$), and for (iii), (R) 1 ($b=2a^2n^2$). The conclusions for _r(") follow from Lemma 17, and those for _s(") from Lemma 3. For (ii) and (iii), factors of $\frac{1}{2}$ arise in Lemma 3 since 1, but additional factors (2 and 3, respectively) come from the log term.

Theorem 51. If G = (V;E) 2 G(a;b) with b 0, let $k = a + \frac{1}{2} + \frac{p}{(a + \frac{1}{2})^2}$ 2b and = maxf(bk c 1)=2;a b=dkeg. If > 2, let = (q) and = = (q). Then, if q > +,

r(") (1) 1 n ln (n="); s (") (1) 1 ln (n="):

Proof. From Theorem 48, we have

r

$$kR k_2 = (R) \frac{(G)}{q} = < 1:$$

The conclusions for r(") now follow from Lemmas 14 and 17, and those for r(") from Lemma 3.

Corollary 52. If G = (V; E), with j V j = n and maximum degree, is

(i) a nonregular connected graph and $q=\,2\,$, then

 $r^{(0)} = 2^{2}n^{3}\ln(n="); \quad s^{(0)} = 2^{2}n^{2}\ln(n=");$ (ii) a graph with the width thand 2t, let $= 2^{p} \frac{1}{t(-1)}$. Then $\begin{cases} (q -)(q -)^{-1}n\ln(n="); \quad \text{if } q > + ; \\ r^{(0)} = 2^{-2}(t(t+1)(-2t))^{-1}n^{2}\ln(n="); \quad \text{if } q = + \text{ and } > 2t; \\ 8(t+1)^{-2}n^{3}\ln(n="); \quad \text{if } q = + \text{ and } > 2t; \\ 8(t+1)^{-2}n^{3}\ln(n="); \quad \text{if } q = + \text{ and } > 2t; \\ 12(t+1)(-2t))^{-1}n\ln(n="); \quad \text{if } q = + \text{ and } > 2t; \\ 12(t+1)^{-2}n^{2}\ln(n="); \quad \text{if } q = + \text{ and } > 2t; \\ 12(t+1)^{-2}n^{2}\ln(n="); \quad \text{if } q = + \text{ and } > 2t; \\ 12(t+1)^{-2}n^{2}\ln(n="); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{(12(t+1))} \ln(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ 12(t-1)^{-2}n^{2}\ln(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{(12(t-1))} \ln(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{(12(t-1))} \ln(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } > 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } = 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } = 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } = 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } = 2t; \\ r^{(0)} = 2^{-2}(12(t-1))^{-2}n(n=m); \quad \text{if } q = + \text{ and } = 2t; \\ r^{(0)$

Proof. This follows directly from Lemma 39 and Theorems 50 and 51.

Remark 17. Corollary 52 (i) bounds the mixing time of heat bath G lauber dynamics for sampling proper q-colourings of a nonregular graph G with maximum degree when q = 2. (We can bound the mixing time for a disconnected graph G by considering the components.) It is also possible to extend the mixing time result for nonregular graphs to regular graphs using the decomposition method of M artin and R andall [19]. See [10, Section 5] for details about how to do this. The use of our C orollary 52 (i) in proves Theorem 5 of [10] by a factor of n.

References

[1] H.Bodlaender, A partial k-arboretum of graphs with bounded treewidth, Theoretical Com – puter Science 209 (1998), 1{45.

- [2] M.Bordew ich and M.D yer, Path coupling without contraction, Journal of Discrete A lgorithm s, doi:10.1016/j.jda.2006.04.001, 2006, in press.
- [3] R.Bubley and M.D yer, Path coupling: a technique for proving rapid mixing in M arkov chains, in Proc. 38th Annual IEEE Symposium on Foundations of Computer Science, IEEE Computer Society, 1997, pp. 223{231.
- [4] S.Cioaba, D.Gregory and V.Nikiforov, Extrem e eigenvalues of nonregular graphs, Journal of Combinatorial Theory, Series B, to appear.
- [5] G. Chartrand and L. Lesniak, Graphs and Digraphs, 4th edn., Chapm an and Hall/CRC, 2005.
- [6] R.Diestel, Graph theory, 3rd edn., Springer, Berlin, 2006.
- [7] R.Dobrushin, Prescribing a system of random variables by conditional distributions, Theory of Probability and its Applications 15 (1970), 458{486.
- [8] R. Dobrushin and S. Shlosman, Constructive criterion for the uniqueness of a Biggs eld, in Statisticalm echanics and dynam ical systems (J.Fritz, A.Ja e and D.Szasz, eds.), Birkhauser, Boston, 1985, pp. 347 (370.
- [9] W . Doeblin, Expose de la theorie des cha^nes simple constantes de Marko a un nombre ni d'etats, Revue Mathematique de l'Union Interbalkanique 2 (1938), 77{105.
- [10] M. Dyer, L. Goldberg and M. Jernum, Dobrushin conditions and systematic scan, Electronic Collequium on Computational Complexity, http://eccc.hpi-web.de/eccc/, TR05-075, 2005.
- [11] M. Dyer, L. Goldberg and M. Jerrum, Dobrushin conditions and systematic scan, in Proc. 10th InternationalW orkshop on Random ization and Computation, Lecture Notes in Computer Science 4110, Springer, 2006, pp. 327{338.
- [12] M. Dyer and C. Greenhill, Random walks on combinatorial objects, in Surveys in Combinatorics 1999 (J.Lamb and D. Preece, eds), LM S Lecture Note Series 267, Cambridge University Press, Cambridge, 1999, pp. 101{136.
- [13] H. Follmer, A covariance estimate for G ibbs measures, Journal of Functional Analysis 46 (1982), 387{395.
- [14] A. Frank and A. Gyarfas, How to orient the edges of a graph?, in Combinatorics (Proc. 5th Hungarian Colloquium, Keszthely, 1976), Vol. I, Colloquia M athematics Societatis Janos Bolyai 18, North-Holland, Am sterdam, 1978, pp. 353{364.
- [15] A.Goldberg, Finding a maximum density subgraph, Tech. Rep. UCB/CSD 84/171, University of California, Berkeley, 1984.

- [16] T. Hayes, A simple condition implying rapid mixing of single-site dynamics on spin systems, in Proc. 47th Annual IEEE Symposium on Foundations of Computer Science, IEEE Computer Society, 2006, pp. 39{46.
- [17] R. Horn and C. Johnson, Matrix analysis, Cambridge University Press, Cambridge, 1990.
- [18] M. Jerrum, A very simple algorithm for estimating the number of k-colorings of a low-degree graph, Random Structures and Algorithm s 7 (1995), 157{165.
- [19] R. Martin and D. Randall, Sampling adsorbing staircase walks using a new Markov chain decomposition method, in Proc. 41st Annual IEEE Symposium on Foundations of Computer Science, IEEE Computer Society, 2000, pp. 492{502.
- [20] M.M itzenm acher and E.Upfal, Probability and computing, Cambridge, 2005.
- [21] K. Pedersen, PhD thesis, University of Liverpool, in preparation.
- [22] J. Salas and A. Sokal, Absence of phase transition for antiferrom agnetic Potts models via the D obrushin uniqueness theorem, Journal of Statistical Physics 86 (1997), 551 (579.
- [23] E. Seneta, Non-negative matrices and Markov chains, rev. 2nd edn., Springer, New York, 2006.
- [24] B. Sim on, The statistical mechanics of lattice gases, Princeton Series in Physics, vol. I, Princeton University Press, Princeton, 1993.
- [25] D. Stevanovic, Bounding the largest eigenvalue of trees in terms of the largest vertex degree, Linear Algebra and its Applications 360 (2003), 35{42.
- [26] D. Stevanovic, The largest eigenvalue of nonregular graphs., Journal of C om binatorial Theory, Series B, 91 (2004), 143{146.
- [27] D.Weitz, Combinatorial criteria for uniqueness of Gibbs measures, Random Structures and Algorithms 27 (2005), 445{475.
- [28] X. Zhang, Eigenvectors and eigenvalues of non-regular graphs, Linear Algebra and its Applications 409 (2005), 79{86.