CORNER VIEW ON THE CROWN DOMAIN

BERNHARD KROTZ

Date:M ay 26, 2019.

1. Introduction

Our concern is with the crown domain, henceforth denoted by . We recall that is an equivariant complexi cation of a Riemannian symmetric space X = G=K of the non-compact type. Most naturally one de ness by the theory of unitary K -spherical representations of the symmetry group G (see the introduction of [3]). Geometrically, one can de ne as the maximal G-invariant domain in the a necomplexient cation $X_c = G_c = K_c$ which can be equivariantly embedded into the tangent bundle TX.

As a complex manifold, has the property that bounded holom orphic functions separate points. Therefore we may de ne its distinguished (i.e. Shilov) boundary Q_d as the smallest closed subset of the topological boundary Q_d on which bounded plurisubharm onic functions on cl() attend their maximum. We know by [] and [3] that Q_d is a nite (and explicite) union of G-orbits, say

$$Q_d = O_1 q ::: q O_s :$$

From now on we shall identify each O_j with a homogeneous space: G=H_j. The main result of [1] was:

If $G = H_j$ is a symmetric space, then it is a non-compactly causal symmetric space. Moreover, every non-compactly causal symmetric space Y = G = H appears in the distinguished boundary of the corresponding crown domain for X = G = K.

The aim of this paper is to understand this result better. To be m ore concise: what is the reason that precisely non-com pactly causal (NCC) sym m etric spaces appear in the boundary?

NCC-spaces are very special am ong all sem is in ple sym metric spaces. We recall their de nition. We assume the Lie algebra of G to be simple and write q for the tangent space of Y at the standard base point $y_o = H 2 Y$. We note that q is a linear H -m odule. Now, non-com pactly causalm eans that q admits an non-empty open H -invariant convex cone, say C, which is hyperbolic and does not contain any a ne lines.

The theme of this paper is to view from the corner point $y_0 2 Y$ and not as a thickening of X as custom ary. Now a slight precision is of need. In general Q_d has several connected components. If this happens to be the case, then we shrink to a G-dom ain _H whose distinguished boundary is precisely Y, see [2].

2

For C being the m inim alcone we form in the tangent bundle $TY = G_{H} q$ the cone-subbundle

 $C = G _{H} C$

and with that its boundary cone-bundle

$$QC = G_{H} QC$$
:

In this context we ask the following

Question: Is there a G -equivariant, generically in jective, proper continuous surjection p: $C! Q_H$?

In other words, we ask if there exists an equivariant "resolution" of the boundary in terms of the geometrically simple boundary cone bundle @C.

In this paper we give an a mative answer to this question if X is a Herm itian tube domain. In this simplied situation the crown domain is $= X - \overline{X}$ with \overline{X} denoting X but endowed with the opposite complex structure (i.e., if X is already complex, then the crown is the complex double). On top of that $\ell_d = Y$ is connected, i.e. = H.

I wish to point out that the presented method of proof will not generalize. In order to advance one has to understand more about the structure of the minimal cone C; one might speculate that some sort of "H \setminus K -invariant theory" for C could be useful.

Acknow ledgem ent: The origin of this paper traces back to my productive stay at the RIM S in 2005/2006. I am happy to express my gratitude to my form er host Toshiyuki Kobayashi. A lso I would like to thank Toshihiko M atsuki for som e useful intuitive conversations arround this topic.

2. M ain part

Let X = G = K be a Herm itian symmetric space of tube type. This means that there is an Euclidean (or formally real) Jordan algebra V with positive cone W V such that

$$X = V + iW V_C$$
:

The action of G is by fractional linear transform ation and our choice of K is such it is the base point $x_0 = ie w$ is the 2 V the identity element of the Jordan algebra.

It is no loss of generality if we henceforth restrict ourselves to the basic case of G = Sp(n;R) { the more general case is obtained by using standard dictionary which can be found in text books.

For our speci c choice, the Jordan algebra is V = Sym(n;R) and W = V is the cone of positive de nite symmetric matrices. The identity element e is I_n , the n n identity matrix. The group G acts on X by standard fractional linear transformations: $g = \begin{bmatrix} a & b \\ c & d \end{bmatrix} 2$ G with appropriate a;:::;d 2 M (n;R) acts as

$$g z = (az + b)(cz + d)^{\perp}$$
 (z 2 X):

The maxim alcompact subgroup K identi es with U (n) under the standard em bedding

It is then clear that K = U(n) is the stabilizer of $x_0 = iI_n$. In the sequel we consider V_c as the a nepiece of the projective variety L of Lagrangians in C^{2n} ; the embedding is given by

$$V_{\rm C}$$
 ? L; T ? $L_{\rm T} = f(T(v);v) jv 2 C^{\rm n}g$:

It is then clear that $G_c = Sp(n;C)$ acts on L; in symbols: $g = a b \\ c d \\ 2 G_c$ with appropriate a;:::;d 2 M (n;C) acts as

g L = f(av + bw; cv + dw) j(v; w) 2 $C^{n} = C^{2n}g$ (L 2 L):

The space L is hom ogeneous under G $_{\rm C}$. If we choose the base point

$$x_0 \$$
 L₀ = f(iv;v) jv 2 Cⁿg;

then the stabilizer of x_0 in G_C is the Siegel parabolic

 $S^+ = K_C n P^+$ and $P^+ = 1 +$ in u ju 2 V_C :

Thus we have

$$L = G_C \quad I_0 ' \quad G_C = S^+$$
:

Sometimes it is useful to take the conjugate base point $\overline{x_0} = iI_n$. Then the stabilizer of $\overline{L_0}$ in L is the oppposite Siegel parabolic

.

$$S = K_c n P$$
 and $P = 1 + \frac{u}{iu} ju 2 V_c$

and

4

$$L = G_C \quad \overline{I}_{\theta} ' G_C = S :$$

Next we come to the realization of the a ne complexi cation of $X_c = G_c = K_c \cdot W$ e consider the G_c -equivariant embedding

$$X_{C}$$
 ! L L; gK_{C} ? (g b; g b):

It is not hard to see that

$$X_{C} = f(L;L^{0}) 2 L L jL + L^{0} = C^{2n}g;$$

i.e., X_{c} is the a nevariety of pairs of transversal Lagrangians.

Set X = V iW and note that the map z ? \overline{z} identi es X with X in a G-equivarinat, but antiholom orphic manner.

Next we come to the subject matter, the crown domain of X :

$$= X \overline{X} X_{C}$$
:

Let us denote by @ the topological boundary of in X_c . The goal is to resolve @ by a cone bundle over the a ne sym metric space Y = G = H where H = G l(n; R) is the structure group of the Euclidean Jordan algebra V.

We de ne an involution on G by

$$(g) = I_{n,n} g I_{n,n} \quad \text{where} \quad I_{n,n} = I_n \quad I_n$$

The xed point set of is

$$H = a_{t} ja 2 Gl(n;R) = Gl(n;R):$$

We write h for the Lie algebra of H and denote by as well the derived involution on g. The -eigenspace decom position on g shall be denoted by

$$g = h + q$$
 where $q = \begin{pmatrix} 0 & V \\ V & 0 \end{pmatrix}$:

Write $q^+ = \begin{pmatrix} 0 & V \\ 0 & 0 \end{pmatrix}$ and $q = \begin{pmatrix} 0 & 0 \\ V & 0 \end{pmatrix}$ and note that $q = q^+ \quad q$

is the splitting of q into two inequivalent irreducible H -m odules.

The a ne space Y = G = H adm its (up to sign) a unique H -invariant convex open cone C q, containing no a ne lines and consisting of hyperbolic elements. Explicitly:

 $C = \bigcup_{W=0}^{0 W} = W \quad W \quad q^+ \quad q :$

W e form the cone bundle

$$C = G H W$$

and note that there is a natural G-equivariant m ap

P:G_HC!;
$$[g;(y_1;y_2)]$$
? $g(iy;(iy_2)^{\perp})$:

Let us verify that this map is in fact de ned. For that one needs to check that for h 2 H and $y_1; y_2 2 W$, the elements $(h; y_1; y_2)$ and $(1; hy_1 h^t; h^t y_2 h^{-1})$ have the same im age. Indeed,

h $(iy; (iy_2)^{-1}) = (ihy_1h^t; h(iy_2)^{-1}h^t) = (ihy_1h^t; (ih^{-t}y_2h^{-1})^{-1})$

which was asserted.

Lem m a 2.1. The m ap P :C ! is onto.

Proof. Write A for the group of diagonal matrices in G with positive entries. Note that the Lie algebra a of A is a maximal at in $p = g \setminus Sym(2n;R)$. In general, we know that p = Ad(K)a. Furtherm ore, if W_d denotes the diagonal part of W, then $iW_d = A$ x. From G = KAK it mow follows that for any two points (z;w) 2 X there exist a g 2 G such that g $(z;w) = (x;w^0)$ with $w^0 2$ iW_d. As a consequence we obtain that

$$= G (iW_{1}; iI_{n}):$$

C learly the right hand side is contained in the image of P and this nishes the proof.

R em ark 2.2. (a) The map P is not injective. We shall give two di erent arguments for this assertion, beginning with an abstract one. If P were injective, then P establishes an homeoporphism between and $C = G_H C$. In particular is homotopy equivalent to Y = G = H. But we know that is contractible; a contradiction.

More concretely for k 2 K; k \notin 1, the elements [k; (iI_n ; iI_n)] \notin [1; (iI_n ; iI_n)] have the same image in . It should be remarked however, that the map is generically injective.

(b) As H acts properly on C, it follows that G acts properly on the cone-bundle G $_{\rm H}$ C. Further it is not hard to see that the map P is proper.

W e need a more invariant formulation of the map ${\tt P}$. For that, note that the rational map

$$V_{\rm C}$$
 ! $V_{\rm C}$; z ? z ¹

belongs to K . Its extension to L , shall be denoted by s_0 and is given by

 $s_0(L) = f(w;v) 2 C^{2n} j(v;w) 2 Lg:$

A lso, the anti-sym plectic m ap V_c ! V_c ; z ? z has a natural extension to L given by

In this way, we can rewrite P as

P:G_HC!;
$$[g; (y_1; y_2)]$$
 7 g ($iy_1; s_0 (iy_2)$)

and we see that P extends to a continuous m ap

 $P^{\sim}:G_{H} q ! L L; [g; (y_{1}; y_{2})] ? g (iy; s_{0}(iy_{2})):$

We restrict P to G $_{\rm H}$ @C and call this restriction p. It is clear that in p is contained in the boundary of in L $\,$ L.But even m ore is true: the following proposition constitutes a G-equivariant \resolution" of @ .

Proposition 2.3. im p @ and the G-equivariant map

p:G_H @C ! @ ; [g; (y₁; y₂)] 7 g (iy; s₀ (iy₂))

is onto and proper.

Proof. We rest show that imp @. This means that imp X_c . So we have to verify that for $y_1; y_2 \ge cl(W)$ the Lagrangians

$$L_1 = f(iy_1v_iv) jv 2 C^n g$$
 and $L_2 = f(w_iv_2w) jw 2 C^n g$

are transversal. We use the structure group H to bring y_1 in norm al form

$$y_1 = \text{diag}(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}; 0; \dots; 0):$$

p times

Thus $(iy_1v;v) = (w; iy_2w)$ for some $v; w \ge C^n$ means explicitly that

$$(iv_1; iv_2; \dots; iv_p; 0; \dots; 0; v_1; \dots; v_n) = (w_1; \dots; w_n; iy_2 (w))$$
:

We conclude that $w_{p+1} = :::= w_n = 0$. If p = 0, then we are nished. So let us assume that p > 0. But then

$$y_2 = \frac{l_p}{l_p}$$

and this contradicts the fact that y_2 is positive sem ide nite. We turn our attention to the onto-ness of p. First we note that

Next we note that the closure cl(X) in L equals the geodesic com – pacti cation. As a result QX = K (iQW) = K (iQW). Likewise $Q\overline{X} = K$ (iQW). Observe that

We rst show that X $(X \text{ in } p, even m \text{ ore precisely } p(G_H (W (W))) = X (X \text{ . In fact,})$

 $X \quad (\overline{X} = G \quad (\overline{i}I; K \quad \overline{i}OW) = G \quad (\overline{i}IOW)$

and the claim is implied by (2.1). In the manner one veri es that $@X \quad \overline{X} \quad \text{im p.}$

In order to conclude the proof it is now enough to show that p is proper. This is because proper maps are closed and we have already seen that im p contains the dense piece X QX = QXХ 0 . Now to see that p is proper, it is enough to show that inverse in ages of com pact subsets in $[0X \quad 0X] \setminus X_c$ are compact. For the other pieces in 0 this is more or less autom atic: U se that G acts properly on X, resp. \overline{X} which implies that G acts properly on X QX resp.QX X ; likew ise Gacts properly on G_H (W @W) and G _H (@WW). Thus we are about to show that preim ages of compacta in [QX $QX] \setminus X_{C}$ are again compact. But this is more or less im mediate from transversality; I allow myself to skip the details.

R em ark 2.4. For n = 1 the map p is in fact a hom comorphism which we showed in [3]. If n > 1, the map p fails to be injective by the same computational reason shown in the preceeding remark. However, we emphasize that the map is generically injective and that p_{jec} is injective.

References

- S.G indikin and B.K rotz, Complex crowns of Riemannian symmetric spaces and non-compactly causal symmetric spaces, Trans. Amer. M ath. Soc. 354 (8), 3299{3327
- [2] S.G indikin, B.K rotz and G.O lafsson, Holom orphic H-spherical distribution vectors in principal series representations, Invent. m ath. 158, 643 (682 (2004))
- [3] B.Krotz and E.M. Opdam, Analysis on the crown domain, MPIM preprint 2006 (71); to appear in GAFA

8

CORNER VIEW S

M ax-Planck-Institut fur M athematik, V ivatsgasse 7, D-53111 Bonn, email: kroetz@mpim-bonn.mpg.de