# UNIQUENESS OF H-SURFACES IN $\mathbb{H}^2 \times \mathbb{R}$ , $|H| \le 1/2$ , WITH BOUNDARY ONE OR TWO PARALLEL HORIZONTAL CIRCLES

B. NELLI, R. SA EARP, W. SANTOS AND E. TOUBIANA

ABSTRACT. We prove that a H-surface M in  $\mathbb{H}^2 \times \mathbb{R}$ ,  $H \leq \frac{1}{2}$ , inherits the symmetries of its boundary  $\partial M$ , when  $\partial M$  is either a horizontal curve with curvature greater than one or two parallel horizontal curves with curvature greater than one, whose distance is greater or equal to  $\pi$ . Furthermore we prove that the asymptotic boundary of a surface with mean curvature bounded away from zero consists of parts of straight lines, provided it is sufficiently regular.

#### 1. Introduction

An old question in classical Differential Geometry in Euclidean space concerns the influence of the boundary on the behavior of a H-surface. A similar question can also be asked when the ambient space is a homogeneous 3-manifold.

In this paper we consider this problem when the ambient space is the product  $\mathbb{H}^2 \times \mathbb{R}$  and the boundary of the H-surface M is either a horizontal curve (Theorem 2.2) or two parallel horizontal curves (Theorem 3.1) with curvature greater than one. By parallel curves we mean congruent up to a vertical translation. The main result is that M inherits the symmetries of its boundary. In particular, if the boundary curve is a horizontal circle or two parallel horizontal circles with distance greater or equal to  $\pi$ , then M is rotational.

We recall the principal related results in Euclidean and hyperbolic 3-space.

It has been conjectured that a connected compact embedded Hsurface in  $\mathbb{R}^3$  with boundary a round circle is spherical. Of course, this
conjecture can be posed in hyperbolic space  $\mathbb{H}^3$  and in the product
space  $\mathbb{H}^2 \times \mathbb{R}$ . In  $\mathbb{R}^3$  this is still an open problem. F. Braga Brito, W.

<sup>1991</sup> Mathematics Subject Classification. 53C42.

The second and the third authors would like to thank CNPq, PRONEX of Brazil and Accord Brasil-France, for partial financial support.

Meeks, H. Rosenberg and R. Sa Earp proved the conjecture provided the surface is transverse to the plane containing the circle, along the circle ([8]). F. Braga Brito and R. Sa Earp proved that, if the radius of the circle and the mean curvature are equal to one, then the conjecture is true assuming only that the surface is immersed ([6]). In fact they also proved analogous characterizations of a spherical cap for f-surfaces (special Weingarten surfaces) of disk type ([7]). We remark that N. Kapouleas announced examples of immersed H-surfaces with genus  $g \geq 3$ , with boundary a circle ([15]).

In hyperbolic space L. Barbosa and R. Sa Earp proved the following sharp result: if a compact connected immersed surface in  $\mathbb{H}^3$ , with boundary a round circle, has constant mean curvature smaller or equal to one, then it is totally umbilical ([3], [4] and [5]). B. Nelli and H. Rosenberg proved the same result in the embedded case ([18]). In [27], R. Sa Earp and E. Toubiana generalized the above result of [18] to f-surfaces in hyperbolic 3-space satisfying  $f^2 \leq 1$ . They also proved the following: if M has constant mean curvature one, is embedded into  $\mathbb{H}^3$  with  $\partial M = C_1 \cup C_2$ , where  $C_1$  and  $C_2$  are two parallel circles and  $\operatorname{dist}(C_1, C_2)$  great enough, then M is a piece of a catenoid cousin ([27]).

For further results, the reader is referred to [2], [16] and [24] for the Euclidean space and [21] and [26] for the hyperbolic space.

Now, we describe our results for H-surfaces in  $\mathbb{H}^2 \times \mathbb{R}$ .

First, assume M is compact, immersed into  $\mathbb{H}^2 \times \mathbb{R}$ , with mean curvature 1/2, with boundary a round circle in a horizontal slice. Then M is part of the rotational surface with vanishing Abresch-Rosenberg holomorphic quadratic differential Q. An analogous result holds if M has constant mean curvature less than 1/2 (Theorem 2.2 and 5.1). In both situations M is part of an entire rotational vertical graph. More generally, we prove that if  $\partial M$  is a horizontal curve with curvature greater than one, then M is a vertical graph, in particular M has genus zero and inherits the symmetries of its boundary (Theorem 2.2). Assume now M is compact, embedded and has constant mean curvature  $\leq 1/2$ . Assume also that  $\partial M = C_1 \cup C_2$ , where  $C_1$  and  $C_2$  are parallel horizontal curves with curvature greater than one and  $dist(C_1, C_2) \geq \pi$ . Then M inherits the symmetries of  $C_1 \cup C_2$ . Consequently, if  $C_1$  and  $C_2$  are two parallel circles then M is part of an embedded complete rotational H-annulus. The last assertion follows from the geometric classification of rotational H-surfaces (Section 5).

Finally, we consider surfaces M in  $\mathbb{H}^2 \times \mathbb{R}$  which are regular up to the asymptotic boundary and whose mean curvature is bounded away from zero. In [25] R. Sa Earp gives examples suggesting that the asymptotic boundary of such surfaces must lie on a vertical line in  $\partial_{\infty}(\mathbb{H}^2) \times \mathbb{R}$ .

We prove that if the asymptotic boundary  $\partial_{\infty} M$  in  $\partial_{\infty}(\mathbb{H}^2) \times \mathbb{R}$  is  $C^1$  and the surface is  $C^1$  up to  $\partial_{\infty} M$ , then each component of  $\partial_{\infty} M$  is part of a vertical line. As far as we know, the same question is open, if one relax the regularity up to the asymptotic boundary. Observe that in the minimal case many other possibilities can occur ([10], [19], [25] and [28]). In particular, in [19] B. Nelli and H. Rosenberg solve the Dirichlet Problem in  $\mathbb{H}^2 \times \mathbb{R}$  for any Jordan curve  $\gamma$  in the asymptotic boundary  $\partial_{\infty}(\mathbb{H}^2) \times \mathbb{R}$  that is a vertical graph.

The paper is organized as follows. Section 2 and 3 deal with compact H-surfaces with boundary either one curve or two curves, with curvature greater than one. In Section 4, we study the behavior of the asymptotic boundary of a H-surface with strictly positive mean curvature. Finally, in the Appendix we discuss the geometry of rotational H-surfaces for any  $H \in \mathbb{R}$ , since we will need to use these surfaces throughout the whole paper. Rotational surfaces in  $\mathbb{H}^2 \times \mathbb{R}$  have been studied in [1], [14], [17], [23] and [28]. Further results on H-surfaces in  $\mathbb{H}^2 \times \mathbb{R}$  are in [12], [20].

## 2. H-surfaces with boundary a curve with curvature greater than one

In this section we discuss existence and uniqueness of compact H-surfaces,  $|H| \leq 1/2$ , with boundary a curve on a horizontal slice, with curvature greater than one. Then, we study under which conditions a compact H-surface with boundary a planar curve inherits the symmetries of its boundary. In particular we deal with the circle boundary case.

Let  $H \in (0, 1/2]$ . Denote by t the third coordinate in  $\mathbb{H}^2 \times \mathbb{R}$  and by  $\sigma$  the origin in  $\mathbb{H}^2$  (in the disk model of  $\mathbb{H}^2$  we have  $\sigma = 0$ ). We denote by  $S^H$  the simply connected embedded surface in  $\mathbb{H}^2 \times \mathbb{R}$  with constant mean curvature H, invariant by rotation about the axis  $\{\sigma\} \times \mathbb{R}$ , tangent to  $\mathbb{H}^2 \times \{0\}$  at  $\sigma$ . Recall that  $S^H$  is an entire vertical graph (Section 5).

Let us recall the Convex Hull Lemma ([22]). Let K be a compact set in  $\mathbb{H}^2 \times \mathbb{R}$ . For any  $H \in (0, \frac{1}{2}]$ , we define  $\mathcal{F}_K^H$  as follows. A surface B belongs to  $\mathcal{F}_K^H$  if K is contained in the mean convex side of B and if it is obtained from  $S^H$  either by vertical and horizontal translations or by symmetry with respect to a horizontal slice.

Using the maximum principle one can prove the following.

LEMMA 2.1. (Convex Hull Lemma) [22] Let M be a compact surface immersed in  $\mathbb{H}^2 \times \mathbb{R}$  with constant mean curvature  $H \in (0, \frac{1}{2}]$ . Then M is contained in the convex hull of the family  $\mathcal{F}_{\partial M}^H$ .

We recall the Flux Formula ([3], Appendix B in [5] and [13]). Let  $\Sigma$  and Q be two compact, smooth, not necessarily connected surfaces in  $\mathbb{H}^2 \times \mathbb{R}$  such that their boundaries coincide. Assume that there exists a compact domain U in  $\mathbb{H}^2 \times \mathbb{R}$  such that the boundary of U is  $\partial U = \Sigma \cup Q$  and it is orientable. Notice that the boundary of U is smooth except perhaps on  $\partial \Sigma = \partial Q$ .

Let  $n_{\Sigma}$ ,  $n_Q$  be the unit normal fields to  $\Sigma$  and Q respectively that point inside U. Denote by  $\nu$  the unit conormal to  $\Sigma$  along  $\partial \Sigma$ , pointing inside  $\Sigma$ . Finally assume that  $\Sigma$  is a compact surface with constant mean curvature H. Let Y be a Killing vector field in  $\mathbb{H}^2 \times \mathbb{R}$ . One can prove the  $Flux\ Formula$ 

$$\int_{\partial \Sigma} \langle Y, \nu \rangle = 2H \int_{Q} \langle Y, n_{Q} \rangle. \tag{1}$$

In  $\mathbb{H}^2 \times \mathbb{R}$  there is a natural notion of vertical graph ( [12], [22] or [25]): let  $\Omega$  be a subset of  $\mathbb{H}^2$  and let  $u:\Omega \longrightarrow \mathbb{R}$  be a  $C^2$  function. The vertical graph of u is the subset of  $\mathbb{H}^2 \times \mathbb{R}$  given by

$$\{(x, y, t) \in \Omega \times \mathbb{R} \mid t = u(x, y)\}.$$

We choose the unit normal vector field to the graph of u with positive third component and we compute the mean curvature (of the graph) with respect to it.

The graph of a function  $u: \mathbb{H}^2 \longrightarrow \mathbb{R}$  has the function H as mean curvature if and only if u satisfies the following partial differential equation.

$$\operatorname{div}_{\mathbb{H}}\left(\frac{\nabla_{\mathbb{H}}u}{W_{u}}\right) = 2H,\tag{2}$$

where  $\operatorname{div}_{\mathbb{H}}$ ,  $\nabla_{\mathbb{H}}$  are the hyperbolic divergence and gradient respectively and  $W_u = \sqrt{1 + |\nabla_{\mathbb{H}} u|_{\mathbb{H}}^2}$ , being  $|\cdot|_{\mathbb{H}}$  the norm in  $\mathbb{H}^2$ .

Consider the halfspace model for  $\mathbb{H}^2$ , with Euclidean coordinates x, y, y > 0. In this model, equation (2) takes the following form

$$\operatorname{div}\left(\frac{\nabla u}{W_u}\right) = \frac{2H}{y^2},\tag{3}$$

where div is the Euclidean divergence and  $W_u = \sqrt{1 + y^2(u_x^2 + u_y^2)}$ .

The following Theorem is a consequence of a result proved by L. Hauswirth, H. Rosenberg and J. Spruck ([11], Th. 3.2). We sketch a proof for completeness.

THEOREM 2.1. Let  $\Omega$  be a domain in  $\mathbb{H}^2$  such that  $\partial\Omega$  is  $C^{2,\alpha}$  and has curvature greater than one. For any  $H \in [-\frac{1}{2}, \frac{1}{2}]$ , there exists a vertical graph  $G_H$  with constant mean curvature H and boundary  $\partial\Omega$ .

#### Proof.

By formula (3), one has to solve the following Dirichlet problem

$$\begin{cases} F[u] = \operatorname{div}\left(\frac{\nabla u}{W_u}\right) - \frac{2H}{y^2} = 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

In order to prove existence, we use the continuity method. For every  $t \in [0, 1]$ , consider the Dirichlet problem

$$\begin{cases} F^{t}[u] = \operatorname{div}\left(\frac{\nabla u}{W_{u}}\right) - \frac{2Ht}{y^{2}} = 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$
 (4)

Let  $S = \{t \in [0,1] \mid \text{ there exists a solution of } (4)\}$ . Observe that  $u \equiv 0$  is a solution of (4) for t = 0, hence  $0 \in S$ . If one proves that S is open and closed, then  $1 \in S$  and the desired solution is a solution of (4) for t = 1.

That S is open follows from the Implicit Function Theorem. S closed follows from  $C^{2,\alpha}$  a-priori estimates for solutions of (4). By Schauder's theory ([9]),  $C^{2,\alpha}$  a-priori estimates follow from  $C^1$  a-priori estimates. The Convex Hull Lemma guarantees  $C^0$  estimates and boundary gradient bounds on solutions of (4). Therefore, we infer with Theorem 3.1 in [29] that boundary gradient bounds implies interior gradient bounds.

Theorem 2.2. Let M be a compact surface immersed in  $\mathbb{H}^2 \times \mathbb{R}$  with boundary a  $C^{2,\alpha}$  Jordan curve C with curvature greater than one, contained in the slice  $\mathbb{H}^2 \times \{0\}$ . Assume that M has constant mean curvature  $H \in (0, \frac{1}{2}]$ . Then M is a vertical graph (given by Theorem 2.1). In particular M has genus zero and inherits the same symmetries of its boundary. If C is a circle, then M is a part of the simply connected rotational surface containing C with constant mean curvature H.

#### Proof.

Denote by  $P = \{t = 0\}$  the slice containing the boundary curve C and denote by  $\Omega$  the domain in P, bounded by C. Consider the convex hull of the family  $\mathcal{F}_C^H$ . As C has curvature greater than one, for any

point of  $p \in C$  there is a surface of the family  $\mathcal{F}_C^H$  tangent to C at p. By Lemma 2.1, M is contained in the convex hull of the family  $\mathcal{F}_C^H$ , hence  $M \setminus \partial M$  is entirely contained in the vertical cylinder over C and M does not meet P outside  $\Omega$ .

By Theorem 2.1 there exists a graph  $G_H$  on  $\Omega$ , with boundary C and constant mean curvature H. We can choose  $G_H$  to be contained in the halfspace  $t \geq 0$  with mean curvature vector pointing downwards. We will prove that M is contained in one of the two halfspaces determined by the slice P.

First assume that M is embedded. Lift up  $G_H$  to be above M, then move  $G_H$  down towards M: by the maximum principle, one can not touch M till the boundary of M and the boundary of  $G_H$  coincide. Hence M lies below  $G_H$ . Then  $G_H \cup M$  bounds a domain U in  $\mathbb{H}^2 \times \mathbb{R}$  and the mean curvature vector of M points either inside U or outside U.

Assume, by contradiction, that M has points in both halfspaces  $\{t>0\}$ ,  $\{t<0\}$ . By the maximum principle, the mean curvature vector at an highest point of M points downwards, i.e. outside U, while the mean curvature vector at a lowest point of M points upwards, i.e. inside U. This is a contradiction. Then M is contained in a halfspace, say  $\{t>0\}$ ,  $M\cup\Omega$  bounds a domain W in  $\mathbb{H}^2\times\mathbb{R}$  and the mean curvature vector of M points inside W. We already know that M lies below  $G_H$ . Now move down  $G_H$  to be disjoint from M, then lift  $G_H$  up towards M. At a first interior interior contact point the mean curvature vectors of M and  $G_H$  coincide. Contradiction by the maximum principle. Then one can lift  $G_H$  till the boundaries of M and  $G_H$  coincide, that is M is above  $G_H$ . This implies that  $G_H \equiv M$ , as desired.

Now, by applying Alexandrov reflection method with vertical geodesic planes, one obtains that M has all the symmetries of C. In particular, if C is a circle, then M is part of the simply connected rotational surface  $S^H$  containing C. Observe that if M is embedded a simpler alternative argument based on Alexandrov Reflection Principle, using the horizontal slices, yields that M is a vertical graph.

Now let M be an immersed surface (not necessarily embedded). First assume that C is a circle.

By Lemma 2.1, M is contained in the convex hull of the family  $\mathcal{F}_C^H$ . As C is a circle, this convex hull is the domain bounded by the compact part of  $S^H$  containing C, say  $B_C^1$ , and its symmetry with respect to the slice P, say  $B_C^2$ .

Let  $\nu_3^1$ ,  $\nu_3$ ,  $\nu_3^2$  the third components of the inward unit conormal along C of  $B_C^1$ , M,  $B_C^2$  respectively. As M is between  $B_C^1$  and  $B_C^2$ , then at any point of C

$$\nu_3^2 \le \nu_3 \le \nu_3^1. \tag{5}$$

Consider the Flux Formula for M, with  $\Omega$  equals to the planar domain bounded by C, Y = (0,0,1) and  $n_{\Omega} = (0,0,\pm 1)$  according to the orientation given by M. In order to fix ideas, assume that  $n_{\Omega} = (0,0,1)$ . Formula (1) yields

$$\int_{C} \nu_3 = 2HArea(\Omega). \tag{6}$$

Now, consider the Flux Formula for  $B_C^1$  and  $B_C^2$ , with  $\Omega$  equal to the planar domain bounded by C, Y = (0, 0, 1). By formula (1)

$$\int_{C} \nu_3^1 = 2HArea(\Omega) = -\int_{C} \nu_3^2. \tag{7}$$

Then, equalities (6) and (7) yield

$$-\int_{C} \nu_{3}^{2} = \int_{C} \nu_{3} = \int_{C} \nu_{3}^{1}.$$
 (8)

If, in equation (5), the inequalities are strict at every point of C then, one has a contradiction by equation (8).

Then, there is at least one point p in C such that  $\nu_3$  agrees with either  $\nu_3^2$  or  $\nu_3^1$  at p. Therefore, by the boundary maximum principle, M coincides with either  $B_C^2$  or  $B_C^1$ .

If the boundary of M is an embedded curve C with curvature greater than one the proof is analogous. It is enough to replace the caps  $B_C^i$ , i = 1, 2 by the graph  $G_H$  of mean curvature H and boundary C and its symmetry with respect to the slice P.

REMARK 1. Notice that one can prove the analogous results for a surface M, whose mean curvature function H(x,y) satisfies for any  $(x,y) \in \overline{\Omega}$ :  $0 < |H(x,y)| \le \frac{1}{2}$ .

### 3. H-surfaces with boundary two parallel curves with curvature greater than one

We say that  $C_1$  and  $C_2$  are parallel curves, if they are congruent up to a vertical translation.

THEOREM 3.1. Let M be a compact embedded surface in  $\mathbb{H}^2 \times \mathbb{R}$ , with boundary two parallel, embedded  $C^{2,\alpha}$  curves  $C_a \subset P_a = \{t = a\}$ , and  $C_{-a} \subset P_{-a} = \{t = -a\}$ , with curvature greater than one. Assume that M has constant mean curvature H, with  $|H| \leq 1/2$ . Then M is symmetric with respect to the horizontal slice  $\{t = 0\}$ . If  $2a \geq \pi$ , then

M is contained in the closed slab  $\{-a \leq t \leq a\}$ , with  $M \cap (P_a \cup P_{-a}) = C_a \cup C_{-a}$ . Furthermore, M inherits the symmetries of  $C_a \cup C_{-a}$ .

#### Proof.

Let  $D_a$  be the bounded domain in  $P_a$  with boundary  $C_a$ . Let  $\operatorname{ext}(D_a) = P_a \setminus D_a$ . By the Convex Hull Lemma,  $M \cap \operatorname{ext}(D_a) = C_a$  and  $M \cap \operatorname{ext}(D_{-a}) = C_{-a}$ . For any point  $p \in C_a$ , there exists a circle containing  $D_a$  in its interior, tangent to  $C_a$  at p. Let  $Z_p$  be the cylinder over such a circle. Denote by  $Z_p^+$  the mean convex open domain of  $\mathbb{H}^2 \times \mathbb{R}$  bounded by  $Z_p$ . Each  $Z_p$  has mean curvature greater than  $\frac{1}{2}$  hence, by the maximum principle, M is contained in  $\cap_{p \in C_a} Z_p^+$ . Notice that  $\partial(\cap_{p \in C_a} Z_p^+)$  is the cylinder over  $C_a$ . We call it Z and we have that  $M \cap Z = C_a \cap C_{-a}$ . Then, we apply Alexandrov reflection with horizontal slices to infer that the slice  $\{t=0\}$  is a plane of symmetry for M.

Now, assume that  $2a \geq \pi$ .

By Theorem 2.1, there exists a graph  $S_1$  with boundary  $C_a$  with mean curvature H and mean curvature pointing downward, and a graph  $S_2$  with boundary  $C_{-a}$ , with mean curvature H and mean curvature vector pointing upward. By the maximum principle M lies below  $S_1$  and above  $S_2$ , hence  $M \cup S_1 \cup S_2$  is a closed embedded surface, not smooth along  $C_a \cup C_{-a}$ . Let U be the domain in  $\mathbb{H}^2 \times \mathbb{R}$  bounded by  $M \cup S_1 \cup S_2$ . We claim that the mean curvature vector at any point of M points towards U. Consider the family of minimal catenoids (Section 5). Let us recall their shape. For any  $t \in (0, \frac{\pi}{2})$  there exists a catenoid bounded by two circles at infinity at height  $\pm t$ . When  $t \to 0$ , the catenoids tend to the double covering of the slice  $\{t=0\}$  and when  $t \to \frac{\pi}{2}$ , the catenoids tend to infinity. Then, one comes with catenoids from infinity towards M. Let p be the first contact point between M and one of the catenoids. By the maximum principle, the mean curvature vector of M at p points inside U, then it points inside U at any point of M.

Now we prove that M is contained in the closed slab  $\{-a \leq t \leq a\}$ , with  $M \cap \{t = \pm a\} = C_a \cup C_{-a}$ .

Assume by contradiction that M has some points above the slice t = a, and let q be a highest point of M. As  $S_1$  is above M, the mean curvature vector at q points upwards. This gives a contradiction by comparing M with a horizontal slice. Now, assume by contradiction that M has some points below the slice t = -a, and let q be a lowest point of M. As  $S_2$  is below M, the mean curvature vector at q points downwards. And this gives a contradiction by comparing M with a horizontal slice.

By applying the Alexandrov reflection method with vertical geodesic planes, we obtain that M inherits the symmetries of  $C_a \cup C_{-a}$ .

COROLLARY 3.1. Let M be a compact embedded surface in  $\mathbb{H}^2 \times \mathbb{R}$  with boundary two parallel circles. Assume that M has constant mean curvature H, with  $|H| \leq 1/2$ . Let d be the distance of the two boundary curves. If  $d \geq \pi$ , then M is part of an embedded complete rotational surface of constant mean curvature H.

#### Proof.

By the previous Theorem, M is part of a rotational surface with mean curvature H. From the proof of Lemma 3.1, we infer that the mean curvature vector of M points towards the interior of  $M \cup D_a \cup D_{-a}$ , where  $D_a \subset P_a$ ,  $D_{-a} \subset P_{-a}$  are the domains bounded by  $C_a$  and  $C_{-a}$  respectively. By the geometric classification of the rotational surface with constant mean curvature H, M must be a part of an embedded complete rotational surface.

#### 4. Surfaces with nonempty asymptotic boundary

In [25] R. Sa Earp describes many examples of complete H-surfaces with H > 0. When the asymptotic boundary is nonempty, then it consists of parts of straight lines.

THEOREM 4.1. Let M be a surface in  $\mathbb{H}^2 \times \mathbb{R}$  with mean curvature satisfying  $0 < \delta \leq H(p)$  at any point  $p \in M$ . Assume that the asymptotic boundary of M in  $\partial_{\infty}(\mathbb{H}^2) \times \mathbb{R}$  is a  $C^1$  curve and that M is  $C^1$  up to the asymptotic boundary (eventually M has nonempty finite boundary).

Then, each connected component of the asymptotic boundary of M is part of a vertical straight line in  $\partial_{\infty}(\mathbb{H}^2) \times \mathbb{R}$ .

#### Proof.

Let C be a connected component of the asymptotic boundary of M,  $C \subset \partial_{\infty} M \subset \partial_{\infty}(\mathbb{H}^2) \times \mathbb{R}$ . We show that C is vertical at the point p for any  $p \in C$ . We can assume that  $p = 1 \in \partial_{\infty}(\mathbb{H}^2) \times \{0\}$ .

Suppose that C is not vertical at p. Then, an open neighborhood of p in C is a graph  $(e^{i\theta}, t(\theta))$  for  $\theta$  in an open interval around 0  $(p = 1 = e^{i0})$  with t(0) = 0. Let  $\varepsilon > 0$  be a small number. There exists a real number  $\nu = \nu(\varepsilon) > 0$  such that  $|t(\theta)| < \varepsilon/2$  for any  $\theta \in [-\nu, \nu]$ . We set  $p_1 := (e^{i\nu}, t(\nu))$  and  $p_2 := (e^{-i\nu}, t(-\nu))$  and we call  $C_{\varepsilon} \subset C$  the closed subarc of C bounded by  $p_1$  and  $p_2$  and containing p

$$C_{\varepsilon} \subset \partial_{\infty}(\mathbb{H}^2) \times (-\varepsilon/2, \varepsilon/2), \ \partial C_{\varepsilon} = \{p_1, p_2\}.$$

Since M is  $C^1$  up to the asymptotic boundary, there exists a simple arc  $C'_{\varepsilon} \subset M$  with asymptotic boundary  $p_1$  and  $p_2$  such that the connected component  $M_{\varepsilon}$  of  $M \setminus C'_{\varepsilon}$  containing  $C_{\varepsilon}$  in its asymptotic

boundary satisfies  $M_{\varepsilon} \subset \mathbb{H}^2 \times (-\varepsilon, \varepsilon)$ . Summarizing, we have

$$M_{\varepsilon} \subset M \cap \mathbb{H}^2 \times (-\varepsilon, \varepsilon), \ \partial M_{\varepsilon} = C'_{\varepsilon}, \ \partial_{\infty} M_{\varepsilon} = C_{\varepsilon} \subset \partial_{\infty}(\mathbb{H}^2) \times (-\varepsilon/2, \varepsilon/2).$$

Let  $\Pi: \overline{\mathbb{H}^2} \times \mathbb{R} \to \overline{\mathbb{H}^2}$  be the projection on the first two coordinates. For  $\varepsilon$  small enough,  $\Pi(C_{\varepsilon}) \subset \partial_{\infty} \mathbb{H}^2$  is an open arc with end points  $\Pi(p_1)$  and  $\Pi(p_2)$  and containing 1 = p in its interior. Moreover  $\Pi(M_{\varepsilon})$  is an open subset of  $\mathbb{H}^2$  with asymptotic boundary  $\Pi(C_{\varepsilon})$ .

Let  $\Gamma \in \mathbb{H}^2$  be the geodesic (-1,1) and let  $T_s, s > 0$ , be the hyperbolic translation along  $\Gamma$  defined as follows: any point  $x \in \Gamma$  is sent to the point of  $\Gamma$  between x and -1 whose hyperbolic distance from x is s. Then, extend  $T_s$  to  $\mathbb{H}^2 \times \mathbb{R}$  by vertical translation.

Let  $\eta > 0$  be a small number. For s great enough and  $\varepsilon$  small enough, the curve  $T_s(\Pi(C'_{\varepsilon}))$  is inside the open euclidean ball centered at -1 with radius  $\eta$ . Thus  $T_s(\Pi(M_{\varepsilon}))$  is the connected component of  $\mathbb{H}^2 \backslash T_s(\Pi(C'_{\varepsilon}))$  containing 1 in the asymptotic boundary.

The vertical projection of the surface  $T_s(M_{\varepsilon})$  covers a large part of  $\mathbb{H}^2$  in the euclidean sense. Its boundary is the simple arc  $T_s(C'_{\varepsilon})$ . Let  $\gamma_s \subset \partial_{\infty}(\mathbb{H}^2) \times (-\varepsilon, \varepsilon)$  be a  $C^1$  arc with end points  $T_s(p_1)$  and  $T_s(p_2)$  such that  $\gamma_s \cup T_s(C_{\varepsilon})$  is a Jordan curve which projects one to one onto  $\partial_{\infty}(\mathbb{H}^2) \times \{0\}$ .

Finally, let  $R_s \subset \mathbb{H}^2 \times \mathbb{R}$  be any embedded smooth disk, disjoint from the interior of  $T_s(M_{\varepsilon})$ , with finite boundary  $T_s(C'_{\varepsilon})$  and asymptotic boundary  $\gamma_s$ :

$$R_s \cap T_s(M_\varepsilon) = T_s(C'_\varepsilon),$$
  
 $\partial R_s = T_s(C'_\varepsilon),$   
 $\partial_\infty R_s = \gamma_s.$ 

Then  $R_s \cup T_s(M_{\varepsilon})$  is an embedded simply connected surface, it is  $C^0$  along  $T_s(C'_{\varepsilon})$  and smooth everywhere else. The surface  $R_s \cup T_s(M_{\varepsilon})$  separates  $\mathbb{H}^2 \times \mathbb{R}$  in two connected components and its asymptotic boundary is the Jordan curve  $\gamma_s \cup T_s(C_{\varepsilon}) \subset \partial_{\infty}(\mathbb{H}^2) \times [-\varepsilon, \varepsilon]$ .

First, we assume that the mean curvature vector of  $T_s(M_{\varepsilon})$  points towards the connected component containing  $\mathbb{H}^2 \times (\varepsilon, +\infty)$ .

We can assume that  $\delta < 1/2$ . Let us consider the simply connected H-surface  $S^{\delta}$  given in Proposition 5.2 with  $H = \delta$  and with rotational axis equals to the vertical geodesic  $\{0\} \times \mathbb{R}$ . Now, lift up  $S^{\delta}$  to be above  $R_s \cup T_s(M_{\varepsilon})$ , then move  $S^{\delta}$  down. By our construction and the geometry of  $S^{\delta}$  the first contact with  $R_s \cup T_s(M_{\varepsilon})$  will be at an interior point of  $T_s(M_{\varepsilon})$ . This gives a contradiction with the maximum principle. If the mean curvature vector of  $T_s(M_{\varepsilon})$  points towards the other component, one does the same reasoning with the surface obtained from  $S^{\delta}$  by

symmetry with respect to the slice  $\mathbb{H}^2 \times \{0\}$ . Therefore, the asymptotic boundary C is vertical at any point  $p \in C$ .

### 5. Appendix: Geometric behavior of the rotational H-surfaces in $\mathbb{H}^2 \times \mathbb{R}$

In the Appendix we describe in details the geometric behavior of rotational H-surfaces. Our discussion is based on formulae founded in [28]. We recall that rotational surfaces in  $\mathbb{H}^2 \times \mathbb{R}$  have been studied in [1], [14], [17], [23] and [28].

We work with the disk model for  $\mathbb{H}^2$ , so that

$$\mathbb{H}^2 = \{(x, y) \in \mathbb{R}^2, \ x^2 + y^2 < 1\},\$$

and the metric is

$$ds_{\mathbb{H}}^2 = \left(\frac{2}{1 - (x^2 + y^2)}\right)^2 (dx^2 + dy^2).$$

Therefore the product metric on  $\mathbb{H}^2 \times \mathbb{R}$  reads as follows

$$d\tilde{s}^{2} = \left(\frac{2}{1 - (x^{2} + y^{2})}\right)^{2} (dx^{2} + dy^{2}) + dt^{2},$$

where  $(x,y) \in \mathbb{H}^2$  and  $t \in \mathbb{R}$ . We consider the following particular geodesic of  $\mathbb{H}^2$ 

$$\Gamma = \{(x,0), x \in (-1,1)\} \subset \mathbb{H}^2.$$

Up to ambient isometry, we can assume the rotational surfaces are generated by curves in the vertical geodesic plane  $P = \Gamma \times \mathbb{R} \subset \mathbb{H}^2 \times \mathbb{R}$  and that the rotational axis is the vertical geodesic  $R := \{(0,0)\} \times \mathbb{R}$ .

On the geodesic  $\Gamma$  we denote by  $\rho \in \mathbb{R}$  the signed distance to the origin (0,0), thus  $x = \tanh \rho/2$ . Therefore the metric on P is

$$ds^2 = \left(\frac{2}{1-x^2}\right)^2 dx^2 + dt^2 = d\rho^2 + dt^2.$$

Let us consider a curve in P which is a vertical graph:  $c(\rho) = (\rho, \lambda(\rho))$  where  $\lambda$  is a smooth real function defined for  $\rho \geq 0$ . On the rotational surface generated by c we consider the orientation given by the unit normal field pointing up. It is shown in [28] (formula (21)) that the curve c generates a rotational surface with constant mean curvature H if and only if the function  $\lambda$  is given by

$$\lambda(\rho) = \int_{*}^{\rho} \frac{d + 2H \cosh r}{\sqrt{\sinh^2 r - (d + 2H \cosh r)^2}} dr,\tag{9}$$

where d is a real parameter and \* is the minimum such that the condition  $\sinh^2 r - (d + 2H \cosh r)^2 \ge 0$  is satisfied. Using the isometry

 $(x, y, t) \mapsto (x, y, -t)$  we can assume that  $H \ge 0$ . We will analyze consecutively the cases  $H = 0, H \in (0, 1/2]$  and H > 1/2.

PROPOSITION 5.1. (Minimal rotational surfaces) For each  $d \geq 0$  there exists a complete minimal rotational surface  $\mathcal{M}_d$ . The surface  $\mathcal{M}_0$  is the horizontal slice  $\{t=0\}$ . For d>0 the rotational surface  $\mathcal{M}_d$  (called catenoid) is embedded and homeomorphic to an annulus. The distance between the rotational axis and the "neck" of  $\mathcal{M}_d$  is arcsinh d. The asymptotic boundary of  $\mathcal{M}_d$  is two horizontal circles in  $\partial_\infty(\mathbb{H}^2) \times \mathbb{R}$  and the vertical distance between them is a nondecreasing function h(d) satisfying  $\lim_{d\to 0} h(d) = 0$  and  $\lim_{d\to +\infty} h(d) = \pi$ . Therefore  $\mathcal{M}_d$  converges to the double covering of the slice  $\{t=0\}$  when d goes to 0.

Moreover any minimal rotational surface is, up to an ambient isometry, a part of a complete surface  $\mathcal{M}_d$ .

#### Proof.

If the graph of a function  $\lambda$  generates a minimal surface we deduce from formula (9) that

$$\lambda(\rho) = \int_{\operatorname{arcsinh}(d)}^{\rho} \frac{d}{\sqrt{\sinh^2 r - d^2}} dr.$$

Thus we have  $\lambda \equiv 0$  for d=0. For d>0 the function  $\lambda$  is defined for  $\rho \geq \operatorname{arcsinh}(d) > 0$  and the graph c has a vertical tangent at  $\rho = \operatorname{arcsinh} d$ . It is clear that  $\lim_{\rho \to +\infty} \lambda(\rho)$  exists and is finite. Let us call  $\widetilde{c}$  the union of the curve c with its symmetry with respect to the horizontal geodesic  $\{y=0\}$  of P. Therefore  $\widetilde{c}$  is a complete curve which generates a rotational complete and embedded minimal surface homeomorphic to an annulus.

As in formula (41) of [28] we introduce the new coordinate s setting  $ds = \sqrt{1 + \lambda'^2} d\rho$  (s is the arclength of the graph  $c(\rho) = (\rho, \lambda(\rho))$ ). Then using the formulae (49), (36) and (37) of [28] we get

$$\rho(s) = \int_0^s \frac{\sinh t}{\sqrt{(1+d^2)\cosh^2 t - 1}} dt + \operatorname{arcosh} \sqrt{1+d^2},$$
$$= \operatorname{arcosh}(\sqrt{1+d^2}\cosh s).$$

and

$$\lambda \circ \rho(s) = \int_0^s \frac{d}{\sqrt{(1+d^2)\cosh^2 t - 1}} dt,$$

for  $s \geq 0$ . Therefore we have

$$h(d) = 2 \int_0^{+\infty} \frac{d}{\sqrt{(1+d^2)\cosh^2 t - 1}} dt.$$

Consider the positive function defined by  $f(t,d) = \frac{d}{\sqrt{(1+d^2)\cosh^2 t - 1}}$  for  $t,d \geq 0$ . Let  $d_0 > 0$  be any positive real number. Clearly the integral h(d) is convergent for any  $d \in [d_0, +\infty)$ . Moreover we have

$$\frac{\partial}{\partial d}f(t,d) = \frac{\sinh t}{((1+d^2)\cosh^2 t - 1)^{3/2}}.$$

We deduce that the integral

$$\int_0^{+\infty} \frac{\partial}{\partial d} f(t, d) dt,$$

is uniformly convergent for  $d \geq d_0 > 0$ . Consequently the function h(d) is differentiable on  $[d_0, +\infty)$  and

$$h'(d) = 2 \int_0^{+\infty} \frac{\partial}{\partial d} f(t, d) dt, \tag{10}$$

$$=2\int_0^{+\infty} \frac{\sinh t}{((1+d^2)\cosh^2 t - 1)^{3/2}} dt.$$
 (11)

As this is true for  $d \ge d_0 > 0$  for any  $d_0 > 0$ , then h is differentiable for d > 0 and its derivative is given by (11). We deduce that h(d) is nondecreasing and we have  $\lim_{d\to 0} h(d) = 0$ .

Finally, from the inequalities

$$\frac{d}{\sqrt{(1+d^2)}} \frac{1}{\cosh t} \le \frac{d}{\sqrt{(1+d^2)\cosh^2 t - 1}} \le \frac{1}{\cosh t},$$

for any d, t > 0, we get

$$\lim_{d \to +\infty} h(d) = 2 \int_0^{+\infty} \lim_{d \to +\infty} \frac{d}{\sqrt{(1+d^2)\cosh^2 t - 1}} dt,$$

$$= 2 \int_0^{+\infty} \frac{dt}{\cosh t},$$

$$= 2 \int_0^{+\infty} \frac{du}{u^2 + 1}, \quad (u = \sinh t),$$

$$= \pi.$$

This concludes the proof.

For later use we define the functions  $g(\rho)$  and  $f(\rho)$  setting for  $d \in \mathbb{R}$  and H > 0,

$$g(\rho) = d + 2H \cosh \rho,$$
  

$$f(\rho) = \sinh^2 \rho - (d + 2H \cosh \rho)^2,$$
  

$$= (1 - 4H^2) \cosh^2 \rho - 4dH \cosh \rho - 1 - d^2,$$

so that  $\lambda'(\rho) = g(\rho)/\sqrt{f(\rho)}$ .

Lemma 5.1. Assume 0 < H < 1/2. We have  $f(\rho) \ge 0$  if and only if  $\cosh \rho \ge \frac{2dH + \sqrt{1 - 4H^2 + d^2}}{1 - 4H^2}$ . Let  $\rho_1 \ge 0$  such that  $\cosh \rho_1 = \frac{2dH + \sqrt{1 - 4H^2 + d^2}}{1 - 4H^2}$ , then  $f(\rho_1) = 0$  and  $\rho_1 = 0$  if and only if d = -2H.

- (1) If d > -2H, then  $\frac{-d}{2H} < \cosh \rho_1$ . Consequently the function  $\lambda$  is nondecreasing for  $\rho \ge \rho_1 > 0$  and has a nonfinite derivative at  $\rho_1$ .
- (2) If d = -2H, then  $\lambda'(\rho) = \frac{2H\sqrt{\cosh\rho-1}}{\sqrt{(1-4H^2)\cosh\rho+4H^2+1}}$ . Therefore the function  $\lambda$  is defined for  $\rho \geq 0$ , it has a zero derivative at 0 and is nondecreasing for  $\rho > 0$ .
- (3) If d < -2H, then there exists  $\rho_0 > \rho_1 > 0$  such that  $\frac{-d}{2H} = \cosh \rho_0$ . Consequently the function  $\lambda$  is defined for  $\rho \geq \rho_1 > 0$  with a nonfinite derivative at  $\rho_1$ , it is nonincreasing for  $\rho_1 < \rho < \rho_0$ , has a zero derivative at  $\rho_0$  and it is nondecreasing for  $\rho > \rho_0$ .
- (4) For any d we have  $\lim_{\rho \to +\infty} \lambda(\rho) = +\infty$ .

Next Lemma, is analogous to Lemma 5.1 in the case H=1/2. We observe that, in this case, the set  $\{\rho>0\mid f(\rho)>0\}$  is nonempty if and only if d<0.

LEMMA 5.2. Assume H=1/2 and d<0. Then  $f(\rho)\geq 0$  if and only if  $\cosh\rho\geq \frac{1+d^2}{-2d}$ . Let  $\rho_1\geq 0$  such that  $\cosh\rho_1=\frac{1+d^2}{-2d}$ , then  $f(\rho_1)=0$  and  $\rho_1=0$  if and only if d=-1.

- (1) If  $d \in (-1,0)$ , then  $\frac{-d}{2H} < \cosh \rho_1$ . Consequently the function  $\lambda$  is nondecreasing for  $\rho \ge \rho_1 > 0$  and has a nonfinite derivative at  $\rho_1$ .
- (2) If d = -1, then  $\lambda'(\rho) = \frac{1}{\sqrt{2}} \sqrt{\cosh \rho 1}$ . Therefore the function  $\lambda$  is defined for  $\rho \geq 0$ , it has a zero derivative at 0 and is nondecreasing for  $\rho > 0$ .
- (3) If d < -1 there exists  $\rho_0 > \rho_1 > 0$  such that  $\frac{-d}{2H} = \cosh \rho_0$ . Consequently the function  $\lambda$  is defined for  $\rho \geq \rho_1 > 0$  with a nonfinite derivative at  $\rho_1$ , it is nonincreasing for  $\rho_1 < \rho < \rho_0$ , has a zero derivative at  $\rho_0$  and it is nondecreasing for  $\rho > \rho_0$ .

(4) For any d we have  $\lim_{\rho \to +\infty} \lambda(\rho) = +\infty$ .

The proof of Lemma 5.1 and 5.2 is a straightforward computation taking into account Formula (9). As a consequence of Lemma 5.1 and 5.2 we have the following results.

PROPOSITION 5.2. (Rotational H-surfaces with  $|H| \le 1/2$ )

Assume  $0 < H \le 1/2$ . There exists a one-parameter family  $\mathcal{H}_d$ ,  $d \in \mathbb{R}$  for H < 1/2 and d < 0 for H = 1/2, of complete rotational H-surfaces.

- (1) For d > -2H, the surface  $\mathcal{H}_d$  is a properly embedded annulus, symmetric with respect to the slice  $\{t = 0\}$ , the distance between the "neck" and the rotational axis  $R = \{(0,0)\} \times \mathbb{R}$  is  $\operatorname{arcosh}(\frac{2dH + \sqrt{1 4H^2 + d^2}}{1 4H^2})$  for H < 1/2 and  $\operatorname{arcosh}(\frac{1 + d^2}{2d})$  for H = 1/2.
- (2) For d = -2H, the surface  $\mathcal{H}_{-2H}$  is an entire vertical graph, denoted by  $S^H$ . Moreover  $S^H$  is contained in the halfspace  $\{t \geq 0\}$  and it is tangent to the slice  $\mathbb{H}^2 \times \{0\}$  at the point (0,0,0).
- (3) For d < -2H, the surface  $\mathcal{H}_d$  is a properly immersed (and nonembedded) annulus, it is symmetric with respect to the slice  $\{t=0\}$ , the distance between the "neck" and the rotational axis R is  $\operatorname{arcosh}(\frac{2dH+\sqrt{1-4H^2+d^2}}{1-4H^2})$  for H < 1/2 and  $\operatorname{arcosh}(\frac{1+d^2}{-2d})$  for H = 1/2.
- (4) In each of the previous case the surface is unbounded in the t-coordinate. When d tends to -2H with either d > -2H or d < -2H, then the surfaces  $\mathcal{H}_d$  tends towards the union of  $S^H$  and its symmetry with respect to the slice  $\{t = 0\}$ . Furthermore, any rotational H-surface with  $0 < H \le 1/2$  is, up to an ambient isometry, a part of a surface of the family  $\mathcal{H}_d$ .

#### Proof.

The result is a straightforward consequence of Lemma 5.1 and 5.2. For d = -2H,  $\mathcal{H}_{-2H}$  is the rotational surface generated by the graph of the function  $\lambda$ .

For  $d \neq -2H$ , let  $\gamma$  be the union of the graph of  $\lambda$  joint with its symmetry with respect to the slice  $\{t = 0\}$ . Then  $\mathcal{H}_d$  is the rotational surface generated by the curve  $\gamma$ .

Observe that, for H > 1/2, the set  $\{\rho > 0 \mid f(\rho) > 0\}$  is nonempty if and only if  $d < -\sqrt{4H^2 - 1}$ .

- LEMMA 5.3. Let H and d satisfying H > 1/2 and  $d < -\sqrt{4H^2 1}$ . Then, there exist two numbers  $0 \le \rho_1 < \rho_2$  such that  $\cosh \rho_1 = \frac{2dH + \sqrt{1 - 4H^2 + d^2}}{1 - 4H^2}$  and  $\cosh \rho_2 = \frac{2dH - \sqrt{1 - 4H^2 + d^2}}{1 - 4H^2}$ . Therefore,  $f(\rho) > 0$  if and only if  $\rho_1 < \rho < \rho_2$  and  $f(\rho_1) = f(\rho_2) = 0$ .
  - (1) If d < -2H, then  $\rho_1 > 0$  and there exists a unique number  $\rho_0 \in (\rho_1, \rho_2)$  satisfying  $g(\rho_0) = 0$ . Furthermore  $g \leq 0$  on  $[\rho_1, \rho_0)$  and  $g \geq 0$  on  $(\rho_0, \rho_2]$ . Consequently, the function  $\lambda$  is defined on  $[\rho_1, \rho_2]$ , has a nonfinite derivative at  $\rho_1$  and  $\rho_2$ , has a zero derivative at  $\rho_0$ , is nonincreasing on  $(\rho_1, \rho_0)$  and nondecreasing on  $(\rho_0, \rho_2)$ .
  - (2) If d = -2H, then  $\rho_1 = 0$  and  $\lambda'(\rho) = \frac{2H\sqrt{\cosh\rho-1}}{\sqrt{(1-4H^2)\cosh\rho+4H^2+1}}$ . Consequently, the function  $\lambda$  is defined on  $[0, \rho_2]$ , is nondecreasing, has a zero derivative at 0 and a nonfinite derivative at  $\rho_2$ .
  - (3) If  $-2H < d < -\sqrt{4H^2 1}$ , then  $\rho_1 > 0$  and  $g \ge 0$  on  $[\rho_1, \rho_2]$ . Therefore the function  $\lambda$  is defined on  $[\rho_1, \rho_2]$ , is nondecreasing and has nonfinite derivative at  $\rho_1$  and  $\rho_2$ .

The proof of Lemma 5.3 follows from formula (9) by a computation. In the following Proposition we assume the notations of Lemma 5.3

PROPOSITION 5.3. (Rotational surfaces with H > 1/2)

Assume H > 1/2. There exists a one-parameter family  $\mathcal{D}_d$  of complete rotational H-surfaces,  $d \leq -\sqrt{4H^2 - 1}$ .

- (1) For d < -2H, the surface  $\mathcal{D}_d$  is an immersed (and nonembedded) annulus, invariant by a vertical translation and is contained in the closed region bounded by the two vertical cylinders  $\rho = \rho_1$  and  $\rho = \rho_2$ . Furthermore  $\rho_1 \to +\infty$  and  $\rho_2 \to +\infty$  when  $d \to -\infty$  and  $\rho_1 \to 0$  and  $\rho_2 \to \operatorname{arcosh}(\frac{4H^2+1}{4H^2-1})$  when  $d \to -2H$ . Such surfaces are analogous to the nodoids of Delaunay in  $\mathbb{R}^3$ .
- (2) For d = -2H, the surface  $\mathcal{D}_{-2H}$  is an embedded sphere and the maximal distance from the rotational axis is  $\rho_2 = \operatorname{arcosh}(\frac{4H^2+1}{4H^2-1})$ .
- (3) For  $-2H < d < -\sqrt{4H^2 1}$ , the surface  $\mathcal{D}_d$  is an embedded annulus, invariant by a vertical translation and is contained in the closed region bounded by the two vertical cylinders  $\rho = \rho_1$  and  $\rho = \rho_2$ . Furthermore  $\rho_1 \to 0$  and  $\rho_2 \to \operatorname{arcosh}(\frac{4H^2 + 1}{4H^2 1})$  when  $d \to -2H$  and both  $\rho_1, \rho_2 \to \operatorname{arcosh}(\frac{2H}{\sqrt{4H^2 1}})$  when  $d \to -\sqrt{4H^2 1}$ . Moreover  $\rho_1 < \operatorname{arcosh}(\frac{2H}{\sqrt{4H^2 1}}) < \rho_2$ . Such surfaces are analogous to the undoloids of Delaunay in  $\mathbb{R}^3$ .
- (4) For  $d = -\sqrt{4H^2 1}$ , the surface  $\mathcal{D}_{-\sqrt{4H^2 1}}$  is the vertical cylinder over the circle with hyperbolic radius  $\operatorname{arcosh}(\frac{2H}{\sqrt{4H^2 1}})$ .

#### Proof.

For  $d = -\sqrt{4H^2 - 1}$  we get the limit case of a vertical cylinder given by  $\cosh \rho = \frac{2H}{\sqrt{4H^2 - 1}}$ . A straightforward computation shows the mean curvature of such a cylinder is H.

For the other cases the proof is a straightforward consequence of Lemma 5.3. Let  $\gamma$  be the union of the graph of  $\lambda$  joint with its symmetries with respect to the horizontal slices on which  $\lambda$  is vertical. When d = -2H,  $\gamma$  is a compact arc and for  $d \neq -2H$ ,  $\gamma$  is periodic and complete, embedded only when d > -2H.

From Proposition 5.1, 5.2, 5.3 and Proposition 26 in [28], we infer the following classification of rotational H-surfaces with vanishing Abresch-Rosenberg holomorphic quadratic differential. The classification of H-surfaces with vanishing Abresch-Rosenberg holomorphic quadratic differential is established in [1].

Theorem 5.1. Let M be a rotational H-surface,  $H \ge 0$ , with vanishing Abresch-Rosenberg holomorphic quadratic differential. We have up to congruence:

- (1) If H = 0 then M is a slice  $\mathbb{H}^2 \times \{t\}$ .
- (2) If H > 1/2 then M is an embedded two-sphere.
- (3) If H = 1/2 then M is the entire vertical graph  $S^{1/2}$ .
- (4) If H < 1/2 then M is either the entire vertical graph  $S^H$  or the embedded annulus  $\mathcal{H}_{2H}$ .

#### References

- [1] U. ABRESCH AND H. ROSENBERG. A Hopf differential for constant mean curvature surfaces in  $\mathbb{S}^2 \times \mathbb{R}$  and  $\mathbb{H}^2 \times \mathbb{R}$ , Acta Math. 193, N° 2, 141-174 (2004).
- [2] J. L. Barbosa: Constant mean curvature surfaces bounded by a plane curve. Mat. Contemp. 1, 3-15 (1991).
- [3] J. L. Barbosa and R. Sa Earp: New results on prescribed mean curvature hypersurfaces in Space Forms. An Acad. Bras. Ci. 67, No 1,1-5 (1995).
- [4] J. L. BARBOSA AND R. SA EARP: Prescribed mean curvature hypersurfaces in  $H^{n+1}(-1)$  with convex planar boundary, I. Geom. Dedicata, **71**, 61-74 (1998).
- [5] J.L. BARBOSA AND R. SA EARP: Prescribed Mean Curvature Hypersurfaces in  $\mathbb{H}^{n+1}$  with Convex Planar Boundary, II. Séminaire de Théorie Spectrale et Géométrie, Grenoble 16 (1998) 43-79.
- [6] F. Braga Brito and R. Sa Earp: Geometric configurations of constant mean curvature surfaces with planar boundary. An. Acad. Bras. Ci. **6**3,No 1, 5-19 (1991).

- [7] F. Braga Brito and R. Sa Earp: On the Structure of certain Weingarten surfaces with boundary a circle. An. Fac. Sci. Toulouse VI, No 2, 243-255 (1997).
- [8] F. Brito, W. Meeks III, H. Rosenberg and R. Sa Earp: Structure Theorems for Constant Mean Curavture Surfaces Bounded by a Planar Curve. Indiana Univ. Math. Jour. 40 (1) 333-343 (1991).
- [9] D. GILBARG AND N.S. TRUDINGER: Elliptic Partial Differential Equations of Second Order. Springer- Verlag (1998).
- [10] L. Hauswirth: Generalized Riemann examples in three-dimensional manifolds. Pacific Journal of Math. 224, No. 1, 91-117 (2006).
- [11] L. HAUSWIRTH, H. ROSENBERG AND J. SPRUCK: Infinite boundary value problems for constant mean curvature graphs in  $\mathbb{H}^2 \times \mathbb{R}$  and  $\mathbb{S}^2 \times \mathbb{R}$ . http://www.math.jhu.edu/~js/jsfinal.pdf
- [12] L. HAUSWIRTH, R. SA EARP AND E. TOUBIANA: Associate and conjugate minimal immersions in  $\mathbb{M} \times \mathbb{R}$ . Preprint.
- [13] D. HOFFMAN, J. DE LIRA AND H. ROSENBERG: Constant Mean Curvature Surfaces in  $M^2 \times \mathbb{R}$ . Trans. Amer. Math. Soc. 358 n.2 (2006), 491-507.
- [14] W.T. HSIANG AND W.Y. HSIANG: On the uniqueness of isoperimetric solutions and embedded soap bubbles in noncompact symmetric spaces I. Invent. Math 98, 39-58 (1989).
- [15] N. KAPOULEAS: Compact constant mean curvature surfaces in Euclidean three-space. J. Diff. Geom. 33, 683-715 (1991).
- [16] M. Koiso: Symmetry of hypersurfaces of constant mean curvature with symmetric boundary. Math. Z. 191, 567-574 (1986).
- [17] MONTALDO AND I. ONNIS: Invariant CMC surfaces in  $\mathbb{H}^2 \times \mathbb{R}$ . Glasg. Math. J. 46 No. 2, 311-321 (2004).
- [18] B. Nelli and H. Rosenberg: Some remarks on embedded hypersurfaces in Hyperbolic Space of constant mean curvature and spherical boundary. Ann. Glob. An. and Geom. 13, 23-30, (1995).
- [19] B. Nelli and H. Rosenberg: Minimal surfaces in  $\mathbb{H}^2 \times \mathbb{R}$ . Bull. Braz. Soc. 33, 263-292 (2002).
- [20] B. Nelli and H. Rosenberg: Global properties of constant mean curvature surfaces in  $\mathbb{H}^2 \times \mathbb{R}$ . Pacific Journ. Math. **226**, No 1, 137-152 (2006).
- [21] B. Nelli and R. Sa Earp: Some Properties of Hypersurfaces of Prescribed Mean Curvature in  $H^{n+1}$ . Bull. Sc. Math. 120, No 6, 537-553 (1996).
- [22] B. Nelli and R. Sa Earp: Vertical Ends of Constant Mean Curvature  $H = \frac{1}{2}$  in  $\mathbb{H}^2 \times \mathbb{R}$ . Preprint.
- [23] R.H.L. Pedrosa and M. Ritore: Isoperimetric domains in the Riemannian product of a circle with a simply connected space form and applications to free boundary problems. Indiana Univ. Math. J. 48 No. 4, 1357-1394 (1999).

- [24] H. ROSENBERG AND R. SA EARP: Some Structure Theorems for Complete Constant Mean Curvature Surfaces with Boundary a Convex Curve. Proc. Amer. Math. Soc. 113, No 4, 1045-1053 (1991).
- [25] R. SA EARP: Parabolic and Hyperbolic Screw motion in  $\mathbb{H}^2 \times \mathbb{R}$ . http://www.mat.puc-rio.br/ $\sim$ earp/pscrew.pdf, to appear in Journ. Austra. Math. Soc..
- [26] R. SA EARP, E. TOUBIANA: Some Applications of Maximum Principle to Hypersurfaces in Euclidean and Hyperbolic Space. New Approaches in Nonlinear Analysis, Themistocles M. Rassias Editor, Hardonic Press (1999) 183-202.
- [27] R. SA EARP AND E. TOUBIANA: Symmetry of properly embedded special Weingarten surfaces in  $H^3$ . Trans. Amer. Math. Soc. **352**, No 12, 4693-4711 (1999).
- [28] R. SA EARP AND E. TOUBIANA: Screw Motion Surfaces in  $\mathbb{H}^2 \times \mathbb{R}$  and  $\mathbb{S}^2 \times \mathbb{R}$ , Illinois Jour. of Math. 49 n.4 (2005) 1323-1362.
- [29] J. Spruck: Interior Gradient Estimates and Existence Theorems for Constant Mean Curvature Graphs in  $M \times \mathbb{R}$ . http://www.math.jhu.edu/ $\sim$ js/grad2.pdf

DIPARTIMENTO DI MATEMATICA, UNIVERSITÁ DI L'AQUILA, VIA VETOIO - LOC. COPPITO, 67010 (L'AQUILA), ITALY E-mail address: nelli@univaq.it

DEPARTAMENTO DE MATEMÁTICA, PONTIFÍCIA UNIVERSIDADE CATÓLICA DO RIO DE JANEIRO, RIO DE JANEIRO, 22453-900 RJ, BRAZIL

E-mail address: earp@mat.puc-rio.br

Universidade Federal do Rio de Janeiro, Instituto de Matemtica, Av. Brigadeiro Trompowsky, s/n Cidade Universitria, Ilha do Fundão, Caixa Postal 68530, 21945-970 Rio de Janeiro RJ, Brazil

E-mail address: walcy@im.ufrj.br

Institut de Mathématiques de Jussieu, Université Paris VII, Denis Diderot, Case 7012, 2 Place Jussieu, 75251 Paris Cedex 05, France *E-mail address*: toubiana@math.jussieu.fr