CONJECTURE OF TITS TYPE FOR COMPLEX VARIETIES AND THEOREM OF LIE-KOLCHIN TYPE FOR A CONE

JONGHAE KEUM, KEIJI OGUISO, AND DE-QI ZHANG

ABSTRACT. First, we shall formulate and prove Theorem of Lie-Kolchin type for a cone and derive some algebro-geometric consequences. Next, inspired by a recent result of Dinh and Sibony we pose a conjecture of Tits type for a group of automorphisms of a complex variety and verify its weaker version. Finally, applying Theorem of Lie-Kolchin type for a cone, we shall confirm the conjecture of Tits type for complex tori, hyperkähler manifolds, surfaces, and minimal threefolds.

1. INTRODUCTION

We often study a group G of automorphisms of a projective variety X through its action on cohomological spaces such as on the Néron-Severi group NS(X). Let

$$G^* = \operatorname{Im}(G \longrightarrow \operatorname{GL}(\operatorname{NS}(X))).$$

Then Tits' Theorem [Ti] implies that

- (i) either G^* contains a subgroup isomorphic to $\mathbf{Z} * \mathbf{Z}$ (highly noncommutative);
- (ii) or G^* contains a connected solvable subgroup of finite index.

Here, connected means that its Zariski closure in $GL(NS(X)_{\mathbb{C}})$ is connected.

This property already provides a useful tool in studying groups of automorphisms of projective varieties (see [Og2], [Zh1] for some concrete applications). Another important feature of G^* is that it preserves the ample cone $\operatorname{Amp}(X)$ and the nef cone $\overline{\operatorname{Amp}}(X)$, the closure of the ample cone, both of which encode much information on geometry of X.

Keeping these two features in mind, we shall prove first the following Theorem of Lie-Kolchin type (see also Theorem 2.1 and Corollary 2.3), and derive some direct consequences for groups of automorphisms of projective varieties or compact Kähler manifolds (Theorems 3.1-3.5):

Theorem 1.1. (Theorem of Lie-Kolchin type for a cone) Let V be a finite dimensional real vector space, and let $C \neq \{0\}$ be a strictly convex closed cone of V. Let G be a connected (in the sense above) solvable subgroup of GL(V) such that $G(C) \subseteq C$. Then G has a common eigenvector in the cone C.

This theorem is purely algebraic and *a priori* has nothing to do with algebraic geometry or complex geometry. In our proof, we use Birkhoff-Perron-Frobenius' Theorem [Bir] and the proof of Dinh-Sibony [DS] for the case where G is abelian.

²⁰⁰⁰ Mathematics Subject Classification. 14J50, 14E07, 32H50.

Key words and phrases. automorphism, hyperkähler, Lie-Kolchin, Tits, topological entropy.

The first named author was supported by Korea Research Foundation Grant (KRF-). The second named author was supported by JSPS.

In the same paper, Dinh and Sibony proved the following interesting result:

Theorem 1.2. ([DS]) Let M be an n-dimensional compact Kähler manifold and let G be an abelian subgroup of Aut (M) such that each element of $G \setminus \{id\}$ is of positive entropy. Then G is a free abelian group of rank at most n-1. Furthermore, the rank estimate is optimal.

See for instance [ibid] or (4.1) for the notion of entropy of an automorphism.

In view of Tits' Theorem and Dinh-Sibony's Theorem, it is natural to pose the following conjecture:

Conjecture 1.3. (Conjecture of Tits type) Let X be an n-dimensional compact Kähler manifold or an n-dimensional complex projective variety with at most rational \mathbf{Q} -factorial singularities. Let G be a subgroup of Aut (X). Then, one of the following two assertions holds:

- (1) G contains a subgroup isomorphic to $\mathbf{Z} * \mathbf{Z}$.
- (2) There is a finite index subgroup G_1 of G such that the subset

 $N(G_1) = \{ g \in G_1 \mid g \text{ is of null entropy} \}$

of G_1 is a normal subgroup of G_1 and the quotient group $G_1/N(G_1)$ is a free abelian group of rank at most n-1.

It turns out that the crucial part of the conjecture is the rank estimate in the statement (2), as a weaker version of the conjecture (Theorem 4.3) can be easily verified. We shall also verify the conjecture for some basic cases, say, for surfaces (Theorem 4.4), hyperkähler manifolds (Theorem 4.5), complex tori (Theorem 4.6), and minimal threefolds (Theorem 5.1). Except for the case of complex tori, Theorem of Lie-Kolchin type (Theorem 1.1) plays a crucial role in our proof.

It would be interesting to verify/deny the conjecture for higher dimensional Calabi-Yau manifolds and manifolds with Kodaira dimension $-\infty$.

Acknowledgement. We would like to express our thanks to Professor Yujiro Kawamata for his encouragement and valuable suggestions. The first two authors would like to thank Department of Mathematics, National University of Singapore, and staff members there for financial support and hospitality. The main part of this work was done during their stay in Singapore.

2. Theorem of Lie-Kolchin type for a cone

(2.1) We call a group virtually solvable if it has a solvable subgroup of finite index. A solvable subgroup G of $GL(n, \mathbb{C})$ is called *connected* if its Zariski closure \overline{G} in $GL(n, \mathbb{C})$ is connected. Note that given a virtually solvable subgroup G of $GL(n, \mathbb{C})$, one can always find a connected solvable, finite index subgroup H of G. Indeed, choose a solvable finite index subgroup G_1 of G and take the identity component $\overline{G_1}^0$ of the Zariski closure $\overline{G_1}$. Then the group $H := \overline{G_1}^0 \cap G_1$ is a connected solvable finite index subgroup of G.

(2.2) Let V be an r-dimensional real vector space. We regard V as an r-dimensional euclidean space with the natural topology. Let C be a subset of V. We call C a strictly convex closed cone of V if C is closed in V, closed under addition and multiplication by a non-negative scalar, and contains no 1-dimensional linear space.

(2.3) In this section, we shall prove the following theorem:

Theorem 2.1. (Theorem of Lie-Kolchin type for a cone) Let V be an rdimensional real vector space, and let $C \neq \{0\}$ be a strictly convex closed cone of V. Let G be a virtually solvable subgroup of GL(V) such that $G(C) \subseteq C$. Then there are a finite-index subgroup H of G and a real vector $v \in C \setminus \{0\}$ such that $H(\mathbf{R}_{\geq 0}v) = \mathbf{R}_{\geq 0}v$. In other words, a suitable finite-index subgroup of G has a common eigenvector in the cone C. Moreover, if G is connected and solvable, then one may choose H = G.

Before starting the proof, we recall several relevant results and state one special but useful version.

Remark 2.2.

(1) Tits' Theorem [Ti] states that any subgroup G of $\operatorname{GL}(V_{\mathbf{C}})$ is either virtually solvable or contains a free subgroup isomorphic to $\mathbf{Z} * \mathbf{Z}$. In the latter case, the conclusion in Theorem 2.1 is not true in general even in the case $V = \mathbf{Z}^r \otimes \mathbf{R}$ and $G \subset \operatorname{GL}(\mathbf{Z}^r)$. In this sense, our restriction on the group G is reasonable.

(2) The original version of Lie-Kolchin Theorem (see eg. [Hu]) states that any connected solvable subgroup G of $\operatorname{GL}(V_{\mathbf{C}})$ has a common eigenvector. Note that it is, in general, far from being real, even in the case $V = \mathbf{Z}^r \otimes \mathbf{R}$ and $G \subset \operatorname{GL}(\mathbf{Z}^r)$.

(3) Birkhoff-Perron-Frobenius' Theorem [Bir] states that any $g \in GL(V)$ such that $g(C) \subset C$ has an eigenvector v_g in C. Note that g acts on the real vector space $W = \langle C \rangle$. So, one can choose v_g so that $g(v_g) = \rho(g|W) v_g$, where $\rho(g|W)$ is the spectral radius of g|W (see [ibid]).

(4) Dinh-Sibony ([DS] Proposition 4.1) proved Theorem 2.1 when G is abelian. So, our theorem is not only a refinement of Lie-Kolchin Theorem but also an optimal generalization of Birkhoff-Perron-Frobenius' Theorem [Bir] and Dinh-Sibony's version [DS] in view of Tits' Theorem.

For algebro-geometric applications, the following arithmetical version will be most useful:

Corollary 2.3. Let L be a free **Z**-module of finite rank $r \ge 1$, and let $C \ne \{0\}$ be a strictly convex closed cone of $L_{\mathbf{R}} := L \otimes \mathbf{R}$. Let H be a connected solvable subgroup of Aut $L = \operatorname{GL}(L)$ such that $H(C) \subseteq C$. Then there is a real vector $v \in C \setminus \{0\}$ such that $H(\mathbf{R}_{>0}v) = \mathbf{R}_{>0}v$.

This version together with the next lemma or its proof will be frequently used in the sequel. In this lemma, the condition "the action of H is defined over \mathbb{Z} " is very crucial.

Lemma 2.4. Let L be a free **Z**-module of finite rank $r \ge 1$ and let H be a connected solvable subgroup of GL(L). Then, the following set

$$N(H) := \{h \in H \,|\, \rho(h) = 1\}$$

is a normal subgroup of H and H/N(H) is a free abelian group of rank at most r-1. Here, $\rho(h)$ is the spectral radius of h on $L_{\mathbf{C}}$.

Proof. By Lie-Kolchin Theorem, H is a subgroup of the group T(r) of upper triangle matrices of $GL(r, \mathbb{C})$ under the identification $GL(L_{\mathbb{C}}) = GL(r, \mathbb{C})$ with respect to a suitable basis of $L_{\mathbb{C}}$.

Regarding $H \subset T(n)$, we write the (i, i)-th entry of the matrix $h \in H$ by $\chi_i(h)$ $(1 \leq i \leq r)$. Note that $\{\chi_i(h)\}_{i=1}^r$ is the set of eigenvalues of h (listed with multiplicity) and

$$\chi_i(h_1h_2) = \chi_i(h_1)\chi_i(h_2),$$

 $|\chi_1(h)\chi_2(h)\cdots\chi_r(h)| = 1.$

The last equality holds because $h \in GL(L)$ and L is a free **Z**-module. Thus, we have a group homomorphism defined as follows:

$$\chi : H \longrightarrow \mathbf{R}^{r-1} = \{ (x_i)_{i=1}^r \in \mathbf{R}^r | \sum_{i=1}^r x_i = 0 \} \subset \mathbf{R}^r,$$
$$\chi(h) := (\log |\chi_1(h)|, \ \log |\chi_2(h)|, \ \cdots, \ \log |\chi_r(h)|) .$$

By the definition of N(H), we have

$$N(H) = \operatorname{Ker} \chi.$$

Thus N(H) is a normal subgroup and its quotient group $H/N(H) \simeq \chi(H)$ is a subgroup of \mathbf{R}^{r-1} .

It now suffices to show that $\chi(H)$ is discrete in the additive group \mathbf{R}^{r-1} with respect to the usual topology. Or equivalently, it suffices to show that the identity element $(0)_{i=1}^r$ is an isolated point of $\chi(H)$. Here the fact that H is defined over \mathbf{Z} is very crucial.

Let δ be any positive real number. Consider the set of all elements $h \in H$ satisfying the inequalities

$$\log |\chi_i(h)| < \delta \ , \ i = 1, \cdots, r$$

For such an h, $\chi_i(h)$ are all bounded, and hence the coefficients of the characteristic polynomial $\Phi_h(x)$ are all bounded. Since $\Phi_h(x) \in \mathbf{Z}[x]$, there are only finitely many characteristic polynomials for all such h. Thus the set of vectors $(\log |\chi_i(h)|)_{i=1}^r$ for all such h is also finite. Hence $(0)_{i=1}^m$ is an isolated point of $\chi(H)$.

In the rest of this section, we shall prove Theorem 2.1. First, we observe the following:

Lemma 2.5. Let H be a connected solvable, possibly discrete, subgroup of $GL(r, \mathbb{C})$. Let [H, H] denote the commutator subgroup of H. Then the eigenvalues of every element of [H, H] are all equal to 1.

Proof. By Lie-Kolchin Theorem, H is a subgroup of the group T(r) of upper triangle matrices of $\operatorname{GL}(r, \mathbb{C})$. From this, it is easy to see that the diagonal entries of any commutator $g_1g_2g_1^{-1}g_2^{-1}$ are all equal to 1. So are the diagonal entries of any product of commutators.

Proof of Theorem 2.1. By replacing G by a connected solvable subgroup of finite index, we may assume that G itself is connected and solvable.

We consider the derived series of G:

$$G = G^{(0)} > G^{(1)} > \dots > G^{(k-1)} > A := G^{(k)} > G^{(k+1)} = \{ \text{ id} \}.$$

Here, the series has length $k \ge 0$, $G^{(l)}$ are normal subgroups of G and $[G^{(l)}, G^{(l)}] = G^{(l+1)}$ for $l = 0, 1, \dots, k$. Note that $A := G^{(k)}$ is an abelian group.

We shall prove Theorem 2.1 by induction on the length k.

If k = 0, then G = A, i.e. G itself is abelian, and the result follows from the result of Dinh-Sibony (Remark 2.2(4)).

Assume that $k \ge 1$. Assuming that Theorem 2.1 is true when the length $\le k-1$, we shall show that Theorem 2.1 is also true when the length = k.

First note that $A \subseteq [G, G] = G^{(1)}$. By Lemma 2.5, the eigenvalues on $V_{\mathbf{C}}$ of every element of A are all equal to 1.

For each $a \in A$, we write the associated real eigenspace by V(a, 1):

 $V(a, 1) := \{ x \in V \mid a(x) = x \} \subset V .$

Let us consider the linear subspace W of V,

 $W = \bigcap_{a \in A} V(a, 1) \,.$

Claim 2.6.

- (1) $C_W := C \cap W \neq \{0\}$.
- (2) For each $g \in G$, g(W) = W. In particular, $G(C_W) \subseteq C_W$.

Proof. (1) Since Theorem 2.1 is true for the abelian group A, there is a common eigenvector in C for A and hence (1) follows.

(2) Let $0 \neq w \in W$. Since A is a normal subgroup of G, we can write $g^{-1}ag = a_g$ for some $a_g \in A$. Thus, regardless of the choice of w we have

$$(g^{-1}ag)(w) = a_g(w) = w$$
.

Hence

So

$$a(g(w)) = g(w).$$

W.

$$g(W) \subseteq \bigcap_{a \in A} V(a, 1) =$$

Since g is invertible, this implies g(W) = W.

From now on, let us consider the action of G on W and on C_W . Set

$$G_W^{(l)} := \operatorname{Im} \left(G^{(l)} \longrightarrow \operatorname{GL}(W) \right)$$
.

Then, $G_W := G_W^{(0)}$ is connected solvable, and the new series

$$G_W = G_W^{(0)} \ge G_W^{(1)} \ge \dots \ge G_W^{(k-1)} \ge A_W := G_W^{(k)} \ge G_W^{(k+1)} = \{ \operatorname{id}_W \}$$

gives the derived series, possibly redundant, of G_W .

Since $W = \bigcap_{a \in A} V(a, 1)$ and hence $A_W = \{ id_W \}$, the new series is in fact redundant. Now we can apply induction hypothesis to the series of length $\leq k - 1$ and we obtain the result.

This completes the proof of Theorem 2.1.

3. Some algebro-geometric consequences of Theorem of Lie-Kolchin type for a cone

In this section, we shall derive some direct consequences of Theorem of Lie-Kolchin type for a cone (Theorem 2.1 or Corollary 2.3) for groups of automorphisms or birational automorphisms of algebraic varieties and compact Kähler manifolds.

(3.1) Let $f: X \longrightarrow S$ be a projective surjective morphism of normal varieties defined over a field k with connected fibers of positive dimension. Assume that S is quasi-projective. Special important cases are the case where S = Spec k (abstract

case) and the case where $f := \Phi_{|mK_X|} : X \longrightarrow S$ is the relatively minimal Kodairalitaka fibration of X. We note that the relative canonical class $K_{X/S}$ is trivial in NS(X/S) (cf. (3.3)) in the second case.

(3.2) By Bir (X) and by Aut (X), we denote the group of birational transformations of X and the group of biregular transformations of X respectively. Let:

$$\operatorname{Bir}(X/S) := \{\varphi \in \operatorname{Bir}(X) \mid f = f \circ \varphi\}$$

$$\operatorname{Bir}^{s}(X/S) := \{\varphi \in \operatorname{Bir}(X/S) \mid \varphi \text{ is isomorphic in codimension one} \},\$$

$$\operatorname{Aut}(X/S) := \{\varphi \in \operatorname{Aut}(X) \mid f = f \circ \varphi\}.$$

Note that $\operatorname{Bir}^{s}(X) = \operatorname{Bir}(X)$ when X is a (terminal) minimal model.

(3.3) Let NS(X/S) be the set of numerical equivalence classes of Cartier divisors on X over S. Note that NS(X/S) is the free part of the "Néron-Severi" group. It is well known that NS(X/S) is a free Z-module of finite rank ([Kl]). We define the relative nef cone $\overline{Amp}(X/S)$ to be the closure of the relative ample cone Amp(X/S)in $NS(X/S)_{\mathbf{R}}$ (under the usual topology). Similarly, the relative pseudo-effective cone $\overline{Big}(X/S)$ is defined to be the closure of the relative big cone Big(X/S) in $NS(X/S)_{\mathbf{R}}$. Both spaces form closed convex cones of NS(X/S). Moreover, since the dimension of fibers of f is assumed to be positive, both cones are strict. For more details on these cones and their roles in birational algebraic geometry, we refer the readers to [Ka2].

(3.4) Let G be a subgroup of Aut (X/S). We call G numerically virtually solvable if its image in GL (NS(X/S)) is virtually solvable. By Tits' Theorem (Remark 2.2(1)), G satisfies either one of the following two, which are not in general mutually exclusive:

- (i) G is numerically virtually solvable, or
- (ii) G contains a subgroup isomorphic to $\mathbf{Z} * \mathbf{Z}$.

Applying Corollary 2.3 to the action of G on $(NS(X/S), \overline{Amp}(X/S))$, we have the following:

Theorem 3.1. Let G be a numerically virtually solvable subgroup of $\operatorname{Aut}(X/S)$. Then there are a finite-index subgroup G_1 of G and a nonzero real vector $v \in \operatorname{Amp}(X/S)$ such that $G_1(\mathbf{R}_{\geq 0}v) = \mathbf{R}_{\geq 0}v$. In particular, if G is a numerically virtually solvable subgroup of $\operatorname{Aut}(X)$, then a suitable finite-index subgroup of G admits a common real nef eigenvector.

We know that the relative canonical divisor class $K_{X/S}$ is always fixed by the group Aut (X/S). However, the class $K_{X/S}$ is relatively trivial over S, if f is a relatively minimal Kodaira-Iitaka fibration or if X has a numerically trivial canonical class (X a Calabi-Yau variety in a wider sense) and $S = \operatorname{Spec} k$.

The next theorem will be applied in Section 5.

Theorem 3.2. Let X be an n-dimensional projective manifold whose (n-1)-th Chern class $c_{n-1}(X)$ is in the boundary of the cone of effective 1-cycles $\overline{NE}(X)$. Let G be a numerically virtually solvable subgroup of $\operatorname{Aut}(X)$. Then there are a finiteindex subgroup G_1 of G and a nonzero real nef vector v such that $G_1(\mathbf{R}_{\geq 0}v) = \mathbf{R}_{\geq 0}v$ and $(v.c_{n-1}(X)) = 0$. The same is true for a complex projective minimal threefold (see §5) whose second Chern class $c_2(X)$ is not in the interior of $\overline{NE}(X)$. *Proof.* Since $c_{n-1}(X)$ is stable under Aut (X), the free submodule M of NS(X) defined by

$$M := \{ x \in NS(X) \mid (x, c_{n-1}(X)) = 0 \}$$

is also $\operatorname{Aut}(X)$ -stable. We set

$$C := \overline{\operatorname{Amp}}(X) \cap M_{\mathbf{R}}.$$

Since $c_{n-1}(X)$ is in the boundary of $\overline{\operatorname{NE}}(X)$, it follows that $C \neq \{0\}$. So, we obtain the first assertion by applying Corollary 2.3 for the action of G on (M, C). The last statement follows from a result of Miyaoka [Mi], which says that $c_2(X)$ is well-defined and is always in $\overline{\operatorname{NE}}(X)$ for a minimal projective complex threefold. \Box

(3.5) For several reasons, even $\operatorname{Bir}^{s}(X/S)$ does not act on $\operatorname{NS}(X/S)$ in general. We quote the following result of Kawamata [Ka2, the proof of Lemma 1.1].

Theorem 3.3. Assume that char k = 0 and X admits at most \mathbf{Q} -factorial rational singularities. Let $N^1(X/S)$ be the group of numerical equivalence classes of Weil divisors on X over S. Then $N^1(X/S)$ is a free \mathbf{Z} -module of finite rank and satisfies $[N^1(X/S) : NS(X/S)] < \infty$. Moreover, $\operatorname{Bir}^s(X/S)$ acts naturally and linearly on $N^1(X/S)$ and preserves $\overline{\operatorname{Big}}(X/S) (\subset N^1(X/S)_{\mathbf{R}})$.

Let G be a subgroup of Bir^s (X/S). We call G numerically virtually solvable if its image in GL $(N^1(X/S))$ is virtually solvable. Since $[N^1(X/S) : NS(X/S)] < \infty$, the definitions here and in (3.4) are consistent if $G \subset Aut(X/S)$.

Now, applying Corollary 2.3 to the action G on $(N^1(X/S), \overline{Big}(X/S))$, we obtain the following:

Theorem 3.4. Assume that char k = 0 and X admits at most \mathbf{Q} -factorial rational singularities. Let G be a numerically virtually solvable subgroup of $\operatorname{Bir}^{s}(X/S)$. Then there are a finite-index subgroup G_{1} of G and a nonzero real vector $v \in \operatorname{Big}(X/S)$ such that $G_{1}(\mathbf{R}_{\geq 0}v) = \mathbf{R}_{\geq 0}v$. In particular, if X is minimal and if G is a numerically virtually solvable subgroup of $\operatorname{Bir}(X)$, then a suitable finite-index subgroup of G admits a common real pseudo-effective eigenvector.

In the above Theorem, we do not need the rationality of singularities, because **Q**-factoriality already implies that $NS(X/S)_{\mathbf{Q}} = N^1(X/S)_{\mathbf{Q}}$. However, in order to guarantee the discreteness as in Lemma 2.4, the action of G on some **Z**-module structure will be needed.

(3.6) Let X be a compact Kähler manifold and let $\overline{\mathcal{K}}(X)$ be the closure of the Kähler cone $\mathcal{K}(X)$ in $H^2(X, \mathbf{R})$. Let G be a subgroup of Aut (X). We call G cohomologically virtually solvable if its image in GL $(H^2(X, \mathbf{Z})/\text{torsion})$ is virtually solvable. When X is projective, G is numerically virtually solvable if G is cohomologically virtually solvable.

Applying Corollary 2.3 to the action of G on $(H^2(X, \mathbb{Z})/\text{torsion}, \overline{\mathcal{K}}(X))$, we obtain the following:

Theorem 3.5. If G is a cohomologically virtually solvable subgroup of Aut (X), then there are a finite-index subgroup G_1 of G and a nonzero real vector $v \in \overline{\mathcal{K}}(X)$ such that $G_1(\mathbf{R}_{\geq 0}v) = \mathbf{R}_{\geq 0}v$.

4. Entropy and Conjecture of Tits type

In this section, we verify Conjecture of Tits type (Conjecture 1.3) posed in Introduction for some basic varieties, i.e., for surfaces, complex tori and hyperkähler manifolds. In the next section, we shall also verify the conjecture for minimal threefolds. In our proof, Theorem 2.1 or Corollary 2.3 also plays an important role.

(4.1) We recall a definition of *entropy* of an automorphism of a compact Kähler manifold from [DS] and generalize it to a complex normal projective variety.

For the references to the following important results, see Dinh-Sibony [DS] and Guedj[Gu, (1.2), (1.6)].

Theorem 4.1. ([DS]) Let M be a compact Kähler manifold of dimension n and let g be an automorphism of M. By $\rho(g^*|W)$ we denote the spectral radius of the action of g^* on a g^* -stable subspace W of the total cohomology group $H^*(M, \mathbb{C})$. Then we have:

(1) $\rho(g^*|H^*(M, \mathbf{C})) \ge 1$, and $\rho(g^*|H^*(M, \mathbf{C})) = 1$ (resp. > 1) if and only if $\rho(g^*|H^2(M, \mathbf{C})) = 1$ (resp. > 1).

Moreover, $\rho(g^*|H^*(M, \mathbf{C})) = 1$ (resp. > 1) if and only if so is for g^{-1} . (2) $\rho(g^*|H^2(M, \mathbf{C})) = \rho(g^*|H^{1,1}(M))$.

- Moreover, if M is projective, then this value is also equal to $\rho(g^*|NS(M))$.
- (3) The spectral radius $\rho(g^*|H^{i,i}(X, \mathbb{C}))$ equals the *i*-th dynamical degree $d_i(g)$. Further, there are integers $m \leq m'$ such that

$$1 = d_0(g) < \dots < d_m(g) = \dots = d_{m'}(g) > \dots > d_n(g) = 1.$$

(4) $\rho(g^*|H^*(X, \mathbf{C})) = \max_{0 \le i \le n} d_i(g) = e^{h(g)}$, with h(g) the topological entropy of g.

Definition 4.2.

(1) Let M be a compact Kähler manifold and let g be an automorphism of M. We call g of null entropy (resp. of positive entropy) if and only if the spectral radius $\rho(g^*|H^2(M, \mathbb{C})) = 1$ (resp. > 1). By (4.1), g is of null entropy (resp. of positive entropy) if and only if $\rho(g^*|\text{NS}(M)) = 1$ (resp. > 1) when X is projective.

(2) Let V be a complex normal projective variety. Taking account of the last characterization in (1), we call an automorphism g of V of null entropy (resp. of positive entropy) if the spectral radius $\rho(g^*|NS(V))$ on the free part NS(V) of the Néron-Severi group is 1 (resp. > 1). If V is **Q**-factorial, then $g \in Aut(V)$ is of positive entropy if and only if so is the lifting of g to an automorphism on an equivariant resolution of V; see the proof of [Zh2, Lemma 2.6].

(4.2) In Introduction, inspired by Tits' Theorem and Dinh-Sibony's Theorem (Theorem 1.2) we posed Conjecture of Tits type (Conjecture 1.3). The following is a weaker version of the conjecture, which can be easily verified.

Theorem 4.3. Let L be a free \mathbb{Z} -module of rank $r \geq 1$ and let H be a subgroup of $\operatorname{GL}(L)$. Assume that H does not contain a subgroup isomorphic to $\mathbb{Z} * \mathbb{Z}$. Then, there is a connected solvable finite-index subgroup H_1 of H such that the subset

$$N(H_1) := \{ h \in H_1 \mid \rho(h) = 1 \}$$

of H_1 is a normal subgroup of H_1 and $H_1/N(H_1)$ is a free abelian group of rank at most r-1.

Applying this theorem to the action of G on $H^2(X, \mathbb{Z})$ or on NS(X), we see that the conjecture is true *except the rank estimate*. In other words, the rank estimate is the most essential part of the conjecture.

Proof. By Tits' Theorem, H is virtually solvable. Thus H has a finite-index subgroup H_1 which is connected solvable. Then, the result follows from Lemma 2.4. \Box

(4.3) In the rest of this section, we verify Conjecture of Tits' type for surfaces (Theorem 4.4), hyperkähler manifolds (Theorem 4.5), and complex tori (Theorem 4.6). We also study Conjecture of Tits type for coverings and fibrations (Propositions 4.7, 4.8).

Theorem 4.4. Conjecture of Tits type is true if dim $X \leq 2$.

Proof. There is nothing to prove when $\dim X = 0$ or 1.

Assume dim X = 2. If X is smooth and projective, the conjecture has been proved by [Zh1]. Therefore, when X is projective, it is possible to reduce to the smooth projective case by considering a minimal resolution \tilde{X} of X and the induced action on \tilde{X} . It is also possible to argue by using classification of surfaces (see eg. [BHPV]) when X is Kähler. Instead of such a case-by-case proof, we shall give here a unified, classification-free proof.

Define

$$L := H^{2}(X, \mathbf{Z}) \text{ (resp.} = \mathrm{NS}(X)),$$
$$C := \overline{\mathcal{K}}(X) \text{ (resp.} = \overline{\mathrm{Amp}}(X))$$

when X is a compact Kähler surface (resp. X is a normal projective surface). By the Hodge index theorem, the real linear subspace

$$W := \langle C \rangle$$

of $L_{\mathbf{R}}$ is hyperbolic, i.e. of signature (1, *) with respect to the intersection form. This is because

$$W = H^{1,1}(M, \mathbf{R})$$
 or $W = NS(X)_{\mathbf{R}}$

according to the two cases above.

Let us consider the image

$$G^* := \operatorname{Im} \left(G \longrightarrow \operatorname{GL}(L) \right)$$
.

We may assume that G^* is connected solvable. Let $v \in C$ be a common eigenvector of G^* found by Corollary 2.3. Then, we can write $g^*(v) = \chi(g)v$.

Claim: (1)
$$\rho(g^*|W) = \rho(g^*|L)$$
. (2) $\chi(g) = \rho(g^*|L)$ or $\frac{1}{\rho(g^*|L)}$.

Proof of Claim. (1) In the Kähler surface case, the intersection form on the subspace $(H^{2,0}(X) \oplus H^{0,2}(X)) \cap H^2(X, \mathbf{R})$ is positive definite, so the eigenvalues of g^* on this subspace all have absolute value 1.

(2) By Birkhoff-Perron-Frobenius' Theorem (Remark 2.2(3)) applied to the action of g^* on (W, C), there is a nonzero vector $v_g \in C$ such that $g^*(v_g) = \rho(g^*|W) v_g$. Note that

$$(v.v_g) = (g^*(v).g^*(v_g)) = \chi(g)\rho(g^*|W)(v.v_g)$$
.

If $\chi(g) \neq \frac{1}{\rho(g^*|W)}$, then we must have $(v, v_g) = 0$. Since $v, v_g \in C$ and since W is hyperbolic by the Hodge index theorem, it follows that v is parallel to v_g , so $\chi(g) = \rho(g^*|W)$. Now, (2) follows from (1).

Resuming the proof of the theorem, we consider the group homomorphism

$$\chi: G \longrightarrow \mathbf{R} ; g \mapsto \log |\chi(g)| = \log \chi(g) .$$

By Claim, we see that $\chi(g) = \rho(g^*|L) = 1$ if and only if g is of null entropy. This means that

$$\operatorname{Ker} \chi = N(G),$$

and hence

$$G/N(G) \simeq \chi(G).$$

Furthermore, by Claim again, we have

$$\chi(G) \subset \{\pm \log \rho(g^*|L) \mid g \in G\} .$$

Note here that if $\rho(g^*|L)$ is bounded, then all eigenvalues of $g^*|L$ are also bounded. We also note that the action G^* is defined over the **Z**-module L. Thus, $\chi(G)$ is discrete in **R** as we have seen in the proof of Lemma 2.4. Hence $\chi(G) \simeq \mathbf{Z}^s$ with $s \leq 1$.

By a hyperkähler manifold M, we mean a simply-connected compact Kähler manifold admitting an everywhere non-degenerate holomorphic 2-form σ_M such that $H^0(M, \Omega_M^2) = \mathbf{C}\sigma_M$. According to the Bogomolov decomposition Theorem [Be1], hyperkähler manifolds form one of the three building blocks of compact Kähler manifolds with vanishing first Chern class. The other two building blocks are complex tori and Calabi-Yau manifolds (in the narrow sense).

Theorem 4.5. Conjecture of Tits type is true for hyperkähler manifolds.

Proof. By Beauville [Be1], $H^2(M, \mathbb{Z})$ of a hyperkähler manifold M admits an integral symmetric bilinear form (*, **) of signature $(3, b_2(M) - 3)$. This form is invariant under Aut (M) and is of signature $(1, h^{1,1}(M) - 1)$ on $H^{1,1}(M, \mathbb{R})$. A bit more precisely, $(\eta, \eta') > 0$ for any Kähler classes η and η' . Thus, the same proof as in Theorem 4.4 shows the result. Furthermore, in the statement (2) of the conjecture, we have $G_1/N(G_1) \simeq \mathbb{Z}^s$ with $s \leq 1$.

Theorem 4.6. Conjecture of Tits type is true for complex tori.

Proof. Let X be a complex torus of dimension n and let G be a subgroup of Aut (X). Note that

$$H^{1}(X, \mathbf{Z}) \otimes \mathbf{C} = H^{0}(X, \Omega^{1}_{X}) \oplus H^{0}(X, \Omega^{1}_{X}) ,$$

$$H^{*}(X, \mathbf{Z}) = \bigoplus_{k=0}^{2n} \wedge^{k} H^{1}(X, \mathbf{Z}) .$$

Thus $g \in G$ is of null entropy if and only if the spectral radius $\rho(g^*|H^0(X, \Omega^1_X)) = 1$. Let us consider the action of G on $H^0(X, \Omega^1_X) \simeq \mathbb{C}^n$, and define

$$G^* := \operatorname{Im} \left(G \longrightarrow \operatorname{GL} \left(H^0(X, \Omega^1_X) \right) \right)$$

As in the proof of Theorem 4.4, we may assume that G^* is connected solvable. Then, as in Lemma 2.4, choosing a suitable basis of $H^0(X, \Omega^1_X)$, one can embed G^* into the subgroup T(n) of upper triangle matrices of $\operatorname{GL}(n, \mathbb{C}) = \operatorname{GL}(H^0(X, \Omega^1_X))$, and we have a group homomorphism

$$\chi: G \longrightarrow \mathbf{R}^{n-1} = \{(x_i)_{i=1}^n \in \mathbf{R}^n \mid \sum_{i=1}^n x_i = 0\} \subset \mathbf{R}^n$$

which is defined by

$$\chi(g) = (\log |\chi_1(g)|, \ \log |\chi_2(g)|, \ \cdots, \ \log |\chi_n(g)|) .$$

Here $\chi_i(g)$ is the (i, i)-th entry of the matrix g (regarded as an element of T(n)). Thus

$$N(G) = \operatorname{Ker} \chi$$

Note that the eigenvalues of $g^*|H^1(X, \mathbb{Z})$ are $\chi_i(g)$ and $\overline{\chi_i(g)}$ $(1 \le i \le n)$. Thus, one can apply the same argument as in Lemma 2.4 to get

$$G/N(G) \simeq \operatorname{Im} \chi \simeq \mathbf{Z}^s$$

for some $s \leq n - 1$.

It is highly interesting to verify the conjecture for Calabi-Yau manifolds (in the narrow sense). Unfortunately, we could not yet do it. All we can say now is that the conjecture is true for Calabi-Yau threefold, as a special case of Theorem 5.1 in Section 5. See also Proposition 4.9 for a relevant result.

Next two propositions will be used in Section 5.

Proposition 4.7. Let G be a subgroup of Aut X and let $\pi : \tilde{X} \longrightarrow X$ be a generically finite surjective morphism between compact Kähler manifolds or between complex normal projective varieties. Assume that G lifts to a subgroup $\tilde{G} \simeq G$ of Aut (\tilde{X}) equivariantly. Then Conjecture of Tits type is true for (X, G) if so is for (\tilde{X}, \tilde{G}) .

Proof. Let dim $X = \dim \tilde{X} = n$. If G contains a subgroup isomorphic to $\mathbb{Z} * \mathbb{Z}$, then we are done. Therefore we may assume that $\tilde{G} \simeq G$ does not contain a subgroup isomorphic to $\mathbb{Z} * \mathbb{Z}$. Then, by Tits' Theorem, any linear action of \tilde{G} on a free \mathbb{Z} module of finite rank is virtually solvable. Furthermore, by replacing $G \simeq \tilde{G}$ by a suitable finite-index subgroup, we may assume that $\tilde{G}|H^2(\tilde{X}, \mathbb{Z})$ and $G|H^2(X, \mathbb{Z})$ or $\tilde{G}|\mathrm{NS}(\tilde{X})$ and $G|\mathrm{NS}(X)$ are connected solvable. Since $\pi^*: H^2(X, \mathbb{Z}) \longrightarrow H^2(\tilde{X}, \mathbb{Z})$ or $\pi^*: \mathrm{NS}(X) \longrightarrow \mathrm{NS}(\tilde{X})$ is injective, there is then a surjective homomorphism ϕ : $\tilde{G}/N(\tilde{G}) \longrightarrow G/N(G)$. Since G/N(G) is free by Lemma 2.4, the result follows. \Box

If both X and \hat{X} are **Q**-factorial, then the group homomorphism ϕ is indeed an isomorphism (see [NZ, Appendix, Lemma A.8] and also [Zh2, Lemma 2.6]), hence the converse of the conclusion is also true.

Thanks to Proposition 4.7, we may freely replace the variety X by its G-equivariant resolution, which always exists.

Proposition 4.8. Let $f : X \longrightarrow Y$ be a surjective morphism with connected fibres. Assume that X is smooth. Let F be a general fiber. Let G be a subgroup of Aut (X/Y) (so that G acts faithfully on F). Then, Conjecture of Tits type is true for G with the rank bound dim F - 1 if and only if so is for the action of G on F.

Proof. We write by G_F the action of G on F. As in the previous proposition, we may assume that G does not contain $\mathbf{Z} * \mathbf{Z}$. Replacing G by a finite-index subgroup, we may assume that both

 $G^* := \operatorname{Im}(\varphi_X : G \longrightarrow \operatorname{GL}(\operatorname{NS}(X)) \text{ and } G_F^* := \operatorname{Im}(\varphi_F : G_F \longrightarrow \operatorname{GL}(\operatorname{NS}(F)))$

are connected solvable. The groups N(G) and $N(G_F)$ are then well-defined normal subgroups of $G = G_F$ (cf. Lemma 2.4). Then, by [Zh2] (2.1) Remark (11) or [NZ, Appendix, Theorem D], we have $N(G_F) = N(G)$ as subgroups of G. Thus we have the identification:

$$G_F/N(G_F) = G/N(G)$$
.

This implies the result.

As an application, we have the following:

Proposition 4.9. Conjecture of Tits type is true for compact Kähler manifolds M with $c_1(M) = 0$ in $H^2(M, \mathbf{R})$, provided that Conjecture of Tits type is true for Calabi-Yau manifolds (in the narrow sense).

Proof. Let $\pi : \tilde{M} \longrightarrow M$ be the minimal splitting cover of M in the sense of [Be2]. \tilde{M} is the product of complex torus T, finitely many hyperkähler manifolds V_i and finitely many Calabi-Yau manifolds W_j . The subgroup G of Aut (M) lifts equivariantly to the subgroup $\tilde{G} \simeq G$ of Aut (\tilde{M}) and \tilde{G} preserves each factor T, V_i, W_j [ibid]. By the Künneth formula, $H^2(\tilde{M}, \mathbb{Z})$ is naturally isomorphic to the direct sum of $H^2(T, \mathbb{Z}), H^2(V_i, \mathbb{Z})$ and $H^2(W_j, \mathbb{Z})$. Thus, Conjecture of Tits type is true for \tilde{G} by Theorems 4.5, 4.6 and by our assumption for Calabi-Yau manifolds. Now the result follows from Proposition 4.7.

Remark 4.10. Furthermore, the proof of Proposition 4.9 together with Theorem 5.1 below implies that Conjecture of Tits type is true for compact Kähler manifolds M with $c_1(M) = 0$ in $H^2(M, \mathbf{R})$ if its minimal splitting cover does not have a Calabi-Yau manifold of dimension ≥ 4 as a factor.

5. Conjecture of Tits type for minimal threefolds

Let X be a normal complex projective variety. It is called *minimal* if X admits at most Q-factorial terminal singularities and the canonical class K_X is nef.

In this section, we shall show the following:

Theorem 5.1. Conjecture of Tits type is true for minimal threefolds.

Our proof is mostly classification free, but unfortunately, we could not avoid using some classification result of Calabi-Yau threefolds of very special type at the final step.

Proof. By the abundance theorem for threefolds [Ka3], the linear system $|mK_X|$ for some m > 0 defines a surjective morphism $\varphi : X \longrightarrow W$ from X to the canonical model W of X (Iitaka-Kodaira fibration). The morphism φ is Aut (X)-equivariant and the induced action of Aut X on W is finite by Deligne-Nakamura-Ueno [Ue, Theorem 14.10]. Thus, replacing G by a finite-index subgroup, we may assume that $G \subset \text{Aut}(X/W)$. Since the conjecture of Tits type is true when dim ≤ 2 by Theorem 4.4, the conjecture is also true if dim $W \geq 1$, i.e. if K_X is not numerically trivial. This can be seen by applying Proposition 4.7 to a G-equivariant resolution $\pi : \tilde{X} \longrightarrow X$, and then Proposition 4.8 to the composition $\varphi \circ \pi : \tilde{X} \longrightarrow W$.

Let us consider the case where K_X is numerically trivial. By Proposition 4.7 applied to the global index-one cover of X, we may assume that K_X is linearly equivalent to 0.

If $q(X) := h^1(\mathcal{O}_X) > 0$, then the albanese morphism $a : X \longrightarrow Alb(X)$ is an étale fiber bundle by [Ka1]. Thus, X is smooth and its minimal splitting cover is either the product of an elliptic curve and a K3 surface or an abelian threefold. In this case, the result follows from Proposition 4.9.

If $c_2(X) = 0$ as a linear form on NS(X), then X is an étale quotient of an abelian threefold by [SW]. So, X is smooth and its minimal splitting cover is an abelian threefold. Therefore the result also follows from Proposition 4.9.

Thus, we may assume further that q(X) = 0 and $c_2(X) \neq 0$.

Lemma 5.2. Let X be a minimal threefold such that $K_X \sim 0$, q(X) = 0 and $c_2(X) \neq 0$ (or more generally $K_X \sim_{\mathbf{Q}} 0$ and $c_2(X) \neq 0$). Let $G \subset \text{Aut}(X)$. Then, either one of the following holds:

- (1) Conjecture of Tits type is true for (X, G).
- (2) There is a nef and big Cartier divisor D such that $(c_2(X).D) = 0$.

Proof. Set $G^* := \text{Im} (G \longrightarrow \text{GL}(\text{NS}(X)))$. If $\mathbf{Z} * \mathbf{Z} \subset G^*$, then (1) holds. So, by Tits' theorem, we may assume that G^* is connected solvable (after replacing G by a finite-index subgroup).

Let

$$M := \{x \in \mathrm{NS}(X) \mid (c_2(X).x) = 0\}, \quad C := \overline{\mathrm{Amp}}(X) \cap M_{\mathbf{R}}.$$

For $g \in G$, we denote by $\rho(g)$ the spectral radius of $g^*|NS(X)$, i.e.

$$\rho(g) := \rho(g^* | \mathrm{NS}(X)) \,.$$

If each element of G is of null entropy, then we are done. So we may assume that $\rho(g_1) > 1$ for some $g_1 \in G$. Then, by Birkhoff-Perron-Frobenius' Theorem, there is a real vector $v_{g_1} \in \overline{\operatorname{Amp}}(X) \setminus \{0\}$ such that $g_1^* v_{g_1} = \rho(g_1) v_{g_1}$. Thus $c_2(X) v_{g_1} = 0$ as shown below. Hence $C \neq \{0\}$ and we can apply Theorem 3.2 to obtain a real vector $v_0 \in C \setminus \{0\}$ such that $G(\mathbf{R}_{\geq 0}v_0) = \mathbf{R}_{\geq 0}v_0$. Thus, for each $g \in G$ we can write $g^*v_0 = \chi(g)v_0$, and hence we have the group homomorphism:

$$\chi: G \longrightarrow \mathbf{R} ; g \mapsto \log \chi(g)$$

Note that $\rho(g) > 1$ if and only if $\rho(g^{-1}) > 1$ (for det $g^*|\text{NS}(X) = \pm 1$). Furthermore, if $\rho(g) = 1$ then $\rho(g) = \chi(g) = \rho(g^{-1}) = 1$. This implies that we have either one of the following two cases:

Case(i) There is an element $g \in G$ such that $\rho(g) > 1$ and the three positive real numbers $\chi(g)$, $\rho(g)$, $1/\rho(g^{-1})$ are mutually distinct.

Case(ii) For each $g \in G$, either $\rho(g) = 1$, or $\chi(g) = \rho(g) > 1$, or $\chi(g) = 1/\rho(g^{-1}) < 1$.

We deal with Case (i) first. By Birkhoff-Perron-Frobenius' Theorem, there is a real vector $v_g \in \overline{\text{Amp}}(X) \setminus \{0\}$ such that

$$g^* v_g = \rho(g) v_g$$
.

Since $g^*c_2(X) = c_2(X)$, we have

$$(c_2(X).v_g) = (g^*c_2(X).g^*v_g) = \rho(g)(c_2(X).v_g)$$

Thus

$$(c_2(X).v_q) = 0.$$

Similarly, since $\rho(g^{-1}) > 1$, there is a real vector $v_{g^{-1}} \in \overline{\operatorname{Amp}}(X) \setminus \{0\}$ such that $(g^{-1})^* v_{g^{-1}} = \rho(g^{-1}) v_{g^{-1}}$ and $(c_2(X).v_{g^{-1}}) = 0$.

Note that
$$c_2(X)$$
 gives an integer valued linear form on NS(X), so $v_g, v_{g^{-1}} \in M_{\mathbf{R}}$
Hence

 $v_g, v_{g^{-1}} \in C \setminus \{0\}.$

Let

$$D' := v_0 + v_a + v_{a^{-1}}$$

Then

$$(D'.c_2(X)) = 0.$$

Since v_0, v_g and $v_{g^{-1}}$ are eigenvectors of g^* with distinct eigenvalues $\chi(g), \rho(g)$ and $1/\rho(g^{-1})$, we have

$$(v_0.v_q.v_{q^{-1}}) \neq 0$$

by [DS], Lemma 4.4. Since v_0 , v_g and $v_{g^{-1}}$ are all nef, we have then

 $((D')^3) > 0$

and hence D' is nef and big. Since $M_{\mathbf{R}}$ is a rational hypersurface in NS(X)_{**R**} and the nef cone is locally rational polyhedral away from the cubic cone by [Ka2] or [Wi], we can then find a rational nef and big divisor D such that $(D.c_2(X)) = 0$. Therefore, the assertion (2) in the lemma holds.

Next we deal with Case (ii). First note that the image Im χ of the group homomorphism χ defined above is discrete in **R**. This follows from the same argument as in the last part of the proof of Theorem 4.4. Thus

Im
$$\chi \simeq \mathbf{Z}^s$$
 with $s \leq 1$.

Also, by the case assumption it is easy to see that $\operatorname{Ker} \chi = N(G)$. Therefore, $G/N(G) \simeq \operatorname{Im} \chi$ and the assertion (1) in the lemma holds. This proves the lemma.

Now the following lemma completes the proof of Theorem 5.1.

Lemma 5.3. Let X be a minimal threefold such that $K_X \sim 0$, q(X) = 0 and $c_2(X) \neq 0$. Assume that there is a nef and big Cartier divisor D such that $(c_2(X).D) = 0$. Then, Conjecture of Tits type is true for X.

Proof. Let us write $\zeta_n := e^{2\pi i/n}$. Let l = 3 or 7. By [Og1] or [OS], the universal cover of X is either one of X_l (l = 3, 7). Here X_3 is the unique crepant resolution of the quotient threefold $\overline{X}_3 := A_3/\langle g_3 \rangle$, where A_3 is the 3-times self product of the elliptic curve E_3 of period ζ_3 and $g_3 := \text{diag}(\zeta_3, \zeta_3, \zeta_3)$; X_7 is the unique crepant resolution of the quotient threefold $\overline{X}_7 := A_7/\langle g_7 \rangle$, where A_7 is the Jacobian threefold of the Klein quartic curve and $g_7 := \text{diag}(\zeta_7, \zeta_7^2, \zeta_7^4)$. Note that the singular locus of \overline{X}_l consists of finitely many points.

By Proposition 4.7, we may assume that X is X_l . In each case, we denote by $\nu_l : X_l \longrightarrow \overline{X}_l$ the unique crepant resolution. By [OS], this ν_l is also the unique birational contraction which contracts the class $c_2(X_l)$ in the cone of effective 1-cycles $\overline{NE}(X_l)$. Thus Aut (X_l) acts equivariantly on \overline{X}_l

Let $G \subset \operatorname{Aut}(X_l)$. Then G acts on \overline{X}_l equivariantly. Replacing G by a finiteindex subgroup, we may assume that each ν_l -exceptional divisor is G-stable. Thus, it suffices to show the assertion for the action of G on \overline{X}_l . Let \overline{X}_l^0 be the smooth locus of \overline{X}_l . Clearly \overline{X}_l^0 is G-stable. Moreover, by the shape of g_l , we see that $\pi_1(\overline{X}_l^0) = \pi_1(A_l).\langle g_l \rangle$ (a semi-direct product). We regard $\pi_1(\overline{X}_l^0)$ as a deck transformation group of the universal cover $U_l = \mathbb{C}^3 \setminus B_l$ of \overline{X}_l^0 . Here B_l is a discrete set of points ('lying over' the singular points of \overline{X}_l). Then $\pi_1(\overline{X}_l^0)$ is an affine transformation subgroup of \mathbb{C}^3 , in which the factor $\pi_1(A_l)$ forms the group of parallel translations and the factor $\langle g_l \rangle$ forms the group of linear transformations defined by the matrix g_l . From this, we can see that, in $\pi_1(\overline{X}_l^0)$,

$$\pi_1(A_l) = \{ \sigma \in \pi_1(\overline{X}_l^0) \mid \operatorname{ord} \sigma = \infty \} \cup \{ \operatorname{id} \} .$$

Thus the (equivariant) action of G on the universal cover U_l normalizes not only $\pi_1(\overline{X}_l^0)$, but also $\pi_1(A_l)$. Hence the action of G on \mathbb{C}^3 descends to an action on A_l equivariantly. Now the result follows from Theorem 4.6 and Proposition 4.7.

This completes the proof of the theorem.

References

- [BHPV] W. Barth, K. Hulek, C. Peters, A. Van de Ven, *Compact complex surfaces*, Second edition. Springer-Verlag, Berlin, 2004.
- [Be1] A. Beauville, Variété Kähleriennes dont la premiére classe de chern est nulle, J. Differential Geom. 18 (1983) 755–782.
- [Be2] A. Beauville, Some remarks on Kähler manifolds with $c_1 = 0$:In Classification of algebraic and analytic manifolds (Katata, 1982), Progr. Math. Birkhauser Boston, Boston, MA **39** (1983) 1–26.
- [Bir] G. Birkhoff, Linear transformations with invariant cones, Amer. Math. Month. 74 (1967) 274–276.
- [DS] T.-C. Dinh, N. Sibony, Groupes commutatifs d'automorphismes d'une variété kählerienne compacte, Duke Math. J. 123 (2004) 311–328.
- [Gu] V. Guedj, Ergodic properties of rational mappings with large topological degree, Ann. of Math. (2) 161 (2005), no. 3, 1589–1607.
- [Hu] J. E. Humphreys, *Linear algebraic groups*, Graduate Texts in Mathematics 21, Springer-Verlag, New York-Heidelberg (1975).
- [Kl] S. Kleiman, Toward a numeriacl theory of ampleness, Ann. of Math. 84 (1966) 293–344.
- [Ka1] Y. Kawamata, Minimal models and the Kodaira dimension of algebraic fiber space, J. Reine Angew. Math. 363 (1985) 1–46.
- [Ka2] Y. Kawamata, Crepart blowing-ups of 3-dimensional canonical singularities and its application to degeneration of surfaces, Ann. of Math. 127 (1988) 93–163.
- [Ka3] Y. Kawamata, Abundance theorem for minimal threefolds, Invent. Math. 108 (1992) 229–246.
- [Mi] Y. Miyaoka, The Chern classes and Kodaira dimension of a minimal variety, Adv. Stud. Pure Math. 10 North-Holland, Amsterdam (1987) 449–476.
- [NZ] N. Nakayama, D. -Q. Zhang, Building blocks of étale endomorphisms of complex projective manifolds, RIMS preprint 1577, Res. Inst. Math. Sci. Kyoto Univ. 2007. http://www.kurims.kyoto-u.ac.jp/preprint/index.html
- [Og1] K. Oguiso, On the complete classification of Calabi-Yau threefolds of type III₀, In the book: Higher-dimensional Complex Varieties (Trento, 1994), de Gruyter (1996) 329– 339.
- [Og2] K. Oguiso, Tits alternative in hypekähler manifolds, Math. Research. Letters 13 (2006) 307–316.
- [OS] K. Oguiso, J. Sakurai: Calabi-Yau threefolds of quotient type, Asian J. Math. 5 (2001) 43–77.
- [SW] N. I. Shepherd-Barron, P.M.H. Wilson, Singular threefolds with numerically trivial first and second chern class, J. Algebraic Geom. 3 (1994) 265–281.
- [Ti] J. Tits, Free subgroups in linear groups, J. Algebra **20** (1972) 250–270.
- [Ue] K, Ueno, Classification theory of alebraic varieties and compact complex spaces, Lecture Notes in Mathematics 439 Springer-Verlag, Berlin-New-York (1975).
- [Wi] P.M.H. Wilson, The Kähler cone on Calabi-Yau threefolds, Invent. math. 107 (1992) 561–583.
- [Zh1] D. -Q. Zhang, Automorphism groups and aniti-pluricnonical curves, preprint (2006). www.math.nus.edu.sg/~matzdq
- [Zh2] D. -Q. Zhang, Dynamics of automorphism on projective complex manifolds, preprint (2006); also Max Planck Institute Preprint, MPIM2007-19.

School of Mathematics, Korea Institute for Advanced Study, Dongdaemun-gu, Seoul 130-722, Korea

E-mail address: jhkeum@kias.re.kr

DEPARTMENT OF ECONOMICS, KEIO UNIVERSITY, HIYOSHI, KOHOKU-KU, YOKOHAMA, JAPAN AND SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, DONGDAEMUN-GU, SEOUL 130-722, KOREA

 $E\text{-}mail\ address:\ \texttt{oguiso@hc.cc.keio.ac.jp}$

Department of Mathematics, National University of Singapore, 2 Science Drive 2, Singapore 117543, Singapore

 $E\text{-}mail\ address: \verb"matzdq@math.nus.edu.sg"}$