

HILBERT POLYNOMIALS AND POWERS OF IDEALS

JÜRGEN HERZOG, TONY J. PUTHENPURAKAL AND JUGAL K. VERMA

ABSTRACT. The growth of Hilbert coefficients for powers of ideals are studied. For a graded ideal I in the polynomial ring $S = K[x_1, \dots, x_n]$ and a finitely generated graded S -module, the Hilbert coefficients $e_i(M/I^k M)$ are polynomial functions. Given two families of graded ideals $(I_k)_{k \geq 0}$ and $(J_k)_{k \geq 0}$ with $J_k \subset I_k$ for all k with the property that $J_k J_\ell \subset J_{k+\ell}$ and $I_k I_\ell \subset I_{k+\ell}$ for all k and ℓ , and such that the algebras $A = \bigoplus_{k \geq 0} J_k$ and $B = \bigoplus_{k \geq 0} I_k$ are finitely generated, we show the function $k \mapsto \ell(I_k/J_k)$ is of quasi-polynomial type, say given by the polynomials P_0, \dots, P_{g-1} . If $J_k = J^k$ for all k then we show that all the P_i have the same degree and the same leading coefficient. As one of the applications it is shown that $\lim_{k \rightarrow \infty} \ell(\Gamma_{\mathfrak{m}}(S/I^k))/k^n \in \mathbb{Q}$. We also study analogous statements in the local case.

INTRODUCTION

This paper is inspired by a result of Cutkosky, Ha, Srinivasan and Theodorescu. In their paper [9] they showed that for a graded ideal I in a polynomial ring $K[x_1, \dots, x_n]$ over a field K of characteristic 0, the limit $\lim_{k \rightarrow \infty} \ell((I^k)^{\text{sat}}/I^k)/k^n$ exists, but need not to be a rational number. Here I^{sat} denotes the saturation of I and $\ell(M)$ the length of a module. Our primary question then was whether this limit is a rational number when I is a monomial ideal. This is indeed the case, as we show in Corollary 2.6.

More generally, one may consider the j th symbolic power of an ideal I with respect to an ideal J , namely the ideal $I_k(J) = I^k : J^\infty$. The case of saturated powers is the special case in which J is the graded maximal ideal of S . Again, if I and J are monomials we can show that $\lim_{k \rightarrow \infty} e_0(I_k(J)/I^k)/k^{n-d}$ exists and is a rational number. Of course, $I_k(J)/I^k$ is not a finite length module in general, so that in this limit formula we had to replace the length of $I_k(J)/I^k$ by the multiplicity $e_0(I_k(J)/I^k)$ of this module. Moreover in the denominator we can replace k^n by k^{n-d} , where d is the limit dimension, that is, the dimension of $I_k(J)/I^k$ for $k \gg 0$, as it can be shown that $\dim I_k(J)/I^k$ is constant for large k . For the proof of this result we use the fact that the graded S -algebra $\bigoplus_{k \geq 0} I_k(J)$ is finitely generated if I and J are monomial ideals, as shown in [11, 3.2].

1991 *Mathematics Subject Classification*. Primary 13H15, 13D40 ; Secondary 13A30.

Key words and phrases. multiplicity, regularity, reduction, Hilbert polynomial, Rees polynomial.

Jugal Verma and Tony J. Puthenpurakal thank Universität Duisburg-Essen for hospitality during Sept-Dec 2006. The authors thanks DFG for financial support, which made this visit possible.

The result described in the preceding paragraph is again a special case of the following more general situation: given two families $(I_k)_{k \geq 0}$ and $(J_k)_{k \geq 0}$ of graded ideals of S with the property that $J_k J_\ell \subset J_{k+\ell}$ and $I_k I_\ell \subset I_{k+\ell}$ for all k and ℓ , and such that the algebras $A = \bigoplus_{k \geq 0} J_k$ and $B = \bigoplus_{k \geq 0} I_k$ are finitely generated. We also assume that $J_k \subset I_k$ for all k and that $I_0 = J_0 = S$.

What can be said about $e_0(I_k/J_k)$ as a function of k ? We first notice in Proposition 2.4 is that $e_0(I_k/J_k)$ is a quasi-polynomial for $k \gg 0$. Indeed, this can be deduced from the fact that $e_0(I^k M/J^k M)$ is quasi-polynomial for $k \gg 0$ for any finitely generated graded S -module M and graded ideal $J \subset I \subset S$, see Proposition 2.1. For the proof of this proposition we use Theorem 1.1, where we consider a finitely generated graded S -module N . The Hilbert polynomial $P_N(x)$ of N can be written in the form

$$(1) \quad P_N(x) = \sum_{i=0}^d (-1)^i e_i(N) \binom{x+d-i}{d-i}$$

with integer coefficients $e_i(N)$, called the *Hilbert coefficients* of N . Now Theorem 1.1 states the following: Let M be a graded S -module, and $I \subset S$ a graded ideal with $\dim M/IM = d$. Then for all $i = 0, \dots, d$, the Hilbert coefficient $e_i(M/I^k M)$ as a function of k is of polynomial type of degree $\leq n - d + i$. The assertion that $e_i(M/I^k M)$ is a polynomial follows easily from [12, 4.3] (also see proof of [15, Theorem 1]). However our degree bound is new.

In Section 5 we give a different proof of Proposition 2.1, not referring to Theorem 1.1. This alternative proof is also valid for ideals $J \subset I$ in a local ring. In fact, we do not have a local version of Theorem 1.1. In other words, we do not know whether for an ideal I in a local ring (R, \mathfrak{m}) the Hilbert coefficients $e_i(R/I^k)$ are polynomial functions for large k . This is well-known and easy to prove for $i = 0$, but seems to be unknown for $i > 1$. Actually we expect that $e_i(R/I^k)$ may not be of polynomial type for $i > 1$.

For any quasi-polynomial numerical function like $e_0(I_k/J_k)$ (say of period g and given by polynomials P_0, \dots, P_{g-1}) a natural question is:

(*) Do the polynomials P_0, \dots, P_{g-1} have the same degree and the same leading term?

To put this question in the right historical framework let us consider the following: Let P be a d -dimensional rational convex polytope in \mathbb{R}^n and let $E(P, k) = \# kP \cap \mathbb{Z}^n$ be the Ehrhart function. It is well-known that $E(P, k)$ is of quasi-polynomial type cf. [4, 6.3.11], say given by Q_0, \dots, Q_{g-1} . We state a simplified version of a conjecture due to Ehrhart: *If the affine span of every $d - 1$ dimensional face of P contains a point with integer co-ordinates then Q_0, \dots, Q_{g-1} have the same degree and the same leading term.*

A more general version of this was proved independently by McMullen (see [16]) and Stanley ([24], Theorem 2.8). For a recent proof see [5, 5].

In our case, since we know that $e_0(I_k/J_k)$ is of quasi-polynomial type, we also would like to know whether the polynomials describing this quasi-polynomial function are all of

same degree and all have the same leading term. Then this would imply at once that $\lim_{k \rightarrow \infty} e_0(I_k/J_k)/k^{n-d}$ exists and is a rational number.

The right set-up to study these questions is to consider the case of homogeneous inclusion of Noetherian algebras $A = \bigoplus_{k \geq 0} A_k \subseteq B = \bigoplus_{k \geq 0} B_k$, with $A_0 = B_0$. We assume A is standard graded, while B need not be. Furthermore A_0 is either the polynomial ring $S = K[x_1, \dots, x_n]$ or a local Noetherian ring (R, \mathfrak{m}) with residue field $K = R/\mathfrak{m}$. In the case when $A_0 = S$ we also assume that A, B are bigraded. In this general set-up one can similarly prove that $k \mapsto e_0(B_k/A_k)$ is of quasi-polynomial-type (in the local case we take Hilbert-Samuel multiplicity with respect to \mathfrak{m}). We consider the question (*) for this function.

Let $M = \bigoplus_{k \geq 0} M_k$ be a graded but not necessarily finitely generated A -module (it is bigraded if A is). We say M is a quasi-finite A -module if $H_{A_+}^0(M)_i = 0$ for $i \gg 0$. Here $H_{A_+}^0(M)$ denotes the 0th local cohomology of M with respect to the ideal $A_+ = \bigoplus_{k > 0} A_k$. Our main result (see Theorem 2.5 and Theorem 3.3) is that when B/A is quasi-finite A -module, then indeed the function $k \mapsto e(B_k/A_k)$ satisfies (*). The module B/A is quasi-finite under the mild hypothesis that $\text{grade}(A_1 B, B) > 0$. This is trivially satisfied in the cases when A is the Rees algebra $\mathcal{R}(I) = \bigoplus_{n \geq 0} I^n t^n$ of an ideal I and $B = \bigoplus_{k \geq 0} I_k$, where $(I_k)_k \geq 0$ is a family of ideals such that $I^{\bar{k}} \subseteq I_k$ for all k . In the case when $A_0 = R$ is local, $A = \mathcal{R}(J)$ and $B = \mathcal{R}(\mathcal{F})$, then note that $\text{grade}(A_1 B, B) > 0$ if $\text{grade}(J, R) > 0$.

The main technical tool required is that if M is a quasi-finite A -module and if K is uncountable, then M has a filter-regular element $x \in A_1$. Filter-regular elements in the case when M is finitely generated have a lot of applications in the study of blow-up algebras [26]. An early version of quasi-finite modules is in [18, 4.7]. However it was somewhat unexpected to us that the existence of filter-regular elements for quasi-finite modules have implications in question (*) above.

In section four, we specialise the above set-up to the case where $A = \mathcal{R}(J) \subset B = \mathcal{R}(I)$ for ideals $J \subseteq I$ of a d -dimensional local ring (R, \mathfrak{m}) and $\ell(I/J) < \infty$. Amao [1] showed that the numerical function $n \mapsto H(I/J, n) := \ell(I^n/J^n)$ is a polynomial function of degree at most d . We call $H(I/J, n)$ the Rees function of the pair (J, I) and the corresponding polynomial $P(I/J, n)$, the Rees polynomial of the pair (J, I) . Our objective is to identify the degree of the Rees polynomial. Rees [21] showed that if (R, \mathfrak{m}) is quasi-unmixed, then J is a reduction of I if and only if $\deg P(I/J, n) < d$. When I and J are \mathfrak{m} -primary, identification of the degree of $P(I/J, n)$ amounts to knowing the number of Hilbert-Samuel coefficients of I and J which coincide. This problem was considered in [23] by K. Shah. He introduced coefficient ideals of an \mathfrak{m} -primary ideal which help in identification of the degree of the Rees polynomial. We show in Theorem 4.4 that coefficient ideals exist even in the general case. Computation of coefficient ideals seems to be a difficult problem at present. We show that when J is a reduction of I then the degree of the Rees polynomial of the

pair (J, I) is $\dim \mathcal{R}(I)/\mathcal{R}(J) - 1$. As a consequence it follows that if R is quasi-unmixed, $J \neq I$ and $\mathcal{R}(J)$ satisfies Serre's condition S_2 , then $\deg P(I/J, n) = d - 1$.

Recall that an ideal is called normal if all its powers are integrally closed. It is rather surprising that for distinct normal ideals $J \subset I$, the problem of determination of the degree of the Rees polynomial of the pair (J, I) can be completely solved. Let $J \subset I$ be ideals of an analytically unramified d -dimensional local ring R such that $\ell(I/J)$ is finite and J is not a reduction of I . In Theorem 4.11 we show that $f(n) := \ell(\overline{I^n}/\overline{J^n})$ is a polynomial function of degree d . It follows that if J and I are normal then $\deg P(I/J, n) = d$.

In Section 5, (R, \mathfrak{m}) is a Noetherian local ring, I an ideal in R , M a finitely generated R -module and J a reduction of I with respect to M , i.e., $I^{m+1}M = JI^mM$ for some m . If $\dim M = r$, then for $i = 0, 1, \dots, r$ let $e_i(M)$ be the i th Hilbert coefficient of M (with respect to \mathfrak{m}). In other words,

$$\ell(M/\mathfrak{m}^{n+1}M) = \sum_{i=0}^r (-1)^i e_i(M) \binom{n+r-i}{r-i} \quad \text{for all } n \gg 0.$$

By a result due to Brodmann [2], $\text{Ass } M/I^k M$ is stable for $k \gg 0$. It follows that $\dim M/I^k M$ is stable for $k \gg 0$. We denote this stable value by $\theta_I(M)$. It is clear that $\bigoplus_{k \geq 0} I^k M/I^{k+1}M$ and $\bigoplus_{k \geq 0} I^k M$ are finitely generated modules over $\mathcal{R}(I)$. Since $\bigoplus_{k \geq 0} I^k M/J^k M$ is a finitely generated module over $\mathcal{R}(J)$, $\dim I^k M$, $\dim I^k M/I^{k+1}M$ and $\dim I^k M/J^k M$ stabilize for $k \gg 0$. We denote these stable values by $\alpha_I(M)$, $\beta_I(M)$ and $\gamma_J^I(M)$, respectively.

In view of Theorem 1.1 one may ask the following question: Is it true that the numerical functions

- (1) $e_i(M/I^k M)$, $i = 0, \dots, \theta_I(M)$,
- (2) $e_i(I^k M/I^{k+1}M)$, $i = 0, \dots, \alpha_I(M)$,
- (3) $e_i(I^k M)$, $i = 0, \dots, \beta_I(M)$, and
- (4) $e_i(I^k M/J^k M)$, $i = 0, \dots, \gamma_J^I(M)$

are polynomial functions in k ?

The graded versions of Question (1),(2) and (3) are equivalent, while (4) follows from (2). However in the local case all these questions are different. In Section 5, we settle Question (2), (3) and (4) (see Corollary 5.4). As we mentioned already, the answer to Question (1) is still open. It can be easily checked that if x^* , the initial form of x in the associated graded $\text{gr}_{\mathfrak{m}}(R)$ of R , is $\text{gr}_{\mathfrak{m}}(R)$ -regular then $e_i(R/(x)^k)$ is a polynomial in k . We do not know of any other cases where this question has a positive answer.

1. THE HILBERT COEFFICIENTS OF $M/I^k M$ AS A FUNCTION OF k

Let K be a field, $S = K[x_1, \dots, x_n]$ the polynomial ring in n variables, and let N be any graded S -module of dimension d . Then for $i \gg 0$, the numerical function $H(N, i) = \sum_{j \leq i} \dim_K N_j$ is a polynomial function of degree d , see [4, 4.1.6]. In other words, there

exists a polynomial $P_N(x) \in \mathbb{Z}[x]$ such that

$$H(N, i) = P_N(i) \quad \text{for all } i \gg 0.$$

The polynomial $P_N(x)$ is called the *Hilbert polynomial* of N . It can be written in the form

$$(2) \quad P_N(x) = \sum_{i=0}^d (-1)^i e_i(N) \binom{x+d-i}{d-i}$$

with integer coefficients $e_i(N)$, called the *Hilbert coefficients* of N .

Let $I \subset S$ a graded ideal and M a graded S -module with $\dim M/IM = d$. We are interested in the Hilbert coefficients $e_i(M/I^k M)$ of $M/I^k M$ as functions in k .

We say that a function $f: \mathbb{N} \rightarrow \mathbb{Q}$ is of polynomial type of degree d , if there exists a polynomial $h \in \mathbb{Q}[x]$ of degree d such that $f(k) = h(k)$ for all $k \gg 0$.

As the main result of this section we prove

Theorem 1.1. *Let M be a graded S -module, and $I \subset S$ a graded ideal with $\dim M/IM = d$. Then for all $i = 0, \dots, d$, the Hilbert coefficient $e_i(M/I^k M)$ as a function of k is of polynomial type of degree $\leq n - d + i$.*

Proof. For the proof we proceed by induction on d . If $d = 0$, then $P_{M/I^k M}(x) = \ell(M/I^k M)$, and hence the only Hilbert coefficient of $M/I^k M$ is $e_0(M/I^k M) = \ell(M/I^k M)$. This length, as a function of k , is known to be of polynomial type of degree $\leq n$, see [4, 4.1.6].

Now let $d > 0$. By a theorem of Brodmann [2] the set $\bigcup_k \text{Ass}(M/I^k M)$ is finite. Hence if $|K| = \infty$, as we may assume, there exists a linear form $y \in S$ such that the kernel of the multiplication map $y: (M/I^k M)(-1) \rightarrow M/I^k M$ is of finite length for all k . Set $N = M/yM$, then $N/I^k N = (M/I^k M)/y(M/I^k M)(-1)$, and it follows that

$$P_{N/I^k N}(j) = P_{M/I^k M}(j) - P_{M/I^k M}(j-1) \quad \text{for all } j.$$

From this we conclude that $e_i(M/I^k M) = e_i(N/I^k N)$ for $i = 0, \dots, d-1$. By induction hypothesis $e_i(N/I^k N)$ is of polynomial type of degree $\leq (n-1) - (d-1) + i = n-d+i$. Thus it remains to prove that $e_d(M/I^k M)$ is of polynomial type of degree $\leq n$.

We note that for all integers a one has

$$e_d(M/I^k M) = P_{M/I^k M}(ak) - \sum_{i=0}^{d-1} (-1)^i e_i(M/I^k M) \binom{ak+d-i}{d-i},$$

and that each of the summands $e_i(M/I^k M) \binom{ak+d-i}{d-i}$ is of polynomial type of degree $\leq n$. Thus it suffices to show that $P_{M/I^k M}(ak)$ is of polynomial type of degree $\leq n$.

To this end, we consider the sequence

$$0 \longrightarrow I^k M \longrightarrow M \longrightarrow M/I^k M \longrightarrow 0,$$

from which we deduce that

$$P_{M/I^k M}(ak) = P_M(ak) - P_{I^k M}(ak).$$

Since $P_M(x)$ is a polynomial of degree $\leq n$ it follows that $P_M(ak)$ is a polynomial in k of degree $\leq n$. Thus the assertion follows once it is shown that for suitable integer $a > 0$, $P_{I^k M}(ak)$ is of polynomial type of degree $\leq n$.

We consider the Hilbert series

$$\text{Hilb}_{I^k M}(t) = \sum_{i \geq 0} H(I^k M, i) t^i = \frac{Q_k(t)}{(1-t)^{n+1}}$$

of $I^k M$. Here $Q_k(t)$ is a polynomial. Indeed let $\beta_{ij} = \dim_K \text{Tor}_i^S(I^k M, K)_j$ be the graded Betti-numbers of $I^k M$. Then

$$Q_k(t) = \sum_{i=0}^n (-1)^i \beta_{ij} t^j,$$

see [4, 4.1.13]. It follows that $\deg Q_k(t) \leq \text{reg}(I^k M) + n$. As shown in [9, 2.4] and [15, 1] one has $\text{reg}(I^k M) \leq ek + f$ for all k and certain integers e and f with $e > 0$. Therefore, $\deg \text{Hilb}_{I^k M}(t) = ek + f - 1 < ak$ for all k and any integer $a > \min\{e, e + f - 1\}$. By [4, 4.1.12] we have $H(I^k M, i) = P_{I^k M}(i)$ for all $i > \deg \text{Hilb}(t)$. In particular, $H(I^k M, ak) = P_{I^k M}(ak)$ for all k . Thus it remains to show that $H(I^k M, ak)$ is of polynomial type of degree $\leq n$ for a suitable a .

Let $W = M \otimes_S S[x_{n+1}]$; then $(I^k W)_j = \bigoplus_{i \leq j} (I^k M)_i x_{n+1}^{j-i}$, so that $H(I^k M, ak) = \dim_K (I^k W)_{ak}$ for all k . Hence we must show that $\dim_K (I^k W)_{ak}$ is of polynomial type of degree $\leq n$ for a suitable a . Since $I^k S[x_{n+1}]W = I^k W$, we may replace I^k by its extension $I^k S[x_{n+1}] = (IS[x_{n+1}])^k$ which, by an abuse of notation, we again denote by I^k .

Let c_k be the highest degree of a generator of $I^k W$, c the highest degree of a generator of I and d the highest degree of a generator of W . Then one has $c_k \leq kc + d$ for all k . Let $J = I_{\geq c} = \bigoplus_{j \geq c} I_j$. We choose $a > c + d$; then

$$\dim_K (J^k W)_{ak} = \dim_K ((J^k W)_{\geq c_k})_{ak} = \dim_K ((I^k W)_{\geq c_k})_{ak} = \dim_K (I^k W)_{ak}.$$

Thus we may replace I by J . Since J is generated in one degree, we may as well assume that I is generated in one degree, say c . In this case the Rees algebra $\mathcal{R}(I) = \bigoplus_{j \geq 0} I^j$ is naturally standard bigraded, the bigraded components being $\mathcal{R}(I)_{(g,h)} = (I^h)_{g+ch}$. Moreover, $V = \bigoplus_{k \geq 0} I^k W$ is a finitely generated bigraded $\mathcal{R}(I)$ -module with $V_{(g,h)} = (I^h W)_{g+ch}$.

If R is any standard bigraded K -algebra, and e and f are two natural numbers. Then the subalgebra $R_\Delta = \bigoplus_{k \geq 0} R_{(ke, kf)}$ is called the (e, f) -diagonal of R . It is easily seen that any such diagonal is a standard graded K -algebra. Moreover, if N is a finitely generated bigraded R -module, then the diagonal submodule $N_\Delta = \bigoplus_{k \geq 0} N_{(ek, fk)}$ is a finitely generated graded R_Δ -module.

We now consider the $(a - c, 1)$ -diagonal Δ of $\mathcal{R}(I)$. Then $V_\Delta = \bigoplus_{k \geq 0} (I^k W)_{ak}$ is a finitely generated graded A -module where A is the standard graded K -algebra with $A_k = (I^k)_{ak}$ for all k . It follows that $\dim_K(I^k W)_{ak} = \dim_K(V_\Delta)_k$ is of polynomial type of degree $\dim A - 1$. Since $\dim A < \dim \mathcal{R}(I) = n + 2$ (see [6, 1.1(i)]), the proof of the theorem is completed. \square

We close this section with the following result whose proof follows the same line of arguments as in the part of the proof of Theorem 1.1 where it is shown that $P_{I^k M}(ak)$ is of polynomial type of degree $\leq n$.

Let $A = K[x_1, \dots, x_n, y_1, \dots, y_m]$ be the standard bigraded polynomial ring, that is, we have degree $x_i = (1, 0)$ and degree $y_j = (d_j, 1)$ for all i and j . Furthermore, let E be a finitely generated graded bigraded S -module. For all k we set $E_k = \bigoplus_i E_{i,k}$. Then each E_k is a finitely generated graded S -module, where $S = K[x_1, \dots, x_n]$, and we may consider its Hilbert polynomial $P_{E_k}(x)$ and its Hilbert coefficients $e_i(E_k)$.

Theorem 1.2. *For $k \gg 0$ the dimension of E_k is constant, say d . Then for $i = 0, \dots, d$, the i th Hilbert coefficient $e_i(E_k)$ as a function of k is of polynomial type of degree $\leq n + m - d + i$.*

2. ASYMPTOTIC BEHAVIOR OF MULTIPLICITY; THE GRADED CASE

Let $J \subset I \subset S$ be graded ideals, M graded S -modules and $N \subset M$ a graded submodule of M . Then $J^k N \subset I^k M$. Since by Theorem 1.1 the Hilbert coefficients of $M/I^k M$ and $N/J^k N$ as functions of k are of polynomial type, it follows from the equation

$$(3) \quad P_{I^k M/J^k N}(x) = P_{M/N}(x) + P_{N/J^k N}(x) - P_{M/I^k M}(x)$$

that the degree of $P_{I^k M/J^k N}(x)$ is constant for large k . In other words, there exists an integer k_0 such that $\dim(I^k M/J^k N) = d$ for all $k \geq k_0$.

More generally, if $(M_k)_{k \geq 0}$ is family of finitely generated S -modules with the property that there exists an integer k_0 such that $\dim M_k$ is constant for all $k \geq k_0$, then we call $\dim M_{k_0}$ the *limit dimension* of $(M_k)_{k \geq 0}$.

Let $(a_k)_{k=1,2,\dots}$ be a sequence of real numbers. We write $\lim_{k \rightarrow \infty} a_k \in \mathbb{Q}$ to indicate that $\lim_{k \rightarrow \infty} a_k$ exists and that the limit is a rational number.

Proposition 2.1. *Let d be the limit dimension of $(I^k M/J^k N)_{k \geq 0}$. Then*

$$\lim_{k \rightarrow \infty} \frac{e_0(I^k M/J^k N)}{k^{n-d}} \in \mathbb{Q}.$$

Proof. Let $d_1 = \dim M/IM$, $d_2 = \dim N/JN$ and $c = \dim M/N$. Then $\dim M/I^k M = d_1$ and $\dim N/J^k N = d_2$ for all k . It follows from equation (3) that

$$e_0(I^k M/J^k N) = e_{c-d}(M/N) + e_{d_2-d}(N/J^k N) - e_{d_1-d}(M/I^k M) \quad \text{for } k \gg 0.$$

Hence Theorem 1.1 implies that $e_0(I^k M/J^k N)$ is of polynomial type of degree $\leq n - d$, and this implies that $\lim_{k \rightarrow \infty} e_0(I^k M/J^k N)/k^{n-d}$ exists and is a rational number. \square

Let \overline{J} denote the integral closure of an ideal J . As an immediate consequence of Proposition 2.1 we obtain

Corollary 2.2. *Let $I \subset S$ be a graded ideal. Then $(\overline{I^k}/I^k)_{k \geq 0}$ admits a limit dimension, say d , and*

$$\lim_{k \rightarrow \infty} \frac{e_0(\overline{I^k}/I^k)}{k^{n-d}} \in \mathbb{Q}.$$

Proof. Since $A = \bigoplus_{k \geq 0} \overline{I^k}$ is equal to the integral closure of the Rees algebra $\mathcal{R}(I)$, it follows that A is a finitely generated graded $\mathcal{R}(I)$ -module. In particular it follows that there exists an integer k_0 such that $A_k = I^{k-k_0} A_{k_0}$ for all $k \geq k_0$. In other words, we have

$$\overline{I^k}/I^k = I^{k-k_0} \overline{I^{k_0}}/I^{k-k_0} I^{k_0} \quad \text{for all } k \geq k_0.$$

Thus the assertion follows from Proposition 2.1. \square

We give another application of Proposition 2.1.

Corollary 2.3. *Let $I \subset S$ be a monomial ideal and let $\mathfrak{m} = (x_1, \dots, x_n)$ be the graded maximal ideal of S . Then $((I^k : \mathfrak{m})/I^k)_{k \geq 0}$ admits a limit dimension, say d , and*

$$\lim_{k \rightarrow \infty} \frac{\ell((I^k : \mathfrak{m})/I^k)}{k^{n-d}} \in \mathbb{Q}.$$

Proof. We will show that $M = \bigoplus_{k \geq 0} I^k : \mathfrak{m}$ is a finitely generated $\mathcal{R}(I)$ -module. Then we proceed as in the proof of Corollary 2.2.

We first notice that $I^k : \mathfrak{m} = \bigcap_{i=1}^n (I^k : x_i)$. This implies that $M = \bigcap_{i=1}^n M_i$ where $M_i = \bigoplus_{k \geq 0} I^k : x_i$. Since $\mathcal{R}(I)$ is Noetherian it suffices therefore to show that each M_i is a finitely generated $\mathcal{R}(I)$ -module. In fact, we will show that M_i is generated by its degree 1 component. In other words, we will show that $I^k : x_i = (I^{k-1} : x_i)I$ for all $k > 1$.

Since I is a monomial ideal we can write $I = Jx_i + L$ where J and L are again monomial ideals, and where none of the generators of L is divisible by x_i . Then one has

$$I^k = \sum_{j=1}^k J^j L^{k-j} x_i^j + L^k, \quad \text{and hence} \quad I^k : x_i = \sum_{j=1}^k J^j L^{k-j} x_i^{j-1} + L^k.$$

Similarly we have $I^{k-1} : x_i = \sum_{j=1}^{k-1} J^j L^{k-1-j} x_i^{j-1} + L^{k-1}$. It follows that

$$\begin{aligned} (4) \quad (I^{k-1} : x_i)I &= \left(\sum_{j=1}^{k-1} J^j L^{k-1-j} x_i^{j-1} + L^{k-1} \right) (Jx_i + L) \\ &= A + B + L^{k-1} Jx_i + L^k, \end{aligned}$$

with

$$A = \sum_{j=1}^{k-1} J^{j+1} L^{k-1-j} x_i^j \quad \text{and} \quad B = \sum_{j=1}^{k-1} J^j L^{k-j} x_i^{j-1}.$$

Observe that $L^{k-1} J x_i$ appears as a multiple of a summand in B , so we may omit it in the right hand sum of (4). Next we may rewrite the sum A as $\sum_{j=2}^k J^j L^{k-j} x_i^{j-1}$. Then it follows from (4) that

$$(I^{k-1} : x_i) I = \sum_{j=1}^k J^j L^{k-j} x_i^{j-1} + L^k = I^k : x_i,$$

as desired. \square

Let I and J be graded ideals in S . The k th symbolic power of I with respect to J is defined to be the graded ideal $I_k(J) = I^k : J^\infty$.

We want to single out two cases of interest: in the first case, let J be the graded maximal ideal of S . Then $I_k(J)$ is the k th saturated power of I , denoted $(I^k)^{\text{sat}}$.

In the second case, consider the set $A^*(I) = \bigcup_k \text{Ass}(S/I^k)$ of asymptotic prime ideals of I . This set is known to be finite (see [2]). Of course the set on minimal prime ideals $\text{Min}(I)$ of I is a subset of $A^*(I)$. Let

$$J = \bigcap_{P \in \text{Ass}^*(I) \setminus \text{Min}(I)} P.$$

For this choice of J one obtains the ordinary symbolic powers $I^{(k)}$ of I .

We are interested in the limit behavior of $e_0((I_k(J)/I^k))$. In [8, 2.2] there is given an example that shows that $\ell((I^k)^{\text{sat}}/I^k)$ need not to be of polynomial type, and not even of quasi-polynomial type. However we will see that if I and J are monomial ideals, then the limit behavior of $e_0(I_k(J)/I^k)$ is quite nice. This is mainly due to the fact that in this situation, the graded ring $\bigoplus_{k \geq 0} I_k(J)$ is a finitely generated S -algebra, as shown in [11, 3.2].

More generally, suppose that we are given two families $(I_k)_{k \geq 0}$ and $(J_k)_{k \geq 0}$ of graded ideals of S with the property that $J_k J_\ell \subset J_{k+\ell}$ and $I_k I_\ell \subset I_{k+\ell}$ for all k and ℓ , and such that the algebras $A = \bigoplus_{k \geq 0} J_k$ and $B = \bigoplus_{k \geq 0} I_k$ are finitely generated. We also assume that $J_k \subset I_k$ for all k and that $I_0 = J_0 = S$.

We first show

Proposition 2.4. *The numerical function $e_0(I_k/J_k)$ is of quasi-polynomial type.*

Proof. Since A and B are finitely generated graded S -algebras, there exist integers s and t such that the Veronese algebra $A^{(s)}$ of A and the Veronese subalgebra $B^{(t)}$ of B is standard graded, see for example [11, 2.1]. We may assume that $s = t$. Otherwise we replace s as well as t by st .

As an $A^{(s)}$ -module the algebra A is a direct sum of the graded $A^{(s)}$ -modules $A^{(s;i)} = \bigoplus_{k \geq 0} A_{ks+i}$. Indeed, $A = \bigoplus_{i=0}^{s-1} A^{(s;i)}$. Similarly, $B = \bigoplus_{i=0}^{s-1} B^{(s;i)}$.

Since $A^{(s;i)}$ is a finitely generated $A^{(s)}$ -module, there exist an integer v_i such that $A_{(k+v_i)s+i} = A_{ks}A_{v_i s+i}$ for all $k \geq 0$. Let $v = \max\{v_i : i = 0, \dots, s-1\}$. Replacing each v_i by v we may assume that all v_i are equal to v . Similarly there exists an integer w such that $B_{(k+w)s+i} = B_{ks}B_{ws+i}$ for all $k \geq 0$ and all $i = 0, \dots, s-1$. Taking the maximum of v and w we may as well assume that $v = w$.

Thus we may assume that A and B are standard graded and have to show that $e_0(I_1^k M/J_1^k N)$ is of polynomial type, where M and N are graded S -modules. But this has already been shown in Proposition 2.1. \square

A priori it is not clear that $\dim I_k/J_k$ is constant for $k \gg 0$. But if we assume that A is standard, that is $J_k = J^k$ for all k where $J = J_1$, then we have

Theorem 2.5. *With the notation introduced, suppose that A is standard graded. Let P_0, \dots, P_{g-1} be the polynomials such that $e_0(I_{mg+i}/J^{mg+i}) = P_i(m)$ for $m \gg 0$. Then*

- (a) $\dim I_k/J^k$ is constant for $k \gg 0$.
- (b) $\deg P_i = \deg P_0$ for all i and all of them have the same leading coefficient.

Corollary 2.6. *Let I and J be monomial ideals, and let d be the limit dimension of $(I_k(J)/I^k)_{k \geq 0}$. Then*

$$\lim_k \frac{1}{k^{n-d}} e_0 \left(\frac{I_k(J)}{I^k} \right) \in \mathbb{Q}.$$

In particular,

$$\lim_k \frac{1}{k^n} \ell \left(\frac{(I^k)^{\text{sat}}}{I^k} \right) \in \mathbb{Q}.$$

In order to prove Theorem 2.5 we need to introduce some definitions. Let $M = \bigoplus_{k \geq 0} M_k$ be a bigraded $A = \bigoplus_k J^k$ -module over the bigraded algebra A such that each M_k is a finitely generated S -module. All modules considered in the sequel will be of this kind.

The module M is called *quasi-finite*, if $H_{A_+}^0(M)_i = 0$ for $i \gg 0$. Here $H_{A_+}^0(M)$ denotes the 0th local cohomology of M with respect to the ideal $A_+ = \bigoplus_{k > 0} A_k$. We consider quasi-finite modules because in the application we have in mind the modules $\bigoplus_{k \geq 0} I_k(J)/I^k$ are in general not finitely generated $\bigoplus_k I_k$ modules, but only quasi-finite, as we shall see.

We say an element $x \in A_1$ is *M -filter regular* if $(0 :_M x)_i = 0$ for $i \gg 0$. We do not require that x is bihomogeneous. The existence of an M -filter regular element is useful due to the following:

Proposition 2.7. *Let $M = \bigoplus_{k \geq 0} M_k$ be an A -module such that $e_0(M_k)$ is a quasi-polynomial. Let P_0, \dots, P_{g-1} be the polynomials such that $e_0(M_{mg+i}) = P_i(m)$ for $m \gg 0$. If there exists an element in A_1 which is M -filter regular, then*

- (a) $\dim M_k$ is constant for $k \gg 0$.
- (b) $\deg P_i = \deg P_0$ for all i and all of them have the same leading coefficient.

Proof. (a) Suppose for $m \geq s$ we have $e_0(M_{mg+i}) = P_i(m)$ for $i = 0, \dots, g-1$. We choose $x \in A_1$ which is M -filter regular. Then the map $M_j \rightarrow M_{j+1}$ induced by multiplication with x is injective for all $j \gg 0$. This implies that for $m \gg 0$ we have

$$\dim M_{mg} \leq \dim M_{mg+1} \leq \dots \leq \dim M_{mg+g-1} \leq \dim M_{(m+1)g}.$$

The assertion follows.

(b) We assert that $e_0(M_k) \leq e_0(M_{k+1})$ for all $k \gg 0$. Let d be the limit dimension of the family $(M_k)_{k \geq 0}$. If $d = 0$, then $e_0(M_k) = \ell(M_k)$ and the assertion is obvious, because the maps $M_k \rightarrow M_{k+1}$ induced by multiplication by x are injective for $k \gg 0$.

So now let $d > 0$, and let c be the initial degree of the element x in the graded module A_1 . Then the maps $M_k \rightarrow M_{k+1}$ induce maps $M_{k,i \geq s} \rightarrow M_{k+1,i \geq s+c}$ for all s , where $M_{k,i \geq t} = \bigoplus_{i \geq t} M_{k,i}$.

Notice that $e_0(M_k) = e_0(M_{k,i \geq s}) \forall s$ and that both modules, $M_{k,i \geq s}$ and $M_{k+1,i \geq s+c}$ are generated in a single degree for s large enough. We choose such an s . Let \mathfrak{m} be the graded maximal ideal of S . Since $M_{k,i \geq s}$ is generated in a single degree, the associated graded module $\text{gr}_{\mathfrak{m}}((M_{k,i \geq s})_{\mathfrak{m}})$ is isomorphic to $M_{k,i \geq s}$, up to a shift. Hence $e_0(\mathfrak{m}, (M_{k,i \geq s})_{\mathfrak{m}}) = e_0(M_{k,i \geq s})$. On the other hand we have that $(M_{k,i \geq s})_{\mathfrak{m}} \rightarrow (M_{k+1,i \geq s+c})_{\mathfrak{m}}$ is injective for all $k \gg 0$. Since both modules have the same dimension, the inequality in the next display follows from [4, 4.7.7].

$$e_0(M_k) = e_0(\mathfrak{m}, (M_{k,i \geq s})_{\mathfrak{m}}) \leq e_0(\mathfrak{m}, (M_{k+1,i \geq s+c})_{\mathfrak{m}}) = e_0(M_{k+1}).$$

Thus for $m \geq s$ we have that

$$(5) \quad P_0(m) \leq P_1(m) \leq \dots \leq P_i(m) \leq \dots \leq P_{g-1}(m) \leq P_0(m+1).$$

The inequalities in (5) imply that all P_i have the same degree. Let s_i be the leading coefficient of P_i . As all P_i have the same degree note that (5) implies that

$$s_0 \leq s_1 \leq \dots \leq s_{g-1} \leq s_0.$$

Thus $s_0 = s_1 = \dots = s_{g-1}$. □

Next we discuss the existence of a filter regular element for a quasi-finite module. Set $N = M/H_{A_+}^0(M)$. If M is finitely generated as an A -module, then $x \in A_1$ is M -filter regular precisely when x is N -regular, see [25, 2.1]. Thus M -filter regular elements always exists if K is infinite. For quasi-finite modules we have

Lemma 2.8. *Let $M = \bigoplus_{k \geq 0} M_k$ be a quasi-finite A -module. Set $N = M/H_{A_+}^0(M)$. Then*

- (a) $\text{Ass}_A N \cap \text{Var}(A_+) = \emptyset$.
- (b) $\text{Ass}_A N$ is a countable set.
- (c) If $x \in A_1$ is N -regular, then it is M -filter regular.

(d) If K is uncountable, then there exists $x \in A_1$ which is N regular and so M -filter regular.

Proof. (a) Suppose there exists $\mathfrak{p} \in \text{Ass}_A N \cap \text{Var}(A_+)$. Let $\mathfrak{p} = (0: \bar{x})$ where $\bar{x} = x + H_{A_+}^0(M)$ is a nonzero homogeneous element of N . As $\mathfrak{p}\bar{x} = 0$ we have $E = \mathfrak{p}x \subseteq H_{A_+}^0(M)$. Notice E is finitely generated as an A -module. So $A_+^m E = A_+^m \mathfrak{p}x = 0$. Since $\mathfrak{p} \supseteq A_+$ we have that $A_+^{m+1}x = 0$. Therefore $x \in H_{A_+}^0(M)$, a contradiction.

(b) For $i \geq 0$, set D_i to be the A -submodule of N generated by N_0, \dots, N_i . As each N_i is a finitely generated S -module, it follows that each D_i is a finitely generated A -module. The assertion follows since $N = \bigcup_{i \geq 0} D_i$.

(c) Let x be N -regular. As M is quasi-finite we have $M_i = N_i$ for $i \gg 0$. It follows that $(0: {}_M x)_i = 0$ for $i \gg 0$. Thus x is M -filter regular.

(d) By (b) we get that $\text{Ass}_{A_+} N$ is countable. Say $\text{Ass}_A N = \{Q_1, Q_2, \dots\}$. Then for each $i \geq 1$ we have that $Q_i \cap A_1$ is a proper graded S -submodule of A_1 . Since K is uncountable, there exists $x \in A_1 \setminus \bigcup_{i \geq 1} Q_i$. Clearly x is N -regular. \square

Proof of 2.5. We may assume K is uncountable. If not, we choose a field extension K' of K which is uncountable, and set $S' = S \otimes_K K'$, $A' = A \otimes_S S'$ and $B' = B \otimes_S S'$. It can be checked that $e_0(B'_k/A'_k) = e_0(B_k/A_k)$ for all $k \geq 1$.

Set $M = B/A$. The exact sequence $0 \rightarrow A \rightarrow B \rightarrow B/A \rightarrow 0$ yields an exact sequence

$$0 \rightarrow H_{A_+}^0(A) \rightarrow H_{A_+}^0(B) \rightarrow H_{A_+}^0(B/A) \rightarrow H_{A_+}^1(A) \rightarrow$$

We assert that B/A is a quasi-finite A -module. In fact, $H_{A_+}^0(B) = H_{A_+B}^0(B) = 0$ because B is a domain, and $H_{A_+}^1(A)_k = 0$ for $k \gg 0$ in general [3, 15.1.5]. Thus the long exact cohomology sequence implies that $H_{A_+}^0(B/A)_k = 0$ for $k \gg 0$, as desired. By Lemma 2.8(d) we have that there exists M -filter regular elements. So by Proposition 2.7 the result follows. \square

3. ASYMPTOTIC BEHAVIOR OF MULTIPLICITY IN THE LOCAL CASE

Let (R, \mathfrak{m}) be a local ring. If M is a finitely generated R -module then $e_0(M)$ denotes the Hilbert-Samuel multiplicity of M with respect to \mathfrak{m} . Let $\mathcal{F} = \{I_n\}_{n \geq 0}$ be a Noetherian filtration on R i.e., a chain of ideals

$$R = I_0 \supseteq I_1 \supseteq \dots \supseteq I_i \supseteq I_{i+1} \supseteq \dots \text{ such that } I_i I_j \subseteq I_{i+j} \text{ for all } i, j \text{ and}$$

$\mathcal{R}(\mathcal{F}) = \bigoplus_{k \geq 0} I_k$ is a graded (not necessarily standard) Noetherian R -algebra.

Example 3.1. Let $I^{(k)}$ denote the k^{th} symbolic power of I_1 . There are many cases known when $\mathcal{F} = \{I^{(k)}\}_{k \geq 0}$ is Noetherian; see for example [7] [13] and [10]. The symbolic power filtration need not be Noetherian in general [22].

The main result of this section is the local analogue of Theorem 2.5

Theorem 3.2. *Let J be an ideal in R with positive grade such that $J \subseteq I_1$. Then*

- (a) $\dim I_k/J^k$ is constant for $k \gg 0$.
- (b) There exists polynomials P_0, \dots, P_{g-1} such that $P_i(m) = e_0(I_{mg+i}/J^{mg+i})$ for all $m \gg 0$. In other words, $e_0(I_k/J^k)$ is of quasi-polynomial type of period g .
- (c) $\deg P_i = \deg P_0$ for all i and all of them have the same leading coefficient.

It is convenient to prove Theorem 3.2 by considering the following more general setup: let $A \subseteq B$ be a homogeneous inclusion of Noetherian graded algebras with $R = A_0 = B_0$ a local ring with maximal ideal \mathfrak{m} . We assume A is a standard graded R -algebra, while B is *not necessarily* a standard graded R -algebra.

A typical example of this situation is when $B = \mathcal{R}(\mathcal{F})$ and $A = \mathcal{R}(J)$ with $J \subset I_1$.

In analogy to the graded case we call a graded A -module $M = \bigoplus_{k \geq 0} M_k$ *quasi-finite* if each M_k is a finitely generated R -module and $H_{A_+}^0(M)_j = 0$ for all $j \gg 0$.

Note that B is trivially a quasi-finite A -module if $\text{grade}(A_1 B, B) > 0$. Notice if A_n, B_n are ideals in R with $\text{grade}(A_1, R) > 0$, then $\text{grade}(A_1 B, B) > 0$, and so B is a quasi-finite A -module.

In this more general frame Theorem 3.2 reads as follows:

Theorem 3.3. *Assume that B is quasi-finite A -module. Then*

- (a) $\dim B_k/A_k$ is constant for $k \gg 0$.
- (b) There exists polynomials P_0, \dots, P_{g-1} such that $P_i(m) = e_0(B_{mg+i}/A_{mg+i})$ for all $m \gg 0$. In other words, $e_0(B_k/A_k)$ is of quasi-polynomial type of period g .
- (c) $\deg P_i = \deg P_0 \leq \dim B + d - 1$ for all i and all of them have the same leading coefficient. Here d is the limit dimension of $(B_k/A_k)_{k \geq 0}$.

We give an example which shows that the conclusion of the theorem would fail without the quasi-finite condition on B .

Example 3.4. Let $R = K[[x]]$ be the power series ring over a field K . Set $A = R[u, v]$ and $B = R[u, v, w]/(wu, wv)$. We give w degree 2. Thus $wB \subseteq H_{A_+}^0(B)$. Thus B is not a quasi-finite A -module. It can be easily seen that $B_k/A_k = 0$ if k is odd and $Rw^{k/2}$ if k is even. So $g = 2$ and the multiplicity polynomial is $(0, 1)$.

One of the chief problems in trying to prove the above theorem is that B/A is not a B -module. It is also, in general, not a finitely generated A -module. Rees's technique [19], in this case is to consider the Rees ring $S = \bigoplus_{k \geq 0} (A_+ B)^k$ and the associated graded ring $G = \text{gr}_{A_1 B} B$ of the ideal $A_1 B$ in B .

Observe that $(A_+ B)^k = \bigoplus_{i \geq 0} A_k B_i$, so that $S = \bigoplus_{i, j} A_i B_{j-i}$. Hence S is a naturally bigraded R -algebra with $S_{ij} = A_i B_{j-i}$ for all i, j . We have $S_{i, j} = 0$ if $j < i$ and $S_{i, i} = A_i$.

Notice that G inherits a bigraded structure from S , so that $G = \bigoplus_{i \geq 0, j \geq 0} G_{i, j}$ with

$$G_{i, j} = \frac{S_{i, j}}{S_{i, j+1}} = \frac{B_{i-j} A_j}{B_{i-j-1} A_{j+1}}.$$

G also has the property that $G_{i,j} = 0$ if $j < i$ and $G_{i,i} = A_i$. If we consider A as bigraded algebra with $A_{i,j} = 0$ if $i \neq j$ and $A_{i,i} = A_i$ then A becomes a bigraded R -subalgebra of G . Finally let $L = \bigoplus_{j>i} G_{i,j}$. Notice that L is an ideal of G .

We now consider G, L singly graded by summing across columns, i.e, set $G = \bigoplus_{i \geq 0} G_i$, $L = \bigoplus_{i \geq 0} L_i$ where

$$G_i = \bigoplus_{j \geq 0} G_{i,j} = \bigoplus_{j=0}^i G_{i,j} \quad \text{and} \quad L_i = \bigoplus_{j \geq 0} L_{i,j}.$$

The advantage of Rees's technique is due to the following observation: consider the filtration of R -modules

$$(6) \quad A_i \subseteq B_1 A_{i-1} \subseteq B_2 A_{i-2} \subseteq \dots \subseteq B_{i-1} A_1 \subseteq B_i.$$

One can check that

$$(7) \quad L_i = \bigoplus_{j=0}^{i-1} \frac{B_{i-j} A_j}{B_{i-(j+1)} A_{j+1}}.$$

This leads to the following

Lemma 3.5. *With the notation introduced we have an inclusion of graded algebras $A \subseteq G$ and a finitely generated G -module L with*

- (a) $\dim_R B_k/A_k = \dim_R L_k$ for all $k \geq 1$.
- (b) $e_0(B_k/A_k) = e_0(L_k)$ for all $k \geq 1$.
- (c) $\dim(G) = \dim(B)$.
- (d) $e_0(B_k/A_k) = e_0(L_k)$ is of quasi-polynomial type.

Proof. The first two statements follow from (6) and (7). For the third statement see [4, 4.5.6] Finally, statement (d) follows from Proposition 5.5. \square

Note that assertion (b) of Theorem 3.3 follows from statement (d) above. The remaining assertions of Theorem 3.3 depend on the existence of a B/A -filter regular element in A_1 . One proves its existence just as in the graded case by using a local version of Proposition 2.7 and Proposition 2.8.

Proof of 3.3 (a) and (c). If $K = R/\mathfrak{m}$ is not uncountable then we do the following standard trick. First we complete R . Then we consider $R' = \widehat{R}[[x]]_{\mathfrak{m}[[x]]}$ where \widehat{R} is the completion of R . Notice that R' has residue field $K((x))$, which is uncountable. Set $A' = A \otimes_R R'$ and $B' = B \otimes_R R'$. Since A' is a flat extension of A it follows that B' is also a quasi-finite A' -module. It can be checked that $e_0(B'_k/A'_k) = e_0(B_k/A_k)$ for all $k \geq 1$.

Similarly as in the proof of Theorem 2.5 it follows that B/A is a quasi-finite A -module. By the local version of Proposition 2.8(d) there exists a B/A -filter regular element in A_1 . So by the local version of Proposition 2.7 the results follow, except the degree bound for the P_i . But this is a consequence of Proposition 5.5. \square

Our main theorem is an easy corollary of Theorem 3.3.

Proof of Theorem 3.2. Let $A = \mathcal{R}(J)$ and $B = \mathcal{R}(\mathcal{F})$. Since $J \subseteq I_1$ we have an inclusion of graded algebras $A \subseteq B$. Since $\text{grade}_R J > 0$ we get that $\text{grade}_B A_1 B > 0$. So $H_{A_+}^0(B) = H_{A_1 B}^0(B) = 0$. In particular, B is a quasi-finite A -module. The result now follows from Theorem 3.3. \square

4. THE REES POLYNOMIAL OF I AND J

Throughout this section, let (R, \mathfrak{m}) be a local ring of dimension d . For an \mathfrak{m} -primary ideal I , put $H(I, n) = \ell(R/I^n)$ where ℓ denotes length. Let $P(I, n)$ denote the Hilbert polynomial corresponding to the Hilbert function $H(I, n)$. We write $P(I, n)$ in the form:

$$P(I, n) = e_0(I) \binom{n+d-1}{d} - e_1(I) \binom{n+d-2}{d-1} + \cdots + (-1)^d e_d(I).$$

Recall that an ideal $J \subset I$ is called a reduction of I if $JI^n = I^{n+1}$ for some positive integer n . The Rees multiplicity theorem asserts that if R is a quasi-unmixed local ring then J is a reduction of I if and only if $e_0(I) = e_0(J)$. The Rees multiplicity theorem has been generalized for ideals that are not \mathfrak{m} -primary by a number of authors. In [21] Rees provided one such generalization which we now state. Let $J \subset I$ be ideals of R such that $\ell(I/J)$ is finite. Then the numerical function $H(I/J, n) = \ell(I^n/J^n)$ is given by a polynomial $P(I/J, n)$ for large n , [1]. We call $H(I/J, n)$ and $P(I/J, n)$ the *Rees function* and the *Rees polynomial* of the pair (J, I) respectively. Rees showed that if (R, \mathfrak{m}) is quasi-unmixed then J is a reduction of I if and only if $\deg P(I/J, n) < d$. It is natural to ask for the exact degree of $P(I/J, n)$. In case I and J are \mathfrak{m} -primary it is easy to see that $\deg P(I/J, n) < d - k$ if and only if $e_i(I) = e_i(J)$ for $i = 0, 1, \dots, k$ for all $k = 0, 1, \dots, d$. K. Shah [23] introduced coefficient ideals which characterize the degree of $P(I/J, n)$ when I and J are \mathfrak{m} -primary.

Theorem 4.1 (K. Shah). *Let (R, \mathfrak{m}) be a quasi-unmixed local ring of positive dimension d with infinite residue field R/\mathfrak{m} . Let I be an \mathfrak{m} -primary ideal. Then there exist unique largest ideals I_1, I_2, \dots, I_d containing I such that $e_i(I) = e_i(I_k)$ for $i = 0, 1, \dots, k$ and $I \subset I_d \subset I_{d-1} \subset \cdots \subset I_1 \subset I_0 = \bar{I}$. The ideal I_k is called the k th-coefficient ideal of I .*

We will prove an analogue of the above theorem for non- \mathfrak{m} -primary ideals which will help us in determining the degree of $P(I/J, n)$. The proof is similar to the proof of Shah's theorem. First we need the following

Definition 4.2. *Let (R, \mathfrak{m}) be a local ring and let I be an ideal in R . Let $I^{\text{sat}} = \bigcup_{n \geq 1} (I : \mathfrak{m}^n)$ be the saturation of I . We call $\mathfrak{q}(I) = \bar{I} \cap I^{\text{sat}}$ the relative integral closure of I .*

In several results in this section we consider ideals $J \subseteq I$ so that J is a reduction of I and $\ell(I/J) < \infty$. For these cases we have:

Proposition 4.3. *Let (R, \mathfrak{m}) be a local ring and I an ideal of R . Then*

- (1) I is a reduction of $\mathfrak{q}(I)$ and $\ell(\mathfrak{q}(I)/I) < \infty$.
- (2) If I is a reduction of K with $\ell(K/I) < \infty$ then $K \subseteq \mathfrak{q}(I)$.
- (3) If $J \subseteq I$ then $\mathfrak{q}(J) \subseteq \mathfrak{q}(I)$.
- (4) $\mathfrak{q}(\mathfrak{q}(I)) = \mathfrak{q}(I)$.

Proof. The assertions (1), (2), (3) are easy to show. We prove (4). If $\mathfrak{q}(I)$ is a reduction of K with $\ell(K/\mathfrak{q}(I)) < \infty$ then note that K is also a reduction of I and also $\ell(K/I) < \infty$. So by (2) we have $K \subseteq \mathfrak{q}(I)$. Thus $K = I$. By taking $K = \mathfrak{q}(\mathfrak{q}(I))$ we get the desired result. \square

We now give an analogue of Shah's Theorem.

Theorem 4.4. *Let (R, \mathfrak{m}) be a d -dimensional quasi-unmixed local ring with infinite residue field R/\mathfrak{m} . Let I be any ideal of analytic spread a . Then for $k = 1, 2, \dots, a$, there exist unique largest ideals I_k with $\ell(I_k/I) < \infty$, and $I \subseteq I_a \subseteq I_{a-1} \subseteq \dots \subseteq I_1 \subseteq \mathfrak{q}(I)$, such that $\deg P(I_k/I, n) < a - k$ for $k = 1, 2, \dots, a$.*

Proof. For each $k = 1, 2, \dots, a$, consider the sets

$$V_k = \{L \mid L \text{ is an ideal of } R \text{ with } \ell(L/I) < \infty, \deg P(L/I, n) < a - k\}.$$

If $L \in V_k$, then $\deg P(L/I, n) < d - 1$. Hence by Rees's theorem, I is a reduction of L . Hence $L \subseteq \mathfrak{q}(I)$.

Since $I \in V_k$ and R is Noetherian, V_k has a maximal member, say J . We show that J is unique. Let $L \in V_k$ and $x \in L$. Since $I \subset (I, x) \subset L$, for all $n \geq 1$, $\ell((I, x)^n/I^n) \leq \ell(L^n/I^n) < \infty$. Hence $\deg P((I, x)/I, n) \leq \deg P(L/I, n) < a - k$. By Rees's theorem, I is a reduction of (I, x) . Let $I(I, x)^t = (I, x)^{t+1}$ for some integer t . Then $x^{t+1} \in I(I, x)^t \subset J(J, x)^t$. Hence $J(J, x)^t = (J, x)^{t+1}$. It follows that for all $n \geq t$, $J^{n-t}(J, x)^t = (J, x)^n$. Since $\ell((I, x)/I) < \infty$, there exists an r such that $x\mathfrak{m}^r \subset I \subset J$. Hence $\ell((J, x)/I) < \infty$. Now we show that $\deg P((J, x)/I, n) < a - k$. For all $n \geq t$, we have:

$$\begin{aligned} \ell((J, x)^n/I^n) &= \ell(J^{n-t}(J, x)^t/I^n) = \ell((J^n, J^{n-1}x, \dots, x^t J^{n-t})/I^n) \\ &\leq \sum_{i=1}^t \ell((J^n + x^i J^{n-i})/I^n) \leq \sum_{i=1}^t [\ell(J^n/I^n) + \ell((I^n + x^i J^{n-i})/I^n)] \\ &= \sum_{i=1}^t \left[\ell(J^n/I^n) + \ell((x^i I^{n-i} + I^n)/I^n) + \ell\left(\frac{x^i J^{n-i} + I^n}{x^i I^{n-i} + I^n}\right) \right] \\ &\leq \sum_{i=1}^t \ell(J^{n-i}/I^{n-i}) + \ell((I, x)^n/I^n) + \ell(J^n/I^n). \end{aligned}$$

Hence $\deg P((J, x)/I, n) < d - k$. Therefore $(J, x) \in V_k$. By maximality of J , we conclude that $x \in J$. Hence $L \subset J$ and therefore J is unique. \square

Corollary 4.5. *Let (R, \mathfrak{m}) be a quasi-unmixed local ring. Let $I \subset J$ be ideals of R such that $\ell(J/I) < \infty$. Then $\deg P(J/I, n) = a - k$ if and only if $J \subset I_{k-1}$ but $J \not\subset I_k$ for $k = 1, 2, \dots, a$.*

As a consequence of the next result, we identify the degree of the Rees polynomial of J and I .

Proposition 4.6. *Let $R = \bigoplus_{i=0}^{\infty} R_n$ be a standard graded Noetherian ring over a Noetherian local ring (R_0, \mathfrak{m}) . Let $M = \bigoplus_{i=0}^{\infty} M_n$ be a finitely generated graded R -module with $\ell(M_n) < \infty$ for all $n \geq 0$. Let $H(M, n) = \ell_{R_0}(M_n)$ be the Hilbert function of M . Then $\dim M = \dim M/\mathfrak{m}M$ and $H(M, n)$ is a polynomial function of degree $\dim M - 1$.*

Proof. Since M is a finitely generated R -module and $\ell(M_n) < \infty$, for all $n \geq 0$, there exists an r such that $\mathfrak{m}^r M_n = 0$ for all $n \geq 0$. Thus $\mathfrak{m}^r R \subset \text{ann}_R(M)$. Since $\text{Supp } M/\mathfrak{m}M = \text{Supp } M \cap V(\mathfrak{m}R)$, it follows that $\text{Supp } M = \text{Supp } M/\mathfrak{m}M$ and hence $\dim M = \dim M/\mathfrak{m}M$. Let $q = \bigoplus_{n=0}^{\infty} q_n = \text{ann}_R(M)$. Then each M_n is an R_0/q_0 -module. Let $M = \sum_{i=1}^g R m_i$ where $\deg m_i = d_i$ for $i = 1, 2, \dots, g$. Define the R -module homomorphism $\varphi : R \rightarrow M^{\oplus g}$ by $\varphi(r) = (rm_1, rm_2, \dots, rm_g)$. Then $\text{Ker } \varphi = \text{ann } M$. Restricting φ to R_0 , we see that R_0/q_0 is a submodule of an R_0 -module of finite length. Hence q_0 is \mathfrak{m} -primary. Let $S = R/\text{Ann } M = \bigoplus_{n=0}^{\infty} S_n$. Then M is a finitely generated S -module where S_0 is an Artin local ring. Hence by Hilbert-Serre theorem, $H(M, n)$ is a polynomial function of degree $\dim M - 1$. \square

Corollary 4.7. *Let (R, \mathfrak{m}) be a local ring of dimension d . Let $J \subset I$ be ideals of R such that J is a reduction of I and $\ell(I/J) < \infty$. Let $M = \mathcal{R}(I)/\mathcal{R}(J) = \bigoplus_{n=0}^{\infty} I^n/J^n$. Then $H(I/J, n)$ is a polynomial function of degree $\dim M - 1$.*

Proof. Since J is a reduction of I , $\mathcal{R}(I)$ and M are finite $\mathcal{R}(J)$ -module. Now apply the above proposition to M . \square

The maximum degree of the Rees polynomial $P(I/J, n)$ of a reduction J of an ideal I is $d - 1$. In the next theorem we give a sufficient condition so that this degree is $d - 1$.

Theorem 4.8. *Let (R, \mathfrak{m}) be a d -dimensional quasi-unmixed local domain. Let J be a proper reduction of an ideal I of R with $\ell(I/J) < \infty$. If the Rees ring $\mathcal{R}(J)$ satisfies Serre's condition S_2 , then $a(I) = d$ and $\deg P(I/J, n) = d - 1$.*

Proof. Let $A = \mathcal{R}(J)$ and $B = \mathcal{R}(I)$. Then A and B have same quotient field and B is a finite integral extension of A . Since A satisfies the S_2 condition, the conductor $A : B$ is a height one unmixed ideal. Indeed, let $B = Ab_1 + Ab_2 + \dots + Ab_n$ for some $b_j \in B$ for $j = 1, 2, \dots, n$. Let $b_j = x_j/y_j$ for some $x_j, y_j \in A$, for $j = 1, 2, \dots, n$. Then $A : B = \bigcap_{j=1}^n A : b_j = \bigcap_{j=1}^n (y_j : x_j)$. Since A satisfies the S_2 condition, principal ideals in A have

no embedded primary components. Hence $A : B$ is a height one unmixed ideal. Since $\text{Ann}_A M = A : B$, and A is universally catenary, we have $\dim M = \dim \mathcal{R}(J)/\text{Ann } M = d + 1 - 1 = d$. Hence $\deg P(I/J, n) = \dim M - 1 = \dim M/\mathfrak{m}M - 1 = d - 1 \leq a(J) - 1 \leq d - 1$. Therefore $a(J) = a(I) = d$. \square

Remark 4.9. *The relative integral closure can equal I if either $\overline{I} = I$ or $I^{\text{sat}} = I$. We call these as trivial cases. The following is a nice consequence to Theorem 4.8*

Corollary 4.10. *Let (R, \mathfrak{m}) be a d -dimensional quasi-unmixed local domain and let I be an ideal in R . If $a(I) < d$ and $\mathcal{R}(I)$ is S_2 then $\mathfrak{q}(I) = I$.*

Proof. If $\mathfrak{q}(I) \neq I$ then note that I is a proper reduction of $\mathfrak{q}(I)$ and $\ell(\mathfrak{q}(I)/I)$ is finite. So by Theorem 4.8 we get $a(\mathfrak{q}(I)) = a(I) = d$ which is a contradiction. \square

The degree of the Rees polynomial of a pair of normal ideals can be completely determined in analytically unramified quasi-unmixed local rings. This follows from the next theorem.

Theorem 4.11. *Let (R, \mathfrak{m}) be an analytically unramified quasi-unmixed local ring of dimension d . Let $J \subset I$ be ideals of R such that J is not a reduction of I and $\ell(I/J) < \infty$. Let $f(n) = \ell(\overline{I^n}/\overline{J^n})$. Then $f(n)$ is a polynomial function of degree d .*

Proof. Since $\ell(I/J) < \infty$, there is a positive integer r such that $\mathfrak{m}^r I \subseteq J$. Hence $\mathfrak{m}^r \overline{I} \subseteq \overline{J}$. Therefore $\ell(\overline{I}/\overline{J}) < \infty$. Since $\ell(I^n/J^n) < \infty$ for all n , it follows that $f(n) < \infty$ for all n . Consider the filtration of ideals

$$\overline{J^n} \subseteq \overline{I J^{n-1}} \subseteq \overline{I^2 J^{n-2}} \subseteq \dots \subseteq \overline{I^{n-1} J} \subseteq \overline{I^n}.$$

Then $f(n) = \sum_{i=0}^{n-1} \ell\left(\overline{J^i I^{n-i}}/\overline{J^{i+1} I^{n-i-1}}\right)$. The integral closure of the bigraded Rees algebra $\mathcal{R} := \mathcal{R}(I, J) = \bigoplus_{r,s \geq 0} I^r J^s u^r v^s$ in the polynomial ring $R[u, v]$ is the bigraded ring $\overline{\mathcal{R}} := \bigoplus_{r,s \geq 0} \overline{I^r J^s} u^r v^s$.

Since R is analytically unramified, $\overline{\mathcal{R}}$ is a finite \mathcal{R} -module. Indeed, by Theorem 1.4 of [20], there exists an integer t such that $\overline{I^r J^s} \subset I^{r-t} J^{s-t}$ for all $r, s \geq t$. For positive integers r and s , let $V_s = \bigoplus_{r=0}^{\infty} \overline{I^r J^s} \subset \bigoplus \overline{I^r}$ and $H_r = \bigoplus_{s=0}^{\infty} \overline{I^r J^s} \subset \bigoplus \overline{J^s}$. Then V_s is finite $\mathcal{R}(I)$ -module and H_r is a finite $\mathcal{R}(J)$ -module by Corollary 9.2.1 of [14]. Thus $\overline{\mathcal{R}}$ is a finite \mathcal{R} -module.

Now consider the ideals $A = \bigoplus_{r \geq 1, s \geq 0} \overline{I^r J^s}$ and $B = \bigoplus_{r \geq 1, s \geq 0} \overline{I^{r-1} J^{s+1}}$ of $\overline{\mathcal{R}}$. Then $C := A/B$ is a finitely generated bigraded \mathcal{R} -module. Since each bigraded component of C has finite length, it follows that $H(C; x, y) := \sum_{r \geq 1, s \geq 0} \ell(C_{rs}) x^r y^s$ is a rational function of the form $H(C; x, y) = h(x, y)/(1-x)^p(1-y)^q$ for some polynomial $h(x, y) \in \mathbb{Z}[x, y]$.

Note that $f(n) = \sum_{i=0}^{n-1} \ell(C_{(n-i,i)})$. Hence

$$\begin{aligned} \sum_{n=0}^{\infty} f(n)z^n &= \sum_{n=0}^{\infty} \sum_{i=0}^{n-1} \ell(C_{(n-i,i)})z^n \\ &= h(z, z)/(1-z)^{p+q}. \end{aligned}$$

Hence $f(n)$ is a polynomial function. It remains to show that its degree is d . Since R is analytically unramified, there exists an integer l such that $(\overline{I^l})^n = \overline{I^{ln}}$ and $(\overline{J^l})^n = \overline{J^{ln}}$ for all $n \geq 1$, by Corollary 9.2.1 of [14]. Since J is not a reduction of I , $\overline{J^l}$ is not a reduction of $\overline{I^l}$. Hence $\ell\left(\frac{(\overline{I^l})^n}{(\overline{J^l})^n}\right) = f(nl)$ is a polynomial function in n of degree d by [21]. Hence $f(n)$ is a polynomial function of degree d . \square

The following example shows that we cannot hope for a *relative minimal reduction* in general.

Example 4.12. *Let (R, \mathfrak{m}) be a d -dimensional Cohen-Macaulay ring and infinite residue field. Let I be an ideal with a minimal reduction $W = (u_1, \dots, u_s)$ which is generated by a regular sequence. Assume $0 < s < d$. Then if J is a reduction of I with $\ell(I/J)$ finite then there exists a reduction K of I with $K \subseteq J$, $K \neq J$ and $\ell(I/K) < \infty$.*

Proof. Let L be a minimal reduction (in the usual sense) of I such that $L \subseteq J$. Note that L is also generated by a regular sequence. Notice as R/L is Cohen-Macaulay of positive dimension we get in particular $L^{sat} = L$. In particular $\mathfrak{q}(L) = L$ and so $\ell(I/L) = \infty$.

As $\ell(I/J)$ is finite there exists c such that $\mathfrak{m}^c I \subseteq J$. Consider the sequence of ideals $J_n = L + \mathfrak{m}^n I$ where $n \geq c$. Then $J_n \subseteq J$, J_n is a reduction of I and $\ell(I/J_n) < \infty$. Since $\bigcap_{n \geq c} J_n = L$, it follows that atleast one $J_m \neq J$. Put $K = J_m$. \square

5. HILBERT-SAMUEL COEFFICIENTS OF $I^k M / I^{k+1} M$ AS A FUNCTION OF k

In this section (R, \mathfrak{m}) is a Noetherian local ring, I is an ideal in R , M is a finitely generated R -module and J is a reduction of I with respect to M , i.e., $I^{m+1} M = J I^m M$ for some m . If $\dim M = r$, then for $i = 0, 1, \dots, r$ let $e_i(M)$ be the i th Hilbert coefficient of M (with respect to \mathfrak{m}). In Corollary 5.4, we prove that $e_i(I^k M / I^{k+1} M)$, $e_i(I^k M)$ and $e_i(I^k M / J^k M)$ are polynomial functions in k for $k \gg 0$.

We might also ask similar questions for Hilbert coefficients with respect to an \mathfrak{m} -primary ideal in R . Our proofs for Question (1) and (3) in the introduction also covers this slightly more case. However for notational convenience and as a matter of taste we stick to the maximal ideal.

As in Section 3, it is convenient to consider standard graded R -algebras, $A = \bigoplus_{n \geq 0} A_n$ with $A_0 = R$, and finitely generated graded A -modules. To ensure the existence of limit dimensions and a result regarding annihilators we show

Proposition 5.1. *Let $A = \bigoplus_{n \geq 0} A_n$ be a standard graded R -algebra with $A_0 = R$. Let $M = \bigoplus_{k \geq 0} M_k$ be a finitely generated A -module. Then $\text{ann}_R M_k = \text{ann}_R M_{k-1}$ for all $k \gg 0$. In particular $\dim M_k$ is constant for all $k \gg 0$.*

Proof. Notice that $M_k = A_1 M_{k-1}$ for all $k \gg 0$. It follows that $\text{ann}_R M_{k-1} \subseteq \text{ann}_R M_k$ for all $k \gg 0$.

To prove the other inclusion we first assume that the residue field $K = R/\mathfrak{m}$ is infinite. Then by [25, 2.1,3.1] there exists $u \in A_1$ which is M -filter regular. So the R -linear maps $M_{k-1} \rightarrow M_k$ given by multiplication by u are injective for $k \gg 0$. In particular $\text{ann}_R M_k \subseteq \text{ann}_R M_{k-1}$ for $k \gg 0$. Thus the result holds when K is infinite.

If K is finite then we do the standard trick, i.e., we consider $R' = R[x]_{\mathfrak{m}R[x]}$. The residue field of R' is $K(x)$, which is infinite. Set $A' = A \otimes_R R'$ and $M'_k = M \otimes_R R'$ for $k \geq 0$. Notice that $M' = M \otimes_R R' = \bigoplus_{k \geq 0} M'_k$ is a finitely generated A' -module and A' is a finitely generated R' -algebra. Thus as proved before one has $\text{ann}_{R'} M'_k = \text{ann}_{R'} M'_{k-1}$ for all $k \gg 0$.

As each M_k is a finitely generated R -module we get

$$\text{ann}_{R'} M'_k = \text{ann}_R M_k \otimes_R R' \quad \text{for each } k \geq 1.$$

In the first line of this proof we had shown that $\text{ann}_R M_{k-1} \subseteq \text{ann}_R M_k$ for all $k \gg 0$ without any assumptions on K . Therefore

$$\frac{\text{ann}_R M_k}{\text{ann}_R M_{k-1}} \otimes_R R' = \frac{\text{ann}_{R'} M'_k}{\text{ann}_{R'} M'_{k-1}} = 0 \quad \text{for all } k \gg 0.$$

R' is a faithfully flat R -algebra. So $\text{ann}_R M_k = \text{ann}_R M_{k-1}$ for all $k \gg 0$. \square

We call the K -algebra $A/\mathfrak{m}A$ the *fiber* of A and $\dim A/\mathfrak{m}A = s(A)$, the *spread* of A . These definitions are influenced by the case when $A = \mathcal{R}(I)$. Then $A/\mathfrak{m}A = F(I)$ is called the fiber-cone of I and $s(I) = \dim F(I)$ is the analytic spread of I .

If $K = R/\mathfrak{m}$ is infinite, then by the same argument as those that are used in the proof of the existence of minimal reductions of an ideal, we can show that there exists $u_1, \dots, u_c \in A_1$ (here $c = s(A)$) such that $A_k = (u_1, \dots, u_c)A_{k-1}$ for all $k \gg 0$.

Remark 5.2. Set $B = R[y_1, \dots, y_c]$. Consider the ring homomorphism $\varphi: B \rightarrow A$ which sends y_i to u_i for each i . Then A , considered as a B -module via φ , is a finitely generated B -module.

Theorem 5.3. *With hypothesis as in Proposition 5.1 let d be the limit dimension of the family $\{M_k\}_{k \geq 0}$. Then $e_i(M_k)$ is a polynomial of degree $\leq d + s(A) - 1 - i$ for $k \gg 0$.*

Proof. We will prove the theorem in three steps.

Step I: We first consider the function $f(k, n) = \ell(M_k/\mathfrak{m}^{n+1}M_k)$ and show

$$\sum_{k, n \geq 0} f(k, n) z^n w^k = \frac{h(z, w)}{(1-z)^{p+1}(1-w)^q} \quad \text{where } h(z, w) \in \mathbb{Z}[z, w].$$

Consider the Rees ring $S = \bigoplus_{n \geq 0} (\mathfrak{m}A)^n$ and the associated graded ring $T = \text{gr}_{\mathfrak{m}A} A$ of the ideal $\mathfrak{m}A$ in A . Observe that $(\mathfrak{m}A)^n = \bigoplus_{k \geq 0} \mathfrak{m}^n A_k$. So $S = \bigoplus_{k, n \geq 0} \mathfrak{m}^n A_k$ is a naturally standard bigraded R -algebra with $S_{k, n} = \mathfrak{m}^n A_k$ and $S_{0, 0} = R$. It can also be easily checked that $T = \bigoplus_{k, n \geq 0} T_{k, n}$, where $T_{k, n} = \mathfrak{m}^n A_k / \mathfrak{m}^{n+1} A_k$ is a standard bigraded K -algebra.

Next consider the Rees module $E = \mathcal{R}(\mathfrak{m}A, M) = \bigoplus_{n \geq 0} (\mathfrak{m}A)^n M$ and the associated graded module $N = \text{gr}_{\mathfrak{m}A} M$ of M with respect to the ideal $\mathfrak{m}A$. Clearly E and N are finitely generated bigraded S and T modules, respectively. Notice that $N = \bigoplus_{k, n \geq 0} \mathfrak{m}^n M_k / \mathfrak{m}^{n+1} M_k$.

By a bigraded version of the Hilbert-Serre theorem, cf. [17, 8.20] we get that

$$\sum_{k, n \geq 0} \ell \left(\frac{\mathfrak{m}^n M_k}{\mathfrak{m}^{n+1} M_k} \right) z^n w^k = \frac{h(z, w)}{(1-z)^p (1-w)^q} \quad \text{where } h(z, w) \in \mathbb{Z}[z, w].$$

Therefore

$$\sum_{k, n \geq 0} \ell \left(\frac{M_k}{\mathfrak{m}^{n+1} M_k} \right) z^n w^k = \frac{h(z, w)}{(1-z)^{p+1} (1-w)^q}.$$

Step II: $e_i(M_k)$ is of polynomial type in k of degree $\leq p + q - 1 - i$.

By Step 1. we get that there exists k_0 and $n_0 \gg 0$, such that

$$f(n, k) = \sum_{0 \leq i+j \leq p+q-1} a_{i,j} \binom{n+i}{i} \binom{k+j}{j} \quad \text{for all } n \geq n_0 \text{ and } k \geq k_0.$$

Here $a_{i,j} \in \mathbb{Z}$. Since $\dim M_k$ is d for $k \gg 0$ we get that $a_{i,j} = 0$ for all $i > d$. Fix $k \geq k_0$. It follows that for all $n \gg 0$ we have

$$\ell \left(\frac{M_k}{\mathfrak{m}^{n+1} M_k} \right) = f(n, k) = \sum_{i=0}^d \left(\sum_{0 \leq j \leq p+q-1-i} a_{i,j} \binom{k+j}{j} \right) \binom{n+i}{i}$$

Hence

$$e_i(M_k) = \sum_{0 \leq j \leq p+q-1-i} a_{i,j} \binom{k+j}{j} \quad \text{for } i = 0, \dots, d.$$

This implies that $e_i(M_k)$ is a polynomial in k of degree $\leq p + q - 1 - i$ for $k \gg 0$.

Step III: We show that $p \leq d$ and $q \leq s(A)$.

We first do several reduction steps.

Reduction step (a): We may assume $d = \dim M_k = \dim R$ for all $k \geq 0$. In fact, by Proposition 5.1, there exists r such that $\text{ann}_R M_k = \text{ann}_R M_r = \mathfrak{q}$ (say) for $k \geq r$. We consider the submodule $D = \bigoplus_{k \geq r} M_k$ of M . Notice that D is a $A/\mathfrak{q}A$ -module and $s(A/\mathfrak{q}A) \leq s(A)$. Thus (a) follows.

Reduction step (b): We may assume K is infinite. Indeed, if this is not the case we may apply a suitable base field extension.

Reduction step (c): We may assume $A = R[y_1, \dots, y_c]$, where $c = s(A)$. This follows from Remark 5.2.

Now let $J = (x_1, \dots, x_d)$ be a minimal reduction of \mathfrak{m} . Then JA is a reduction of $\mathfrak{m}A$. It follows that $S = \bigoplus_{n \geq 0} (\mathfrak{m}A)^n$ is finitely generated as a $S' = \bigoplus_{n \geq 0} (JA)^n$ -module. Therefore $N = E \otimes_R R/\mathfrak{m}$ is a finitely generated $S' \otimes_R R/\mathfrak{m}$ -module. Since $T' = S' \cong \mathcal{R}(J)[y_1, \dots, y_c]$, we get $S' \otimes_R R/\mathfrak{m} \cong F(J)[y_1, \dots, y_c]$. (Here $\mathcal{R}(J)$ is the Rees algebra of J and $F(J)$ is the fiber-cone of J). Since J is a minimal reduction of \mathfrak{m} , it follows that $F(J)$ is a polynomial ring over K in d -variables, say t_1, \dots, t_d . Notice that

$$(8) \quad T' \cong k[t_1, \dots, t_d, y_1, \dots, y_c] \quad \text{where } \deg t_j = (0, 1) \text{ and } \deg y_i = (1, 0), \text{ and that}$$

$$(9) \quad N = \bigoplus_{k, n \geq 0} \mathfrak{m}^n M_k / \mathfrak{m}^{n+1} M_k \text{ is a finitely generated, bigraded } T' \text{-module.}$$

It follows that $p \leq d$ and $q \leq c \leq s(A)$, see [17, 8.20]. This completes Step III.

Our result follows from Steps I, II and III. \square

Corollary 5.4. *Let M be a finitely generated R -module. Let I be an ideal in R and let J be a reduction of I with respect to M . Then for $k \gg 0$,*

- (a) $e_i(I^k M / I^{k+1} M)$ is a polynomial in k of degree $\leq \alpha_I(M) + s(I) - i - 1$, where $0 \leq i \leq \alpha_I(M)$.
- (b) $e_i(I^k M)$ is a polynomial in k of degree $\leq \beta_I(M) + s(I) - i - 1$, where $0 \leq i \leq \beta_I(M)$.
- (c) $e_i(I^k M / J^k M)$ polynomial in k of degree $\leq \gamma_J^I(M) + s(J) - i - 1$, where $i = 0, \dots, \gamma_J^I(M)$.

Proof. The assertions (a) and (b) follow from Theorem 5.3, since $\bigoplus_{k \geq 0} I^k M / I^{k+1} M$ and $\bigoplus_{k \geq 0} I^k M$ are finitely generated modules over $\mathcal{R}(I)$. The assertion (c) follows from Theorem 5.3, since $\bigoplus_{k \geq 0} I^k M / J^k M$ is a finitely generated modules over $\mathcal{R}(J)$. \square

The following result regarding multiplicities of modules over not-necessarily standard graded algebras is needed in Section 3.

Proposition 5.5. *Let $B = \bigoplus_{k \geq 0} B_k$ be a finitely generated R -algebra with $B_0 = R$. Let $M = \bigoplus_{k \geq 0} M_k$ be a finitely generated B -module. Then there exists $g \geq 1$ such that*

- (a) $\dim M_{mg+i}$ is constant (say $d(i)$) for $m \gg 0$ for each $i = 0, 1, \dots, g-1$.
- (b) The function $f(k) = e_0(M_k)$ is quasi-polynomial of period g for $k \gg 0$.
- (c) If P_0, \dots, P_{g-1} are polynomials such that for each i we have $P_i(m) = e_0(M_{mg+i})$ for $m \gg 0$, then $\deg P_i \leq \dim B/\mathfrak{m}B + d(i) - 1$.

Proof. We choose g such that the Veronese subring $A = B^{(g)}$ is standard graded. Then $M^{(g;i)} = \bigoplus_{k \geq 0} M_{kg+i}$ is a finitely generated A -module for each $i = 0, \dots, g-1$. Also $M = \bigoplus_{i=0}^{g-1} M^{(g;i)}$. The assertions (a) and (b) follow from Theorem 5.3. To get the degree estimate notice that

$$\left(\frac{B}{\mathfrak{m}B} \right)^{(g)} = \frac{B^{(g)}}{\mathfrak{m}B^{(g)}} = \frac{A}{\mathfrak{m}A}.$$

So $s(A) = \dim A/\mathfrak{m}A = \dim B/\mathfrak{m}B$. □

REFERENCES

- [1] J. O. Amao, On a certain Hilbert polynomial. *J. London Math. Soc. (2)* 14 (1976), no. 1, 13–20.
- [2] M. Brodmann, Asymptotic stability of $\text{Ass}(M/I^n M)$. *Proc. Amer. Math. Soc.* **74** (1979), no. 1, 16–18.
- [3] M. Brodmann and R. Y. Sharp, *Local Cohomology: An algebraic introduction with geometric applications*, Cambridge U. Press, 1998.
- [4] W. Bruns and J. Herzog, *Cohen-Macaulay Rings*, revised edition, Cambridge Studies in Advanced Mathematics, 39. Cambridge University Press, 1998.
- [5] W. Bruns and B. Ichim, On the coefficients of Hilbert quasipolynomials, *Proc. Amer. Math. Soc.* 135 (2007) no. 5, 1305–1308.
- [6] A. Conca, J. Herzog, N. V. Trung and G. Valla, Diagonal subalgebras of bigraded algebras and embeddings of blow-ups of projective spaces, *Amer. J. Math.* 119 (1997), 859–901
- [7] D. Cutkosky, Symbolic algebras of monomial primes, *Journal für die reine und angewandte Mathematik* 416 (1991), 71–89.
- [8] S. D. Cutkosky, H. T. Ha, H. Srinivasan and E. Theodorescu, Asymptotic behaviour of the length of local cohomology *Canad. J. Math.* 57 (2005), no. 6, 1178–1192.
- [9] S. D. Cutkosky, J. Herzog and N. V. Trung, Asymptotic behaviour of the Castelnuovo-Mumford regularity, *Compositio Math.* 118 (1999), no. 3, 243–261.
- [10] S. Goto, K. Nishida and K. Watanabe, Non-Cohen-Macaulay symbolic Rees algebras for space monomial curves and counterexamples to Cowsik’s question, *Proc. Amer. Math. Soc.* 120 (1994), 383–392.
- [11] J. Herzog, T. Hibi and N. V. Trung, Symbolic powers of monomial ideals and vertex cover algebras, *Advances in Math.* 210 (2007), 304–322.
- [12] N. D. Hoang and N. V. Trung, Hilbert polynomials of non-standard bigraded algebras, *Math. Z.* 245 (2003), no. 2, 309–334.
- [13] C. Huneke, On the finite generation of symbolic blow-ups, *Math. Z.* 179 (1982), 465–472.
- [14] C. Huneke and I. Swanson, *Integral closures of ideals, rings and modules*, London Math. Society lecture Note Series 336, Cambridge University Press, 2006.
- [15] V. Kodiyalam, Asymptotic behaviour of Castelnuovo-Mumford regularity, *Proc. Amer. Math. Soc.* 128 (2000), no. 2, 407–411.
- [16] P. McMullen. *Lattice invariant valuations on rational polytopes*. *Arch. Math.* 31 (1978/79), 509–516.
- [17] E. Miller and B. Sturmfels, *Combinatorial commutative algebra*, Graduate Texts in Mathematics, 227, Springer-Verlag, New York, 2005.
- [18] T. J. Puthenpurakal, Ratliff-Rush Filtration, regularity and depth of higher associated graded modules: Part I. *J. Pure Appl. Algebra.* 208 (2007), 159–176.
- [19] D. Rees, \mathfrak{A} -transforms of local rings and a theorem on multiplicities of ideals, *Proc. Cambridge Philos. Soc. (2)* 57 (1961) 8–17.
- [20] D. Rees, A note on analytically unramified local rings, *J. London Math. Soc.* 36 (1961), 24–28.
- [21] D. Rees, Amao’s theorem and reduction criteria, *J. London Math. Soc. (2)* 32 (1985), no. 3, 404–410.
- [22] P. C. Roberts, A prime ideal in a polynomial ring whose symbolic blow-up is not Noetherian, *Proc. Amer. Math. Soc.* 94 (1985), no. 4, 589–592.
- [23] K. Shah, Coefficient ideals, *Trans. Amer. Math. Soc.* 327 (1991), no. 1, 373–384.

- [24] R. Stanley. *Decompositions of rational convex polytopes*. Ann. Discr. Math. 6, (1980), 333–342.
- [25] N. V. Trung, Reduction exponent and degree bound for the defining equations of graded rings, Proc. Amer. Math. Soc. **101** (1987) 229-236.
- [26] N. V. Trung, The Castelnuovo Regularity of the Rees algebra and the associated graded ring, Trans. Amer. Math. Soc. 350 (1998), 2813-2832

JÜRGEN HERZOG, FACHBEREICH MATHEMATIK UND INFORMATIK, UNIVERSITÄT DUISBURG-ESSEN,
CAMPUS ESSEN, 45117 ESSEN, GERMANY

E-mail address: `juergen.herzog@uni-essen.de`

TONY J. PUTHENPURAKAL, DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY,
MUMBAI, INDIA

E-mail address: `tputhen@math.iitb.ac.in`

JUGAL K. VERMA, DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY, MUMBAI,
INDIA

E-mail address: `jkv@math.iitb.ac.in`