

ON FIBERS OF ALGEBRAIC INVARIANT MOMENT MAPS

IVAN V. LOSEV

Abstract. In this paper we study some properties of fibers of the invariant moment map for a Hamiltonian action of a reductive group on an affine symplectic variety. We prove that all fibers have equal dimension. Further, under some additional restrictions, we show that the quotients of fibers are irreducible normal schemes.

Contents

1. Introduction	1
2. Preliminaries	4
2.1. Hamiltonian actions	4
2.2. Conical Hamiltonian varieties	7
2.3. Local structure of Hamiltonian actions	8
2.4. Some results concerning $\mathbb{A}^1_{G/X}; \mathbb{C}_{G/X}$	10
3. Dimensions of fibers	12
3.1. A Hamiltonian version of the Luna-Richardson theorem	13
3.2. A stratification of a fiber of $\mathbb{A}^1_{G/X} = G$	14
3.3. The proof of Theorem 3.1	15
4. Some results concerning Weyl groups	17
4.1. Some technical propositions	18
4.2. The structure of Weyl groups of affine Hamiltonian varieties	21
4.3. Examples of computation of Weyl groups	25
5. Fibers of $\mathbb{A}^1_{G/X}$ and untwisted varieties	28
5.1. Reducedness of fibers of $\mathbb{A}^1_{G/X}$	28
5.2. Proof of Theorem 1.4	31
5.3. Some classes of untwisted varieties	32
5.4. Some counterexamples	34
6. Some open problems	34
7. Notation and conventions	35
References	36

1. Introduction

Let K be a connected compact Lie group acting on a symplectic real manifold M by symplectomorphisms. Suppose there exists a moment map $\mu : M \rightarrow \mathfrak{k}^*$ (see, for instance, [GS] for the definition of moment maps). It is an important problem in symplectic geometry

Key words and phrases: reductive groups, Hamiltonian actions, moment maps, irreducibility of fibers, Weyl groups.

2000 Mathematics Subject Classification. 14R20, 53D20.

to study properties of μ . In fact, usually one studies not the map μ itself, but some coarser map, which we call the invariant moment map. It is constructed as follows. One chooses a Weyl chamber $C \subset \mathfrak{k}$. The inclusion $C \hookrightarrow \mathfrak{k}$ induces a homeomorphism $C \cong \mathfrak{k}/K$ of topological spaces. By definition, the invariant moment map μ^{inv} is the composition of $\mu: M \rightarrow \mathfrak{k}$ and the quotient map $\mathfrak{k} \rightarrow \mathfrak{k}/K$. It turns out that the map μ^{inv} has the following amazing properties:

- (a) The image of μ^{inv} is a compact convex polytope in C .
- (b) All fibers of μ^{inv} are connected.
- (c) μ^{inv} is an open map.

(a) and (b) were proved by Kirwan in [K1], (c) is due to Knop, [Kn5]. Since μ is K -equivariant, one can extract some information about the image of μ from (a). From (b) one derives that all fibers of μ are connected. Hamiltonian K -manifolds satisfying (a)–(c) were called convex in [Kn5]. In fact, all interesting classes of Hamiltonian manifolds are convex, see [Kn5] for details.

An algebraic analog of the category of smooth manifolds with an action of a compact Lie group is the category of smooth affine varieties acted on by a reductive algebraic group. Similarly to the case of compact groups one can define the notion of a Hamiltonian action of a reductive group, see Subsection 2.1. It is an interesting problem to understand:

- (1) what are algebraic analogs of properties (a)–(c)?
- (2) what varieties satisfy these properties?

Below all groups and varieties are defined over \mathbb{C} . First of all, we need to define the invariant moment map in the algebraic category. Let X be a symplectic algebraic variety and G a reductive algebraic group acting on X in a Hamiltonian way. Fix a moment map $\mu_X: X \rightarrow \mathfrak{g}$ for this action. In the sequel it will be convenient to identify \mathfrak{g} and \mathfrak{g}^* by means of a nondegenerate invariant symmetric form on \mathfrak{g} and consider μ_X as a morphism $X \rightarrow \mathfrak{g}^*/G$. By the invariant moment map for X we mean the morphism $\mu_X := \mu_X \circ \pi_X$, where π_X denotes the quotient morphism $\mathfrak{g} \rightarrow \mathfrak{g}^*/G$ for the adjoint action $G \curvearrowright \mathfrak{g}$. Note that the relation between μ_X and μ_X is more loose than in the case of compact groups. For example, one cannot determine μ_X by μ_X .

It turns out that the morphism μ_X does have some good properties.

Theorem 1.1. The morphism μ_X is equidimensional (i.e., any component of any its nonempty fiber has the same dimension equal, obviously, to $\dim X - \dim \mu_X^{-1}(0)$).

In fact, a more precise result holds, see Theorem 3.1.

However, μ_X does not seem to have other good properties. For example, even its general fiber need not be connected, see [Kn4], Introduction. Therefore one needs to modify the morphism μ_X .

To this end we introduce a kind of the Stein factorization of μ_X . Namely, let A denote the integral closure of the subalgebra $\mu_X^{-1}(\mathbb{C}[\mathfrak{g}^*])$ in $\mathbb{C}[X]$. Set $C_{\mu_X} := \text{Spec}(A)$. There are the natural G -invariant morphism $e_{\mu_X}: X \rightarrow C_{\mu_X}$ and the finite morphism $\mu_X: C_{\mu_X} \rightarrow \mathfrak{g}^*/G$ such that $\mu_X \circ e_{\mu_X} = \mu_X$. Note that at least general fibers of $e_{\mu_X}: X \rightarrow C_{\mu_X}$ are connected whenever G is connected. The idea to replace μ_X with e_{μ_X} belongs to F. Knop, see [Kn4].

In [Kn4] Knop proved that any fiber of e_{μ_X} is connected provided X is the cotangent bundle of some smooth irreducible (not necessarily affine) G -variety. On the other hand,

he constructed an example of a four-dimensional affine Hamiltonian \mathbb{C}^* -variety X such that $e_{G,X}$ has a disconnected fiber.

On the other hand, Theorems 1.6, 1.8 from [Lo2] describe the image of $e_{G,X}$. This description is particularly easy when X satisfies some additional condition that can be described as a presence of a grading on $\mathbb{C}[X]$ compatible with the structure of a Hamiltonian variety.

Definition 1.2. An affine Hamiltonian G -variety X equipped with an action $\mathbb{C}^* : X \rightarrow X$ commuting with the action of G is said to be conical if the following two conditions are fulfilled

- (Con1) The morphism $\mathbb{C}^* : X \rightarrow G \backslash X \cong G \backslash (t \cdot_{G,X}(x)) \cong G \backslash (tx)$; can be extended to a morphism $\mathbb{C}^* : X \rightarrow G \backslash X \cong G$.
- (Con2) There exists a positive integer k (called the degree of X) such that $!_{tx} = t^k !_x$ and $_{G,X}(tx) = t^k \cdot_{G,X}(x)$ for all $t \in \mathbb{C}^*; x \in X$.

For example, a symplectic G -module and the cotangent bundle of a smooth affine G -variety are conical.

If X is conical, then $C_{G,X}$ is a quotient of a vector space by a finite group and $e_{G,X}$ is surjective, see [Lo2], Theorem 1.8. More precisely, there is a subspace $\mathfrak{a} \subset \mathfrak{g}$ (called the Cartan space of X) and a subgroup $W \subset N_G(\mathfrak{a}) = Z_G(\mathfrak{a})$ (the Weyl group) such that $C_{G,X} = \mathfrak{a}/W$ and the finite morphism $_{G,X} : C_{G,X} \rightarrow \mathfrak{g}/G$ is induced by the embedding $\mathfrak{a} \hookrightarrow \mathfrak{g}$. So the subspace $\mathfrak{a} \subset \mathfrak{g}$ and the group W encode the difference between $e_{G,X}$ and $_{G,X}$.

We have no examples of conical Hamiltonian G -varieties, where $e_{G,X}$ has a disconnected fiber. We conjecture that in this case all fibers of $e_{G,X}$ are connected and, more precisely, that X enjoys the following property:

- (Irr) Any fiber of $e_{G,X} = G \backslash X \rightarrow G \backslash C_{G,X}$ is irreducible.

We are able to prove (Irr) only under another restriction on X .

Definition 1.3. An affine Hamiltonian G -variety X is said to be untwisted if

- (Utw1) $C_{G,X}$ is smooth.
- (Utw2) The morphism $e_{G,X}$ is smooth in codimension 1 (that is, the complement to the set of smooth points of $e_{G,X}$ has codimension at least 2).

Theorem 1.4. Let G be connected and X a conical Hamiltonian G -variety.

- (1) If X is untwisted, then any fiber of $e_{G,X} = G \backslash X \rightarrow G \backslash C_{G,X}$ is a normal Cohen-Macaulay scheme.
- (2) If X satisfies (Utw1) and all fibers of $e_{G,X} = G \backslash X \rightarrow G \backslash C_{G,X}$ are normal (as schemes), then X satisfies (Irr).
- (3) Suppose X is algebraically simply connected. If X satisfies (Irr), then X is untwisted.

We recall that a smooth irreducible variety X is called algebraically simply connected if a finite étale morphism $\gamma : Y \rightarrow X$ is the identity whenever Y is irreducible.

Note that a fiber of $e_{G,X} = G \backslash X \rightarrow G \backslash C_{G,X}$ can be thought as an algebraic analog of a Marsden-Weinstein reduction, [MW].

Remark 1.5. In fact, assertion 3 can be generalized to non simply connected varieties. Namely, suppose X satisfies (Irr). Then there exists an untwisted conical Hamiltonian G -variety \mathcal{X} and a free action of a finite group Γ on \mathcal{X} by Hamiltonian automorphisms (see Definition 2.12) such that $X = \mathcal{X}/\Gamma$ and $_{G,X} : \mathcal{X} \rightarrow X$ is a Hamiltonian morphism. The proof of this claim is similar to that of assertion 3 in Subsection 5.2.

This remark partially justifies the term "untwisted".

Now let us describe some classes of conical untwisted Hamiltonian G -varieties. Knop showed in [Kn3] that the cotangent bundle of any smooth irreducible affine variety is untwisted. In the present paper we sketch an alternative proof of this claim and prove that a symplectic G -module is untwisted.

Let us briefly describe the content of the paper. In Section 2 we recall some known results concerning Hamiltonian actions in the algebraic setting. Section 3 is devoted to the proof of Theorem 1.1 (in fact, of a more precise statement). In Section 4 we prove some results concerning the Weyl groups of Hamiltonian actions (see above). These results are used in the proof of Theorem 1.4. Besides, they play a crucial role in the computation of Weyl groups and root lattices of affine G -varieties, the former is done in the preprint [Lo4]. Section 5 is devoted to the proof of Theorem 1.4. We also present there some classes of untwisted varieties. In Section 6 we discuss some open problems related to the subject of the paper. Finally, Section 7 contains conventions and the list of notation we use. In the beginning of Sections 2-5 their content is described in more detail.

This paper is an expanded version of Section 5 and Subsection 7.2 of the preprint [Lo4].

Acknowledgements. Part of the work on this paper was done during my visit to Ruhr University, Bochum, in July, 2005, in the framework of Euler program. I would like to thank this institution and especially Professor H. Flenner for hospitality. I also express my gratitude to Professor F. Knop for his kind permission to use his counterexample in Subsection 5.4.

2. Preliminaries

In this section G is a reductive algebraic group and X is a variety equipped with a regular symplectic form ω and an action of G by symplectomorphisms.

In Subsection 2.1 we recall the definition of a Hamiltonian action and give some examples. Subsection 2.2 is devoted to conical Hamiltonian varieties introduced in [Lo2]. In Subsection 2.3 we study a local structure of Hamiltonian actions. At first, we recall the theory of cross-sections of Hamiltonian actions (Proposition 2.20) tracing back to Guillemin-Stenberg, [GS]. Next, in this subsection we recall the symplectic slice theorem from [Lo3]. These two results are key ingredients of most proofs in this paper. Finally, in Subsection 2.4 we recall some results from [Lo2], [Lo5]. The most important ones are Propositions 2.31, 2.32.

2.1. Hamiltonian actions. Let U be an open subset of X and f a regular function on U . The skew-gradient $v(f)$ of f is, by definition, a regular vector field on U given by the equality

$$\omega_x(v(f); \cdot) = \text{hd}_x f; \cdot; \forall x \in U; \forall \cdot \in T_x X :$$

For $f, g \in C[U]$ one defines their Poisson bracket $\{f, g\} \in C[U]$ by

$$\{f, g\} = \omega(v(f); v(g)) :$$

Clearly, $\{f, g\} = L_{v(f)}g$, where L denotes the Lie derivative.

To any element $\lambda \in \mathfrak{g}$ one associates the velocity vector field $v(\lambda)$. Suppose there is a linear map $\gamma : \mathfrak{g} \rightarrow C[X]; \lambda \mapsto \gamma(\lambda)$; satisfying the following two conditions:

(H1) The map $\gamma : \mathfrak{g} \rightarrow C[X]$ is G -equivariant.

(H2) $v(\gamma(\lambda)) = v(\lambda)$.

Definition 2.1. The action $G : X$ equipped with a linear map $\gamma : \mathfrak{g} \rightarrow C[X]$ satisfying (H1), (H2) is said to be Hamiltonian and X is called a Hamiltonian G -variety.

Remark 2.2. Very often the definition of a Hamiltonian action is given in a slightly different way. Namely, for a connected group G condition (H1) is replaced by the condition $\langle H, g \rangle = H_{[1,2]}$. However, these two conditions are equivalent provided (H2) is fulfilled. We also remark that the definition given above is a special case of the definition from [Lo2].

For a Hamiltonian action $G \curvearrowright X$ we define the morphism $\mu_{G,X} : X \rightarrow \mathfrak{g}^*$ by the formula

$$\langle \mu_{G,X}(x), \xi \rangle = \langle H(x), \xi \rangle, \quad \xi \in \mathfrak{g}, x \in X.$$

This morphism is called the moment map of the Hamiltonian G -variety X .

Conditions (H1), (H2) are equivalent, respectively, to

(M1) $\mu_{G,X}$ is G -equivariant.

(M2) $\langle d\mu_{G,X}(x)(v), \xi \rangle = \langle \xi, x \cdot v \rangle$ for all $x \in X, v \in T_x X, \xi \in \mathfrak{g}$.

Any two maps $\mu_{G,X} : X \rightarrow \mathfrak{g}^*$ satisfying conditions (M1), (M2) differ by an element of \mathfrak{g}^G . Moreover, $H_{[1,2]} = \langle H, g \rangle = \langle \mu_{G,X}, \xi \rangle$ (see, for example, [V2]). Conversely, for any $\xi \in \mathfrak{g}^G$ there exists a unique Hamiltonian G -variety X coinciding with X as a symplectic G -variety and such that $\mu_{G,X} = \mu_{G,X} + \xi$.

Let us choose some effective G -module V and put $\langle \xi, \eta \rangle = \text{tr}_V(\xi \eta)$. The form $\langle \xi, \eta \rangle$ is G -invariant, symmetric and its restriction to the Lie algebra of any reductive subgroup of G is nondegenerate. Using this form, we identify \mathfrak{g} and \mathfrak{g}^* . In particular, we may consider $\mu_{G,X}$ as a morphism from X to \mathfrak{g} .

Let us now give some examples of Hamiltonian G -varieties.

Example 2.3 (Cotangent bundles). Let X_0 be a smooth G -variety, $X = T^*X_0$ the cotangent bundle of X_0 . X is a symplectic algebraic variety (the symplectic form is presented, for example, in [V2]). The action of G on X is Hamiltonian. The moment map is given by $\langle \mu_{G,X}(y, \eta), \xi \rangle = \langle \xi, y \cdot \eta \rangle$. Here $y \in X_0, \eta \in T_y^*X_0, \xi \in \mathfrak{g}$.

Example 2.4 (Symplectic vector spaces). Let V be a vector space equipped with a non-degenerate skew-symmetric bilinear form ω . Then V is a symplectic variety. Let G act on V by linear symplectomorphisms. Then the action $G \curvearrowright V$ is Hamiltonian. The moment map $\mu_{G,V}$ is given by $\langle \mu_{G,V}(v), \xi \rangle = \frac{1}{2} \omega(\xi v, v), \quad \xi \in \mathfrak{g}, v \in V$.

Example 2.5 (Model varieties). This example generalizes the previous one. Let H be a reductive subgroup of $G, \xi \in \mathfrak{g}^H, V$ a symplectic H -module. Put $U = \{z_g(\xi) = h\}$. Let us equip the homogeneous vector bundle $X = G/H \times_H (U \times V)$ with a certain closed 2-form. Choose an \mathfrak{sl}_2 -triple (n, h, f) in $z_g(\xi)^H$ (where h is semisimple and f is nilpotent). The H -module U can be identified with $z_g(f) \setminus h^2$. Fix a point $x = [1; (u, v)] \in X$. The tangent space $T_x X$ is naturally identified with $h^2 \oplus U \oplus V$, where $U \oplus V$ is the tangent space to the fiber of the projection $G/H \times_H (U \times V) \rightarrow G/H$ and the embedding $h^2 \hookrightarrow T_x X$ is given by $\eta \mapsto x \cdot \eta$. Put

$$\langle \eta, (u_1 + v_1 + \eta_1, u_2 + v_2 + \eta_2) \rangle = \langle \eta, (v_1, v_2) \rangle + \langle \eta_1, u_2 \rangle - \langle \eta_2, u_1 \rangle + \langle \eta + u + \mu_{G,V}(v), [\eta_1, \eta_2] \rangle;$$

$$u_1, u_2 \in U; v_1, v_2 \in V; \eta_1, \eta_2 \in h^2.$$

The corresponding section $\omega \in H^2(G/H \times_H (U \times V); \mathbb{C}^2(h^2 \oplus U \oplus V))$ is H -invariant. Thus ω can be extended to a unique G -invariant 2-form on X , which is denoted also by ω . It turns out that ω is closed and nondegenerate in any point of the zero section $G=H$, [Lo3], assertion 1 of Proposition 1. If ξ is nilpotent, then ω is nondegenerate on the whole variety X . In the general case the subset $X_r = \{x \in X \mid \mu_{G,X}(x) = \xi\}$ is nondegenerate in x is a ne.

The action $G : X_r$ is Hamiltonian. The moment map is given by (see [Lo3], assertion 3 of Proposition 1)

$$G_{\mathcal{X}_r}([g; (u; v)]) = \text{Ad}(g)(\theta + u + \theta_H(v)):$$

We denote the Hamiltonian variety X_r by $M_G(H; \theta; V)$ and call it a model variety.

Remark 2.6. The Hamiltonian structure on $M_G(H; \theta; V)$ depends on the choice of an \mathfrak{sl}_2 -triple $(n; h; f)$ in $\mathfrak{z}_g(\mathfrak{s})^H$ (if $n \neq 0$). However, Hamiltonian varieties corresponding to different choices of $h; f$ are isomorphic (see Remark 1 from [Lo3]). In the sequel we say that $(n; h; f)$ is an \mathfrak{sl}_2 -triple generating $M_G(H; \theta; V)$.

Remark 2.7. For $\theta \in \mathfrak{g}^G$ the Hamiltonian G -varieties $M_G(H; \theta + \theta_0; V); M_G(H; \theta; V)_0$ are naturally identified. They even coincide as subsets in $G/H \times (U \times V)$.

Now we consider two constructions with Hamiltonian varieties.

Example 2.8 (Restriction to a subgroup). Let H be a reductive subgroup of G and X a Hamiltonian G -variety. Then X is a Hamiltonian H -variety with the moment map $\theta_H|_X = p|_X$. Here p denotes the restriction $m \mapsto h$.

Example 2.9 (Products). Suppose $X_1; X_2$ are Hamiltonian G -varieties. Being the product of symplectic varieties, the variety $X_1 \times X_2$ has the natural symplectic structure. The action $G_1 : X_1 \times X_2$ is Hamiltonian. The moment map is given by the formula $G_{\mathcal{X}_1 \times \mathcal{X}_2}(x_1; x_2) = G_{\mathcal{X}_1}(x_1) + G_{\mathcal{X}_2}(x_2)$ for $x_1 \in X_1; x_2 \in X_2$.

Remark 2.10. It follows directly from the construction of a model variety that if $(H; \theta; V)$ is the same as in Example 2.5 and V_0 is a trivial symplectic H -module, then the Hamiltonian G -varieties $M_G(H; \theta; V \oplus V_0) = M_G(H; \theta; V) \times V_0$ are isomorphic (the action $G : V_0$ is assumed to be trivial).

Now we define some important numerical invariants of an irreducible Hamiltonian G -variety X . For an action of G on an algebraic variety Y we denote by $m_G(Y)$ the maximal dimension of a G -orbit on Y . The number $m_G(X) = m_G(\overline{m_G(X)})$ is called the defect of X and is denoted by $\text{def}_G(X)$. The number $\dim X - \text{def}_G(X) = m_G(X)$ is called the corank of X and is denoted by $\text{cork}_G(X)$. Equivalently, $\text{cork}_G(X) = \text{tr:deg } C(X)^G - \text{def}_G(X)$. An irreducible Hamiltonian G -variety X such that $\text{cork}_G(X) = 0$ is called coisotropic.

It follows from the standard properties of a moment map (see, for example, [V2]) that the defect and the corank of X coincide, respectively, with $\dim \ker !j_x, \text{rk} !j_x$ for a point $x \in X$ in general position. Further, the following statement holds, see [Lo2], Proposition 3.7.

Lemma 2.11. $\dim C_{G, \mathcal{X}} = \dim \overline{m_G(X)} = \text{def}_G(X)$.

Definition 2.12. Let $X_1; X_2$ be Hamiltonian G -varieties. A morphism $\theta' : X_1 \rightarrow X_2$ is called Hamiltonian if it is an étale G -equivariant symplectomorphism intertwining the moment maps.

Note that a Hamiltonian morphism $\theta' : X_1 \rightarrow X_2$ induces a unique morphism $\theta'_0 : C_{G, \mathcal{X}_1} \rightarrow C_{G, \mathcal{X}_2}$ such that $e_{G, \mathcal{X}_2} \circ \theta' = \theta'_0 \circ e_{G, \mathcal{X}_1}$.

Remark 2.13. One can similarly define Hamiltonian actions on complex analytic manifolds. The definitions of the corank and the defect can be extended to this case without any noticeable modifications.

2.2. Conical Hamiltonian varieties. The definition of a conical Hamiltonian variety was given in Introduction, Definition 1.2.

Example 2.14 (Cotangent bundles). Let X_0/X be such as in Example 2.3. The variety X is a vector bundle over X_0 . The action $C : X$ by the fiberwise multiplication turns X into a conical variety of degree 1.

Example 2.15 (Symplectic vector spaces). The symplectic G -module V equipped with the action $C : V$ given by $(t; v) \mapsto tv$ is conical of degree 2.

Example 2.16 (Model varieties). This example generalizes the previous one. Let $H; \cdot; V$ be such as in Example 2.5 and $X = M_G(H; \cdot; V)$. Suppose that \cdot is nilpotent. Here we define an action $C : X$ turning X into a conical Hamiltonian variety of degree 2. Let $(\cdot; h; f)$ be the \mathfrak{sl}_2 -triple in \mathfrak{g}^H generating X . As a G -variety, $X = G_H(U \cup V)$, where $U = z_g(f) \setminus h^2$. Note that h is an image of a coroot under an embedding of Lie algebras. In particular, there exists a one-parameter subgroup $\cdot : C \rightarrow G$ with $\frac{d}{dt}\cdot|_{t=0} = h$. Since $[h; h] = 0; [h; f] = -2f$, we see that $\cdot(t)(h^2) = h^2; \cdot(t)(U) = U$. Define a morphism $C : X \rightarrow X$ by formula

$$(2.1) \quad (t; [g; (u; v)]) \mapsto [g; t^2 \cdot(t)^{-1}u; tv]; t \in C; g \in G; u \in U; v \in V:$$

One checks directly that morphism (2.1) is well-defined and determines the action of C on X commuting with the action of G . Let us check that X with this action is a conical Hamiltonian variety. The action of C on $X = G$ coincides with that induced by the action $C : X$ given by

$$(2.2) \quad (t; [g; (u; v)]) \mapsto [g; t^2 \cdot(t)^{-1}u; tv]:$$

The eigenvalues of $\text{ad}(h)$ on $z_g(f)$ are not positive. Thus morphism (2.2) can be extended to a morphism $C : X \rightarrow X$. This yields (Con1). (Con2) for $k = 2$ is verified directly using the construction of Example 2.5.

Remark 2.17. Let X be such as in the previous example. The action $C : X$ induces the non-negative grading on $C[X]^g$. In the notation of the previous example $C[X]^g = C[U \cup V]^g$, where $U = z_g(f) \setminus h^2$. The grading on $C[U \cup V]^g$ is induced from the following grading on $C[U \cup V]$:

elements of $V \subset C[U \cup V]$ have degree 1. The H -module U is naturally identified with $z_g(\cdot) \setminus h^2$. Put $g_i = f - 2g[jh; \cdot] = i \cdot g$. An element from $z_g(\cdot) \setminus h^2 \setminus g_i$ has degree $i + 2$.

Lemma 2.18 ([Lo2], Lemma 3.27). Let X be a conical Hamiltonian G -variety of degree k . Then

- (1) $0 \in \text{in}_{G, X}$.
- (2) Assume X is irreducible and normal. Then the subalgebra $K[C_{G, X}] \subset K[X]^g$ is C -stable. The morphisms $e_{G, X} : X \rightarrow C_{G, X}; g_{G, X} : C_{G, X} \rightarrow g = G$ are C -equivariant, where the action $C : g = G$ is induced from the action $C : g$ given by $(t; x) \mapsto t^k x; t \in C; x \in g$.
- (3) Under the assumptions of assertion 2, there is a unique point $o \in C_{G, X}$ such that $g_{G, X}(o) = 0$. For any point $\cdot \in C_{G, X}$ the limit $\lim_{t \rightarrow 0} t \cdot$ exists and is equal to o .

Remark 2.19. It follows easily from Lemma 2.18 that $0 \in a_{G, X}^{(\alpha_L)}$ provided X is conical.

2.3. Local structure of Hamiltonian actions. Firstly, we review the algebraic variant of the Guillemin–Stenberg local cross-section theory, see [Kn3], Section 5, [Lo2] Subsection 5.1. Let L be a Levi subgroup of G and \mathfrak{l} the corresponding Lie algebra. Put $\mathbb{P}^r = \mathbb{A}^2 \setminus \mathbb{A}^1 \setminus \{0\}$.

Proposition 2.20 ([Kn3], Theorem 5.4 and [Lo2], Corollary 5.3, Propositions 5.2, 5.4, 5.7). Let $x \in X$; $\mathfrak{l} = \mathfrak{z}_G(\mathfrak{g}_x)$; $Y = \mathbb{P}^r$. Then

- (1) $T_y X = \mathfrak{l}^* y \oplus T_y Y$ is a skew-orthogonal direct sum for any $y \in Y$. In particular, Y is a smooth subvariety of X and the restriction of ω to Y is nondegenerate. Thus Y is equipped with the symplectic structure.
- (2) The action $N_G(L) : Y \rightarrow Y$ is Hamiltonian with the moment map $\mathbb{A}^1 \setminus \{0\}$.
- (3) The natural morphism $G/N_G(L) \times Y \rightarrow X$ is étale. Its image is saturated.
- (4) If x is in general position, then the natural morphism $G/N_G(L) \times Y \rightarrow X$ is an open embedding and $N_G(L)$ permutes the connected components of Y transitively.

A subset Z^0 of a G -variety Z is said to be saturated if there exist a G -invariant morphism $\nu : Z \rightarrow Z^0$ and a subset $Z_0^0 \subset Z^0$ such that $Z^0 = \nu^{-1}(Z_0^0)$.

Definition 2.21. An irreducible (= connected) component of \mathbb{P}^r equipped with the structure of a Hamiltonian L -variety obtained by restriction of the Hamiltonian structure from \mathbb{P}^r is called an L -cross-section of X .

Definition 2.22. The Levi subgroup $L = \mathfrak{z}_G(\mathfrak{g}_x)$, where $x \in X$ is in general position, is said to be the principal centralizer of X .

Note that the principal centralizer is determined uniquely up to G -conjugacy.

Lemma 2.23. Let L be the principal centralizer and X_L an L -cross-section of X . Then the following conditions are equivalent:

- (1) $m_G(X) = \dim G$.
- (2) $\text{def}_G(X) = \text{rk } G$.
- (3) $\overline{\text{im}}_{G/X} = \mathfrak{g}$.
- (4) L is a maximal torus in G and $m_L(X_L) = \text{def}_L(X_L) = \text{rk } G$.

Under these conditions, $\text{cork}_G(X) = \dim X - \dim G = \text{rk } G$.

Proof. The equivalence of conditions (1)–(4) was proved in [Lo5], Lemma 4.5. The equality for $\text{cork}_G(X)$ follows from (1) and (2).

Lemma 2.24. Let L be the principal centralizer and X_L an L -cross-section of X . Suppose that the stabilizer in general position L_0 for the action $L : X_L$ is reductive and that $0 \in \overline{\text{im}}_{G/X}$. Then $\overline{\text{im}}_{G/X} = G \setminus \{1\}$.

Proof. By Theorem 4.1 from [Lo2], $(L; L) \subset L_0$. Now the assertion stems easily from Proposition 2.20.

Now we turn to the problem of describing the structure of an affine Hamiltonian G -variety in some neighborhood of a point with closed G -orbit. A neighborhood is taken with respect to a usual complex topology (in the sequel we call such neighborhoods analytical).

At first, we define some invariants of the triple $(G; X; x)$. Put $H = G_x$; $\mathfrak{h} = \mathfrak{g}_x$. The subgroup $H \subset G$ is reductive and $\mathfrak{h} \subset \mathfrak{g}^H$. Put $V = (\mathfrak{g}_x)^\perp = (\mathfrak{g}_x \setminus \mathfrak{g}_x)^\perp$. This is a symplectic H -module. We say that $(H; V)$ is the determining triple of X at x . For

example, the determining triple of $X = M_G(H; \cdot; V)$ in $x = [1; (0; 0)]$ is $(H; \cdot; V)$, see [Lo3], assertion 4 of Proposition 1.

As the name suggests, a determining triple should determine the structure of the Hamiltonian G -variety X near x . In fact, a slightly stronger claim takes place.

Definition 2.25. Let X_1, X_2 be affine Hamiltonian G -varieties, $x_1 \in X_1, x_2 \in X_2$ be points with closed G -orbits. The pairs $(X_1; x_1); (X_2; x_2)$ are called analytically equivalent, if there are saturated open analytical neighborhoods O_1, O_2 of $x_1 \in X_1, x_2 \in X_2$, respectively, that are isomorphic as complex-analytical Hamiltonian G -manifolds.

Remark 2.26. An open saturated analytical neighborhood in X is the inverse image of an open analytical neighborhood in $X = G$ under π_X . See, for example, [Lo3], Lemma 5.

Proposition 2.27 (Symplectic slice theorem, [Lo3]). Let X be an affine Hamiltonian G -variety, $x \in X$ a point with closed G -orbit, $(H; \cdot; V)$ the determining triple of X at x . Then the pair $(X; x)$ is analytically equivalent to the pair $(M_G(H; \cdot; V); [1; (0; 0)])$.

Now we prove two lemmas, which will be used in Subsection 4.1.

We have two approaches to the local study of affine Hamiltonian varieties: the cross-sections theory and the symplectic slice theorem. Let us establish a connection between them.

Lemma 2.28. Let $x \in X$ be a point with closed G -orbit and $(H; \cdot; V)$ the determining triple of X at x . Put $M = Z_G(\cdot_s)$. Denote by X_M a unique M -cross-section of X containing x . Then the following assertions hold

- (1) Mx is closed in X_M and $(H; \cdot; V)$ is the determining triple of X_M at x .
- (2) There exists an affine saturated open (with respect to Zariski topology) neighborhood $X_M^0 \subset X_M$ of x such that the following conditions are satisfied:
 - (a) the natural morphism $X_M^0 \xrightarrow{\pi} X = G; \pi_{X_M}(z) \mapsto \pi_X(z)$ is étale;
 - (b) for any $z \in X_M^0$ the orbit Mz is closed in X_M^0 (equivalently, in X_M) if Gz is closed in X .

Proof. The morphism $\pi' : G \times_M X_M \rightarrow X; [g; x] \mapsto gx$ is étale (assertion 3 of Proposition 2.20). Since Gx is closed in X , we see that $G[1; x]$ is closed in $G \times_M X_M$, equivalently, Mx is closed in X_M . Since $G_x = Z_G(\pi_{X_M}^{-1}(z)) = Z_G(\pi_X^{-1}(z)_s) = M$, we have $G_z = M_z$ for $z \in X_M$. By construction of π_{X_M} , $\pi_{X_M}(z) = \pi_X(z)$. Assertion 1 will follow if we check that the H -modules $g \cdot x^\vee = (g \cdot x^\vee \setminus g \cdot x)$ and $m \cdot x^\vee = (m \cdot x^\vee \setminus m \cdot x)$ are isomorphic. Here the skew-orthogonal complement to $g \cdot x$ (resp., to $m \cdot x$) is taken in $T_x X$ (resp., in $T_x X_M$). The existence of an isomorphism stems from $g \cdot x = m^2 \cdot x$ and assertion 1 of Proposition 2.20.

By above, the orbits $G[1; x]; Gx$ are closed and π' induces their isomorphism. It follows from Luna's fundamental lemma, [Lu], that for some open affine neighborhood U of the point $\pi_{X_M}^{-1}(x)$ in $X_M \xrightarrow{\pi} M = (G \times_M X_M) \xrightarrow{\pi} G$ the morphism $\pi' = G : U \rightarrow X = G$ is étale and

$$(2.3) \quad \pi_{G \times_M X_M}^{-1}(U) = U \times_{X=G} X :$$

Clearly, $\pi_{G \times_M X_M}^{-1}(U) = G \times_M \pi_{X_M}^{-1}(U)$. Thanks to (2.3), we see that for all $z \in X_M^0 \xrightarrow{\pi} M \xrightarrow{\pi} G$ the orbit $G[1; z]$ is closed in $G \times_M \pi_{X_M}^{-1}(U)$ if Gz is closed in X .

The next lemma studies the behavior of determining triples under replacing G with some connected subgroup $G^0 \subset G$ containing $(G; G)$.

Lemma 2.29. Let $x \in X$ be a point with closed G -orbit and $(H; \cdot; V)$ the determining triple of X at x . Then G^0x is closed in X and the determining triple of the Hamiltonian G^0 -variety X at x has the form $(H \setminus G^0; \cdot_0; V - V_0)$, where V_0 is a trivial $H \setminus G^0$ -module and \cdot_0 is the projection of \cdot to g^0 .

Proof. Since G^0 is a normal subgroup of G , we see that all G^0 -orbits in Gx have the same dimension whence closed. Obviously, $G_x^0 = G^0 \setminus H; \cdot_{G^0x}(x) = \cdot_0$. Clearly, $g^0x \subset gx$ and $gx \setminus g^0x \subset g^0x \setminus g^0x$. Therefore we have the natural embedding $gx \setminus (gx \setminus g^0x) \hookrightarrow g^0x \setminus (g^0x \setminus g^0x)$ and the natural projection $gx \setminus (gx \setminus g^0x) \rightarrow g^0x \setminus (g^0x \setminus g^0x)$. The cokernel of the former is a quotient of the $H \setminus G^0$ -module $g^0x \setminus gx \setminus (gx \setminus g^0x) = (gx \setminus g^0x)$, while the kernel of the latter is a submodule in $gx \setminus g^0x$. Since $gx \setminus g^0x$ is a trivial $H \setminus G^0$ -module, we are done.

2.4. Some results concerning $e_{G,x}; C_{G,x}$. Let us, at first, define two important invariants of a Hamiltonian variety: its Cartan space and Weyl group. The proofs of the facts below concerning these invariants can be found in [Lo2], Subsection 5.2.

Let L be the principal centralizer and X_L an L -cross-section of a Hamiltonian G -variety X . It turns out that $\overline{z(L); x_L}$ is an affine subspace in $z(L)$. We denote this affine subspace by $a_{G,x}^{(X_L)}$ and call it the Cartan space of X . It intersects the Lie algebra of the inefficiency kernel for the action $z(L) : X_L$ in a unique point. Taking this point for zero in $a_{G,x}^{(X_L)}$ we may (and will) consider $a_{G,x}^{(X_L)}$ as a vector space. In fact, if $0 \in \overline{z(L); x_L}$ (this is the case when X is conical), then $a_{G,x}^{(X_L)}$ is a vector subspace in $z(L)$.

The group $N_G(L; X_L)$ acts linearly on $a_{G,x}^{(X_L)}$. We denote the image of $N_G(L; X_L)$ in $GL(a_{G,x}^{(X_L)})$ by $W_{G,x}^{(X_L)}$ and call it the Weyl group of X . If G is connected, then $W_{G,x}^{(X_L)}$ is naturally identified with $N_G(L; X_L) = L$.

Note that, in a suitable sense, the pair $(a_{G,x}^{(X_L)}; W_{G,x}^{(X_L)})$ does not depend up to G -conjugacy from the choice of $L; X_L$. When a particular choice of $L; X_L$ does not matter, we write $a_{G,x}^{()}$ for $a_{G,x}^{(X_L)}$ and $W_{G,x}^{()}$ for $W_{G,x}^{(X_L)}$.

Note that in $\overline{z(L); x_L} \subset a_{G,x}^{(X_L)} \subset z(L)$. There is a unique G -invariant morphism $b_{G,x} : X \rightarrow a_{G,x}^{()} = W_{G,x}^{()}$ coinciding with $\overline{z(L); x_L}$ on X_L . The morphism $\pi_{G,x} : X \rightarrow g=G$ is the composition of $b_{G,x}$ and the finite morphism $a_{G,x}^{()} = W_{G,x}^{()} \rightarrow g=G$ induced by the embedding $a_{G,x}^{(X_L)} \hookrightarrow g$. So $b_{G,x}$ factors through $e_{G,x}$ and the respective morphism $C_{G,x} \rightarrow a_{G,x}^{()} = W_{G,x}^{()}$ is finite and dominant.

The following lemma describes the behavior of $b_{G,x}$ under some simple modifications of the pair $(G; X)$.

Lemma 2.30. Let G be a reductive group, X an irreducible affine Hamiltonian G -variety, L the principal centralizer and X_L an L -cross-section of X .

- (1) Recall ([Lo5], Lemma 6.9) that $a_{G,x}^{(X_L)} = a_{G,x}^{(X_L)}$ and $W_{G,x}^{(X_L)}$ is a normal subgroup of $W_{G,x}^{(X_L)}$. The morphism $b_{G,x}$ is the composition of $b_{G,x}$ and the natural morphism of quotients $a_{G,x}^{(X_L)} = W_{G,x}^{(X_L)} \rightarrow a_{G,x}^{(X_L)} = W_{G,x}^{(X_L)}$ induced by the inclusion $W_{G,x}^{(X_L)} \subset W_{G,x}^{(X_L)}$.
- (2) Suppose G is connected. Let M be a Levi subgroup of G containing L . Recall, [Lo5], Lemmas 4.6, 6.10, that there exists a unique M -cross-section X_M of X containing

X_L and that $a_M^{(X_L)} = a_{G \backslash X}^{(X_L)}; W_M^{(X_L)} = W_{G \backslash X}^{(X_L)} \setminus M = L$. Then the following diagram is commutative.

$$\begin{array}{ccccc} X_M & \xrightarrow{\hat{M \backslash X_M} -} & a_M^{()} & \xrightarrow{W_M^{(X_L)} -} & m=M \\ \downarrow ? & & \downarrow ? & & \downarrow ? \\ X & \xrightarrow{\hat{G \backslash X} -} & a_{G \backslash X}^{()} & \xrightarrow{W_{G \backslash X}^{(X_L)} -} & g=G \end{array}$$

Here the morphism $X_M \rightarrow X$ is the embedding, the morphism $a_M^{()} \rightarrow a_{G \backslash X}^{()}; W_M^{(X_L)} \rightarrow W_{G \backslash X}^{(X_L)}$ is given by $W_M^{(X_L)} \rightarrow W_{G \backslash X}^{(X_L)}$, and the morphism $m=M \rightarrow g=G$ is induced by the restriction of functions from g to m .

- (3) Suppose G is connected and $0 \leq \dim G^0$. Let G^0 be a connected subgroup of G containing $(G; G)$. Recall, [Lo5], Lemma 4.6, that $L^0 = L \setminus G^0$ is the principal centralizer and X_L is an L^0 -cross-section of the Hamiltonian G^0 -variety X . Further, by [Lo5], Lemma 6.13, $a_{G \backslash X}^{(X_L)} \setminus g^0 = a_{G^0 \backslash X}^{(X_L)}$, the groups $W_{G \backslash X}^{(X_L)}; W_{G^0 \backslash X}^{(X_L)}$ are naturally identified, and the orthogonal projection $g \rightarrow g^0$ induces the $W_{G \backslash X}^{(X_L)}$ -equivariant epimorphism $a_{G \backslash X}^{(X_L)} \rightarrow a_{G^0 \backslash X}^{(X_L)}$. Then the following diagram is commutative.

$$\begin{array}{ccc} X & & \\ \downarrow \text{?} & & \\ a_{G \backslash X}^{(X_L)} & \xrightarrow{W_{G \backslash X}^{(X_L)} -} & a_{G^0 \backslash X}^{(X_L)} \end{array}$$

- (4) Let X^0 be an affine irreducible Hamiltonian G -variety and $\iota: X \rightarrow X^0$ a generically finite dominant G -equivariant morphism such that $\iota_{G \backslash X^0} = \iota_{G \backslash X}$. By Lemma 4.6 from [Lo5], L is the principal centralizer of X^0 and there exists a unique L -cross-section X_L^0 of X^0 such that $\iota(X_L) = X_L^0$. Further, by Lemma 6.11 from [Lo5], $a_{G \backslash X}^{(X_L)} = a_{G \backslash X^0}^{(X_L^0)}; W_{G \backslash X}^{(X_L)} = W_{G \backslash X^0}^{(X_L^0)}$. Then the following diagram is commutative.

$$\begin{array}{ccc} X & \xrightarrow{\iota} & X^0 \\ \downarrow \text{?} & & \downarrow \text{?} \\ a_{G \backslash X}^{(X_L)} & \xrightarrow{W_{G \backslash X}^{(X_L)} -} & a_{G \backslash X^0}^{(X_L^0)} \end{array}$$

- (5) Suppose G is connected and X satisfies the equivalent conditions of Lemma 2.23. Let $G = Z(G) \times G_1 \times \dots \times G_k$ be the decomposition of G into the locally direct product of the unit component of the center and simple normal subgroups. Put $T = L; T_i = L \setminus G_i$. Recall, [Lo5], Lemma 4.6, that T_i is the principal centralizer of the Hamiltonian G_i -variety X and there is a unique T_i -cross-section X_{T_i} of X containing $(\prod_{j \neq i} G_j)X_T$. Further, Lemma 6.14 implies that $a_{G_i \backslash X}^{(X_{T_i})} = t_i; W_{G_i \backslash X}^{(X_{T_i})} = \prod_{i=1}^k W_{G_i \backslash X}^{(X_{T_i})}$ and the projection of $W_{G \backslash X}^{(X_T)}$ to $GL(t_i)$ coincides with $W_{G_i \backslash X}^{(X_{T_i})}$. Then the following diagram, where the map $a_{G \backslash X}^{(X_T)} \rightarrow a_{G_i \backslash X}^{(X_{T_i})}; W_{G \backslash X}^{(X_T)} \rightarrow W_{G_i \backslash X}^{(X_{T_i})}$ is induced by the natural epimorphism $g \rightarrow g_i$, is commutative.

$$\begin{array}{c}
 X \\
 \downarrow \mathcal{C} \\
 \downarrow \mathcal{C} \\
 \downarrow \mathcal{C} \\
 a_{G/X}^{(\cdot)} = W_{G/X}^{(\cdot)} \xrightarrow{\quad} a_{G_1/X}^{(\cdot)} = W_{G_1/X}^{(\cdot)}
 \end{array}$$

Proof. The proofs of assertions 1,3,4 follow directly from the definition of $b_{G/X}$.

Let us prove assertion 2. The commutativity of the right square of the diagram follows directly from the definition of $b_{G/X}$. To prove the commutativity of the left square we note that the both morphisms $X_M \rightarrow a_{G/X}^{(\cdot)} = W_{G/X}^{(\cdot)}$ from the diagram are M -invariant and their restrictions to X_L coincide with $b_{N_G(L/X_L)/X_L}$. To complete the proof it remains to recall that $M \backslash X_L$ is dense in X_M .

Proceed to assertion 5. The morphism $b_{G_1/X} \circ j_1$ is $Z(G) \times_{j \in i} G_j$ -invariant. It follows that $b_{G_1/X}$ is G -invariant. It remains to note that the restrictions of both morphisms $X \rightarrow a_{G_1/X}^{(\cdot)} = W_{G_1/X}^{(\cdot)}$ coincide on X_T .

Now we are going to quote some properties of $C_{G/X}$; $e_{G/X}$; $b_{G/X}$ proved in [Lo2].

Proposition 2.31. The morphism $b_{G/X} = G : X \rightarrow G \rightarrow a_{G/X}^{(\cdot)} = W_{G/X}^{(\cdot)}$ is equidimensional and open. Further, for any closed subvariety Y in $b_{G/X}$ and any irreducible component Z of $b_{G/X} = G^{-1}(Y)$ the subset $b_{G/X} = G(Z)$ is dense in Y .

Proof. Note that $a_{G/X}^{(\cdot)} = W_{G/X}^{(\cdot)}$ is a normal variety of dimension $\text{def}_G(X)$. Thanks to Theorem 1.3 from [Lo2], $b_{G/X} = G$ is equidimensional. The openness stems from [Ch]. The last assertion of the proposition is an easy corollary of the fact that $b_{G/X} = G$ is equidimensional.

Proposition 2.32 ([Lo2], Theorem 1.8). Suppose X is conical. Then $C_{G/X} = a_{G/X}^{(\cdot)} = W_{G/X}^{(\cdot)}$ and $e_{G/X} = b_{G/X}$. If X is, in addition, simply connected, then $C_{G/X}$ is an affine space, or, equivalently, $W_{G/X}^{(\cdot)}$ is generated by reflections.

3. Dimensions of fibers

Throughout the section G is a connected reductive group and X is a Hamiltonian G -variety with the symplectic form ω .

In Subsection 3.1 we prove a variant of the Luna-Richardson restriction theorem ([LR]) for Hamiltonian varieties. This allows us to reduce a general affine Hamiltonian G -variety to one satisfying the equivalent conditions of Lemma 2.23.

Subsection 3.2 deals with a stratification of a fiber of the morphism $b_{G/X} = G : X \rightarrow G \rightarrow a_{G/X}^{(\cdot)} = W_{G/X}^{(\cdot)}$. A stratum consists of the images of all points with closed G -orbit and the same determining triple. The main results of the subsection are the proof that any stratum is smooth and the formula for the dimensions of the strata (Proposition 3.5).

The main part of this section is Subsection 3.3. There we prove the following result that strengthens Theorem 1.1.

Theorem 3.1. The morphisms $C_{G/X}$; $e_{G/X}$; $b_{G/X}$ are equidimensional. The morphisms $b_{G/X}$; $e_{G/X}$ are open. For any closed irreducible subvariety Y in $b_{G/X}$ and any irreducible

component $\mathcal{P} = b_{G/X}^{-1}(Y)$ the subvariety $_{G/X}(\mathcal{P}) = X=G$ is an irreducible component of $b_{G/X}^{-1}(Y)$.

The proof uses the stratification introduced in Subsection 3.2 and the estimate on dimensions of fibers of $_{G/X}$ obtained in Proposition 3.7.

3.1. A Hamiltonian version of the Luna-Richardson theorem. Let H be a reductive subgroup of G . The subvariety $X^H = X$ is smooth (see [PV], Subsection 6.5) and $N_G(H)$ -stable. Let us equip X^H with a structure of a Hamiltonian $N_G(H)$ -variety.

Proposition 3.2. (1) \dot{j}_H is nondegenerate, thus X^H is equipped with the symplectic structure.

(2) The action $N_G(H) : X^H$ is Hamiltonian with the moment map $_{N_G(H)/X^H} = _{G/X} \dot{j}_H$.

Proof. For a symplectic vector space V and a reductive subgroup $H \subset \text{Sp}(V)$ the H -modules V^H and $V = (V^H)^\perp$ are isomorphic. Thus \dot{j}_H is nondegenerate. Since $T_x(X^H) = (T_x X)^H$, see [PV], Subsection 6.5, we see that \dot{j}_H is nondegenerate.

Note that the Lie algebra of $N_G(H)$ coincides with $\mathfrak{g}^H + \mathfrak{h}$. Since $_{G/X}$ is G -equivariant, we have $_{G/X}(X^H) = \mathfrak{g}^H$. Clearly, $_{G/X} \dot{j}_H$ is $N_G(H)$ -equivariant. It remains to check that

$$(3.1) \quad v(H \dot{j}_H)_x = -x$$

for all $x \in \mathfrak{g}^H + \mathfrak{h}; x \in X^H$. Obviously, $v(H \dot{j}_H)_x = -x = 0$ for all $x \in \mathfrak{h}; x \in X^H$. Thus (3.1) holds for $x \in \mathfrak{h}$. Now let $x \in \mathfrak{g}^H$. Then $H \subset C[X]^H$, and $v(H \dot{j}_H)_x$ is an H -invariant vector for $x \in X^H$. It follows from the construction of the symplectic form on X^H that $v(H \dot{j}_H)_x = v(H \dot{j}_H)_x$.

Now we will apply the previous construction to a special choice of H .

Let L be the principal centralizer of X and X_L an L -cross-section. By Corollary 4.9 from [Lo2], the restriction of $_{(L;L)/X_L} : X_L \rightarrow X_L = (L;L)$ to $X_L^{(L;L)} = X_L$ is an isomorphism. Denote by L_0 the unit component of the identity kernel of the action $L : X_L = (L;L) = X_L^{(L;L)}$. It follows from Theorem 4.7, [Lo2], that $L_0 = (L;L)T_0$, where T_0 is the unit component of the identity kernel for the action $Z(L) : X_L$. Let X_0 be a unique connected component of X^{L_0} containing $X_L^{(L;L)}$. Put $\mathcal{G}_0 = N_G(L_0; X_0); G_0 = \mathcal{G}_0 = L_0$. We identify \mathfrak{g}_0 with $\mathfrak{g}^{L_0} \setminus \mathfrak{l}_0^2$. It follows from Lemma 3.2 that the action $\mathcal{G}_0 : X_0$ is Hamiltonian with the moment map $_{G_0/X_0} \dot{j}_0$. By Remark 3.3 from [Lo2], the action $G_0 : X_0$ is Hamiltonian with the moment map $_{G_0/X_0} = p_{\tilde{G}_0/X_0}$, where p denotes the natural projection $\mathfrak{g}_0 \rightarrow \mathfrak{g}_0$.

The following proposition is what we mean by a "Hamiltonian version of the Luna-Richardson theorem".

Proposition 3.3. In the notation introduced above the following statements hold.

- (1) The morphism $X_0 = G_0 \rightarrow X = G$ induced by the restriction of functions is an isomorphism.
- (2) $m_{G_0}(X_0) = \dim G_0$, $\text{def}_{G_0}(X_0) = \text{def}_G(X)$; $\text{cork}_G(X) = \text{cork}_{G_0}(X_0)$.
- (3) $L = L_0$ is the principal centralizer of X_0 . The subvariety X^{L_0} is dense in a unique $L = L_0$ -cross-section X_{0L} of X_0 , $a_{G_0/X_0}^{(X_{0L})} = a_{G/X}^{(X_{0L})} = 0$, where $0 \in \mathfrak{l}_0 \setminus a_{G/X}^{(X_{0L})}$; and $W_{G_0/X_0}^{(X_{0L})} = W_{G/X}^{(X_{0L})}$.
- (4) $b_{G/X} \dot{j}_0 = b_{G_0/X_0}$.

In the proof we will use the notion of the principal isotropy group for an action of a reductive group on an affine variety. Namely, let Y be an irreducible affine variety acted on by a reductive group H . It is known, see [PV], Theorem 7.12, that there exists an open subset $Y_0 \subset Y=H$ such that for any $y \in Y_0$ the closed orbit in $H^1_{H,Y}(y)$ is isomorphic to $H=C$, where C is a reductive subgroup of H .

Definition 3.4. Such a subgroup C (determined uniquely up to H -conjugacy) is called the principal isotropy subgroup for the action $H : Y$.

Proof of Proposition 3.3. The action $Z(L) : X_L^{(L;L)} = X_L = (L;L)$ is stable ([Lo2], Proposition 4.18). Thus L_0 is the unit component of the principal isotropy subgroup for the action $L : X_L$. Since the natural morphism $G \rightarrow L \rightarrow X_L \rightarrow X$ is étale and its image is saturated, we see that the group L_0 is the unit component of the principal isotropy subgroup for the action $G : X$ and that the morphism $X_0 = G_0 \rightarrow X = G$ is dominant. By the Luna-Richardson theorem ([LR]), the morphism $X_0 = G_0 \rightarrow X = G$ is an isomorphism and the action of G_0 on X_0 is locally effective. The latter yields $\text{def}_{G_0}(X_0) = \text{rk } G_0 = \text{rk } G - \text{rk } L_0 = \text{def}_G(X)$. By Theorem 1.10 from [Lo2], $C(X)^G = \text{Quot}(C[X]^G); C(X_0)^{G_0} = \text{Quot}(C[X_0]^{G_0})$. So

$$\text{cork}_G(X) = \text{tr:deg } C(X)^G \quad \text{def}_G(X) = \text{tr:deg } C(X_0)^{G_0} \quad \text{def}_{G_0}(X_0) = \text{cork}_{G_0}(X_0):$$

Proceed to assertion 3. Since $m_{G_0}(X_0) = \dim G_0$, $L=L_0$ is the principal centralizer of X_0 (see Lemma 2.23) and $a_{G_0/X_0}^{(X_{0L})} = 1 \setminus 1_0^2 = a_{G/X}^{(X_L)}|_0$ for any $L=L_0$ -cross-section X_{0L} of X_0 . The natural morphism $X_L=L \rightarrow X=G$ is dominant and quasi-finite, therefore so is the natural morphism $(X_L^{L_0})=(L=L_0) \rightarrow X_0=G_0$. It follows from [Lo2], Theorem 1.10, that $\dim X_L^{L_0} = \dim X_0 = G_0 + \dim L=L_0 = \dim X_0 - \dim G_0 + \dim L=L_0$. Since $_{G/X}(X_L^{L_0}) \rightarrow \mathbb{P}^r$, we see that $X_L^{L_0}$ lies in a unique $L=L_0$ -cross-section X_{0L} of X_0 . Comparing the dimensions, we see that $X_L^{L_0}$ is dense in X_{0L} . The equality for the Weyl groups stems from $N_G(L;X_L)=L_0$ $G_0; N_G(L;X_L^{L_0}) = N_G(L;X_L)$.

Finally, both morphisms in assertion 4 are G_0 -invariant and their restrictions to $X_L^{L_0}$ are equal to the restriction of $b_{N_G(L;X_L)/X_L}$.

3.2. A stratification of a fiber of $_{G/X}=G$. In this subsection we introduce a stratification of a fiber of the morphism $_{G/X}=G : X=G \rightarrow g=G$. We consider fibers of $_{G/X}=G$ as algebraic varieties. Namely, let $g \in g$, H be a reductive subgroup of G and V a symplectic H -module. We put

$$S_{G/X}(H; g; V) = \{x \in X \mid x \text{ is closed; } (H; g; V) \text{ is the determining triple of } X \text{ at } xg\}$$

Clearly, $S_{G/X}(H_1; g_1; V_1) = S_{G/X}(H_2; g_2; V_2)$ if there is $g \in G$ and a linear isomorphism $V_1 \xrightarrow{\sim} V_2$ such that $\text{Ad}(g)g_1 = g_2$, $gH_1g^{-1} = H_2$ and $(ghg^{-1})(v) = g(hv)$ for all $h \in H_1$.

The main result of this subsection is the following

Proposition 3.5. Let $X; G; H; g; V$ be as above, $g = g_{X/g}()$. Then $S_{G/X}(H; g; V)$ is a locally-closed smooth subvariety of pure codimension $\text{cork}_G(X) - \dim V^H$ in $_{G/X}=G^{-1}()$.

Proof. Firstly, we show that $S_{G/X}(H; g; V)$ is a locally-closed subvariety of X . Denote by Y the set of all points $x \in X$ with closed G -orbit, $G_x = H$ and $T_x X = g \cdot x = V$ ($g = h$). It follows from the Luna slice theorem applied to any point of Y that Y is a locally-closed subvariety in X . Therefore $Y = Y \setminus \bigcup_{g \in G} (\text{Ad}(G))$ is a locally closed subvariety of X . Since all orbits in Y are closed in X , we see that Y is an open saturated subvariety of \overline{Y} . Thus $S_{G/X}(H; g; V) = g_{X/g}(Y)$ is open in $\overline{Y} = G$.

Applying Proposition 2.27, we reduce the codimension and smoothness claims to the case $X = M_G(H; \cdot; V)$. Put $s = z_g(s)$. Choose an \mathfrak{sl}_2 -triple $(n; h; f)$ in \mathfrak{s}^H generating $M_G(H; \cdot; V)$. Denote by U the H -module $z_s(f) \setminus \mathfrak{h}^\perp$.

Lemma 3.6. In the above notation \bar{x} is an isolated point of $(\bar{x} + z_m(f)) \setminus \overline{\text{Ad}(G)}$.

Proof of Lemma 3.6. Note that $T(\bar{x} + z_m(f)) = z_m(f); T\overline{\text{Ad}(G)} = [\mathfrak{g}; \cdot]$. It is enough to show $z_m(f) \setminus [\mathfrak{g}; \cdot] = f0\mathfrak{g}$. The equality $m = z_g(s)$ yields $[\mathfrak{g}; \cdot] = [\mathfrak{m}^\perp; \cdot] + [\mathfrak{m}; \cdot] = \mathfrak{m}^\perp + [\mathfrak{m}; \cdot]$. Thanks to the representation theory of \mathfrak{sl}_2 , $[\mathfrak{m}; \cdot] \setminus z_m(f) = 0$ whence the required inequality.

In virtue of Remark 2.10, it is enough to assume that $V^H = f0\mathfrak{g}$. Put $x = [1; (0; 0)]$. Everything will follow if we check that ${}_{G, X}(x)$ is an isolated point in $S_{G, X}(H; \cdot; V)$. Indeed, by Lemma 2.31, $\text{cork}_G(X) = \dim X = \dim \text{def}_G(X) = \dim {}_{G, X}(x) - \dim \overline{{}_{G, X}(x)} = 1$.

There exists a neighborhood O^0 of \bar{x} in $\bar{x} + z_s(f)$ such that $O^0 \setminus \overline{\text{Ad}(G)} = \emptyset$. Replacing O^0 with $H O^0$, if necessary, we may assume that O^0 is H -stable. Set $O = f[\mathfrak{g}; (u; v)] \cap M_G(H; \cdot; V) \cap u + {}_{H, V}(v) \cap O^0$. By definition, O is an open G -subvariety of X containing x . It is enough to show that any point $x_1 \in O$ with closed G -orbit and the determining triple $(H; \cdot; V)$ is G -conjugate to x . Assume the converse. Put $x_1 = [\mathfrak{g}; (u; v)]$, $u \in U$; $v \in V$; $(u; v) \notin 0$. Recall that ${}_{G, X}(x_1) = \text{Ad}(g)(\bar{x} + u + {}_{H, V}(v))$. Since ${}_{G, X}(x_1) = \emptyset$, Lemma 3.6 implies that $u + {}_{H, V}(v) = 0$. Since $U \setminus \mathfrak{h} = f0\mathfrak{g}$, $u = 0$. The subgroup $H_v \subset H$ is conjugate to H in G . Thus $v \in V^H = f0\mathfrak{g}$. Contradiction.

3.3. The proof of Theorem 3.1. At first, we obtain an estimate for the dimension of a fiber of ${}_{G, X}$.

Proposition 3.7. The dimension of any fiber of ${}_{G, X} : X \rightarrow X/G$ does not exceed $\dim X - \text{def}_G(X) - \frac{\text{cork}_G(X)}{2}$.

Proof. The proof is carried out in two steps. Firstly we consider the case when X satisfies equivalent conditions of Lemma 2.23 and then deduce the general case from this one.

Step 1. Suppose X satisfies equivalent conditions of Lemma 2.23. Then

$$\text{def}_G(X) + \frac{\text{cork}_G(X)}{2} = \frac{\dim X - \dim G + \text{rk } G}{2};$$

Let $y \in X/G$, x a point from a unique closed G -orbit in ${}_{G, X}^{-1}(y)$, $H = G_x$, $\mathfrak{h} = {}_{G, X}(x)$, $U = (z_g(\cdot) = \mathfrak{h})$; $V = (\mathfrak{g} \setminus x)^\perp = (\mathfrak{g} \setminus x)^\perp$. The H -modules $U \subset V$ and $T_x X/G$ are isomorphic.

Using the Luna slice theorem, we see that it is enough to check

$$(3.2) \quad \dim {}_{H, U \cup V}^{-1}(0) \leq \dim U + \dim V - \frac{\dim X - \dim G + \text{rk } G}{2}$$

Lemma 3.8 ([Sch2], Proposition 2.10). Let H be a reductive group, T_H its maximal torus, and V a self-dual H -module. Then

$$\dim {}_{H, V}^{-1}(0) \leq \frac{1}{2}(\dim V - \dim V^{T_H} + \dim H - \dim T_H):$$

Lemma 3.9. $U \subset V$ is a self-dual H -module.

Proof of Lemma 3.9. Replacing g with $z_g(s)$ if necessary, we may assume that $s \in z(\mathfrak{g})$ and easily reduce the proof to the case when \mathfrak{s} is nilpotent. The representation $H : V$ is symplectic and thus self-dual. Choose an \mathfrak{sl}_2 -triple $(\cdot; h; f) \subset \mathfrak{g}^H$ and consider the grading

$g = \sum_{i \in \mathbb{Z}} g_i$, where $g_i = f \otimes g_j h$; $j = i - 1$. Put $z_g(\lambda)_i = z_g(\lambda) \setminus g_i$. It is well known that $z_g(\lambda) = \sum_{i \geq 0} z_g(\lambda)_i$ and that $z_g(\lambda)_0$ is a reductive subalgebra of g . The latter implies that $z_g(\lambda)_0 = h \oplus U$ is an orthogonal H -module. Now it is enough to show that the H -module $z_g(\lambda)_i$ is self-dual.

Choose a connected subgroup $S \subset G$ corresponding to the \mathfrak{sl}_2 -triple (h, e, f) , and let w be an element of $N_S(h)$ acting on h by id . For $x, y \in z_g(\lambda)_i$ put $(x, y)_w = (x, wy)$. Since $w \in Z_G(H)$, the form $(\cdot, \cdot)_w$ is H -invariant. It follows from the representation theory of \mathfrak{sl}_2 that

$$[g, \lambda] \cdot z_g(f) = g \cdot w(z_g(\lambda)_i) = z_g(w(\lambda)) \setminus g_i = z_g(f) \setminus g_i.$$

Note that $z_g(\lambda)^2 = [g, \lambda] \cdot (g_i, g_j) = 0$ for $i + j \neq 0$. Hence the pairing of $z_g(\lambda)_i$ and $z_g(f) \setminus g_i$ induced by $(\cdot, \cdot)_w$ is nondegenerate. Equivalently, $(\cdot, \cdot)_w$ is nondegenerate on $z_g(\lambda)_i$.

We see that the H -module $U \oplus V$ satisfies the assumptions of Lemma 3.8. Let T_H be a maximal torus of H . Let us show that $\dim U^{T_H} > \text{rk } g - \text{rk } h$. Since $\dim h^{T_H} = \text{rk } h$, it is enough to show that $\dim z_g(\lambda)^{T_H} > \text{rk } g$ for any $\lambda \in g^H$. It is enough to check the last inequality for $\lambda \in g^H$ in general position. But in this case $z_g(\lambda)$ is semisimple. Hence $z_g(\lambda)$ is a Levi subalgebra of g and everything is clear.

By Lemma 3.8, we have the following inequalities

$$(3.3) \quad \begin{aligned} \dim_{H \cdot U \oplus V}^1(0) &\leq \frac{1}{2} (\dim U + \dim V - \dim U^{T_H} - \dim V^{T_H} + \dim h - \text{rk } h) \\ &\leq \frac{1}{2} (\dim U + \dim V - (\text{rk } g - \text{rk } h) + \dim h - \text{rk } h) \end{aligned}$$

It is checked directly that the last expression in (3.3) coincides with the r.h.s of (3.2).

Step 2. Now we consider the general case. Let X_0, G_0 be such as in Subsection 3.1.

By Proposition 3.3, $\text{cork}_G(X) = \text{cork}_{G_0}(X_0)$; $\text{def}_G(X) = \text{def}_{G_0}(X_0)$. The proposition will follow if we show that

$$(3.4) \quad \text{codim}_X^1(X) > \text{codim}_{X_0}^1(X_0);$$

for any $y \in X = G$. It follows from Proposition 3.3 that $X_0 = G_0 = X = G$, $\text{codim}_{X_0}^1(X_0) = \text{codim}_X^1(X) \setminus X_0$. Now (3.4) stems from the following general fact of Algebraic geometry:

$\dim_x Y \setminus Z > \dim_x Y + \dim_x Z - \dim X$ for any smooth subvarieties Y, Z of an irreducible smooth variety and $x \in Y \setminus Z$.

Proof of Theorem 3.1. Clearly, $e_{G/X}; G/X$ are equidimensional provided $b_{G/X}$ is. Any equidimensional morphism to a normal variety is open (see [Ch]).

To prove the theorem it remains to check that for all $\lambda \in g = G$ and any irreducible component Z of $\text{im}_{G/X}^1(\lambda)$ the equality $\dim_{G/X}(Z) = \dim X = G - \text{def}_G(X)$ and the inequality $\dim Z \leq \dim X - \text{def}_G(X)$ take place (the opposite inequality holds automatically, since $\text{def}_G(X) = \dim \text{im}_{G/X}^1(\lambda)$). The equality $\dim_{G/X}(Z) = \dim X = G - \text{def}_G(X)$ will imply

$$(3.5) \quad \dim_{G/X}(Z) = \dim X = G - \text{def}_G(X) + \dim Y$$

for an irreducible component Z of $\text{im}_{G/X}^1(Y)$, where $Y = \text{im}_{G/X}^1$ is an arbitrary closed irreducible subvariety (recall that, by Proposition 2.31, $\text{im}_{G/X}^1 = \text{im}_{G/X}^1 = G$ is an open subvariety in $a_{G/X}^{(1)} = W_{G/X}^{(1)}$). Thanks to Proposition 2.31, (3.5) holds if $\text{im}_{G/X}^1(Z)$ is an irreducible component in $\text{im}_{G/X}^1 = G(Y)$.

Choose a subvariety $S_{G/X}(H; \cdot; V) \subset G/X = G^{(1)}(\cdot)$ (see Subsection 3.2) such that $S_{G/X}(H; \cdot; V)$ is dense (and so, in virtue of Proposition 3.5, open) in $\text{im}_{G/X}^1(Z)$. Further,

choose a point $x \in Z \setminus \pi_{G,X}^{-1}(S_{G,X}(H; ;V))$ with closed G -orbit. Applying Proposition 2.27 to x , we may replace X with $M_G(H; ;V)$. Thanks to Remark 2.10, we may assume that $V^H = 0$. From Proposition 3.5 it follows that $\pi_{G,X}(Z)$ is a point. By Proposition 3.7, $\dim Z \leq \dim X - \text{def}_G(X) - \frac{1}{2} \text{cork}_G(X)$. It follows that $\text{cork}_G(X) = 0; \dim \pi_{G,X}^{-1}(x) = 0; \dim Z = \dim X - \text{def}_G(X)$. This verifies the claim in the beginning of the previous paragraph and completes the proof.

Corollary 3.10. For any $z \in \pi_{G,X}$ and any irreducible component Z of $\pi_{G,X}^{-1}(z)$ there exists an open subset $Z_0 \subset Z = G \cdot z$ such that Z_0 is smooth (as a variety), $\text{codim}_{Z=G \cdot z} Z_0 > 2$, and for any $z' \in Z_0$ and any point $x \in \pi_{G,X}^{-1}(z')$ with closed G -orbit the following condition holds:

(*) $M_G(H; ;V=V^H)$ is coisotropic, where $(H; ;V)$ is the determining triple of X at x . Moreover, $M_G(H; ;V=V^H)$ does not depend (up to an isomorphism) on the choice of z .

Proof. (*) is equivalent to $\text{cork}_G(X) = \text{cork}_G(M_G(H; ;V)) = \dim V^H$. It follows from Theorem 3.1 that Z maps dominantly whence, by the standard properties of quotient morphisms, surjectively onto some irreducible component of $\pi_{G,X}^{-1}(z)$. The required claims follow now from Proposition 3.5.

Corollary 3.11. Let Y be a closed irreducible subvariety in $\pi_{G,X}$. Then $\overline{\pi_{G,X}^{-1}(\mathcal{F})} = Y$ for any irreducible component \mathcal{F} of $\pi_{G,X}^{-1}(Y)$.

Proof. According to Theorem 3.1, $\pi_{G,X}^{-1}(\mathcal{F})$ is an irreducible component of $\pi_{G,X}^{-1}(Y) = X = G \cdot x$. It remains to apply Proposition 2.31.

Corollary 3.12. A simply connected affine conical Hamiltonian G -variety satisfies (Utw1).

Proof. Thanks to Proposition 2.32, $\pi_{G,X}^2 : C_{G,X} \rightarrow a_{G,X}^{(\cdot)} = W_{G,X}^{(\cdot)}$ is an isomorphism. By Theorem 3.1, the morphism $e_{G,X} : X \rightarrow C_{G,X}$ is equidimensional. Since G is connected, the subalgebra $C[X]^G$ is integrally closed in $C[X]$. Thus $C[C_{G,X}]$ is integrally closed in $C[X]$. In other words, a general fiber of $e_{G,X}$ is connected. Summarizing, we see that $e_{G,X}$ is an equidimensional morphism with a connected general fiber from a simply connected variety X to $C_{G,X} = a_{G,X}^{(\cdot)} = W_{G,X}^{(\cdot)}$. The proof of the proposition is based on an idea of Panyushev [Pa] and is completely analogous to that given in [Kn7], Theorem 7.2.

4. Some results concerning Weyl groups

Throughout the section $G;X;!$ have the same meaning as in the previous section.

In this section we study the structure of the Weyl group $W_{G,X}^{(\cdot)}$. Subsection 4.1 contains three technical propositions, which play a crucial role in the subsequent exposition. Propositions 4.1, 4.3 allows one to reduce the study of an arbitrary affine Hamiltonian G -variety to the study of a coisotropic conical model variety. Proposition 4.6 describes the behavior of Weyl groups under this reduction.

Using results of Subsection 4.1, in Subsection 4.2 we establish some properties of Weyl groups of varieties satisfying the equivalent conditions of Lemma 2.23. In particular, we get some restrictions on varieties with a "small" Weyl group (Proposition 4.9, Corollary 4.14) and show that a Weyl group cannot be "too small" (Corollary 4.16). As a consequence of Corollary 4.16 we get some explicit restrictions on Weyl groups for simple G of types A – E in Proposition 4.17, Corollary 4.19.

Finally, in Subsection 4.3 we compute the Weyl groups of linear actions of simple groups satisfying some additional restrictions. This computation will be used in Subsection 5.3 to check that any symplectic G -module is an untwisted Hamiltonian variety.

4.1. Some technical propositions.

Proposition 4.1. Let L be the principal centralizer and X_L an L -cross-section of X , $a_{G,X}^{(X_L)} = W_{G,X}^{(X_L)} a_{G,X}^{(X_L)}(\cdot)$, $M = Z_G(\cdot)$. Suppose $z \in b_{G,X}^1$. Choose an irreducible component Z of $b_{G,X}^1(\cdot)$. Then there exists $x \in X$ possessing the following properties:

- (a) $x \in Z$.
- (b) $_{G,X}(x)_s \in z(m) \setminus m^{\text{pr}}$.
- (c) A unique M -cross-section X_M of X containing x contains X_L and $b_{M,X_M}(x) = W_{M,X_M}^{(X_L)} a_{M,X_M}^{(X_L)}(\cdot)$.
- (d) Gx is closed in X . Let \mathcal{G} be a connected subgroup of M such that $(M; M)H = \mathcal{G}$. Automatically, $\mathcal{G}x$ is closed in X_M .
- (e) The Hamiltonian \mathcal{G} -variety $\mathcal{X} = M_{\widehat{G}}(H \setminus \mathcal{G}; \mathfrak{n}; V=V^H)$ is coisotropic, where $(H; \mathfrak{n}; V)$ is the determining triple of X_M (or, equivalently, of X) at x .

Remark 4.2. If X satisfies equivalent conditions of Lemma 2.23, then so does the Hamiltonian \mathcal{G} -variety \mathcal{X} . This stems easily from Proposition 2.27.

Proof of Proposition 4.1. Choose a point $z \in Z$ with closed G -orbit. Let us show that gz satisfies (b), (c) for some $g \in G$. Put $M_1 = Z_G(\cdot)_{G,X}(z)_s$. Since $_{G,X}(\cdot)_{G,X}(z)_s = _{G,X}(\cdot)$, we have $M_1 \subset G M$. Let X_{M_1} be an M_1 -cross-section of X containing z , L_1 be the principal centralizer and X_{L_1} an L_1 -cross-section of X_{M_1} . Replacing z with gz for an appropriate element $g \in G$, we may assume that $L_1 = L$; $X_{L_1} = X_L$. Next, replacing z with mz for some $m \in M_1$, one obtains $_{G,X}(z) \in a_{M_1,X_{M_1}}^{(X_L)} = a_{G,X}^{(X_L)}$. By the commutative diagram of assertion 2 of Lemma 2.30, for some $n \in N_G(L; X_L)$ the following equality holds

$$(4.1) \quad b_{M_1,X_{M_1}}^1(z) = W_{M_1,X_{M_1}}^{(X_L)} a_{M_1,X_{M_1}}^{(X_L)}(n):$$

Note that $_{M_1,X_{M_1}}(z) \in z(m_1) \neq m_1 = M_1$. From (4.1) it follows that $_{M_1,X_{M_1}}(n) \in z(m_1) \neq m_1 = M_1$ whence $n \in z(m_1)$. On the other hand, $n \in z(\text{Ad}(n)m) \setminus (\text{Ad}(n)m)^{\text{pr}}$ and so $m_1 \subset \text{Ad}(n)m$. We have seen above that $M_1 \subset G M$ whence $M_1 = nM n^{-1}$. Replacing z with $n^{-1}z$, we get the point z satisfying (a)–(c). Put $\cdot^0 = W_{M,X_M}^{(X_L)} a_{M,X_M}^{(X_L)}(\cdot)$.

According to Lemma 2.28, there exists an open M -saturated subvariety $X_M^0 \subset X_M$ containing z such that for any $x \in X_M^0$ the orbit Gx is closed in X if Mx is closed in X_M . Further, by Lemma 2.29, $\mathcal{G}x \subset X_M$ is closed whenever Mx is closed.

From assertion 2 of Lemma 2.30, Theorem 3.1 and the fact that the natural morphism $G_M X_M \rightarrow X$ is étale we get $\dim Z \setminus X_M = \dim b_{M,X_M}^1(\cdot^0)$. Hence there is an irreducible component Z^0 of $b_{M,X_M}^1(\cdot^0)$ containing z and contained in $Z \setminus X_M$. By Corollary 3.10, there is an open subset $Y^0 \subset b_{M,X_M}^1(Z^0)$ such that any point $x \in Y^0$ with closed M -orbit satisfies (a)–(d) and (e) for $\mathcal{G} = M$. When $\mathcal{G} \subsetneq M$, there is a covering $T_0 \rightarrow \mathcal{G}$ and a finite Hamiltonian morphism $T \rightarrow T_0$ $\mathcal{X} \rightarrow M_M(H \setminus \mathcal{G}; \mathfrak{n}; V=V^H)$, where T_0 is a torus. Since $H \subset \mathcal{G}$, we are done.

Proposition 4.3. Let $X; L; X_L$ be such as in Proposition 4.1, T_0 denote the unit component of the inefficiency kernel of the action $Z(L) : X_L, 0 \in a_{G/X}^{(X_L)}, M = Z_G(0)$. Suppose $0 \in \text{im } b_{G/X}$. Put $z = z(m) \setminus a_{G/X}^{(X_L)}; \underline{z} = \bigcap_{W_{G/X}^{(X_L)} a_{G/X}^{(X_L)}}(z)$. Choose an irreducible component \mathcal{F} of $b_{G/X}^{-1}(\underline{z})$. Let $z \in \mathcal{F}$ be a point in general position. Then there is a component Z of $b_{G/X}^{-1}(\bigcap_{W_{G/X}^{(X_L)} a_{G/X}^{(X_L)}}(z))$ lying in \mathcal{F} and a point $x \in Z$ satisfying the conditions (b)–(e) of Proposition 4.1 and

(f) $G_x = (M; M)T_0$ (where X_M is such as in (c)).

Remark 4.4. Under the assumptions of Proposition 4.3 one may assume that \mathcal{F} defined in (d) coincides with $(M; M)T_0$. If X satisfies the equivalent conditions of Lemma 2.23, then one can take $(M; M)$ for \mathcal{F} .

Proof of Proposition 4.3. The morphism $b_{G/X}$ is open, Theorem 3.1. So $Z; \mathcal{F}$ do exist. Choose a point $z \in Z$ satisfying conditions (a)–(e) and such that $b_{G/X}^{-1}(\mathcal{F})$ is the only component of $b_{G/X}^{-1}(G(\underline{z}))$ (see Theorem 3.1) containing $b_{G/X}(z)$.

By the choice of z , any irreducible component \mathcal{F}^0 of $b_{M/X_M}^{-1}(z)$ containing z is contained in $\mathcal{F} \setminus X_M$ (compare with the proof of Proposition 4.1). As in the proof of Proposition 4.1, there is an open subset $Y^0 \subset b_{M/X_M}^{-1}(\mathcal{F}^0)$ such that any $x \in b_{M/X_M}^{-1}(Y^0)$ with closed M -orbit satisfies conditions (a)–(e) (for appropriate λ).

It remains to prove that $M_x = (M; M)T_0$ for a general point $x \in \mathcal{F}^0$ with closed M -orbit. Recall (see the discussion preceding Proposition 3.3) that $L_0 = (L; L)T_0$ is the unit component of the principal isotropy group for the action $M : X_M$. Let C denote the principal isotropy subgroup for the action $M : \mathcal{F}^0$, so $L_0 \subset C$. By the definition of C , there exists an irreducible component X_1 of X_M^C such that $b_{M/X_M}^{-1}(X_1 \setminus \mathcal{F}^0)$ is dense in $b_{M/X_M}^{-1}(\mathcal{F}^0)$.

By Lemma 3.2, the action $N_M(C; X_1) : X_1$ is Hamiltonian with the moment map $\overline{N_M(C; X_1)} : X_1 \rightarrow \mathfrak{g}_1$. Since $0 \in \text{im } b_{G/X}$, we get $0 \in b_{M/X_M}^{-1}(\mathcal{F}^0)$, equivalently, $b_{M/X_M}^{-1}(X_1)$ contains a nilpotent element. Since C acts trivially on X_1 , we get

$$(4.2) \quad b_{M/X_M}^{-1}(X_1) \subset m^C \setminus (m + c^2)$$

for any $\lambda \in b_{M/X_M}^{-1}(X_1)$. Since there is a nilpotent element in $b_{M/X_M}^{-1}(X_1)$, we see that the r.h.s. of (4.2) coincides with $m^C \setminus c^2$. For brevity, put $s = m^C \setminus c^2$. This is an ideal in m^C .

Choose $x \in \mathcal{F}^0 \setminus X_1$ and put $\lambda = b_{M/X_M}^{-1}(x)$. Then $\lambda \in s$ and $(\lambda_s) + \lambda_n \in s$. Clearly, $c^2 \subset z(m^C)$. Thus $[\lambda_s; \lambda_n] = 0$ whence $\lambda_s = (\lambda_s)_s$ and

$$(4.3) \quad b_{M/X_M}^{-1}(x)_s \in s \setminus c^2; \lambda \in \mathcal{F}^0 \setminus X_1;$$

Lemma 4.5. $m = z + t_0 + [m; m]$.

Proof. It is enough to check that

$$(4.4) \quad t = z + t_0 + t_1; t_1 = t \setminus [m; m];$$

where t denotes a Cartan subalgebra of \mathfrak{l} . Recall that

$$\begin{aligned} t &= z(m) \setminus t_1; \\ z &= z(m) \setminus a_{G/X}^{(X_L)} = z(m) \setminus (z(\mathfrak{l}) \setminus t_0^2) = z(m) \setminus t_0^2; \end{aligned}$$

Since $z(\mathfrak{m}); t_1; t_0$ are the Lie algebras of algebraic groups, we see that $(\cdot; \cdot)$ is nondegenerate on $z(\mathfrak{m}); t_1; t_0; z$. To prove (4.4) it is enough to note that $t_0 + t_1 = z^2$.

If $c \in [\mathfrak{m}; \mathfrak{m}] + t_0$, then, thanks to Lemma 4.5, the r.h.s. of (4.3) is a proper subspace in z . Hence ${}_{M \times M}(\mathcal{Z}^0) = {}_{M \times M}(\mathcal{Z}^0 \setminus X_1)$ is not dense in z . Since $z \setminus \text{im } {}_{M \times M}$ is an open subset in z , we get a contradiction with Corollary 3.11.

Proposition 4.6. Let $X; L; X_L; M; X_M; \mathcal{G}$ be such as in Proposition 4.1, $\mathcal{B} = L \setminus \mathcal{G}$. Let $x \in X$ satisfy conditions (a)–(d) of Proposition 4.1 (for some Z) and \mathcal{X} be the model variety constructed by x as in (e). Then \mathcal{B} is the principal centralizer of \mathcal{X} and there is an \mathcal{B} -cross-section \mathcal{X}_L of \mathcal{X} such that $a_{\widehat{G \times X}}^{(\widehat{X_L})}$ is $W_{G \times X}^{(X_L)} \setminus M = L$ -stable subspace of $a_{G \times X}^{(X_L)}$ and $W_{\widehat{G \times X}}^{(\widehat{X_L})}$ lies in the image of $W_{G \times X}^{(X_L)} \setminus M = L$ in $GL(a_{\widehat{G \times X}}^{(\widehat{X_L})})$.

Proof. Recall, see Lemma 2.30, that \mathcal{B} is the principal centralizer and X_L is an \mathcal{B} -cross-section of the Hamiltonian \mathcal{G} -variety X_M . Let $(H; \cdot; V)$ denote the determining triple of X at x . Thanks to Lemmas 2.28, 2.29, $(H \setminus \mathcal{G}; \cdot; V = V^H \oplus V_0)$ is the determining triple of the Hamiltonian \mathcal{G} -variety X_M at x , where V_0 is a trivial $H \setminus \mathcal{G}$ -module. Put $\mathcal{X}^0 := M_{\widehat{G}}(H \setminus \mathcal{G}; \cdot; V = V^H \oplus V_0) = \mathcal{X} \oplus V_0$. It is enough to prove the analogue of the assertion of the proposition for \mathcal{X}^0 .

By Proposition 2.27, there is a \mathcal{G} -saturated analytical open neighborhood O of $[L; (0; 0)]$ in \mathcal{X}^0 , that is isomorphic (as a Hamiltonian \mathcal{G} -manifold) to a saturated analytical neighborhood of x in X_M . One may assume additionally that O is connected. By [Lo3], Lemma 5, $O_1 := \widehat{G \times X^0}(O)$ is an open neighborhood of $\widehat{G \times X^0}([L; (0; 0)])$ in $\mathcal{X}^0 = \mathcal{G}$. Further, according to Example 2.16, \mathcal{X}^0 is a conical Hamiltonian variety. Replacing O with a smaller neighborhood, we may assume that $tO \subset O$ for $0 \leq t \leq 1$. Note that \mathcal{B} is the principal centralizer of the Hamiltonian \mathcal{G} -variety \mathcal{X}^0 . Since $\mathcal{G}X_L$ is an open subvariety of X_M (in Zariski topology), we have $X_L \setminus O \neq \emptyset$. Choose an \mathcal{B} -cross-section \mathcal{X}_L^0 of \mathcal{X}^0 such that some connected component of $X_L \setminus O$ is contained in $\mathcal{X}_L^0 \setminus O$.

Lemma 4.7. The manifold $\mathcal{X}_L^0 \setminus O$ is connected.

Proof of Lemma 4.7. Let $(\cdot; h; f)$ be an \mathfrak{sl}_2 -triple in $\mathfrak{h}^H \setminus \widehat{G}$ generating the model variety \mathcal{X}^0 . Note that the action $C : \mathcal{X}^0$ preserves \mathcal{X}_L^0 . Let $Y^0; Y^1$ be two distinct connected components of $\mathcal{X}_L^0 \setminus O$, $y^i \in Y^i$, $i = 0, 1$; and y^t , $0 \leq t \leq 1$; a continuous curve connecting $y^0; y^1$ in \mathcal{X}_L^0 . There is a positive real $\epsilon < 1$ such that $y^t \notin O$ for all t ; $0 \leq t \leq 1$. Finally, note that $y^i \in Y^i$ for all real t such that $-\epsilon \leq t \leq \epsilon$ and $i = 0, 1$. Therefore $t \mapsto y^t$ is a continuous curve in $\mathcal{X}_L^0 \setminus O$ connecting points from $Y^0; Y^1$. Contradiction.

One easily deduces Proposition 2.27 that $a_{\widehat{G \times X^0}}^{(\widehat{X_L^0})} = a_{\widehat{G \times X_M}}^{(X_L)}$. The equalities $W_{\widehat{G \times X_M}}^{(X_L)} = W_{M \times M}^{(X_L)} = W_{G \times X}^{(X_L)} \setminus M = L$ hold, see [Lo5], Lemmas 6.10, 6.13. By Lemma 4.7, $N_{\widehat{G}}(\mathcal{B}; X_L^0) = N_{\widehat{G}}(\mathcal{B}; X_L^0 \setminus O)$. It remains to recall that $X_L^0 \setminus O \not\subset X_L$ whence $N_{\widehat{G}}(\mathcal{B}; X_L^0 \setminus O) = N_{\widehat{G}}(\mathcal{B}; X_L)$.

$$\begin{array}{ccccccc}
x^0 & \xrightarrow{\quad} & x^0 & \circ & \overline{x_M} & \xrightarrow{\quad} & \overline{x} \\
B & & B & & @ & & | \\
B & b_{G;x^0} & b_{G;x^0} & & @ & b_{M;x_M} & | \\
B & \hat{x} & \hat{x} & & @ & @ & b_{G;x} \\
B & & & & @ & @ & | \\
B & & & & @ & @ & ? \\
B & & & & @ & @ & ? \\
a_{G;\hat{x}}^{(\hat{x}_L)} = W_{G;\hat{x}}^{(\hat{x}_L)} & \xrightarrow{\quad} & a_{G;x_M}^{(x_L)} = W_{G;x_M}^{(x_L)} & \xrightarrow{\quad} & a_{M;x_M}^{(x_L)} = W_{M;x_M}^{(x_L)} & \xrightarrow{\quad} & a_{G;x}^{(x_L)} = W_{G;x}^{(x_L)}
\end{array}$$

Proposition 4.9. Let G be a connected reductive group, H its reductive subgroup, χ an H -invariant nilpotent element of G , and V a symplectic H -module. Suppose $X := M_G(H; \chi; V)$ satisfies the equivalent conditions of Lemma 2.23 and $\text{cork}_G(X) = 0$.

- (1) If $W_{G,X}^{(\cdot)} = \text{flg}$, then
 (*) $= 0; (G; G) = H, (G; G) = G_1 \quad \dots \quad G_k$ for some k , where $G_i = \text{SL}_2$. Finally,
 the G -modules $V = V^{(G; G)}$ and $V_1 \oplus V_2 \oplus \dots \oplus V_k$ are isomorphic, where V_i is the
 direct sum of two copies of the two-dimensional irreducible G_i -module.
 (2) Conversely, if G is semisimple, and X satisfies (*), then $W_{G,X}^{(\cdot)} = \text{flg}$.

Proof. Suppose, at first, that G is semisimple. Let us prove the first assertion.

It follows from Proposition 2.32 that $b_{G,X} = G : X = G ! a_{G,X}^{()} = W_{G,X}^{()}$ is an isomorphism. The equality $W_{G,X}^{()} = \text{flg}$ is equivalent to the condition that $C[X]^G$ is generated by elements of degree 2 (the grading on $C[X]^G$ is introduced in Remark 2.17).

The morphism $M_G(H; ; V) ! M_G(H; ; V); [g; (u; v)] \mapsto [g; (u; v)]$ satisfies the assumptions of assertion 4 of Lemma 2.30. Thus $W_{G, M_G(H; ; V)}^{()} = \text{flg}$ and we may assume that H is connected.

Put $U = (z_g() \setminus h^2)$. Let us equip the algebra $C[U \cup V]$ with the grading described in Remark 2.17. The algebra $C[U \cup V]^H$ is generated by elements of degree 2. Recall that $2U$ has degree 4. Therefore $U \cap h^2 = 0$. Any element from $U \cap h^2$ has degree 2. Therefore $U \cap h^2 = U \cap H = U^H$, equivalently, $C[U \cap U^H]^H = C$. But the H -module $U \cap U^H$ is orthogonal (that is, possesses a nondegenerate H -invariant symmetric form) because U is orthogonal. Hence $U \cap U^H = U^H$. Equivalently, $g = h + g^h$. In particular, h is an ideal of g . By Lemma 2.23,

$$(4.5) \quad \dim X = \dim g + \text{rk } g;$$

But $\dim X = 2 \dim g - 2 \dim h + \dim V$. Note that $m_H(V) = \dim H$, since $m_G(X) = \dim G$ and $g \cap h$ is a trivial h -module. Since V is a symplectic H -module, we have

$$(4.6) \quad \dim V = \dim h + \text{rk } h + \text{cork}_H(V);$$

From (4.5), (4.6) it follows that

$$(4.7) \quad \dim g + \text{rk } g = \dim X = 2 \dim G = H + \dim V > 2 \dim g - \dim h + \text{rk } h;$$

We deduce from (4.7) that $\dim g - \text{rk } g \leq \dim h - \text{rk } h$. Since h is an ideal of g , the last inequality is equivalent to $g = h$.

Let $G = G_1 \times \dots \times G_k$ be the decomposition into the locally direct product of simple subgroups. According to Lemma 6.14 from [Lo5], $W_{G_i, V}^{()} = \text{flg}$. By Propositions 4.1, 4.3, there exists a point $x \in \pi_{G_i, V}^{-1}(0)$ satisfying the conditions (a)–(f) of those propositions (with $G; X$ replaced with $G_i; V$). Let $(H_0; 0; V_0)$ be the determining triple of the G_i -variety V at x . Thanks to Proposition 4.6, $W_{G_i, M_{G_i}(H_0; 0; V_0)}^{()} = \text{flg}$. By assertion 1, $0 = 0; H_0 = G_i$ whence $V_0 = V$. Further, $V = V^{G_i}$ is a coisotropic G_i -module of dimension $\dim g_i + \text{rk } g_i$. Using the classification of coisotropic modules obtained in [Lo1], [Kn6], we get $G_i = \text{SL}_2$. Since $W_{G_i, V = V^{G_i}}^{()} = \text{flg}$, we see that $C[V = V^{G_i}]$ is generated by an element of degree 2. One easily deduces from this that $V = V^{G_i}$ is isomorphic to the direct sum of two copies of the irreducible 2-dimensional SL_2 -module.

Since $V = V^{G_i} = V^{G_i \setminus}$ is a symplectic G -module and $\text{Sp}(V = V^{G_i})^{G_i}$ is a torus, the group $\prod_{j \neq i} G_j$ acts trivially on $V = V^{G_i}$. Note that $(\prod_{i=1}^k V^{G_i} \setminus) \setminus = \prod_{i=1}^k V^{G_i} = V^G = 0$. The last equality holds because $\text{cork}_G(V) = 0$. To complete the proof of assertion 1 note that $\prod_{i=1}^k G_i$ acts on V effectively. It follows that the natural epimorphism $\prod_{i=1}^k G_i \rightarrow G$ is an isomorphism.

Now suppose that X is of the form indicated in (*). It is enough to check the equality $W_{G,X}^{()} = \text{flg}$ for $k = 1$. Here the equality follows from the observation that $C[X]^G$ is generated by an element of degree 2.

Proceed to the case when G is not necessarily semisimple. Let x be a point of X satisfying conditions (a)–(f) of Propositions 4.1, 4.6 for $M = G$, $\mathcal{G} = (G; G)$ and \mathcal{X} be the model variety constructed by $\mathcal{G}; x$ in (e). By Proposition 4.6, $W_{\widehat{G}, \widehat{\mathcal{X}}}^{()} = \text{flg}$. Therefore $(G; G) = \prod_{i=1}^k G_i; \mathcal{X} = \prod_{i=1}^k V_i$. Since any stabilizer of a point with closed G -orbit is conjugate to a

subgroup of H , we have $(G; G) \cap H; = 0$. By above, there is a point $x \in V^{(G; G)} \cap X$ such that $V = g \cdot x = \sum_{i=1}^k V_i$. This observation completes the proof.

Now we are going to obtain a sufficient condition for $W_{G; X}^{(\cdot)}$ to intersect any subgroup of $W(g)$ conjugate to a certain fixed subgroup. To state the corresponding assertion we need some definitions.

Definition 4.10. A subset $A \subset W(g)$ is called completely perpendicular if the following two conditions take place:

- (1) If $x \in A$, then $-x \in A$.
- (2) $\text{Span}_{\mathbb{R}}(A) \cap W(g) = A \cap W(g)$.

For example, any one-element subset of $W(g)$ is completely perpendicular.

Definition 4.11. A pair $(h; V)$, where h is a reductive subalgebra of \mathfrak{g} and V is an h -module, is said to be a g -stratum. Two g -strata $(h_1; V_1), (h_2; V_2)$ are called equivalent if there exists $g \in G$ and a linear isomorphism $\varphi: V_1 = V_1^{h_1} \rightarrow V_2 = V_2^{h_2}$ such that $\text{Ad}(g)h_1 = h_2$ and $(\text{Ad}(g))'(\varphi(v_1)) = \varphi(v_1)$ for all $v_1 \in V_1 = V_1^{h_1}$.

Definition 4.12. Let Y be a smooth affine variety and $y \in Y$ a point with closed G -orbit. The pair $(g_y; T_y Y = g_y)$ is called the g -stratum of y . We say that $(h; V)$ is a g -stratum of Y if $(h; V)$ is equivalent a g -stratum of a point of Y . In this case we write $(h; V) \subset_g Y$.

Remark 4.13. Let us justify the terminology. Pairs $(h; V)$ do define some stratification of $Y = G \backslash Y$ by varieties with quotient singularities. Besides, analogous objects were called "strata" in [Sch3], where the term is borrowed from.

Let A be a nonempty completely perpendicular subset of $W(g)$. By $S^{(A)}$ we denote the g -stratum $(g^{(A)}; \sum_{2A} V)$, where V is, by definition, the direct sum of two copies of the two-dimensional irreducible $g^{(A)} = g^{(\text{Anf } g)}$ -module.

Corollary 4.14. If $W_{G; X}^{(\cdot)} \setminus W(g^{(A)}) = \text{flg}$, then $S^{(A)} \subset_g X$.

Proof. Put $M = Z_G(\sum_{2A} \ker)$. We remark that $G^{(A)} = (M; M)$. Choose a point $x \in X$ satisfying conditions (a)–(f) of Propositions 4.1, 4.3 for general $\lambda \in z(\mathfrak{m})$. Let $(H; ; V)$ be the determining triple of X at x and $\mathfrak{X} = M_{G^{(A)}}(H \setminus G^{(A)}; ; V = V^H)$. By Proposition 4.6, $W_{G^{(A)}; X}^{(\cdot)} = \text{flg}$. Using Proposition 4.9, we see that $S^{(A)}$ is equivalent to the g -stratum of x .

Now we obtain some restriction on $W_{G; X}^{(\cdot)}$, namely, we check that $W_{G; X}^{(\cdot)}$ is large in the sense of the following definition.

Definition 4.15. A subgroup $W(g)$ is said to be large if for any roots $\alpha; \beta \in W(g)$ such that $\alpha; \beta; (\alpha; \beta) \neq 0$ there exists $s \in \mathbb{R}^+ + \mathbb{R}^-$ with $s \neq 0$.

Corollary 4.16. The subgroup $W_{G; X}^{(\cdot)} \cap W(g)$ is large.

Proof. Assume the converse. Choose $\alpha; \beta \in W(g)$ such that $\alpha; \beta; (\alpha; \beta) \neq 0$ but $s \in \mathbb{R} \setminus \{0\}$ for all $s \in W(g) \setminus (\mathbb{R}^+ + \mathbb{R}^-)$. Put $M = Z_G(\ker \setminus \ker)$. Note that $(M; M) = G^{(\cdot)}$. Let $x \in X$ satisfy conditions (a)–(f) of Propositions 4.1, 4.3 for general $\lambda \in z(\mathfrak{m})$. Let $(H; ; V)$ be the determining triple of X at x . Put $\mathfrak{X} = M_{G^{(\cdot)}}(H \setminus G^{(\cdot)}; ; V = V^H)$. It follows from Proposition 4.6 that $W_{G^{(\cdot)}; X}^{(\cdot)}$ does not contain a reflection. Let \mathcal{C} denote

the simply connected covering of $G^{(\cdot)}$. It is a simple simply connected group of rank 2. Further, denote by \mathbb{H} the connected normal subgroup of \mathbb{G} corresponding to h . Put $\mathbb{X} = M_{\tilde{G}}(\mathbb{H}; n; V=V^H)$. It is a coisotropic variety. There is a natural morphism $\mathbb{X} \rightarrow \mathbb{P}^1$ satisfying the assumptions of the forth assertion of Lemma 2.30. Therefore the group $W_{\tilde{G}, \tilde{\mathbb{X}}}^{(\cdot)}$ does not contain a reflection. On the other hand, by Corollary 3.12, the group $W_{\tilde{G}, \tilde{\mathbb{X}}}^{(\cdot)}$ is generated by reflections. Therefore $W_{\tilde{G}, \tilde{\mathbb{X}}}^{(\cdot)} = \text{flg}$. This contradicts Proposition 4.9.

Now let us describe large subgroups of $W(g)$ for simple groups G of types $A \dots E$.

Firstly, we consider the situation when g is simple and has type $A; D; E$, in other words, when all elements of (g) are of the same length.

Recall the classification of maximal proper root subsystems in (g) (see [D]). We fix a system $\alpha_1; \dots; \alpha_r \in (g)$ of simple roots. Let α_0 be the minimal root and $n_1; \dots; n_r$ (uniquely determined) nonnegative integers satisfying $\alpha_0 + n_1\alpha_1 + \dots + n_r\alpha_r = 0$. A proper root subsystem $\beta \subset (g)$ is maximal if it is $W(g)$ -conjugate to one of the following root subsystems.

- (a) $\text{Span}_{\mathbb{Z}}(\alpha_1; \dots; \alpha_{i-1}; \alpha_{i+1}; \dots; \alpha_r) \setminus (g)$ for $n_i = 1$.
- (b) $\text{Span}_{\mathbb{Z}}(\alpha_0; \alpha_1; \dots; \alpha_{i-1}; \alpha_{i+1}; \dots; \alpha_r) \setminus (g)$ for prime n_i .

The number n_i depends only on α_0 . We will call this number the characteristic of β .

For a proper subgroup $\beta \subset W(g)$ let β denote the set of all $\alpha \in (g)$ such that $s_\alpha \in \beta$.

Proposition 4.17. Let g be a simple Lie algebra of type $A; D; E$, $\text{rk } g > 1$, and β a proper subgroup in $W(g)$. Then β is large if β is a maximal proper root subsystem in (g) of characteristic 1 or 2.

Lemma 4.18. Let g be a simple Lie algebra of type $A; D; E$. Then β is a root subsystem in (g) for any subgroup $\beta \subset W(g)$.

Proof. Let $\alpha \in (g)$. Since all roots of (g) are of the same length, we see that $\alpha + 2\alpha \in (g)$, (resp., $\alpha - 2\alpha \in (g)$) if $l(\alpha) < 0$, (resp., $l(\alpha) > 0$).

By its definition, β has the following property:

- (A) β coincides with the set of all roots $\alpha \in (g)$ such that s_α is contained in the subgroup of $W(g)$ generated by elements of β .

We need to check that $\alpha \in \beta$ implies $\alpha + 2\alpha \in \beta$ and that $\alpha \in \beta; \alpha + 2\alpha \in (g)$ imply $\alpha + 2\alpha \in \beta$. The first implication follows directly from (A). To prove the second one we note that $\alpha + 2\alpha = s_{\alpha+\alpha}$ whenever $\alpha \in \beta; \alpha + 2\alpha \in (g)$.

Proof of Proposition 4.17. The subgroup $\beta \subset W(g)$ is big if

- (B) $\alpha \in \beta; \alpha + g \setminus \beta \neq \emptyset$ for all $\alpha \in (g)$ such that $\alpha + 2\alpha \in (g)$.

One checks directly that a maximal root subsystem $\beta \subset (g)$ of characteristic 1 or 2 satisfies (B).

Now let β be a root subsystem of (g) satisfying (B). At first, assume that β is not maximal. Let β_1 be a maximal proper root subsystem of (g) containing β . Choose $\alpha \in \beta_1 \setminus \beta$. We see that $\alpha \in \beta$ for all $\alpha \in \beta_1$. Otherwise $\alpha \in g \setminus \beta_1 = ?$. Analogously, $\alpha \in (g)$. Therefore $l(\alpha) = 0$. In particular, $\alpha \in [f, g]$ is a root subsystem of (g) . By the choice of β_1 , $\alpha \in [f, g] = (g)$. This is impossible because g is simple.

It remains to show that the characteristic of \mathbb{F} is less than 3. Assume that $\mathbb{F} = \text{Span}_{\mathbb{Z}} \{f_0, \dots, f_{i-1}, f_{i+1}, \dots, f_n\} \setminus (g)$, where $n_i > 2$. Let \bar{i} denote the dual fundamental weight corresponding to i . The subset $\mathcal{W}_i(g)$ coincides with the set of all $\lambda \in \mathfrak{h}^*$ such that $n_i \langle \lambda, \bar{i} \rangle \in \mathbb{Z}$. So it is enough to check that there are $\lambda \in \mathcal{W}_i(g)$ such that $\langle \lambda, \bar{i} \rangle = 1$ and $\langle \lambda, \bar{i} \rangle = 2$. There is $\lambda \in \mathcal{W}_i(g)$ with $\langle \lambda, \bar{i} \rangle = 1$. Choose such an element $\lambda = \sum_{j=1}^r m_j \bar{j}$ such that m_j is minimal possible. One sets $\lambda = \bar{i}$; $\lambda = \bar{i} + 2\bar{j}$ (g).

Corollary 4.19. Suppose g is a simple classical Lie algebra. Then $\mathcal{W}_i(g)$ is large if i is listed in Table 4.1.

Table 4.1: Subsets $\mathcal{W}_i(g)$ for large subgroups $\mathcal{W}_i(g)$ when g is classical

g	
$A_1; l > 2$	$f_{i-1} - f_{i+1}; j \geq 2 \text{ I or } i; j \notin I; I \cap \{f_1, \dots, f_{n+1}\} \neq \emptyset$
$B_1; l > 3$	(a) $f_{i-1} - f_{i+1}; j \geq 2 \text{ I or } i; j \notin I; I \cap \{f_1, \dots, f_n\} \neq \emptyset$ (b) $f_{i-1} - f_{i+1}; j \geq 2 \text{ I or } i; j \notin I; I \cap \{f_1, 2f_2, \dots, f_n\} \neq \emptyset$ (c) $f_{i-1} - f_{i+1}; j \geq 2 \text{ I or } i; j \notin I; I \cap \{(f_1 + f_2); i \geq 2 \text{ I; } j \notin I\} \neq \emptyset$
$C_1; l > 2$	(a) $f_{i-1} - f_{i+1}; j \geq 2 \text{ I or } i; j \notin I; I \cap \{2f_1, \dots, f_n\} \neq \emptyset$ (b) $f_{i-1} - f_{i+1}; j \geq 2 \text{ I or } i; j \notin I; I \cap \{2f_1, 2f_2, \dots, f_n\} \neq \emptyset$ (c) $f_{i-1} - f_{i+1}; j \geq 2 \text{ I or } i; j \notin I; I \cap \{(f_1 + f_2); i \geq 2 \text{ I; } j \notin I\} \neq \emptyset$
$D_1; l > 3$	(a) $f_{i-1} - f_{i+1}; j \geq 2 \text{ I or } i; j \notin I; I \cap \{f_1, \dots, f_n\} \neq \emptyset$ (b) $f_{i-1} - f_{i+1}; j \geq 2 \text{ I or } i; j \notin I; I \cap \{(f_1 + f_2); i \geq 2 \text{ I; } j \notin I\} \neq \emptyset$

Note that some subsets appear in Table 4.1 more than once.

Proof. For g of type A_1 or D_1 the required assertion stems directly from Proposition 4.17.

Suppose $g = \mathfrak{sp}_{2l}; l > 2$. If $l = 2$, then $\mathcal{W}_i(g)$ is large if $i \in \{1, 2\}$. All nonempty subsets $\mathcal{W}_i(\mathfrak{sp}_4)$ do appear in Table 4.1. Now suppose $l > 2$. Let \mathcal{W}_0 denote the subset of all short roots in (g) and W_0 the subgroup of $W(g)$ generated by $s_i; i \geq 2$. Note that W_0 is the Weyl group of the root system D_{l-1} . By the definition of a large subgroup, the subgroup \mathcal{W}_0 generated by $s_i; i \geq 2$ is large in W_0 . If $\mathcal{W}_0 \setminus \mathcal{W}_i$ is of type (a) (see Table 4.1), then \mathcal{W}_i is large in $W(g)$ if $\mathcal{W}_0 \setminus \mathcal{W}_i$ contains a long root. If $\mathcal{W}_0 \setminus \mathcal{W}_i$ is of type (b) or (c), then \mathcal{W}_0 is large in $W(g)$. Since $N_{W(g)}(\mathcal{W}_0) = W_0$, we see that large subgroups in $W(g)$ are precisely those presented in Table 4.1.

The proof for $g = \mathfrak{so}_{2l+1}; l > 2$; follows easily from the duality between the root systems $B_l; C_l$.

4.3. Examples of computation of Weyl groups. In this subsection we classify pairs $(G; V)$, where G is a simple algebraic group, and V is a symplectic G -module such that $\text{def}_G(V) = \text{rk } G$, $W_{G,V}^{(\cdot)} \in W(g)$. The computation for $V = U \oplus U$ (and, more generally, $X = T(G \ltimes V)$) is made in [Lo4], Section 7, so here we consider only the case $X \in U \oplus U$.

Lemma 4.20. Let G be a simple group, $X = M_G(H; \cdot; V)$, where H is nilpotent, satisfy the equivalent conditions of Lemma 2.23. If $s \notin W_{G,X}^{(\cdot)}$ for some $s \in W(g)$, then there exist a

subalgebra $s_{\mathfrak{h}}$ such that $s_{\mathfrak{g}} \mathfrak{g}^{(\cdot)}$ and

$$(4.8) \quad \frac{\mathrm{tr}_{U \oplus V}(\mathfrak{h}^2)}{\mathrm{tr}_{\mathfrak{h}}(\mathfrak{h}^2)} = 1 - \frac{4}{\mathrm{tr}_{\mathfrak{h}}(\mathfrak{h}^2)} :$$

Here $U = z_{\mathfrak{g}}(\mathfrak{h})$ and \mathfrak{h} is a coroot in $s_{\mathfrak{h}}$.

Proof. By Corollary 4.14, $S^{(\cdot)} = \mathfrak{g} \times X$. Equivalently, there is a subalgebra $s_{\mathfrak{h}}$ such that $s_{\mathfrak{g}} \mathfrak{g}^{(\cdot)}$ and $(s_{\mathfrak{g}} \mathbb{C}^2 \oplus \mathbb{C}^2) \cap U \oplus V$. The last condition implies that the s -modules $\mathfrak{h} = s_{\mathfrak{g}}(\mathbb{C}^2)^2$ and $U \oplus V$ differ by a trivial summand. Comparing the traces of \mathfrak{h}^2 on these two modules, we get the claim.

Here is the main result of this subsection.

Proposition 4.21. Let G be a simple group and V a symplectic G -module satisfying the equivalent conditions of Lemma 2.23 such that $V \not\subset U \oplus U$ for any G -module U . Then $W_{G/V}^{(\cdot)} \not\subset W(\mathfrak{g})$ if V is contained in Table 4.2. The group $W_{G/V}^{(\cdot)}$ is presented in the forth column of the table.

Table 4.2: G -modules V such that $W_{G/V}^{(\cdot)} \not\subset W(\mathfrak{g})$

N	\mathfrak{g}	V	$W_{G/V}^{(\cdot)}$
1	$\mathfrak{g} = \mathfrak{sl}_6$	$V = V(\mathfrak{e}_3) \oplus V(\mathfrak{e}_1)^2 \oplus V(\mathfrak{e}_5)^2$	$A_1 \oplus A_3$
2	$\mathfrak{g} = \mathfrak{sp}_4$	$V = V(\mathfrak{e}_1) \oplus V(\mathfrak{e}_2)^2$	$C_1 \oplus C_1$
3	$\mathfrak{g} = \mathfrak{sp}_6$	$V = V(\mathfrak{e}_3) \oplus V(\mathfrak{e}_1)^2$	$C_1 \oplus C_2$
4	$\mathfrak{g} = \mathfrak{so}_{11}$	$V = V(\mathfrak{e}_5) \oplus V(\mathfrak{e}_1)^4$	$B_1 \oplus B_4$
5	$\mathfrak{g} = \mathfrak{so}_{13}$	$V = V(\mathfrak{e}_6) \oplus V(\mathfrak{e}_1)^2$	$B_2 \oplus B_4$

In the fourth column we indicate the type of a root subsystem in (\mathfrak{g}) such that the reflections corresponding to its roots generate $W_{G/V}^{(\cdot)}$. By B_1 (resp., C_1) we mean a root subsystem containing two opposite short (resp., long) roots in B_n (resp., C_n). Root subsystems indicated in column 4 are determined uniquely up to $W(\mathfrak{g})$ -conjugacy.

Proof of Proposition 4.21. By Corollary 4.14, $S^{(\cdot)} = \mathfrak{g} \times V$ for some $\mathfrak{e} \in 2(\mathfrak{g})$. There is an SL_2 -stable prime divisor D^0 on $\mathbb{C}^2 \oplus \mathbb{C}^2$ such that $m_{SL_2}(D^0) = 2$. Applying the Luna slice theorem, we see that there is a prime G -stable divisor D on V such that $m_G(D) < \dim G$. All G -modules V with $m_G(V) = \dim G$ possessing such a divisor D were classified by Knop and Littelman, [KnL]. All such symplectic modules V such that $V \not\subset U \oplus U$ are presented in Table 4.2. Let us show that for these modules the inequality $W_{G/V}^{(\cdot)} \not\subset W(\mathfrak{g})$ does hold.

Case 1. $\mathfrak{g} = \mathfrak{sl}_6$; $V = V(\mathfrak{e}_3) \oplus V(\mathfrak{e}_1)^2 \oplus V(\mathfrak{e}_5)^2$. We can consider V as a symplectic $\mathfrak{G} = SL(6) \times SL(2)$ -module, where $SL(2)$ acts on $V(\mathfrak{e}_1)^2 \oplus V(\mathfrak{e}_5)^2$ as on $\mathbb{C}^2 \oplus (V(\mathfrak{e}_1) \oplus V(\mathfrak{e}_5))$. This module has the discrete stabilizer in general position and is coisotropic, see [Kn6], [Lo1]. The Weyl group $W_{\tilde{G}/V}^{(\cdot)}$ was computed in [Kn6], Table 12, it corresponds to the root system $A_1 \oplus A_1 \oplus A_3$. By assertion 2 of Lemma 6.14 from [Lo5], $W_{\tilde{G}/V}^{(\cdot)} = W_{SL_2/V}^{(\cdot)} \times W_{G/V}^{(\cdot)}$. It follows that $W_{G/V} = A_1 \oplus A_3$.

Case 2. $\mathfrak{g} = \mathfrak{sp}_4$; $V = V(\mathfrak{e}_1) \oplus V(\mathfrak{e}_2)^2$. We can consider V as a symplectic $\mathfrak{G} = Sp(4) \times C$ -module. Again, this module is coisotropic and has the discrete stabilizer in general

position. Using results of Knop, [Kn6], we see that $W_{\tilde{G}/V}^{(\cdot)} = A_1 \oplus A_1$. But $W_{\tilde{G}/V}^{(\cdot)} = W_{\tilde{G}/V}^{(\cdot)}$, see assertion 3 of Lemma 2.30. Using Lemma 4.20, we see that $s \in W_{\tilde{G}/V}^{(\cdot)}$ for all long roots α .

Case 3. $\mathfrak{g} = \mathfrak{sp}_6$; $V = V(\alpha_3) \oplus V(\alpha_1)^2$. One argues exactly as in the previous case.

Before proceeding to the remaining two cases let us make some remarks.

Firstly, $s \in W_{\tilde{G}/V}^{(\cdot)}$ for all short roots α . One checks this using Lemma 4.20 (the fraction in the l.h.s. of (4.8) (the index of the G -module V) can be computed using the Table 1 of [AEV]).

Since $s \in W_{\tilde{G}/V}^{(\cdot)}$ for any short root α , it follows from Corollary 4.16, Proposition 4.17 that $W_{\tilde{G}/V}^{(\cdot)}$ is either the whole Weyl group $W(\mathfrak{g})$ or is maximal among all proper subgroups generated by reflections. The latter is the case if $C[V]^G = Z(C[V]^G)$ contains two linearly independent elements of degree 4.

Case 4. $\mathfrak{g} = \mathfrak{so}_{11}$; $V = V(\alpha_5) \oplus V(\alpha_1)^4$.

Note that $\mathrm{Sp}(V)^G = \mathrm{Sp}(4)$. We consider the \mathbb{Z}^2 -grading on $C[V]$ induced by the degrees with respect to $V(\alpha_5)$ and $V(\alpha_1)^4$. By [Sch2], $C[V]^G$ is freely generated by a 21-dimensional subspace $U \subset C[V]^G$ such that

- (1) U is an $\mathrm{Sp}(4)$ -submodule in $C[V]^G$.
- (2) There is the decomposition $U = U_1 \oplus U_2 \oplus U_3 \oplus U_4$, where $U_1 = S^2 C^4$; $U_2 = V^2 C^4$; $U_3 = C^4$; $U_4 = C$ (isomorphisms of $\mathrm{Sp}(4)$ -modules).
- (3) U_i ; $i = \overline{1, 4}$; is homogeneous with respect to the \mathbb{Z}^2 -grading on $C[V]$. The degrees of $U_1; U_2; U_3; U_4$ are $(2; 0); (2; 2); (1; 2); (0; 4)$, respectively.

Let us check that $U_4 \subset Z(C[V]^G)$. Since $V(\alpha_5)$ and $V(\alpha_1)^4$ are skew-orthogonal, we get $fU_4; U_1g = 0$. Suppose $fU_4; U_2g \notin f0g$. Since $U_4 \subset C[V]^G \subset \mathrm{Sp}(4)$, we see that $fU_4; U_2g$ is isomorphic (as an $\mathrm{Sp}(4)$ -module) to $V^2 C^4$. But $fU_4; U_2g$ consists of homogeneous elements of degree $(2, 4)$ whence $fU_4; U_2g \subset U_1U_4 + U_3^2$. The both summands are isomorphic to $S^2 C^4$. This contradicts $fU_4; U_2g \notin f0g$. Finally, the degree of $fU_3; U_4g$ equals $(1; 4)$ whence $fU_3; U_4g = 0$.

Let q denote a homogeneous element in $C[V]^G$ corresponding to an invariant nondegenerate form. Then $W_{\tilde{G}/V}(q)$ is a homogeneous element of $C[V]^G$ of degree 4. It remains to check that $W_{\tilde{G}/V}(q) \notin U_4$. One checks easily that in $W_{\tilde{G}/V}(\alpha_1)^2$ contains an element α such that $(\alpha; \alpha) \neq 0$. If $v \in V(\alpha_1)^2$, $\alpha \cdot v$ is such that $W_{\tilde{G}/V}(\alpha \cdot v) = \alpha$, then $W_{\tilde{G}/V}(q(v)) \neq 0$; $f(v) = 0$ for any $f \in U_4$.

By Proposition 4.17, $W_{\tilde{G}/V}^{(\cdot)}$ corresponds either to $B_1 \oplus B_4$ or to $B_2 \oplus B_3$. Thanks to Corollary 4.14, it remains to prove that $S^{(A)} \notin \mathfrak{g}_V$, where $A = f''_1 \oplus f''_2 \oplus f''_3 \oplus f''_4 g$. If $S^{(A)} \in \mathfrak{g}_V$, then

$$(4.9) \quad \dim V^{g^{(A)}} + \dim \mathfrak{g} = \dim \mathfrak{g}_V(g^{(A)}) = \dim V - 8:$$

But $\dim V(\alpha_1)^{g^{(A)}} = 3$; $\dim V(\alpha_5)^{g^{(A)}} = 8$ (recall that the weight system of $V(\alpha_5)$ consists of all weights of the form $\frac{1}{2}(\mu_1 \oplus \dots \oplus \mu_5)$ without multiplicities; $V(\alpha_5) \subset V(\alpha_5)^{g^{(A)}}$ if $(\mu_1 \oplus \mu_2) = (\mu_3 \oplus \mu_4) = 0$), $\dim \mathfrak{g}_V(g^{(A)}) = 15$. So (4.9) does not hold.

Step 5. $\mathfrak{g} = \mathfrak{so}_{13}$; $V = V(\alpha_6) \oplus V(\alpha_1)^2$. By [Sch2], the algebra $C[V]^G$ is freely generated by 12 elements $f_{(2;0;0)}; f_{(1;1;0)}; f_{(0;2;0)}; f_{(0;0;4)}; f_{(0;0;8)}; f_{(1;0;4)}; f_{(0;1;4)}; f_{(2;0;4)}; f_{(1;1;4)}; f_{(0;2;4)}; f_{(1;1;2)}; f_{(1;1;6)}$, where the lower index indicates the grading with respect to the decomposition $V = V(\alpha_1) \oplus V(\alpha_1) \oplus V(\alpha_6)$. Note that $\mathrm{Sp}(V)^G = \mathrm{SL}_2$. The elements $f_{(0;0;4)}; f_{(0;0;8)}; f_{(1;1;2)}; f_{(1;1;6)}$ are SL_2 -invariant, $\mathrm{Span}_C(f_{(2;0;0)}; f_{(1;1;0)}; f_{(0;2;0)})$; $\mathrm{Span}_C(f_{(2;0;4)}; f_{(1;1;4)}; f_{(0;2;4)}) = S^2 C^2$; $\mathrm{Span}(f_{(1;0;4)}; f_{(0;1;4)}) = C^2$.

Analogously to the previous case (i.e., using the grading and the SL_2 -module structure of $C[V^{\mathbb{F}}]$), we check that $f_{(0;0;4)}; f_{(0;0;8)} \in z(C[V^{\mathbb{F}}])$. Let $\varphi_1; \varphi_2$ denote the moment maps for the actions $G \curvearrowright V(\varphi_6); G \curvearrowright V(\varphi_1)^2; V$. Clearly, $\varphi = \varphi_1 + \varphi_2$. Further, put $f_2(\varphi) = \text{tr}(\varphi^2); f_4(\varphi) = \text{tr}(\varphi^4)$; $\varphi \in \mathfrak{g}$ (the traces are taken in the tautological so_{13} -module). We have shown that $\varphi_1(f_2); \varphi_1(f_4) \in z(C[V^{\mathbb{F}}])$. On the other hand, $\varphi(f_2); \varphi(f_4) \in z(C[V^{\mathbb{F}}])$. Let us check that $\varphi_1(f_2); \varphi_1(f_4); \varphi(f_2); \varphi(f_4)$ are algebraically independent. Analogously to the previous case one checks that $\varphi_1(f_2); \varphi(f_2)$ are independent. It remains to check that the equality

$$(4.10) \quad a \text{tr}(\varphi_1^4) + b \text{tr}((\varphi_1 + \varphi)^4) + c \text{tr}(\varphi_1^2)^2 + d \text{tr}(\varphi_1^2) \text{tr}((\varphi_1 + \varphi)^2) + e \text{tr}((\varphi_1 + \varphi)^2)^2 = 0;$$

$$\varphi_1 \in \overline{\text{in}}_2; i=1;2;$$

implies $a = b = 0$. The stable subalgebras in general positions for $V(\varphi_6); V(\varphi_1)^2$ are $\mathfrak{g}^{(1;2;4;5)}; \mathfrak{g}^{(2;3;6)}$, respectively. By Lemma 2.24, $\overline{\text{in}}_1 = G \text{Span}_{\mathbb{C}}(\varphi_1 + \varphi_2 + \varphi_3; \varphi_4 + \varphi_5 + \varphi_6);$
 $\overline{\text{in}}_2 = G \text{Span}_{\mathbb{C}}(\varphi_1)$. Since $\text{tr}(\varphi_1^4)$ and $\text{tr}(\varphi_1^2)^2$ are not proportional, we get $a + b = 0$. Writing down the terms of (4.10) of bidegree $(3,1)$ with respect to $(\varphi_1; \varphi_2)$, we get

$$4b \text{tr}(\varphi_1^3 \varphi_2) + 2d \text{tr}(\varphi_1^2) \text{tr}(\varphi_1 \varphi_2) + 4e \text{tr}(\varphi_1^2) \text{tr}(\varphi_1 \varphi_2) = 0;$$

Putting $\varphi_1 = \varphi_1 + \varphi_2 + \varphi_3 + i(\varphi_4 + \varphi_5 + \varphi_6); \varphi_2 = \varphi_1$, we see that $b = 0$.

To prove that the group $W_{G/X}^{(\varphi)}$ has the form indicated in Table 4.2 it is enough to check that $S^{(A)} \subset \mathfrak{g} \curvearrowright V$ for $A = f\varphi_1 - \varphi_2; \varphi_3 - \varphi_4; \varphi_5 - \varphi_6$. This is done analogously to the previous case.

5. Fibers of $b_{G/X}$ and untwisted varieties

Throughout this section $G; X$ are such as in the previous one.

The goal of this section is to prove Theorem 1.4 and establish some examples of untwisted varieties. Subsection 5.1 contains some technical results used in the proof of Theorem 1.4. The proof itself is given in Subsection 5.2. In Subsection 5.3 we describe some classes of untwisted Hamiltonian varieties. We state a result by Knop that the cotangent bundle of an affine variety is untwisted and show that any symplectic module is untwisted. Finally, in Subsection 5.4 we give two counterexamples: of a Hamiltonian variety not satisfying (Irr) and of a conical coisotropic model variety not satisfying (Utw2). The former counterexample is due to F. Knop.

5.1. Reducedness of fibers of $b_{G/X}$.

Proposition 5.1. Let $\varphi \in \mathfrak{a}_{G/X}^{(\varphi)}$ be such that $(W_{G/X}^{(\varphi)}) = \text{flg}$. Then the fiber $b_{G/X}^{-1}(\varphi) = W_{G/X}^{(\varphi)} \curvearrowright \mathfrak{a}_{G/X}^{(\varphi)}$ is reduced.

Proof. We preserve the notation of Proposition 4.1 and Remark 4.8 and put $\mathfrak{b} = M$. It is enough to prove that $b_{G/X}$ is smooth at any point $x \in X$ satisfying conditions (a)–(e) of Proposition 4.1 for any component Z of the fiber in interest. By Remark 4.8, $W_{M/X_M}^{(\varphi)} = \text{flg}$. Using the commutative diagram of Remark 4.8, we see that it is enough to prove the proposition in the case when $\varphi = 0$ and $X = M_G(H; V)$, where φ is nilpotent, $\text{cork}_G(X) = 0$, and $W_{G/X}^{(\varphi)} = \text{flg}$. In this case $\mathfrak{a}_{G/X}^{(\varphi)} = W_{G/X}^{(\varphi)} = X = G$. Let $X_L; L; G_0; X_0; L_0$ be such as in Proposition 3.3. An element of $N_G(L_0)$ acting trivially on $\mathfrak{a}_{G/X}^{(X_L)}$ lies in $Z_G(\mathfrak{a}_{G/X}^{(X_L)}) = L$. Thus (see [Lo5], Lemma 6.9) $W_{G_0/X_0}^{(\varphi)} = W_{G_0/X_0}^{(\varphi)} = G_0 = G_0$. So G_0 is connected. There is a point

$y \in X_0$ of the form $[g; 0]$, we may assume that $g = e$. It follows directly from Example 2.5 that $X_0 = M_{G_0}(N_H(L_0) = L_0; \cdot; V^{L_0})$. Note that e is a nilpotent element of \mathfrak{g}^{L_0} so $e \in \mathfrak{g}^{L_0} \setminus \mathfrak{l}_0^\perp = \mathfrak{g}_0$. By assertion 1 of Proposition 3.3, it is enough to check that the fiber $\pi_{G_0; X_0}^{-1}(0)$ is generically reduced. Replacing $(G; X)$ with $(G_0; X_0)$ we may assume, in addition, that X satisfies the equivalent conditions of Lemma 2.23. It follows that X satisfies condition (*) of Proposition 4.9.

We may replace G with a covering and assume that $G = T_0 \ltimes H$, where T_0 is a torus. Further, by assertion 4 of Lemma 2.30, $\widehat{W}_{G; X}^{(\cdot)} = \text{flg}$, where $X^\circ = M_G(H; \cdot; V)$. So we may replace X with X° and assume that H is connected. Since in this case $X = T(T_0) \ltimes V$, we reduce to the case $H = G; X = V$. Changing G by a covering again, we may assume that $G = (G; G) \ltimes Z$, where Z is a torus.

Recall that $a_{G; V} = V = G$. The required claim will follow if we show that the zero fiber of the morphisms $\pi_{(G; G); V} : Z \times V \rightarrow (G; G)$ are reduced. For the former morphism this stems easily from the decomposition $V = V^{(G; G)} \oplus \bigoplus_{i=1}^k V_i$. Put, for brevity, $U_1 = V^{(G; G)}; U_2 = \bigoplus_{i=1}^k V_i$. Note that $\text{Sp}(U_2)^{(G; G)}$ is a torus of dimension k acting trivially on $U_2 = (G; G)$. So it remains to prove that $\pi_{Z; U_1}^{-1}(0)$ are reduced. Since $\text{cork}_G(V) = 0$, we have $\dim V = 4k + 2 \dim Z$. It follows that $\dim U_2 = 2 \dim Z$. Further, by above, $k + \dim Z = \dim V = G = \dim U_1 = Z + \dim U_2 = (G; G)$, whence $\dim U_1 = Z = \dim Z$. Since $C(U_1)^Z = \text{Quot}(C[U_1]^Z)$ (see [Lo2], Theorem 1.10, for the proof in the general case), it follows that Z acts on U_1 locally effectively. Thus the weight system of Z in U_1 coincides with $\epsilon_1; \dots; \epsilon_r; -\epsilon_1; \dots; -\epsilon_r$, where $\epsilon_1; \dots; \epsilon_r$ form a basis in \mathfrak{t} . Now the claim is easy.

Proposition 5.2. Suppose $0 \in \pi_{G; X}^{-1}$, so that $0 \in a_{G; X}^{(\alpha_L)}$. Let $s \in W_{G; X}^{(\cdot)}$ be a reflection. Put $s = (a_{G; X}^{(\cdot)})^s, D_s = \pi_{W_{G; X}^{(\cdot)}; a_{G; X}^{(\cdot)}}^{-1}(s)$. Let \mathcal{F} be an irreducible component of $b_{G; X}^{-1}(D_s)$. Let x be a general point in $s, M = Z_G(\cdot)$, and $x \in \mathcal{F}$ satisfy conditions (a)–(f) of Propositions 4.1, 4.3. In the notation of those propositions put $\mathcal{P} = T_0(M; M)$ and let \mathcal{X} be such as in (e) of Proposition 4.1. Then the multiplicity of \mathcal{F} in $b_{G; X}^{-1}(D_s)$ equals 1 or 2 and the latter is the case if $\widehat{W}_{G; X}^{(\cdot)} = \text{flg}$.

Proof. We use the notation of Propositions 4.1, 4.3 and Remark 4.8. From the choice of M it follows that $s \in W_{G; X_M}^{(\alpha_L)}$. Put $s_0 = (a_{G; X_M}^{(\alpha_L)})^s$; and let D_s^0 be the image of s_0 in $a_{G; X_M}^{(\widehat{\alpha_L})} = W_{G; X_M}^{(\widehat{\alpha_L})}$. Let \mathcal{F}_M be an irreducible component of $\mathcal{F} \setminus X_M$ containing x . Also it is an irreducible component of $b_{G; X_M}^{-1}(\pi_{W_{G; X_M}^{(\alpha_L)}; a_{G; X_M}^{(\alpha_L)}}^{-1}(s_0))$. Let \mathcal{P}^0 denote an irreducible component of $b_{G; X_M}^{-1}(D_s^0)$ containing a connected component of $0 \setminus \mathcal{F}_M$. Clearly, $\mathcal{P}^0 = \mathcal{P} \cap V^H$ for some subvariety $\mathcal{P} \subset \mathcal{X}$, which is an irreducible component of $b_{G; X}^{-1}(D_s^0)$. From the commutative diagram of Remark 4.8 one deduces that precisely one of the following possibilities holds:

- (1) $\widehat{W}_{G; X}^{(\cdot)} = Z_2$. The multiplicity of \mathcal{F} in $b_{G; X}^{-1}(D_s)$ equals the multiplicity of \mathcal{P} in $b_{G; X}^{-1}(D_s^0)$.
- (2) $\widehat{W}_{G; X}^{(\cdot)} = \text{flg}$. By Proposition 5.1, the multiplicity of \mathcal{P} in $b_{G; X}^{-1}(D_s^0)$ is one. Further, the multiplicity of \mathcal{F} in $b_{G; X}^{-1}(D_s)$ is 2.

It remains to consider the first possibility. Note that, by definition of M , one gets $s_z(m)$ whence $\frac{0}{s} = z(\mathfrak{g})$. Since $W_{\widehat{G}; \widehat{X}}^{(\alpha_L)} \notin \text{flg}$, it follows that $\dim a_{\widehat{G}; \widehat{X}}^{(\alpha_L)} \setminus [\mathfrak{g}; \mathfrak{g}] = 1$. Further, note that T_0 acts trivially on $\widehat{\mathcal{X}}_L$. Since $a_{\widehat{G}; \widehat{X}}^{(\alpha_L)} = z(\mathfrak{h}) \setminus t_0^2$ and t_0 projects surjectively to $z(\mathfrak{g})$ (recall that $\mathfrak{g} = t_0 + [\mathfrak{m}; \mathfrak{m}]$), we see that $a_{\widehat{G}; \widehat{X}}^{(\alpha_L)} \setminus z(\mathfrak{g}) = f_0 \mathfrak{g}$. Therefore $a_{\widehat{G}; \widehat{X}}^{(\alpha_L)}[\mathfrak{g}; \mathfrak{g}]; \dim a_{\widehat{G}; \widehat{X}}^{(\alpha_L)} = 1; Z(\mathcal{C}) = T_0$. From the last inclusion it follows that $Z(\mathcal{C})$ acts trivially on $\widehat{\mathcal{X}}$.

Replacing $(G; X)$ with $((\mathcal{C}; \mathcal{C}); \widehat{\mathcal{X}})$, we reduce the problem to the proof of following claim.

(**) Suppose that $X = M_G(H; V)$, G is semisimple, \mathfrak{h} is nilpotent, $\text{cork}_G(X) = 0$, $\dim a_{G; X}^{(\alpha)} = 1; W_{G; X}^{(\alpha)} = \text{flsg}$, where s is a reflection. Then the fiber $b_{G; X}^{-1}(0)$ is reduced.

As in the proof of Proposition 5.1, we see that $b_{G; X} = \frac{1}{G; X}$. So it is enough to check that $\frac{1}{H; U \cup V}(0)$ is reduced, where $U = (z_{\mathfrak{g}}(\cdot) = \mathfrak{h})$. Note that $C[U \cup V]^{\mathfrak{h}}$ is generated by an element of degree 4. Recall that $\frac{2}{U}(\cdot)^{\mathfrak{h}} \subset C[U \cup V]^{\mathfrak{h}}$ is of degree 4. So if $\mathfrak{g} \notin \mathfrak{h}$, we are done. If $\mathfrak{g} \in \mathfrak{h}$, then there is an element $q \in C[\mathfrak{g} = \mathfrak{h}]^{\mathfrak{h}}$ corresponding to an H -invariant quadratic form on $\mathfrak{g} = \mathfrak{h} = \mathfrak{h}^2$. It has degree 4. Any fiber of q is reduced, since $\dim \mathfrak{g} = \mathfrak{h} > 1$. Therefore it remains to consider the case $H = G; X = V$. The reducedness of fibers in this case follows from the observation that a homogeneous generator of $C[V]^{\mathfrak{g}}$ is irreducible.

Proposition 5.3. Again, we keep the notation of Proposition 4.1 and Remark 4.8. Suppose X is untwisted and $0 \in b_{G; X}$. Let x be a point satisfying conditions (a)–(d) of Proposition 4.1 for a point $o \in a_{G; X}^{(\alpha_L)}$. Then the Hamiltonian \mathcal{C} -variety $\widehat{\mathcal{X}}$ is untwisted and $W_{\widehat{G}; \widehat{X}}^{(\alpha_L)} = (W_{G; X}^{(\alpha_L)})_o$.

Proof. At first we consider the case when $\mathcal{C} = M$. Let s be a reflection lying in $(W_{G; X}^{(\alpha_L)})_o$. Let $\frac{s}{s}; D_s$ have the same meaning as in Proposition 5.2 and D_s^0 denote the image of $\frac{s}{s}$ in $a_{M; X}^{(\alpha_L)} = W_{M; X}^{(\alpha_L)}$ (we write $\widehat{\mathcal{X}}_L$ instead of $\widehat{\mathcal{X}}_{\widehat{L}}$ because $L = \mathfrak{h}$). By Proposition 5.1, it is enough to show that $b_{M; X}^{-1}(D_s^0)$ is reduced. Note that the last scheme is non-empty because $b_{M; X}^{-1}(x) \in D_s^0$. Choose a component \mathcal{Z} of $b_{M; X}^{-1}(D_s^0)$. Note that $b_{G; X}(\mathcal{Z} \setminus 0) \subset D_s$. Choose a component Z of $b_{G; X}^{-1}(D_s)$ containing a connected component of $\mathcal{Z} \setminus 0$. Since $\mathcal{C} = M$, we see that the morphism $a_{M; X}^{(\alpha_L)} = W_{M; X}^{(\alpha_L)} \rightarrow a_{\widehat{G}; \widehat{X}}^{(\alpha_L)} = W_{\widehat{G}; \widehat{X}}^{(\alpha_L)}$ from the commutative diagram of Remark 4.8 can be inverted. So one can consider the morphism $b : \widehat{\mathcal{X}} \rightarrow a_{G; X}^{(\alpha_L)} = W_{G; X}^{(\alpha_L)}$ from this diagram. It follows from the diagram that the multiplicities of \mathcal{Z} in $b^{-1}(D_s)$ and of Z in $b_{G; X}^{-1}(D_s)$ coincide. By (Utw2) the latter is one. It follows that the multiplicity of \mathcal{Z} in $b_{M; X}^{-1}(D_s^0)$ is 1 and that the morphism $a_{M; X}^{(\alpha_L)} = W_{M; X}^{(\alpha_L)} \rightarrow a_{G; X}^{(\alpha_L)} = W_{G; X}^{(\alpha_L)}$ is unramified in D_s^0 . The latter is equivalent to $s \in W_{M; X}^{(\alpha_L)}$. Since $W_{G; X}^{(\alpha_L)}$ is generated by reflections, so is $(W_{G; X}^{(\alpha_L)})_o$. This completes the proof when $\mathcal{C} = M$.

Proceed to the general case.

Lemma 5.4. Let $X_1; X_2$ be Hamiltonian G -varieties and $\gamma : X_1 \rightarrow X_2$ a Hamiltonian G -morphism. If X_2 is untwisted, then so is X_1 and the natural morphism $\gamma_0 : C_{G; X_1} \rightarrow C_{G; X_2}$ induced by γ is étale.

Changing M by a covering, we may assume that $M = \mathcal{B} \times T_0$, where T_0 is a torus. There is a natural Hamiltonian morphism $\pi: T \times (T_0) \rightarrow M_M(H; V)$. By Lemma 5.4, $T \times (T_0) \rightarrow M_M(H; V)$ is untwisted and the natural morphism $C_{M, T \times (T_0)} \xrightarrow{\pi^*} C_{M, M_M(H; V)}$ is étale. From Proposition 2.32 it follows that the Weyl groups of $M_M(H; V)$ and $T \times (T_0) \rightarrow M_M(H; V)$ coincide. This implies all required claims.

Lemma 5.5. *If X is an affine Hamiltonian variety satisfying (Utw1), then all fibers of $e_{G,X} = G$ are Cohen-Macaulay (as schemes).*

Proof of assertion 1. Recall that $C_{G/\mathbb{A}^1} = a_{G/\mathbb{A}^1}^{(1)} = W_{G/\mathbb{A}^1}^{(1)}$, Proposition 2.32. Thanks to Lemma 5.5, it remains to prove that any fiber of $b_{G/\mathbb{A}^1} = G$ is smooth in codimension 1. Since X is conical (of degree, say, k), there are actions $C \curvearrowright X = G$; $a_{G/\mathbb{A}^1}^{(1)} = W_{G/\mathbb{A}^1}^{(1)}$ such that the former extends to a morphism $C \curvearrowright X = G \rightarrow X = G$, the latter is induced by $C \curvearrowright a_{G/\mathbb{A}^1} \rightarrow a_{G/\mathbb{A}^1}$; $(t; v) \mapsto t^k v$, and the morphism $b_{G/\mathbb{A}^1} = G$ is C -equivariant. Applying a standard argument, we see that it is enough to prove that $b_{G/\mathbb{A}^1} = G^{-1}(0)$ is smooth in codimension 1.

$$V^H \quad \begin{array}{c} \text{X}=\text{G} \quad \text{---} \quad \text{O}=\text{G} \quad \text{---} \quad \text{X}=\text{G} \\ \text{a} \\ \text{pr}_2 \\ \text{a} \\ \text{a}_{\text{G};\text{X}}^{(\cdot)} = \text{w}_{\text{G};\text{X}}^{(\cdot)} \end{array} \quad \text{b}_{\text{G};\text{X}}=\text{G}$$

Proof of assertion 2. Choose a nonzero point $2 \in a_{G/X}^{(1)}$ and put $a_0 := C$. Further, put $Y := (X=G) \cap a_{G/X}^{(1)} = W_{G/X}^{(1)} \cap a_{G/X}^{(1)}$; $Y_0 := (X=G) \cap a_{G/X}^{(1)} = W_{G/X}^{(1)} \cap a_0$ (here a_0 maps to $a_{G/X}^{(1)} = W_{G/X}^{(1)}$ via the composition $a_0 \xrightarrow{\iota} a_{G/X}^{(1)} \xrightarrow{\iota} a_{G/X}^{(1)} = W_{G/X}^{(1)}$).

Let us check that Y is normal (as a scheme) and Cohen-Macaulay. Indeed, the morphism $a_{G/X}^{(1)} : a_{G/X}^{(0)} \rightarrow W_{G/X}^{(1)}$ is flat, since X satisfies (Utw1). Therefore the morphism $Y \rightarrow X = G$ is flat. But, as we have already remarked, $X = G$ is Cohen-Macaulay. By Corollary 18.17 from

$[E]$, Y is Cohen-Macaulay. Note that $X = G$ is smooth in codimension 1 over $a_{G,X}^{(1)} = W_{G,X}^{(1)}$. Hence Y is smooth in codimension 1 over $a_{G,X}^{(1)}$ hence normal.

Again, being a complete intersection in a Cohen-Macaulay variety, Y_0 is Cohen-Macaulay. Similarly to the previous paragraph, Y_0 is normal.

Let us show that $C[a_0]$ is integrally closed in $C[Y_0]$. Let \mathfrak{a}_0 denote the spectrum of the integral closure of $C[a_0]$ in $C[Y]$. There is the action of C on Y_0 lifted from $C : X = G$. The morphism $Y_0 \rightarrow \mathfrak{a}_0$ is C -equivariant. Therefore there is an action $C \curvearrowright \mathfrak{a}_0$ contracting \mathfrak{a}_0 to a unique point over $0 \in \mathfrak{a}_0$. It follows that $\mathfrak{a}_0 = C$. Since the zero fiber of the morphism $Y \rightarrow \mathfrak{a}_0$ is normal, we see that the morphism $\mathfrak{a}_0 \rightarrow \mathfrak{a}_0$ is étale in 0. From the C -equivariance it follows that it is an isomorphism.

Thus a general fiber of the morphism $Y_0 \rightarrow \mathfrak{a}_0$ is irreducible. Thanks to the presence of C -action, the same is true for any fiber but the zero one. It follows easily from (Con1), (Con2) that $b_{G,X} = G^{-1}(0)$ is connected. Since $b_{G,X}^1 = G^{-1}(0)$ is normal, it is irreducible.

Proof of assertion 3. By Proposition 2.32, X satisfies (Utw1). Assume that X does not satisfy (Utw2). By Proposition 5.1, there is $s \geq 2$ $W_{G,X}^{(1)}$ such that some irreducible component $\mathcal{E} \subset b_{G,X}^1(D_s)$ (where, as above, D_s denotes the image of $(a_{G,X}^{(1)})^s$ in $a_{G,X}^{(1)} = W_{G,X}^{(1)}$) is of multiplicity 2. Put $Y = b_{G,X}^{-1}(\mathcal{E})$. By Proposition 2.31, Y is an irreducible component of $b_{G,X} = G^{-1}(D_s)$ and the set of closed orbits of any two components $\mathcal{E}_1, \mathcal{E}_2 \subset b_{G,X} = G^{-1}(D_s)$ is the same. It follows from (Irr) that $Y = b_{G,X} = G^{-1}(D_s)$. By Proposition 5.2, the multiplicity of any component \mathcal{E}_1 in $b_{G,X}^1(D_s)$ is 2. Let $f \in C[a_{G,X}]^{W_{G,X}}$ be such that $(f) = D_s$. Let us remark that f is not a square in $C[X]$. Assume the converse, let $f = f_1^2$; $f_1 \in C[X]$. Then $f_1 \in C[C_{G,X}] = C[a_{G,X}^{(1)}]^{W_{G,X}^{(1)}}$ which is absurd.

Put $\mathcal{A} = C[X][t] = (t^2 - f)$. There is the natural morphism of schemes $\pi : \text{Spec}(\mathcal{A}) \rightarrow X$. This morphism is unramified over $X \setminus b_{G,X}^1(D_s)$. Note also that the group Z_2 acts on $\text{Spec}(\mathcal{A})$ so that π is the quotient for this action. Hence the restriction of π to any irreducible component of $\text{Spec}(\mathcal{A})$ is dominant. Since f is not a square in $C[X]$, we see that $\text{Spec}(\mathcal{A})$ is irreducible. Recall that $(f) = 2D$ for some D . It follows that π is unramified over $b_{G,X}^1(D_s)$. So π is étale in codimension 1. It follows from the Zariski-Nagata theorem that π is étale. So it is an isomorphism.

5.3. Some classes of untwisted varieties.

Proposition 5.6. Let X be coisotropic, simply connected and conical (e.g. a symplectic vector space). Then X is untwisted.

Proof. Thanks to Proposition 2.32, X satisfies (Utw1). Furthermore, X obviously satisfies (Irr). Applying assertion 3 of Theorem 1.4, we complete the proof.

Theorem 5.7. Let X_0 be a smooth irreducible affine G -variety. Then T^*X_0 is an untwisted Hamiltonian G -variety.

Proof. (Utw1) is checked in Satz 6.6 of [Kn1]. (Utw2) follows from [Kn3], Corollary 7.6.

For a sketch of an alternative proof see Remark 5.11.

Theorem 5.8. Let V be a symplectic G -module. Then V is an untwisted Hamiltonian G -variety.

We will prove this theorem after some auxiliary considerations.

Proposition 5.9. Let X be a conical Hamiltonian G -variety and $G_0; X_0$ be such as in the discussion preceding Proposition 3.3. If the Hamiltonian G_0 -variety X_0 satisfies (Irr), then so does X .

Proof. The action $C : X$ preserves X_0 and so gives rise to the structure of a conical Hamiltonian G_0 -variety on X_0 . By Proposition 3.3, the following diagram, where the horizontal arrows are quotient morphisms for the actions $G_0 = G_0$ on $X_0 = G_0; a_{G_0; X_0}^{()} = W_{G_0; X_0}^{()}$, is commutative.

$$\begin{array}{ccc} X_0 = G_0 & \xrightarrow{\quad} & X = G \\ \downarrow b_{G_0; X_0} & & \downarrow b_{G; X} \\ a_{G_0; X_0}^{()} = W_{G_0; X_0}^{()} & \xrightarrow{\quad} & a_{G; X}^{()} = W_{G; X}^{()} \end{array}$$

Choose $\alpha \in a_{G; X}^{()} = W_{G; X}^{()}$ and a point $\beta \in a_{G_0; X_0}^{()} = W_{G_0; X_0}^{()}$ mapping to α . By the previous commutative diagram, $b_{G; X} = G^{-1}(\alpha)$ is the quotient of $b_{G_0; X_0} = G_0^{-1}(\beta)$ by some finite group. In particular, $b_{G; X} = G^{-1}(\alpha)$ is irreducible.

Proposition 5.10. Let X be an irreducible conical affine Hamiltonian G -variety such that $\dim a_{G; X}^{()} = \text{rk } G$. Let $G = Z(G) \cdot G_1 \cdots G_k$ be the decomposition of G into the locally direct product of simple normal subgroups and the unit component of the center. If X satisfies (Utw1) and is untwisted as a Hamiltonian G_i -variety for any i , then X is untwisted as a Hamiltonian G -variety. Conversely, if X is untwisted as a Hamiltonian G -variety, then so is it as a Hamiltonian G_i -variety.

Proof. This follows easily from assertion 5 of Lemma 2.30.

Proof of Theorem 5.8. Applying Proposition 5.9, we reduce the proof to the case when V satisfies the equivalent conditions of Lemma 2.23. Further, thanks to Proposition 5.10, we may (and will) assume that G is simple.

Suppose V is not untwisted. By Proposition 2.32, $b_{G; V}$ is not smooth in codimension 1. By Proposition 5.2, in the notation of that proposition, for some $s \in W_{G; V}^{()}$ there is a point $x \in b_{G; V}^{-1}(D_s)$ satisfying conditions (a)–(f) with $W_{G; X}^{()} = \text{flg}$. From Remark 4.4 it follows that one can take $(M; M)$ for \mathfrak{g} . Note that $\mathfrak{g} = \mathfrak{sl}_2$. By Proposition 4.9, $g_x = \mathfrak{sl}_2; T_x V = (\mathfrak{g} \cdot x + T_x V^{g_x}) = C^2 \oplus C^2$ (here C^2 denotes the irreducible two-dimensional \mathfrak{sl}_2 -module). All modules V containing such a point x are presented in Table 4.2, see the proof of Proposition 4.21.

Let us choose a point $x \in b_{G; V}^{-1}(0)$ satisfying conditions (a)–(e) of Proposition 4.1. Let \mathcal{X} be the corresponding model G -variety. Note that $b_{G; V} = G^{-1}(0)$ is irreducible (even if V does not satisfy (Irr), see the end of the proof of assertion 3 of Theorem 1.4) whence \mathcal{X} does not depend on the choice of x . If $S^{(1)} \subset \mathfrak{g} \cdot \mathcal{X}$ for some $s \in \mathfrak{g}$, then $S^{(1)} \subset V$. From the proof of Proposition 4.21 it follows that $S^{(1)} \subset \mathfrak{g} \cdot V$ implies $s_w \in W_{G; V}^{()}$ for some $w \in W(\mathfrak{g})$. It is not difficult to deduce from Proposition 4.17 that $W_{G; X}^{()} = W_{G; V}^{()}$. It follows from the commutative diagram of Remark 4.8 that the fiber $b_{G; V} = G^{-1}(0)$ is smooth in x .

By Proposition 3.3, this fiber is smooth in codimension 1. Proceeding as in the proof of assertion 1 of Theorem 1.4, we see that any fiber of $b_{G,V} = G$ is normal. Applying assertions 2,3 of Theorem 1.4, we see that V is untwisted.

Remark 5.11. This remark contains a sketch of an alternative proof of Theorem 5.7. Let $X = T X_0$, where X_0 is a smooth affine G -variety. Applying Propositions 5.9, 5.10 we see that it is enough to assume that G is simple and $a_{G,X} = t$. Let $x_0 \in X_0$ be a point with closed G -orbit. Put $X_0^0 = G_{x_0} (T_{x_0} X_0 = g_{x_0})$; $X^0 = T X_0^0$. By Satz 6.5 from [Kn1], $a_{G,X}^{(t)} = a_{G,X^0}^{(t)}$; $W_{G,X}^{(t)} = W_{G,X^0}^{(t)}$. Applying the Luna slice theorem, we see that the statements of Theorem 5.7 for $X; X^0$ are equivalent. So we may assume that X_0 is a homogeneous vector bundle. It follows from Theorem 7.1.2 in [Lo4] that $S^{(t)}_{g,X} \cong S^{(t)}_{g,X^0}$ for some $w \in W$. Now the proof is analogous to the normal part of the proof of Theorem 5.8.

5.4. Some counterexamples. The following example of a Hamiltonian variety not satisfying (Irr) belongs to F. Knop.

Example 5.12. Put $G = C^2$; $X = C^2 \times C^2 \times C^2$. Choose coordinates x_1, \dots, x_4 on X corresponding to the above decomposition. Define the action $G : X$ by $t(x_1; x_2; x_3; x_4) = (tx_1; t^{-1}x_2; x_3; x_4)$. Put $\omega = (x_1x_2^2 - x_1^{-1}x_3^2)dx_1 + x_4dx_3$. Clearly, ω is G -invariant. Put

$$\omega = d\theta = 2x_1x_2dx_1 \wedge dx_2 - 2x_1^{-1}x_3dx_1 \wedge dx_3 - dx_3 \wedge dx_4.$$

One checks directly that ω is nondegenerate. The action $G : X$ is Hamiltonian with $\theta_{G,X}(x) = h; \frac{\partial}{\partial t} h_x = x_1^2x_2^2 - x_3^2$. It is clear that $S^{(t)}_{G,X}(a)$ is irreducible whenever $a \notin f_0g$. It follows that $C_{G,X} = g$. On the other hand, $S^{(t)}_{G,X}(0)$ has two connected components.

We remark that the Hamiltonian variety in Example 5.12 is the smallest one in the sense that both group and variety have the smallest possible dimensions.

Now let us present an example of a coisotropic conical model variety $X = M_G(H; \cdot; V)$ that is not untwisted. An example, where the group $W_{G,X}^{(t)}$ is not generated by reflections, can be found in [Lo2], Subsection 5.11. In the following example $W_{G,X}^{(t)}$ is generated by reflections but (Utw2) does not hold. Note that this example is very similar to that from [Lo2].

Example 5.13. Put $G = SL_2 \times C^2$; $X = M_G(\mathbb{Z}_2 \times SL_2; 0; C^2 \times C^2)$, where C^2 denotes the two-dimensional irreducible SL_2 -module with the symplectic form given by $(u;v) \mapsto \det(u;v)$ and the nontrivial element $\gamma \in \mathbb{Z}_2 \times C^2$ acts on $C^2 \times C^2$ as follows: $(v_1; v_2) \mapsto (v_2; -v_1)$. One easily checks that $W_{G,X}^{(t)} = N_G(T) = T = \mathbb{Z}_2$ and (Utw2) does not hold.

6. Some open problems

Firstly, we state two conjectures concerning property (Irr). Below G is a connected reductive group.

Conjecture 6.1. Any conical irreducible Hamiltonian G -variety X satisfies (Irr).

The following conjecture is a weaker version of the first one.

Conjecture 6.2. $X = M_G(H; \cdot; V)$, where γ is nilpotent, satisfies (Irr).

In virtue of the local cross-section and symplectic slice theorems (Propositions 2.20, 2.27) one can deduce from Conjecture 6.2 that any fiber of $b_{G,X}$ is normal (as a variety).

Unlike the first conjecture, the second one can be reduced to some case-by-case consideration. Let us sketch the scheme of this reduction.

At first, one reduces the problem to the case when X satisfies the equivalent conditions of Lemma 2.23 and then to the case when X is algebraically simply connected. Here one should check that X satisfies (Utw2). This will follow if one verifies the following assertion:

(*) for any $g \in G$ such that $S^{(1)}_g X$ there is $w \in W(g)$ such that $s_w \notin W_{G/X}^{(1)}$.

Finally, it is enough to check (*) only for some special quadruples $(G; H; \cdot; V)$. By analogy with Section 7 of [Lo4], we call such triples quasiessential. By definition, a quadruple $(G; H; \cdot; V)$ is quasiessential if $M_G(H; \cdot; V)$ satisfies the equivalent conditions of Lemma 2.23 and for any ideal $h_1 \subset h$ there is $g \in G$ such that $S^{(1)}_g M_G(H; \cdot; V)$ but $S^{(1)}_g M_G(H_1; \cdot; V)$, where H_1 is a subgroup of H corresponding to h_1 . It is not very difficult to show that if $(G; H; \cdot; V)$ is quasiessential, then G is simple and H is semisimple.

The next conjecture strengthens assertion 1 of Theorem 1.4.

Note that any fiber of $e_{G/X} = G$ has the natural structure of a Poisson variety. The open stratum (in the sense of Subsection 3.2) is symplectic.

Conjecture 6.3. Let $X = M_G(H; \cdot; V)$, where \cdot is nilpotent, be untwisted. Then any fiber Y of $e_{G/X} = G$ has symplectic singularities. This means that there is a resolution of singularities $\tilde{Y} \rightarrow Y$ such that the symplectic form on the smooth part of Y is extended to some regular form on \tilde{Y} .

Finally, we would like to propose a conjecture giving an estimate on dimensions of fibers of $e_{G/X}$.

Conjecture 6.4. Let X be an irreducible affine Hamiltonian G -variety. Then $\dim_{G/X}^{(1)}(\cdot) \leq \dim X - (\dim_G(X) + \text{def}_G(X) + \dim G) = 2$.

If X is the cotangent bundle of a (not necessarily affine) G -variety X_0 this conjecture follows from Vinberg's theorem on the modality of the action of a Borel subgroup of G on X_0 , see [V1].

7. Notation and conventions

For an algebraic group denoted by a capital Latin letter we denote its Lie algebra by the corresponding small German letter. For roots and weights of semisimple Lie algebras we use the notation of [PV].

G	the equivalence relation induced by an action of a group G .
$\text{cork}_G(X)$	the corank of a Hamiltonian G -variety X .
$\text{def}_G(X)$	the defect of a Hamiltonian G -variety X .
e	a nonzero element of the root subspace $\mathfrak{g} = \mathfrak{g}$.
(f)	the zero divisor of a rational function f .
$(G; G)$ (resp., $[\mathfrak{g}; \mathfrak{g}]$)	the commutant of a group G (resp., of a Lie algebra \mathfrak{g})
G	the connected component of unit of an algebraic group G .
$G_H V$	the homogeneous bundle over G/H with a fiber V .
$[g; v]$	the equivalence class of $(g; v)$ in $G_H V$.
G_x	the stabilizer of $x \in X$ under an action $G : X$.

\mathfrak{g}	the root subspace of \mathfrak{g} corresponding to a root α .
$\mathfrak{g}^{(A)}$ (resp., $G^{(A)}$)	the subalgebra \mathfrak{g} generated by \mathfrak{g}^α ; $2A \cap A$ (resp., the corresponding connected subgroup of G).
$m_G(X)$	$= \max_{x \in X} \dim Gx$.
$N_G(H)$, (resp., $N_G(\mathfrak{h}); n_{\mathfrak{g}}(\mathfrak{h})$)	the normalizer of an algebraic subgroup H in an algebraic group G (resp., of a subalgebra $\mathfrak{h} \subset \mathfrak{g}$ in G , of a subalgebra $\mathfrak{h} \subset \mathfrak{g}$ in \mathfrak{g}).
$\text{Quot}(A)$	the fraction field of A .
$\text{rk}(G)$	the rank of an algebraic group G .
s	the reflection in a Euclidian space corresponding to a vector α .
$\text{Span}_F(A)$	the linear span of a subset A of a module over a field F .
$\text{tr.deg } A$	the transcendence degree of an algebra A .
U^\perp	the skew-orthogonal complement to a subspace $U \subset V$ of a symplectic vector space V .
$V^{\mathfrak{g}}$	$= \{v \in V \mid \mathfrak{g}v = 0\}$, where \mathfrak{g} is a Lie algebra and V is a \mathfrak{g} -module.
$V(\lambda)$	the irreducible module of the highest weight λ over a reductive algebraic group or a reductive Lie algebra.
$W(\mathfrak{g})$	the Weyl group of a reductive Lie algebra \mathfrak{g} .
X^G	the fixed point set for an action $G : X$.
$X // G$	the categorical quotient for an action $G : X$, where G is a reductive group and X is an affine G -variety.
$Z(G)$ (resp., $z(\mathfrak{g})$)	the center of an algebraic group G (resp., a Lie algebra \mathfrak{g}).
$Z_G(H)$, (resp., $Z_G(\mathfrak{h}); z_{\mathfrak{g}}(\mathfrak{h})$)	the centralizer of a subgroup H in an algebraic group G (resp., of a subalgebra $\mathfrak{h} \subset \mathfrak{g}$ in G , of a subalgebra $\mathfrak{h} \subset \mathfrak{g}$ in a Lie algebra \mathfrak{g}).
$-\alpha$	the dual root of $\alpha \in \Delta(\mathfrak{g})$.
$\Phi(\mathfrak{g})$	the root system of a reductive Lie algebra \mathfrak{g} .
$G \curvearrowright X$	the moment map for a Hamiltonian G -variety X .
s (resp., n)	the semisimple (the nilpotent) part of an element of an algebraic Lie algebra.
$v(\alpha)$	the velocity vector field associated with $\alpha \in \Delta(\mathfrak{g})$.
$\Phi(\mathfrak{g})$	the system of simple roots for a reductive Lie algebra \mathfrak{g} .
$G \curvearrowright X$	the (categorical) quotient morphism $X \dashrightarrow X // G$.
$' = G$	the morphism of (categorical) quotients induced by a G -equivariant morphism $'$.
$'$	the homomorphism $C[X_2] \rightarrow C[X_1]$ induced by a morphism $' : X_1 \rightarrow X_2$.

References

- [AEV] E.M. Andreev, E.B. Vinberg, A.G. Elashvili, Orbits of highest dimension of semisimple linear Lie groups. *Func. anal. priblizh.*, 1(1967), N1, 3-7 (in Russian). English translation: *Funct. anal. appl.* v.1 (1968), p. 257-261.
- [Ch] C. Chevalley. *Fondements de la géométrie algébrique*. Paris, 1958.

- [D] E.B. Dynkin. Semisimple subalgebras of semisimple Lie algebras. *Mat. Sbornik*, 30 (1952), 349–462 (in Russian). English trans.: *Amer. Math. Soc. Transl. (Ser. 2)* v. 6 (1957), 111–245.
- [E] D. Eisenbud, *Commutative algebra with a view towards algebraic geometry*. GTM 150, Springer Verlag, 1995.
- [GS] V. Guillemin, Sh. Stenberg. *Symplectic techniques in physics*. Cambridge University Press, 1984.
- [Ki] F. Kirwan, Convexity properties of the moment mapping III. *Invent. Math.* 77 (1984), 547–552.
- [KL] F. Klop, P. Littmann, Der Grad erzeugender Funktionen von Invariantenringen. *Math. Z.* 196 (1987), 211–229.
- [Kn1] F. Klop. Weylgruppe und Momentabbildung. *Invent. Math.* 1990. V. 99. p. 1–23.
- [Kn2] F. Klop. The asymptotic behaviour of invariant collective motion. *Invent. Math.* 1994. V. 114. p. 309–328.
- [Kn3] F. Klop. Weyl groups of Hamiltonian manifolds, I. Preprint (1997). dg-ga/9712010.
- [Kn4] F. Klop. A connectedness property of algebraic moment maps. *J. Algebra*, v. 258 (2002), 122–136.
- [Kn5] F. Klop. Convexity of Hamiltonian manifolds. *J. Lie Theory*, 12 (2002), 571–582.
- [Kn6] F. Klop. Classification of multiplicity free symplectic representations. *J. Algebra*, 301 (2006), 531–553.
- [Kn7] F. Klop. Invariant functions on symplectic representations. Preprint (2005), arXiv math AG /0506171.
- [Kr] H. Kraft.: *Geometrische Methoden in der Invariantentheorie*. Vieweg, Braunschweig/Wiesbaden (1985).
- [Lo1] I.V. Losev. Coisotropic representations of reductive groups. *Trudy Moscov. Mat. Obshch.* 66 (2005), 157–181 (in Russian). English translation in: *Trans. Moscow Math. Soc.* (2005), 143–168.
- [Lo2] I.V. Losev. Algebraic Hamiltonian actions. Preprint (2006), arXiv math AG /0601023, 46 pages.
- [Lo3] I.V. Losev. Symplectic slices for reductive groups. *Mat. Sbornik* 197 (2006), N 2, 75–86 (in Russian). English translation in: *Sbornik Math.* 197 (2006), N 2, 213–224.
- [Lo4] I.V. Losev. Computation of Weyl groups of G -varieties. Preprint (2006), arXiv math AG /0612559, 83 pages.
- [Lo5] I.V. Losev. Combinatorial invariants of algebraic Hamiltonian actions. Preprint (2007), arXiv math AG /0701823, 23 pages.
- [LR] D. Luna, R.W. Richardson. A generalization of the Chevalley restriction theorem. *Duke Math. J.*, v. 46 (1980), N 3, p. 487–496.
- [Lu] D. Luna. Slices etales. *Bull. Soc. Math. France*, 33 (1973), p. 81–105.
- [MW] J. Marsden, A. Weinstein. Reduction of symplectic manifolds with symmetry. *Rep. Math. Phys.* 5 (1974), p. 121–130.
- [OV] A.L. Onishchik, E.B. Vinberg. *Seminar on Lie groups and algebraic groups*. Moscow, Nauka 1988 (in Russian). English translation: Berlin, Springer, 1990.
- [Pa] D.I. Panyushev. On the orbit spaces of finite and connected linear groups. *Izv. Akad. Nauk SSSR, Ser. Math.* 47 (1982), N 1, 95–99 (in Russian). English translation in: *Math. USSR, Izv.* 20 (1983), 97–101.
- [PV] V.L. Popov, E.B. Vinberg. Invariant theory. *Itogi nauki i tekhniki. Sovr. probl. matem. Fund. napr.*, v. 55. Moscow, VINITI, 1989, 137–309 (in Russian). English translation in: *Algebraic geometry 4, Encyclopaedia of Math. Sciences*, vol. 55, Springer Verlag, Berlin, 1994.
- [Sch1] G.W. Schwarz. Representations of simple Lie groups with regular rings of invariants. *Invent. Math.*, 49 (1978), pp. 167–191.
- [Sch2] G.W. Schwarz. Representations of simple groups with a free module of covariants. *Invent. Math.*, 50 (1978), p. 1–12.
- [Sch3] G.W. Schwarz. Lifting smooth homotopies of orbit spaces. *Publ. Math. IHES*, N 51 (1980), pp. 37–135.
- [V1] E.B. Vinberg. Complexity of an action of a reductive group. *Funktsional. Anal. i Prilozhen.*, 20 (1986), N 1, 1–13 (in Russian). English translation in: *Func. Anal. Appl.*, 20 (1986), p. 1–11.
- [V2] E.B. Vinberg.: Commutative homogeneous spaces and coisotropic symplectic actions. *Usp. Mat. Nauk* 56 (2001), 3–62 (in Russian). English translation in: *Russ. Math. Surveys* 56 (2001), 1–60.

Chair of Higher Algebra, Department of Mechanics and Mathematics, Moscow State University.

E-mail address: ivanlosev@yandex.ru